

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

185,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Non-classical Signalling Mechanisms in Stem Cells

Alice Pébay^{1,2}, Hitesh Peshavariya¹,
Raymond C.B. Wong³ and Gregory J Dusting^{1,4}

¹O'Brien Institute, Fitzroy, VIC 3065,

²Centre for Neuroscience, Department of Pharmacology, University of Melbourne,
Parkville VIC 3010,

³Department of Biological Chemistry, University of California Irvine, Irvine, CA 92617,

⁴Department of Surgery, University of Melbourne, Parkville VIC 3010,

^{1,2,4}Australia

³USA

1. Introduction

There are many gaps in our understanding of stem cell biology that need to be explored in order to realize the full potential of stem cells for therapies. Understanding how stem cells communicate within their microenvironment is important to be able to manipulate their functions. Indeed, the stem cell microenvironment is critical for their maintenance, and communication between neighbouring cells plays an important part in determining cell fate. Most studies of the stem cell niche focus on paracrine or juxtacrine cell interactions, particularly the influence of cytokines and various G-protein coupled receptor ligands, influences which have been reviewed elsewhere (Kobayashi et al 2010). Other signalling mechanisms that have received less attention include oxidant signalling through various protein kinase pathways, and intercellular communication through gap junctions. Not surprisingly gap junctional intercellular communication (GJIC) has been implicated in regulating crucial biological events in many stem cells, including proliferation, differentiation and apoptosis. Understanding and modulating GJIC in stem cells could potentially lead to the development of novel methods for expanding stem cells *in vitro* and directing their differentiation into functional mature cells. In this review we have summarized current knowledge on the identified roles of gap junctions and of reactive oxygen species (ROS) in stem cells, and speculate on how these may be exploited to develop the therapeutic potential of stem cells.

2. Gap junctions and gap junction intercellular communication

The classical background on gap junctions and GJIC has been covered extensively in a number of excellent recent reviews and therefore will only be briefly discussed here (Martin and Evans, 2004). Gap junctions are intercellular junctions found to be either in an opened or a closed conformation. They are the only intercellular junctions that allow direct transfer

of signalling molecules and metabolites to adjacent cells (Alexander and Goldberg, 2003; Kumar and Gilula, 1996). Gap junctions are hydrophilic channels consisting of two connexons, hemichannels localized in the membrane of adjacent cells, each of them consisting of six connexins (Cx) (Kumar and Gilula, 1996; Sosinsky and Nicholson, 2005) which can be assembled from either a single type or multiple types of connexins. Importantly, the connexin constitution of the gap junctions will define the pore size of the junction, hence allowing different permeability for the transfer of molecules (Saez et al., 2003). It is however generally accepted that only molecules less than 1-1.5kDa diffuse through gap junctions (De Maio et al., 2002; Evans et al., 2006). It is now also suggested that unpaired connexon hemichannels can mediate intercellular communication without forming gap junctions (Ebihara 2003; Goodenough and Paul 2003; Evans, De Vuyst et al. 2006).

3. Gap junctions in undifferentiated stem cells

3.1 Mouse embryonic stem cells

Mouse embryonic stem cells (mESC) display functional GJIC, express mRNA transcripts of various connexins: Cx26, Cx30.3, Cx31, Cx32 and Cx37, Cx43 and Cx45 but only Cx31, Cx43 and Cx45 proteins, suggesting a translational regulation of connexins in mESC (Nishi et al., 1991; Oyamada et al., 1996; Worsdorfer et al., 2008). Studies so far seem to indicate that Cx45 does not play a fundamental role in mESC regulation of pluripotency and differentiation and suggest a role of Cx43, although still unclear, in these processes. Indeed, Cx45-null mESC are able to differentiate into cells of the three germ layers following embryoid body formation (Egashira et al., 2004). In contrast, some studies showed that Cx43-knock down mESC display decreased cell proliferation, down-regulation of several stem cell markers and up-regulation of differentiation markers and inability to form embryoid bodies (Todorova et al., 2008; Worsdorfer et al., 2008). However, other data suggest the opposite where the down-regulation of Cx43 resulted in Src phosphorylation and increase in cell proliferation (Kim et al., 2010). In the same study, the adenosine analogue 5'-N-ethylcarboxamide (NECA) was shown to stimulate Cx43 phosphorylation and to inhibit GJIC in mESC, through the activation of PI3K/Akt, PKC, MAPK and NFκB signalling pathways. The closure of GJIC would subsequently lead to Src-induced cell proliferation and migration (Kim et al., 2010). Lastly, Cx43-null mESC show not only a defective differentiation to oligodendrocytes but also an increase in differentiation to astrocytes (Parekkadan et al., 2008). Hence there are conflicting findings on the role of GJIC in mESC, with some data suggesting that Cx43 and open gap junctions are necessary for mESC proliferation, for survival and for maintaining them in the pluripotent state, while other data suggest that the closure of gap junctions is in fact a signal for proliferation and migration. Clearly further work is required to properly define the roles GJIC in mESC under different conditions.

3.2 Human embryonic stem cells

Human embryonic stem cells (hESC) are also coupled through functional gap junctions (Bhattacharya et al., 2004; Carpenter et al., 2004; Wolvetang et al., 2007; Wong et al., 2006; Wong et al., 2004) (Figure 1) and express almost all transcripts of human connexin isoforms, except Cx40.1 and Cx50 (Huettnner et al., 2006). Functional GJIC seems to be a common characteristic of hESC maintained in different culture conditions (Wong et al., 2006; Wong et al., 2004). So far, only a few factors have been shown to inhibit GJIC in hESC such as bone morphogenetic protein (BMP)-4 (Wong et al., 2006), endothelin (ET) 1 and ET-2 (Wong et al.,




Fig. 1. GJIC in hESC. (A, D, G, J): Light and fluorescence micrographs with Lucifer yellow (B, E, H, K) and rhodamine-dextran (C, F, I, L) in HES-3 cells. Rhodamine-dextran was used as a negative control, showing no dye transfer across to the neighboring cell. Cells were incubated in the presence (A-C) or absence (D-F) of $\text{Ca}^{2+}\text{Mg}^{2+}$ or in the presence of phorbol 12-myristate 13-acetate (G-I) or U0126 (J-L). Scale bars = 100 μm . Reproduced, with permission, from (Wong et al., 2004).

2009). Similar to mESC, studies suggest that gap junctions play an important role in the regulation of hESC pluripotency and differentiation. Indeed, long-term chemical inhibition of GJIC in hESC can increase apoptosis (Wong et al., 2006) and in pro-differentiation conditions (such as the absence of a feeder cell layer, and non-conditioned medium), hESC do not communicate through gap junctions (Wong et al., 2006). Furthermore, the pro-differentiation factor BMP-4 (Pera et al., 2004; Xu et al., 2002) inhibits GJIC in hESC, an effect that can be prevented by the addition of the BMP antagonist noggin (Wong et al., 2006). The inhibitory effect of ET-1 is short-lasting and is not associated with changes in colony size, morphology or stem cell marker expression (Wong et al., 2009). These studies suggest that open gap junctions are required for the maintenance of hESC pluripotency and that closure of gap junctions is associated with differentiation or cell death. Further work is however needed to understand the precise implications of GJIC for maintaining hESC pluripotency.

3.3 Somatic stem cells

The inhibition of GJIC in somatic stem cells prevents regeneration of the planarian flatworm, suggesting a conserved role of gap junctions in regulating stem cell fate (Oviedo and Levin, 2007). However, not all somatic stem cells communicate through functional gap junctions. Some somatic stem and putative progenitor/stem somatic cells lack connexin expression and/or functional GJIC : keratinocyte stem cells (Matic et al., 2002), corneal epithelial stem cells (Matic et al., 1997), pancreatic ductal epithelial stem cells (Tai et al., 2003), neural-glial stem cells (Dowling-Warriner and Trosko, 2000), bovine mammary gland progenitor cells (Holland et al., 2003), human breast epithelial stem cells (Kao et al., 1995) and human kidney epithelial stem cells (Chang et al., 1987). Other somatic stem cells, in particular hematopoietic stem cells (HSC), mesenchymal stem cells (MSC) and neural stem/progenitor cells (NS/PC) express connexins and/or possess functional gap junctions that seem to play a role in regulating their homeostasis, pluripotency and/or differentiation. Current knowledge is however sparse. It has been proposed that Cx32, Cx43 and GJIC play a role in HSC maintenance and differentiation (Hirabayashi et al., 2007b; Ploemacher et al., 2000; Rosendaal et al., 1997). Indeed, Cx32 knockout mice exhibit more undifferentiated HSC and fewer progenitor cells, suggesting a role of Cx32 in maturation of HSC to progenitor cells (Hirabayashi et al., 2007a; Hirabayashi et al., 2007b). Moreover, Cx43-deficient mice demonstrate defects in blood cell formation (Montecino-Rodriguez et al., 2000). Cx43 mRNA is not expressed in undifferentiated and quiescent HSC (Montecino-Rodriguez et al., 2000) but is up-regulated in adult mouse bone marrow upon stem cell division (Rosendaal et al., 1994). These data thus suggest an important role of Cx43 in hematopoiesis, but the precise molecular mechanisms remain to be elucidated. Human MSC can communicate through GJIC and express Cx40, Cx43 and Cx45 (Lin et al., 2007; Valiunas et al., 2004). Moreover, human MSC have been demonstrated to form Cx43-mediated GJIC with umbilical vein endothelial cells, which is of importance for their osteogenic differentiation (Villars et al., 2002). Lastly, rat brain derived NS/PC express Cx43 and Cx45 and communicate through GJIC, which is essential for their survival and proliferation (Cai et al., 2004). In these cells, Cx32 and Cx43 are upregulated during differentiation (Yang et al., 2005). Similar data were observed in mouse fetal NS/PC, where the closure of gap junctions decreases cell proliferation and reduces cell survival (Cheng et al., 2004; Duval et al., 2002). Interestingly the overexpression of Cx43 stimulates proliferation of these cells (Cheng et al., 2004), findings somewhat different to the observations in rat-derived NS/PC. Furthermore, in

mouse embryonic NS/PC, Cx43 and GJIC allow interkinetic nuclear migration, an apical basal movement of the nucleus observed during cell cycle and necessary for corticogenesis (Liu et al., 2010). In addition, NS/PC from other species have also been demonstrated to express Cx43 and communicate via GJIC (Wen et al., 2008) (Russo et al., 2008), suggesting a conserved and critical role of GJIC in NS/PC self-renewal and differentiation. Finally, it is notable that gap junctions appear to be important for establishing functional interactions between grafted NS/PC and host (Jaderstad et al., 2010).

4. What goes through gap junctions and how can this modify stem cell fate?

GJIC refers to the passive diffusion of intracellular molecules through gap junctions to a neighboring cell (Kumar and Gilula, 1996). Numerous cytoplasmic molecules can diffuse through gap junction channels, including small ions (Na^+ , K^+ , Ca^{2+} , H^+ , Cl^-), second messengers (cyclic nucleotides, inositol triphosphate), amino acids (glycine, glutamate), cellular metabolites (glucose, glutathione, adenosine, AMP, ADP, ATP), short interfering RNA (siRNA) and peptides involved in cross-presentation of major histocompatibility complex class I molecules (Alexander and Goldberg, 2003; Krysko et al., 2005; Neijssen et al., 2005; Valiunas et al., 1997). In adult cells and some tissue systems, GJIC has long been known as crucial for certain cellular functions, such as electrical synchronization, intercellular buffering of cytoplasmic ions, cell metabolism, control of cell migration and cell fate including carcinogenesis (De Maio et al., 2002; Krysko et al., 2005; Mesnil et al., 2005; Parekkadan et al., 2008; Todorova et al., 2008; Vine and Bertram, 2002). However, few studies have addressed the importance of gap junctions for promulgating intercellular Ca^{2+} waves in stem or progenitor cells. Early studies suggested that IP_3 can induce intracellular Ca^{2+} release from endoplasmic-reticulum, and both Ca^{2+} and IP_3 can permeate gap junction channels to the neighbouring cells (Boitano et al., 1992; Saez et al., 1989). Such diffusion of IP_3 and Ca^{2+} through gap junctions effectively forms a positive feedback loop allowing intercellular communication between distant cells. Alternatively, other studies have demonstrated that such intercellular Ca^{2+} waves can also be maintained by the paracrine messenger ATP (Guthrie et al., 1999). It was demonstrated that IP_3 can trigger ATP release to the extracellular space through connexon hemichannels, and act on G-protein coupled receptors on neighbouring cells, leading to phospholipase C activation, IP_3 production and subsequent Ca^{2+} release in the neighbouring cells (Braet et al., 2003; Ebihara et al., 2003; Guthrie et al., 1999). Gap junction-mediated transmission of Ca^{2+} waves has been inferred in some progenitor cells. In NS/PC cells, transmission via Ca^{2+} waves appears to control their proliferation in the ventricular zone (Weissman et al., 2004) as well as in retinal neural progenitor cells (Pearson et al., 2005). Moreover, transient increases in intracellular Ca^{2+} can also stimulate differentiation and neurite outgrowth of different NS/PC cells, but whether such Ca^{2+} signalling occurs through gap junction-mediated waves remains to be determined (Carey and Matsumoto, 1999; Gomez and Spitzer, 1999; Gu and Spitzer, 1995). Recent transplantantion studies by Jaderstad et al. (2010) showed that establishment of GJIC that allows synchronized Ca^{2+} waves is important for grafted NS/PC to integrate functionally to the host neural circuitry. Such GJIC between grafted NS/PC and host cells thus provided a neuroprotective effect in mouse models of neurodegeneration (Jaderstad et al., 2010). Interestingly, diffusion of Ca^{2+} via gap junctions has also been suggested to modulate differentiation of other somatic stem cells. Previous studies by Muller-Borer et al. (2004) suggested that synchronized Ca^{2+} signalling via GJIC co-cultured with neonatal

cardiomyocytes induces rat liver stem cells to express cardiac transcription factors and acquire a phenotype resembling cardiomyocytes (Anderson et al., 2007; Muller-Borer et al., 2004). Similarly, we have recently shown gap junctions established between human adipose-derived MSC and neonatal rat cardiomyocytes in co-culture, which induces the expression of cardiac genes and spontaneous cardiomyocyte-like contractions in the human cells (Choi et al., 2010a) (Figure 2). All this points to a growing appreciation of the role of gap-junction-mediated Ca^{2+} waves in modulating gene expression and promoting differentiation of stem cells. Other metabolites that can diffuse through gap junction include cyclicAMP and glutamate, which were previously shown to act as signals for cell death (Amsterdam et al., 1996; Ozog et al., 2002). Although it remains to be proven, it is possible that cyclicAMP and glutamate might play a role in propagation of death signals in stem cells that possess GJIC, such as hESC and NS/PC. Finally, interesting candidates amongst many that might yet be exposed to diffuse through gap junctions, are small RNAs. Previous studies provided evidence that exogenous siRNA can diffuse through gap junction to the neighbouring cells




Fig. 2. Functional gap junctions formed between human adipose-derived MSC and rat neonatal cardiomyocytes co-cultured for 24 hours. The human stem cells are labeled red with DiI, which is unable to transfer between cells, and the rat cardiomyocytes were stained by calcein-AM which can transfer from the cytosol of one cell to the adjacent ones through gap junctions. After co-culture, double labeled cells (human cells – cardiomyocyte-like-differentiated from MSC) can be seen as indicated by the arrows (Choi, Dusting, Dilley et al – unpublished, similar to Choi et al 2010a)

(Valiunas et al., 2005; Wolvetang et al., 2007) to silence genes therein. Thus, it is possible that this mechanism may also apply to endogenous microRNAs that provide post-transcriptional regulation of a diverse array of genes. Given the emerging role of microRNA in regulating various physiological processes in hESC and other somatic stem cells (Liu and Zhao, 2009; Mallanna and Rizzino, 2010; Navarro and Lieberman, 2010), future work is likely to explore gap junction-mediated transfer of microRNA in modulating the fate of adjacent stem cells.

5. Oxidant signalling and NADPH oxidase in proliferation and survival of vascular cells

Over-production of reactive oxygen species (ROS) and diminished antioxidant systems (e.g. superoxide dismutase, catalase, glutathione peroxidase and glutathione) may lead to oxidative stress, and this is known to contribute to the pathogenesis of several diseases. These include ischemic-reperfusion injury (e.g. heart attack and stroke), atherosclerosis, hypertension, ischemic heart disease, cancers and neurodegeneration. However, given that all types of cells generate low but detectable amounts of ROS under different circumstances, it is likely that ROS serve as important mediators under physiological conditions. In fact, the production of ROS is tightly regulated by antioxidant systems, which maintain redox homeostasis within the cellular environment. As a consequence, ROS have distinct functional effects, which are dependent on a number of factors such as the type of cell within which ROS are generated, and the type and ultimate concentration of ROS at sub-cellular sites where they may modulate enzyme activity and influence gene expression. One of the most important ROS in the vasculature is superoxide anion, formed enzymatically and non-enzymatically, by the univalent reduction of oxygen. The best characterized source of superoxide is the mitochondrial electron transport chain, but many other intracellular enzymes such as xanthine oxidase (XO), cyclooxygenase (COX), nitric oxide synthase (NOS), cytochrome P₄₅₀ oxidase and NADPH oxidase are capable of producing this radical. All these enzymes, save NADPH oxidase, have important cellular functions apart from superoxide generation, whereas the NADPH oxidase enzyme complex is the only known enzyme dedicated to production of ROS, using intracellular NADPH as the “substrate” and electron donor. Since the mid 1990’s it has become evident that many cells produce superoxide constitutively by an enzyme with all the characteristics of the NADPH oxidase previously shown to be present in dedicated phagocytic or inflammatory cells. Although constitutively active, NADPH oxidases in blood vessels can be further activated by stimuli such as angiotensin-II (Ang-II), tumour necrosis factor- α (TNF α), TGF β , thrombin, platelet-derived growth factor (PDGF) and by specific ROS themselves (Barry-Lane et al., 2001; Lassegue et al., 2001; Li and Shah, 2003; Moe et al., 2006; Patterson et al., 1999; Suh et al., 1999). The NADPH oxidase is comprised of a membrane-bound heterodimeric unit called flavocytochrome *b558*, composed of small subunit p22phox and gp91phox (aka Nox2). The catalytic moiety of gp91phox (Nox2) contains flavin-adenine dinucleotide (FAD) binding site, two heme components and one NADPH binding site. In the presence of stimuli such as phorbol ester, bacterial lipopolysaccharides or formyl-methionyl-leucyl-phenylalanine (fMLP), protein kinase C (PKC) causes phosphorylation of p47phox and initiates the translocation of p47phox, and its associated proteins p67phox, p40phox and small G-protein Rac1 to the membrane to bind to the cytochrome *b558* complex. The fully assembled complex allows NADPH to bind to gp91phox on the cytoplasmic side of the membrane to initiate a series of electron transfers starting from NADPH to FAD then to

heme and finally to oxygen to produce two molecules of superoxide anion radical. To date five isoforms of the catalytic subunit Nox have been identified (Nox1 to Nox5). In addition two homologs of the associated intracellular proteins p47phox and p67phox known as NoxA1 (NADPH oxidase activator1) and NoxO1 (NADPH oxidase organizer), respectively, have been identified (Babior et al., 1973; Lassegue and Clempus, 2003; Li and Shah, 2004). We and others have shown that NADPH oxidase-derived ROS have important functions in survival and proliferation of vascular cells (Ago et al., 2004; Chen et al., 2008; Peshavariya et al., 2009; Petry et al., 2006). Petry et al. showed that suppression of either Nox2 or Nox4 reduce endothelial cell proliferation in vitro, whereas over-expression of these isoforms increased cell proliferation (Petry et al., 2006). Similarly, we have shown that suppression of Nox4 only reduces proliferation, whereas suppression of Nox2 increases apoptosis and therefore also effectively promotes proliferation of endothelial cells (Peshavariya et al., 2009). Vascular smooth muscle cells (VSMC) from different sources express highly both Nox4 and Nox1, but do not express Nox2 (Chan et al., 2009). It is well documented that several growth factors increase Nox1 expression or activity or both, and they also enhance proliferation of VSMC (Lassegue et al., 2001; Suh et al., 1999). However the role of Nox4 in VSMC proliferation is complicated. For instance, it has been shown that transforming growth factor-beta (TGF β) - induced Nox4 is important in pulmonary smooth muscle cell proliferation (Sturrock et al., 2006). In contrast, the expression of Nox4 increased under quiescent (serum-deprived) conditions and this leads to aortic VSMC differentiation rather than proliferation under these conditions (Clempus et al., 2007). Despite such inconsistencies in the literature regarding the role of Nox4-derived ROS in proliferation versus differentiation, almost all studies indicate that Nox4 is involved in migration of VSMC and endothelial cells (Datla et al., 2007; Lyle et al., 2009; Sturrock et al., 2006). Taken together these findings suggest that NADPH oxidase and its isoforms have distinct roles in vascular cell survival and proliferation, and the disparity of functions may be due to the different ligand receptor couplings and sub-cellular localization of the NADPH oxidase complexes involved.

6. Intracellular kinase pathways induced by oxidant signalling promote cell survival or proliferation.

There is compelling evidence that low levels of oxidants activate several cell signalling pathways and regulate cell survival and proliferation, whereas high levels of oxidants stimulate stress-activated signalling pathways leading to cell death. For example, nanomolar to sub-micromolar concentrations of hydrogen peroxide (H₂O₂) stimulate the proliferation of several cell types, but higher concentrations of H₂O₂ (>100 μ M) leads to cell death (Giorgio et al., 2007; Kim et al., 2009; Stone and Yang, 2006). However, the effect of H₂O₂ is cell type dependent: H₂O₂ (100 μ M) increased VSMC proliferation whereas the same concentration inhibits the proliferation of endothelial and fibroblast cells (Rao and Berk, 1992). It is important to note, however, that provision of exogenous H₂O₂ may produce very different effects from stimulation of endogenous H₂O₂ release at particular subcellular sites (Forman, 2007). Ligand-receptor interactions also result in the generation of H₂O₂ by mammalian cells, but these may have different effects on their downstream signalling pathways and thus exert distinct effects on cell survival, proliferation and differentiation (Chen et al., 2008; Datla et al., 2007; Suh et al., 1999; Wang et al., 2000). ROS have several targets such as transcription factors, phosphatases, and enzymes of the receptor kinase family. It has

become evident that ROS regulate activity of MAP kinases (MAPKs), a family of serine/threonine kinases, including extracellular signal-regulated kinases 1 and 2 (ERK1/2), c-Jun N-terminal kinases (JNKs, also termed stress activated protein kinase; SAPKs) and p38MAPK (Figure 3) Several growth factors (Vascular endothelial growth factor, VEGF; epidermal growth factor, EGF; PDGF and thrombin) and cytokines (Ang-II and TNF- α) induce proliferation of endothelial and VSMC via ROS-mediated activation of MAPK family members (Chen et al., 2008; Datla et al., 2007; Lassegue et al., 2001; Li et al., 2005; Li and Shah, 2003; Park et al., 2009; Patterson et al., 1999; Suh et al., 1999; Ushio-Fukai et al., 1998; Ushio-Fukai et al., 1999; Ushio-Fukai et al., 2002). For example, VEGF, EGF, and TNF- α stimulate proliferation or angiogenesis of endothelial cells which is dependent upon ROS-induced phosphorylation of ERK (Chen et al., 2004; Chen et al., 2008; Datla et al., 2007; Li et al., 2005; Ushio-Fukai et al., 2002). The effect of TNF- α on endothelial cell proliferation is concentration dependent: lower concentrations of TNF- α induce angiogenesis whereas higher concentrations induce apoptosis of endothelial cells (Chen et al., 2004; Deshpande et al., 2000). Similarly, Ang-II and PDGF induced the phosphorylation of p38MAPK, JNK and ERK, but only the phosphorylation of p38MAPK and JNK are ROS sensitive and lead to VSMC growth (Lassegue et al., 2001). Furthermore, in human aortic VSMC 7-ketocholesterol induces ROS via Nox4 at the endoplasmic reticulum and activates pJNK, but this leads to cell death (Pedruzzi et al., 2004). In contrast to the MAPK, Akt/protein kinase B has been identified as an important component of a pro-survival signalling pathway. Addition of either exogenous H₂O₂ or receptor-mediated intracellular H₂O₂ leads to the phosphorylation of Akt (Esposito et al., 2003; Ushio-Fukai et al., 1999). Recently, it has been demonstrated that ROS-mediated activation of the PI3kinase/Akt pathway increased the production of




Fig. 3. NADPH oxidase and ROS signalling. Extracellular ligand-activated membrane receptors (including GPCRs) linked to NADPH oxidase produce ROS intracellularly (superoxide anion or H₂O₂ in the case of Nox4), which in turn, lead to the phosphorylation and activation of the MAP kinases indicated. The different kinases have different effects in

adult vascular smooth muscle and endothelial cells, as indicated, some of which are shared in stem cells (see text).

nitric oxide and again promoted survival of endothelial cells (Bodiga et al.; Dhanasekaran et al., 2009). Previously, we showed that the proliferative state of endothelial cells exhibits higher ROS production and phosphorylation of Akt compared to quiescent cells, and inhibition of either ROS production or the PI3kinase/Akt pathway reduces endothelial cell proliferation (Peshavariya et al., 2009). ROS-mediated activation of Phospho-Akt (pAkt) promoting cell survival has also been reported in other cell types, such as HeLa, NIH3T3 cells (Wang et al., 2000) and hepatocytes (Kim et al., 2008). Thus it emerges that activation of ROS-dependent signalling pathways are influenced by several factors such as ligand-receptor interaction, the type of cell, sub-cellular localisation of ROS producing enzymes and the antioxidant status of cells. Therefore, ROS mediated signalling pathways are fine-tuned to differential functions such as proliferation, survival or apoptosis in different cell types under different conditions.

7. Intracellular kinase pathways promoting survival of stem cells and oxidant signalling inducing differentiation

Given that ROS signalling clearly has important roles in the proliferation, survival and differentiation of several cell types, it should be considered whether or not this applies to stem cells. In hESC, the PI3k/Akt and ERK1/2 pathways are constitutively active in most culture media used for the maintenance of hESC (Lin et al., 2007). Moreover, the pro-maintenance factors bFGF, neutrophins, S1P and PDGF activate these kinase signalling pathways, suggesting an essential role of these pathways in hESC maintenance and pluripotency. Furthermore, inhibition of either pathway results in differentiation of hESC or cell death (Armstrong et al., 2006; McLean et al., 2007; Wong et al., 2007). Recently the link to ROS activation of these pathways has been demonstrated in hESC, where ROS stimulation leads to differentiation to mesodermal cells (Ji et al., 2010). Recently we have explored intracellular pathways promoting cell survival after hypoxic pre-conditioning of adipose-derived mesenchymal stem cells (MSC). In these cells it was clear that VEGFA was the major cytoprotective cytokine released during hypoxia, and again VEGF acted via Akt1 phosphorylation to protect these MSC from apoptosis during subsequent severe ischaemia, for the protective effect of the preconditioning was blocked by a VEGF antibody and the PI3 kinase inhibitor LY294002 (Stubbs et al., 2010). Interestingly this paracrine protective effect could be imparted to endothelial cells subjected to hypoxia, revealing an interesting way that adipose-derived MSC that we have utilised in tissue engineering of cardiac tissue, could promote the growth of any complex tissue that requires a vasculature (Chan et al., 2009; Choi et al., 2010b). Several other studies have suggested that ROS signalling can determine the fate of stem cells (Lee et al., 2009; Li et al., 2006; Li and Marban, 2010). For example an early study showed that mESC generate intracellular ROS, and addition of exogenous H₂O₂ promotes their differentiation to cardiomyocytes. This study suggested that PI3 kinase/Akt pathway is upstream of ROS, for inhibition of PI3 kinase reduced ROS formation and cardiac cell differentiation (Sauer et al., 2000; Sauer and Wartenberg, 2005). The enzymatic source of ROS and their downstream signalling pathways have been further explored in differentiation of ESC down the cardiac lineage. Interestingly, Nox4 emerged as the main source of ROS involved in cardiac differentiation, and it appears to regulate

phosphorylation of p38MAPK and the cardiac differentiation markers Nkx2.5 and myocyte enhancer factor 2C (MEF2C; (Li et al., 2006)). Mechanical strain induces cardiac differentiation of mESC and both ROS and their downstream signalling pathways were shown to be involved. Antioxidants N-(2-mercapto-propionyl-glycine (NMPG) and vitamin E suppress mechanical strain-induced ROS and reduces the critical downstream signalling pathways p38MAPK, ERK and JNK and also compromises cardiac differentiation and vasculogenesis. Thus it seems that mechanical strain activates these three members of the MAPK family via ROS and there is no specificity at this level (Schmelter et al., 2006). Intracellular ROS derived from mESC not only enhanced the differentiation to cardiomyocytes but also increased their proliferation, underlining the importance of ROS in cardiomyogenesis (Buggisch et al., 2007). Finally, ESC-derived ROS signalling is not limited to cardiac differentiation but is also involved in differentiation to VSMC and vascular stabilization. Xiao et al (2009) demonstrated that Nox4 over-expressing ESC showed enhanced differentiation to VSMC, involving transcription factors including serum response factor (SRF) and myocardin.

8. Conclusion

We are just beginning to understand the signalling that occurs between stem cells, both adult mesenchymal and embryonic, and soon studies will be focused on these mechanisms in induced pluripotent stem cells. There is emerging evidence on the importance of intercellular and intracellular signalling mechanisms that use gap junctions and oxidant signalling to regulate maintenance of stemness and proliferation, or alternatively trigger differentiation or apoptosis to grow new tissues. Defining these mechanisms will lead to greater efficiency in developing stem cell therapies for the clinic, and we amongst others are using this to build larger, more robust constructs from stem cells through tissue engineering (eg Chan et al 2009). The fruits of this signalling research will enhance approaches to regenerative medicine in many fields, and hopefully allow the great promise of stem cells to realise its full potential.

9. Acknowledgements

The authors' work and Fellowship (GJD) was supported by grants from the National Health and Medical Research Council of Australia.

RCB Wong is currently supported by the California Institute of Regenerative Medicine (RC1-00110-1).

10. References

- Ago, T., Kitazono, T., Ooboshi, H., Iyama, T., Han, Y. H., Takada, J., Wakisaka, M., Ibayashi, S., Utsumi, H., Iida, M. (2004). Nox4 as the major catalytic component of an endothelial NAD(P)H oxidase. *Circulation*. 109, 227-33.
- Alexander, D. B., Goldberg, G. S. (2003). Transfer of biologically important molecules between cells through gap junction channels. *Curr Med Chem*. 10, 2045-58.
- Amsterdam, A., Keren-Tal, I., Aharoni, D. (1996). Cross-talk between cAMP and p53-generated signals in induction of differentiation and apoptosis in steroidogenic granulosa cells. *Steroids*. 61, 252-6.

- Anderson, P. A., Muller-Borer, B. J., Esch, G. L., Coleman, W. B., Grisham, J. W., Malouf, N. N. (2007). Calcium signals induce liver stem cells to acquire a cardiac phenotype. *Cell Cycle*. 6, 1565-9.
- Armstrong, L., Hughes, O., Yung, S., Hyslop, L., Stewart, R., Wappler, I., Peters, H., Walter, T., Stojkovic, P., Evans, J., Stojkovic, M. & Lako, M. (2006). The role of PI3K/AKT, MAPK/ERK and NFkappabeta signalling in the maintenance of human embryonic stem cell pluripotency and viability highlighted by transcriptional profiling and functional analysis, *Hum Mol Genet* Vol.(11): 1894-913.
- Babior, B. M., Kipnes, R. S., Curnutte, J. T. (1973). Biological defense mechanisms. The production by leukocytes of superoxide, a potential bactericidal agent. *J Clin Invest*. 52, 741-4.
- Barry-Lane, P. A., Patterson, C., van der Merwe, M., Hu, Z., Holland, S. M., Yeh, E. T., Runge, M. S. (2001). p47phox is required for atherosclerotic lesion progression in ApoE(-/-) mice. *J Clin Invest*. 108, 1513-22.
- Bhattacharya, B., Miura, T., Brandenberger, R., Mejido, J., Luo, Y., Yang, A. X., Joshi, B. H., Ginis, I., Thies, R. S., Amit, M., Lyons, I., Condie, B. G., Itskovitz-Eldor, J., Rao, M. S., Puri, R. K. (2004). Gene expression in human embryonic stem cell lines: unique molecular signature. *Blood*. 103, 2956-64.
- Bodiga, S., Gruenloh, S. K., Gao, Y., Manthathi, V. L., Dubasi, N., Falck, J. R., Medhora, M., Jacobs, E. R. 20-HETE-induced nitric oxide production in pulmonary artery endothelial cells is mediated by NADPH oxidase, H₂O₂, and PI3-kinase/Akt. *Am J Physiol Lung Cell Mol Physiol*. 298, L564-74.
- Boitano, S., Dirksen, E. R., Sanderson, M. J. (1992). Intercellular propagation of calcium waves mediated by inositol trisphosphate. *Science*. 258, 292-5.
- Braet, K., Vandamme, W., Martin, P. E., Evans, W. H., Leybaert, L. (2003). Photoliberating inositol-1,4,5-trisphosphate triggers ATP release that is blocked by the connexin mimetic peptide gap 26. *Cell Calcium*. 33, 37-48.
- Buggisch, M., Ateghang, B., Ruhe, C., Strobel, C., Lange, S., Wartenberg, M., Sauer, H. (2007). Stimulation of ES-cell-derived cardiomyogenesis and neonatal cardiac cell proliferation by reactive oxygen species and NADPH oxidase. *J Cell Sci*. 120, 885-94.
- Cai, J., Cheng, A., Luo, Y., Lu, C., Mattson, M. P., Rao, M. S., Furukawa, K. (2004). Membrane properties of rat embryonic multipotent neural stem cells. *J Neurochem*. 88, 212-26.
- Carey, M. B., Matsumoto, S. G. (1999). Spontaneous calcium transients are required for neuronal differentiation of murine neural crest. *Dev Biol*. 215, 298-313.
- Carpenter, M. K., Rosler, E. S., Fisk, G. J., Brandenberger, R., Ares, X., Miura, T., Lucero, M., Rao, M. S. (2004). Properties of four human embryonic stem cell lines maintained in a feeder-free culture system. *Dev Dyn*. 229, 243-58.
- Chan, E. C., Jiang, F., Peshavariya, H. M., Dusting, G. J. (2009). Regulation of cell proliferation by NADPH oxidase-mediated signaling: potential roles in tissue repair, regenerative medicine and tissue engineering. (Review). *Pharmacology Therapeutics*. 122, 97-108.

- Chang, C. C., Trosko, J. E., el-Fouly, M. H., Gibson-D'Ambrosio, R. E., D'Ambrosio, S. M. (1987). Contact insensitivity of a subpopulation of normal human fetal kidney epithelial cells and of human carcinoma cell lines. *Cancer Res.* 47, 1634-45.
- Chen, J. X., Chen, Y., DeBusk, L., Lin, W., Lin, P. C. (2004). Dual functional roles of Tie-2/angiopoietin in TNF-alpha-mediated angiogenesis. *Am J Physiol Heart Circ Physiol.* 287, H187-95.
- Chen, K., Kirber, M. T., Xiao, H., Yang, Y., Keaney, J. F., Jr. (2008). Regulation of ROS signal transduction by NADPH oxidase 4 localization. *J Cell Biol.* 181, 1129-39.
- Cheng, A., Tang, H., Cai, J., Zhu, M., Zhang, X., Rao, M., Mattson, M. P. (2004). Gap junctional communication is required to maintain mouse cortical neural progenitor cells in a proliferative state. *Dev Biol.* 272, 203-16.
- Choi, Y. S., Dusting, G. J., Stubbs, S., Arunothayaraj, S., Han, X. L., Collas, P., Morrison, W. A., Dilley, R. J. (2010a). Differentiation of human adipose-derived stem cells into beating cardiomyocytes. *J Cell Mol Med.* 14, 878-889.
- Choi, Y. S., Matsuda, K., Dusting, G. J., Morrison, W. A., Dilley, R. J. (2010b). Engineering cardiac tissue in vivo from human adipose-derived stem cells. *Biomaterials.* 31, 2236-42.
- Clempus, R. E., Sorescu, D., Dikalova, A. E., Pounkova, L., Jo, P., Sorescu, G. P., Schmidt, H. H., Lassegue, B., Griendling, K. K. (2007). Nox4 is required for maintenance of the differentiated vascular smooth muscle cell phenotype. *Arterioscler Thromb Vasc Biol.* 27, 42-8.
- Datla, S. R., Peshavariya, H., Dusting, G. J., Mahadev, K., Goldstein, B. J., Jiang, F. (2007). Important role of Nox4 type NADPH oxidase in angiogenic responses in human microvascular endothelial cells in vitro. *Arterioscler Thromb Vasc Biol.* 27, 2319-24.
- De Maio, A., Vega, V., Contreras, J. (2002). Gap junctions, homeostasis, and injury. *J Cell Physiol.* 191, 269-282.
- Deshpande, S. S., Angkeow, P., Huang, J., Ozaki, M., Irani, K. (2000). Rac1 inhibits TNF-alpha-induced endothelial cell apoptosis: dual regulation by reactive oxygen species. *Faseb J.* 14, 1705-14.
- Dhanasekaran, A., Bodiga, S., Gruenloh, S., Gao, Y., Dunn, L., Falck, J. R., Buonaccorsi, J. N., Medhora, M., Jacobs, E. R. (2009). 20-HETE increases survival and decreases apoptosis in pulmonary arteries and pulmonary artery endothelial cells. *Am J Physiol Heart Circ Physiol.* 296, H777-86.
- Dowling-Warriner, C. V., Trosko, J. E. (2000). Induction of gap junctional intercellular communication, connexin43 expression, and subsequent differentiation in human fetal neuronal cells by stimulation of the cyclic AMP pathway. *Neuroscience.* 95, 859-68.
- Duval, N., Gomes, D., Calaora, V., Calabrese, A., Meda, P., Bruzzone, R. (2002). Cell coupling and Cx43 expression in embryonic mouse neural progenitor cells. *J Cell Sci.* 115, 3241-51.
- Ebihara, L., Liu, X., Pal, J. D. (2003). Effect of external magnesium and calcium on human connexin46 hemichannels. *Biophys J.* 84, 277-86.

- Egashira, K., Nishii, K., Nakamura, K., Kumai, M., Morimoto, S., Shibata, Y. (2004). Conduction abnormality in gap junction protein connexin45-deficient embryonic stem cell-derived cardiac myocytes. *Anat Rec A Discov Mol Cell Evol Biol.* 280, 973-9.
- Esposito, F., Chirico, G., Montesano Gesualdi, N., Posadas, I., Ammendola, R., Russo, T., Cirino, G., Cimino, F. (2003). Protein kinase B activation by reactive oxygen species is independent of tyrosine kinase receptor phosphorylation and requires SRC activity. *J Biol Chem.* 278, 20828-34.
- Evans, W. H., De Vuyst, E., Leybaert, L. (2006). The gap junction cellular internet: connexin hemichannels enter the signalling limelight. *Biochem J.* 397, 1-14.
- Forman, H. J. (2007). Use and abuse of exogenous H₂O₂ in studies of signal transduction. *Free Radic Biol Med.* 42, 926-32.
- Giorgio, M., Trinei, M., Migliaccio, E., Pelicci, P. G. (2007). Hydrogen peroxide: a metabolic by-product or a common mediator of ageing signals? *Nat Rev Mol Cell Biol.* 8, 722-8.
- Gomez, T. M., Spitzer, N. C. (1999). In vivo regulation of axon extension and pathfinding by growth-cone calcium transients. *Nature.* 397, 350-5.
- Gu, X., Spitzer, N. C. (1995). Distinct aspects of neuronal differentiation encoded by frequency of spontaneous Ca²⁺ transients. *Nature.* 375, 784-7.
- Guthrie, P. B., Knappenberger, J., Segal, M., Bennett, M. V., Charles, A. C., Kater, S. B. (1999). ATP released from astrocytes mediates glial calcium waves. *J Neurosci.* 19, 520-8.
- Hirabayashi, Y., Yoon, B. I., Tsuboi, I., Huo, Y., Kodama, Y., Kanno, J., Ott, T., Trosko, J. E., Inoue, T. (2007a). Membrane channel connexin 32 maintains Lin(-)/c-kit(+) hematopoietic progenitor cell compartment: analysis of the cell cycle. *J Membr Biol.* 217, 105-13.
- Hirabayashi, Y., Yoon, B. I., Tsuboi, I., Huo, Y., Kodama, Y., Kanno, J., Ott, T., Trosko, J. E., Inoue, T. (2007b). Protective role of connexin 32 in steady-state hematopoiesis, regeneration state, and leukemogenesis. *Exp Biol Med (Maywood).* 232, 700-12.
- Holland, M. S., Tai, M. H., Trosko, J. E., Griffin, L. D., Stasko, J. A., Cheville, N. C., Holland, R. E. (2003). Isolation and differentiation of bovine mammary gland progenitor cell populations. *Am J Vet Res.* 64, 396-403.
- Huettner, J. E., Lu, A., Qu, Y., Wu, Y., Kim, M., McDonald, J. W. (2006). Gap junctions and connexon hemichannels in human embryonic stem cells. *Stem Cells.* 24, 1654-67.
- Jaderstad, J., Jaderstad, L. M., Li, J., Chintawar, S., Salto, C., Pandolfo, M., Ourednik, V., Teng, Y. D., Sidman, R. L., Arenas, E., Snyder, E. Y., Herlenius, E. (2010). Communication via gap junctions underlies early functional and beneficial interactions between grafted neural stem cells and the host. *Proc Natl Acad Sci U S A.* 107, 5184-9.
- Ji, A. R., Ku, S. Y., Cho, M. S., Kim, Y. Y., Kim, Y. J., Oh, S. K., Kim, S. H., Moon, S. Y., Choi, Y. M. (2010). Reactive oxygen species enhance differentiation of human embryonic stem cells into mesendodermal lineage. *Exp Mol Med.* 42, 175-86.
- Kao, C. Y., Nomata, K., Oakley, C. S., Welsch, C. W., Chang, C. C. (1995). Two types of normal human breast epithelial cells derived from reduction mammoplasty:

- phenotypic characterization and response to SV40 transfection. *Carcinogenesis*. 16, 531-8.
- Kim, H. S., Loughran, P. A., Rao, J., Billiar, T. R., Zuckerbraun, B. S. (2008). Carbon monoxide activates NF-kappaB via ROS generation and Akt pathways to protect against cell death of hepatocytes. *Am J Physiol Gastrointest Liver Physiol*. 295, G146-G152.
- Kim, J. H., Choi, W., Lee, J. H., Jeon, S. J., Choi, Y. H., Kim, B. W., Chang, H. I., Nam, S. W. (2009). Astaxanthin inhibits H₂O₂-mediated apoptotic cell death in mouse neural progenitor cells via modulation of P38 and MEK signaling pathways. *J Microbiol Biotechnol*. 19, 1355-63.
- Kim, M. O., Lee, Y. J., Han, H. J. (2010). Involvement of Cx43 phosphorylation in 5'-N-ethylcarboxamide-induced migration and proliferation of mouse embryonic stem cells. *J Cell Physiol*. 224, 187-94.
- Krysko, D. V., Leybaert, L., Vandenabeele, P., D'Herde, K. (2005). Gap junctions and the propagation of cell survival and cell death signals. *Apoptosis*. 10, 459-69.
- Kumar, N. M., Gilula, N. B. (1996). The gap junction communication channel. *Cell*. 84, 381-8.
- Lassegue, B., Clempus, R. E. (2003). Vascular NAD(P)H oxidases: specific features, expression, and regulation. *Am J Physiol Regul Integr Comp Physiol*. 285, R277-97.
- Lassegue, B., Sorescu, D., Szocs, K., Yin, Q., Akers, M., Zhang, Y., Grant, S. L., Lambeth, J. D., Griendling, K. K. (2001). Novel gp91(phox) homologues in vascular smooth muscle cells : nox1 mediates angiotensin II-induced superoxide formation and redox-sensitive signaling pathways. *Circ Res*. 88, 888-94.
- Lee, S. H., Na, S. I., Heo, J. S., Kim, M. H., Kim, Y. H., Lee, M. Y., Kim, S. H., Lee, Y. J., Han, H. J. (2009). Arachidonic acid release by H₂O₂ mediated proliferation of mouse embryonic stem cells: involvement of Ca²⁺/PKC and MAPKs-induced EGFR transactivation. *J Cell Biochem*. 106, 787-97.
- Li, J., Stouffs, M., Serrander, L., Banfi, B., Bettiol, E., Charnay, Y., Steger, K., Krause, K. H., Jaconi, M. E. (2006). The NADPH oxidase NOX4 drives cardiac differentiation: Role in regulating cardiac transcription factors and MAP kinase activation. *Mol Biol Cell*. 17, 3978-88.
- Li, J. M., Fan, L. M., Christie, M. R., Shah, A. M. (2005). Acute tumor necrosis factor alpha signaling via NADPH oxidase in microvascular endothelial cells: role of p47phox phosphorylation and binding to TRAF4. *Mol Cell Biol*. 25, 2320-30.
- Li, J. M., Shah, A. M. (2003). Mechanism of endothelial cell NADPH oxidase activation by angiotensin II. Role of the p47phox subunit. *J Biol Chem*. 278, 12094-100.
- Li, J. M., Shah, A. M. (2004). Endothelial cell superoxide generation: regulation and relevance for cardiovascular pathophysiology. *Am J Physiol Regul Integr Comp Physiol*. 287, R1014-30.
- Li, T. S., Marban, E. (2010). Physiological levels of reactive oxygen species are required to maintain genomic stability in stem cells. *Stem Cells*. 28, 1178-85.
- Lin, T. M., Chang, H. W., Wang, K. H., Kao, A. P., Chang, C. C., Wen, C. H., Lai, C. S., Lin, S. D. (2007). Isolation and identification of mesenchymal stem cells from human lipoma tissue. *Biochem Biophys Res Commun*. 361, 883-9.

- Liu, C., Zhao, X. (2009). MicroRNAs in adult and embryonic neurogenesis. *Neuromolecular Med.* 11, 141-52.
- Liu, X., Hashimoto-Torii, K., Torii, M., Ding, C., Rakic, P. (2010). Gap junctions/hemichannels modulate interkinetic nuclear migration in the forebrain precursors. *J Neurosci.* 30, 4197-209.
- Lyle, A. N., Deshpande, N. N., Taniyama, Y., Seidel-Rogol, B., Pounkova, L., Du, P., Papaharalambus, C., Lassegue, B., Griendling, K. K. (2009). Poldip2, a novel regulator of Nox4 and cytoskeletal integrity in vascular smooth muscle cells. *Circ Res.* 105, 249-59.
- Mallanna, S. K., Rizzino, A. (2010). Emerging roles of microRNAs in the control of embryonic stem cells and the generation of induced pluripotent stem cells. *Dev Biol.* 344, 16-25.
- Martin, P. E., Evans, W. H. (2004). Incorporation of connexins into plasma membranes and gap junctions. *Cardiovasc Res.* 62, 378-87.
- Matic, M., Evans, W. H., Brink, P. R., Simon, M. (2002). Epidermal stem cells do not communicate through gap junctions. *J Invest Dermatol.* 118, 110-6.
- Matic, M., Petrov, I. N., Chen, S., Wang, C., Dimitrijevic, S. D., Wolosin, J. M. (1997). Stem cells of the corneal epithelium lack connexins and metabolite transfer capacity. *Differentiation.* 61, 251-60.
- McLean, A. B., D'Amour, K. A., Jones, K. L., Krishnamoorthy, M., Kulik, M. J., Reynolds, D. M., Sheppard, A. M., Liu, H., Xu, Y., Baetge, E. E. & Dalton, S. (2007). Activin efficiently specifies definitive endoderm from human embryonic stem cells only when phosphatidylinositol 3-kinase signaling is suppressed, *Stem Cells* Vol.(1): 29-38.
- Mesnil, M., Crespín, S., Avanzo, J. L., Zaidan-Dagli, M. L. (2005). Defective gap junctional intercellular communication in the carcinogenic process. *Biochim Biophys Acta.* 1719, 125-45.
- Moe, K. T., Aulia, S., Jiang, F., Chua, Y. L., Koh, T. H., Wong, M. C., Dusting, G. J. (2006). Differential upregulation of Nox homologues of NADPH oxidase by tumor necrosis factor- α in human aortic smooth muscle and embryonic kidney cells. *J Cell Mol Med.* 10, 231-9.
- Montecino-Rodriguez, E., Leathers, H., Dorshkind, K. (2000). Expression of connexin 43 (Cx43) is critical for normal hematopoiesis. *Blood.* 96, 917-24.
- Muller-Borer, B. J., Cascio, W. E., Anderson, P. A., Snowwaert, J. N., Frye, J. R., Desai, N., Esch, G. L., Brackham, J. A., Bagnell, C. R., Coleman, W. B., Grisham, J. W., Malouf, N. N. (2004). Adult-derived liver stem cells acquire a cardiomyocyte structural and functional phenotype ex vivo. *Am J Pathol.* 165, 135-45.
- Navarro, F., Lieberman, J. (2010). Small RNAs guide hematopoietic cell differentiation and function. *J Immunol.* 184, 5939-47.
- Neijssen, J., Herberts, C., Drijfhout, J. W., Reits, E., Janssen, L., Neefjes, J. (2005). Cross-presentation by intercellular peptide transfer through gap junctions. *Nature.* 434, 83-8.
- Nishi, M., Kumar, N. M., Gilula, N. B. (1991). Developmental regulation of gap junction gene expression during mouse embryonic development. *Dev Biol.* 146, 117-30.


- Oviedo, N. J., Levin, M. (2007). *smadinx-11* is a planarian stem cell gap junction gene required for regeneration and homeostasis. *Development*. 134, 3121-31.
- Oyamada, Y., Komatsu, K., Kimura, H., Mori, M., Oyamada, M. (1996). Differential regulation of gap junction protein (connexin) genes during cardiomyocytic differentiation of mouse embryonic stem cells in vitro. *Exp Cell Res*. 229, 318-26.
- Ozog, M. A., Siushansian, R., Naus, C. C. (2002). Blocked gap junctional coupling increases glutamate-induced neurotoxicity in neuron-astrocyte co-cultures. *J Neuropathol Exp Neurol*. 61, 132-41.
- Parekkadan, B., Berdichevsky, Y., Irimia, D., Leeder, A., Yarmush, G., Toner, M., Levine, J. B., Yarmush, M. L. (2008). Cell-cell interaction modulates neuroectodermal specification of embryonic stem cells. *Neurosci Lett*. 438, 190-195.
- Park, B. C., Thapa, D., Lee, J. S., Park, S. Y., Kim, J. A. (2009). Troglitazone inhibits vascular endothelial growth factor-induced angiogenic signaling via suppression of reactive oxygen species production and extracellular signal-regulated kinase phosphorylation in endothelial cells. *J Pharmacol Sci*. 111, 1-12.
- Patterson, C., Ruef, J., Madamanchi, N. R., Barry-Lane, P., Hu, Z., Horaist, C., Ballinger, C. A., Brasier, A. R., Bode, C., Runge, M. S. (1999). Stimulation of a vascular smooth muscle cell NAD(P)H oxidase by thrombin. Evidence that p47(phox) may participate in forming this oxidase in vitro and in vivo. *J Biol Chem*. 274, 19814-22.
- Pearson, R. A., Dale, N., Llaudet, E., Mobbs, P. (2005). ATP released via gap junction hemichannels from the pigment epithelium regulates neural retinal progenitor proliferation. *Neuron*. 46, 731-44.
- Pedruzzi, E., Guichard, C., Ollivier, V., Driss, F., Fay, M., Prunet, C., Marie, J. C., Pouzet, C., Samadi, M., Elbim, C., O'Dowd, Y., Bens, M., Vandewalle, A., Gougerot-Pocidalo, M. A., Lizard, G., Ogier-Denis, E. (2004). NAD(P)H oxidase Nox-4 mediates 7-ketocholesterol-induced endoplasmic reticulum stress and apoptosis in human aortic smooth muscle cells. *Mol Cell Biol*. 24, 10703-17.
- Pera, M. F., Andrade, J., Houssami, S., Reubinoff, B., Trounson, A., Stanley, E. G., Ward-van Oostwaard, D., Mummery, C. (2004). Regulation of human embryonic stem cell differentiation by BMP-2 and its antagonist noggin. *J Cell Sci*. 117, 1269-80.
- Peshavariya, H., Dusting, G. J., Jiang, F., Halmos, L. R., Sobey, C. G., Drummond, G. R., Selemidis, S. (2009). NADPH oxidase isoform selective regulation of endothelial cell proliferation and survival. *Naunyn Schmiedeberg's Arch Pharmacol*. 380, 193-204.
- Petry, A., Djordjevic, T., Weitnauer, M., Kietzmann, T., Hess, J., Gorlach, A. (2006). NOX2 and NOX4 mediate proliferative response in endothelial cells. *Antioxid Redox Signal*. 8, 1473-84.
- Ploemacher, R. E., Mayen, A. E., De Koning, A. E., Krenacs, T., Rosendaal, M. (2000). Hematopoiesis: Gap Junction Intercellular Communication is Likely to be Involved in Regulation of Stroma-dependent Proliferation of Hemopoietic Stem Cells. *Hematology*. 5, 133-147.
- Rao, G. N., Berk, B. C. (1992). Active oxygen species stimulate vascular smooth muscle cell growth and proto-oncogene expression. *Circ Res*. 70, 593-9.

- Rosendaal, M., Green, C. R., Rahman, A., Morgan, D. (1994). Up-regulation of the connexin43+ gap junction network in haemopoietic tissue before the growth of stem cells. *J Cell Sci.* 107 (Pt 1), 29-37.
- Rosendaal, M., Mayen, A., de Koning, A., Dunina-Barkovskaya, T., Krenacs, T., Ploemacher, R. (1997). Does transmembrane communication through gap junctions enable stem cells to overcome stromal inhibition? *Leukemia.* 11, 1281-9.
- Russo, R. E., Reali, C., Radmilovich, M., Fernandez, A., Trujillo-Cenoz, O. (2008). Connexin 43 delimits functional domains of neurogenic precursors in the spinal cord. *J Neurosci.* 28, 3298-309.
- Saez, J. C., Berthoud, V. M., Branes, M. C., Martinez, A. D., Beyer, E. C. (2003). Plasma Membrane Channels Formed by Connexins: Their Regulation and Functions. *Physiol Rev.* 83, 1359-1400.
- Saez, J. C., Connor, J. A., Spray, D. C., Bennett, M. V. (1989). Hepatocyte gap junctions are permeable to the second messenger, inositol 1,4,5-trisphosphate, and to calcium ions. *Proc Natl Acad Sci U S A.* 86, 2708-12.
- Sauer, H., Rahimi, G., Hescheler, J., Wartenberg, M. (2000). Role of reactive oxygen species and phosphatidylinositol 3-kinase in cardiomyocyte differentiation of embryonic stem cells. *FEBS Lett.* 476, 218-23.
- Sauer, H., Wartenberg, M. (2005). Reactive oxygen species as signaling molecules in cardiovascular differentiation of embryonic stem cells and tumor-induced angiogenesis. *Antioxid Redox Signal.* 7, 1423-34.
- Schmelter, M., Ateghang, B., Helmig, S., Wartenberg, M., Sauer, H. (2006). Embryonic stem cells utilize reactive oxygen species as transducers of mechanical strain-induced cardiovascular differentiation. *Faseb J.* 20, 1182-4.
- Sosinsky, G. E., Nicholson, B. J. (2005). Structural organization of gap junction channels. *Biochim Biophys Acta.* 1711, 99-125.
- Stone, J. R., Yang, S. (2006). Hydrogen peroxide: a signaling messenger. *Antioxid Redox Signal.* 8, 243-70.
- Stubbs, S., Dilley, R., Peshavariya, H., Dusting, G. (2010). Hypoxic Preconditioning Improves Survival of Human Adipose-derived Stem Cells for Cardiac Tissue Engineering. *Basic & Clinical Pharmacology & Toxicology.* 107 (s), 128.
- Sturrock, A., Cahill, B., Norman, K., Huecksteadt, T. P., Hill, K., Sanders, K., Karwande, S. V., Stringham, J. C., Bull, D. A., Gleich, M., Kennedy, T. P., Hoidal, J. R. (2006). Transforming growth factor-beta1 induces Nox4 NAD(P)H oxidase and reactive oxygen species-dependent proliferation in human pulmonary artery smooth muscle cells. *Am J Physiol Lung Cell Mol Physiol.* 290, L661-L673.
- Suh, Y. A., Arnold, R. S., Lassegue, B., Shi, J., Xu, X., Sorescu, D., Chung, A. B., Griendling, K. K., Lambeth, J. D. (1999). Cell transformation by the superoxide-generating oxidase Mox1. *Nature.* 401, 79-82.
- Tai, M. H., Olson, L. K., Madhukar, B. V., Linning, K. D., Van Camp, L., Tsao, M. S., Trosko, J. E. (2003). Characterization of gap junctional intercellular communication in immortalized human pancreatic ductal epithelial cells with stem cell characteristics. *Pancreas.* 26, e18-26.

- Todorova, M. G., Soria, B., Quesada, I. (2008). Gap junctional intercellular communication is required to maintain embryonic stem cells in a non-differentiated and proliferative state. *J Cell Physiol.* 214, 354-62.
- Ushio-Fukai, M., Alexander, R. W., Akers, M., Griendling, K. K. (1998). p38 Mitogen-activated protein kinase is a critical component of the redox-sensitive signaling pathways activated by angiotensin II. Role in vascular smooth muscle cell hypertrophy. *J Biol Chem.* 273, 15022-9.
- Ushio-Fukai, M., Alexander, R. W., Akers, M., Yin, Q., Fujio, Y., Walsh, K., Griendling, K. K. (1999). Reactive oxygen species mediate the activation of Akt/protein kinase B by angiotensin II in vascular smooth muscle cells. *J Biol Chem.* 274, 22699-704.
- Ushio-Fukai, M., Tang, Y., Fukai, T., Dikalov, S. I., Ma, Y., Fujimoto, M., Quinn, M. T., Pagano, P. J., Johnson, C., Alexander, R. W. (2002). Novel role of gp91(phox)-containing NAD(P)H oxidase in vascular endothelial growth factor-induced signaling and angiogenesis. *Circ Res.* 91, 1160-7.
- Valiunas, V., Bukauskas, F. F., Weingart, R. (1997). Conductances and selective permeability of connexin43 gap junction channels examined in neonatal rat heart cells. *Circ Res.* 80, 708-19.
- Valiunas, V., Doronin, S., Valiuniene, L., Potapova, I., Zuckerman, J., Walcott, B., Robinson, R. B., Rosen, M. R., Brink, P. R., Cohen, I. S. (2004). Human mesenchymal stem cells make cardiac connexins and form functional gap junctions. *J Physiol.* 555, 617-26.
- Valiunas, V., Polosina, Y. Y., Miller, H., Potapova, I. A., Valiuniene, L., Doronin, S., Mathias, R. T., Robinson, R. B., Rosen, M. R., Cohen, I. S., Brink, P. R. (2005). Connexin-specific cell-to-cell transfer of short interfering RNA by gap junctions. *J Physiol.* 568, 459-68.
- Villars, F., Guillotin, B., Amedee, T., Dutoya, S., Bordenave, L., Bareille, R., Amedee, J. (2002). Effect of HUVEC on human osteoprogenitor cell differentiation needs heterotypic gap junction communication. *Am J Physiol Cell Physiol.* 282, C775-85.
- Vine, A. L., Bertram, J. S. (2002). Cancer chemoprevention by connexins. *Cancer Metastasis Rev.* 21, 199-216.
- Wang, X., McCullough, K. D., Franke, T. F., Holbrook, N. J. (2000). Epidermal growth factor receptor-dependent Akt activation by oxidative stress enhances cell survival. *J Biol Chem.* 275, 14624-31.
- Weissman, T. A., Riquelme, P. A., Ivic, L., Flint, A. C., Kriegstein, A. R. (2004). Calcium waves propagate through radial glial cells and modulate proliferation in the developing neocortex. *Neuron.* 43, 647-61.
- Wen, C. M., Cheng, Y. H., Huang, Y. F., Wang, C. S. (2008). Isolation and characterization of a neural progenitor cell line from tilapia brain. *Comp Biochem Physiol A Mol Integr Physiol.* 149, 167-80.
- Wolvetang, E. J., Pera, M. F., Zuckerman, K. S. (2007). Gap junction mediated transport of shRNA between human embryonic stem cells. *Biochem Biophys Res Commun.* 363, 610-5.

- Wong, R. C., Davidson, K. C., Leung, J., Pera, M. F., Pebay, A. (2009). Acute effect of endothelins on intracellular communication of human embryonic stem cells. *Journal of Stem Cells*. 4, 47-55.
- Wong, R. C., Dottori, M., Koh, K. L., Nguyen, L. T., Pera, M. F., Pebay, A. (2006). Gap junctions modulate apoptosis and colony growth of human embryonic stem cells maintained in a serum-free system. *Biochem Biophys Res Commun*. 344, 181-8.
- Wong, R. C., Pebay, A., Nguyen, L. T., Koh, K. L., Pera, M. F. (2004). Presence of functional gap junctions in human embryonic stem cells. *Stem Cells*. 22, 883-9.
- Worsdorfer, P., Maxeiner, S., Markopoulos, C., Kirfel, G., Wulf, V., Auth, T., Urschel, S., von Maltzahn, J., Willecke, K. (2008). Connexin expression and functional analysis of gap junctional communication in mouse embryonic stem cells. *Stem Cells*. 26, 431-9.
- Xu, R. H., Chen, X., Li, D. S., Li, R., Addicks, G. C., Glennon, C., Zwaka, T. P., Thomson, J. A. (2002). BMP4 initiates human embryonic stem cell differentiation to trophoblast. *Nat Biotechnol*. 20, 1261-4.
- Yang, S. R., Cho, S. D., Ahn, N. S., Jung, J. W., Park, J. S., Jo, E. H., Hwang, J. W., Jung, J. Y., Kim, T. Y., Yoon, B. S., Lee, B. H., Kang, K. S., Lee, Y. S. (2005). Role of gap junctional intercellular communication (GJIC) through p38 and ERK1/2 pathway in the differentiation of rat neuronal stem cells. *J Vet Med Sci*. 67, 291-4.

IntechOpen



Embryonic Stem Cells: The Hormonal Regulation of Pluripotency and Embryogenesis

Edited by Prof. Craig Atwood

ISBN 978-953-307-196-1

Hard cover, 672 pages

Publisher InTech

Published online 26, April, 2011

Published in print edition April, 2011

Pluripotency is a prerequisite for the subsequent coordinated differentiation of embryonic stem cells into all tissues of the body. This book describes recent advances in our understanding of pluripotency and the hormonal regulation of embryonic stem cell differentiation into tissue types derived from the ectoderm, mesoderm and endoderm.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Alice Pébay, Hitesh Peshavariya, Raymond C.B. Wong and Gregory J Dusting (2011). Non-classical Signalling Mechanisms in Stem Cells, Embryonic Stem Cells: The Hormonal Regulation of Pluripotency and Embryogenesis, Prof. Craig Atwood (Ed.), ISBN: 978-953-307-196-1, InTech, Available from: <http://www.intechopen.com/books/embryonic-stem-cells-the-hormonal-regulation-of-pluripotency-and-embryogenesis/non-classical-signalling-mechanisms-in-stem-cells>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2011 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike-3.0 License](https://creativecommons.org/licenses/by-nc-sa/3.0/), which permits use, distribution and reproduction for non-commercial purposes, provided the original is properly cited and derivative works building on this content are distributed under the same license.

IntechOpen

IntechOpen