

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

185,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Management Factors Affecting Fertility in Sheep

Pilar Santolaria, Inmaculada Palacin and Jesús Yániz
*Instituto Universitario de Ciencias Ambientales y Departamento de Producción Animal y
 Ciencia de los Alimentos. Escuela Politécnica Superior, Huesca
 Universidad de Zaragoza
 Spain*

1. Introduction

An Arab horse breeder in the early 13th century carried out the first insemination reported, by trapping stallion semen in wool placed in the vagina of a mare and transferring this to the vagina of another mare (Heape, 1898). Later, in 1780, an Italian priest and physiologist named Lazzaro Spallanzani performed artificial insemination with dog semen, and revolutionised the way scientists thought. Since then, scientist and farmers have striven to improve this technology, motivated by the benefits that could be achieved. Sheep is one of the species subsequently linked to this technology and in which many questions still remain to be resolved to improve fertility. However, the potential impact of this technique on the genetic progress of sheep is high and further studies are needed to improve its efficiency.

In Spain, artificial insemination programmes in sheep are linked to the genetic selection schemes of the breeds, but it has not been successfully integrated with reproductive technology on farms as happen in sows or cows. The technical difficulty and weak fertility, ranged between 15 to 60 % for pregnancy rate, limits its application. In Spain, most commercial programmes use refrigerated semen (15 degrees C) by superficial intracervical deposition (cervical), whereas the use of frozen-thawed semen by intrauterine deposition (laparoscopy) is more restricted. Cervical insemination with fresh semen is the main method chosen due to its simplicity and satisfactory results. With the aim of improving its efficiency, this paper focuses on identifying the main management effects affecting AI results when this technology is applied.

2. Female associated factors

Management factors associated with artificial insemination in the ewe can modify fertility. In reproductive planning, intervals between lambings, season, age of ewe, heat stress, nutrition state or breed are some of the factors which have a great effect on fertility results. David et al. (2008), using a joint model combining two main traits, one relative to female and the other relative to the male, reported that the main variation factors of AI success were relative to non-sex-specific effects and to female effect, suggesting that choosing females to inseminate might slightly improve the AI results.

2.1 Season

Seasonal variations are described as a limiting factor in sheep reproduction. In natural conditions, seasonality, which is mediated by photoperiod, modifies hormonal balance and causes seasonal reproductive variations in sheep (Karsch, et al. 1984; Yeates, 1949), giving rise to a decrease in reproductive activity during long days (anoestrous season). Photoperiodic information is translated into neuroendocrine changes through variations in melatonin secretion from the pineal gland (Bittman, et al., 1983). Melatonin, secreted in pineal gland, triggers variations in the secretion of luteinising hormone-releasing hormone (GnRH), luteinising hormone (LH) and follicle stimulating hormone (FSH) (Arendt, et al., 1983, Karsch, et al., 1984). In any case, seasonal changes in reproductive activity are clearly defined in sheep breeds from high latitudes ($>40^\circ$) (Pelletier, et al., 1987), where the differences in daylight duration between short days and long days are more notable.

As in natural mating, season affects fertility after AI, although hormonal treatment is used to synchronise and induce oestrus. Windsor (1995) reported low cervical AI fertility rates in non-breeding season in Merino ewes, a shallow seasonal breed. According to this, Anel et al. (2005) found a season effect on the AI fertility in Churra ewes, which was more important in cervical than laparoscopic artificial insemination. In cervical AI, semen is deposited in the external portion of the cervix and the sperm transport is affected by cervical mucus quality. These authors suggest that photoperiod could alter progestagens and so cervical mucus characteristics, making it scarcer and more viscous. In consequence, sperm transport in the cervix can be interfered with. It is important to note that seasonality affects the ram reproductive parameters in the same way and changes in seminal quality during anoestrous season may decrease the fertility results after AI.

Moreover, subcutaneous melatonin implants are widely used to bring the breeding season forward and improve reproductive performance in non-breeding season (Chemineau, et al., 1991; Haresign, et al., 1990). Melatonin treatments act by mimicking a short-day-like response (O'Callaghan, et al., 1991) and induce oestrus during the non-breeding season. Not only an increase in the percentage of pregnant ewes (fertility) has been described after melatonin implant treatment in anoestrous season, but also the number of lambs born per ewe (litter size) (Abecia, et al., 2007; Arrebola, et al., 2009; Chemineau, et al., 1992). This improvement could be due to a higher rate of embryonic survival, an improvement in luteal function or a reduction in the antiluteolytic mechanisms (Abecia, et al., 2008). In AI, a fertility rate increase has been reported with melatonin implant treatments (Laliotis, et al., 1998; Legaz, et al., 2000) after cervical AI.

2.2 Age

Another important factor affecting fertility after cervical AI is ewe age. In comparison with adult ewes, young and maiden ewes have lower fertility, probably due to impaired sperm transport combined with low mucus production in the cervical canals during oestrus (Selaive-Villarreal & Kennedy, 1983a, b). After that, most studies have described a decrease in AI success with increasing female age. Shackell et al. (1990), predicted 2-3% fertility reduction per year of age for different breeds. Esmailizadeh et al., (2009) reported that as the age of the ewes increased from 2 to 7 years, the proportion of barren ewes significantly decreased from 29 to 5%. In a more recent study in the Spanish Churra dairy breed, Anel et al. (2005), described how, after 1.5 years of age, the lambing rate decreased by 1.74% per year for cervical AI. The highest rates of fertility declining with age were described in the Lacaine breed by Colas et al. (1973), who reported a drop in fertility of 15% per year.

Paulenz et al. (2007), observed that the age of the ewes had a significant effect on the non-return rate, but not on lambing rate, whereas in Fukui et al. (2010), both the pregnancy and lambing rates in the ewes significantly declined as age increased. The detrimental effect of increased fertility age could be explained by the fact that aged ewes have increased risks of reproductive disorders and decreased ovulation rates with quality ovulated oocytes compared with younger ewes.

The question is: what is the optimal age for cervical AI? Alabart et al. (2002), studied the influence of age on fertility in a total of 3819 Rasa Aragonesa ewes aged 1 to 12 years. In this study, maximum fertility (56.7%) was observed at 3 years, and ewes aged from 2 to 5 years (79.5% of the inseminated ewes) had mean fertility values above 50%. These results partially confirm the observations by Colas et al. (1973), who reported a significant decrease in the fertility of ewes inseminated when over 3.5 years old, and Gabiña and Folch (1987), who observed a strong fall in the fertility of ewes inseminated at 4 or more years of age. Anel et al. (2005) recorded the best fertility rates in ewes aged between 1.5 and 4.5 years; beyond this age, fertility declined remarkably. Fertility also decreased depending on the number of previous parturitions. In other studies, the maximum fertility was obtained at around 2 years of age, with a progressive fall afterwards (Fantova, et al., 1998). It may be concluded that insemination groups should be made up of 2 to 5 year old ewes while younger and older ewes should be used for natural mating.

2.3 Lambing-AI interval

The need for a resting period for the ewe after lambing to allow uterine involution is well known. However, sometimes the increasing reproductive rate imposed by the demanding production system involves short resting periods from lambing to AI, which affects fertility in a negative way. According to Bodin et al. (1999), reducing the lambing-AI interval to below 40-50 days induces a significant decrease in fertility, even after natural mating. Most authors recommend not inseminating ewes any sooner than 50 days post-partum (Anel, et al., 2005).

2.4 Breed

Ewe breed is also a significant source of variation in fertility after AI (Donovan, et al., 2004; Fukui, et al., 2007; Papadopoulos, et al., 2005; Salamon & Maxwell, 1995). Differences in the mean time of ovulation and ovulation rates in different breeds of ewes at different locations may explain the variation in fertility (Salamon & Maxwell, 1995).

Alternatively, variation may be due to differences in the morphometric characteristics of the cervix (Eppleston, et al., 1994). In this sense, Kaabi et al. (2006), carried out a morphometric study in four ovine breeds (Assaf, Churra, Castellana and Merino) showing important differences in length, width, number of folds and distance between folds, which originates breed variations in the depth of catheter penetration into the cervix during AI. In this study, the breeds yielding lower fertility after AI resulted in higher cervical complexity, and achieved a lesser degree of cervical penetration of the catheter during cervical AI. As explained below, different studies have found a positive correlation between cervical AI depth and fertility (Kaabi, et al., 2006).

2.5 Body weight and body condition score

For an adequate response in a breeding programme, ewes must be suitably nourished and maintained in good body condition. Clearly, ewes with a good nutrient intake respond most

rapidly to the onset of the breeding season and continue to respond with an increase in ovulation rate (Keisler & Buckrell, 1997). Flushing is understood as the rapid increase in ovulation rate of ewes receiving a nutrient supplementation before mating. Under harsher nutritional conditions in the semi-arid southern Mediterranean region, where regular food supply is not guaranteed, lambing and twinning rates were shown to be boosted following nutritional flushing (Younis, et al., 1978; Landau & Molle, 1997; Branca, et al., 2000) or when the live weights of the ewes were higher at mating (Gunn & Doney, 1979, Thomson & Bahhady, 1988).

However, Lassoued et al. (2004), showed important interactions between genotype and level of nutrition. In this sense, in highly prolific ewes like D'Man breed, higher levels of nutrition prior to and during mating were associated with improved reproductive performance, but in low prolific breeds such as Queue Fine de l'Ouest, neither ovulation rate nor lambing rate were affected by the dietary treatment. In a recent work by Fukui et al. (2010) body weight did not significantly affect fertility.

Body condition score (BCS) has proved useful as a management tool for subjectively assessing the nutritional status of ewes. In this way, body condition directly affects hypothalamic activity and GnRH secretion, but not pituitary sensitivity to GnRH, and these effects on reproductive performance are also mediated through changes in ovarian hormones or in hypothalamo-pituitary sensitivity to ovarian hormones (Rhind, et al., 1989). The effect of body condition on the ovulation rate of ewes has been extensively reported (Ducker & Boyd, 1977; Morley, et al., 1978; Adalsteinsson, 1979; Gunn & Doney, 1979; Gonzalez, et al., 1997). High body condition score has been associated with an increase of ovulation (Rhind, et al., 1989; Xu, et al., 1989), especially in Mediterranean breeds at the beginning of the breeding season (Forcada, et al., 1992). Most authors recommend a BCS of 2.5 to 3.0 either for natural or artificially breeding (Contreras-Solis, et al., 2009; Husein & Ababneh, 2008). In a study carried out in inseminated Rasa Aragonesa ewes (Bru, et al., 1995), the lowest pregnancy rates (32,7 %) were obtained in sheep with a BCS<2, the average values (48.3%) with BCS between 3 and 2 and the higher values (58.8%) when BCS was >3. The importance of BCS in fertility has been also reported in Spanish Manchega breed (Montoro, 1995).

Fukui et al. (2010), , concluded that body nutritional condition is an important factor, next to ewe age, influencing the fertility of ewes after AI regardless of body weight. Nulliparous ewes less than 3 years old and with a BCS of more than 3.0 are expected to have higher fertility than other types of ewes.

2.6 Farm/Herd

Different ewe farms have different management practices and this may have an impact on fertility after AI. Reproductive planning (intervals between lambings, season, age of first mating, AI technique, etc.) and animal handling (feeding, health, preparation of AI lots, etc.) have a great effect on fertility results (Anel, et al., 2005). The significant effect of the farm has been described in several studies (Anel, et al., 2005, Fantova, et al., 1999, Paulenz, et al., 2002). Thus, in order to improve fertility results, handling conditions on farms must be improved, along with more widespread use of AI techniques.

The geographical area where the farm is located may also have an influence on the success of AI. In a recent study carried out in north-eastern Spain (Palacin et al., 2011), data from 18.528 AI in Rasa Aragonesa ewes belonging to a selection scheme with similar management

were recorded in order to analyse the effect of farm geographical location and the theoretical time distance between the farm and insemination centre on the fertility after cervical AI. An average fertility of 54.3% was observed, with significant differences among the 14 regions studied. Fertility rates higher than 60% were found in the northern regions near the Pyrenees Mountains and the lowest results (38.5-48.3%) were obtained in the southern regions. The average time distance of these regions did not differ. The regions nearest to the insemination centre, with similar climatic conditions, showed medium fertility rate (54.0-57.8%). These results showed a huge variability after insemination, taking into account the geographical location of the farm.

2.7 Heat stress

It has been reported that in tropical and sub-tropical areas the local sheep show restricted sexual activity in the summer months (Marai et al., 2004). Marai et al., 2007 reviewed how exposure to high ambient temperature causes impairment of reproductive functions in sheep. The heat effect is aggravated when heat stress is accompanied with high ambient humidity (Marai et al., 2000, 2004, 2006, 2007). Heat stress evokes a series of drastic changes in animal biological functions, which include a decrease in feed intake efficiency and use, disturbances in the metabolism of water, protein, energy and mineral balances, enzymatic reactions, hormonal secretions and blood metabolites. (Shelton, 2000; Marai et al., 2006).

2.8 Synchronization treatment

Oestrous synchronisation and ovulation induction treatments are widely spread in AI in order to control the optimum insemination time. Oestrous behaviour in small ruminants is not clearly shown, so treatments are needed to prevent an asynchrony between ovulation and insemination time, which may be the commonest cause of failure of artificial insemination programmes (Jabbour & Evans, 1991).

Synchronisation treatments are a useful tool not only for AI programmes but also in natural mating, particularly to ensure lambing in anoestrous season. However, many studies report that synchronisation causes reduced fertility after cervical AI (Robinson, et al., 1970; Hawk, 1971) and after natural mating (Quinlivan & Robinson, 1969; Hawk & Conley, 1975; Allison & Kelly, 1978). Different hormonal treatments have been used in the control of sheep reproduction, but progestagen analogues (fluorogestone acetate and medroxyprogesterone acetate) are the most commonly used to induce and synchronise oestrus in natural mating or AI of small ruminants (Lunstra & Christenson, 1981; Pearce & Robinson, 1985; Langford, et al., 1982; Baril, et al., 1993; Greyling, et al., 1997). Progestagens produce a mimetic effect of the luteal phase of the oestrous cycle and a sudden progestagen removal followed by administration of an equine chorionic gonadotropin (eCG) dose (FSH- and LH-like stimulation) induces oestrous activity. Currently, intravaginal progestagen-impregnated devices (sponges or CIDR) for 12-14 days followed by administration of 250-500IU eCG has been proposed (Abecia, et al., 2011) as a synchronised treatment in sheep, in which insemination can be performed from 47h (intrauterine) to 55 (cervical) hours after device removal.

Although long-term progestagen treatment results in efficient oestrous synchronisation, high variability has been reported in fertility (Vinoles, et al., 2001; Menchaca & Rubianes, 2004). Progestagen treatment appears to result in an asynchrony between oestrus and ovulation (Scaramuzzi, et al., 1988; Sirois & Fortune, 1990) and reduce sperm transport

through the cervix (Killen & Caffery, 1982; Armstrong & Evans, 1984; Pearce & Robinson, 1985). Other studies have reported lower fertility rate caused by the negative effect of long-term progesterone treatment on oocyte development (Vinoles, et al., 2001; Menchaca & Rubianes, 2004) related to subluteal progesterone levels.

According to this, recent research efforts are focused on shortening synchronisation treatments. Shortening treatment (5-6days) with different progestagen devices seems to be enough to achieve efficient oestrous synchronisation in natural mounting both during the anoestrous season (Ungerfeld & Rubianes, 1999) and in breeding season (Vinoles, et al., 1999; Ustuner, et al., 2007) with a similar fertility rate to long-term treatment. Nevertheless, the interval between progestagen device withdrawal and the onset of oestrus was shortened (Ungerfeld & Rubianes, 1999; Vinoles, et al., 2001; Zeleke, et al., 2005; Ustuner, et al., 2007) due to a delay in corpus luteus regression in cyclic ewes (Menchaca & Rubianes, 2004). This asynchrony could decrease the success of artificial insemination programmes, so a treatment that ensures an acceptable luteolysis seems to be necessary to enhance the oestrous synchronisation (Ustuner, et al., 2007).

The luteolytic effect of prostaglandin $F_{2\alpha}$ ($PGF_{2\alpha}$) (McCracke, et al., 1972) and its analogues has been used to control corpus luteus activity. In goats, the use of a short-term progestagen protocol combined with $PGF_{2\alpha}$ administered at progestagen sponge insertion time was successful in artificial insemination with frozen-thawed semen (Corteel, et al., 1988). In sheep, Beck et al (1993) reported acceptable oestrous synchronisation and fertility results in natural mating using $PGF_{2\alpha}$ treatment combined with short-term progestagen. Nowadays, $PGF_{2\alpha}$ treatment use in small ruminant reproduction control has increased because of narrow restrictions on the use of progestagens in animal production both in the United States and the European Union (Martin, et al., 2004; Menchaca & Rubianes, 2004). $PGF_{2\alpha}$ is quickly metabolised with a minimum residue level (Light, et al., 1994). In some South American countries, short progestagen treatment with $PGF_{2\alpha}$ administration is used in goat management (Menchaca & Rubianes, 2004). However, the effectiveness of $PGF_{2\alpha}$ or its analogues depends on the ovary status (Houghton, et al., 1995) and is not very useful during the anoestrous season (Acritopoulou & Haresign, 1980). Moreover, a varied response and low pregnancy rates in AI have been reported with single $PGF_{2\alpha}$ treatment or progestagen combined (Boland, et al., 1978; Hackett, et al., 1981; Olivera-Muzante, et al., 2011), so with the current methods it is not an appropriate synchronisation treatment for AI unless previous oestrous detection is carried out.

Another factor affecting fertility after cervical AI related with hormonal synchronisation is the development of anti-eCG antibodies. Females involved in repeated treatment throughout their reproductive life, particularly those involved in AI genetic programmes or intensive lambing systems, develop anti-eCG antibodies (Baril, et al., 1996; Bodin, et al., 1997; Roy, et al., 1999). These antibodies are associated with low reproductive rates, especially in fixed time AI in sheep (Maurel, et al., 2003) and goats (Baril, et al., 1996), and in multiple ovulated ewes (Forcada, et al., 2011). This is less pronounced when repeated treatment is combined with natural mating. High concentrations of anti-eCG antibodies are reported with a lack or delay in oestrus and pre-ovulatory LH surge (Baril, et al., 1993; Roy, et al., 1999; Maurel, et al., 2003), decreasing the fertility after AI.

Alternatives for reproductive activity control that ensure optimum oestrous synchronisation and successful AI results are needed in sheep, in line with current demands in public and animal welfare.

3. Male associated factors

The ram may greatly influence fertility results after cervical AI. It has been reported that variation in fertility of ram ejaculates exists independently of the sperm quality (Choudhry, et al., 1995; Paulenz, et al., 2002). Variations in the fertility of rams have been reported after cervical inseminations with fresh semen (Anel, et al., 2005; Paulenz, et al., 2002), with frozen semen (Colas, 1979; Windsor, 1997; Soderquist, et al., 1999; Paulenz, et al., 2005, 2007) and after laparoscopic inseminations with frozen semen (Eppleston et al., 1986; Maxwell, 1986; Eppleston, et al., 1991; Eppleston & Maxwell, 1995). In a large scale epidemiological study, Anel et al. (2005) observed that the male factor significantly influenced fertility. Despite the restrictions in the choice of ejaculates, the authors found important differences in fertility among rams, particularly when cervical AI with cooled semen was used. Salamon and Maxwell (1995) proposed that ram differences in fertility could be both genetic and environmental, whereas ejaculate differences are probably due to nutrition, management and previous frequency of ejaculation.

Whereas differences in fertility have been demonstrated among fertile males in different species, the causes of these differences remain unclear (Ostermeier, et al., 2001). Saacke et al. (1988, 1994) have suggested in bulls that factors associated with semen quality which affect fertility can be classified as either compensable or non-compensable. It was suggested that the effects of compensable factors on fertility might be sensitive to the number of sperm inseminated, whereas those of non-compensable factors were not. As the number of sperm inseminated increases, fertility increases until a plateau is reached (den Daas, 1992). At this point, compensable factors no longer have an effect on fertility. Commercial insemination of ovine in Mediterranean Countries provides at least the plateau number of sperm in an insemination dose. It is thus the non-compensable factors that contribute most to the fertility level of a ram. A non-compensable defect in sperm would be one in which a sperm reaches the fertilisation site and initiates the egg activation process, but fails to sustain zygotic, embryonic, or foetal development (Ostermeier, et al., 2001). Evidence of such defects in sperm has been described in bulls with fertility differences (Eid, et al., 1994). Likely candidates for non-compensable factors would be incorrectly assembled chromatin or damaged DNA within the sperm nucleus. It seems logical to assume that the transfer of a complete and intact DNA molecule from sperm to ovum is crucial to obtain fertilisation with certain prospects of success. It is well-known that the presence of defects in the genetic material, such as anomalies in chromatin condensation related with the sperm maturation process, the integrity of the DNA molecule associated with the presence of breaks both of single and double DNA strands, or the presence of chromosomal anomalies, are closely associated with infertility (Aravindan, et al., 1997).

Season

Although seasonality is less marked in male than in female, changes in testicular volume, hormonal profiles, sexual behaviour and semen quality that affect the reproductive performance of rams have been reported (Casao, et al., 2010a). In this sense, the treatment of rams with slow release implants of melatonin during the non-breeding season accounted for increased scrotal diameter and improved the reproductive performance of ewes inseminated during anoestrus with semen from these melatonin-implanted males. A direct beneficial action of melatonin on sperm motility (Casao, et al., 2010c) and on other ram sperm characteristics during the non-breeding season has recently been demonstrated, with decreased apoptotic-like changes and modulating capacitation and fertilisation rates (Casao, et al., 2010b).

Nutrition

Several studies on nutrition in rams have demonstrated that diet may have an effect on testis size and sperm production (Brown, 1994). It also has been described that specific components of the diet, such as Vitamin E, may have a positive effect in increasing semen quality and quantity (Yue, et al., 2010). The effect of diet of the rams on the reproductive success of ewes after AI remains to be determined.

4. Artificial insemination-associated techniques

Inadequate semen preservation and difficulty in passing through the cervix during AI are the major obstacles to the extensive use of cooled or cryopreserved semen in sheep AI programmes (Yaniz, et al., 2010, 2011). Exo-cervical deposition of diluted liquid ram semen, preserved at 15° C for less than 8 h, is currently the AI technique predominantly used in the Mediterranean countries (Lopez-Saez, et al., 2000; Yaniz, et al., 2010, 2011).

4.1 Semen collection

4.1.1 Semen collection frequency

Semen collection frequency may have an impact on sperm quality. Long abstinence periods (Pascual, 1993) and successive ejaculations (Ollero et al., 1994) have been associated with membrane alterations of spermatozoa. A decrease in semen volume and sperm concentration with successive ejaculations has been reported in several studies (Ollero et al., 1996; Kaya et al., 2002). In the study by Ollero et al. (1996), the maximum proportion of viable cells was obtained in the second ejaculate after an abstinence period of 3 days. The authors concluded that the use of the second and/or a mixture of second and third ejaculates would improve the results in artificial insemination. The general recommendation is to establish a routine of semen collection, for example of two-three times per week (two collections per day/per ram) on different and non-consecutive days, independently of the use of the semen obtained. Increased semen collection frequency may have an effect on sperm quality and the composition of the seminal plasma (Kaya et al., 2002), although it remains to be determined whether this has an impact on field fertility. In this sense, the procedure of taking one or two collections per day from each ram during the working week (Monday-Friday), with a 2-day rest period during the weekend, has been described in Ireland (Gordon, 1997).

4.1.2 Hygienic conditions

Semen collection in farm animal species is not a sterile procedure, and some degree of contamination with bacteria cannot be avoided (Clément et al., 1995; Varner et al., 1998; Althouse et al., 2000; Thibier and Guerin, 2000; Althouse and Lu, 2005; Aurich and Spengler, 2007; Bielanski, 2007; Yániz et al., 2010). In rams, semen is usually collected with an open-ended artificial vagina, which may be contaminated with bacteria from the surface of the penis and prepuce, collection area, equipment and people. As a consequence, bacteria might compromise semen quality during storage and contaminate the female's reproductive tract. We have recently described that ram semen is normally colonised by a variety of microorganisms that may reduce semen preservation and fertility (Yániz et al., 2010). In particular, the contamination of ram semen with enterobacterial species reduced sperm quality during storage at 15 °C in a concentration-dependent manner.

Different strategies may be taken to minimise the effects of bacterial contamination on extended semen, as the bacterial concentration remains below a threshold level, so fertility is not affected (Althouse et al., 2000). The first and most viable option is to enhance the hygienic measures during semen collection and processing. Dilution of the ejaculates with sterile diluents will further decrease the concentration of contaminants (Bielanski, 2007), although this aspect has low influence in ovine because of the high sperm concentration employed for AI. Finally, control of bacterial growth is usually performed by the use of semen extenders containing antibiotics with broad-spectrum bactericidal or bacteriostatic activity (Maxwell and Salamon, 1993; Salamon and Maxwell, 2000). Perhaps too much reliance is often placed on this method of bacterial control in ovine semen. In this species, necessarily short storage periods for semen determine that the control of bacterial multiplication may be less important than in other animal species in which successful long-life semen extenders have been developed. Interestingly, in a recent study (Yániz et al., 2010), it was observed that 13% of identified bacteria were simultaneously resistant to penicillin and streptomycin, the most common preservative antibiotic combination used in ovine semen extenders, whereas *E. coli*, the bacteria with the highest impact on sperm quality, was frequently resistant to both antibiotics (31.7 %, 13/41). Antibiotics with higher antimicrobial activities were gentamycin and ceftiofur, and their inclusion in ram semen diluents should be considered.

4.2 Semen evaluation

Semen evaluation is a useful tool in the selection of males and ejaculates for assisted reproduction. Traditional evaluation techniques, based on the subjective assessment of parameters such as sperm motility and morphology, semen volume or concentration, have long been employed in the diagnosis of male subfertility and sterility (Verstegen et al., 2002). An *in vitro* system that could accurately predict field fertility would facilitate stricter selection of AI rams with regard to the semen quality and would provide a valuable tool for increasing conception rate (Donovan et al., 2004). However, finding a laboratory test reliable enough to predict the potential fertility of a given semen sample or a given sire for AI is still considered utopian, as indicated by the modest correlations seen between results obtained *in vitro* and field fertility (Rodríguez-Martínez, 2003). Male fertility is complex, and depends upon a heterogeneous population of spermatozoa interacting at various levels of the female genital tract, the vestments of the oocyte, and the oocyte itself (Rodríguez-Martínez, 2003). For this reason, laboratory assessment of semen must include the testing of as many relevant sperm attributes for fertilisation and embryo development as possible, not only in individual spermatozoa but also within a large sperm population (Rodríguez-Martínez, 2003). In practice, routine sperm analysis requires fast, objective and accessible methods of assessing different aspects of sperm viability (Yániz et al., 2008). In this sense, the common use of computer-assisted sperm analysis (CASA) methods for sperm motility, of image analysis for the evaluation of membrane integrity (Yániz et al., 2008), of DNA fragmentation with sperm chromatin diffusion techniques (SCD), (Gosalvez et al., 2008; López-Fernández et al., 2008), or fluorimetry, etc., would theoretically improve the predictive capacity of semen analysis, although more studies are needed to determine the utility of these techniques in the practical use of AI.

4.3 Sperm number per AI dose

The difficulty in passing through the cervix during AI due to the type of cervical canal found in this species determines that semen can only be deposited inside the cervix, usually

in the external portion. The retention capacity of the ovine cervix is very low (0,1-0,2 ml) (Gordon , 1997), whereas a large sperm number per dose is required to compensate the huge barrier effect of the cervix (around 400×10^6 sperm/ dose in Spanish AI).

It is important to determine the minimal sperm dose per insemination in order to maximise the genetic diffusion of males, without decreasing AI success. In the study by Langford and Marcus (1982), fertility after insemination of 400 or 200×10^6 spermatozoa was similar to that observed after natural service at progestagen-induced oestrus. However, when less than or equal to 100×10^6 spermatozoa were inseminated, fertility fell markedly and the number of lambs per ewe inseminated decreased. These data indicate that insemination of 200×10^6 spermatozoa, i.e. less than 10% of the number in a single ram ejaculate, allows normal conception rates in progestagen-treated ewes. It seems that the minimal sperm dose may be between more than 100 and 200×10^6 (120 million sperm in Australian works, Gordon, 1997), although a breed-effect should not be discarded, as differences in the cervix anatomy have been described.

4.4 Semen preservation

4.4.1 Diluents

Despite numerous past efforts to improve semen diluents, few new additives have been introduced in the extender composition for ram semen (Yániz et al., 2005). Biological components such as milk or egg yolk in the diluent have not really been effective in the applied use for AI. So, for example, skimmed milk, a complex and variable biological component, is still the main diluent used to preserve sheep semen at 15 °C for AI in numerous countries (López-Sáez et al., 2000; Yaniz et al., 2011). The basic components of semi-synthetic diluents for the liquid storage of ram semen (buffers combined with sugars and egg yolk), have changed little since those first used in the early 20th century (Maxwell and Salamon, 1993).

The predominant empirical approach of most of the studies could partially explain the slow advances made in the development of chemically defined extenders for ram semen storage. An individual study on the effect of each extender component on the viability of the sperm cell could greatly contribute to the development of a more rational synthetic diluent. For example, in a recent work (Yániz et al., 2011) we studied the effect of different buffer systems included in the composition of well-defined and synthetic diluents (in vitro), on sperm quality parameters during storage at 15°C. TRIS caused drastic modifications of certain sperm kinematic parameters during storage at 15°C although, along with citrate, it is one of the main buffers included in the composition of semi-synthetic ram semen diluents (López et al., 1999; López-Sáez et al., 2000; Salamon and Maxwell, 2000; Paulenz et al., 2003; Martí et al., 2003; Yaniz et al., 2008).

With the available diluents, semen preservation in the non-frozen state is limited to 6-8h without reducing fertility (Maxwell and Salamon (1993); Salamon and Maxwell, 2000). Irrespective of the diluent, dilution rate, storage temperature or conditions, the sperm deteriorated as the storage duration increased. The main changes that occurred during storage included reduction in motility and morphological integrity of sperm. These changes may be attributed to the accumulation of the toxic products of metabolism, mainly of reactive oxygen species (ROS) that cause lipid peroxidation of the sperm plasma membranes. The above events are accompanied by a decline in transport and survival of spermatozoa in the female reproductive tract and reduction in fertility (Salamon and

Maxwell, 2000). When longer period of storage are required, the use of reduced temperature (about 4°C) and egg-yolk-based media is recommended (Gordon, 1997), although in this case the insemination time should also be considered (Fernandez-Abella et al., 2003)

4.4.2 Cold shock

Severe changes in temperature are a common feature of semen storage protocols for assisted reproduction, but are not biological traits to which species have become adapted (López-Fernández et al., 2008). In the ram, as in other mammals, there is a loss of semen quality when cooled semen samples are used for assisted reproduction. Cooled semen undergoes a decrease in sperm quality, which includes reduction in motility, destabilisation of sperm membranes and DNA integrity impairment of sperm function (López-Fernández et al., 2008; Muiño-Blanco et al., 2008). It is well known that ram spermatozoa are more sensitive to cold-shock stress than those of other species (Muiño-Blanco et al., 2008). In fact, the ram exhibits a faster DNA degradation under similar conditions than other species studied (Gosálvez et al., 2007). Temperature excursion episodes in spermatozoa are associated with oxidative stress induced by the generation of reactive oxygen species, which promote DNA fragmentation (López-Fernández et al., 2008). Reactive oxygen species, produced by dead spermatozoa during a sperm temperature reduction, give rise to sperm membrane alterations with the subsequent release into the media of free active enzymes. Then, the accumulation of toxic metabolic products and active free enzymes, such as those contained in the acrosome, is higher in the media as spermatozoa disintegrate. This could facilitate intact sperm degradation in an exponential fashion (López-Fernández et al., 2008). This loss of quality, accompanied by a decline in sperm survival in the female reproductive tract, gives rise to a reduction in fertility and increased embryonic loss (Paulenz et al., 2002). In ovine species, ovulation may take place several hours post-insemination (Cheminau et al., 1992; Romano, 2004) and, consequently, the time that a spermatozoon is able to survive after AI is of critical importance to achieve pregnancy. Consequently, obtained semen samples must be used as quickly as possible with the diluents currently available.

4.4.3 Time from semen recovery to dilution

The role of seminal plasma (SP) in mammalian sperm function remains largely a matter of speculation as both inhibitory and stimulating effects have been found (Muiño-Blanco et al., 2008). It has been reported that exposure of ram spermatozoa to seminal plasma causes a reduction of fertility (Dott et al., 1979), although some components of the seminal plasma, such as certain proteins, seem to have the important function of maintaining the stability of the membrane up to the process of capacitation, and are able to protect and repair the cold-shock damage to sperm (Muiño-Blanco et al., 2008). However, in practice, semen dilution in the ram has to be done as soon as possible after recovery to avoid the inhibitory effects of seminal plasma.

4.5 Insemination technique

4.5.1 Semen deposition site

In mammals, establishing an adequate sperm reservoir in the caudal isthmus and utero-tubaric junction is very important after mounting or AI, as spermatozoa may ascend to the fertilisation site from this reservoir. In ovine species, the cervix is the main anatomical and physiological barrier to the ascent of spermatozoa after mounting or AI. This is particularly

relevant after cervical insemination as, in comparison to fresh spermatozoa, a relatively small proportion of the stored cells penetrates the cervical canal and migrates through the uterus of the ewe to the oviducts (Salamon and Maxwell, 2000). Spermatozoa functionally affected during liquid storage may not migrate, or may migrate slowly, and their survival in the female tract is also reduced to about half that of fresh spermatozoa (Salamon and Maxwell, 2000). Attempts to improve the transport of spermatozoa from the posterior cervix to the oviducts of oestrous ewes by prostaglandins added to stored semen have given conflicting results (Maxwell and Salamon, 1993).

In the non-pregnant ewe, the funnel-shaped rings of the cervix, which average around five in number, are not concentrically aligned, and their openings are constricted in most instances to less than 3 mm (King et al., 2004). As explained above, breed is an important determinant of the morphology of the cervix, and that could at least partially explain differences in fertility after cervical AI (Kaabi et al., 2006). Many studies have found a positive correlation between the depth of cervical AI and fertility (Kaabi et al., 2006). In consequence, numerous efforts have been made to develop new methods to deposit the semen as deep as possible into the uterus. Studies based on the use of modified pipettes, or hormones such as oxytocin to dilate the cervical canal, have shown that cervical penetration can be improved. However, fertility results have been very variable (Kaabi et al., 2006). Special care should be taken to avoid cervical trauma with the catheter during AI, as it has been associated with reductions in pregnancy and lambing rates (Kaabi et al., 2006). Secondary effects of oxytocin may also have an adverse effect on fertility (King, 2004).

4.5.2 Technician

The ability of the inseminator may be another important source of variation of the outcome of sheep AI (Gordon, 1997; Anel et al., 2005). Cervical penetration rates are influenced by operator skill (Eppleston and Maxwell, 1993), and the establishment of training programmes is highly recommended.

4.5.3 Stress around AI

There is some evidence that nutritional or management stress inflicted upon the ewe around AI can markedly reduce fertility by interfering with fertilisation or by increasing early embryo mortality rates (Gordon, 1997). Ewes and rams should be handled with the minimum of disturbance and receive good nutrition around oestrus and the first weeks after AI.

5. Acknowledgements

The authors thank Neil Macowan for assistance with the English translation. This work was supported by the Araid Foundation (grant OTRI 2010-0464), Spanish MICINN (grant IPT-010000-2010-33) and OVIARAGON S.C.L. (grant OTRI OTRI 2010-0465).

6. References

- Abecia, J.A., Forcada, F. & Gonzalez-Bulnes, A. (2011). Pharmaceutical Control of Reproduction in Sheep and Goats. *Veterinary Clinics of North America-Food Animal Practice*, Vol.27, No.1, (Mar), pp. 67-+, 0749-0720

- Abecia, J.A., Forcada, F., Casao, A. & Palacin, I. (2008). Effect of exogenous melatonin on the ovary, the embryo and the establishment of pregnancy in sheep. *Animal*, Vol.2, No.3, pp. 399-404, 1751-7311
- Abecia, J.A., Valares, J.A., Forcada, F., Palacín, I., Martín, S. & Martino, A. (2007). The effect of melatonin on the reproductive performance of three sheep breeds in Spain. *Small Ruminant Research*, Vol.69, No.1-3, (May), pp. 10-16, 0921-4488
- Acritopoulou, S. & Haresign, W. (1980). Response of ewes to a single injection of an analog of PGF-2alpha given at different stages of the estrous-cycle. *Journal of Reproduction and Fertility*, Vol.58, No.1, pp. 219-223, 0022-4251
- Adalsteinsson, S. (1979). Independent effects of liveweight and body condition on fecundity and productivity of Icelandic ewes. *Animal Production*, Vol.28, No.FEB, pp. 13-23, 0003-3561
- Alabart, J.L., Folch, J., Ciudad, M.A., Fantova, E., Sevilla, E. & Quintin, F.J. (2002) Efecto de la edad de la oveja rasa aragonesa sobre la fertilidad en la inseminación artificial (I.A.) dentro del esquema de mejora de la UPRA-OVIARAGÓN S.C.L, Proceedings of: XXVII Congreso de la Sociedad Española de Ovinotecnia y Caprinotecnia, Valencia (España)
- Allison, A.J. & Kelly, R.W. (1978). Synchronization of estrus and fertility in sheep treated with progestagen-impregnated implants, and prostaglandins with or without intra-vaginal sponges and subcutaneous pregnant mare serum. *New Zealand Journal of Agricultural Research*, Vol.21, No.3, pp. 389-393, 0028-8233
- Althouse, G.C., Kuster, C.E., Clark, S.G. & Weisiger, R.M. (2000). Field investigations of bacterial contaminants and their effects on extended porcine semen. *Theriogenology* Vol. 53, pp. 1167-1176, 0093-691X
- Althouse, G.C. & Lu, K.G. (2005). Bacteriospermia in extended porcine semen. *Theriogenology* Vol. 63, pp. 573-584, 0093-691X
- Anel, L., Alvarez, M., Martinez-Pastor, F., Garcia-Macias, V., Anel, E. & De Paz, P. (2006). Improvement strategies in ovine artificial insemination. *Reproduction in Domestic Animals* 41 (SUPPL. 2), pp. 30-42, 0936-6768
- Anel, L., Kaabi, M., Abroug, B., Alvarez, M., Anel, E., Boixo, J.C., de la Fuente, L.F. & de Paz, P. (2005). Factors influencing the success of vaginal and laparoscopic artificial insemination in churra ewes: a field assay. *Theriogenology*, Vol.63, No.4, (Mar), pp. 1235-1247, 0093-691X
- Aravindan, G.R., Bjordahl, J., Jost, L.K. & Evenson, D.P. (1997). Susceptibility of human sperm to in situ DNA denaturation is strongly correlated with DNA strand breaks identified by single-cell electrophoresis. *Experimental Cell Research*, Vol.236, No.1, (Oct), pp. 231-237, 0014-4827
- Arendt, J., Symons, A.M., Laud, C.A. & Pryde, S.J. (1983). Melatonin can induce early onset of the breeding-season in ewes. *Journal of Endocrinology*, Vol.97, No.3, pp. 395-400, 0022-0795
- Armstrong, D.T. & Evans, G. (1984). Intrauterine insemination enhances fertility of frozen-semen in superovulated ewes. *Journal of Reproduction and Fertility*, Vol.71, No.1, pp. 89-94, 0022-4251
- Arrebola, F.A., Abecia, J.A., Forcada, F., García, A., Martín, R.A. & Mesa, O. (2009). Effects of annual rainfall and farm on lamb production after treatment with melatonin

- implants in Merino sheep: A 4-year study. *New Zealand Veterinary Journal*, Vol.57, No.3, pp. 141-145, 0048-0169
- Aurich C. & Spergser J. (2007). Influence of bacteria and gentamicin on cooled-stored stallion spermatozoa. *Theriogenology*, Vol. 67, pp. 912-918, 0093-691X
- Baril, G., Leboeuf, B. & Saumande, J. (1993). Synchronization of estrus in goats: The relationship between time of occurrence of estrus and fertility following artificial-insemination. *Theriogenology*, Vol.40, No.3, (Sep), pp. 621-628, 0093-691X
- Baril, G., Remy, B., Leboeuf, B., Beckers, J.F. & Saumande, J. (1996). Synchronization of estrus in goats: The relationship between eCG binding in plasma time of occurrence of estrus and fertility following artificial insemination. *Theriogenology*, Vol.45, No.8, (Jun), pp. 1553-1559, 0093-691X
- Beck, N.F.G., Davies, B. & Williams, S.P. (1993). Estrous synchronization in ewes. The effect of combining a prostaglandin analog with a 5-day progestagen treatment. . *Animal Production*, Vol.56, (Apr), pp. 207-210, 0003-3561
- Bielanski A. (2007). Disinfection procedures for controlling microorganisms in the semen and embryos of humans and farm animals. *Theriogenology*, Vol. 68, pp.1-22, 0093-691X
- Bittman, E.L., Dempsey, R.J. & Karsch, F.J. (1983). Pineal melatonin secretion drives the reproductive response to daylength in the ewe. *Endocrinology*, Vol.113, No.6, pp. 2276-2283, 0013-7227
- Bodin, L., Drion, P.V., Remy, B., Brice, G., Cognie, Y. & Beckers, J.F. (1997). Anti-PMSG antibody levels in sheep subjected annually to oestrus synchronisation. *Reproduction Nutrition Development*, Vol.37, No.6, (Nov-Dec), pp. 651-660, 0926-5287
- Bodin, L., Elsen, J.M., Hanocq, E., Francois, D., Lajous, D., Manfredi, E., Mialon, M.M., Boichard, D., Foulley, J.L., San-Cristobal-Gaudy, M., Teyssier, J., Thimonier, J. & Chemineau, P. (1999). Genetics of reproduction in ruminants. *Productions Animales*, Vol.12, No.2, (May), pp. 87-100, 0990-0632
- Boland, M.P., Gordon, I. & Kelleher, D.L. (1978). Effect of treatment by prostaglandin analog (ICI-80,996) or progestagen (SC-9880) on ovulation and fertilization in cycles ewes. *Journal of Agricultural Science*, Vol.91, No.DEC, pp. 727-730, 0021-8596
- Branca, A., Molle, G., Sitzia, M., Decandia, M. & Landau, S. (2000). Short-term dietary effects on reproductive wastage after induced ovulation and artificial insemination in primiparous lactating Sarda ewes (vol 58, pg 59, 2000). *Animal Reproduction Science*, Vol.63, No.1-2, (Oct), pp. 123-123, 0378-4320
- Brown, B.W. (1994). A review of nutritional influences on reproduction in boars, bulls and rams. *Reproduction Nutrition Development*, Vol.34, No.2, pp. 89-114, 0926-5287
- Bru, C., Fantova, E., Sevilla, E., Quintin, F.J. & Alabart, J.L. (1995) Resultados de inseminación artificial de las ovejas Rasa aragonesa de las ganaderías de Carne Aragón, S.C.L. Influencia de la condición corporal, Proceedings of: XX *Jornadas Científicas de la Sociedad Española de Ovinotecnia y Caprinotecnia.*, Madrid (España)
- Buckrell, B.C., Buschbeck, C., Gartley, C.J., Kroetsch, T., McCutcheon, W., Martin, J., Penner, W.K. & Walton, J.S. (1994). Further development of a transcervical technique for artificial insemination in sheep using previously frozen semen. *Theriogenology*, Vol.4, No.4, pp. 601-611, 0093-691X
- Casao, A., Cebrian, I., Asumpcao, M.E., Perez-Pe, R., Abecia, J.A., Forcada, F., Cebrian-Perez, J.A. & Muino-Blanco, T. (2010a). Seasonal variations of melatonin in ram

- seminal plasma are correlated to those of testosterone and antioxidant enzymes. *Reproductive Biology and Endocrinology*, Vol.8, (Jun), pp. 1477-7827, 1477-7827.
- Casao, A., Mendoza, N., Perez-Pe, R., Grasa, P., Abecia, J.A., Forcada, F., Cebrian-Perez, J.A. & Muino-Blanco, T. (2010b). Melatonin prevents capacitation and apoptotic-like changes of ram spermatozoa and increases fertility rate. *Journal of Pineal Research*, Vol.48, No.1, (Jan), pp. 39-46, 0742-3098
- Casao, A., Vega, S., Palacin, I., Perez-Pe, R., Lavina, A., Quintin, F.J., Sevilla, E., Abecia, J.A., Cebrian-Perez, J.A., Forcada, F. & Muino-Blanco, T. (2010c). Effects of Melatonin Implants During Non-Breeding Season on Sperm Motility and Reproductive Parameters in Rasa Aragonesa Rams. *Reproduction in Domestic Animals*, Vol.45, No.3, (Jun), pp. 425-432, 0936-6768
- Chemineau, P., Malpaux, B., Delgadillo, J.A., Guerin, Y., Ravault, J.P., Thimonier, J. & Pelletier, J. (1992). Control of sheep and goat reproduction - Use of light and melatonin. *Animal Reproduction Science*, Vol.30, No.1-3, (Nov), pp. 157-184, 0378-4320
- Chemineau, P., Vandaele, E., Brice, G. & Jardon, C. (1991). The use of melatonin in improving reproduction performance in sheep. *Recueil De Medecine Veterinaire*, Vol.167, No.3-4, pp. 227-239, 0034-1843
- Choudhry, T.M., Berger, T. & Dally, M. (1995). In vitro fertility evaluation of cryopreserved ram semen and its correlation with relative in vivo fertility. *Theriogenology*, Vol.43, No.7, (May), pp. 1195-1200, 0093-691X
- Clément, F., Vidament M. & Guérin B. (1995). Microbial contamination of stallion semen. *Biology of Reproduction Monographic*, Vol. 1, pp. 779-786.
- Cognié, Y., Bodin, L. & Terqui, M. (1984). The control of the time of ovulation in relation to the use of artificial insemination. In *Insémination artificielle et amélioration génétique: Bilan et perspectives critiques* ed. JM Elsen and JL Foulley. *Institut National de la Recherche Agronomique*, Paris, France, pp 77-95.
- Colas, G. & Brice, G. (1976). Seasonal variations of the fertilizing capacity of the deep frozen ram semen. In: *8th International Congress of Animal Reproduction Artificial Insemination*, Vol.4, pp.977-980.
- Colas, G. (1979). Fertility in the ewe after artificial insemination with fresh and frozen semen at the induced oestrous, and influence of the photoperiod on the semen quality of the ram. *Livestock Production Science*, Vol.6, No.2, pp. 153-166, 0301-6226
- Colas, G., Thimonier, J., Courot, M. & Ortavant, R. (1973). Fertility and reproductive efficiency during the breeding season of ewes artificially inseminated after treatment with flugestone acetate. *Annales De Zootechnie*, Vol.22, No.4, pp. 441-451, 0003-424X
- Contreras-Solis, I., Vasquez, B., Diaz, T., Letelier, C., Lopez-Sebastian, A. & Gonzalez-Bulnes, A. (2009). Efficiency of estrous synchronization in tropical sheep by combining short-interval cloprostenol-based protocols and "male effect". *Theriogenology*, Vol.71, No.6, (Apr), pp. 1018-1025, 0093-691X
- Corteel, J.M., Leboeuf, B. & Baril, G. (1988). Artificial breeding of adult goats and kids induced with hormones to ovulate outside the breeding season. *Small Ruminant Research*, Vol.1, No.1, pp. 19-35, 0921-4488
- David, I., Robert-Granie, C., Manfredi, E., Lagriffoul, G. & Bodin, L. (2008). Environmental and genetic variation factors of artificial insemination success in French dairy sheep. *Animal*, Vol.2, No.7, (Jul), pp. 979-986, 1751-7311

- den Daas, N. (1992). Laboratory assessment of semen characteristics. *Animal Reproduction Science*, Vol.28, No.1-4, pp. 87-94, 0378-4320
- Donovan, A., Hanrahan, J.P., Kummer, E., Duffy, P. & Boland, M.P. (2004). Fertility in the ewe following cervical insemination with fresh or frozen-thawed semen at a natural or synchronised oestrus. *Animal Reproduction Science*, Vol.84, No.3-4, (Sep), pp. 359-368, 0378-4320
- Dott, H.M., Harrison, R.A.P., & Foster, G.C.A. (1979). The maintenance of motility and the surface properties of epididymal spermatozoa from bull rabbit and ram in homologous seminal and epididymal plasma. *Journal of Reproduction Fertility*, Vol.55, pp. 113-124, 0022-4251
- Ducker, M.J. & Boyd, J.S. (1977). The effect of body size and body condition on the ovulation rate of ewes *Animal Production*, Vol.24, No.JUN, pp. 377-385, 0003-3561
- Eid, L.N., Lorton, S.P. & Parrish, J.J. (1994). Paternal influence on S-phase in the first cell cycle of the bovine embryo. *Biology of Reproduction*, Vol.51, No.6, (Dec), pp. 1232-1237, 0006-3363
- Eppleston, J. & Maxwell, W.C.M. (1993). Recent attempts to improve the fertility of frozen ram semen inseminated into the cervix. *Wool Sheep Breed*, Vol.41, pp 291-302
- Eppleston, J. & Maxwell, W.M.C. (1995). Sources of variation in the reproductive performance of ewes inseminated with frozen-thawed ram semen by laparoscopy. *Theriogenology*, Vol.43, No.4, (Mar), pp. 777-788, 0093-691X
- Eppleston, J., Evans, G. & Roberts, E.M. (1991). Effect of time of PMSG and GnRH on the time of ovulation, LH secretion and reproductive performance after intrauterine insemination with frozen ram semen. *Animal Reproduction Science*, Vol.26, No.3-4, (Dec), pp. 227-237, 0378-4320
- Eppleston, J., Maxwell, W.M.C. , Battye K.M. & Roberts, E.M. (1986). Effect of thawed motility and intrauterine motility of motile sperm on fertility of ewes. *Proceedings of Australian Society of Reproduction Biology*, Vol. 18, pp.19.
- Eppleston, J., Salamon, S., Moore, N.W. & Evans, G. (1994). The depth of cervical insemination and site of intrauterine insemination and their relationship to the fertility of frozen-thawed ram semen. *Animal Reproduction Science*, Vol.36, No.3-4, (Sep), pp. 211-225, 0378-4320
- Esmailizadeh, A.K., Dayani, O. & Mokhtari, M.S. (2009). Lambing season and fertility of fat-tailed ewes under an extensive production system are associated with liveweight and body condition around mating. *Animal Production Science*, Vol.49, No.12, pp. 1086-1092, 1836-5787
- Fantova, E., Bru, R., Sevilla, E., Quintin, F.J., Congost, S., Alabart, J.L. & Folch, J. (1998). Resultados de inseminación artificial en el marco del esquema de selección por prolificidad en las ganaderías de Carne Aragón. ITEA. 94 A (2), 93-100. ITEA, Vol.94, No.2, pp. 93-100,
- Fantova, E., Ciudad, M.A., Vijil, E., Sevilla, E., Quintin, F.J., Folch, J., Sin, E., Alabart, J.L. & Jurado, J.J. (1999) Evolution of the artificial insemination program in the genetic improvement Schedule of UPRA-Carne oviaragón. Study of flock location and insemination personal effects, Proceedings of: XXIV Jornadas Científicas de la Sociedad Española de Ovinotecnia y Caprinotecnia, Soria (España)

- Fernandez-Abella, D., Preve, M.O., & Villegas N. (2003). Insemination time and dilution rate of cooled and chilled ram semen affects fertility. *Theriogenology*, Vol. 60, pp.21-26, 0093-691X.
- Forcada, F., Abecia, J.A. & Sierra, I. (1992). Seasonal-changes in estrus activity and ovulation rate in Rasa-Aragonesa ewes maintained at 2 different body condition levels. *Small Ruminant Research*, Vol.8, No.4, (Sep), pp. 313-324, 0921-4488
- Forcada, F., Ait Amer-Meziane, M., Abecia, J.A., Maurel, M.C., Cebrián-Pérez, J.A., Muiño-Blanco, T., Asenjo, B., Vázquez, M.I. & Casao, A. (2011). Repeated superovulation using a simplified FSH/eCG treatment for in vivo embryo production in sheep. *Theriogenology*, Vol.75, No.4, pp. 769-776, 0093-691X
- Fukui, Y., Kohno, H., Okabe, K., Katsuki, S., Yoshizawa, M., Togari, T. & Watanabe, H. (2010). Factors affecting the fertility of ewes after intrauterine insemination with frozen-thawed semen during the non-breeding season. *Journal of Reproduction and Development*, Vol.56, No.4, (Aug), pp. 460-466, 0916-8818
- Fukui, Y., Kohno, H., Togari, T. & Hiwasa, M. (2007). Fertility of ewes inseminated intrauterinally with frozen semen using extender containing bovine serum albumin. *Journal of Reproduction and Development*, Vol.53, No.4, (Aug), pp. 959-962, 0916-8818
- Gabiña, D. & Folch, J. (1987) La inseminación artificial ovina. Resultados de su aplicación en un programa de selección en la raza Rasa Aragonesa, Proceedings of: ITEA, Zaragoza (España)
- Gonzalez, R.E., Labuonora, D. & Russel, A.J.F. (1997). The effects of ewe live weight and body condition score around mating on production from four sheep breeds in extensive grazing systems in Uruguay. *Animal Science*, Vol.64, (Feb), pp. 139-145, 1357-7298.
- Gonzalez, R.E., Labuonora, D., & Russel, A.J.F. (1997). The effects of ewe live weight and body condition score around mating on production from four sheep breeds in extensive grazing systems in Uruguay. *Animal Science*, Vol.64, No.1, pp. 139-145, 1357-7298.
- Gordon I. (1997). Controlled Reproduction in Sheep and Goats. *Cab International* (U.S.A.).
- Gosálvez, J., Fernández, J.L., Gosálbez, A., Arrollo, A., Agarwal, A. & López-Fernández A. (2007). Dynamics of sperm DNA fragmentation in mammalian species as assessed by the SCD methodology. *Fertility and Sterility*, Vol. 88, pp. 365-1365, 1556-5653.
- Gosálvez, J., Gosálbez, A., Arroyo, F., Fernández, J.L., & López-Fernández, C. (2008). Assessing sperm DNA fragmentation in the field: an adaptation of sperm chromatin dispersion technology. *Biotechnology and Histochemistry*, Vol.83, No.5, pp 247-52, 1473-7760.
- Greyling, J.P.C., Erasmus, J.A., Taylor, G.J. & van der Merwe, S. (1997). Synchronization of estrus in sheep using progestagen and inseminating with chilled semen during the breeding season. *Small Ruminant Research*, Vol.26, No.1-2, (Dec), pp. 137-143, 0921-4488
- Gunn, R.G. & Doney, J.M. (1979). Fertility in Cheviot ewes. 1. the effect of body condition at mating on ovulation and early embryo mortality in North and South Country Cheviot ewes. *Animal Production*, Vol.29, No.AUG, pp. 11-16, 0003-3561
- Gunn, R.G. (1983). The Influence of Nutrition on the Reproductive Performance of Ewes. In *Sheep production*, ed. Haresign, W., Butterworths, London.

- Gunn, R.G., & Doney, J.M. (1979). Fertility in Cheviot ewes. 1. The effect of body condition at mating on ovulation and early embryo mortality in North and South Country Cheviot ewes. *Animal Production*, Vol. 29, pp. 111-121, 0003-3561
- Hackett, A.J., Langford, G.A. & Robertson, H.A. (1981). Rertility of ewes after synchronization of estrus with prostaglandin-f2-alpha and artificial-insemination. *Theriogenology*, Vol.15, No.6, pp. 599-603, 0093-691X
- Haresign, W., Peters, A.R. & Staples, L.D. (1990). The effect of melatonin implants on breeding activity and litter size in commercial sheep flocks in the Uk. *Animal Production*, Vol.50, (Feb), pp. 111-121, 0003-3561
- Hawk, H.W. & Conley, H.H. (1975). Involvement of cervix in sperm transport failures in reproductive-tract of ewe. *Biology of Reproduction*, Vol.13, No.3, pp. 322-328, 0006-3363
- Hawk, H.W. (1971). Sperm destruction in sheep vagina. *Journal of Animal Science*, Vol.33, No.1, pp. 255-258, 0021-8812
- Heape, W. (1898). On the artificial insemination of mares. *Veterinarian*, Vol.71, pp. 202-212,
- Houghton, J.A.S., Liberati, N., Schrick, F.N., Townsend, E.C., Dailey, R.A. & Inskeep, E.K. (1995). day of estrous-cycle affects follicular dynamics after induced luteolysis in ewes. *Journal of Animal Science*, Vol.73, No.7, (Jul), pp. 2094-2101, 0021-8812
- Husein, M.Q. & Ababneh, M.M. (2008). A new strategy for superior reproductive performance of ewes bred out-of-season utilizing progestagen supplement prior to withdrawal of intravaginal pessaries. *Theriogenology*, Vol.69, No.3, (Feb), pp. 376-383, 0093-691X
- Jabbour, H.N. & Evans, G. (1991). Fertility of superovulated ewes following intrauterine or oviducal insemination with fresh or frozen-thawed semen. *Reproduction Fertility and Development*, Vol.3, No.1, pp. 1-7, 1031-3613
- Kaabi, M., Alvarez, M., Anel, E., Chamorro, C.A., Boixo, J.C., de Paz, P. & Anel, L. (2006). Influence of breed and age on morphometry and depth of inseminating catheter penetration in the ewe cervix: A postmortem study. *Theriogenology*, Vol.66, No.8, (Nov), pp. 1876-1883, 0093-691X
- Karsch, F.J., Bittman, E.L., Foster, D.L., Goodman, R.L., Legan, S.J. & Robinson, J.E. (1984). Neuroendocrine basis of seasonal reproduction. *Recent Progress in Hormone Research*, Vol.40, pp. 185-232, 0079-9963
- Kaya A., Aksoy M. & Tekeli, T. (2002). Influence of ejaculation frequency on sperm characteristics, ionic composition and enzymatic activity of seminal plasma in rams. *Small Ruminant Research*, Vol.44, pp 153-158, 0921-4488.
- Keisler, DH. & Buckrell, B.C. (1997). Chapter 88. Breeding strategies. In: *Current therapy in large animal theriogenology*. Edited by Robert s. Youngquist. 1st. edition. ISBN: 0-7216-5396-0.
- Killen, I.D. & Caffery, G.J. (1982). Uterine insemination of ewes with the aid of a laparoscope. *Australian Veterinary Journal*, Vol.59, No.3, pp. 95-95, 0005-0423
- King, M.E., McKelvey, W.A., Dingwall, W.S., Matthews, K.P., Gebbie, F.E., Mylne, M.J., Stewart, E. & Robinson, J.J. (2004). Lambing rates and litter sizes following intrauterine or cervical insemination of frozen/thawed semen with or without oxytocin administration. *Theriogenology*. Vol.62, pp.1236-44, 0093-691X
- Laliotis, V., Vosniakou, A., Zafrakas, A., Lymberopoulos, A. & Alifakiotis, T. (1998). The effect of melatonin on lambing and litter size in milking ewes after advancing the

- breeding season with progestagen and PMSG followed by artificial insemination. *Small Ruminant Research*, Vol.31, No.1, (Dec), pp. 79-81, 0921-4488.
- Landau, S. & Molle, G. (1997). Nutrition effects on fertility in small ruminants with an emphasis on Mediterranean sheep breeding systems. *Proc. FAOrCIHEAM Meeting on the Nutrition of Small Ruminants*, 24-26 October 1996, Rabat, Morocco.
- Langford, G.A. & Marcus, G.J. (1982). Influence of sperm number and seminal plasma on fertility of progestagen-treated sheep in confinement. *Journal of Reproduction Fertility*, Vol.65, pp.325-329, 0022-4251.
- Langford, G.A., Ainsworth, L. & Wolynetz, M.S. (1982). Reproductive response of progestogen-treated sheep in confinement to a single and double insemination. *Journal of Animal Science*, Vol.54, No.1, pp. 12-17, 0021-8812
- Lassoued, N., Rekik, M., Mahouachi, M. & Ben Hamouda, M. (2004). The effect of nutrition prior to and during mating on ovulation rate, reproductive wastage, and lambing rate in three sheep breeds. *Small Ruminant Research*, Vol.52, No.1-2, (Apr), pp. 117-125, 0921-4488
- Legaz, E., Deletang, F. & Martino, A. (2000). Efecto de la utilización de implantes de melatonina sobre los resultados de inseminación artificial en ovejas Manchega x Assaf. *Producción Ovina y Caprina*, Vol.SEOC Teruel, XXV, pp. 555-558,
- Light, J.E., Silvia, W.J. & Reid, R.C. (1994). Luteolytic effect of prostaglandin-f2-alpha and 2 metabolites in ewes. *Journal of Animal Science*, Vol.72, No.10, (Oct), pp. 2718-2721, 0021-8812
- López, A., Soderquist, L. & Rodriguez-Martinez, H. (1999). Sperm viability in ram semen diluted and stored in three different extenders. *Acta Veterinaria Scandinavica*, Vol.40, pp. 1-9, 0044-605X
- López-Fernández, C., Fernández, J.L., Gosálbez, A., Arroyo, F., Vázquez, J.M., Holt, W.V. & Gosálvez, J. (2008). Dynamics of sperm DNA fragmentation in domestic animals III. Ram. *Theriogenology*. Vol.70, pp.898-908, 0093-691X
- Lopez-Saez, A., Ortiz, N., Gallego, L. & Garde, J.J. (2000). Liquid storage (5 degrees C) of ram semen in different diluents. *Archives of Andrology*, Vol.44, No.2, (Mar-Apr), pp. 155-164, 0148-5016
- Lunstra, D.D. & Christenson, R.K. (1981). Fertilization and embryonic survival in ewes synchronized with exogenous hormones during the anestrus and estrus seasons. *Journal of Animal Science*, Vol.53, No.2, pp. 458-466, 0021-8812
- Marai, I.F.M., El-Darawany, A.A., Abou-Fandoud, E.I. & Abdel-Hafez, M.A.M. (2004). Reproductive traits and the physiological background of the seasonal variations in Egyptian suffolk ewes under the conditions of Egypt. *Annals of Arid Zone*, Vol. 43, No.2, (June), pp. 177-183
- Marai, I.F.M., El-Darawany, A.A., Fadiel, A. & Abdel-Hafez, M.A.M. (2007). Physiological traits as affected by heat stress in sheep-A review. *Small Ruminant Research* Vol. 71, No.1-3, (August), pp 1-12
- Marai, I.F.M., Bahgat, L.B., Shalaby, T.H. & Abdel-Hafez, M.A. (2000). Response of Male Lambs to Concentrate Mixtures given with or without Natural Clay under Egypt Conditions. *Annals of Arid Zone*, Vol.39, No.4, pp. 449-460
- Marai, I.F.M., El-Darawany, A.A., Abou-Fandoud, E.I. & Abdel-Hafez, M.A.M. (2006). Serum blood components during pre-oestrus, oestrus and pregnancy phases in Egyptian

- Suffolk as affected by heat stress, under the conditions of Egypt. *Egypt Journal of Sheep and Goats Desertic Animal Science*, Vol.1, No.1, pp. 47-62
- Martí, J.I., Martí, E., Cebrián-Pérez, E. & Muiño-Blanco, T. (2003). Survival rate and antioxidant enzyme activity of ram spermatozoa after dilution with different extenders or selection by a dextran swim-up procedure. *Theriogenology* Vol.60, pp. 1025-1037, 0093-691X
- Martin, G.B., Milton, J.T.B., Davidson, R.H., Hunzicker, G.E.B., Lindsay, D.R. & Blache, D. (2004). Natural methods for increasing reproductive efficiency in small ruminants. *Animal Reproduction Science*, Vol.82-3, (Jul), pp. 231-246, 0378-4320
- Maurel, M.C., Roy, F., Hervé, V., Bertin, J., Vaiman, D., Cribiu, E., Manfredi, E., Bouvier, F., Lantier, I., Boue, P. & Guillou, F. (2003). Réponse immunitaire à la eCG utilisée dans le traitement de l'induction d'ovulation chez la chèvre et la brebis. Immune response to equine Chorionic Gonadotropin used for the induction of ovulation in goats and ewes. *Gynécologie Obstétrique & Fertilité*, Vol.31, No.9, pp. 766-769, 1297-9589
- Maxwell, W.M.C. & Salamon S. (1993). Liquid storage of ram semen-a review. *Reproduction, Fertility and Development*, Vol. 5, pp. 613-638, 1031-3613.
- Maxwell, W.M.C. (1986). Artificial insemination of ewes with frozen-thawed semen at a synchronized oestrus. 2. Effect of dose of spermatozoa and site of intrauterine insemination on fertility. *Animal Reproduction Science*, Vol.10, No.4, (Apr), pp. 309-316, 0378-4320
- McCracke, J.A., Samuelss, B., Goding, J.R., Glew, M.E., Green, K., Carlson, J.C. & Baird, D.T. (1972). Prostaglandin-f2-alpha identified as a luteolytic hormone in sheep. *Nature-New Biology*, Vol.238, No.83, pp. 129-135, 0090-0028.
- Menchaca, A. & Rubianes, E. (2004). New treatments associated with timed artificial insemination in small ruminants. *Reproduction Fertility and Development*, Vol.16, No.4, pp. 403-413, 1031-3613
- Montoro, V. (1995). La inseminación artificial con semen refrigerado en el esquema de selección de la raza ovina manchega. *Tesis Doctoral*, pp. 148,
- Morley, F.H.W., White, D.H., Kenney, P.A. & Davis, I.F. (1978). Predicting ovulation rate from liveweight in ewes *Agricultural Systems*, Vol.3, No.1, pp. 27-45, 0308-521X
- Muiño-Blanco, T., Pérez-Pé, R. & Cebrián-Pérez, J.A. (2008). Seminal plasma proteins and sperm resistance to stress. *Reproduction in Domestic Animals*, Vol.4, pp.18-31, 1439-0531.
- O'Callaghan, D., Karsch, F.J., Boland, M.P. & Roche, J.F. (1991). Role of short days in timing the onset and duration of reproductive activity in ewes under artificial photoperiods. *Biology of Reproduction*, Vol.44, No.1, (January 1, 1991), pp. 23-28, 0006-3363.-3363
- Olivera-Muzante, J., Fierro, S., López, V. & Gil, J. (2011). Comparison of prostaglandin- and progesterone-based protocols for timed artificial insemination in sheep. *Theriogenology*, Vol.In Press, Corrected Proof, pp. 0093-691X
- Ollero, M., Muiño-Blanco, T., López-Pérez, M.J. & Cebrián-Pérez, J.A. (1996). Viability of ram spermatozoa in relation to the abstinence period and successive ejaculations. *International Journal of Andrology*, Vol.19, pp. 287-92, 1365-2605
- Ollero, M., Pascual, M.L., Muiño-Blanco, T., Cebrián-Pérez, J.A. & López-Pérez, M.J. (1994). Revealing surface changes associated with maturation of ram spermatozoa by


- centrifugal counter-current distribution in an aqueous two-phase system. *Journal of Chromatography*, Vol. 668, pp.173-178, 0021-9673.
- Ostermeier, G.C., Sargeant, G.A., Yandell, B.S., Evenson, D.P. & Parrish, J.J. (2001). Relationship of bull fertility to sperm nuclear shape. *Journal of Andrology*, Vol.22, No.4, (Jul-Aug), pp. 595-603, 0196-3635
- Palacin, I., Abecia, J.A., Forcada, F., Casao, A., Cebrian, J.A., Muino, T., Palacios, C. & Pontes, J.M. (2008). Effects of exogenous melatonin treatment on out-of-season ram fertility. *Italian Journal of Animal Science*, Vol.7, No.2, pp. 199-206, 1594-4077
- Palacín, I., Santolaria, P., Fantova, E., Blasco, M.E., Quintín-Casorrán, F.J., Sevilla-Mur, E., & Yániz, J.L. (2011). Efecto de la localización geográfica sobre los resultados de fertilidad en el programa de inseminación artificial de la unión de productores de raza aragonesa (UPRA). *ITEA*, en prensa
- Papadopoulos, S., Hanrahan, J.P., Donovan, A., Duffy, P., Boland, M.P. & Lonergan, P. (2005). In vitro fertilization as a predictor of fertility from cervical insemination of sheep. *Theriogenology*, Vol.63, No.1, (Jan), pp. 150-159, 0093-691X
- Pascual, M.L., Muino-Blanco, T., Cebrian-Perez, J.A., & Lopez-Perez, M.J. (1993). Sperm cell heterogeneity revealed by centrifugal counter-current distribution in an aqueous two-phase system. *Journal of Chromatography - Biomedical Applications* Vol. 617, No.1, pp. 51-57, 1570-0232.
- Paulenz, H., Söderquist, L., Adnøy, T., Fossen, O.H., & Berg, K.A. (2003). Effect of milk- and TRIS-based extenders on the fertility of sheep inseminated vaginally once or twice with liquid semen. *Theriogenology*, Vol.60, pp. 759-766, 0093-691X
- Paulenz, H., Adnøy, T. & Soderquist, L. (2007). Comparison of fertility results after vaginal insemination using different thawing procedures and packages for frozen ram semen. *Acta Veterinaria Scandinavica*, Vol.49, (Sep), pp. 0044-605X
- Paulenz, H., Adnøy, T., Fossen, O.H., Soderquist, L. & Berg, K.A. (2002). Effect of deposition site and sperm number on the fertility of sheep inseminated with liquid semen. *Veterinary Record*, Vol.150, No.10, (Mar), pp. 299-+, 0042-4900
- Paulenz, H., Soderquist, L., Adnøy, T., Soltun, K., Saether, P.A., Fjellsoy, K.R. & Berg, K.A. (2005). Effect of cervical and vaginal insemination with liquid semen stored at room temperature on fertility of goats. *Animal Reproduction Science*, Vol.86, No.1-2, (Mar), pp. 109-117, 0378-4320
- Pearce, D.T. & Robinson, T.J. (1985). Plasma progesterone concentrations, ovarian and endocrinological responses and sperm transport in ewes with synchronized estrus. *Journal of Reproduction and Fertility*, Vol.75, No.1, pp. 49-62, 0022-4251
- Pelletier, J., Chemineau, P., Thimonier, J. & Vollandnail, P. (1987). Environment and changes in reproductive activity in sheep and goat. *Ed. P.Pevet. Karger Basel. Proc. 8th ESPC Conf. Strasbourg*, Vol.3, pp. 121-135,
- Quinliva, T.D. & Robinson, T.J. (1969). Numbers of spermatozoa in genital tract after artificial insemination of progestagen-treated ewes. *Journal of Reproduction and Fertility*, Vol.19, No.1, pp. 73-&, 0022-4251
- Rhind, S.M., McMillen, S., McKelvey, W.A.C., Rodriguezherrejon, F.F. & McNeilly, A.S. (1989). Effect of the body condition of ewes on the secretion of LH and FSH and the pituitary response to gonadotrophin-releasing hormone. *Journal of Endocrinology*, Vol.120, No.3, (Mar), pp. 497-502, 0022-0795

- Robinson, T.J., Moore, N.W., Lindsay, D.R., Fletcher, I.C. & Salamon, S. (1970). Fertility following synchronization of oestrus in sheep with intravaginal sponges .1. effects of vaginal douche, supplementary steroids, time of insemination, and numbers and dilution of spermatozoa. *Australian Journal of Agricultural Research*, Vol.21, No.5, pp. 767-&, 0004-9409
- Rodríguez-Martínez, H. (2003). Laboratory Semen Assessment and Prediction of Fertility: still Utopia?. *Reproduction in Domestic Animals* Vol.38, pp. 312-318, 0936-6768.
- Romano, J.E. (2004). Synchronization of estrus using CIDR. FGA or MAP intravaginal pessaries during the breeding season in Nubian goats. *Small Ruminant Research*, Vol.55, pp.15-19, 0921-4488
- Roy, F., Combes, B., Vaiman, D., Cuibiu, E.P., Pobel, T., Deletang, F., Combarnous, Y., Guillou, F. & Maurel, M.C. (1999). Humoral immune response to equine chorionic gonadotropin in ewes: Association with major histocompatibility complex and interference with subsequent fertility. *Biology of Reproduction*, Vol.61, No.1, (Jul), pp. 209-218, 0006-3363
- Saacke, R.G., Bame, J., Karabinus, D.S., Mulins, J. & Whitman, S. (1988). Transport of abnormal sperm in the artificially inseminated cow based upon accessory sperm in the zona pellucida, Proceedings of: *Proceedings of the 11th International Congress on Animal Reproduction and Artificial Insemination*. Dublin, Ireland, Dublin, Ireland
- Saacke, R.G., Nadir, S., Dalton, J., Bame, J., DeJarnette, J.M., Defelos, S. & Nebel, R.L. (1994). Accessory sperm evaluation and bull fertility—an update. , Proceedings of: *Proceedings of the 15th NAAB Technical Conference on Animal Reproduction and Artificial Insemination*, Milwaukee, Wisc
- Salamon, S., & Maxwell, W.M.C. (2000). Storage of ram semen. *Animal Reproduction Science*, Vol.62, pp. 77-111, 0378-4320
- Salamon, S. & Maxwell, W.M.C. (1995). Frozen storage of ram semen I. Processing, freezing, thawing and fertility after cervical insemination. *Animal Reproduction Science*, Vol.37, No.3-4, (Feb), pp. 185-249, 0378-4320
- Scaramuzzi, R.J., Downing, J.A., Campbell, B.K. & Cownie, Y. (1988). Control of fertility and fecundity of sheep by means of hormonal manipulation. *Australian Journal of Biological Sciences*, Vol.41, No.1, pp. 37-45, 0004-9417
- Selaive-Villarroel, A.B. & Kennedy, J.P. (1983a). Fertility studies in young and mature merino ewes. 1. Cervical mucus production. *Theriogenology*, Vol.20, No.5, pp. 537-541, 0093-691X
- Selaive-Villarroel, A.B. & Kennedy, J.P. (1983b). Fertility studies in young and mature merino ewes. 2. Sperm transport. *Theriogenology*, Vol.20, No.5, pp. 543-547, 0093-691X
- Shackell GH, Kyle B, Littlejohn RP. Factors influencing the success of a large scale artificial insemination programme in sheep. *Proc. New Zeal Soc Anim Prod* 1990;50:427-430
- Shelton, M. (2000). Reproductive performance of sheep exposed to hot environments. Malik R.C., Razzaque M.A., and Al-Nasser A.Y. (Eds), Published by the Kuwait Institute for Scientific Research. *Sheep Production in Hot and Arid Zones*, pp. 155-162.
- Sirois, J. & Fortune, J.E. (1990). Lengthening the bovine estrous-cycle with low-levels of exogenous progesterone - a model for studying ovarian follicular dominance. *Endocrinology*, Vol.127, No.2, (Aug), pp. 916-925, 0013-7227

- Soderquist, L., Lundeheim, N. & Nilsson, B. (1999). Assessment of fertility after using different procedures to thaw ram spermatozoa frozen in mini straws. *Reproduction in Domestic Animals*, Vol.34, No.2, (May), pp. 61-66, 0936-6768
- Thibier, M. & Guerin, B. (2000). Hygienic aspects of storage and use of semen for artificial insemination. *Animal Reproduction Science*, Vol.62, pp. 233-251, 0378-4320
- Thiéry, J.C., Chemineau, P., Hernandez, X., Migaud, M. & Malpaux, B. (2002). Neuroendocrine interactions and seasonality. *Domestic Animal Endocrinology*, Vol.23, No.1-2, pp 87-100
- Thomson, E.F. & Bahhady, F.A. (1988). A note on the effect of live weight at mating on fertility of Awassi ewes in semi-arid northwest Syria. *Animal Production*, Vol.47, (Dec), pp. 505-508, 0003-3561
- Ungerfeld, R. & Rubianes, E. (1999). Effectiveness of short-term progestogen primings for the induction of fertile oestrus with eCG in ewes during late seasonal anoestrus. *Animal Science*, Vol.68, (Apr), pp. 349-353, 1357-7298
- Ustuner, B., Gunay, U., Nur, Z. & Ustuner, H. (2007). Effects of long and short-term progestagen treatments combined with PMSG on oestrus synchronization and fertility in awassi ewes during the breeding season. *Acta Veterinaria Brno*, Vol.76, No.3, (Sep), pp. 391-397, 0001-7213
- Vázquez, M.I., Abecia, J.A., Forcada, F. & Casao, A. (2010). Effects of exogenous melatonin on in vivo embryo viability and oocyte competence of undernourished ewes after weaning during the seasonal anestrus. *Theriogenology*, Vol. 74, pp. 618-26, 0093-691X
- Verstegen, J., Iguar-Ouada, M. & Onclin, K. (2002). Computer assisted semen analyzers in andrology research and veterinary practice. *Theriogenology*. Vol.57, pp. 49-79, 0093-691X
- Vinoles, C., Forsberg, M., Banchero, G. & Rubianes, E. (2001). Effect of long-term and short-term progestagen treatment on follicular development and pregnancy rate in cyclic ewes. *Theriogenology*, Vol.55, No.4, (Mar), pp. 993-1004, 0093-691X
- Vinoles, C., Meikle, A., Forsberg, M. & Rubianes, E. (1999). The effect of subluteal levels of exogenous progesterone on follicular dynamics and endocrine patterns during the early luteal phase of the ewe. *Theriogenology*, Vol.51, No.7, (May), pp. 1351-1361, 0093-691X
- Windsor, D.P. (1995). Factors influencing the success of transcervical insemination in merino ewes. *Theriogenology*, Vol.43, No.6, (Apr), pp. 1009-1018, 0093-691X
- Windsor, D.P. (1997). Variation between ejaculates in the fertility of frozen ram semen used for cervical insemination of Merino ewes. *Animal Reproduction Science*, Vol.47, No.1-2, (May), pp. 21-29, 0378-4320
- Xu, Z., McDonald, M.F. & McCutcheon, S.N. (1989). The effects of nutritionally-induced liveweight differences on follicular development, ovulation rate, oestrus activity and plasma follicle-stimulating hormone levels in the ewe. *Animal Reproduction Science*, Vol.19, pp 67-78, 0378-4320
- Yániz, J., Marti, J.I., Silvestre, M.A., Folch, J., Santolaria, P., Alabart, J.L. & Lopez-Gatius, F. (2005). Effects of solid storage of sheep spermatozoa at 15 degrees C on their survival and penetrating capacity. *Theriogenology*, Vol 64, pp. 1844-1851, 0093-691X
- Yániz, J.L., Santolaria, P., Marco-Aguado, M.A. & López-Gatius, F. (2008). Use of image analysis to assess the plasma membrane integrity of ram spermatozoa in different diluents. *Theriogenology* Vol.15, No.70(2), pp.192-8, 0093-691X

- Yaniz, J.L., Marco-Aguado, M.A., Mateos, J.A. & Santolaria, P. (2010). Bacterial contamination of ram semen, antibiotic sensitivities, and effects on sperm quality during storage at 15 degrees C. *Animal Reproduction Science*, Vol.122, No.1-2, (Oct), pp. 142-149, 0378-4320
- Yaniz, J.L., Mateos, J.A. & Santolaria, P. (2011). Zwitterionic buffers preserve ram semen quality more efficiently than TRIS during storage at 15 degrees C. *Small Ruminant Research*, Vol.95, No.1, (Jan), pp. 54-60, 0921-4488
- Yeates, N.T.M. (1949). The breeding season of the sheep with particular reference to its modification by artificial means using light. *Journal of Agricultural Science*, Vol.39, No.1, pp. 1-&, 0021-8596
- Younis, A.A., Alkamali, A.A. & Eltawil, E.A. (1978). Effect of flushing on fertility of Awassi and Hamdani ewes. *World Review of Animal Production*, Vol.14, No.2, pp. 41-48, 0043-8979
- Yue, D., Yan, L., Luo, H., Xu, X. & Jin, X. (2010). Effect of Vitamin E supplementation on semen quality and the testicular cell membranal and mitochondrial antioxidant abilities in Aohan fine-wool sheep. *Animal Reproduction Science*, Vol.118, No.2-4, pp. 217-222, 0378-4320
- Zelege, M., Greyling, J.P.C., Schwalbach, L.M.J., Muller, T. & Erasmus, J.A. (2005). Effect of progestagen and PMSG on oestrous synchronization and fertility in Dorper ewes during the transition period. *Small Ruminant Research*, Vol.56, No.1-3, (Jan), pp. 47-53, 0921-4488

IntechOpen



Artificial Insemination in Farm Animals

Edited by Dr. Milad Manafi

ISBN 978-953-307-312-5

Hard cover, 300 pages

Publisher InTech

Published online 21, June, 2011

Published in print edition June, 2011

Artificial insemination is used instead of natural mating for reproduction purposes and its chief priority is that the desirable characteristics of a bull or other male livestock animal can be passed on more quickly and to more progeny than if that animal is mated with females in a natural fashion. This book contains under one cover 16 chapters of concise, up-to-date information on artificial insemination in buffalos, ewes, pigs, swine, sheep, goats, pigs and dogs. Cryopreservation effect on sperm quality and fertility, new method and diagnostic test in semen analysis, management factors affecting fertility after cervical insemination, factors of non-infectious nature affecting the fertility, fatty acids effects on reproductive performance of ruminants, particularities of bovine artificial insemination, sperm preparation techniques and reproductive endocrinology diseases are described. This book will explain the advantages and disadvantages of using AI, the various methodologies used in different species, and how AI can be used to improve reproductive efficiency in farm animals.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Pilar Santolaria, Inmaculada Palacin and Jesus Yaniz (2011). Management Factors Affecting Fertility in Sheep, Artificial Insemination in Farm Animals, Dr. Milad Manafi (Ed.), ISBN: 978-953-307-312-5, InTech, Available from: <http://www.intechopen.com/books/artificial-insemination-in-farm-animals/management-factors-affecting-fertility-in-sheep>

INTech
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2011 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike-3.0 License](https://creativecommons.org/licenses/by-nc-sa/3.0/), which permits use, distribution and reproduction for non-commercial purposes, provided the original is properly cited and derivative works building on this content are distributed under the same license.

IntechOpen

IntechOpen