We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Effects of Impurities on a Noncentrosymmetric Superconductor - Application to CePt₃Si

Heshmatollah Yavari
University of Isfahan
Iran

1. Introduction

In the past two decades, a number of novel superconducting materials have been discovered where order parameter symmetries are different from an *s*-wave spin singlet, predicted by the Bardeen-Cooper-Schrieffer (BCS) theory of electron-phonon mediated pairing. From the initial discoveries of unconventional superconductivity in heavy-fermion compounds, the list of examples has now grown to include the high- T_c cuprate superconductors, ruthenates, ferromagnetic superconductors, and possibly organic materials.

In most of these materials, there are strong indications that the pairing is caused by the electron correlations, in contrast to conventional superconductors such as Pb, Nb, etc. Nonphononic mechanisms of pairing are believed to favor a nontrivial spin structure and orbital symmetry of the Cooper pairs. For example, the order parameter in the high- T_c superconductors, where the pairing is thought to be caused by the antiferromagnetic correlations, has the *d*-wave symmetry with lines of zeroes at the Fermi surface. A powerful tool of studying unconventional superconducting states is symmetry analysis, which works even if the pairing mechanism is not known.

In general, the superconducting BCS ground state is formed by Cooper pairs with zero total angular momentum. The electronic states are four-fold degenerate $|k \downarrow\rangle$, and $|-k \uparrow\rangle$ have the same energy $\varepsilon(k)$. The states with opposite momenta and opposite spins are transformed to one another under time reversal operation $K|k \uparrow\rangle = |-k \downarrow\rangle$ and states with opposite momenta are transformed to one another under inversion operation $I|k \uparrow\rangle = |-k \uparrow\rangle$. The four degenerate states are a consequence of space and time inversion symmetries. Parity symmetry is irrelevant for spin-singlet pairing, but is essential for spin-triplet pairing. Time reversal symmetry is required for spin-singlet configuration, but is unimportant for spin-triplet state (Anderson, 1959, 1984).

If this degeneracy is lifted, for example, by a magnetic field or magnetic impurities coupling to the electron spins, then superconductivity is weakened or even suppressed. For spin-triplet pairing, Anderson noticed that additionally inversion symmetry is required to obtain the necessary degenerate electron states. Consequently, it became a widespread view that a material lacking an inversion center would be an unlikely candidate for spin-triplet pairing. For example, the absence of superconductivity in the paramagnetic phase of MnSi close to the quantum critical point to itinerant ferromagnetism was interpreted from this point of

view (Mathur, 1998; Saxena, 2000). Near this quantum critical point the most natural spin fluctuation mediated Cooper pairing would occur in the spin-triplet channel. However, MnSi has the so-called *B20* structure ($P2_1$), without an inversion center, inhibiting spin-triplet pairing.

Unusual properties are expected in superconductors whose crystal structure does not possess an inversion center (Edelstein, 1995; Frigeri et al., 2004; Gor'kov & Rashba, 2001; Samokhin et al., 2004; Sergienko & Curnoe, 2004).

Recent discovery of heavy fermion superconductor CePt₃Si has opened up a new field of the study of superconductivity (Bauer et al., 2004). This is because this material does not have inversion center, which has stimulated further studies (Akazawa et al., 2004; Yogi et al., 2005). Because of the broken inversion symmetry, Rashba-type spin-orbit coupling (RSOC) is induced (Edelstein, 1995; Rashba, 1960; Rashba & Bychkov, 1984), and hence different parities, spin-singlet pairing and spin triplet pairing, can be mixed in a superconducting state (Gor'kov & Rashba, 2001).

From a lot of experimental and theoretical studies, it is believed that the most possible candidate of superconducting state in CePt₃Si is *s+p*-wave pairing (Frigeri et al., 2004; Hayashi et al., 2006). This mixing of the pairing channels with different parity may result in unusual properties of experimentally observed quantities such as a very high upper critical field H_{c2} which exceeds the paramagnetic limit (Bauer et al., 2004; Bauer et al., 2005a, 2005b; Yasuda et al., 2004), and the simultaneous appearance of a coherence peak feature in the NMR relaxation rate T_1^{-1} and low-temperature power-law behavior suggesting line nodes in the quasiparticle gap (Bauer et al., 2005a, 2005b; Yogi et al., 2004). The presence of line nodes in the gap of CePt₃Si is also indicated by measurements of the thermal conductivity (Izawa et al., 2005) and the London penetration depth (Bauer et al., 2005; Bonalde et al., 2005).

It is known that the nonmagnetic as well as the magnetic impurities in the conventional and unconventional superconductors already have been proven to be a useful tool in distinguishing between various symmetries of the superconducting state (Blatsky et al., 2006). For example, in the conventional isotropic *s*-wave superconductor, the single magnetic impurity induced resonance state is located at the gap edge, which is known as Yu-Shiba-Rusinov state (Shiba, 1968). In the case of unconventional superconductor with $d_{x^2-y^2}$ -wave symmetry of the superconducting state, the nonmagnetic impurity-induced bound state appears near the Fermi energy as a hallmark of $d_{x^2-y^2}$ -wave pairing symmetry (Salkalo et al., 1996). The origin of this difference is understood as being due to the nodal structure of two kinds of SC order: in the $d_{x^2-y^2}$ -wave case, the phase of Cooper pairing wave function changes sign across the nodal line, which yields finite density of states (DOS) below the superconducting gap, while in the isotropic *s*-wave case, the density of states is gapped up to energies of about Δ_0 and thus the bound state can appear only at the gap edge. In principle the formation of the impurity resonance states can also occur in unconventional superconductors if the nodal line or point does not exist at the Fermi surface of a superconductor, as it occurs for isotropic nodeless *p*-wave and/or $d_x + id_y$ -wave superconductors for the large value of the potential strength (Wang Q.H. & Wang Z.D, 2004).

In unconventional superconductors non-magnetic impurities act as pair-breakers, similar to magnetic impurities in s-wave superconductors. A bound state appears near an isolated non-magnetic strong (scattering phase shift $\frac{\pi}{2}$, or unitarity) scatterer, at the energy close to the Fermi level. The broadening of this bound state to an impurity band at finite disorder leads to a finite density of states at zero energy, $N(0)$, that increases with increasing impurity concentration (Borokowski & Hirschfeld, 1994). The impurity scattering changes the temperature dependence of the physical quantities below T corresponding to the impurity bandwidth: $\Delta\lambda$ changes the behavior from T to T^2 the NMR relaxation rate changes from T^3 to T , and specific heat $C(T)$ changes from T^2 to T . In other words, the impurities modify the power laws, especially at low temperatures.

The problem of a magnetic impurity in a superconductor has been extensively studied, but is not completely solved because of the difficulty of treating the dynamical correlations of the coupled impurity-conduction electron system together with pair correlations. Generally, the behavior of the system can be characterized by the ratio of the Kondo energy scale in the normal metal to the superconducting transition temperature $\frac{T_K}{T_c}$. For $\frac{T_K}{T_c} \ll 1$, conduction

electrons scatter from classical spins and physics in this regime can be described by the Abrikosov-Gor'kov theory (Abrikosov & Gor'kov, 1961). In the opposite limit, $\frac{T_K}{T_c} \gg 1$, the

impurity spin is screened and conduction electrons undergo only potential scattering. In this regime s-wave superconductors are largely unaffected by the presence of Kondo impurities due to Anderson's theorem. Superconductors with an anisotropic order parameter, e.g. p-wave, d-wave etc., are strongly affected, however and the potential scattering is pair-breaking. The effect of pair breaking is maximal in s-wave superconductors in the intermediate region, $T_K \sim T_c$, while in the anisotropic case it is largest for $\frac{T_K}{T_c} \rightarrow \infty$

(Borkowski & Hirschfeld, 1992).

In the noncentrosymmetric superconductor with the possible coexistence of s-wave and p-wave pairing symmetries, it is very interesting to see what the nature of the impurity state is and whether a low energy resonance state can still occur around the impurity through changing the dominant role played by each of the pairing components. Previously, the effect of nonmagnetic impurity scattering has been studied in the noncentrosymmetric superconductors with respect to the suppression of T_c and the behavior of the upper critical field (Frigeri et al., 2004; Mineev & Samokhin, 2007).

This in turn stimulates me to continue studying more properties. My main goal in this chapter is to find how the superconducting critical temperature, magnetic penetration depth, and spin-lattice relaxation rate of a noncentrosymmetric superconductor depend on the magnetic and nonmagnetic impurity concentration and also discuss the application of our results to a model of superconductivity in CePt₃Si. I do these by using the Green's function method when both s-wave and p-wave Cooper pairings coexist.

The chapter is organized as follows. In Sect. 2, the disorder averaged Green's functions in the superconducting states are calculated and the effect of impurity is treated via the self-

energies of the system. In Sect. 3, the equations for the superconducting gap functions renormalized by impurities are used to find the critical temperature T_c .

In Sect. 4, by using linear response theory I calculate the appropriate correlation function to evaluate the magnetic penetration depth. In this system the low temperature behavior of the magnetic penetration depth is consistent with the presence of line nodes in the energy gap.

In Sect. 5, the spin-lattice relaxation rate of nuclear magnetic resonance (NMR) in a superconductor without inversion symmetry in the presence of impurity effect is investigated.

In the last two cases I assume that the superconductivity in CePt₃Si is most likely unconventional and our aim is to show how the low temperature power law is affected by nonmagnetic impurities.

Finally sect. 6 contains the discussion and conclusion remarks of my results.

2. Impurity scattering in normal and superconducting state

By using a single band model with electron band energy ξ_k measured from the Fermi energy where electrons with momentum k and spin s are created (annihilated) by operators $C_{k,s}^\dagger$ ($C_{k,s}$), the Hamiltonian including the pairing interaction can be written as

$$H = \sum_{k,s} \xi_k C_{k,s}^\dagger C_{k,s} + \frac{1}{2} \sum_{k,k'} \sum_{s,s'} V_{k,k'} C_{k,s}^\dagger C_{-k,s'}^\dagger C_{-k',s'} C_{k',s} \quad (1)$$

This system possesses time reversal and inversion symmetry ($\xi_k = \xi_{-k}$) and the pairing interaction does not depend on the spin and favors either even parity (spin-singlet) or odd parity (spin-triplet) pairing as required. The absence of inversion symmetry is incorporated through the antisymmetric Rashba-type spin-orbit coupling

$$H_{so} = \sum_{k,s,s'} \alpha \vec{g}_k \cdot \vec{\sigma}_{s,s'} C_{ks}^\dagger C_{ks'} \quad (2)$$

which removes parity but conserves time-reversal symmetry, i.e., $I H_{so} I^{-1} = -H_{so}$ and $T H_{so} T^{-1} = H_{so}$. In Eq. (2), σ denotes the Pauli matrices (this satisfies the above condition $I \sigma I^{-1} = -\sigma$ and $T \sigma T^{-1} = \sigma$), g_k is a dimensionless vector [$g_k = -g_{-k}$ to preserve time reversal symmetry], and $\alpha (> 0)$ denotes the strength of the spin-orbit coupling. The antisymmetric spin-orbit coupling (ASOC) term $\alpha \vec{g}_k \cdot \vec{\sigma}$ is different from zero only for crystals without an inversion center and can be derived microscopically by considering the relativistic corrections to the interaction of the electrons with the ionic potential (Frigeri et al., 2004; Dresselhaus, 1995). For qualitative studies, it is sufficient to deduce the structure of the g -vector from symmetry arguments (Frigeri et al., 2004) and to treat α as a parameter. I set $\langle g_k^2 \rangle_k = 1$, where $\langle \dots \rangle$ denotes the average over the Fermi surface. The ASOC term lifts the spin degeneracy by generating two bands with different spin structure.

In the normal state the eigenvalues of the total Hamiltonian ($H + H_{so}$) are

$$\xi_{k\pm} = \varepsilon_k - \mu \mp \alpha |g_k| \quad (3)$$

where $\varepsilon_k = \frac{k^2}{2m}$ and μ is the chemical potential.

It is obvious from here that the time reversal symmetry is lost and the shape of the Fermi surfaces does not obey the mirror symmetry.

Due to the big difference between the Fermi momenta we neglected the pairing of electronic states from different bands. The structure of theory is now very similar to the theory of ferromagnetic superconductors with triplet pairing (Mineev, 2004).

Effects of disorder are described by potential scattering of the quasiparticles, which in real-space representation is given by

$$H_{imp} = \sum_i \int \psi_s^\dagger(\vec{r}) U_{imp} \psi_s(\vec{r}) d\vec{r} \quad (4)$$

where $U_{imp} = U_n + U_m$, U_n is the potential of a non-magnetic impurity, which we consider rather short-ranged such that s-wave scattering is dominant and $U_m = J(\vec{r}) \vec{S} \cdot \vec{\sigma}$ is the potential interaction between the local spin on the impurity site and conduction electrons, here J is the exchange coupling and S is the spin operator.

2.1 Impurity averaging in superconducting state

Let us calculate the impurity-averaged Green's functions in the superconducting state. The Gor'kov equations with self-energy contributions are formally analogous to those obtained for system with inversion symmetry (Abrikosov et al., 1975).

$$(i\omega_n - \xi_{k\pm} - \tilde{\Sigma}_G(i\omega_n)) \tilde{\mathcal{G}}_\pm(k, i\omega_n) + (\Delta_k + \tilde{\Sigma}_F(i\omega_n)) \tilde{F}_\pm^\dagger(k, \omega_n) = \hat{\sigma}_0 \quad (5)$$

$$(i\omega_n + \xi_{-k\pm} + \tilde{\Sigma}_G^\dagger(i\omega_n)) \tilde{F}_\pm^\dagger(k, \omega_n) + (\Delta_k^\dagger + \tilde{\Sigma}_F^\dagger(i\omega_n)) \tilde{\mathcal{G}}_\pm(k, \omega_n) = 0 \quad (6)$$

where $\omega_n = (2n+1)\pi T$ are the Matsubara Fermionic frequencies, $\hat{\sigma}_0$ is the unit matrix in the spin state, and the impurity scattering enters the self-energy of the Green's function of the normal, $\tilde{\Sigma}_G$, and the anomalous type, $\tilde{\Sigma}_F$, their mathematical expressions read

$$\tilde{\Sigma}_G(i\omega_n) = (n_n |U_n|^2 + n_m |U_m|^2) \int \frac{d\vec{k}'}{(2\pi)^3} \tilde{\mathcal{G}}(\vec{k}', i\omega_n) \quad (7)$$

$$\tilde{\Sigma}_F(i\omega_n) = (n_n |U_n|^2 + n_m |U_m|^2) \int \frac{d\vec{k}'}{(2\pi)^3} \tilde{F}(\vec{k}', i\omega_n) \quad (8)$$

here n_n and n_m are the concentrations of nonmagnetic and magnetic impurities, respectively.

The equations for each band are only coupled through the order parameters given by the self-consistency equations

$$\Delta_{k\pm} = -T \sum_{k', \nu, n} V_{\pm\nu}(\vec{k}, \vec{k}') F_\nu(\vec{k}', \omega_n) \quad (9)$$

where $\nu = \pm$.

Solving the Gor'kov equations one obtains the following expressions for the disorder-averaged Green's functions

$$\tilde{\mathfrak{S}}_{\pm}(\vec{k}, \omega_n) = \begin{pmatrix} \mathfrak{S}_{\pm}(\vec{k}, i\omega_n) & F_{\pm}(\vec{k}, i\omega_n) \\ F_{\pm}^{\dagger}(\vec{k}, i\omega_n) & -\mathfrak{S}_{\pm}(-\vec{k}, -i\omega_n) \end{pmatrix} \quad (10)$$

where

$$\mathfrak{S}_{\pm}(\vec{k}, \omega_n) = \frac{i\omega_n + \xi_{-k\pm}}{(i\omega_n - \Sigma_{imp} + \vec{v}_s \cdot \vec{k}_F - \xi_{k\pm})(i\omega_n - \Sigma_{imp} + \vec{v}_s \cdot \vec{k}_F + \xi_{-k\pm}) - \Delta_{k\pm} \Delta_{k\pm}^{\dagger}} \quad (11)$$

$$F_{\pm}(\vec{k}, \omega_n) = \frac{-\Delta_{k\pm}}{(i\omega_n - \Sigma_{imp} + \vec{v}_s \cdot \vec{k}_F - \xi_{k\pm})(i\omega_n - \Sigma_{imp} + \vec{v}_s \cdot \vec{k}_F + \xi_{-k\pm}) - \Delta_{k\pm} \Delta_{k\pm}^{\dagger}} \quad (12)$$

here $\Sigma_{imp} = \Sigma_{imp(n)} + \Sigma_{imp(m)}$ is the self energy due to non magnetic and magnetic impurities. The energies of elementary excitations are given by

$$E_{k\pm} = \frac{\xi_{k\pm} - \xi_{-k\pm}}{2} \pm \sqrt{\left(\frac{\xi_{k\pm} + \xi_{-k\pm}}{2}\right)^2 + \Delta_{k\pm} \Delta_{k\pm}^{\dagger}} \quad (13)$$

The presence of the antisymmetric spin-orbit coupling would suppress spin-triplet pairing. However, it has been shown by Frigeri et al., (Frigeri et al., 2004) that the antisymmetric spin-orbit coupling is not destructive to the special spin-triplet state with the d vector

parallel to $\vec{g}_k$ ($\vec{d}_k \parallel \vec{g}_k$). Therefore, by choosing $\vec{g}_k = \sqrt{\frac{3}{2}} \frac{1}{k_F} (-k_y, k_x, 0)$, one adopts the p-

wave pairing state with parallel $\vec{d}$ vector $d_k = \Delta(-\tilde{k}_y, \tilde{k}_x, 0)$. Here the unit vector $\tilde{k} = (\tilde{k}_x, \tilde{k}_y, \tilde{k}_z) = (\cos\varphi \sin\theta, \sin\varphi \sin\theta, \cos\varphi)$.

By considering this parity-mixed pairing state the order parameter defined in (5) and (6) can be expressed as

$$\Delta(\vec{r}, \vec{k}) = [\Delta_0(\vec{r})\hat{\sigma}_0 + \vec{d}(\vec{k}) \cdot \vec{\sigma}] i\hat{\sigma}_y = [\Delta_0(\vec{r})\hat{\sigma}_0 + \Delta(\vec{r})(-k_y\hat{\sigma}_x + k_x\hat{\sigma}_y)] \quad (14)$$

with the spin-singlet s-wave component $\Delta_0(\vec{r})$ and the $\vec{d}$ vector $\vec{d}_k(\vec{r}) = \Delta(\vec{r})(-\tilde{k}_y, \tilde{k}_x, 0)$, here, the vector $\vec{r}$ indicates the real-space coordinates. While this spin-triplet part alone has point nodes (axial state with two point nodes), the pairing state of Eq. (14) can possess line nodes in a gap as a result of the combination with the s-wave component (Hayashi et al., 2006; Sergienko 2004). In the presence of uniform supercurrent the gap function has the $\vec{r}$ dependence as

$$\Delta(\vec{r}, \vec{k}) = \Delta_k e^{i2m\vec{v}_s \cdot \vec{r}} \quad (15)$$

where m is the bare electron mass.

The particular form of order parameter prevents the existence of interband terms in the Gor'kov equations

$$(i\omega_n - \xi_{k,\pm} - \Sigma_G(i\omega_n))\mathfrak{T}_{\pm}(k, i\omega_n) + (\Delta_{k,\pm} + \Sigma_F(i\omega_n))F_{\pm}^{\dagger}(k, \omega_n) = 1 \quad (16)$$

$$(i\omega_n + \xi_{-k,\pm} + \Sigma_G^{\dagger}(i\omega_n))F_{\pm}^{\dagger}(k, \omega_n) + (\Delta_{k,\pm}^{\dagger} + \Sigma_F^{\dagger}(i\omega_n))\mathfrak{T}_{\pm}(k, \omega_n) = 0 \quad (17)$$

where in this case

$$\tilde{\Sigma}_G(i\omega_n) = (n_n |U_n|^2 + n_m |U_m|^2) \int \frac{d\vec{k}'}{(2\pi)^3} [\mathfrak{T}_+(\vec{k}', i\omega_n) + \mathfrak{T}_-(\vec{k}', i\omega_n)] \quad (18)$$

$$\tilde{\Sigma}_F(i\omega_n) = (n_n |U_n|^2 + n_m |U_m|^2) \int \frac{d\vec{k}'}{(2\pi)^3} [F_+(\vec{k}', i\omega_n) + F_-(\vec{k}', i\omega_n)] \quad (19)$$

and

$$\Delta_{\pm} = \Delta_0 \pm d|g_k| \quad (20)$$

I consider the superconducting gaps $|\Delta_0 + \Delta \sin \theta|$ and $|\Delta_0 - \Delta \sin \theta|$ on the Fermi surfaces I and II, respectively (such as superconductor CePt₃Si). Such a gap structure can lead to line nodes on either Fermi surface I or II (Hayashi et al., 2006). These nodes are the result of the superposition of spin-singlet and spin-triplet contributions (each separately would not produce line nodes). On the Fermi surface I, the gap is $|\Delta_0 + \Delta \sin \theta|$ and is nodeless, (not that we choose $\Delta_0 > 0$ and $\Delta > 0$). On the other hand, the form of the gap on the Fermi surface II is $|\Delta_0 - \Delta \sin \theta|$, where line nodes can appear for $\Delta_0 < \Delta$ (Hayashi et al., 2006).

3. Effects of impurities on the transition temperature of a noncentrosymmetrical superconductor

In the case of large SO band splitting, the order parameter has only intraband components and the gap equation (Eq. (9)) becomes

$$\Delta_{k\pm} = -T \sum_{n,\nu} \int \frac{d^3 k'}{(2\pi)^3} V_{\pm\nu}(\vec{k}, \vec{k}') \frac{\Delta_{k\pm}}{(i\omega_n - \Sigma_{imp} - \xi_{k\pm})(i\omega_n - \Sigma_{imp} + \xi_{-k\pm}) - \Delta_{k\pm} \Delta_{k\pm}^{\dagger}} \quad (21)$$

The coupling constants $V_{\lambda\lambda'}(\vec{k}, \vec{k}')$ I have used in previous considerations can be expressed through the real physical interactions between the electrons naturally introduced in the initial spinor basis where BCS type Hamiltonian has the following form

$$H_{\text{int}} = \frac{1}{4\Omega} \sum_{kk'q} \sum_{\alpha\beta\mu\delta} \left[V_{\alpha\beta\mu\delta}^s(\vec{k}, \vec{k}') + V_{\alpha\beta\mu\delta}^t(\vec{k}, \vec{k}') + V_{\alpha\beta\mu\delta}^m(\vec{k}, \vec{k}') \right] \times c_{\vec{k}+\vec{q},\lambda}^{\dagger} c_{-\vec{k},\lambda}^{\dagger} c_{-\vec{k}',\lambda} c_{\vec{k}+\vec{q},\lambda'} \quad (22)$$

where the pairing interaction is represented as a sum of the k-even, k-odd, and mixed-parity terms: $V = V^s + V^t + V^m$. The even contribution is

$$V_{\alpha\beta\mu\delta}^s(\vec{k}, \vec{k}') = V^s(\vec{k}, \vec{k}') (i\sigma_2)_{\alpha\beta} (i\sigma_2)_{\mu\delta}^\dagger \quad (23)$$

The odd contribution is

$$V_{\alpha\beta\mu\delta}^t(\vec{k}, \vec{k}') = V_{ij}^t(\vec{k}, \vec{k}') (i\sigma_i\sigma_2)_{\alpha\beta} (i\sigma_j\sigma_2)_{\mu\delta}^\dagger \quad (24)$$

here the amplitudes $V^s(\vec{k}, \vec{k}')$ and $V_{ij}^t(\vec{k}, \vec{k}')$ are even and odd with respect to their arguments correspondingly.

Finally, the mixed-parity contribution is

$$V_{\alpha\beta\mu\delta}^m(\vec{k}, \vec{k}') = V_i^m(\vec{k}, \vec{k}') (i\sigma_i\sigma_2)_{\alpha\beta} (i\sigma_2)_{\mu\delta}^\dagger + V_i^m(\vec{k}', \vec{k}) (i\sigma_2)_{\alpha\beta} (i\sigma_i\sigma_2)_{\mu\delta}^\dagger \quad (25)$$

The first term on the right-hand side of Eq. (25) is odd in k and even in k', while the second term is even in k and odd in k'.

The pairing interaction leading to the gap function [Eq. (14)] is characterized by three coupling constants, V_s , V_t , and V_m . Here, V_s , and V_t result from the pairing interaction within each spin channel (s : singlet, t : triplet). V_m is the scattering of Cooper pairs between those two parity channels, present in systems without inversion symmetry. The linearized gap equations acquire simple algebraic form

$$\Delta_0 = V_s \pi T \sum_n \langle E_+ \rangle + V_m \pi T \sum_n \langle \sin \theta E_- \rangle \quad (26)$$

$$\Delta = V_t \pi T \sum_n \langle \sin \theta E_- \rangle + V_m \pi T \sum_n \langle E_+ \rangle \quad (27)$$

where the angular brackets denote the average over the Fermi surface, assuming the spherical Fermi surface for simplicity, $E_\pm = \frac{E_I \pm E_{II}}{2}$, $E_{I,II} = \frac{\Delta_0 \pm \Delta \sin \theta}{B_{I,II}}$, and

$$B_{I,II} = \left[\left(\omega_n + i \sum_{imp} \right)^2 + |\Delta_0 \pm \Delta \sin \theta|^2 \right]^{1/2} \quad (28)$$

From Eqs. (26) and (27) one obtains then the following expression for the critical temperature

$$\ln \frac{T_{c0}}{T_c} = (1 - X) \left[\Psi \left(\frac{1}{2} + \frac{1}{4\pi T \tau_m} \right) - \Psi \left(\frac{1}{2} \right) \right] + X \left\{ \Psi \left[\frac{1}{2} + \frac{1}{4\pi T_c} \left(\frac{1}{\tau_n} + \frac{1}{\tau_m} \right) \right] - \Psi \left(\frac{1}{2} \right) \right\} \quad (29)$$

where

$$\frac{1}{\tau_n} = 2\pi n_n N_0 |U_n|^2 \quad \frac{1}{\tau_m} = 2\pi n_m N_0 |U_m|^2 \quad (30)$$

$\Psi(x)$ is the digamma function, $N_0 = (N_+ + N_-)/2$, $N_{\pm}$ are the densities of state (DOS) of the two bands at the Fermi level, and T_{c0} is the critical temperature of the clean superconductor.

The coefficient $X = 1 - \frac{\langle \Delta(p) \rangle_{FS}^2}{\langle \Delta^2(p) \rangle_{FS}}$ quantifies the degree of anisotropy of the order parameter

on the Fermi surface (FS), where the angular brackets $\langle \dots \rangle_{FS}$ stand for a FS average.

For isotropic *s*-wave pairing $\langle \Delta(p) \rangle_{FS}^2 = \langle \Delta^2(p) \rangle_{FS}$ ($X = 0$) and for any pairing state with angular momentum $l > 1$, e.g., for *p*-wave and *d*-wave states ($l = 1, 2$), $\left(X = 1, \frac{1}{\tau_m} = 0 \right)$ Eq. (29) reduces to the well-known expressions (Abrikosov, 1993; Abrikosov, A. A. & Gor'kov, 1959).

$$\ln \frac{T_{c0}}{T_c} = \Psi \left(\frac{1}{2} + \frac{1}{4\pi T_c \tau_m} \right) - \Psi \left(\frac{1}{2} \right) \quad (31)$$

$$\ln \frac{T_{c0}}{T_c} = \Psi \left(\frac{1}{2} + \frac{1}{4\pi T_c \tau_n} \right) - \Psi \left(\frac{1}{2} \right) \quad (32)$$

For mixing of *s*-wave state with some higher angular harmonic state, e.g., for example *s* + *p* and *s* + *d*, $\left(0 < X < 1, \frac{1}{\tau_m} = 0 \right)$, Eq. (29) becomes

$$\ln \frac{T_{c0}}{T_c} = X \left[\Psi \left(\frac{1}{2} + \frac{1}{4\pi T_c \tau_n} \right) - \Psi \left(\frac{1}{2} \right) \right] \quad (33)$$

At $\tau_n T_{c0} \gg 1$ and $\tau_m T_{c0} \gg 1$ (weak scattering) one has from Eq. (29):

$$T_{c0} - T_c \approx \frac{\pi}{4} \left[\frac{X}{2} \frac{1}{\tau_n} + \frac{1-X}{2} \frac{1}{\tau_m} \right] \quad (34)$$

In two particular cases of (i) both nonmagnetic and magnetic scattering in an isotropic *s*-wave superconductor ($X = 0$) and (ii) nonmagnetic scattering only in a superconductor with arbitrary anisotropy of $\Delta(p)$ ($\frac{1}{\tau_m} = 0, 0 < X < 1$), the Eq. (34) reduces to well-known expressions

$$T_{c0} - T_c \approx \frac{\pi}{4\tau_m} \quad (35)$$

$$T_{c0} - T_c \approx \frac{\pi X}{8\tau_n} \quad (36)$$

In the strong scattering limit ($\tau_n T_c \ll 1$, $\tau_m T_c \ll 1$), by using

$$\Psi\left(\frac{1}{2} + \frac{1}{4\pi T_c \tau}\right) - \Psi\left(\frac{1}{2}\right) \approx \ln\left(\frac{\gamma}{\pi \tau T_c}\right) + \frac{2\pi^2}{3}(\tau T)^2 + O(\tau T)^3 \quad (37)$$

From Eq. (29) one finds

$$\left(\frac{1}{\tau_m}\right)^{1-X} \left(\frac{1}{\tau_n} + \frac{1}{\tau_m}\right)^X = \frac{\pi}{\gamma} T_{c0} 2^{X-1} \quad (38)$$

One can see that the left hand side of Eq. (38) increases monotonically with both $\frac{1}{\tau_n}$ and $\frac{1}{\tau_m}$ for any value of X , with the exception of the case $X = 0$ which does not depend on magnetic impurities.

For strongly anisotropic gap parameter ($X \sim 1$), Eq. (38) reduces to

$$\frac{1}{\tau_n} + \frac{1}{\tau_m} = \frac{\pi}{\gamma} T_{c0} \quad (39)$$

i.e., the contribution of magnetic and nonmagnetic impurities to pairing breaking is about the same.

For strongly isotropic case ($X \ll 1$), one has

$$\frac{1}{\tau_m} = \frac{\pi}{2\gamma} T_{c0} \quad (40)$$

and T_c is determined primarily by magnetic impurities.

For the case of $s + p$ wave pairing in the absence of magnetic impurities, one has

$$\left(\frac{1}{\tau_n}\right)^X = \frac{\pi}{\gamma} 2^{X-1} T_{c0} \quad (41)$$

In this case the value of T_c asymptotically goes to zero as τ_n^{-1} increase, whereas T_c of a d or p wave superconductor with $X = 1$ vanishes at a critical value $\frac{1}{\tau_n^c} = \frac{\pi T_{c0}}{\gamma}$.

In the absence of nonmagnetic impurities one obtains

$$\left(\frac{1}{\tau_m}\right) = \frac{\pi}{\gamma} 2^{X-1} T_{c0} \quad (42)$$

And for the s -wave superconductor with $X = 0$ one has $\frac{1}{\tau_m^c} = \frac{\pi}{2\gamma} T_{c0}$.

Application of these results to real noncentrosymmetric materials is complicated by the lack of definite information about the superconducting gap symmetry and the distribution of the pairing strength between the bands.

As far as the pairing symmetry is concerned, there is strong experimental evidence that the superconducting order parameter in CePt₃Si has lines of gap nodes (Yasuda et al., 2004; Izawa et al., 2005; Bonalde et al., 2005). The lines of nodes are required by symmetry for all nontrivial one-dimensional representations of C_{4v} (A_2 , B_1 , and B_2), so that the superconductivity in CePt₃Si is most likely unconventional. This can be verified using the measurements of the dependence of T_c on the impurity concentration: For all types of unconventional pairing, the suppression of the critical temperature is described by the universal Abrikosov-Gor'kov function, see Eq. (32).

It should be mentioned that the lines of gap nodes can exist also for conventional pairing (A_1 representation), in which case they are purely accidental. While the accidental nodes would be consistent with the power-law behavior of physical properties observed experimentally, the impurity effect on T_c in this case is qualitatively different from the unconventional case. In this case in the absence of magnetic impurities one obtains the following equation for the critical temperature:

$$\ln \frac{T_{c0}}{T_c} = X \left[\Psi \left(\frac{1}{2} + \frac{1}{4\pi T_c \tau_n} \right) - \Psi \left(\frac{1}{2} \right) \right] \quad (43)$$

In the low ($\tau_n T_c \gg 1$) and dirty ($\tau_n T_{c0} \ll 1$) limit of impurity concentration one has

$$T_{c0} - T_c \approx \frac{X\pi}{8\tau_n} \quad \tau T_{c0} \gg 1 \quad (44)$$

$$T_c = T_{c0} \left(\frac{\pi \tau_n T_{c0}}{e^C} \right)^{\frac{X}{1-X}} \quad \tau T_{c0} \ll 1 \quad (45)$$

This means that anisotropy of the conventional order parameter increases the rate at which T_c is suppressed by impurities. Unlike the unconventional case, however, the superconductivity is never completely destroyed, even at strong disorder.

4. Low temperature magnetic penetration depth of a superconductor without inversion symmetry

To determine the penetration depth or superfluid density in a superconductor without inversion symmetry one calculates the electromagnetic response tensor $K(\vec{q}, \vec{v}_s, T)$, relating the current density $\vec{J}$ to an applied vector potential $\vec{A}$

$$\vec{J}(\vec{q}) = -K(\vec{q}, \vec{v}_s, T) \vec{A}(\vec{q}) \quad (46)$$

The expression for the response function can be obtained as

$$K(\vec{q}, \vec{v}_s, T, \omega_m) = \frac{ne^2}{mc} \left(1 + \frac{2\pi}{m} \left\langle T \sum_{n,k} \hat{k}_{\parallel}^2 \mathfrak{I}_{\pm}(\vec{k}_+, \omega_n) \mathfrak{I}_{\pm}(\vec{k}_-, \omega_n, \omega_m) \right\rangle \right) \quad (47)$$

where $k_{\pm} = k \pm \frac{q}{2}$, $\hat{k}_{\parallel}^2$ is the direction of the supercurrent and $\langle \dots \rangle$ represents a Fermi surface average.

By using the expression of Green's function into Eq. (47) one obtains

$$K(\vec{q}, \vec{v}_s, T, \omega_n) = \frac{ne^2}{mc} \left(1 + \frac{2\pi T}{m} \sum_n \int \frac{d^2k}{(2\pi)^2} \hat{k}_{\parallel}^2 \frac{(i\omega_n - \Sigma_{imp} + \vec{v}_s \cdot \vec{k}_F)^2 + \xi_{k_+, \pm} \xi_{k_-, \pm} + \Delta_{k_+} \Delta_{k_-}}{\left[(i\omega_n - \Sigma_{imp} + \vec{v}_s \cdot (\vec{k} + \frac{\vec{q}}{2}))^2 - E_{k_+, \pm}^2 \right] \left[(i\omega_n - \Sigma_{imp} + \vec{v}_s \cdot (\vec{k} - \frac{\vec{q}}{2}))^2 - E_{k_-, \pm}^2 \right]} \right) \quad (48)$$

Now we separate out the response function as

$$K(\vec{q}, \vec{v}_s, T) = K(0, 0, 0) + \delta K(\vec{q}, \vec{v}_s, T) \quad (49)$$

where $K(0, 0, 0) = \frac{c}{4\pi\lambda^2(0)}$ ($\lambda(0) = \left(\frac{mc^2}{4\pi ne^2} \right)^{1/2}$ is the zero temperature London penetration depth).

Doing the summation over Matsubara frequencies for each band one gets

$$\begin{aligned} \delta K(\vec{q}, \vec{v}_s, T) &= -\frac{2n_+ e^2}{mc} \left\langle \hat{k}_{\parallel}^2 \int_0^{\infty} d\omega \operatorname{Re} \frac{[f(\omega - \vec{v}_s \cdot \vec{k}_F) - f(-\omega - \vec{v}_s \cdot \vec{k}_F) \Delta_{k_+, +}^2]}{\sqrt{(\omega - i\Sigma_{imp})^2 - \Delta_{k_+, +}^2} \left[\Delta_{k_+, +}^2 + \left(\frac{\vec{q} \cdot \vec{k}_F}{2m} \right)^2 - (2\alpha g_k)^2 - (\omega - i\Sigma_{imp})^2 \right]} \right\rangle \\ &= -\frac{2n_- e^2}{mc} \left\langle \hat{k}_{\parallel}^2 \int_0^{\infty} d\omega \operatorname{Re} \frac{[f(\omega - \vec{v}_s \cdot \vec{k}_F) - f(-\omega - \vec{v}_s \cdot \vec{k}_F) \Delta_{k_-, -}^2]}{\sqrt{(\omega - i\Sigma_{imp})^2 - \Delta_{k_-, -}^2} \left[\Delta_{k_-, -}^2 + \left(\frac{\vec{q} \cdot \vec{k}_F}{2m} \right)^2 - (2\alpha g_k)^2 - (\omega - i\Sigma_{imp})^2 \right]} \right\rangle \\ &= -\frac{n_+ e^2}{mc} \left\{ 1 - 2 \left\langle \hat{k}_{\parallel}^2 \frac{\sinh^{-1} \left(\frac{\vec{q} \cdot \vec{k}_F}{2m\Delta_{k_+, +}} \right)}{\left(\frac{\vec{q} \cdot \vec{k}_F}{2m\Delta_{k_+, +}} \right) \sqrt{1 + \left(\frac{\vec{q} \cdot \vec{k}_F}{2m\Delta_{k_+, +}} \right)^2 + \left(\frac{2\alpha g_k}{\Delta_{k_+, +}} \right)^2}} \right\rangle \right. \\ &\quad + \left\langle \hat{k}_{\parallel}^2 \ln \frac{\sqrt{\Delta_{k_+, +}^2 + \left(\frac{\vec{q} \cdot \vec{k}_F}{2m} \right)^2} - \left(1 + \left(\frac{\vec{q} \cdot \vec{k}_F}{2m\Delta_{k_+, +}} \right)^2 - \left(\frac{2\alpha g_k}{\Delta_{k_+, +}} \right)^2 \right) \Sigma_{imp} - \left(\frac{\vec{q} \cdot \vec{k}_F}{2m\Delta_{k_+, +}} \right) \sqrt{1 + \left(\frac{\vec{q} \cdot \vec{k}_F}{2m\Delta_{k_+, +}} \right)^2 - \left(\frac{4m\Delta_{k_+, +} \alpha g_k}{\vec{q} \cdot \vec{k}_F} \right)^2} \sqrt{\Sigma_{imp}^2 - \Delta_{k_+, +}^2}}{\sqrt{\Delta_{k_+, +}^2 + \left(\frac{\vec{q} \cdot \vec{k}_F}{2m} \right)^2} + \left(1 + \left(\frac{\vec{q} \cdot \vec{k}_F}{2m\Delta_{k_+, +}} \right)^2 - \left(\frac{2\alpha g_k}{\Delta_{k_+, +}} \right)^2 \right) \Sigma_{imp} - \left(\frac{\vec{q} \cdot \vec{k}_F}{2m\Delta_{k_+, +}} \right) \sqrt{1 + \left(\frac{\vec{q} \cdot \vec{k}_F}{2m\Delta_{k_+, +}} \right)^2 - \left(\frac{4m\Delta_{k_+, +} \alpha g_k}{\vec{q} \cdot \vec{k}_F} \right)^2} \sqrt{\Sigma_{imp}^2 - \Delta_{k_+, +}^2}} \right\rangle \\ &\quad \left. + 2 \left\langle \hat{k}_{\parallel}^2 \int_0^{\infty} d\omega \operatorname{Re} \frac{[f(\omega - \vec{v}_s \cdot \vec{k}_F) + f(\omega + \vec{v}_s \cdot \vec{k}_F)]}{\sqrt{(\omega - i\Sigma_{imp})^2 - \Delta_{k_+, +}^2} \left[\Delta_{k_+, +}^2 + \left(\frac{\vec{q} \cdot \vec{k}_F}{2m} \right)^2 - (2\alpha g_k)^2 - (\omega - i\Sigma_{imp})^2 \right]} \right\rangle \right\} \quad (50) \end{aligned}$$

$$\begin{aligned}
& \frac{-n_- e^2}{mc} \left\{ 1 - 2 \left\langle \frac{\hat{k}_{\parallel}^2 \sinh^{-1} \left(\frac{\vec{q} \cdot \vec{k}_F}{2m\Delta_{k,-}} \right)}{\left(\frac{\vec{q} \cdot \vec{k}_F}{2m\Delta_{k,-}} \right) \sqrt{1 + \left(\frac{\vec{q} \cdot \vec{k}_F}{2m\Delta_{k,-}} \right)^2 + \left(\frac{2\alpha g_k}{\Delta_{k,-}} \right)^2}} \right\rangle \right. \\
& + \left\langle \frac{\hat{k}_{\parallel}^2 \ln \left[\frac{\sqrt{\Delta_{k,-}^2 + \left(\frac{\vec{q} \cdot \vec{k}_F}{2m} \right)^2} - \left(1 + \left(\frac{\vec{q} \cdot \vec{k}_F}{2m\Delta_{k,-}} \right)^2 - \left(\frac{2\alpha g_k}{\Delta_{k,-}} \right)^2 \right) \Sigma_{imp} - \left(\frac{\vec{q} \cdot \vec{k}_F}{2m\Delta_{k,-}} \right) \sqrt{1 + \left(\frac{\vec{q} \cdot \vec{k}_F}{2m\Delta_{k,-}} \right)^2 - \left(\frac{4m\Delta_{k,-}\alpha g_k}{\vec{q} \cdot \vec{k}_F} \right)^2} \sqrt{\Sigma_{imp}^2 - \Delta_{k,-}^2}}}{\sqrt{\Delta_{k,-}^2 + \left(\frac{\vec{q} \cdot \vec{k}_F}{2m} \right)^2} + \left(1 + \left(\frac{\vec{q} \cdot \vec{k}_F}{2m\Delta_{k,-}} \right)^2 - \left(\frac{2\alpha g_k}{\Delta_{k,-}} \right)^2 \right) \Sigma_{imp} - \left(\frac{\vec{q} \cdot \vec{k}_F}{2m\Delta_{k,-}} \right) \sqrt{1 + \left(\frac{\vec{q} \cdot \vec{k}_F}{2m\Delta_{k,-}} \right)^2 - \left(\frac{4m\Delta_{k,-}\alpha g_k}{\vec{q} \cdot \vec{k}_F} \right)^2} \sqrt{\Sigma_{imp}^2 - \Delta_{k,-}^2}} \right] \right\rangle \\
& \left. + 2 \left\langle \frac{\hat{k}_{\parallel}^2 \int d\omega \operatorname{Re} \frac{\left[f(\omega - \vec{v}_s \cdot \vec{k}_F) + f(\omega + \vec{v}_s \cdot \vec{k}_F) \right]}{\sqrt{(\omega - i\Sigma_{imp})^2 - \Delta_{k,-}^2} \left[\Delta_{k,-}^2 + \left(\frac{\vec{q} \cdot \vec{k}_F}{2m} \right)^2 - (2\alpha g_k)^2 - (\omega - i\Sigma_{imp})^2 \right]} \right\rangle \right\} \quad (50)
\end{aligned}$$

The factor αg_k characterizes and quantifies the absence of an inversion center in a crystal lattice. This is the main result of my work i.e. nonlocality, nonlinearity, impurity and nonsentrosymmetry are involved in the response function. The first two terms in Eq. (50) represent the nonlocal correction to the London penetration depth and the third represents the nonlocal and impure renormalization of the response while the forth combined nonlocal, nonlinear, and impure corrections to the temperature dependence.

I consider a system in which a uniform supercurrent flows with the velocity $\vec{v}_s$, so all quasiparticles Matsubara energies modified by the semiclassical Doppler shift $\vec{v}_s \cdot \vec{k}_F$.

The specular boundary scattering in terms of response function can be written as (Kosztin & Leggett, 1997)

$$\frac{\Delta\lambda_{spec}(T)}{\lambda_0} = \frac{2}{\pi} \int_0^\infty d\tilde{q} \frac{\delta K(\tilde{q}, \vec{v}_s, T)}{(\tilde{q}^2 + 1)^2} \quad (51)$$

In the pure case there are four relevant energy scales in the low energy sector in the Meissner state: T , E_{nonlin} , E_{nonloc} , and αg_k . The first two are experimentally controlled parameters while the last two are intrinsic one.

In low temperatures limit the contribution of the fully gap ($|\Delta_0 + \Delta \sin \theta|$) Fermi surface I decrease and the effect of the gap $|\Delta_0 - \Delta \sin \theta|$ Fermi surface II is enhanced. I consider geometry where the magnetic field is parallel to c axis and thus $\vec{v}_s$ and the penetration direction $\vec{q}$ are in the ab plane, and in general, $\vec{v}_s$ makes an angle φ with the axis. There are two effective nonlinear energy scales $E_{nonlin}^+ = v_s k_F u_{\phi l_1}$ and $E_{nonlin}^- = v_s k_F u_{\phi l_2}$. where $u_{\phi l} = |\cos \varphi + l \sin \varphi|$ and $l_1, l_2 = \pm 1$.

In the nonlocal ($q \neq 0$), linear ($v_s \rightarrow 0$) limit, i.e., in the range of temperature where $E_{nonlin} \ll T \ll E_{nonloc}$ one gets

$$\delta K(q, 0, T) = \begin{cases} \frac{-c}{4\pi\lambda_0^2} (2\ln 2) \frac{T}{\Delta_0} & \alpha' w_{\phi l} \ll T \\ \frac{-c}{4\pi\lambda_0^2} \sum_l u_{\phi l}^2 \left(\frac{\pi}{4} \frac{\alpha' w_{\phi l}}{\Delta_0} + \frac{3}{2} \xi(3) \frac{T^3}{\Delta_0 \alpha'^2 w_{\phi l}^2} \right) & \alpha' w_{\phi l} \gg T \end{cases} \quad (52)$$

where $w_{\theta l} = |\sin \theta - l \cos \theta|$, $u_{\theta l} = |\cos \theta + l \sin \theta|$, and $\alpha' = \frac{qv_F}{2\sqrt{2}} - \sqrt{2}\alpha g_k$.

Depending on the effective nonlocal energy scales $\left(E_{nonloc}^+ = \frac{v_F u_{\theta l_1}}{\lambda_0}, E_{nonloc}^- = \frac{v_F u_{\theta l_2}}{\lambda_0}, l_1, l_2 = \pm 1\right)$ one obtains

$$\frac{\Delta \lambda_{spec}(T)}{\lambda_0} \propto \begin{cases} T & E_{nonloc}^+, E_{nonloc}^- \ll T \\ T & E_{nonloc}^- \ll T \ll E_{nonloc}^+ \\ T^2 & E_{nonloc}^+, E_{nonloc}^- \gg T \end{cases} \quad (53)$$

For CePt₃Si superconductor with $T_c = 0.75K$, the linear temperature dependence would crossover to a quadratic dependence below $T_{nonloc}^* \sim 0.015K$.

Magnetic penetration depth measurements in CePt₃Si did not find a T^2 law as expected for line nodes. I argue that it may be due to the fact that such measurements were performed above 0.015K. On the other hand, it is note that CePt₃Si is an extreme type-II superconductor with the Ginzburg-Landau parameter, $K = 140$, and the nonlocal effect can be safely neglected, and because this system is a clean superconductor, neglect the impurity effect can be neglected (Bauer et al., 2004; Bauer et al., 2005).

In the local, clean, and nonlinear limit ($q \rightarrow 0, v_s \neq 0$) the penetration depth is given by

$$\lambda_{spec}^{(loc)} = \left(\frac{c}{4\pi\delta K(q \rightarrow 0, \vec{v}_s, T)} \right)^{1/2} \quad (54)$$

Where

$$\begin{aligned} \delta K(\vec{q}, \vec{v}_s, T) = & \frac{-n_+ e^2}{mc} \left\{ 1 - 2 \left\langle \hat{k}_{\parallel}^2 \frac{\sinh^{-1} \left(\frac{\vec{q} \cdot \vec{k}_F}{2m\Delta_{k,+}} \right)}{\left(\frac{\vec{q} \cdot \vec{k}_F}{2m\Delta_{k,+}} \right) \sqrt{1 + \left(\frac{2\alpha k_F}{\Delta_{k,+}} \right)^2}} \right\rangle + 2 \left\langle \hat{k}_{\parallel}^2 \int d\omega \operatorname{Re} \frac{\left[f(\omega - \vec{v}_s \cdot \vec{k}_F) + f(\omega + \vec{v}_s \cdot \vec{k}_F) \right]}{\sqrt{\omega^2 - \Delta_{k,+}^2} \left[\Delta_{k,+}^2 - (2\alpha g_k)^2 - \omega^2 \right]} \right\rangle \right\} \\ & \frac{-n_- e^2}{mc} \left\{ 1 - 2 \left\langle \hat{k}_{\parallel}^2 \frac{\sinh^{-1} \left(\frac{\vec{q} \cdot \vec{k}_F}{2m\Delta_{k,-}} \right)}{\left(\frac{\vec{q} \cdot \vec{k}_F}{2m\Delta_{k,-}} \right) \sqrt{1 + \left(\frac{2\alpha k_F}{\Delta_{k,-}} \right)^2}} \right\rangle + 2 \left\langle \hat{k}_{\parallel}^2 \int d\omega \operatorname{Re} \frac{\left[f(\omega - \vec{v}_s \cdot \vec{k}_F) + f(\omega + \vec{v}_s \cdot \vec{k}_F) \right]}{\sqrt{\omega^2 - \Delta_{k,-}^2} \left[\Delta_{k,-}^2 - (2\alpha g_k)^2 - \omega^2 \right]} \right\rangle \right\} \end{aligned} \quad (55)$$

Thus by considering only the second term in the right hand side of Eq. (55) into Eq. (51) one gets

$$\frac{\delta\lambda_{\text{spec}}}{\lambda_0} \approx \begin{cases} \frac{1}{2} \ln 2 \sum_{l=\pm 1} u_{\theta l}^2 \frac{T}{\Delta_0} & E_{\text{nonlin}}^+, E_{\text{nonlin}}^- \ll T \\ \frac{1}{2} \ln 2 u_{\theta l_2}^2 \frac{T}{\Delta_0} + \frac{u_{\theta l_1}^3}{2\sqrt{2}} \left[\frac{v_s k_F}{2\Delta_0} + \frac{2\alpha g_k}{\Delta_0} \right] & E_{\text{nonlin}}^- \ll T \ll E_{\text{nonlin}}^+ \\ \frac{1}{2} \sum_{l=\pm} u_{\theta l}^3 \left[\frac{v_s k_F}{2\Delta_0} + \frac{2\alpha g_k}{\Delta_0} \right] \frac{1}{\sqrt{2}} + o \left(T e^{\frac{-(v_s k_F + 4\alpha g_k)}{\sqrt{2}T}} \right) & E_{\text{nonlin}}^+, E_{\text{nonlin}}^- \gg T \end{cases} \quad (56)$$

The linear temperature dependence of penetration depth is in agreement with Bonalde et al's result (Bonalde et al., 2005).

Thus the T behavior at low temperatures of the penetration depth in Eq. (56) is due to nonlinearity indicating the existence of line nodes in the gap parameter in CePt₃Si compound. A T linear dependence of the penetration depth in the low temperature region is expected for clean, local and nonlinear superconductors with line nodes in the gap function. Now the effect of impurities when both s -wave and p -wave Cooper pairings coexist is considered.

I assume that the superconductivity in CePt₃Si is unconventional and is affected only by nonmagnetic impurities. The equation of motion for self-energy can be written as

$$\Sigma_{\text{imp}(n)}(\vec{p}, i\omega_n) = n_n T(\vec{p}, \vec{p}', i\omega_n) \quad (57)$$

where the T matrix is given by

$$T(\vec{p}, \vec{p}', i\omega_n) = \frac{u\sigma_3}{1 - u\sigma_3 \mathfrak{I}(\vec{p}, \vec{p}', i\omega_n)} \quad (58)$$

here σ_3 is the third Pauli-spin operator.

By using the expression of the Green's function in Eq. (58) one can write

$$T(\vec{p}, \vec{p}', i\omega_n) = \frac{\pi N_0 u_0^2 I}{1 + (\pi N_0 u_0 I)^2} \quad (59)$$

where

$$I = \int_0^{2\pi} \frac{d\Omega}{4\pi} \frac{-\left(\omega + i\Sigma_{\text{imp}(n)}\right)}{\left[\left(\omega + i\Sigma_{\text{imp}(n)}\right)^2 + \Delta_+ \Delta_- \right]^{1/2}} \quad (60)$$

and u_0 is a single s -wave matrix element of scattering potential u . Small u_0 puts us in the limit where the Born approximation is valid, where large u_0 ($u_0 \rightarrow \infty$), puts us in the unitarity limit.

Theoretically it is known that the nodal gap structure is very sensitive to the impurities. If the spin-singlet and triplet components are mixed, the latter might be suppressed by the impurity scattering and the system would behave like a BCS superconductor. For p-wave gap function the polar and axial states have angular structures, $\Delta_k(T) = \Delta_0(T)\cos\theta_k$ and $\Delta_k(T) = \Delta_0(T)\sin\theta_k$ respectively. The electromagnetic response now depends on the mutual orientation of the vector potential $\mathbf{A}$ and $\hat{I}$ (unit vector of gap symmetry), which itself may be oriented by surfaces, fields and superflow. A detailed experimental and theoretical study for the axial and polar states was presented in Ref. (Einzel, 1986). In the clean limit and in the absence of Fermi-Liquid effects the following low-temperature asymptotic were obtained for axial and polar states

$$\frac{\Delta\lambda(T)^{\parallel,\perp}}{\lambda(0)} = a_{\parallel,\perp} \left(\frac{k_B T}{\Delta_0} \right)^{n_{\parallel,\perp}} \quad (61)$$

where in the axial state $n = 2(4)$ and $a = \pi^2 \left(\frac{7\pi^4}{15} \right)$, and in the polar state $n = 3(1)$ and

$$a = \frac{27\pi\xi(3)}{4} \left(\frac{3\pi\ln 2}{2} \right), \text{ for the orientations } \parallel(\perp).$$

The influence of nonmagnetic impurities on the penetration depth of a p-wave superconductor was discussed in detail in Ref (Gross et al., 1986). At very low temperatures, the main contribution will originate from the eigenvalue with the lower temperature exponent n , i.e., for the axial state (point nodes) with T^2 low, and for the polar state (line nodes) the dominating contribution with a linear T . The quadratic dependence in axial state may arise from nonlocality.

The low temperature dependence of penetration depth in polar and axial states used by Einzel et al., (Einzel et al. 1986) to analyze the $\lambda(T) \sim T^2$ behavior of UBe_{13} at low temperatures. The axial $\vec{A} \parallel \hat{I}$ case seems to be the proper state to analyze the experiment because it was favored by orientation effects and was the only one with T^2 dependence. Meanwhile, it has turned out that T^2 behavior is introduced immediately by T-matrix impurity scattering and also by weak scattering in the polar case. The axial state, and according to the Andersons theorem the s-wave value of the London penetration depth are not at all affected by small concentration of nonmagnetic impurities.

Thus, for the polar state, Eq. (60) can be written as

$$I = \int_0^{2\pi} \frac{d\theta}{2\pi} \frac{-\left(\omega + i\sum_{imp(n)}\right)}{\left[\left(\omega + i\sum_{imp(n)}\right)^2 + \Delta_0^2 \cos^2\theta\right]^{1/2}} \quad (62)$$

Doing the angular integration in Eq. (62) and using Eqs. (57) and (59) one obtains

$$\Sigma_{imp(n)} = \frac{-\left[2\tilde{\omega}N(0)n_n u_0^2 / \sqrt{\tilde{\omega}^2 + \Delta_0^2}\right] K\left(\frac{\Delta_0^2}{\sqrt{\tilde{\omega}^2 + \Delta_0^2}}\right)}{1 + \left[4N(0)^2 u_0^2 \tilde{\omega}^2 / (\tilde{\omega}^2 + \Delta_0^2)\right] K^2\left(\frac{\Delta_0^2}{\sqrt{\tilde{\omega}^2 + \Delta_0^2}}\right)} \quad (63)$$

here K is the elliptic integral and $\tilde{\omega} = \omega + i\Sigma_{imp(n)}$. We note that in the impurity dominated gapless regime, the normalized frequency $\tilde{\omega}$ takes the limiting form $\tilde{\omega} \rightarrow \omega + i\gamma$, where γ is a constant depending on impurity concentration and scattering strength.

In the low temperature limit we can replace the normalized frequency $\tilde{\omega}$ everywhere by its low frequency limiting form and after integration over frequency one gets

$$\delta K(\vec{q}, \vec{v}_s, T) = \frac{-N_1 e^2}{mc} \left\{ \frac{4\gamma\pi^2 T^2}{3} \left\langle \hat{k}_{\parallel}^2 \frac{\Delta_k^2}{(\Delta_k^2 + \gamma^2)^{5/2}} \right\rangle \right\} \quad (64)$$

As in the case of d-wave order parameter, from Eqs. (64) and (51) one finds

$$\frac{\delta\lambda(T)}{\lambda(0)} = \frac{\gamma}{4\pi\Delta_0} \ln\left(\frac{4\Delta_0}{\gamma}\right) + \frac{\pi}{24\gamma\Delta_0} T^2 \quad (65)$$

In p -wave cuprates, scattering fills in electronic states at the gap nodes, thereby suppressing the penetration depth at low temperatures and changing T -linear to T^2 behavior.

5. Effect of impurities on the low temperature NMR relaxation rate of a noncentrosymmetric superconductor

I consider the NMR spin-lattice relaxation due to the interaction between the nuclear spin magnetic moment $\gamma_n I$ (γ_n is the nuclear gyro magnetic ratio) and the hyperfine field h , created at the nucleus by the conduction electrons. Thus the system Hamiltonian is

$$H = H_0 + H_{so} + H_n + H_{int} \quad (66)$$

where H_0 and H_{so} are defined by Eqs. (1) and (2), $H_n = -\gamma_n I H$ is the Zeeman coupling of the nuclear spin with the external field $\vec{H}$, and $H_{int} = -\gamma_n I h$ is the hyperfine interaction.

The spin-lattice relaxation rate due to the hyperfine contact interaction of the nucleus with the band electron is given by

$$R = \frac{1}{T_1 T} = -\frac{J^2}{2\pi} \lim_{\omega \rightarrow 0} \frac{\text{Im} K_{+-}^R(\omega)}{\omega} \quad (67)$$

where ω is the NMR frequency, $J = \frac{8\pi}{3} \gamma_n \gamma_e$ (γ_e is the electron geomagnetic ratio) is the hyperfine coupling constant, and $K_{+-}^R(\tau)$, the Fourier transform of the retarded correlation

function of the electron spin densities at the nuclear site, in the Matsubara formalism is given by (in our units $k_B = \hbar = 1$)

$$K_{+-}^R(\tau) = -\langle T_\tau S_+(\tau) S_-(0) \rangle \quad (68)$$

here T_τ is the time order operator, τ is the imaginary time, $S_\pm(\tau) = e^{H\tau} S_\pm e^{-H\tau}$, and

$$S_+(r) = \psi_\uparrow^\dagger(\vec{r}) \psi_\downarrow(\vec{r}) \quad S_-(r) = \psi_\downarrow^\dagger(\vec{r}) \psi_\uparrow(\vec{r}) \quad (69)$$

with $\psi_\sigma^\dagger(\vec{r})$ and $\psi_\sigma(\vec{r})$ being the electron field operators.

The Fourier transform of the correlation function is given by

$$K_{+-}^R(i\omega_n) = \int_0^\beta d\tau e^{i\omega_n \tau} K_{+-}^R(\tau) \quad (70)$$

The retarded correlation function is obtained by analytical continuation of the Matsubara correlation function $K_{+-}^R(\omega) = K_{+-}^R(i\omega_n) \Big|_{i\omega_n \rightarrow \omega + i\delta}$.

From Eqs. (66)-(70), one gets

$$\frac{1}{T_1 T} = -\frac{J^2}{2\pi} \lim_{\omega \rightarrow 0} \frac{1}{\omega} \times \text{Im} \left\{ T \sum_{p, p', \omega_n} \left[\text{Tr}(\mathfrak{I}_\pm(\vec{p}, i\omega_n + i\Omega_m) \mathfrak{I}_\pm(\vec{p}', i\omega_n)) - \text{Tr}(F_\pm(\vec{p}, i\omega_n + i\Omega_m) F_\pm(\vec{p}', i\omega_n)) \right] \right\}_{i\Omega_n \rightarrow \omega + i\delta} \quad (71)$$

where $\Omega_m = 2m\pi T$ are the bosonic Matsubara frequencies. By using Eqs. (11) and (12) into Eq. (71), the final result for the relaxation rate is

$$\frac{1}{T_1 T} = J^2 \int_0^\infty d\omega \left(-\frac{\partial f}{\partial \omega} \right) \{ N_+(\omega) N_-(\omega) + M_+(\omega) M_-(\omega) \} \quad (72)$$

where $f(\omega) = \left(e^{\omega/T} + 1 \right)^{-1}$ is the Fermi Function., $N_\sigma(\omega)$ and $M_\sigma(\omega)$ defined by the retarded Green's functions as

$$N_\sigma(\omega) = - \sum_{p, \nu=\pm} \text{Im} \mathfrak{I}_\nu^R(\vec{p}, \omega) \quad (73)$$

$$M_\sigma(\omega) = - \sum_{p, \nu=\pm} \text{Im} F_\nu^R(\vec{p}, \omega) \quad (74)$$

In low temperatures limit the contribution of the fully gap ($|\Delta_0 + \Delta \sin \theta|$) Fermi surface I decrease and the effect of the gap $|\Delta_0 - \Delta \sin \theta|$ Fermi surface II is enhanced.

As I mentioned above, the experimental data for CePt₃Si at low temperature seem to point to the presence of lines of the gap nodes in gap parameter (In our gap model for $\Delta_0 < \Delta$, $|\Delta_0 - \Delta \sin \theta|$ has line nodes). Symmetry imposed gap nodes exist only for the order parameters which transform according to one of the nonunity representations of the point group. For all such order parameters $M_\sigma = 0$. Thus, Eq. (72) can be written as

$$\frac{1}{T_1 T} = \frac{J^2}{4T} \int_0^\infty \frac{d\omega}{\cosh^2\left(\frac{\omega}{2T}\right)} \{N_+(\omega)N_-(\omega)\} \quad (75)$$

In the clean limit the density of state can be calculated from BCS expression

$$N_\sigma(\omega) = N_0 \operatorname{Re} \left\langle \frac{\omega}{\sqrt{\omega^2 - \Delta_\pm^2}} \right\rangle \quad (76)$$

For the gap parameter with line nodes from Eq. (76) one gets

$$N(\omega) = N_0 \frac{\pi}{2} \frac{\omega}{\Delta_0} \quad (77)$$

Thus from Eq. (75) one has

$$\frac{1}{T_1} = \frac{J^2 \pi^2 N_0^2 T^3}{2\Delta_0^2} \quad (78)$$

Therefore, line nodes on the Fermi surface II lead to the low-temperature T^3 law in T_1^{-1} which is in qualitative agreement with the experimental results.

In the dirty limit the density of state can be written as

$$N_{imp}(\omega) = \int d\Omega \frac{N_{BCS}(\omega, \theta)}{1 + u_0^2 N_{BCS}^2(\omega, \theta)} \quad (79)$$

In the limit, $\Gamma \ll \Delta_0$ where $\Gamma = \frac{n_{imp}}{\pi N_0} \left(\frac{N}{V} \right)$ ($\frac{N}{V}$ is the electron density) the density of state is

$$N_{imp}(\omega) \approx N(0) + ac^2 \omega \quad (80)$$

where $c = \cot g\delta_0$ (δ_0 is the s-wave scattering phase shift), a is a constant, and $N(0)$ the zero energy ($\omega = 0$) quasi-particle density of state is given by

$$N(0) = N_o \left(\frac{\varsigma}{\sqrt{1 + \frac{1}{4}\varsigma^2} + \frac{1}{2}\varsigma} \right)^{1/2} \quad (81)$$

where $\varsigma = \frac{\Gamma}{\Delta}$.

In the unitary limit ($u_0 \rightarrow \infty$), $c = 0$ ($\delta_0 = \pi/2$), from Eqs. (75) and (80) one obtains

$$\frac{1}{T_1} = J^2 N(0)^2 T \quad (82)$$

Thus the power-low temperature dependence of T_1^{-1} is affected by impurities and it changes to linear temperature dependence characteristic of the normal state Koringa relation again is in agreement with the experimental results.

6. Conclusion

In this chapter I have studied theoretically the effect of both magnetic and nonmagnetic impurities on the superconducting properties of a non-centrosymmetric superconductor and also I have discussed the application of my results to a model of superconductivity in CePt₃Si.

First, the critical temperature is obtained for a superconductor with an arbitrary of impurity concentration (magnetic and nonmagnetic) and an arbitrary degree of anisotropy of the superconducting order parameter, ranging from isotropic s wave to p wave and mixed (s+p) wave as particular cases.

The critical temperature is found to be suppressed by disorder, both for conventional and unconventional pairings, in the latter case according to the universal Abrikosov-Gor'kov function.

In the case of nonsentrosymmetrical superconductor CePt₃Si with conventional pairing (A_1 representation with purely accidental line nodes), I have found that the anisotropy of the conventional order parameter increases the rate at which T_c is suppressed by impurities. Unlike the unconventional case, however, the superconductivity is never completely destroyed, even at strong disorder.

In section 4, I have calculated the appropriate correlation function to evaluate the magnetic penetration depth. Besides nonlinearity and nonlocality, the effect of impurities in the magnetic penetration depth when both s-wave and p-wave Cooper pairings coexist, has been considered.

For superconductor CePt₃Si, I have shown that such a model with different symmetries describes the data rather well. In this system the low temperature behavior of the magnetic penetration depth is consistence with the presence of line nodes in the energy gap and a quadratic dependence due to nonlocality may accrue below $T_{nonloc}^* = 0.015K$. In a dirty superconductor the quadratic temperature dependence of the magnetic penetration depth may come from either impurity scattering or nonlocality, but the nonlocality and nodal behavior may be hidden by the impurity effects.

Finally, I have calculated the nuclear spin-lattice relaxation of CePt₃Si superconductor. In the clean limit the line nodes which can occur due to the superposition of the two spin channels lead to the low temperature T^3 law in T_1^{-1} . In a dirty superconductor the linear temperature dependence of the spin-lattice relaxation rate characteristic of the normal state Korringa relation.

7. Acknowledgment

I wish to thank the Office of Graduate Studies and Research Vice President of the University of Isfahan for their support.

8. References

- Abrikosov, A. A. (1993). Influence of the Gap Anisotropy on Superconducting Properties. *Physica C*, Vol. 214, No. 1-2, (1 September 1993), pp.107-110, ISSN 0921-4534
- Abrikosov, A. A. & Gor'kov, L. P. (1959). On the theory of superconducting alloys. *Soviet Physics, JETP*, Vol. 8, pp. 1090. ISSN 0038-5654
- Abrikosov, A. A. & Gor'kov, L. P. (1961). Contribution to the theory of superconducting alloys with paramagnetic impurities. *Soviet Physics, JETP*, Vol. 12, pp. 1243. ISSN 0038-5654
- Abrikosov, A. A; Gor'kov, L. P & Dzyaloshnii, I. E. (1975). *Methods of Quantum Field Theory in Statistical Physics*, Dover Publications Inc, ISBN 0486632288, New York.
- Akazawa, T; Hidaka, H; Fujiwara, T; Kobayashi, T. C; Yamamoto, E; Haga, Y; Settai, R. & Onuki, Y. (2004). Pressure-induced Superconductivity in UIr without inversion symmetry. *Journal of Physics: Condensed Matter*, Vol. 16, No. 4, (16 January 2004), pp. 29-32, ISSN 1361-648X
- Anderson, P. W. (1959). Theory of dirty superconductors. *Journal of Physics and Chemistry of Solids*, Vol. 11, No. 1-2, (September 1959), pp. 26-30, ISSN 0022-3697
- Anderson, P. W. (1984). Structure of triplet superconducting energy gap. *Physical Review B*, Vol. 30, No.7, (1 October 1984), pp. 4000-4002, ISSN 1550-235X
- Bauer, E; Bonalde, I. & Sigrist, M. (2005). Superconductivity and normal state properties of non-centrosymmetric CePt₃Si: a status report. *Low Temp. Phys*, Vol. 31, No. 8, (7 October 2005), pp.748-757,
- Bauer, E; Hilscher, G; Michor, H; Paul, Ch; Scheidt, E. W; Griбанov, A; Seropegin, Yu; Noël, H; Sigrist, M. & Rogl, P. (2004). Heavy Fermion Superconductivity and Magnetic Order in Noncentrosymmetric CePt₃Si. *Physical Review Letters*, Vol. 92, No. 2, (13 January 2004), pp.027003 [4 pages], ISSN 1079-7114
- Bauer, E; Hilscher, G; Michor, H; Sieberer, M; . Scheidt, E. W; Griбанov, A; Seropegin, Yu; Rogl, P; Amato, A; Song, W. Y; Park, J. G; Adroja, D. T; Nicklas, M; Sparn, G; Yogi, M. & Kitaoka, Y. (2005). Unconventional superconductivity and magnetism in CePt₃Si_{1-x}Ge_x. *Physica B*, Vol. 359-361, (30 April 2005), Pp. 360-367, ISSN 0921-4526
- Blatsky, A.V; Vekhter, I. & Zhu, J.X. (2006). Impurity-induced states in conventional and unconventional superconductors. *Reviews of Modern Physics*, Vol. 78, No. 2, (9 May 2006), pp.373-433, ISSN 1539-0756

- Bonalde, I; Brämer-Escamilla, W. & Bauer, E. (2005). Evidence for Line Nodes in the Superconducting Energy Gap of Noncentrosymmetric CePt₃Si from Magnetic Penetration Depth Measurements. *Physical Review Letters*, Vol. 94, No. 20, (23 May 2005), pp.207002 [4 pages], ISSN 1079-7114
- Borkowski, L. S; & Hirschfeld, P. J. (1992). Kondo effect in gapless superconductors. *Physical Review B*, Vol.46, No.14, (1 October 1992), pp.9274–9277, ISSN 1550-235X
- Borkowski, L. S; & Hirschfeld, P. J. (1994). Distinguishing *d*-wave superconductors from highly anisotropic *s*-wave superconductors. *Physical Review B*, Vol. 49, No. 21, (1 June 1994), pp.15404–15407, ISSN 1550-235X
- Dresselhaus, G. (1955). Spin-Orbit Coupling Effects in Zinc Blende Structures. *Phys. Rev.* 100, 580. *Physical Review*, Vol.100, No.2, (15 October 1955), pp.580–586,
- Einzel, D; Hirschfeld, P. J; Gross, F; Chandrasekhar, B.S; Andres, K; Ott, H R; Beuers, J; Fisk, Z. & Smith, J.L. (1986). Magnetic Field Penetration Depth in the Heavy-Electron Superconductor UBe₁₃. *Physical Review Letters*, Vol. 56, No. 23, (9 June 1986), pp.2513–2516, ISSN 1079-7114
- Edelstein, V. M. (1995). Magnetoelectric Effect in Polar Superconductors. *Physical Review Letters*, Vol. 75, No 4, (4 September 1995), pp. 2004–2007, ISSN 1079-7114
- Frigeri, P. A; Agterberg, D.F. & Sigrist, M. (2004). Spin susceptibility in superconductors without inversion symmetry. *New Journal of Physics*, Vol.6, No.1, (2 September 2004), pp.115 [9 pages], ISSN 1367-2630
- Frigeri, P. A; Agterberg, D. F; Koga, A. & Sigrist, M. (2004). Superconductivity without Inversion Symmetry: MnSi versus CePt₃Si. *Physical Review Letters*, Vol.92, No. 9, (3 March 2004), pp.097001 [4 pages], ISSN 1079-7114
- Gor'kov, L. P & Rashba, E. I. (2001). Superconducting 2D System with Lifted Spin degeneracy: Mixed Singlet-Triplet State. *Physical Review Letters*, Vol. 87, No. 3, (16 July 2001), pp.037004 [4 pages], ISSN 1079-7114
- Gross, F; Chandrasekhar, B.S; einzel, D; Andres, K; Hirschfeld, P.J; Ott, H.R; Beuers, J; Fisk, Z. & Smith, J.L. (1986). Anomalous temperature dependence of the magnetic field penetration depth in superconducting UBe₁₃. *Zeitschrift für Physik B: Condensed Matter*, Vol. 64, No. 2, pp.175-188, ISSN 0722-3277
- Hayashi, N; Wakabayashi, K; Frigeri, P. A. & Sigrist, M. (2006). Nuclear magnetic relaxation rate in a noncentrosymmetric superconductor. *Physical Review B*, Vol. 73, No. 9, (17 March 2006), pp.092508 [4 pages], ISSN 1550-235x
- Hayashi, N; Wakabayashi, K; Frigeri, P. A. & Sigrist, M. (2006). Temperature dependence of the superfluid density in a noncentrosymmetric superconductor. *Physical Review B*, Vol. 73, No. 2, (19 January 2006), pp. 024504 [9 pages], ISSN 1550-235X
- Izawa, K; Kasahara, Y; Matsuda, Y; Behnia, K; Yasuda, T; Settai, R. & Onuki, Y. (2005) Line Nodes in the Superconducting Gap Function of Noncentrosymmetric CePt₃Si. *Physical Review Letters*, Vol. 94, No. 19, (16 May 2005), pp.197002 [4 pages], ISSN 1079-7114
- Kosztin, I & Leggett, A.J. (1997) Nonlocal Effects on the Magnetic Penetration Depth in *d*-Wave Superconductors. *Physical Review Letters*, Vol. 79, No. 1, (7 July 1997), pp.135–138, ISSN 1079-7114

- Mathur, N.D; Grosche, F. M; Julian S. R; Walker, I. R; Freye, D. M; Haselwimmer R. K. W & Lonzarich G. G. (1998). Magnetically mediated superconductivity in heavy fermion compounds. *Nature*, Vol. 394, (2 July 1998), pp.39-43 ISSN 0028-0836
- Mineev, V. P. & Samokhin, K. V. (2007). Effects of impurities on superconductivity in noncentrosymmetric compounds. *Physical Review B*, Vol.75, No. 18, (25 May 2007), pp.184529 [14 pages], ISSN 1550-235X
- Mineev, V. P. & Champel, T. (2004) Theory of superconductivity in ferromagnetic superconductors with triplet pairing. *Phys. Physical Review B*, Vol. 69, No. 14, (27 April 2004), pp.144521 [6 pages], ISSN 1550-235X
- Rashba, E. I. (1960). Properties of Semiconductors with an Extremum Loop I. Cyclotron and Combinatorial Resonance in a Magnetic Field Perpendicular to the Plane of the Loop. *Soviet Physics – Solid State*, vol. 2, (Jun. 1960), pp. 1109-1122, ISSN 0038-5654.
- Rashba, E. I & Bychkov, Yu. A. (1984). Oscillatory effects and the magnetic susceptibility of carriers in inversion layers. *Journal of Physics C: Solid State Physics*, Vol. 17, No. 33, (30 November 1984), pp.6039-6045, ISSN 0022-3719
- Salkalo, M.I; Balatsky, A.V. & Scalapino, D.J. (1996) Theory of Scanning Tunneling Microscopy Probe of Impurity States in a D-Wave Superconductor. *Physical Review Letters*, Vol. 77, No. 9, (26 August 1996), pp.1841-1844, ISSN 1079-7114
- Samokhin, K. V; Zijlstra, E. S; & Bose, S. K. (2004). CePt₃Si: An unconventional superconductor without inversion center. *Physical Review B*, Vol. 69, No. 9, (19 March 2004), pp. 094514 [8 pages], ISSN 1550-235X
- Saxena, S. S; Agarwal, P; Ahilan, K; Grosche, F. M; Haselwimmer, R. K. W; Steiner, M. J; Pugh, E; Walker, I. R; Julian, S. R; Monthoux, P; Lonzarich, G. G; Huxley, A; Sheikin, I Braithwaite D & Flouquet, J. (2000). Superconductivity on the border of itinerant-electron ferromagnetism in UGe₂. *Nature*, Vol. 406, No. 6796, (10 August 2000), pp.587-592, ISSN 0028-0836
- Sergienko, I. A. & Curnoe, S. H. (2004). Order parameter in superconductors with nondegenerate bands. *Physical Review B*, Vol.70, No. 21, (7 December 2004), pp.214510 [8 pages], ISSN 1550-235X
- Sergienko, I. A. (2004). Mixed-parity superconductivity in centrosymmetric crystals, *Physical Review B*, Vol. 69, No. 17, (18 May 2004), pp.174502 [5 pages], ISSN 1550-235X
- Shiba, H. (1968). Classical Spins in Superconductors. *Progress of Theoretical Physics*, Vol. 40, No. 3 (12 April 1968), pp. 435-451, ISSN 1347-0481
- Wang, Q.H. & Wang, Z.D. (2004). Impurity and interface bound states in $d_{x^2-y^2}+id_{xy}$ and p_x+ip_y superconductors. *Physical Review B*, Vol. 69, No. 9, (10 March 2004), pp.092502 [4 pages], ISSN 1550-235X
- Yasuda, T; Shishido, H; Ueda, T; Hashimoto, S; Settai, R; Takeuchi, T; Matsuda, T. D; Haga, Y. & Onuki, Y. (2004). Superconducting Property in CePt₃Si under Pressure. *Journal of the Physical Society of Japan*, Vol. 73, No. 7, (13 April 2004) pp. 1657-1660, ISSN: 1347-4073
- Yogi, M; Kitaoka, Y; Hashimoto, S; Yasuda, T; Settai, R; Matsuda, T. D; Haga, Y; Onuki, Y; Rogl, P. & Bauer, E. (2004). Evidence for a Novel State of Superconductivity in

Noncentrosymmetric CePt_3Si : A ^{195}Pt -NMR Study. *Physical Review Letters*, Vol. 93, No. 2, (8 July 2004), pp.027003 [4 pages], ISSN 1079-7114

Yogi, M; Kitaoka, Y; Hashimoto, S; Yasuda, T; Settai, R; Matsuda, T. D; . Haga, Y; Onuki, Y; Rogl, P. & Bauer, E. (2005). Novel superconductivity in noncentrosymmetric heavy-fermion compound CePt_3Si : a ^{195}Pt -NMR study. *Physica B*, Vol.359-361, (30 April 2005), pp. 389-391. ISSN 0921-4526

IntechOpen

IntechOpen

Superconductivity - Theory and Applications

Edited by Dr. Adir Luiz

ISBN 978-953-307-151-0

Hard cover, 346 pages

Publisher InTech

Published online 18, July, 2011

Published in print edition July, 2011

Superconductivity was discovered in 1911 by Kamerlingh Onnes. Since the discovery of an oxide superconductor with critical temperature (T_c) approximately equal to 35 K (by Bednorz and Müller 1986), there are a great number of laboratories all over the world involved in research of superconductors with high T_c values, the so-called “High- T_c superconductors”. This book contains 15 chapters reporting about interesting research about theoretical and experimental aspects of superconductivity. You will find here a great number of works about theories and properties of High- T_c superconductors (materials with $T_c > 30$ K). In a few chapters there are also discussions concerning low- T_c superconductors ($T_c < 30$ K). This book will certainly encourage further experimental and theoretical research in new theories and new superconducting materials.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Heshmatollah Yavari (2011). Effects of Impurities on a Noncentrosymmetric Superconductor: Application to CePt₃Si, Superconductivity - Theory and Applications, Dr. Adir Luiz (Ed.), ISBN: 978-953-307-151-0, InTech, Available from: <http://www.intechopen.com/books/superconductivity-theory-and-applications/effects-of-impurities-on-a-noncentrosymmetric-superconductor-application-to-cept3si>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2011 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike-3.0 License](https://creativecommons.org/licenses/by-nc-sa/3.0/), which permits use, distribution and reproduction for non-commercial purposes, provided the original is properly cited and derivative works building on this content are distributed under the same license.

IntechOpen

IntechOpen