

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

186,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Potential Application of Natural Dietary Components to Target Cancer Stem Cells

Yanyan Li^{1,2}, Steven J. Schwartz² and Duxin Sun¹

¹Department of Pharmaceutical Sciences, University of Michigan,

²Department of Food Science and Technology, Ohio State University, USA

1. Introduction

The emergence of cancer stem cell (CSC) theory has profound implications for cancer prevention and therapy. Although a large majority of chemotherapeutic drugs can considerably shrink tumor sizes (Reya et al. 2001), they often fail to eradicate tumors due to the inability to effectively kill CSCs (Reya et al. 2001; Hambardzumyan et al. 2006; Shafee et al. 2008; Korkaya et al. 2009). The cancer may eventually develop drug resistance and recurrence (Williams et al. 1987; Lippman 2000; Stockler et al. 2000; Reya et al. 2001; Zhou, B. B. et al. 2009). Therefore, the CSC population has become a promising target for cancer prevention and therapy (Zhou, B. B. et al. 2009).

Since a large number of epidemiological studies have demonstrated an association between consumption of fruits and vegetables and the reduced risk of various cancers, naturally-occurring dietary components have received considerable attention for their effects in cancer chemoprevention (Smith-Warner et al. 2003). The anti-cancer activities of many dietary components against various types of cancer have been reported for both *in vitro* and *in vivo* studies (Chinni et al. 2001; Mukhopadhyay et al. 2001; Choudhuri et al. 2002; Lamartiniere et al. 2002; Li and Sarkar 2002; Gupta, S. et al. 2003; Hastak et al. 2003; Li et al. 2003). Recently, a number of studies have found that several dietary components can directly or indirectly affect CSC self-renewal pathways (Kawasaki et al. 2008), and thus may have potential impact on CSCs. This chapter reviews current attempts to target CSCs with bioactive dietary components, with a special emphasis on our work.

2. Self-renewal pathways of CSCs

CSCs produce the tumor mass through continuous self-renewal and differentiation, which may be regulated by similar signaling pathways occurring in normal stem cells (Reya et al. 2001; Liu, S. et al. 2005). Understanding the mechanisms that underlie the self-renewal behavior of CSCs is of greatest importance for discovery and development of agents targeting CSCs. So far, several major pathways including Wnt/ β -catenin, Hedgehog, and Notch have been identified to play pivotal roles in CSC self-renewal (Smalley and Dale 1999; Dontu et al. 2004; Liu, S. et al. 2006).

2.1 Wnt/ β -catenin pathway

Wnt/ β -catenin pathway was demonstrated to modulate cell proliferation, migration, apoptosis, differentiation, and CSC self-renewal (Akiyama 2000; Polakis 2000; Yamaguchi 2001; Turashvili et al. 2006). It has been shown that Wnt/ β -catenin signaling is implicated in the maintenance of CSCs of leukemia (Ysebaert et al. 2006; Khan et al. 2007; Kawaguchi-Ihara et al. 2008), melanoma (Chien et al. 2009), breast (Li et al. 2003; Woodward et al. 2007), colon (Schulenburg et al. 2007), liver (Yang, W. et al. 2008), lung (Teng et al. 2010) cancers. For example, over-expression of β -catenin in stem cell survival pathway was shown to mediate the resistance of mouse mammary stem/progenitor cells to radiation (Woodward et al. 2007). Yang and his colleagues reported that Wnt/ β -catenin signaling promoted expansion of the hepatic progenitor cell population when it is over-expressed in transplanted rat oval cells and when it is transiently expressed in adult mice (Yang, W. et al. 2008). Elimination of β -catenin abrogated the chemo-resistant cell population endowed with progenitor-like features (Yang, W. et al. 2008).

β -Catenin, the essential mediator of canonical Wnt signaling, participates in two distinct functions in the cell, depending on its cellular localization. Membrane-localized β -catenin is sequestered by the epithelial cell-cell adhesion protein E-cadherin to maintain cell-cell adhesion (Nelson and Nusse 2004). On the other hand, cytoplasmic accumulation of β -catenin and its subsequent nuclear translocation, followed by cooperation with the transcription factors T cell factor/lymphoid enhancer factor (TCF/LEF) as a transcription activator, eventually leads to activation of Wnt target genes such as *c-Jun*, *c-Myc*, *fibronectin*, and *cyclin D1* (He et al. 1998; Mann et al. 1999; Orsulic et al. 1999; Tetsu and McCormick 1999; Lin, S. Y. et al. 2000; Liu, S. et al. 2005; Clevers 2006). Binding of Wnt proteins, a family of secreted proteins, to Frizzled receptors results in the cytoplasmic accumulation of β -catenin (Schweizer and Varmus 2003). In the absence of Wnt signaling, β -catenin forms a multi-protein complex with glycogen synthase kinase 3 β (GSK3 β), adenomatous polyposis coli, casein kinase1 α , and axin (Takahashi-Yanaga and Sasaguri 2008). When β -catenin is phosphorylated at Ser33/Ser37/Thr41 by GSK3 β , it is immediately subject to ubiquitin-proteasome degradation (Liu, C. et al. 2002; Takahashi-Yanaga and Sasaguri 2008).

The link between Wnt/ β -catenin and PI3K/Akt pathway has been established by several studies. Activated Akt (i.e., phospho-Akt Ser473) was shown to be able to phosphorylate Ser9 on GSK3 β , which may decrease the activity of GSK3 β , thereby stabilizing β -catenin (Yost et al. 1996; Pap and Cooper 1998; Cohen, P. and Frame 2001). Furthermore, Korkaya et al. demonstrated that PI3K/Akt pathway is important in regulating the mammary stem/progenitor cells by promoting β -catenin downstream events through phosphorylation of GSK3 β (Korkaya et al. 2009).

2.2 Hedgehog pathway

Another critical pathway that is involved in CSC self-renewal is hedgehog signaling pathway (Cohen, M. M., Jr. 2003; Liu, S. et al. 2006; Clement et al. 2007; Charafe-Jauffret et al. 2008). For instance, Liu et al. have demonstrated that the hedgehog pathway plays a crucial role in regulating self-renewal of normal and malignant human mammary stem cells by utilizing both *in vitro* and mouse model systems (Liu, S. et al. 2006). Another recent study revealed the essential role of hedgehog-Gli signaling in controlling the self-renewal behavior of human glioma CSCs and tumorigenicity (Clement et al. 2007).

In the absence of hedgehog ligands (Sonic Hedgehog, Desert Hedgehog, and Indian Hedgehog), their transmembrane receptor Patched (Ptch) associates with Smoothened (Smo) and blocks Smo function (Cohen, M. M., Jr. 2003; Lewis and Veltmaat 2004; Liu, S. et al. 2005). When secreted hedgehog ligands bind to Ptch, Smo is released, triggering dissociation of transcription factors, Gli1, Gli2, and Gli3 from Fused (Fu) and suppressor of Fused (SuFu), leading to transcription of an array of genes, such as *cyclin D*, *cyclin E*, *Myc*, and elements of EGF pathway (Cohen, M. M., Jr. 2003; Pasca di Magliano and Hebrok 2003; Lewis and Veltmaat 2004; Liu, S. et al. 2005).

Sonic hedgehog pathway is also linked to transcription factor NF- κ B signaling. NF- κ B was suggested to be a prominent factor in controlling tumor growth and apoptosis resistance of pancreatic CSCs (Kallifatidis et al. 2009). Over-expression of sonic hedgehog is activated by NF- κ B in pancreatic cancer and pancreatic cancer cell proliferation is accelerated by NF- κ B in part through sonic hedgehog over-expression (Nakashima et al. 2006). Kasperczyk et al. further characterized sonic hedgehog as a novel NF- κ B target gene and mapped minimal NF- κ B consensus site to position +139 of sonic hedgehog promoter (Kasperczyk et al. 2009).

2.3 Notch pathway

Notch signaling is known to control cell proliferation and apoptosis to modulate the development of many organs (Wang, Z. et al. 2009). A number of recent studies have demonstrated that Notch-activated genes and pathways can drive tumor growth through the expansion of CSCs (Wilson and Radtke 2006; Charafe-Jauffret et al. 2008; Fan and Eberhart 2008; Kakarala and Wicha 2008; Peacock and Watkins 2008; Scoville et al. 2008; Wang, Z. et al. 2009). Notch pathway is believed to be dysregulated in CSCs, ultimately leading to uncontrolled CSC self-renewal (Wang, Z. et al. 2009). For example, Notch pathway was shown to play an important role in the self-renewal function of malignant breast cancer CSCs (Dontu et al. 2004; Farnie and Clarke 2007).

Five Notch proteins, Notch-1 to Notch-4, have been identified to express as transmembrane receptors in a variety of stem/progenitor cells (Mumm and Kopan 2000). Binding of surface-bound ligands (Jagged1, Jagged2, Delta-like1, Delta-like3, and Delta-like4) triggers serial cleavage events at the Notch proteins by ADAM protease family and γ -secretase (Wu, J. Y. and Rao 1999; Mumm and Kopan 2000; Borggrefe and Oswald 2009). Subsequently, the intracellular domain of Notch is released and translocates into the nucleus, where it acts as a transcription co-activator of recombination signal sequence-binding protein κ (RBP-J) to activate downstream target genes, e.g., *c-Myc*, *cyclin D1*, *p21*, *NF- κ B* (Oswald et al. 1998; Rangarajan et al. 2001; Ronchini and Capobianco 2001; Satoh et al. 2004; Palomero et al. 2006; Weng et al. 2006; Borggrefe and Oswald 2009).

Notch1 has been reported to be the upstream regulator of NF- κ B pathway in diverse cellular situations (Oswald et al. 1998; Wang, J. et al. 2001; Nickoloff et al. 2002; Dontu et al. 2004; Jang et al. 2004; Wang, Y. et al. 2004; Shin et al. 2006; Wang, Z. et al. 2006; Chen et al. 2007). Specifically, Notch-1 is necessary for expression of several NF- κ B subunits (Cheng et al. 2001; Jang et al. 2004) and stimulates NF- κ B promoter activity (Jang et al. 2004).

3. Sulforaphane

Numerous studies have substantiated the chemopreventive properties of high consumption of cruciferous vegetables, especially broccoli and broccoli sprouts, against various types of

cancer (Clarke et al. 2008). Cruciferous vegetables are characterized by their high content of glucosinolates (Herr and Buchler 2010; Fahey et al. 2001), which are converted to isothiocyanates by the action of myrosinase. The chemopreventive effects have been mostly attributed to the activity of these isothiocyanates (Zhang et al. 1992; Clarke et al. 2008). In particular, sulforaphane (Figure 1) is converted from glucoraphanin, the principal glucosinolate in broccoli and broccoli sprouts (Fahey et al. 1997). Sulforaphane has been shown to be not only effective in preventing chemically induced cancers in animal models, including colon, lung, breast, pancreatic, skin and stomach cancer (Zhang et al. 1994; Fahey et al. 1997; Chung et al. 2000; Fahey et al. 2002; Conaway et al. 2005; Gills et al. 2006; Kuroiwa et al. 2006; Xu et al. 2006), but also inhibit the growth of established tumors (Jackson and Singletary 2004; Singh, A. V. et al. 2004).




Fig. 1. Chemical structure of sulforaphane

Early research focused on inhibition of Phase 1 metabolism enzymes that convert procarcinogens to carcinogens and induction of Phase 2 metabolism enzymes that enhance elimination and excretion of carcinogens by sulforaphane (Clarke et al. 2008), which enhances the detoxification of carcinogens and “blocks” carcinogenesis at the initiation stage of cancer (Juge et al. 2007; Clarke et al. 2008).

Subsequent studies suggest that sulforaphane provides protection against tumor development during the “post-initiation” phase by modulating diverse cellular activities including apoptosis, cell cycle, angiogenesis and metastasis (Zhang and Tang 2007; Clarke et al. 2008). Sulforaphane affects classical molecular targets involved in the apoptosis pathways such as down-regulation of anti-apoptotic Bcl-2 and Bcl-X_L, up-regulation of pro-apoptotic Bax expression, proteolytic activation of caspase-3, and the degradation/cleavage of poly(ADP-ribose) polymerase, induction of apoptotic protease activating factor-1 (Choi et al. 2007; Park, S. Y. et al. 2007). The ability of sulforaphane to induce cell cycle arrest is associated with regulation of many molecules including cyclins, Cdks, and p21 (Singh, S. V. et al. 2004; Herman-Antosiewicz et al. 2007; Zhang and Tang 2007). More recent studies demonstrate that sulforaphane is also capable of suppressing angiogenesis and metastasis, which are associated with transcriptional down-regulation of vascular endothelial growth factor (VEGF), hypoxia-inducible factor-1α (HIF-1α), c-Myc and matrix metalloproteinase-2 (MMP-2), and MMP-9 (Bertl et al. 2006; Zhang and Tang 2007).

Our studies have shown that sulforaphane is effective in targeting breast CSCs *in vitro* and *in vivo* (Li et al. 2010). Mammosphere culture was first used to isolate and expand mammary stem/progenitor cells by Dontu et al., based on the ability of stem/progenitor cells to grow in serum-free, non-adherent suspension as spherical clusters of cells while differentiated cells fail to survive under the same condition (Dontu et al. 2003). By employing this technique, we demonstrated that sulforaphane (0.5-5 μM) significantly suppressed the mammosphere formation of both SUM159 and MCF7 cells. A decrease in the number of sphere-forming cells in the 2nd and 3rd passages indicated a reduced self-renewal capacity of these stem/progenitor cells (Dontu et al. 2003). In breast carcinomas, a cell population

with high aldehyde dehydrogenase (ALDH) activity as assessed by the Aldefluor assay has been demonstrated to enrich tumorigenic stem/progenitor cells (Ginestier et al. 2007). This cell population is capable of self-renewal and generating tumors resembling the parental tumor (Ginestier et al. 2007). We found that sulforaphane (1-5 μ M) was able to significantly decrease the tumor-initiating ALDH-positive cell population of SUM159 by 65% to 80% *in vitro*. Of special note, concentrations of sulforaphane which inhibit stem/progenitor cells in both mammosphere formation assay and Aldefluor assay had only minimal effects on the bulk population of breast cancer cell lines, implying that sulforaphane is likely to preferentially target stem/progenitor cells compared to the differentiated cancer cells.

We further demonstrated that sulforaphane can inhibit breast CSCs *in vivo* (Li et al. 2010). The injection of human breast cancer cells into the mammary fat pad of immune-deficient NOD/SCID mice provides a reliable and sensitive *in vivo* system for studying human breast cancer (Al-Hajj et al. 2003; Dick 2003). Daily injection of sulforaphane for two weeks suppressed tumor growth in primary NOD/SCID mice and reduced ALDH-positive cell population of the tumors by more than 50%. Most importantly, we found that recipient NOD/SCID mice inoculated with tumor cells derived from sulforaphane-treated primary xenografts largely failed to develop tumor re-growth up to 33 days, whereas control tumor cells quickly initiated new tumors upon re-implantation. These results suggest that sulforaphane is able to eliminate breast CSCs *in vivo*, thereby abrogating tumor re-growth after re-implantation of primary tumor cells into the secondary mice.

We also observed a down-regulation of Wnt/ β -catenin self-renewal pathway in sulforaphane-treated breast cancer cells (Li et al. 2010). Park et al. previously reported that β -catenin was down-regulated by sulforaphane in human cervical carcinoma HeLa and hepatocarcinoma HepG2 cells (Park, S. Y. et al. 2007). In consistent with their study, we showed that sulforaphane was able to down-regulate Wnt/ β -catenin self-renewal pathway in breast cancer cells, and sulforaphane-induced β -catenin phosphorylation (Ser33/Ser37/Thr41) and proteasome degradation was possibly through activation of GSK3 β . The down-regulation of Wnt/ β -catenin self-renewal pathway might contribute to the inhibitory effects of sulforaphane on breast CSCs. Further studies are warranted to establish the conclusive role of this down-regulation in inhibition of breast CSCs by sulforaphane.

In addition, our recent work has revealed a new molecular target of sulforaphane. Sulforaphane inhibits heat shock protein 90 (Hsp90) function by blocking the interaction of Hsp90 with its cochaperone p50^{Cdc37}, and we traced this activity to a novel interaction site of Hsp90, which fundamentally differs from the mechanism of other Hsp90 inhibitors (unpublished data). LC-MS peptide mapping identified a covalent adduct of sulforaphane with a short peptide IDIIPNPQER in Hsp90 N-terminal domain. NMR experiment with full-length Hsp90 revealed sulforaphane interaction in sheet 2 and the adjacent loop in Hsp90 N-terminal domain, in which this short peptide resides. Akt is a well-known Hsp90 client protein. Our and several other studies have reported the activity of sulforaphane to down-regulate the protein level of Akt and Akt pathway in ovarian, prostate, and colorectal cancers (Chaudhuri et al. 2007; Shen et al. 2007; Shankar et al. 2008). PI3K/Akt pathway was recently demonstrated to play an important role in regulating breast stem/progenitor cells by promoting β -catenin down-stream events through phosphorylation of GSK3 β (Korkaya et al. 2009). Therefore, inhibition of Hsp90 chaperone function by sulforaphane may contribute to the effect on Akt/GSK3 β / β -catenin pathway.

The resistance of pancreatic cancer toward TRAIL (tumor necrosis factor-related apoptosis-inducing ligand) is due to TRAIL-activated NF- κ B signaling (Ibrahim et al. 2001). Kallifatidis et al. demonstrated that sulforaphane (10 μ M) was able to deplete pancreatic CSCs (CD44⁺CD24⁻) by interfering with NF- κ B binding and abrogating apoptosis resistance (Kallifatidis et al. 2009). They found that the presence of pancreatic TICs correlated with apoptosis resistance towards TRAIL due to the enhanced binding of NF- κ B complexes to DNA. Sulforaphane alone or in combined with TRAIL reduced growth of TIC^{high} tumors *in vivo* without toxicity to normal tissue.

Like TRAIL, sorafenib was also observed to strongly up-regulate NF- κ B activity (Rausch et al. 2010). Sulforaphane (10 μ M) completely abolished sorafenib-induced NF- κ B binding in CSC^{high} cells, thereby synergistically inhibiting pancreatic CSC (CD44⁺CD24⁻) (Rausch et al. 2010). The growth of pancreatic CSC^{high} tumor xenografts was synergistically inhibited by combination of sulforaphane and sorafenib, which involved induction of apoptosis, inhibition of proliferation and angiogenesis, as well as down-regulation of epithelial-mesenchymal transition (EMT) related proteins (vimentin, Zeb-1, and Twist-2). EMT induction in cancer cells results in the acquisition of invasive and metastatic properties (Singh, A. and Settleman 2010; Gupta, P. B. et al. 2009; Klarmann et al. 2009; Sarkar et al. 2009). CSCs undergoing metastasis usually express EMT markers (Tang et al. 2010).

This same group then combined sulforaphane (5 μ M) with several chemotherapeutic drugs and observed increased cytotoxicity toward pancreatic CSCs (CD44⁺CD24⁻) (Kallifatidis et al. 2011). Sulforaphane not only down-regulated basal Notch-1 expression in CSC^{high} cells, but also prevented the gemcitabine-induced Notch-1 up-regulation. There was no tumor growth in mice re-implanted with tumor cells derived from sulforaphane-treated or combination-treated xenografts.

Another recent study also examined the molecular mechanisms by which sulforaphane inhibits growth and induces apoptosis of pancreatic CSCs (Srivastava et al. 2011). They demonstrated that sulforaphane (5-10 μ M) inhibited self-renewal capacity of pancreatic CSCs (CD44⁺CD24⁺ESA⁺). Sulforaphane induced apoptosis by inhibiting the expression of Bcl-2 and XIAP, phosphorylation of FKHR, and activating caspase-3. Moreover, sulforaphane inhibited expression of EMT markers (β -catenin, vimentin, Twist-1, and Zeb-1), suggesting the blockade of early metastasis signaling.

In summary, all of these findings strongly support that combination of sulforaphane or even broccoli/broccoli sprout preparations with chemotherapy may be a promising strategy to eradicate tumors and improve patient survival in different types of cancer.

4. Curcumin

Curcumin (Figure 2) is a dietary polyphenol present in the Indian spice turmeric, which is produced from rhizome of the plant *Curcuma longa* and usually used in preparation of mustard and curry (Park, C. H. et al. 2005). Curcumin has been studied as a chemoprevention agent in several cancer models (Mukhopadhyay et al. 2001; Shao et al. 2002; Lin, J. K. 2007; Anand et al. 2008; Kunnumakkara et al. 2008; Strimpakos and Sharma 2008).

Curcumin has been shown to regulate many cellular pathways (Lin, J. K. 2007; Hatcher et al. 2008; Kunnumakkara et al. 2008; Sa and Das 2008; Ravindran et al. 2009), some of which are associated with self-renewal signaling. Curcumin was suggested to induce caspase-3-mediated cleavage of β -catenin, leading to inactivation of Wnt/ β -catenin signaling in




Fig. 2. Chemical structure of curcumin

HCT116 intestinal cancer cells (Jaiswal et al. 2002). The work of Park et al. strengthened the point that curcumin decreased β -catenin/TCF transcription activity in all tested cancer cell lines, including gastric, colon, and intestinal cancer cells, which was attributed to the reduced amount of nuclear β -catenin and TCF-4 proteins (Park, C. H. et al. 2005). Moreover, analysis of gene transcription profile revealed that the expression of Wnt receptor Frizzled-1 was potently suppressed by curcumin (Yan et al. 2005). Curcumin was also shown to be able to attenuate response of β -catenin to Wnt-3a in colon cancer cells through down-regulation of p300, a positive regulator of Wnt/ β -catenin signaling (Ryu et al. 2008). In addition, Wang and his colleagues demonstrated that curcumin down-regulated Notch-1 mRNA level in pancreatic cancer cells, indicating a transcriptional inactivation of Notch-1 by curcumin (Wang, Z. et al. 2006). AP-1 and NF- κ B signaling pathways were shown to be inhibited by curcumin in glioblastoma cells (Dhandapani et al. 2007). Curcumin-induced inactivation of NF- κ B DNA-binding activity was potentially mediated by Notch-1 signaling pathway (Wang, Z. et al. 2006).

Kakarala et al. demonstrated that curcumin (5-10 μ M) was able to target breast stem/progenitor cells, as evidenced by suppressed mammosphere formation along serial passage and a decrease in the percentage of ALDH1-positive cells (Kakarala et al. 2009). Similar to sulforaphane, the concentrations of curcumin inhibiting mammosphere formation was much lower compared to the concentrations of curcumin having impact on differentiated cells. Results from serial passaging suggest that curcumin interferes with breast CSC self-renewal. By utilizing a TCF-LEF reporter assay system in MCF7 cells, the authors confirmed that the effect of curcumin on breast cancer stem/progenitor cells was mediated through its potent inhibitory effect on Wnt/ β -catenin signaling (Kakarala et al. 2009). These results support the work in other systems showing the ability of curcumin to inhibit Wnt signaling (Jaiswal et al. 2002; Ryu et al. 2008; Prasad et al. 2009). The effects of curcumin was further potentiated by piperine, another dietary polyphenol isolated from black and long peppers (Kakarala et al. 2009). Piperine was suggested to enhance the bioavailability of curcumin through inhibition of P glycoprotein-mediated efflux of curcumin (Shoba et al. 1998; Chearwae et al. 2004; Anand et al. 2007; Limtrakul et al. 2007). Curcumin and piperine, alone or in combination did not cause toxicity to differentiated cells (Kakarala et al. 2009).

Side population (SP) cells, first identified for isolation of murine hematopoietic stem cells from bone marrow (Goodell et al. 1996; Zhou, S. et al. 2001; Hirschmann-Jax et al. 2004), can be used to enrich CSCs (Hadrnagy et al. 2006; Wu, C. and Alman 2008). Curcumin inhibited SP of the rat C6 glioma at low concentration (5 μ M) that had minimal effect on proliferation of C6 cells (Fong et al. 2010). Very recently, a polymeric nanoparticle formulation of curcumin (5-20 μ M) was shown to significantly inhibit clonogenicity and depleted the CD133⁺ stem-

like cell population from brain tumor cultures (Lim et al. 2011). They also found that Gli1 and Ptch1B, two key components of hedgehog signaling, were significantly reduced in embryonal tumor derived cell line DAOY after curcumin treatment.

5. Epigallocatechin-3-gallate (EGCG)

Green tea is one of the most widely consumed beverages in the world. Epidemiological studies suggest an association between green tea consumption and chemopreventive effects against skin, lung, breast, colon, liver, stomach, and prostate cancers (Yang, C. S. et al. 2002; Landis-Piwowar et al. 2007). The various polyphenolic catechins contained in green tea are thought to contribute to its chemoprevention activity.




Fig. 3. Chemical structure of EGCG

In particular, numerous studies indicate that EGCG (Figure 3), the most abundant catechin in green tea, is the primary component for these activities (Fujiki 1999; Nagle et al. 2006). *In vitro* and *in vivo* studies have shown that EGCG modulates a wide array of molecular pathways, resulting in induction of apoptosis and cell cycle arrest, and inhibition of invasion, angiogenesis, and metastasis (Shankar et al. 2007; Shankar et al. 2008).

Some studies have found that EGCG may directly or indirectly affect CSC self-renewal pathways. The basal NF- κ B activity and ATP- or IL-1 β induced activation of NF- κ B were negatively regulated by EGCG (Ahmad et al. 2000; Afaq et al. 2003; Guo, S. et al. 2006; Kim, S. J. et al. 2007; Sarkar et al. 2009). EGCG suppressed Akt activation in both colon cancer cell lines and *in vivo* mouse models (Ju et al. 2005; Shimizu et al. 2005; Peng et al. 2006; Bose et al. 2007). In our previous study, EGCG was shown to inhibit the chaperoning function of Hsp90 by impairing the interaction between Hsp90 with its co-chaperones in pancreatic cancer cells, thereby down-regulating Hsp90 client proteins including Akt (Li et al. 2009). EGCG blocked Wnt signaling by stabilizing mRNA of HBP1, a suppressor of Wnt signaling, thereby reducing breast cancer cell tumorigenic proliferation as well as invasiveness (Kim, J. et al. 2006; Kawasaki et al. 2008). The nuclear import of β -catenin was decreased in adenomas isolated from EGCG-treated Apc^{Min/+} mice, a widely used transgenic model recapitulating human colon cancer that bears an Adenomatous Polyposis Coli (APC) gene mutation (Ju et al. 2005; Bose et al. 2007).

Combination of EGCG and doxorubicin was suggested to eradicate putative prostate CSCs (CD44⁺) (Stearns et al. 2010). EGCG (30 and 60 μ M), either alone or in combination with doxorubicin, reduced the colony-forming capability of human prostate cancer cell line PC-3ML. Relatively low dose of EGCG (57 mg/kg) in combination with nontoxic, sub-therapeutic dosages of doxorubicin can eradicate established prostate tumors derived from CD44^{high} tumor-initiating cells isolated from PCa-20a cells in NOD/SCID mice.

Tang et al. have shown that EGCG either alone or in combination with quercetin can eliminate prostate CSC characteristics (Tang et al. 2010). EGCG inhibited the growth and self-renewal capacity of CD44⁺CD133⁺ CSCs contained in human prostate cancer cell lines and CD44⁺ α 2 β 1⁺CD133⁺ CSCs isolated from human primary prostate tumors, as measured by spheroid and colony formation assay. They also suggested that EGCG was able to induce apoptosis in prostate CSCs. In addition, EGCG was found to suppress EMT by inhibiting the expression of vimentin and nuclear β -catenin, as well as the transcription factors slug and snail which are required for EMT induction. The inhibition of EMT markers by EGCG could retard early metastasis of prostate CSCs.

6. Quercetin

Quercetin (3,3',4',5,7-pentahydroxyflavone) (Figure 4) is a ubiquitous plant polyphenol, naturally occurring in most edible fruits and vegetables, with the high levels being found in apples, cranberries, and blueberries (Androutsopoulos et al. 2010; Guo, W. et al. 2009). Many studies have demonstrated that quercetin possess anti-oxidant, anti-inflammatory and anti-cancer activities (Williamson and Manach 2005; Guo, W. et al. 2009). Quercetin has also been shown to enhance the anti-cancer effects of several chemotherapeutic drugs (Borska et al. 2010; Du et al. 2010; Shih et al. 2010; Wong and Chiu 2010; Limtrakul et al. 2005; Du et al. 2009).




Fig. 4. Chemical structure of quercetin

Several studies have indicated that quercetin may modulate self-renewal pathways. Quercetin was suggested to be a potent inhibitor of β -catenin/TCF signaling in SW480 colon cancer cells, and the reduced β -catenin/TCF transcriptional activity was due to the decreased nuclear β -catenin and TCF-4 proteins (Park, C. H. et al. 2005). The inhibition of colon cancer cell growth by quercetin was related to the inhibition of cyclin D1 and surviving expression through Wnt/ β -catenin signaling pathway (Shan et al. 2009).

Zhou et al. demonstrated that quercetin (100-400 μ M) mediated reduction of self-renewal capacity of pancreatic CSCs, decreased ALDH1 activity, overcame apoptosis resistance of pancreatic CSCs (CD44⁺CD24⁻), and diminished the expression of proteins involved in the EMT (vimentin and Twist-2) in CSC^{high} cells (Zhou, W. et al. 2010). Quercetin strongly reduced rapid growth of CSC-enriched xenografts, while no toxic side effects were observed (Zhou, W. et al. 2010). They further demonstrated that combination of quercetin with sulforaphane led to a synergistic reduction in self-renewal capacity and a complete abrogation of tumor growth in xenograft mouse model. Similarly, in another recent study, quercetin enhanced the inhibitory effects of sulforaphane on self-renewal capacity of pancreatic CSCs (CD44⁺CD24⁺ESA⁺) (Srivastava et al. 2011). Moreover, quercetin (20 μ M) was found to synergize with green tea EGCG in inhibiting prostate CSCs (Tang et al. 2010). Quercetin not only potentiated the inhibitory effects of EGCG on self-renewal, migration and invasion capacities of prostate CSCs isolated from primary tumors, but also synergized with EGCG to induce apoptosis (Tang et al. 2010).

Natural Dietary Compound	Food Origins	Cancer Stem Cells	Potential Molecular Targets
Sulforaphane	Cruciferous vegetables	Pancreatic cancer Breast cancer	Wnt/ β -catenin; NF- κ B binding; EMT markers; Notch-1
Curcumin	Turmeric	Breast cancer Brain tumor	Wnt/ β -catenin; Gli1 and Ptch1B
EGCG	Green tea	Prostate cancer	β -catenin; EMT markers; slug and snail
Quercetin	Ubiquitous, e.g., apple, cranberry, blueberry	Pancreatic cancer Prostate cancer Lung cancer	EMT markers
Piperine	Black and long pepper	Breast cancer	Wnt/ β -catenin; NF- κ B
Genistein	Soy		GSK3 β , β -catenin, Wnt-5a; Notch-2
Resveratrol	Grapes, berries, plums, and peanuts		β -catenin, GSK3 β ; Notch-1
Lycopene	Tomatoes, watermelon, papaya, pink grapefruit		β -catenin
Vitamin D3	Fish, egg yolk, beef liver		TCF-4, E-cadherin

Table 1. Natural dietary components that potentially regulate self-renewal pathways and inhibit CSCs

7. Conclusion

Naturally-occurring dietary components are advantageous in several aspects as chemoprevention agents: (1) they are present in commonly consumed food, which is readily available to most people in daily life; (2) they usually have very low or no toxicity, in contrast to most chemotherapy drugs; (3) many of these compounds have shown potential as an adjunct to chemotherapy drugs in some clinical trials. Although the reports were very limited for dietary components to inhibit CSCs, many of them have been shown to be directly or indirectly involved in modulation of CSC self-renewal pathways. All of these studies stress the need for investigating the efficacy of dietary components against CSCs and elucidating the mechanisms of action. In Table 1, we summarize the compounds discussed in this chapter as well as some others dietary components that may affect the element(s) of self-renewal pathways.

Since CSCs are more resistant to conventional therapies in comparison with differentiated cells constituting the tumor bulk, these studies will provide strong rationale for preclinical and clinical evaluation of the dietary components or potentially their native food extracts combined with chemotherapy. Combination of dietary intervention that are directed against CSCs and conventional chemotherapy would have the potential to eliminate CSCs, overcome tumor resistance, reduce recurrence, and eventually improve patient survival.

8. References

- Afaq, F., V. M. Adhami, N. Ahmad & H. Mukhtar (2003). Inhibition of ultraviolet B-mediated activation of nuclear factor kappaB in normal human epidermal keratinocytes by green tea Constituent (-)-epigallocatechin-3-gallate. *Oncogene*, Vol.22, No.7, pp. 1035-44, ISSN 0950-9232
- Ahmad, N., S. Gupta & H. Mukhtar (2000). Green tea polyphenol epigallocatechin-3-gallate differentially modulates nuclear factor kappaB in cancer cells versus normal cells. *Arch Biochem Biophys*, Vol.376, No.2, pp. 338-46, ISSN 0003-9861
- Akiyama, T. (2000). Wnt/beta-catenin signaling. *Cytokine Growth Factor Rev*, Vol.11, No.4, pp. 273-82, ISSN 1359-6101
- Al-Hajj, M., M. S. Wicha, A. Benito-Hernandez, S. J. Morrison & M. F. Clarke (2003). Prospective identification of tumorigenic breast cancer cells. *Proc Natl Acad Sci U S A*, Vol.100, No.7, pp. 3983-8, ISSN 0027-8424
- Anand, P., A. B. Kunnumakkara, R. A. Newman & B. B. Aggarwal (2007). Bioavailability of curcumin: problems and promises. *Mol Pharm*, Vol.4, No.6, pp. 807-18, ISSN 1543-8384
- Anand, P., C. Sundaram, S. Jhurani, A. B. Kunnumakkara & B. B. Aggarwal (2008). Curcumin and cancer: an "old-age" disease with an "age-old" solution. *Cancer Lett*, Vol.267, No.1, pp. 133-64, ISSN 0304-3835
- Androutsopoulos, V. P., A. Papakyriakou, D. Vourloumis, A. M. Tsatsakis & D. A. (2010). Spandidos Dietary flavonoids in cancer therapy and prevention: substrates and inhibitors of cytochrome P450 CYP1 enzymes. *Pharmacol Ther*, Vol.126, No.1, pp. 9-20, ISSN 1879-016X

- Bertl, E., H. Bartsch & C. Gerhauser (2006). Inhibition of angiogenesis and endothelial cell functions are novel sulforaphane-mediated mechanisms in chemoprevention. *Mol Cancer Ther*, Vol.5, No.3, pp. 575-85, ISSN 1535-7163
- Borggreffe, T. & F. Oswald (2009). The Notch signaling pathway: transcriptional regulation at Notch target genes. *Cell Mol Life Sci*, Vol.66, No.10, pp. 1631-46, ISSN 1420-9071
- Borska, S., M. Sopel, M. Chmielewska, M. Zabel & P. Dziegiel (2010). Quercetin as a potential modulator of P-glycoprotein expression and function in cells of human pancreatic carcinoma line resistant to daunorubicin. *Molecules*, Vol.15, No.2, pp. 857-70, ISSN 1420-3049
- Bose, M., X. Hao, J. Ju, A. Husain, S. Park, J. D. Lambert & C. S. Yang (2007). Inhibition of tumorigenesis in ApcMin/+ mice by a combination of (-)-epigallocatechin-3-gallate and fish oil. *J Agric Food Chem*, Vol.55, No.19, pp. 7695-700, ISSN 0021-8561
- Charafe-Jauffret, E., F. Monville, C. Ginestier, G. Dontu, D. Birnbaum & M. S. Wicha (2008). Cancer stem cells in breast: current opinion and future challenges. *Pathobiology*, Vol.75, No.2, pp. 75-84, ISSN 1423-0291
- Chaudhuri, D., S. Orsulic & B. T. Ashok (2007). Antiproliferative activity of sulforaphane in Akt-overexpressing ovarian cancer cells. *Mol Cancer Ther*, Vol.6, No.1, pp. 334-45, ISSN 1535-7163
- Chearwae, W., S. Anuchapreeda, K. Nandigama, S. V. Ambudkar & P. Limtrakul (2004). Biochemical mechanism of modulation of human P-glycoprotein (ABCB1) by curcumin I, II, and III purified from Turmeric powder. *Biochem Pharmacol*, Vol.68, No.10, pp. 2043-52, ISSN 0006-2952
- Chen, Y., W. Shu, W. Chen, Q. Wu, H. Liu & G. Cui (2007). Curcumin, both histone deacetylase and p300/CBP-specific inhibitor, represses the activity of nuclear factor kappa B and Notch 1 in Raji cells. *Basic Clin Pharmacol Toxicol*, Vol.101, No.6, pp. 427-33, ISSN 1742-7843
- Cheng, P., A. Zlobin, V. Volgina, S. Gottipati, B. Osborne, E. J. Simel, L. Miele & D. I. Gabrilovich (2001). Notch-1 regulates NF-kappaB activity in hemopoietic progenitor cells. *J Immunol*, Vol.167, No.8, pp. 4458-67, ISSN 0022-1767
- Chien, A. J., E. C. Moore, A. S. Lonsdorf, R. M. Kulikauskas, B. G. Rothberg, A. J. Berger, M. B. Major, S. T. Hwang, D. L. Rimm & R. T. Moon (2009). Activated Wnt/beta-catenin signaling in melanoma is associated with decreased proliferation in patient tumors and a murine melanoma model. *Proc Natl Acad Sci U S A*, Vol.106, No.4, pp. 1193-8, ISSN 1091-6490
- Chinni, S. R., Y. Li, S. Upadhyay, P. K. Koppolu & F. H. Sarkar (2001). Indole-3-carbinol (I3C) induced cell growth inhibition, G1 cell cycle arrest and apoptosis in prostate cancer cells. *Oncogene*, Vol.20, No.23, pp. 2927-36, ISSN 0950-9232
- Choi, S., K. L. Lew, H. Xiao, A. Herman-Antosiewicz, D. Xiao, C. K. Brown & S. V. Singh (2007). D,L-Sulforaphane-induced cell death in human prostate cancer cells is regulated by inhibitor of apoptosis family proteins and Apaf-1. *Carcinogenesis*, Vol.28, No.1, pp. 151-62, ISSN 0143-3334
- Choudhuri, T., S. Pal, M. L. Agwarwal, T. Das & G. Sa (2002). Curcumin induces apoptosis in human breast cancer cells through p53-dependent Bax induction. *FEBS Lett*, Vol.512, No.1-3, pp. 334-40, ISSN 0014-5793

- Chung, F. L., C. C. Conaway, C. V. Rao & B. S. Reddy (2000). Chemoprevention of colonic aberrant crypt foci in Fischer rats by sulforaphane and phenethyl isothiocyanate. *Carcinogenesis*, Vol.21, No.12, pp. 2287-91, ISSN 0143-3334
- Clarke, J. D., R. H. Dashwood & E. Ho (2008). Multi-targeted prevention of cancer by sulforaphane. *Cancer Lett*, Vol.269, No.2, pp. 291-304, ISSN 1872-7980
- Clement, V., P. Sanchez, N. de Tribolet, I. Radovanovic & A. Ruiz i Altaba (2007). HEDGEHOG-GLI1 signaling regulates human glioma growth, cancer stem cell self-renewal, and tumorigenicity. *Curr Biol*, Vol.17, No.2, pp. 165-72, ISSN 0960-9822
- Clevers, H. (2006). Wnt/beta-catenin signaling in development and disease. *Cell*, Vol.127, No.3, pp. 469-80, ISSN 0092-8674
- Cohen, M. M., Jr. (2003). The hedgehog signaling network. *Am J Med Genet A*, Vol.123A, No.1, pp. 5-28, ISSN 1552-4825
- Cohen, P. & S. Frame (2001). The renaissance of GSK3. *Nat Rev Mol Cell Biol*, Vol.2, No.10, pp. 769-76, ISSN 1471-0072
- Conaway, C. C., C. X. Wang, B. Pittman, Y. M. Yang, J. E. Schwartz, D. Tian, E. J. McIntee, S. S. Hecht & F. L. Chung (2005). Phenethyl isothiocyanate and sulforaphane and their N-acetylcysteine conjugates inhibit malignant progression of lung adenomas induced by tobacco carcinogens in A/J mice. *Cancer Res*, Vol.65, No.18, pp. 8548-57, ISSN 0008-5472
- Dhandapani, K. M., V. B. Mahesh & D. W. Brann (2007). Curcumin suppresses growth and chemoresistance of human glioblastoma cells via AP-1 and NFkappaB transcription factors. *J Neurochem*, Vol.102, No.2, pp. 522-38, ISSN 0022-3042
- Dick, J. E. (2003). Breast cancer stem cells revealed. *Proc Natl Acad Sci U S A*, Vol.100, No.7, pp. 3547-9, ISSN 0027-8424
- Dontu, G., W. M. Abdallah, J. M. Foley, K. W. Jackson, M. F. Clarke, M. J. Kawamura & M. S. Wicha (2003). In vitro propagation and transcriptional profiling of human mammary stem/progenitor cells. *Genes Dev*, Vol.17, No.10, pp. 1253-70, ISSN 0890-9369
- Dontu, G., K. W. Jackson, E. McNicholas, M. J. Kawamura, W. M. Abdallah & M. S. Wicha (2004). Role of Notch signaling in cell-fate determination of human mammary stem/progenitor cells. *Breast Cancer Res*, Vol.6, No.6, pp. R605-15, ISSN 1465-542X
- Du, G., H. Lin, M. Wang, S. Zhang, X. Wu, L. Lu, L. Ji & L. Yu (2009). Quercetin greatly improved therapeutic index of doxorubicin against 4T1 breast cancer by its opposing effects on HIF-1alpha in tumor and normal cells. *Cancer Chemother Pharmacol*, ISSN 1432-0843
- Du, G., H. Lin, Y. Yang, S. Zhang, X. Wu, M. Wang, L. Ji, L. Lu, L. Yu & G. Han (2010). Dietary quercetin combining intratumoral doxorubicin injection synergistically induces rejection of established breast cancer in mice. *Int Immunopharmacol*, Vol.10, No.7, pp. 819-26, ISSN 1878-1705
- Fahey, J. W., X. Haristoy, P. M. Dolan, T. W. Kensler, I. Scholtus, K. K. Stephenson, P. Talalay & A. Lozniewski (2002). Sulforaphane inhibits extracellular, intracellular, and antibiotic-resistant strains of *Helicobacter pylori* and prevents benzo[a]pyrene-induced stomach tumors. *Proc Natl Acad Sci U S A*, Vol.99, No.11, pp. 7610-5, ISSN 0027-8424

- Fahey, J. W., A. T. Zalcmann & P. Talalay (2001). The chemical diversity and distribution of glucosinolates and isothiocyanates among plants. *Phytochemistry*, Vol.56, No.1, pp. 5-51, ISSN 0031-9422
- Fahey, J. W., Y. Zhang & P. Talalay (1997). Broccoli sprouts: an exceptionally rich source of inducers of enzymes that protect against chemical carcinogens. *Proc Natl Acad Sci U S A*, Vol.94, No.19, pp. 10367-72, ISSN 0027-8424
- Fan, X. & C. G. Eberhart (2008). Medulloblastoma stem cells. *J Clin Oncol*, Vol.26, No.17, pp. 2821-7, ISSN 1527-7755
- Farnie, G. & R. B. Clarke (2007). Mammary stem cells and breast cancer--role of Notch signalling. *Stem Cell Rev*, Vol.3, No.2, pp. 169-75, ISSN 1550-8943
- Fong, D., A. Yeh, R. Naftalovich, T. H. Choi & M. M. (2010). Curcumin inhibits the side population (SP) phenotype of the rat C6 glioma cell line: towards targeting of cancer stem cells with phytochemicals. *Cancer Lett*, Vol.293, No.1, pp. 65-72, ISSN 1872-7980
- Fujiki, H. (1999). Two stages of cancer prevention with green tea. *J Cancer Res Clin Oncol*, Vol.125, No.11, pp. 589-97, ISSN 0171-5216
- Gills, J. J., E. H. Jeffery, N. V. Matusheski, R. C. Moon, D. D. Lantvit & J. M. Pezzuto (2006). Sulforaphane prevents mouse skin tumorigenesis during the stage of promotion. *Cancer Lett*, Vol.236, No.1, pp. 72-9, ISSN 0304-3835
- Ginestier, C., M. H. Hur, E. Charafe-Jauffret, F. Monville, J. Dutcher, M. Brown, J. Jacquemier, P. Viens, C. G. Kleer, S. Liu, A. Schott, D. Hayes, D. Birnbaum, M. S. Wicha & G. Dontu (2007). ALDH1 is a marker of normal and malignant human mammary stem cells and a predictor of poor clinical outcome. *Cell Stem Cell*, Vol.1, No.5, pp. 555-67, ISSN 1934-5909
- Goodell, M. A., K. Brose, G. Paradis, A. S. Conner & R. C. Mulligan (1996). Isolation and functional properties of murine hematopoietic stem cells that are replicating in vivo. *J Exp Med*, Vol.183, No.4, pp. 1797-806, ISSN 0022-1007
- Guo, S., J. Lu, A. Subramanian & G. E. Sonenshein (2006). Microarray-assisted pathway analysis identifies mitogen-activated protein kinase signaling as a mediator of resistance to the green tea polyphenol epigallocatechin 3-gallate in her-2/neu-overexpressing breast cancer cells. *Cancer Res*, Vol.66, No.10, pp. 5322-9, ISSN 0008-5472
- Guo, W., E. Kong & M. Meydani (2009). Dietary polyphenols, inflammation, and cancer. *Nutr Cancer*, Vol.61, No.6, pp. 807-10, ISSN 1532-7914
- Gupta, P. B., C. L. Chaffer & R. A. Weinberg (2009). Cancer stem cells: mirage or reality? *Nat Med*, Vol.15, No.9, pp. 1010-2, ISSN 1546-170X
- Gupta, S., T. Hussain & H. Mukhtar (2003). Molecular pathway for (-)-epigallocatechin-3-gallate-induced cell cycle arrest and apoptosis of human prostate carcinoma cells. *Arch Biochem Biophys*, Vol.410, No.1, pp. 177-85, ISSN 0003-9861
- Hadnagy, A., L. Gaboury, R. Beaulieu & D. Balicki (2006). SP analysis may be used to identify cancer stem cell populations. *Exp Cell Res*, Vol.312, No.19, pp. 3701-10, ISSN 0014-4827
- Hambardzumyan, D., M. Squatrito & E. C. Holland (2006). Radiation resistance and stem-like cells in brain tumors. *Cancer Cell*, Vol.10, No.6, pp. 454-6, ISSN 1535-6108

- Hastak, K., S. Gupta, N. Ahmad, M. K. Agarwal, M. L. Agarwal & H. Mukhtar (2003). Role of p53 and NF-kappaB in epigallocatechin-3-gallate-induced apoptosis of LNCaP cells. *Oncogene*, Vol.22, No.31, pp. 4851-9, ISSN 0950-9232
- Hatcher, H., R. Planalp, J. Cho, F. M. Torti & S. V. Torti (2008). Curcumin: from ancient medicine to current clinical trials. *Cell Mol Life Sci*, Vol.65, No.11, pp. 1631-52, ISSN 1420-682X
- He, T. C., A. B. Sparks, C. Rago, H. Hermeking, L. Zawel, L. T. da Costa, P. J. Morin, B. Vogelstein & K. W. Kinzler (1998). Identification of c-MYC as a target of the APC pathway. *Science*, Vol.281, No.5382, pp. 1509-12, ISSN 0036-8075
- Herman-Antosiewicz, A., H. Xiao, K. L. Lew & S. V. Singh (2007). Induction of p21 protein protects against sulforaphane-induced mitotic arrest in LNCaP human prostate cancer cell line. *Mol Cancer Ther*, Vol.6, No.5, pp. 1673-81, ISSN 1535-7163
- Herr, I. & M. W. Buchler (2010). Dietary constituents of broccoli and other cruciferous vegetables: implications for prevention and therapy of cancer. *Cancer Treat Rev*, Vol.36, No.5, pp. 377-83, ISSN 1532-1967
- Hirschmann-Jax, C., A. E. Foster, G. G. Wulf, J. G. Nuchtern, T. W. Jax, U. Gobel, M. A. Goodell & M. K. Brenner (2004). A distinct "side population" of cells with high drug efflux capacity in human tumor cells. *Proc Natl Acad Sci U S A*, Vol.101, No.39, pp. 14228-33, ISSN 0027-8424
- Ibrahim, S. M., J. Ringel, C. Schmidt, B. Ringel, P. Muller, D. Koczan, H. J. Thiesen & M. Lohr (2001). Pancreatic adenocarcinoma cell lines show variable susceptibility to TRAIL-mediated cell death. *Pancreas*, Vol.23, No.1, pp. 72-9, ISSN 0885-3177
- Jackson, S. J. & K. W. Singletary (2004). Sulforaphane: a naturally occurring mammary carcinoma mitotic inhibitor, which disrupts tubulin polymerization. *Carcinogenesis*, Vol.25, No.2, pp. 219-27, ISSN 0143-3334
- Jaiswal, A. S., B. P. Marlow, N. Gupta & S. Narayan (2002). Beta-catenin-mediated transactivation and cell-cell adhesion pathways are important in curcumin (diferuylmethane)-induced growth arrest and apoptosis in colon cancer cells. *Oncogene*, Vol.21, No.55, pp. 8414-27, ISSN 0950-9232
- Jang, M. S., H. Miao, N. Carlesso, L. Shelly, A. Zlobin, N. Darack, J. Z. Qin, B. J. Nickoloff & L. Miele (2004). Notch-1 regulates cell death independently of differentiation in murine erythroleukemia cells through multiple apoptosis and cell cycle pathways. *J Cell Physiol*, Vol.199, No.3, pp. 418-33, ISSN 0021-9541
- Ju, J., J. Hong, J. N. Zhou, Z. Pan, M. Bose, J. Liao, G. Y. Yang, Y. Y. Liu, Z. Hou, Y. Lin, J. Ma, W. J. Shih, A. M. Carothers & C. S. Yang (2005). Inhibition of intestinal tumorigenesis in Apcmin/+ mice by (-)-epigallocatechin-3-gallate, the major catechin in green tea. *Cancer Res*, Vol.65, No.22, pp. 10623-31, ISSN 0008-5472
- Juge, N., R. F. Mithen & M. Traka (2007). Molecular basis for chemoprevention by sulforaphane: a comprehensive review. *Cell Mol Life Sci*, Vol.64, No.9, pp. 1105-27, ISSN 1420-682X
- Kakarala, M., D. E. Brenner, H. Korkaya, C. Cheng, K. Tazi, C. Ginestier, S. Liu, G. Dontu & M. S. Wicha (2009). Targeting breast stem cells with the cancer preventive compounds curcumin and piperine. *Breast Cancer Res Treat*, ISSN 1573-7217
- Kakarala, M. & M. S. Wicha (2008). Implications of the cancer stem-cell hypothesis for breast cancer prevention and therapy. *J Clin Oncol*, Vol.26, No.17, pp. 2813-20, ISSN 1527-7755

- Kallifatidis, G., S. Labsch, V. Rausch, J. Mattern, J. Gladkich, G. Moldenhauer, M. W. Buchler, A. V. Salnikov & I. Herr (2011). Sulforaphane Increases Drug-mediated Cytotoxicity Toward Cancer Stem-like Cells of Pancreas and Prostate. *Mol Ther*, ISSN 1525-0024
- Kallifatidis, G., V. Rausch, B. Baumann, A. Apel, B. M. Beckermann, A. Groth, J. Mattern, Z. Li, A. Kolb, G. Moldenhauer, P. Altevogt, T. Wirth, J. Werner, P. Schemmer, M. W. Buchler, A. V. Salnikov & I. Herr (2009). Sulforaphane targets pancreatic tumour-initiating cells by NF-kappaB-induced antiapoptotic signalling. *Gut*, Vol.58, No.7, pp. 949-63, ISSN 1468-3288
- Kasperczyk, H., B. Baumann, K. M. Debatin & S. Fulda (2009). Characterization of sonic hedgehog as a novel NF-kappaB target gene that promotes NF-kappaB-mediated apoptosis resistance and tumor growth in vivo. *FASEB J*, Vol.23, No.1, pp. 21-33, ISSN 1530-6860
- Kawaguchi-Ihara, N., I. Murohashi, N. Nara & S. Tohda (2008). Promotion of the self-renewal capacity of human acute leukemia cells by Wnt3A. *Anticancer Res*, Vol.28, No.5A, pp. 2701-4, ISSN 0250-7005
- Kawasaki, B. T., E. M. Hurt, T. Mistree & W. L. Farrar (2008). Targeting cancer stem cells with phytochemicals. *Mol Interv*, Vol.8, No.4, pp. 174-84, ISSN 1534-0384
- Khan, N. I., K. F. Bradstock & L. J. Bendall (2007). Activation of Wnt/beta-catenin pathway mediates growth and survival in B-cell progenitor acute lymphoblastic leukaemia. *Br J Haematol*, Vol.138, No.3, pp. 338-48, ISSN 0007-1048
- Kim, J., X. Zhang, K. M. Rieger-Christ, I. C. Summerhayes, D. E. Wazer, K. E. Paulson & A. S. Yee (2006). Suppression of Wnt signaling by the green tea compound (-)-epigallocatechin 3-gallate (EGCG) in invasive breast cancer cells. Requirement of the transcriptional repressor HBP1. *J Biol Chem*, Vol.281, No.16, pp. 10865-75, ISSN 0021-9258
- Kim, S. J., H. J. Jeong, K. M. Lee, N. Y. Myung, N. H. An, W. M. Yang, S. K. Park, H. J. Lee, S. H. Hong, H. M. Kim & J. Y. Um (2007). Epigallocatechin-3-gallate suppresses NF-kappaB activation and phosphorylation of p38 MAPK and JNK in human astrocytoma U373MG cells. *J Nutr Biochem*, Vol.18, No.9, pp. 587-96, ISSN 0955-2863
- Klarmann, G. J., E. M. Hurt, L. A. Mathews, X. Zhang, M. A. Duhagon, T. Mistree, S. B. Thomas & W. L. Farrar (2009). Invasive prostate cancer cells are tumor initiating cells that have a stem cell-like genomic signature. *Clin Exp Metastasis*, Vol.26, No.5, pp. 433-46, ISSN 1573-7276
- Korkaya, H., A. Paulson, E. Charafe-Jauffret, C. Ginestier, M. Brown, J. Dutcher, S. G. Clouthier & M. S. Wicha (2009). Regulation of mammary stem/progenitor cells by PTEN/Akt/beta-catenin signaling. *PLoS Biol*, Vol.7, No.6, pp. e1000121, ISSN 1545-7885
- Kunnumakkara, A. B., P. Anand & B. B. Aggarwal (2008). Curcumin inhibits proliferation, invasion, angiogenesis and metastasis of different cancers through interaction with multiple cell signaling proteins. *Cancer Lett*, Vol.269, No.2, pp. 199-225, ISSN 1872-7980
- Kuroiwa, Y., A. Nishikawa, Y. Kitamura, K. Kanki, Y. Ishii, T. Umemura & M. Hirose (2006). Protective effects of benzyl isothiocyanate and sulforaphane but not resveratrol against initiation of pancreatic carcinogenesis in hamsters. *Cancer Lett*, Vol.241, No.2, pp. 275-80, ISSN 0304-3835

- Lamartiniere, C. A., M. S. Cotroneo, W. A. Fritz, J. Wang, R. Mentor-Marcel & A. Elgavish (2002). Genistein chemoprevention: timing and mechanisms of action in murine mammary and prostate. *J Nutr*, Vol.132, No.3, pp. 552S-558S, ISSN 0022-3166
- Landis-Piwowar, K. R., C. Huo, D. Chen, V. Milacic, G. Shi, T. H. Chan & Q. P. Dou (2007). A novel prodrug of the green tea polyphenol (-)-epigallocatechin-3-gallate as a potential anticancer agent. *Cancer Res*, Vol.67, No.9, pp. 4303-10, ISSN 0008-5472
- Lewis, M. T. & J. M. Veltmaat (2004). Next stop, the twilight zone: hedgehog network regulation of mammary gland development. *J Mammary Gland Biol Neoplasia*, Vol.9, No.2, pp. 165-81, ISSN 1083-3021
- Li, Y., X. Li & F. H. Sarkar (2003). Gene expression profiles of I3C- and DIM-treated PC3 human prostate cancer cells determined by cDNA microarray analysis. *J Nutr*, Vol.133, No.4, pp. 1011-9, ISSN 0022-3166
- Li, Y. & F. H. Sarkar (2002). Gene expression profiles of genistein-treated PC3 prostate cancer cells. *J Nutr*, Vol.132, No.12, pp. 3623-31, ISSN 0022-3166
- Li, Y., B. Welm, K. Podsypanina, S. Huang, M. Chamorro, X. Zhang, T. Rowlands, M. Egeblad, P. Cowin, Z. Werb, L. K. Tan, J. M. Rosen & H. E. Varmus (2003). Evidence that transgenes encoding components of the Wnt signaling pathway preferentially induce mammary cancers from progenitor cells. *Proc Natl Acad Sci U S A*, Vol.100, No.26, pp. 15853-8, ISSN 0027-8424
- Li, Y., T. Zhang, Y. Jiang, H. F. Lee, S. J. Schwartz & D. Sun (2009). (-)-Epigallocatechin-3-gallate inhibits Hsp90 function by impairing Hsp90 association with cochaperones in pancreatic cancer cell line Mia Paca-2. *Mol Pharm*, Vol.6, No.4, pp. 1152-9, ISSN 1543-8384
- Li, Y., T. Zhang, H. Korkaya, S. Liu, H. F. Lee, B. Newman, Y. Yu, S. G. Clouthier, S. J. Schwartz, M. S. Wicha & D. (2010). Sun Sulforaphane, a dietary component of broccoli/broccoli sprouts, inhibits breast cancer stem cells. *Clin Cancer Res*, Vol.16, No.9, pp. 2580-90, ISSN 1078-0432
- Lim, K. J., S. Bisht, E. E. Bar, A. Maitra & C. G. (2011). Eberhart A polymeric nanoparticle formulation of curcumin inhibits growth, clonogenicity and stem-like fraction in malignant brain tumors. *Cancer Biol Ther*, Vol.11, No.5, pp. 464-73, ISSN 1555-8576
- Limtrakul, P., W. Chearwae, S. Shukla, C. Phisalpong & S. V. Ambudkar (2007). Modulation of function of three ABC drug transporters, P-glycoprotein (ABCB1), mitoxantrone resistance protein (ABCG2) and multidrug resistance protein 1 (ABCC1) by tetrahydrocurcumin, a major metabolite of curcumin. *Mol Cell Biochem*, Vol.296, No.1-2, pp. 85-95, ISSN 0300-8177
- Limtrakul, P., O. Khantamat & K. Pintha (2005). Inhibition of P-glycoprotein function and expression by kaempferol and quercetin. *J Chemother*, Vol.17, No.1, pp. 86-95, ISSN 1120-009X
- Lin, J. K. (2007). Molecular targets of curcumin. *Adv Exp Med Biol*, Vol.595, pp. 227-43, ISSN 0065-2598
- Lin, S. Y., W. Xia, J. C. Wang, K. Y. Kwong, B. Spohn, Y. Wen, R. G. Pestell & M. C. Hung (2000). Beta-catenin, a novel prognostic marker for breast cancer: its roles in cyclin D1 expression and cancer progression. *Proc Natl Acad Sci U S A*, Vol.97, No.8, pp. 4262-6, ISSN 0027-8424

- Lippman, M. E. (2000). High-dose chemotherapy plus autologous bone marrow transplantation for metastatic breast cancer. *N Engl J Med*, Vol.342, No.15, pp. 1119-20, ISSN 0028-4793
- Liu, C., Y. Li, M. Semenov, C. Han, G. H. Baeg, Y. Tan, Z. Zhang, X. Lin & X. He (2002). Control of beta-catenin phosphorylation/degradation by a dual-kinase mechanism. *Cell*, Vol.108, No.6, pp. 837-47, ISSN 0092-8674
- Liu, S., G. Dontu, I. D. Mantle, S. Patel, N. S. Ahn, K. W. Jackson, P. Suri & M. S. Wicha (2006). Hedgehog signaling and Bmi-1 regulate self-renewal of normal and malignant human mammary stem cells. *Cancer Res*, Vol.66, No.12, pp. 6063-71, ISSN 0008-5472
- Liu, S., G. Dontu & M. S. Wicha (2005). Mammary stem cells, self-renewal pathways, and carcinogenesis. *Breast Cancer Res*, Vol.7, No.3, pp. 86-95, ISSN 1465-542X
- Mann, B., M. Gelos, A. Siedow, M. L. Hanski, A. Gratchev, M. Ilyas, W. F. Bodmer, M. P. Moyer, E. O. Riecken, H. J. Buhr & C. Hanski (1999). Target genes of beta-catenin-T cell-factor/lymphoid-enhancer-factor signaling in human colorectal carcinomas. *Proc Natl Acad Sci U S A*, Vol.96, No.4, pp. 1603-8, ISSN 0027-8424
- Mukhopadhyay, A., C. Bueso-Ramos, D. Chatterjee, P. Pantazis & B. B. Aggarwal (2001). Curcumin downregulates cell survival mechanisms in human prostate cancer cell lines. *Oncogene*, Vol.20, No.52, pp. 7597-609, ISSN 0950-9232
- Mumm, J. S. & R. Kopan (2000). Notch signaling: from the outside in. *Dev Biol*, Vol.228, No.2, pp. 151-65, ISSN 0012-1606
- Nagle, D. G., D. Ferreira & Y. D. Zhou (2006). Epigallocatechin-3-gallate (EGCG): chemical and biomedical perspectives. *Phytochemistry*, Vol.67, No.17, pp. 1849-55, ISSN 0031-9422
- Nakashima, H., M. Nakamura, H. Yamaguchi, N. Yamanaka, T. Akiyoshi, K. Koga, K. Yamaguchi, M. Tsuneyoshi, M. Tanaka & M. Katano (2006). Nuclear factor-kappaB contributes to hedgehog signaling pathway activation through sonic hedgehog induction in pancreatic cancer. *Cancer Res*, Vol.66, No.14, pp. 7041-9, ISSN 0008-5472
- Nelson, W. J. & R. Nusse (2004). Convergence of Wnt, beta-catenin, and cadherin pathways. *Science*, Vol.303, No.5663, pp. 1483-7, ISSN 1095-9203
- Nickoloff, B. J., J. Z. Qin, V. Chaturvedi, M. F. Denning, B. Bonish & L. Miele (2002). Jagged-1 mediated activation of notch signaling induces complete maturation of human keratinocytes through NF-kappaB and PPARgamma. *Cell Death Differ*, Vol.9, No.8, pp. 842-55, ISSN 1350-9047
- Orsulic, S., O. Huber, H. Aberle, S. Arnold & R. Kemler (1999). E-cadherin binding prevents beta-catenin nuclear localization and beta-catenin/LEF-1-mediated transactivation. *J Cell Sci*, Vol.112 (Pt 8), pp. 1237-45, ISSN 0021-9533
- Oswald, F., S. Liptay, G. Adler & R. M. Schmid (1998). NF-kappaB2 is a putative target gene of activated Notch-1 via RBP-Jkappa. *Mol Cell Biol*, Vol.18, No.4, pp. 2077-88, ISSN 0270-7306
- Palomero, T., W. K. Lim, D. T. Odom, M. L. Sulis, P. J. Real, A. Margolin, K. C. Barnes, J. O'Neil, D. Neuberg, A. P. Weng, J. C. Aster, F. Sigaux, J. Soulier, A. T. Look, R. A. Young, A. Califano & A. A. Ferrando (2006). NOTCH1 directly regulates c-MYC and activates a feed-forward-loop transcriptional network promoting leukemic cell growth. *Proc Natl Acad Sci U S A*, Vol.103, No.48, pp. 18261-6, ISSN 0027-8424

- Pap, M. & G. M. Cooper (1998). Role of glycogen synthase kinase-3 in the phosphatidylinositol 3-Kinase/Akt cell survival pathway. *J Biol Chem*, Vol.273, No.32, pp. 19929-32, ISSN 0021-9258
- Park, C. H., J. Y. Chang, E. R. Hahm, S. Park, H. K. Kim & C. H. Yang (2005). Quercetin, a potent inhibitor against beta-catenin/Tcf signaling in SW480 colon cancer cells. *Biochem Biophys Res Commun*, Vol.328, No.1, pp. 227-34, ISSN 0006-291X
- Park, C. H., E. R. Hahm, S. Park, H. K. Kim & C. H. Yang (2005). The inhibitory mechanism of curcumin and its derivative against beta-catenin/Tcf signaling. *FEBS Lett*, Vol.579, No.13, pp. 2965-71, ISSN 0014-5793
- Park, S. Y., G. Y. Kim, S. J. Bae, Y. H. Yoo & Y. H. Choi (2007). Induction of apoptosis by isothiocyanate sulforaphane in human cervical carcinoma HeLa and hepatocarcinoma HepG2 cells through activation of caspase-3. *Oncol Rep*, Vol.18, No.1, pp. 181-7, ISSN 1021-335X
- Pasca di Magliano, M. & M. Hebrok (2003). Hedgehog signalling in cancer formation and maintenance. *Nat Rev Cancer*, Vol.3, No.12, pp. 903-11, ISSN 1474-175X
- Peacock, C. D. & D. N. Watkins (2008). Cancer stem cells and the ontogeny of lung cancer. *J Clin Oncol*, Vol.26, No.17, pp. 2883-9, ISSN 1527-7755
- Peng, G., D. A. Dixon, S. J. Muga, T. J. Smith & M. J. Wargovich (2006). Green tea polyphenol (-)-epigallocatechin-3-gallate inhibits cyclooxygenase-2 expression in colon carcinogenesis. *Mol Carcinog*, Vol.45, No.5, pp. 309-19, ISSN 0899-1987
- Polakis, P. (2000). Wnt signaling and cancer. *Genes Dev*, Vol.14, No.15, pp. 1837-51, ISSN 0890-9369
- Prasad, C. P., G. Rath, S. Mathur, D. Bhatnagar & R. Ralhan (2009). Potent growth suppressive activity of curcumin in human breast cancer cells: Modulation of Wnt/beta-catenin signaling. *Chem Biol Interact*, Vol.181, No.2, pp. 263-71, ISSN 1872-7786
- Rangarajan, A., C. Talora, R. Okuyama, M. Nicolas, C. Mammucari, H. Oh, J. C. Aster, S. Krishna, D. Metzger, P. Chambon, L. Miele, M. Aguet, F. Radtke & G. P. Dotto (2001). Notch signaling is a direct determinant of keratinocyte growth arrest and entry into differentiation. *EMBO J*, Vol.20, No.13, pp. 3427-36, ISSN 0261-4189
- Rausch, V., L. Liu, G. Kallifatidis, B. Baumann, J. Mattern, J. Gladkikh, T. Wirth, P. Schemmer, M. W. Buchler, M. Zoller, A. V. Salnikov & I. Herr (2010). Synergistic activity of sorafenib and sulforaphane abolishes pancreatic cancer stem cell characteristics. *Cancer Res*, Vol.70, No.12, pp. 5004-13, ISSN 1538-7445
- Ravindran, J., S. Prasad & B. B. Aggarwal (2009). Curcumin and cancer cells: how many ways can curry kill tumor cells selectively? *AAPS J*, Vol.11, No.3, pp. 495-510, ISSN 1550-7416
- Reya, T., S. J. Morrison, M. F. Clarke & I. L. Weissman (2001). Stem cells, cancer, and cancer stem cells. *Nature*, Vol.414, No.6859, pp. 105-11, ISSN 0028-0836
- Ronchini, C. & A. J. Capobianco (2001). Induction of cyclin D1 transcription and CDK2 activity by Notch(ic): implication for cell cycle disruption in transformation by Notch(ic). *Mol Cell Biol*, Vol.21, No.17, pp. 5925-34, ISSN 0270-7306
- Ryu, M. J., M. Cho, J. Y. Song, Y. S. Yun, I. W. Choi, D. E. Kim, B. S. Park & S. Oh (2008). Natural derivatives of curcumin attenuate the Wnt/beta-catenin pathway through down-regulation of the transcriptional coactivator p300. *Biochem Biophys Res Commun*, Vol.377, No.4, pp. 1304-8, ISSN 1090-2104

- Sa, G. & T. Das (2008). Anti cancer effects of curcumin: cycle of life and death. *Cell Div*, Vol.3, pp. 14, ISSN 1747-1028
- Sarkar, F. H., Y. Li, Z. Wang & D. Kong (2009). Cellular signaling perturbation by natural products. *Cell Signal*, Vol.21, No.11, pp. 1541-7, ISSN 1873-3913
- Sarkar, F. H., Y. Li, Z. Wang & D. Kong (2009). Pancreatic cancer stem cells and EMT in drug resistance and metastasis. *Minerva Chir*, Vol.64, No.5, pp. 489-500, ISSN 0026-4733
- Satoh, Y., I. Matsumura, H. Tanaka, S. Ezoe, H. Sugahara, M. Mizuki, H. Shibayama, E. Ishiko, J. Ishiko, K. Nakajima & Y. Kanakura (2004). Roles for c-Myc in self-renewal of hematopoietic stem cells. *J Biol Chem*, Vol.279, No.24, pp. 24986-93, ISSN 0021-9258
- Schulenburg, A., P. Cech, I. Herbacek, B. Marian, F. Wrba, P. Valent & H. Ulrich-Pur (2007). CD44-positive colorectal adenoma cells express the potential stem cell markers musashi antigen (msi1) and ephrin B2 receptor (EphB2). *J Pathol*, Vol.213, No.2, pp. 152-60, ISSN 0022-3417
- Schweizer, L. & H. Varmus (2003). Wnt/Wingless signaling through beta-catenin requires the function of both LRP/Arrow and frizzled classes of receptors. *BMC Cell Biol*, Vol.4, pp. 4, ISSN 1471-2121
- Scoville, D. H., T. Sato, X. C. He & L. Li (2008). Current view: intestinal stem cells and signaling. *Gastroenterology*, Vol.134, No.3, pp. 849-64, ISSN 1528-0012
- Shafee, N., C. R. Smith, S. Wei, Y. Kim, G. B. Mills, G. N. Hortobagyi, E. J. Stanbridge & E. Y. Lee (2008). Cancer stem cells contribute to cisplatin resistance in Brca1/p53-mediated mouse mammary tumors. *Cancer Res*, Vol.68, No.9, pp. 3243-50, ISSN 1538-7445
- Shan, B. E., M. X. Wang & R. Q. Li (2009). Quercetin inhibit human SW480 colon cancer growth in association with inhibition of cyclin D1 and survivin expression through Wnt/beta-catenin signaling pathway. *Cancer Invest*, Vol.27, No.6, pp. 604-12, ISSN 1532-4192
- Shankar, S., S. Ganapathy, S. R. Hingorani & R. K. Srivastava (2008). EGCG inhibits growth, invasion, angiogenesis and metastasis of pancreatic cancer. *Front Biosci*, Vol.13, pp. 440-52, ISSN 1093-4715
- Shankar, S., S. Ganapathy & R. K. Srivastava (2007). Green tea polyphenols: biology and therapeutic implications in cancer. *Front Biosci*, Vol.12, pp. 4881-99, ISSN 1093-4715
- Shankar, S., S. Ganapathy & R. K. Srivastava (2008). Sulforaphane enhances the therapeutic potential of TRAIL in prostate cancer orthotopic model through regulation of apoptosis, metastasis, and angiogenesis. *Clin Cancer Res*, Vol.14, No.21, pp. 6855-66, ISSN 1078-0432
- Shao, Z. M., Z. Z. Shen, C. H. Liu, M. R. Sartippour, V. L. Go, D. Heber & M. Nguyen (2002). Curcumin exerts multiple suppressive effects on human breast carcinoma cells. *Int J Cancer*, Vol.98, No.2, pp. 234-40, ISSN 0020-7136
- Shen, G., T. O. Khor, R. Hu, S. Yu, S. Nair, C. T. Ho, B. S. Reddy, M. T. Huang, H. L. Newmark & A. N. Kong (2007). Chemoprevention of familial adenomatous polyposis by natural dietary compounds sulforaphane and dibenzoylmethane alone and in combination in ApcMin/+ mouse. *Cancer Res*, Vol.67, No.20, pp. 9937-44, ISSN 0008-5472
- Shih, Y. L., H. C. Liu, C. S. Chen, C. H. Hsu, M. H. Pan, H. W. Chang, C. H. Chang, F. C. Chen, C. T. Ho, Y. Y. Yang & Y. S. Ho (2010). Combination treatment with luteolin

- and quercetin enhances antiproliferative effects in nicotine-treated MDA-MB-231 cells by down-regulating nicotinic acetylcholine receptors. *J Agric Food Chem*, Vol.58, No.1, pp. 235-41, ISSN 1520-5118
- Shimizu, M., A. Deguchi, J. T. Lim, H. Moriwaki, L. Kopelovich & I. B. Weinstein (2005). (-)-Epigallocatechin gallate and polyphenon E inhibit growth and activation of the epidermal growth factor receptor and human epidermal growth factor receptor-2 signaling pathways in human colon cancer cells. *Clin Cancer Res*, Vol.11, No.7, pp. 2735-46, ISSN 1078-0432
- Shin, H. M., L. M. Minter, O. H. Cho, S. Gottipati, A. H. Fauq, T. E. Golde, G. E. Sonenshein & B. A. Osborne (2006). Notch1 augments NF-kappaB activity by facilitating its nuclear retention. *EMBO J*, Vol.25, No.1, pp. 129-38, ISSN 0261-4189
- Shoba, G., D. Joy, T. Joseph, M. Majeed, R. Rajendran & P. S. Srinivas (1998). Influence of piperine on the pharmacokinetics of curcumin in animals and human volunteers. *Planta Med*, Vol.64, No.4, pp. 353-6, ISSN 0032-0943
- Singh, A. & J. Settleman (2010). EMT, cancer stem cells and drug resistance: an emerging axis of evil in the war on cancer. *Oncogene*, Vol.29, No.34, pp. 4741-51, ISSN 1476-5594
- Singh, A. V., D. Xiao, K. L. Lew, R. Dhir & S. V. Singh (2004). Sulforaphane induces caspase-mediated apoptosis in cultured PC-3 human prostate cancer cells and retards growth of PC-3 xenografts in vivo. *Carcinogenesis*, Vol.25, No.1, pp. 83-90, ISSN 0143-3334
- Singh, S. V., A. Herman-Antosiewicz, A. V. Singh, K. L. Lew, S. K. Srivastava, R. Kamath, K. D. Brown, L. Zhang & R. Baskaran (2004). Sulforaphane-induced G2/M phase cell cycle arrest involves checkpoint kinase 2-mediated phosphorylation of cell division cycle 25C. *J Biol Chem*, Vol.279, No.24, pp. 25813-22, ISSN 0021-9258
- Smalley, M. J. & T. C. Dale (1999). Wnt signalling in mammalian development and cancer. *Cancer Metastasis Rev*, Vol.18, No.2, pp. 215-30, ISSN 0167-7659
- Smith-Warner, S. A., D. Spiegelman, S. S. Yaun, D. Albanes, W. L. Beeson, P. A. van den Brandt, D. Feskanich, A. R. Folsom, G. E. Fraser, J. L. Freudenheim, E. Giovannucci, R. A. Goldbohm, S. Graham, L. H. Kushi, A. B. Miller, P. Pietinen, T. E. Rohan, F. E. Speizer, W. C. Willett & D. J. Hunter (2003). Fruits, vegetables and lung cancer: a pooled analysis of cohort studies. *Int J Cancer*, Vol.107, No.6, pp. 1001-11, ISSN 0020-7136
- Srivastava, R. K., S. N. Tang, W. Zhu, D. Meeker & S. Shankar (2011). Sulforaphane synergizes with quercetin to inhibit self-renewal capacity of pancreatic cancer stem cells. *Front Biosci (Elite Ed)*, Vol.3, pp. 515-28, ISSN 1945-0508
- Stearns, M. E., M. D. Amatangelo, D. Varma, C. Sell & S. M. (2010). Goodyear Combination Therapy with Epigallocatechin-3-Gallate and Doxorubicin in Human Prostate Tumor Modeling Studies. Inhibition of Metastatic Tumor Growth in Severe Combined Immunodeficiency Mice. *Am J Pathol*, ISSN 1525-2191
- Stockler, M., N. R. Wilcken, D. Ghersi & R. J. Simes (2000). Systematic reviews of chemotherapy and endocrine therapy in metastatic breast cancer. *Cancer Treat Rev*, Vol.26, No.3, pp. 151-68, ISSN 0305-7372
- Strimpakos, A. S. & R. A. Sharma (2008). Curcumin: preventive and therapeutic properties in laboratory studies and clinical trials. *Antioxid Redox Signal*, Vol.10, No.3, pp. 511-45, ISSN 1523-0864

- Takahashi-Yanaga, F. & T. Sasaguri (2008). GSK-3 β regulates cyclin D1 expression: a new target for chemotherapy. *Cell Signal*, Vol.20, No.4, pp. 581-9, ISSN 0898-6568
- Tang, S. N., C. Singh, D. Nall, D. Meeker, S. Shankar & R. K. Srivastava (2010). The dietary bioflavonoid quercetin synergizes with epigallocatechin gallate (EGCG) to inhibit prostate cancer stem cell characteristics, invasion, migration and epithelial-mesenchymal transition. *J Mol Signal*, Vol.5, pp. 14, ISSN 1750-2187
- Teng, Y., X. Wang, Y. Wang & D. Ma (2010). Wnt/ β -catenin signaling regulates cancer stem cells in lung cancer A549 cells. *Biochem Biophys Res Commun*, ISSN 1090-2104
- Tetsu, O. & F. McCormick (1999). β -catenin regulates expression of cyclin D1 in colon carcinoma cells. *Nature*, Vol.398, No.6726, pp. 422-6, ISSN 0028-0836
- Turashvili, G., J. Bouchal, G. Burkadze & Z. Kolar (2006). Wnt signaling pathway in mammary gland development and carcinogenesis. *Pathobiology*, Vol.73, No.5, pp. 213-23, ISSN 1015-2008
- Wang, J., L. Shelly, L. Miele, R. Boykins, M. A. Norcross & E. Guan (2001). Human Notch-1 inhibits NF- κ B activity in the nucleus through a direct interaction involving a novel domain. *J Immunol*, Vol.167, No.1, pp. 289-95, ISSN 0022-1767
- Wang, Y., S. L. Chan, L. Miele, P. J. Yao, J. Mackes, D. K. Ingram, M. P. Mattson & K. Furukawa (2004). Involvement of Notch signaling in hippocampal synaptic plasticity. *Proc Natl Acad Sci U S A*, Vol.101, No.25, pp. 9458-62, ISSN 0027-8424
- Wang, Z., S. Banerjee, Y. Li, K. M. Rahman, Y. Zhang & F. H. Sarkar (2006). Down-regulation of notch-1 inhibits invasion by inactivation of nuclear factor- κ B, vascular endothelial growth factor, and matrix metalloproteinase-9 in pancreatic cancer cells. *Cancer Res*, Vol.66, No.5, pp. 2778-84, ISSN 0008-5472
- Wang, Z., Y. Li, S. Banerjee & F. H. Sarkar (2009). Emerging role of Notch in stem cells and cancer. *Cancer Lett*, Vol.279, No.1, pp. 8-12, ISSN 1872-7980
- Wang, Z., Y. Zhang, S. Banerjee, Y. Li & F. H. Sarkar (2006). Notch-1 down-regulation by curcumin is associated with the inhibition of cell growth and the induction of apoptosis in pancreatic cancer cells. *Cancer*, Vol.106, No.11, pp. 2503-13, ISSN 0008-543X
- Weng, A. P., J. M. Millholland, Y. Yashiro-Ohtani, M. L. Arcangeli, A. Lau, C. Wai, C. Del Bianco, C. G. Rodriguez, H. Sai, J. Tobias, Y. Li, M. S. Wolfe, C. Shachaf, D. Felsher, S. C. Blacklow, W. S. Pear & J. C. Aster (2006). c-Myc is an important direct target of Notch1 in T-cell acute lymphoblastic leukemia/lymphoma. *Genes Dev*, Vol.20, No.15, pp. 2096-109, ISSN 0890-9369
- Williams, S. D., R. Birch, L. H. Einhorn, L. Irwin, F. A. Greco & P. J. Loehrer (1987). Treatment of disseminated germ-cell tumors with cisplatin, bleomycin, and either vinblastine or etoposide. *N Engl J Med*, Vol.316, No.23, pp. 1435-40, ISSN 0028-4793
- Williamson, G. & C. Manach (2005). Bioavailability and bioefficacy of polyphenols in humans. II. Review of 93 intervention studies. *Am J Clin Nutr*, Vol.81, No.1 Suppl, pp. 243S-255S, ISSN 0002-9165
- Wilson, A. & F. Radtke (2006). Multiple functions of Notch signaling in self-renewing organs and cancer. *FEBS Lett*, Vol.580, No.12, pp. 2860-8, ISSN 0014-5793
- Wong, M. Y. & G. N. (2010). Chiu Simultaneous liposomal delivery of quercetin and vincristine for enhanced estrogen-receptor-negative breast cancer treatment. *Anticancer Drugs*, Vol.21, No.4, pp. 401-10, ISSN 1473-5741

- Woodward, W. A., M. S. Chen, F. Behbod, M. P. Alfaro, T. A. Buchholz & J. M. Rosen (2007). WNT/beta-catenin mediates radiation resistance of mouse mammary progenitor cells. *Proc Natl Acad Sci U S A*, Vol.104, No.2, pp. 618-23, ISSN 0027-8424
- Wu, C. & B. A. Alman (2008). Side population cells in human cancers. *Cancer Lett*, Vol.268, No.1, pp. 1-9, ISSN 0304-3835
- Wu, J. Y. & Y. Rao (1999). Fringe: defining borders by regulating the notch pathway. *Curr Opin Neurobiol*, Vol.9, No.5, pp. 537-43, ISSN 0959-4388
- Xu, C., M. T. Huang, G. Shen, X. Yuan, W. Lin, T. O. Khor, A. H. Conney & A. N. Kong (2006). Inhibition of 7,12-dimethylbenz(a)anthracene-induced skin tumorigenesis in C57BL/6 mice by sulforaphane is mediated by nuclear factor E2-related factor 2. *Cancer Res*, Vol.66, No.16, pp. 8293-6, ISSN 1538-7445
- Yamaguchi, T. P. (2001). Heads or tails: Wnts and anterior-posterior patterning. *Curr Biol*, Vol.11, No.17, pp. R713-24, ISSN 0960-9822
- Yan, C., M. S. Jamaluddin, B. Aggarwal, J. Myers & D. D. Boyd (2005). Gene expression profiling identifies activating transcription factor 3 as a novel contributor to the proapoptotic effect of curcumin. *Mol Cancer Ther*, Vol.4, No.2, pp. 233-41, ISSN 1535-7163
- Yang, C. S., P. Maliakal & X. Meng (2002). Inhibition of carcinogenesis by tea. *Annu Rev Pharmacol Toxicol*, Vol.42, pp. 25-54, ISSN 0362-1642
- Yang, W., H. X. Yan, L. Chen, Q. Liu, Y. Q. He, L. X. Yu, S. H. Zhang, D. D. Huang, L. Tang, X. N. Kong, C. Chen, S. Q. Liu, M. C. Wu & H. Y. Wang (2008). Wnt/beta-catenin signaling contributes to activation of normal and tumorigenic liver progenitor cells. *Cancer Res*, Vol.68, No.11, pp. 4287-95, ISSN 1538-7445
- Yost, C., M. Torres, J. R. Miller, E. Huang, D. Kimelman & R. T. Moon (1996). The axis-inducing activity, stability, and subcellular distribution of beta-catenin is regulated in *Xenopus* embryos by glycogen synthase kinase 3. *Genes Dev*, Vol.10, No.12, pp. 1443-54, ISSN 0890-9369
- Ysebaert, L., G. Chicanne, C. Demur, F. De Toni, N. Prade-Houdellier, J. B. Ruidavets, V. Mansat-De Mas, F. Rigal-Huguet, G. Laurent, B. Payrastre, S. Manenti & C. Racaud-Sultan (2006). Expression of beta-catenin by acute myeloid leukemia cells predicts enhanced clonogenic capacities and poor prognosis. *Leukemia*, Vol.20, No.7, pp. 1211-6, ISSN 0887-6924
- Zhang, Y., T. W. Kensler, C. G. Cho, G. H. Posner & P. Talalay (1994). Anticarcinogenic activities of sulforaphane and structurally related synthetic norbornyl isothiocyanates. *Proc Natl Acad Sci U S A*, Vol.91, No.8, pp. 3147-50, ISSN 0027-8424
- Zhang, Y., P. Talalay, C. G. Cho & G. H. Posner (1992). A major inducer of anticarcinogenic protective enzymes from broccoli: isolation and elucidation of structure. *Proc Natl Acad Sci U S A*, Vol.89, No.6, pp. 2399-403, ISSN 0027-8424
- Zhang, Y. & L. Tang (2007). Discovery and development of sulforaphane as a cancer chemopreventive phytochemical. *Acta Pharmacol Sin*, Vol.28, No.9, pp. 1343-54, ISSN 1671-4083
- Zhou, B. B., H. Zhang, M. Damelin, K. G. Geles, J. C. Grindley & P. B. Dirks (2009). Tumour-initiating cells: challenges and opportunities for anticancer drug discovery. *Nat Rev Drug Discov*, Vol.8, No.10, pp. 806-23, ISSN 1474-1784
- Zhou, S., J. D. Schuetz, K. D. Bunting, A. M. Colapietro, J. Sampath, J. J. Morris, I. Lagutina, G. C. Grosveld, M. Osawa, H. Nakauchi & B. P. Sorrentino (2001). The ABC

transporter Bcrp1/ABCG2 is expressed in a wide variety of stem cells and is a molecular determinant of the side-population phenotype. *Nat Med*, Vol.7, No.9, pp. 1028-34, ISSN 1078-8956

Zhou, W., G. Kallifatidis, B. Baumann, V. Rausch, J. Mattern, J. Gladkich, N. Giese, G. Moldenhauer, T. Wirth, M. W. Buchler, A. V. Salnikov & I. Herr (2010). Dietary polyphenol quercetin targets pancreatic cancer stem cells. *Int J Oncol*, Vol.37, No.3, pp. 551-61, ISSN 1791-2423



Cancer Stem Cells - The Cutting Edge

Edited by Prof. Stanley Shostak

ISBN 978-953-307-580-8

Hard cover, 606 pages

Publisher InTech

Published online 01, August, 2011

Published in print edition August, 2011

Over the last thirty years, the foremost inspiration for research on metastasis, cancer recurrence, and increased resistance to chemo- and radiotherapy has been the notion of cancer stem cells. The twenty-eight chapters assembled in *Cancer Stem Cells - The Cutting Edge* summarize the work of cancer researchers and oncologists at leading universities and hospitals around the world on every aspect of cancer stem cells, from theory and models to specific applications (glioma), from laboratory research on signal pathways to clinical trials of bio-therapies using a host of devices, from solutions to laboratory problems to speculation on cancers' stem cells' evolution. Cancer stem cells may or may not be a subset of slowly dividing cancer cells that both disseminate cancers and defy oncotoxic drugs and radiation directed at rapidly dividing bulk cancer cells, but research on cancer stem cells has paid dividends for cancer prevention, detection, targeted treatment, and improved prognosis.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Yanyan Li, Steven J. Schwartz and Duxin Sun (2011). Potential Application of Natural Dietary Components to Target Cancer Stem Cells, *Cancer Stem Cells - The Cutting Edge*, Prof. Stanley Shostak (Ed.), ISBN: 978-953-307-580-8, InTech, Available from: <http://www.intechopen.com/books/cancer-stem-cells-the-cutting-edge/potential-application-of-natural-dietary-components-to-target-cancer-stem-cells>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2011 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike-3.0 License](https://creativecommons.org/licenses/by-nc-sa/3.0/), which permits use, distribution and reproduction for non-commercial purposes, provided the original is properly cited and derivative works building on this content are distributed under the same license.

IntechOpen

IntechOpen