

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

186,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Residual Vascular Risk in T2DM: The Next Frontier

Michel P. Hermans¹, Sylvie A. Ahn² and Michel F. Rousseau²

¹Endocrinology & Nutrition, Cliniques universitaires St-Luc, Brussels

²Cardiology Division, Cliniques universitaires St-Luc, Brussels
Belgium

1. Introduction

In T2DM, macro- and microvascular complications represent the major cause for morbi-/mortality, decreased quality of life and healthcare costs. Current guidelines for standards of care in T2DM emphasize the significance of multifactorial intervention on standard modifiable variables, in order to achieve recommended levels of blood glucose, LDL-C and blood pressure. T2DM patients achieving such targets represent a minority. Many of those not meeting those targets are exposed to high residual vascular risk (RVR) to develop incident micro- and macrovascular events and/or to suffer from progression of existing complications. Determining RVR in T2DM patients is of major relevance, as a substantial fraction of it is modifiable, including a lipid-related fraction, associated with both LDL and non-LDL lipids and lipoproteins. Atherogenic dyslipidemia (AD) is characterized by raised fasting triglycerides and low HDL-C. AD contributes to RVR of micro- and macrovascular disease in T2DM, even when LDL-C and/or hyperglycemia are controlled. The presence of a metabolic syndrome, or its score, is an additional means to capture modifiable components of RVR for micro- and macroangiopathy. AD is best prevented and addressed by therapeutic lifestyle changes, fibrates, or nicotinic acid. In the ACCORD Lipid trial, RVR of macrovascular events was high despite background simvastatin, and was substantially decreased in patients with AD following bitherapy with fenofibrate plus simvastatin. In FIELD and/or ACCORD trials, fenofibrate also decreased RVR of retinopathy progression, irrespective of baseline lipids, reduced albuminuria incidence and risk of diabetes-related lower-limb amputations. These data suggest a wider role for fenofibrate in the management of multisite microvessel RVR in T2DM. As regards other anti-dyslipidaemic drugs, ongoing trials will establish whether targeting low HDL-C with niacin reduces RVR in high-risk T2DM patients.

2. Standards of care in T2DM and target attainments

In type 2 diabetes mellitus (T2DM), vascular complications represent the major cause for morbi-/mortality, decreased quality of life and healthcare costs. These chronic, long-term complications arise in the setting of elevated residual vascular risk (RVR) factors. These factors may not only damage *macro*vessels, leading to premature-onset coronary artery disease (CAD), cerebrovascular disease (transient ischemic attack (TIA) and/or stroke), or

peripheral arterial disease (PAD), but also affect *microvessels* leading to “diabetic complications”, i.e. diabetes-specific long-term complications of chronic hyperglycemia: retinopathy, nephropathy or neuropathy. Certain vascular diabetic complications such as PAD, TIA/stroke, nephropathy or erectile dysfunction may arise from a combination of macro- and microangiopathies [1-6].

Low-density lipoprotein cholesterol (LDL-C)

Due to the overall efficacy of statins, and the relatively low baseline level of LDL-C in T2DM, LDL-C may be considered as the easiest-modifiable single parameter in diabetic patients [4,5,7-15]. However, in the recent *Centralized Pan-European survey on the under-treatment of hypercholesterolaemia* (CEPHEUS), a primary care survey from 8 European countries, under-treatment of hypercholesterolaemia was highly prevalent, with low total and LDL-C target attainment observed across countries [16,17]. In CEPHEUS, patients with the highest CV risk were T2DM patients with established cardiovascular disease (CVD) (i.e. patients in true secondary CV prevention). Yet, these highest-risk patients achieved the lowest level of LDL-C target attainment, with only 27% attaining the <70 mg/dL target. Moreover, only 58% of T2DM patients without CVD achieved LDL-C target <100 mg/dL. Eight modifiable variables were associated with LDL-C target attainment in CEPHEUS [17]:

1. normal body mass index;
2. being a non-smoker;
3. not having a metabolic syndrome (MetS) phenotype;
4. current treatment with a statin;
5. belonging to a medium-high CVD risk category;
6. good treatment adherence;
7. high patient's awareness of his/her current LDL-C level; and/or
8. frequency of cholesterol reviews.

Six non-modifiable factors were also associated with LDL-C target attainment:

1. age >70 years;
2. male gender;
3. history of diabetes;
4. history of hypertension;
5. absence of PAD; and/or
6. receiving LLD for secondary prevention.

Blood pressure (BP)

Controlling BP values in T2DM is of paramount importance, as it improves both macro- and microvascular outcomes. Such control is nevertheless challenging, as in the common form of T2DM, associated with the MetS and insulin resistance (IR), hypertension is not only highly prevalent, but also responds poorly to BP-lowering monotherapy intervention, and often requires multiple BP-lowering therapies on top of therapeutic lifestyle changes (TLC). In a recent study, only 16% of hypertensive T2DM treated with BP-lowering drug(s) in a tertiary-care setting achieved target BP <130/80 mmHg (<125/75 mmHg in case of proteinuria) [18].

Glycemic control and glycated haemoglobin (HbA_{1c})

Maintaining HbA_{1c} at target is an additional and constant challenge in T2DM, due to relentless loss of β -cell function over time and perpetual requirement for progressive stepping-up of glucose-lowering therapies [3,19,20]. Despite a large choice of oral and

parenteral therapies to lower blood glucose, target attainment as regards glucose control, or that of its surrogate HbA_{1c}, remains suboptimal. Among U.S. adults with diabetes in 1999–2002, only 49.8% had an HbA_{1c} <7.0% [19], a proportion similar to that found 10 years later in the *OPTimal Type 2 dIabetes Management Including Benchmarking and Standard trEatment* (OPTIMISE) trial [21].

Multifactorial intervention

Current guidelines for standards of care in T2DM emphasize the significance of multifactorial intervention on major modifiable RFs to achieve recommended levels of glucose, LDL-C and BP [12–15]. In the Steno study, a multifactorial intervention aimed at achieving recommended levels of critical indicators, including HbA_{1c} as surrogate for contemporary glucose exposure, LDL-C and systolic BP (SBP), was highly effective in reducing (micro)vascular complications [22–24].

As each of these three major modifiable targets have distinct determinants, natural histories and responses to TLC or pharmacotherapy, it comes to no surprise that only a fraction of T2DM patients will reach all three targets in synchrony in real-life conditions, leaving a majority of T2DM patients exposed to incident micro-/macrovascular events over time, or to progression of existing complications [18,21].

Many factors associated with failure to meet critical targets (HbA_{1c}, LDL-C, SBP) in T2DM represent previously identified barriers to chronic diseases management:

- age;
- disease duration;
- ethnicity;
- hyperglycaemia, hypertension and dyslipidaemia are mostly asymptomatic conditions, both in primary or secondary prevention;
- chronic disease misrepresentation;
- faulty perception of risk related to metabolic diseases is frequently observed among many T2DM patients, especially from ethnic minorities;
- patients are often poorly compliant to TLC;
- impractical, conflicting, or competing guidelines;
- many patients are poorly adherent to prescribed treatment regimens and/or to self-monitoring of blood glucose;
- fear of hypoglycaemia;
- concerns about weight gain;
- fear, misperception of the natural history of T2DM and of the risk/benefit ratio of adding exogenous insulin;
- delaying tactics at the time when lifelong insulin supplementation is deemed necessary (insulinophobia);
- reluctance to resort to subcutaneous injections or to perform capillary blood glucose self-testing;
- variations among patients in pharmacological response to antidiabetic, lipid-lowering and/or BP-lowering drugs;
- reluctance of physicians and patients to increasing drug dosage or to switching drugs within classes, or to resorting to combined therapies;
- complex treatment schemes and side-effects;
- wrong perception of potential side-effects;

- insufficient counseling;
- physicians and healthcare providers inertia delaying diagnosis or stepping-up of successive interventions;
- insufficient or unfrequent laboratory follow-up;
- competing T2DM-related co-morbidities and complications: obstructive sleep apnoea syndrome, chronic kidney disease or left ventricular dysfunction;
- lack of patient empowerment and responsibility for self-care;
- social pressures and discrimination related to aspects of diabetes management;
- low socioeconomic or educational status;
- unsupportive/overstretched healthcare systems.

3. Residual vascular risk in T2DM

RVR in T2DM is best defined as “the residual risk of incident vascular events or progression of established vascular damage persisting in patients treated with current evidence-based recommended care, including risk from established risk factors, such as dyslipidemia, high blood pressure, hyperglycemia, inflammation and unhealthy lifestyles, and risk related to emerging or newer risk factors” [4,5].

Determining macro- and micro- RVR in T2DM patients after implementation of standards of care is especially relevant when dealing with a chronic condition in which a substantial fraction of risk remains addressable, eg. by further lowering of exposure levels to standards RFs for micro- and macroangiopathy. Besides those modifiable components to RVR, one should also consider in risk assessment non-gender, non-modifiable components of RVR in T2DM, such as ethnicity, certain polymorphisms, and familial histories for (i) premature-onset CVD; (ii) obesity; and/or (iii) impaired glucose homeostasis or diabetes.

For macrovascular RVR, the usual approach involves single-variable assessment (HbA_{1c}, SBP and LDL-C) and targeting with TLC and/or pharmacotherapy. Due to an overwhelmingly glucocentric approach to T2DM management, the hierarchy of priorities follows a sequence in which hyperglycemia control ranks first, followed on a par with BP control and LDL-C lowering with statin as preferred agent, and then with a needs assessment for aspirin therapy as antiplatelet agent in high-risk patients [12-15].

Combined assessment of the harmful effects of multiple coexisting modifiable variables on RVR is rarely done for an individual T2DM patient, even though multifactorial intervention was demonstrated to be highly-effective in reducing micro- and macrovascular RVR [22-24]. Various calculators were proposed to estimate absolute risk in nondiabetic and diabetic patients [25-31]. At present, the best means to predict macrovascular residual risk of CAD (nonlethal or lethal) and stroke (nonlethal or lethal) in patients in primary macrovascular prevention is the T2DM-specific calculator *United Kingdom Prospective Diabetes Study (UKPDS) Risk Engine* [25,27,28]. It computes risk from the following variables: (i) known T2DM duration, (ii) age, (iii) gender, (iv) ethnicity, (v) smoking status, (vi) atrial fibrillation, (vii) current HbA_{1c} level, (viii) SBP, (ix) total cholesterol and (x) high-density lipoprotein cholesterol (HDL-C).

In a diabetic RVR perspective, such calculations of absolute predicted risk should ideally be performed prior to, and following implementations of standards of care to control hyperglycemia, high BP and hypercholesterolemia, with updated levels as input for estimating the magnitude of risk reduction. Such an approach will yield relevant figures

regarding (i) absolute vascular risk for CAD (nonlethal or lethal) and stroke (nonlethal or lethal), (ii) RVR for CAD (nonlethal or lethal) and stroke (nonlethal or lethal), together with (iii) absolute and (iv) relative decreases in CAD (nonlethal or lethal) and stroke (nonlethal or lethal) risks [25,27,28].

In order to illustrate RVR in T2DM from real-life conditions, we systematically assessed UKPDS Risk in 429 consecutive T2DM outpatients in primary macrovascular prevention. Eighty percent were White Caucasians, with a male-to-female ratio of 59:41. Mean age (1 standard deviation [SD]) was 62 (12) years, and known diabetes duration 12 (8) years. Sixteen percent were current smokers. Major modifiable variables in this cohort receiving standards of care in an academic setting were: total cholesterol: 173 (39) mg/dL; HDL-C: 48 (14) mg/dL; SBP: 137 (18) mmHg; and HbA_{1c}: 7.6 (1.47)%. The 10-year UKPDS Risk Engine estimated RVR values are illustrated in **Figure 1**. For the entire cohort, 10-year risk of CAD was high (almost 20%), a level in accordance with the status of *secondary-prevention equivalent* proposed for T2DM in primary macrovascular prevention [1,2]. According to gender, male T2DM patients had a 62% higher absolute risk for CAD than female patients, although the difference between genders was less marked for fatal CAD or stroke, and abolished for fatal stroke, illustrating the loss of protective effects afforded by the female gender in T2DM. Despite their primary CVD prevention status, those 429 patients had a high prevalence of microangiopathies: 46% (any microangiopathy); 21% (retinopathy); 23% (peripheral neuropathy); and 36% (albuminuria). These figures highlight the complexity of defining RVR in diabetic patients, who may at any time belong to different risk categories according to (i) the micro- vs. macrovascular level of dichotomy, and (ii) which target organs are under scrutiny.




Fig. 1. UKPDS Risk Engine's 10-year prediction estimates for developing non-fatal and fatal coronary artery disease (CAD), fatal CAD, non-fatal and fatal stroke in T2DM patients of both gender (n=429; solid bars), both groups in primary cardiovascular prevention. The UKPDS Risk Engine computes the following variables, i.e. (1) known T2DM duration, (2) age, (3) gender, (4) ethnicity, (5) smoking status, (6) atrial fibrillation, (7) HbA_{1c} level, (8) systolic BP, (9) total cholesterol (C) and (10) HDL-C.

With respect to microangiopathy, individual RVR assessment is limited to an “educated guess” relating *hyperglycemia exposure time* (with known diabetes duration as surrogate) to *hyperglycemia severity* (with current HbA_{1c} as surrogate). Unfortunately, there exists no assigned microvascular calculator available for estimating 10-year absolute risk in major target organs, such as retina, kidney or peripheral nerves destined to diabetic patients or to prediabetic patients with rapid loss of β -cell function and as such at very high risk for new-onset T2DM. Such a limitation in microangiopathic RVR represents a truly unmet need for T2DM management. Such is also the case when it comes to estimating absolute RVR in end-organs at risk for combined micro- and macroangiopathies, such as PAD in lower limbs, cerebrovascular disease, certain subtypes of nephropathy, or erectile dysfunction. Other limitations of current risk calculators for T2DM patients include: (i) the complexity of the underlying pathophysiological processes; (ii) the poor predictive value of (micro)albuminuria as surrogate to diabetes-related nephropathy; (iii) the lack of effective clinical surrogate for early-neuropathy; and (iv) the exclusion of familial histories for diabetes, early-onset CVD, obesity and/or IR and of their mutual metabolic/vascular impacts [25-32].

4. Atherogenic dyslipidemia

A major component of modifiable RVR is lipid-related, both associated with LDL and non-LDL, including atherogenic dyslipidemia (AD). Epidemiological studies and landmark intervention trials clearly established that AD contributes to RVR for macrovascular disease in T2DM, even when LDL-C and/or hyperglycemia are controlled at baseline. The hallmark of AD is raised fasting TG and low HDL-C levels, two routinely available markers of a series of complex and deleterious metabolic abnormalities which affect, in a proatherogenic way, LDL and non-LDL lipoproteins composition and numbers. These abnormalities also diminish, in an antiatherogenic way, the dynamics and magnitude of reverse cholesterol transport [33-45].

The underlying process driving AD in T2DM or nondiabetic patients consists of (i) overproduction by the liver of TG-rich, apolipoprotein B₁₀₀(apoB)-carrying lipoproteins of the very-low density lipoprotein (VLDL) class, as a direct result of (ii) whole body IR and compensatory hyperinsulinemia, leading to (iii) raised portal insulin levels, which affect (iv) VLDL assembly and export by the hepatocyte, as a consequence of preserved insulin sensitivity of select pathways for lipogenesis and TG-rich lipoproteins synthesis [33,34,36,37,41,45].

Screening for AD provides a clinically-relevant and easy means to capture RVR associated with low HDL-C, high TG and their determinants. Such screening is not routinely performed due eg. to lack of general agreement on criteria or cut-offs based on current *vs.* baseline HDL-C and TG values. One way to diagnose AD consists of establishing the combined occurrence of high TG levels and low HDL-C. This seemingly easy estimation is rarely performed as such, due to (i) lack of consensual cut-off values across gender, race/ethnicities or underlying conditions; (ii) the requirement for baseline lipid values (prior to lipid-lowering drugs (LLD)); and (iii) the limiting fact that a *sine qua non* co-occurrence definition does not take into account the linearity of these non-LDL abnormalities, since both HDL-C and TG are continuous CVD risk variables [35,38,39,40,42,44,45]. Such an approach based on co-occurrence may also underestimate AD prevalence and severity in T2DM subpopulations with spontaneously low (Afro-

Americans, sub-Saharan Africans) or elevated TG levels [46-48]. A more rationale approach would be to use ratios between TG and HDL-C in order to incorporate each component's information as continuous variable while increasing the epidemiological potency by having the atherogenic TG variable as *numerator* together with the anti-atherogenic HDL-C as *denominator* [35,38,39,40,45]. We showed that $\log(\text{TG})/\text{HDL-C}$ is a simple means to estimate AD and the residual CV risk it confers to T2DM patients. Thus, this AD surrogate ratio was associated with major cardiometabolic and glucose homeostasis determinants, as well as with poorer metabolic control, and related to macroangiopathy prevalence and estimated UKPDS CAD risk [44].

5. Metabolic syndrome

Low HDL-C and high TG are part of the MetS definition, either as individual AD component or in combination. The presence of a MetS phenotype or its score (from 0/5 to 5/5) is another simple means to capture RVR. Identifying a MetS phenotype may be used as a dichotomic state (presence *vs.* absence). In addition, score ranking within MetS syndrome categories, besides providing a stepwise surrogate for IR/hyperinsulinemia, also provides a simple means to determine increasing CV risk categories (from 1/5 to 5/5 for T2DM patients). While the MetS is not an absolute risk calculator, its presence hints to heightened relative RVR, as a result of exposure to standard CV RFs (underlying the current definition, such as hypertension and hyperglycemia) or due to the presence of AD. MetS ranking is also associated with lesser target achievement for key variables, such as HbA_{1c}, SBP or LDL-C. In addition, the MetS also associates with microangiopathy prevalence in major target organs [18,49-55].

6. New and emerging risk markers and factors

Many candidates RFs were proposed in the last decades to improve CVD risk assessment or RVR appraisal, although few, if any, are globally acknowledged in guidelines as part of standards of care and follow-up. These candidate RFs include eg. biological markers of low-grade subclinical systemic inflammation, markers of plaque instability, of endothelial dysfunction, or of proatherothrombotic conditions. None of these emerging markers/RFs are at present used as input variables in CVD risk calculators. Other emerging RFs for risk assessment in T2DM include potentially modifiable variables contributing to, or associated with RVR, such as the MetS phenotype, IR /hyperinsulinemia, adverse lifestyle habits (excessive caloric intake, Westernized diets, smoking, high ethanol intake, sedentarity), high-cardiometabolic risk anthropometrics (abnormal distribution/expansion of fat tissue, sarcopenia), or other comorbidities increasingly described as associated with the common form of T2DM (sleep-related breathing disorders, chronic kidney disease, left ventricular systolic/diastolic dysfunction, or non-alcoholic fatty liver disease) [56-74].

Table 1 enumerates a non-exhaustive series of non-modifiable and modifiable markers/RFs for micro-/macroangiopathy which may be at play in accruing RVR in T2DM patients, including inflammatory, behavioural/environnemental, and proatherothrombotic, whereas **Table 2** lists markers/RFs related to lipids and lipoproteins or to cardiometabolic factors involved in T2DM-related RVR [45,56-74].

Non-modifiable age male gender ethnicity family history: <i>early-onset CVD</i> <i>overweight / obesity</i> <i>IFG / IGT / T2DM</i> former tobacco smoking small size at birth for gestational age right handedness genes / loci and polymorphisms associated with: <i>CVD</i> <i>β-cell function loss</i> <i>overweight / obesity</i>	Coagulation - haemostasis - platelets platelet activity platelet aggregation platelet size / volume aspirin resistance lipoprotein(a) fibrinogen factor V, VII, and VIII fibrinopeptide A PAI-1 prothrombin fragments 1 + 2 tissue-plasminogen activator von Willebrand factor antigen D-dimer
Inflammatory high-sensitivity C-reactive protein leucocyte count interleukin-6 matrix metalloproteinase 9 serum amyloid A soluble CD ₄₀ ligand vascular / cellular adhesion molecules lipoprotein-associated phospholipase A(2) periodontal disease	Varia cystatin-C asymmetric dimethylarginine nongenetic causes of iron overload hemochromatosis elevated ferritinaemia Nt-proANP and Nt-proBNP endothelin-1 urotensin II <i>Cytomegalovirus</i> , <i>Herpes simplex</i> virus <i>Helicobacter pylori</i> <i>Chlamydia pneumoniae</i> collagen vascular disease non-specific ST-segment ECG changes coronary artery calcifications left ventricular dysfunction obstructive sleep apnoea / hypopnoea syndrome psoriasis rheumatoid arthritis systemic lupus erythematosus HIV infection on highly-active antiretroviral therapy hypoglycemia unawareness
Behavioural / environnemental current tobacco smoking air pollution (including airborne fine particles) sedentary lifestyle (<i>surrogate</i> : TV viewing) physical inactivity quantitative / qualitative sarcopenia psychosocial stress low socioeducative status low income decreased fruit and vegetable consumption	

*: see Table 2 for lipid and cardiometabolic risk factors. ANP: Atrial natriuretic peptide; BNP: brain natriuretic peptide; CVD: cardiovascular disease; ECG: electrocardiogram; eGFR: estimated glomerular filtration rate; HIV: human immunodeficiency virus; IFG: impaired fasting glucose; IGT: impaired glucose tolerance; NAFLD: nonalcoholic fatty liver disease; PAI-1: plasminogen activator inhibitor type 1; T2DM type 2 diabetes mellitus; TV: television.

Table 1. Micro or macrovascular disease risk factors/markers in T2DM patients: standard, emerging and candidate

7. Reducing residual vascular risk in T2DM

a. Targets attainment.

Reducing RVR must be part of a *continuum* in managing individual patients. RVR assessment is the next logical step after implementation of good clinical practices. Despite RFs identification and standards of care provision, all major modifiable RFs rarely attain all recommended targets for a given individual in real-life conditions [12-17,21-24]. This supports a more global approach to further reduce RVR [4-6,75-84]:

Lipids and lipoproteins	Cardiometabolic
total cholesterol	overweight - obesity (<i>surrogate</i> : increased BMI)
LDL-C	central fat distribution (<i>surrogate</i> : enlarged waist)
apolipoprotein B ₁₀₀	hypertension
non-HDL-C	metabolic syndrome (presence <i>vs.</i> absence)
LDL particles number	metabolic syndrome (<i>score</i> : 3/5 - 4/5 - 5/5)
hypo-HDL-cholesterolaemia	insulin resistance / hyperinsulinaemia
decreased apolipoprotein A-I	dysadipokinemia (adiponectin, resistin, <i>etc.</i> ...)
HDL subtypes distribution	non-alcoholic fatty liver / steatohepatitis / NAFLD
fasting hypertriglyceridemia	chronic hyperglycaemia (<i>surrogate</i> : elevated HbA _{1c})
postprandial hypertriglyceridemia	endothelial dysfunction
TG-rich lipoprotein remnants	erectile dysfunction
apolipoprotein CIII(+)-carrying apoB lipoproteins	(micro)albuminuria
oxidized LDL	glomerular hyperfiltration
antibodies to oxidized LDL	CKD - eGFR <60 ml/min/1.73 m ²
small-dense LDL	end-stage renal failure / dialysis
lipoprotein(a)	hyperuricemia
lipoproteins glycation	hyperhomocysteinemia
lipoprotein receptors glycation	vitamin D deficiency
lipid metabolism enzymes glycation	sympathetic nervous system hyperactivity

Apo: apolipoprotein; *BMI*: body mass index; *C*: cholesterol; *CKD*: chronic kidney disease; *CVD*: cardiovascular disease; *eGFR*: estimated glomerular filtration rate; *HbA1c*: glycated haemoglobin; *HDL*: high-density lipoprotein; *LDL*: low-density lipoprotein; *NAFLD*: nonalcoholic fatty liver disease; *T2DM* type 2 diabetes mellitus; *TG*: triacylglycerols).

Table 2. Lipid, lipoproteins and cardiometabolic micro- and macrovascular disease risk factors/ markers in T2DM patients: standard, emerging and candidate

1. continuous strive towards conventional target attainment, including reinforcement of TLC, higher drug dosages, drug switches or combination therapies;
2. addressing identified barriers to chronic diseases management using an individualized patient-based approach;
3. force-driving major modifiable RFs below recommended thresholds or physiological ranges (“the lower is better” paradigm);
4. impacting upon emerging risk factors, but this often means venturing beyond guidelines, or using off-label medications, or resorting to newer therapies not always supported by evidence from randomized clinical trials.

b. Non-LDL dyslipidemia and macroangiopathic RVR
As regards non-LDL dyslipidemia, AD can be improved by TLC, fibrates, or nicotinic acid [45,85-93]. In T2DM, data from the recent *Action to Control Cardiovascular Risk in Diabetes* (ACCORD) Lipid trial reinforce the residual risk hypothesis for macroangiopathy, since despite achieving a mean LDL-C of 80 mg/dl, patients in the pre-specified subgroup with AD (defined as having both baseline TG ≥204 mg/dl and baseline HDL-C ≤34 mg/dl; numbering 17% of the ACCORD cohort) experienced a 70% higher relative rate (7% higher absolute rate) of incident major CV events (composite of CVD death, nonfatal myocardial infarction and nonfatal stroke) over a mean 4.7 year follow-up when compared to patients without AD [94,95].

In ACCORD Lipid, although in the whole T2DM cohort the main macrovascular outcome was negative, a prespecified subgroup of T2DM patients with AD treated with combination LLD therapy [simvastatin *plus* fenofibrate] had a 31% reduction in major CV events, compared to those treated with simvastatin alone (12.4% incidence [simvastatin *plus*

fenofibrate] *vs.* 17.3% [simvastatin monotherapy]), amounting to a 5% absolute RVR reduction. The number-needed-to-treat was low, calculated at 20 T2DM patients with AD receiving LLD biotherapy over 5 years to prevent one major CV event. This number compares very favourably with those from landmark intervention trials with LLD in diabetic or nondiabetic populations. When primary outcome in the whole cohort was analyzed according to gender, another prespecified analysis, data for men showed a strong indication that they benefited from combination therapy (11.2% [fenofibrate] *vs.* 13.3% events [placebo]), while data for women were not as conclusive. Thus, whereas the *P* value for interaction related to gender was significant (0.0106), the confidence interval of the effect in females (9.1% [fenofibrate] *vs.* 6.6% events [placebo]) was not only wide, but also encroached the line of unity.

In ACCORD, AD was present in only one-sixth of the population under study, a subgroup which however markedly benefited in terms of macrovascular RVR reduction from combined therapy [94,95]. Whether in real-life conditions a higher proportion of T2DM patients might benefit from combination therapy [simvastatin *plus* fenofibrate] as a result of having AD as a comorbidity, is a tentative hypothesis. In ACCORD Lipid, the low recorded AD prevalence was a result not only of inclusion criteria, sequential exclusion of tertiles, intrinsic characteristics of the study population, and a relatively high proportion of volunteers of Afro-American ethnicity, who often have naturally low-to-normal triglycerides levels and less prone to suffer from AD [46-48].

We sought to investigate whether AD prevalence was akin to that observed in the ACCORD Lipid trial in an unselected, mostly Caucasian T2DM patients sample ($n=974$), consecutively attending our academic diabetes clinic. As ACCORD Lipid cutoffs, based on study-specific tertiles are unapplicable to other T2DM populations, we selected another approach to define AD, based on the co-occurrence of low HDL-C and high TG, with cutoffs derived from the harmonized definition of the MetS [51,53,55]. We also checked patient's file in order to retrieve pre-LLD HDL-C and TG values in patients on lipid-lowering drug(s), in order to establish the true, unbiased presence of AD (**Table 3**).

With such criteria and stringent use of baseline lipids values, we observed a higher prevalence of AD (35%), similar in both genders. In our survey, mean pre-LLD triglycerides were 203 mg/dl *vs.* 167 mg/dl for current values, representing an average difference of 36 mg/dl (18%), a difference sufficient to otherwise underestimate AD prevalence unless pre-LLD TG values are available. Such a confounding effect of LLD was not observed as regards HDL-C. In this analysis, the expected differences in the general population between genders were observed for HDL-C and apolipoprotein A-I level. Female patients also had significantly higher total cholesterol and significantly lower AD ratio [$\log(\text{TG})/\text{HDL-C}$] levels (**Table 3**).

Whatever AD prevalence data, notwithstanding the beneficial effects of fenofibrate on microvascular residual risk, a substantial proportion of unselected T2DM patients may potentially benefit from combination therapy to substantially decrease a modifiable component of AD-related RVR [94,95]. Such a likely assumption needs however to be confirmed from both epidemiological sources documenting the real-life prevalence of AD, and also from prospective randomised controlled trials, in which only fenofibrate-naïve patients with untreated AD would be included, with AD defined by consensual criteria, such as MetS thresholds for non-LDL dyslipidemia or trial-dependent AD cutoffs derived *a posteriori* on tertiles of non-LDL lipids at study entry such as in ACCORD Lipid [45]. Future trial should also investigate whether triple therapy (simvastatin *plus* fenofibrate *plus*

nicotinic acid), aimed at further correction of most aspects of hypercholesterolemia and AD, will provide proportionate benefit for RVR in T2DM.

		both genders	males	females	p
<i>n</i>		974	637	337	
atherogenic dyslipidemia *	%	35	35	35	NS
pre-LLD total cholesterol	mg.dl ⁻¹	230 (43)	225 (42)	240 (43)	<0.0001
pre-LLD LDL-C	mg.dl ⁻¹	145 (35)	143 (35)	150 (35)	0.0170
pre-LLD HDL-C	mg.dl ⁻¹	47 (13)	45 (12)	52 (15)	<0.0001
pre-LLD TG	mg.dl ⁻¹	203 (167)	201 (155)	208 (193)	NS
anti-dyslipidemic drug(s)	%	65	66	63	NS
statin - fenofibrate - ezetimibe	%	53 - 21 - 3	55 - 22 - 3	51 - 18 - 3	NS
total cholesterol	mg.dl ⁻¹	179 (43)	175 (42)	186 (44)	0.0001
non-HDL-C	mg.dl ⁻¹	132 (42)	130 (41)	135 (42)	NS
apolipoprotein B ₁₀₀	mg.dl ⁻¹	90 (27)	89 (27)	92 (27)	NS
LDL-C	mg.dl ⁻¹	99 (35)	98 (35)	102 (36)	NS
HDL-C	mg.dl ⁻¹	47 (14)	44 (13)	52 (15)	<0.0001
apolipoprotein A-I	mg.dl ⁻¹	149 (30)	143 (28)	163 (31)	<0.0001
triglycerides	mg.dl ⁻¹	167 (120)	171 (128)	161 (105)	NS
log (TG).HDL-C ⁻¹		0.051 (0.025)	0.054 (0.026)	0.046 (0.021)	<0.0001

Mean (1 SD) values; * : atherogenic dyslipidemia was defined as the concurrence of low HDL-C (<40 mg/dl in males; <50 mg/dl in females) and elevated fasting triglycerides (>150 mg/dl in both genders) as defined by the NCEP-ATP III cutoffs used to define the discrete lipid components of the metabolic syndrome score. Lipid values used to define AD were baseline (pre-LLD) lipids levels in patients treated with LLD(s). HDL-C: high-density lipoprotein cholesterol; LDL-C: low-density lipoprotein cholesterol; LLD: lipid-lowering drug(s); T2DM: type 2 diabetes mellitus; TG: triglycerides. p: statistical value from Student's t test; NS: not significant (p value = or > 0.05).

Table 3. Atherogenic dyslipidemia prevalence, lipid-lowering drug(s) therapy, and baseline vs. current lipids and lipoproteins values in unselected T2DM patients

c. Microangiopathic residual vascular risk

Diabetic eye disease includes diabetic retinopathy (DRP) and non-vessel comorbidities, such as cataract or glaucoma [96]. DRP is a microangiopathy affecting the retinal blood vessels resulting predominantly from poor metabolic control, which produces microvascular occlusion and leakage, leading to progressive retinal damage which may end in complete visual loss as a result of maculopathy or ischemic/proliferative retinopathy [96-100]. Few modifiable risk factors are identified with respect to DRP: poor metabolic (i.e. glycemic) control; hypertension, especially if poorly-controlled; LDL and non-LDL dyslipidemia; anemia and possibly, tobacco smoking. The recently published ACCORD Eye substudy highlighted the limits of conventional approaches based on near-normalization of two key modifiable variables (HbA_{1c} and high blood pressure). In ACCORD, DRP outcomes were defined as a ≥3-steps DRP progression on the EDTRS scale, or development of DRP requiring laser therapy or vitrectomy [101-103].

While glycemic control intensification significantly reduced the rate of progression of DRP (by a relative 33% and an absolute 3.1%, respectively), it was associated with serious safety

concerns, as higher all-cause mortality was observed in the intensive glycemic control group [104]. This led to premature termination of the glycemic arm of the ACCORD trial. Another source of disappointment from the ACCORD trial for microvascular RVR reduction was the lack of efficacy of BP normalization to improve primary DRP outcomes [105]. These findings highlights the current pitfalls of current approaches to DRP management based on a “*the lower the better*” paradigm for the two major standard modifiable risk factors, i.e. HbA_{1c} and BP [104].

As regards lipids, landmark statin trials in T2DM or in subgroups with T2DM failed to identify a beneficial effect of LDL- lowering on any type on diabetic microangiopathies. On the other hand, epidemiological evidence links AD with RVR for diabetic retinopathy and/or nephropathy in T2DM [4-6,100,104]. The *REsidual risk, Lipids and Standard Therapies* (REALIST) studies are two *Residual Risk Reduction Initiative*-initiated worldwide epidemiological retrospective case-control surveys designed to assess AD-related macro- and microvascular residual risk. These are performed in non-diabetic and T2DM patients (REALIST MICRO) receiving current standards of care, with LDL-C levels at or near goal, treated or not with a statin. Pilot results from REALIST MICRO show a highly-significant and strong association between AD and microangiopathy incidence, even when LDL-C is controlled [107,108].

In the ACCORD Eye substudy, fenofibrate decreased RVR of DRP progression, assessed using the validated EDTRS scale, and irrespective of baseline lipids. The efficacy of fenofibrate was most obvious in patients with DRP at baseline, i.e. those in secondary DRP prevention [101-104]. Such a beneficial effect confirms the previously documented benefits of fenofibrate on DRP. This, in the *Fenofibrate Intervention and Event Lowering in Diabetes* (FIELD) study, a previous other large randomised controlled trial, a decreased requirement for DRP-related laser therapy was observed in T2DM treated with fenofibrate (*vs.* placebo). Such decreased requirement was however neither a primary outcome endpoint nor a “hard endpoint” for DRP progression assessment [88,104,106].

Results from ACCORD Eye therefore markedly raise the level of clinical evidence for a beneficial effect of this PPAR- α agonist on DRP. The FIELD study also showed that fenofibrate decreased new-onset albuminuria and diabetes-related non-traumatic amputations in T2DM patients [109]. In a recent assessment of fenofibrate's renal effects (FIELD washout sub-study), decreased albuminuria and reduced estimated glomerular filtration rate (eGFR) loss over 5 years were observed in patients allocated to the fenofibrate arm [110]. These observations confirm that fenofibrate not only reduces albuminuria or delay its onset, but also diminishes the natural history of progressive eGFR impairment in T2DM, despite an early and reversible increase in plasma creatinine [104,110-114]. As regards other LLD, future trials will establish whether targeting low HDL-C with nicotinic acid, alone or on top of background statin, reduces RVR of macroangiopathy in high-risk T2DM patients [90-93,115]. Safe and effective combination of LLDs (bi- or tritherapies) targeting most aspects of dyslipidemia are likely to become standards of care in high-risk populations such as patients with T2DM [45,75,91,104,115-118].

8. Benchmarking

Innovative approaches are needed to improve individual and overall target achievement with the aim of reducing RVR. Benchmarking targeting physicians is one of those newer approaches. The goal of the non-interventional randomised OPTIMISE trial was to

investigate in 6 European countries the effect of physician's benchmarking on quality of care, assessed according to the percentage of T2DM patients achieving pre-set targets for three key modifiable variables (HbA_{1c}, LDL-C and SBP), as recommended by international guidelines. At baseline, the percentage of patients achieving targets was highly unsatisfactory: 51% (HbA_{1c}); 27% (SBP); and 35% (LDL-C), with a mere 5% reaching all three targets at study entry. Physicians were randomly assigned to receive either benchmarked feedback or non-benchmarked feedback on their patients' modifiable outcome indicators (HbA_{1c}, fasting glycaemia, total cholesterol, HDL-C, LDL-C and TG). At study end, the percentage of patients achieving all targets almost doubled, suggesting that benchmarking may be an innovative approach to improve target attainment of modifiable variables affecting RVR in T2DM patients [21,119,120].

9. References

- Haffner SM, Lehto S, Rönnemaa T, Pyörälä K, Laakso M. Mortality from coronary heart disease in subjects with type 2 diabetes and in nondiabetic subjects with and without prior myocardial infarction. *N Engl J Med* 1998;339:229-34
- Juutilainen A, Lehto S, Rönnemaa T, Pyörälä K, Laakso M Type 2 diabetes as a 'coronary heart disease equivalent'. An 18-year prospective population-based study in Finnish subjects. *Diabetes Care* 2005;28:2901-7
- Hermans MP. Diabetic macro- and microvascular disease in type 2 diabetes. *Diabetes Vasc Dis Res* 2007;4:S7-11.
- Fruchart J-C, Sacks F, Hermans MP, Assmann G, Brown V, Ceska R, Chapman J, Dodson PM, Fioretto P, Ginsberg HN, Kadowaki T, Lablanche J-M, Marx N, Plutzky J, Reiner Z, Rosenson RS, Staels B, Stock JK, Sy R, Wanner C, Zambon A, Zimmet P. Executive statement: the Residual Risk Reduction Initiative: a call to action to reduce residual vascular risk in dyslipidemic patients. A condensed position paper by the Residual Risk Reduction Initiative (R3i). *Diab Vasc Dis Res* 2008;5:319-35.
- Fruchart J-C, Sacks F, Hermans MP, Assmann G, Brown WV, Ceska R, Chapman MJ, Dodson PM, Fioretto P, Ginsberg HN, Kadowaki T, Lablanche J-M, Marx N, Plutzky J, Reiner Z, Rosenson RS, Staels B, Stock JK, Sy R, Wanner C, Zambon A, Zimmet P for the Residual Risk Reduction Initiative. The Residual Risk Reduction Initiative: a call to action to reduce residual vascular risk in patients with dyslipidemia. *Am J Cardiol* 2008; 102(suppl): 1-34.
- Fioretto P, Dodson PM, Ziegler D, Rosenson RS. Residual microvascular risk in diabetes: unmet needs and future directions. *Nat Rev Endocrinol*. 2010;6:19-25.
- Brown G, Albers JJ, Fisher LD, Schaefer SM, Lin JT, Kaplan C, Zhao XQ, Bisson BD, Fitzpatrick VF, Dodge HT. Regression of coronary artery disease as a result of intensive lipid-lowering therapy in men with high levels of apolipoprotein B. *N Engl J Med* 1990;323:1289-98.
- LaRosa JC, Grundy SM, Waters DD, Shear C, Barter P, Fruchart JC, Gotto AM, Greten H, Kastelein JJ, Shepherd J, Wenger NK. Intensive lipid lowering with atorvastatin in patients with stable coronary disease. *N Engl J Med* 2005;352:1425-35.
- Alsheikh-Ali AA, Lin JL, Abourjaily P, Ahearn D, Kuvin JT, Karas RH. Prevalence of low high-density lipoprotein cholesterol in patients with documented coronary heart disease or risk equivalent and controlled low-density lipoprotein cholesterol. *Am J Cardiol* 2007;100:1499-1501.

- Brunzell JD, Davidson M, Furberg CD, Goldberg RB, Howard BV, Stein JH, Witztum JL. Lipoprotein management in patients with cardiometabolic risk: consensus conference report from the American Diabetes Association and the American College of Cardiology Foundation. *J Am Coll Cardiol* 2008;51:1512-24.
- Forrester JS. Redefining normal low-density lipoprotein cholesterol: a strategy to unseat coronary disease as the nation's leading killer. *J Am Coll Cardiol* 2010;56:630-6.
- Graham I, Atar D, Borch-Johnsen K, Boysen G, Burell G, Cifkova R, Dallongeville J, De Backer G, Ebrahim S, Gjelsvik B, Herrmann-Lingen C, Hoes A, Humphries S, Knapton M, Perk J, Priori SG, Pyorala K, Reiner Z, Ruilope L, Sans-Menendez S, Scholte op Reimer W, Weissberg P, Wood D, Yarnell J, Zamorano JL; Other experts who contributed to parts of the guidelines: Walma E, Fitzgerald T, Cooney MT, Dudina A; European Society of Cardiology (ESC) Committee for Practice Guidelines (CPG), Vahanian A, Camm J, De Caterina R, Dean V, Dickstein K, Funck-Brentano C, Filippatos G, Hellemans I, Kristensen SD, McGregor K, Sechtem U, Silber S, Tendera M, Widimsky P, Zamorano JL, Hellemans I, Altiner A, Bonora E, Durrington PN, Fagard R, Giampaoli S, Hemingway H, Hakansson J, Kjeldsen SE, Larsen ML, Mancina G, Manolis AJ, Orth-Gomer K, Pedersen T, Rayner M, Ryden L, Sammut M, Schneiderman N, Stalenhoef AF, Tokgözoğlu L, Wiklund O, Zampelas A. European guidelines on cardiovascular disease prevention in clinical practice. Fourth Joint Task Force of the European Society of Cardiology and other Societies on Cardiovascular Disease Prevention in Clinical Practice. Executive summary. *Eur Heart J* 2007;28:2375-414.
- Ryden L, Standl E, Bartnik M, Van den Berghe G, Betteridge J, de Boer MJ, Cosentino F, Jonsson B, Laakso M, Malmberg K, Priori S, Ostergren J, Tuomilehto J, Thrainsdottir I, Vanhorebeek I, Stramba-Badiale M, Lindgren P, Qiao Q, Priori SG, Blanc JJ, Budaj A, Camm J, Dean V, Deckers J, Dickstein K, Lekakis J, McGregor K, Metra M, Morais J, Osterspey A, Tamargo J, Zamorano JL, Deckers JW, Bertrand M, Charbonnel B, Erdmann E, Ferrannini E, Flyvbjerg A, Gohlke H, Juanatey JR, Graham I, Monteiro PF, Parhofer K, Pyorala K, Raz I, Schernthaner G, Volpe M, Wood D. Guidelines on diabetes, pre-diabetes, and cardiovascular diseases: executive summary. The Task Force on Diabetes and Cardiovascular Diseases of the European Society of Cardiology (ESC) and of the European Association for the Study of Diabetes (EASD). *Eur Heart J* 2007;28:88-136.
- Genest J, McPherson R, Frohlich J, Anderson T, Campbell N, Carpentier A, Couture P, Dufour R, Fodor G, Francis GA, Grover S, Gupta M, Hegele RA, Lau DC, Leiter L, Lewis GF, Lonn E, Mancini GB, Ng D, Pearson GJ, Sniderman A, Stone JA, Ur E. 2009 Canadian Cardiovascular Society/Canadian guidelines for the diagnosis and treatment of dyslipidemia and prevention of cardiovascular disease in the adult - 2009 recommendations. *Can J Cardiol* 2009;25:567-79.
- American Diabetes Association. Clinical practice recommendations 2010. *Diabetes Care* 2010;33:S1-100.
- Ferrieres J, Tocque-Le Gousse E, Fabry C, Hermans MP on behalf of the French CEPHEUS investigators. Assessment of lipid-lowering treatment in France. *Arch Cardiovasc Dis* 2008; 101:557-63.
- Hermans MP, Castro Cabezas M, Strandberg T, Ferrieres J, Feely J, Elisaff M, Michel G, Sansoy V. Centralized Pan-European survey on the under-treatment of

- hypercholesterolaemia (CEPHEUS): overall findings from eight countries. *Curr Med Res Opin* 2010;26:445-54.
- Hermans MP, Amoussou-Guenou KD, Ahn SA, Rousseau MF, Everaert L, Aerts A. The elusive type 2 diabetes individual achieving tight blood pressure target: A phenotypic study. *Diab Metab Syndr* 2010;4:215-9.
- Resnick HE, Foster GL, Bardsley J, Ratner RE : Achievement of American Diabetes Association clinical practice recommendations among U.S. adults with diabetes, 1999–2002: the National Health and Nutrition Examination Survey. *Diabetes Care* 2006;29:531-7.
- Munoko Th, Hermans MP. Phenotypic characterization of first generation Maghrebian migrants with type 2 diabetes: a gender-based comparison with a reference North-Caucasian Belgian cohort. *Diab Met Syndr* 2008;2:115-24.
- Hermans MP, Muls E, Michel G, Elisaf M, Brotons C, Matthys A. Optimal type 2 diabetes management including benchmarking and standard treatment. Baseline European results. *Atherosclerosis Supplements* 2010;11:44.
- Gaede P, Vedel P, Larsen N, Jensen GV, Parving HH, Pedersen O. Multifactorial intervention and cardiovascular disease in patients with type 2 diabetes. *N Engl J Med* 2003;348:383-93.
- Stratton IM, Cull CA, Adler AI, Matthews DR, Neil HA, Holman RR. Additive effects of glycaemia and blood pressure exposure on risk of complications in type 2 diabetes: a prospective observational study (UKPDS 75). *Diabetologia* 2006;49:1761-9.
- Gaede P, Lund-Andersen H, Parving HH, Pedersen O. Effect of a multifactorial intervention on mortality in type 2 diabetes. *N Engl J Med* 2008;358:580-91.
- Stevens RJ, Kothari V, Adler AI, Stratton IM; United Kingdom Prospective Diabetes Study (UKPDS) Group. The UKPDS Risk Engine: a model for the risk of coronary heart disease in type 2 diabetes. *Clin Science* 2001;101:671-9.
- Assmann G, Cullen P, Schulte H. Simple scoring scheme for calculating the risk of acute coronary events based on the 10-year follow-up of the prospective cardiovascular Münster (PROCAM) study. *Circulation* 2002;105:310-5.
- Kothari V, Stevens RJ, Adler AI, Stratton IM, Manley SE, Neil HA, Holman RR. UKPDS 60: risk of stroke in type 2 diabetes estimated by the UK Prospective Diabetes Study risk engine. *Stroke* 2002;33:1776-81.
- Stevens RJ, Coleman RL, Adler AI, Stratton IM, Matthews DR, Holman RR. UKPDS 66: risk factors for myocardial infarction case fatality and stroke case fatality in type 2 diabetes. *Diabetes Care* 2004;27:201-7.
- Coleman RL, Stevens RJ, Retnakaran R, Holman RR. Framingham, SCORE and DECODE do not provide reliable cardiovascular risk estimates in type 2 diabetes. *Diabetes Care* 2007;30:1292-3.
- Eichler K, Puhon MA, Steurer J, Bachmann LM. Prediction of first coronary events with the Framingham score: a systematic review. *Am Heart J* 2007;153:722-31.
- Wilson PW. Risk scores for prediction of coronary heart disease: an update. *Endocrinol Metab Clin North Am* 2009;38:33-44.
- Ragé M, Van Acker N, Knaapen MW, Timmers M, Streffer J, Hermans MP, Sindic C, Meert T, Plaghki L. Asymptomatic small fiber neuropathy in diabetes mellitus: investigations with intraepidermal nerve fiber density, quantitative sensory testing and laser-evoked potentials. *J Neurol*. 2011 Apr 7. [Epub ahead of print]

- Davis RA. Cell and molecular biology of the assembly and secretion of apolipoprotein B-containing lipoproteins by the liver. *Biochim Biophys Acta* 1999;1440:1-31.
- Brites FD, Bonavita CD, De Geitere C, Cloës M, Delfly B, Yael MJ, Fruchart J, Wikinski RW, Castro GR. Alterations in the main steps of reverse cholesterol transport in male patients with primary hypertriglyceridemia and low HDL-cholesterol levels. *Atherosclerosis* 2000;152:181-92.
- Dobiášová M, Frohlich J. The plasma parameter log (TG/HDL-C) as an atherogenic index: correlation with lipoprotein particle size and esterification rate in apoB-lipoprotein-depleted plasma (FERHDL). *Clin Biochem* 2001;34:583-8.
- Blasiola DA, Davis RA, Attie AD. The physiological and molecular regulation of lipoprotein assembly and secretion. *Mol Biosyst* 2007;3:608-19.
- Adiels M, Olofsson S-O, Taskinen M-R, Borén J. Overproduction of very low-density lipoproteins is the hallmark of the dyslipidemia in the metabolic syndrome. *Arterioscler Thromb Vasc Biol* 2008;28:1225-36.
- da Luz PL, Favarato D, Faria-Neto JR Jr, Lemos P, Chagas AC. High ratio of triglycerides to HDL-cholesterol predicts extensive coronary disease. *Clinics (Sao Paulo)* 2008;63:427-32.
- Bittner V, Johnson BD, Zineh I, Rogers WJ, Vido D, Marroquin OC, Bairey-Merz CN, Sopko G. The triglyceride/high-density lipoprotein cholesterol ratio predicts all-cause mortality in women with suspected myocardial ischemia: a report from the Women's Ischemia Syndrome Evaluation (WISE). *Am Heart J* 2009;157:548-55.
- Cordero A, Andrés E, Ordoñez B, León M, Laclaustra M, Grima A, Luengo E, Moreno J, Bes M, Pascual I, Civeira F, Pocoví M, Alegría E, Casasnovas JA and MESYAS (MEtabolic Syndrome Active Subjects) Study Investigators. Usefulness of triglycerides-to-high-density lipoprotein cholesterol ratio for predicting the first coronary event in men. *Am J Cardiol* 2009;104:1393-7.
- Watts GF, Ooi EMM, Chan DC. Therapeutic regulation of apoB100 metabolism in insulin resistance in vivo. *Pharmacology & Therapeutics* 2009;123:281-91.
- Assmann G, Cullen P, Schulte H. Non-LDL-related dyslipidemia and coronary risk: a case-control study. *Diab Vasc Dis Res* 2010;7:204-12.
- Costet P. Molecular pathways and agents for lowering LDL-cholesterol in addition to statins. *Pharmacol Ther* 2010;126:263-78.
- Hermans MP, Ahn SA, Rousseau MF. log(TG)/HDL-C is related to both residual cardiometabolic risk and β -cell function loss in type 2 diabetes males. *Cardiovasc Diabetol*. 2010;9:88.
- Hermans MP, Fruchart JC. Reducing residual vascular risk in patients with atherogenic dyslipidaemia: where do we go from here? *Clinical Lipidology* 2010;5:811-26.
- Sumner AE, Vega GL, Genovese DJ, Finley KB, Bergman RN, Boston RC. Normal triglyceride levels despite insulin resistance in African Americans: role of lipoprotein lipase. *Metabolism* 2005;54:902-9.
- Dehout F, Haumont S, Gaham N, Amoussou-Guenou KD, Hermans MP. Metabolic syndrome in Bantu subjects with type 2 diabetes from sub-Saharan extraction: prevalence, gender differences and HOMA-hyperbolic product. *Diab Metab Syndr* 2008;2:5-11.

- Paramsothy P, Knopp R, Bertoni AG, Tsai MY, Rue T, Heckbert SR. Combined hyperlipidemia in relation to race/ethnicity, obesity, and insulin resistance in the Multi-Ethnic Study of Atherosclerosis (MESA). *Metabolism* 2009;58:212-9.
- Bonora E, Targher G, Formentini G, Calcaterra F, Lombardi S, Marini F, Zenari L, Saggiani F, Poli M, Perbellini S, Raffaelli A, Gemma L, Santi L, Bonadonna RC, Muggeo M. The metabolic syndrome is an independent predictor of cardiovascular disease in Type 2 diabetic subjects. Prospective data from the Verona Diabetes Complications Study. *Diabet Med* 2004;21:52-8.
- Bruno G, Merletti F, Biggeri A, Barger G, Ferrero S, Runzo C, Prina Cerai S, Pagano G, Cavallo-Perin P; Casale Monferrato Study. Metabolic syndrome as a predictor of all-cause and cardiovascular mortality in Type 2 diabetes: the Casale Monferrato Study. *Diabetes Care* 2004;27:2689-94.
- Grundey SM, Cleeman JI, Daniels SR, Donato KA, Eckel RH, Franklin BA, Gordon DJ, Krauss RM, Savage PJ, Smith SC Jr, Spertus JA, Costa F; American Heart Association; National Heart, Lung, and Blood Institute. Diagnosis and management of the metabolic syndrome: an American Heart Association/National Heart, Lung, and Blood Institute Scientific Statement. *Circulation* 2005;112:2735-52.
- Metascreen Writing Committee. The metabolic syndrome is a risk indicator of microvascular and macrovascular complications in diabetes. Results from Metascreen, a multicenter diabetes clinic-based survey. *Diabetes Care* 2006;29:2701-7.
- Alberti KG, Eckel RH, Grundey SM, Zimmet PZ, Cleeman JI, Donato KA, Fruchart JC, James WP, Loria CM, Smith SC Jr; International Diabetes Federation Task Force on Epidemiology and Prevention; National Heart, Lung, and Blood Institute; American Heart Association; World Heart Federation; International Atherosclerosis Society; International Association for the Study of Obesity. Harmonizing the metabolic syndrome: a joint interim statement of the International Diabetes Federation Task Force on Epidemiology and Prevention; National Heart, Lung, and Blood Institute; American Heart Association; World Heart Federation; International Atherosclerosis Society; and International Association for the Study of Obesity. *Circulation*. 2009;120:1640-5.
- Hermans MP, Ahn SA, Amoussou-Guenou KD, Rousseau MF. Impact of metabolic syndrome on microvascular complications in type 2 diabetes. *Diab Metab Syndr* 2010;4:150-4.
- Sadikot S, Hermans MP. Here we go again ... The metabolic syndrome revisited! *Diab Metab Syndr* 2010;4:111-20.
- Davignon J, Cohn JS. Triglycerides: a risk factor for coronary heart disease. *Atherosclerosis* 1996;124(Suppl):S57-64.
- Fruchart JC, Duriez P. HDL and triglyceride as therapeutic targets. *Curr Opin Lipidol* 2002;13:605-16.
- Lu W, Resnick HE, Jablonski KA, Jones KL, Jain AK, Howard WJ, Robbins DC, Howard BV. Non-HDL cholesterol as a predictor of cardiovascular disease in type 2 diabetes : The Strong Heart Study. *Diab. Care* 2003;26:16-23.
- Fruchart JC, Nierman MC, Stroes ES, Kastelein JJ, Duriez P. New risk factors for atherosclerosis and patient risk assessment. *Circulation* 2004;109(23 Suppl 1):III15-9.

- Yusuf S, Hawken S, Ôunpuu S, Dans T, Avezum A, Lanas F et al. on behalf of the INTERHEART investigators, Effect of potentially modifiable risk factors associated with myocardial infarction in 52 countries (the INTERHEART study): case-control study. *Lancet* 2004;364:937-52.
- Assmann G, Cullen P, Fruchart JC, Greten H, Naruszewicz M, Olsson A, Paoletti R, Riesen W, Stoll M, Tikkanen M, von Eckardstein A; International Task Force for Prevention of Coronary Heart Disease. Implications of emerging risk factors for therapeutic intervention. *Nutr Metab Cardiovasc Dis* 2005;15:373-81.
- Walldius G, Jungner I. The apoB/apoA-I ratio: a strong, new risk factor for cardiovascular disease and a target for lipid-lowering therapy--a review of the evidence. *J Intern Med* 2006;259:493-519.
- Assmann G, Schulte H, Cullen P, Seedorf U. Assessing risk of myocardial infarction and stroke: new data from the Prospective Cardiovascular Münster (PROCAM) study. *Eur J Clin Invest* 2007;37:925-32.
- Bansal S, Buring JE, Rifai N, Mora S, Sacks FM, Ridker PM. Fasting compared with nonfasting triglycerides and risk of cardiovascular events in women. *JAMA* 2007;298:309-16.
- Hermans MP, Ahn SA, Rousseau MF. The non-HDL-C/HDL-C ratio provides cardiovascular risk stratification similar to the ApoB/ApoA1 ratio in diabetics: Comparison with reference lipid markers. *Diab Metab Syndr* 2007;1:23-8.
- Assmann G, Schulte H, Seedorf U. Cardiovascular risk assessment in the metabolic syndrome: results from the Prospective Cardiovascular Munster (PROCAM) Study. *Int J Obes (Lond)* 2008;32(Suppl 2):S11-6.
- Kawakami A, Osaka M, Tani M, Azuma H, Sacks FM, Shimokado K, Yoshida M. Apolipoprotein CIII links hyperlipidemia with vascular endothelial cell dysfunction. *Circulation* 2008;118:731-42.
- Ridker PM, Danielson E, Fonseca FA, Genest J, Gotto AM Jr, Kastelein JJ, Koenig W, Libby P, Lorenzatti AJ, MacFadyen JG, Nordestgaard BG, Shepherd J, Willerson JT, Glynn RJ; JUPITER Study Group. Rosuvastatin to prevent vascular events in men and women with elevated C-reactive protein. *N Engl J Med* 2008;359:2195-207.
- Hermans MP, Ahn SA, Rousseau MF. Cardiometabolic phenotype and UKPDS risk in male type 2 diabetic patients with obstructive sleep apnoea syndrome. *Diabetes Metab Syndr* 2009;3:50-4.
- Hermans MP, Ahn SA, Rousseau MF. Erectile dysfunction, microangiopathy and UKPDS risk in type 2 diabetes. *Diabetes Metab* 2009;35:484-9.
- Mendivil CO, Zheng C, Furtado J, Lel J, Sacks FM. Metabolism of very-low-density lipoprotein and low-density lipoprotein containing apolipoprotein C-III and not other small apolipoproteins. *Arterioscler Thromb Vasc Biol* 2010;30:239-45.
- Sarwar N, Sandhu MS, Ricketts SL, Butterworth AS, Di Angelantonio E, Boekholdt SM, Ouwehand W, Watkins H, Samani NJ, Saleheen D, Lawlor D, Reilly MP, Hingorani AD, Talmud PJ, Danesh J. Triglyceride Coronary Disease Genetics Consortium and Emerging Risk Factors Collaboration. Triglyceride-mediated pathways and coronary disease: collaborative analysis of 101 studies. *Lancet* 2010;375:1634-9.
- Zheng C, Khoo C, Furtado J, Sacks FM. Apolipoprotein C-III and the metabolic basis for hypertriglyceridemia and the dense low-density lipoprotein phenotype. *Circulation* 2010;121:1722-34.


- Hermans MP, Sacks FM, Ahn SA, Rousseau MF. Non-HDL-cholesterol as valid surrogate to apolipoprotein B100 measurement in diabetes: discriminant ratio and unbiased equivalence. *Cardiovasc Diabetol*. 2011;10:20.
- Davidson MH. Reducing residual risk for patients on statin therapy: the potential role of combination therapy. *Am J Cardiol* 2005;96(9A):3K-13K.
- Libby PL. The forgotten majority. Unfinished business in cardiovascular risk reduction. *J Am Coll Cardiol* 2005;46:1225-8.
- Shepherd J, Barter P, Carmena R, Deedwania P, Fruchart JC, Haffner S, Hsia J, Breazna A, LaRosa J, Grundy S, Waters D for the Treating to New Targets Investigators. Effect of lowering LDL cholesterol substantially below currently recommended levels in patients with coronary heart disease and diabetes: The Treating to New Targets (TNT) study. *Diabetes Care* 2006;29:1220-6.
- Barter P, Gotto AM, LaRosa JC, Maroni J, Szarek M, Grundy SM, Kastelein JJ, Bittner V, Fruchart JC. Treating to New Targets Investigators. HDL cholesterol, very low levels of LDL cholesterol, and cardiovascular events. *N Engl J Med* 2007;357:1301-10.
- Friedewald VE Jr, Ballantyne CM, Davidson MH, Guyton JR, Roberts WC. The editor's roundtable: lipid management beyond statins - reducing residual cardiovascular risk. *Am J Cardiol* 2008;102:559-67.
- Jones PH. Expert perspective: reducing cardiovascular risk in metabolic syndrome and type 2 diabetes mellitus beyond low-density lipoprotein cholesterol lowering. *Am J Cardiol* 2008;102:41L-7L.
- Miller M, Cannon CP, Murphy SA, Qin J, Ray KK, Braunwald E; PROVE-IT TIMI 22 Investigators. Impact of triglyceride levels beyond low-density lipoprotein cholesterol after acute coronary syndrome in the PROVE-IT TIMI 22 trial. *J Am Coll Cardiol* 2008;51:724-30.
- Schramm TK, Gislason GH, Køber L, Rasmussen S, Rasmussen JN, Abildstrøm SZ, Hansen ML, Folke F, Buch P, Madsen M, Vaag A, Torp-Pedersen C. Diabetes patients requiring glucose-lowering therapy and nondiabetics with a prior myocardial infarction carry the same cardiovascular risk: a population study of 3.3 million people. *Circulation* 2008;117:1945-54.
- Sulkes D, Brown BG, Krauss RM, Segrest JP, Sniderman AD, Roberts WC. The editor's roundtable: expanded versus standard lipid panels in assessing and managing cardiovascular risk. *Am J Cardiol* 2008;101:828-42.
- Ridker PM, Genest J, Boekholdt SM, Libby P, Gotto AM, Nordestgaard BG, Mora S, MacFadyen JG, Glynn RJ, Kastelein JJ; JUPITER Trial Study Group. HDL cholesterol and residual risk of first cardiovascular events after treatment with potent statin therapy: an analysis from the JUPITER trial. *Lancet* 2010;376:333-9.
- Volek JS, Fernandez ML, Feinman RD, Phinney SD. Dietary carbohydrate restriction induces a unique metabolic state positively affecting atherogenic dyslipidemia, fatty acid partitioning, and metabolic syndrome. *Prog Lipid Res* 2008;47:307-18.
- Katcher HI, Hill AM, Lanford JL, Yoo JS, Kris-Etherton PM. Lifestyle approaches and dietary strategies to lower LDL-cholesterol and triglycerides and raise HDL-cholesterol. *Endocrinol Metab Clin North Am* 2009;38:45-78.
- Fruchart JC. Peroxisome proliferator-activated receptor- α (PPAR α): at the crossroads of obesity, diabetes and cardiovascular disease. *Atherosclerosis* 2009;205:1-8.

- Vergès B. Role for fibrate therapy in diabetes: evidence before FIELD. *Curr Opin Lipidol* 2005;16:648-51.
- Jun M, Foote C, Lv J, Neal B, Patel A, Nicholls SJ, Grobbee DE, Cass A, Chalmers J, Perkovic V. Effects of fibrates on cardiovascular outcomes: a systematic review and meta-analysis. *Lancet* 2010;375:1875-84.
- Taylor AJ, Villines TC, Stanek E, Devine PJ, Griffen L, Miller M, Weissman NJ, Turco M. Extended-release niacin or ezetimibe and carotid intima-media thickness. *New Engl J Med* 2009;361: 2113-22.
- Chapman MJ, Redfern JS, McGovern ME, Giral P. Niacin and fibrates in atherogenic dyslipidemia: pharmacotherapy to reduce cardiovascular risk. *Pharmacol Ther* 2010;126:314-45.
- AIM HIGH: Niacin plus statin to prevent vascular events. *ClinicalTrials.gov* identifier: NCT00120289.
- HPS2-THRIVE. Treatment of HDL to reduce the incidence of vascular events. *ClinicalTrials.gov* Identifier: NCT00461630.
- ACCORD Study Group, Ginsberg HN, Elam MB, Lovato LC, Crouse JR 3rd, Leiter LA, Linz P, Friedewald WT, Buse JB, Gerstein HC, Probstfield J, Grimm RH, Ismail-Beigi F, Bigger JT, Goff DC Jr, Cushman WC, Simons-Morton DG, Byington RP. Effects of combination lipid therapy in type 2 diabetes mellitus. *N Engl J Med* 2010;362:1563-74.
- Fruchart J-C, Sacks FM, Hermans MP. Implications of the ACCORD lipid study : perspective from the Residual Risk Reduction Initiative (R3i). *Curr Med Res Opin* 2010;26:1793-7.
- Hermans MP, Ahn SA, Rousseau MF. Statin therapy and cataract in Type 2 diabetes. *Diabetes Metab* 2011;37:139-43.
- UK Prospective Diabetes Study Group. Tight blood pressure control and risk of macrovascular and microvascular complications in type 2 diabetes. UKPDS 38. *BMJ* 1998;317:703-13.
- Misra A, Kumar S, Kishore Vikram A, Kumar A. The role of lipids in the development of diabetic microvascular complications: implications for therapy. *Am J Cardiovasc Drugs* 2003;3:325-338.
- Dodson PM. Medical treatment for diabetic retinopathy: do the FIELD microvascular study results support a role for lipid lowering? *Practical Diabetes International* 2008;25:76-79.
- Dodson PM. Diabetes and the Eye. In: *Diabetic Retinopathy*, Dodson PM (ed). Oxford University Press, Oxford, pp 33-44, 2009.
- Chew EY, Ambrosius WT, Howard LT, Greven CM, Johnson S, Danis RP, Davis MD, Genuth S, Domanski M; ACCORD Study Group. Rationale, design, and methods of the Action to Control Cardiovascular Risk in Diabetes Eye Study (ACCORD-EYE). *Am J Cardiol* 2007;99(12A):103i-11i.
- ACCORD Study Group; ACCORD Eye Study Group, Chew EY, Ambrosius WT, Davis MD, Danis RP, Gangaputra S, Greven CM, Hubbard L, Esser BA, Lovato JF, Perdue LH, Goff DC Jr, Cushman WC, Ginsberg HN, Elam MB, Genuth S, Gerstein HC, Schubart U, Fine LJ. Effects of medical therapies on retinopathy progression in type 2 diabetes. *N Engl J Med* 2010;363:233-44.

- Matthews DR. Fenofibrate and statin therapy, compared with placebo and statin, slows the development of retinopathy in type 2 diabetes patients of 10 years duration: the ACCORD study. *Evid Based Med* 2011 Apr;16(2):45-6.
- Hermans MP. Prevention of microvascular diabetic complications by fenofibrate: lessons from FIELD and ACCORD. *Diabetes Vasc Dis Res* 2011 (*in press*).
- Cushman WC, Grimm RH, Cutler JA for the ACCORD study group. Rationale and Design for the Blood Pressure Intervention of the Action to Control Cardiovascular Risk in Diabetes (ACCORD) Trial. *Am J Cardiol* 2007;99(suppl):44i-55i.
- Hermans MP. Impact of fenofibrate on type 2 diabetes patients with features of the metabolic syndrome: subgroup analysis from FIELD. *Curr Cardiol Rev* 2010;6:112-8.
- Carey VJ, Bishop L, Laranjo N, Harschfield BJ, Kwiat C, Sacks FM. Contribution of high plasma triglycerides and low high-density lipoprotein cholesterol to residual risk of coronary heart disease after establishment of low-density lipoprotein cholesterol control. *Am J Cardiol* 2010;106:757-63.
- Rajamani K, Colman PG, Li LP, Best JD, Voysey M, D'Emden MC, Laakso M, Baker JR, Keech AC; FIELD study investigators. Effect of fenofibrate on amputation events in people with type 2 diabetes mellitus (FIELD study): a prespecified analysis of a randomised controlled trial. *Lancet* 2009;373:1780-8.
- Millán Núñez-Cortés J, Montoya JP, Salas XP, Hernández Mijares A, Carey VJ, Hermans MP, Sacks FM, Fruchart JC. The REALIST (REsiduAl risk, LIpids and Standard Therapies) study: an analysis of residual risk attributable to lipid profile in acute coronary syndrome. *Endocrinol Nutr* 2011;58:38-47.
- Davis TM, Ting R, Best JD, Donoghoe MW, Drury PL, Sullivan DR, Jenkins AJ, O'Connell RL, Whiting MJ, Glasziou PP, Simes RJ, Kesäniemi YA, GebSKI VJ, Scott RS, Keech AC; Fenofibrate Intervention and Event Lowering in Diabetes Study investigators. Effects of fenofibrate on renal function in patients with type 2 diabetes mellitus: the Fenofibrate Intervention and Event Lowering in Diabetes (FIELD) Study. *Diabetologia* 2011;54:280-90.
- Keech A, Simes RJ, Barter P, Best J, Scott R, Taskinen MR, Forder P, Pillai A, Davis T, Glasziou P, Drury P, Kesaniemi YA, Sullivan D, Hunt D, Colman P, d'Emden M, Whiting M, Ehnholm C, Laakso M. Effects of long-term fenofibrate therapy on cardiovascular events in 9795 people with type 2 diabetes mellitus (the FIELD study): randomised controlled trial. *Lancet* 2005;366:1849-61.
- Scott R, O'Brien R, Fulcher G, Pardy C, D'Emden M, Tse D, Taskinen MR, Ehnholm C, Keech A; Fenofibrate Intervention and Event Lowering in Diabetes (FIELD) Study Investigators. The effects of fenofibrate treatment on cardiovascular disease risk in 9795 people with type 2 diabetes and various components of the metabolic syndrome: the FIELD study. *Diabetes Care* 2009;32:493-8.
- Effect of fenofibrate on progression of coronary-artery disease in type 2 diabetes: the Diabetes Atherosclerosis Intervention Study, a randomised study. *Lancet* 2001;357:905-10.
- Ansquer JC, Foucher C, Rattier S, Taskinen MR, Steiner G; DAIS Investigators. Fenofibrate reduces progression to microalbuminuria over 3 years in a placebo-controlled study in type 2 diabetes: results from the Diabetes Atherosclerosis Intervention Study (DAIS). *Am J Kidney Dis* 2005;45:485-93.

- Polonsky TS, Davidson MH. Reducing the residual risk of 3-hydroxy-3-methylglutaryl coenzyme A reductase inhibitor therapy with combination therapy. *Am J Cardiol* 2008;101(8A):27B-35B.
- Rosenson RS. Management of non-high-density lipoprotein abnormalities. *Atherosclerosis* 2009;207:328-35.
- Rosenson RS, Davidson MH, Pourfarzib R. Underappreciated opportunities for low-density lipoprotein management in patients with cardiometabolic residual risk. *Atherosclerosis* 2010;213/1-7.
- Sacks FM, Carey VJ, Fruchart JC. Combination lipid therapy in type 2 diabetes. *N Engl J Med* 2010;363:692-4.
- Hermans MP, Muls E, Nobels F, Krzentowski G, Claes N, Debacker N, Matthys A. Evaluating benchmarking to optimize management of type 2 diabetes patients: the Belgian data from the OPTIMISE study. *Acta Cardiol* 2011;66:124-5.
- Brotons C, Elisaf M, Hermans MP, Michel G, Muls E, Matthys A. Evaluating benchmarking to optimize management of type 2 diabetic patients: HbA_{1c} control in the European OPTIMISE study. *Diabetes* 2011 (*in press*).

IntechOpen



Recent Advances in the Pathogenesis, Prevention and Management of Type 2 Diabetes and its Complications

Edited by Prof. Mark Zimring

ISBN 978-953-307-597-6

Hard cover, 442 pages

Publisher InTech

Published online 29, August, 2011

Published in print edition August, 2011

Type 2 diabetes “œmellitus” affects nearly 120 million persons worldwide- and according to the World Health Organization this number is expected to double by the year 2030. Owing to a rapidly increasing disease prevalence, the medical, social and economic burdens associated with the microvascular and macrovascular complications of type 2 diabetes are likely to increase dramatically in the coming decades. In this volume, leading contributors to the field review the pathogenesis, treatment and management of type 2 diabetes and its complications. They provide invaluable insight and share their discoveries about potentially important new techniques for the diagnosis, treatment and prevention of diabetic complications.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Michel P. Hermans, Sylvie A. Ahn and Michel F. Rousseau (2011). Residual Vascular Risk in T2DM: The Next Frontier, Recent Advances in the Pathogenesis, Prevention and Management of Type 2 Diabetes and its Complications, Prof. Mark Zimring (Ed.), ISBN: 978-953-307-597-6, InTech, Available from: <http://www.intechopen.com/books/recent-advances-in-the-pathogenesis-prevention-and-management-of-type-2-diabetes-and-its-complications/residual-vascular-risk-in-t2dm-the-next-frontier>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2011 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike-3.0 License](https://creativecommons.org/licenses/by-nc-sa/3.0/), which permits use, distribution and reproduction for non-commercial purposes, provided the original is properly cited and derivative works building on this content are distributed under the same license.

IntechOpen

IntechOpen