

# We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

186,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index  
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?  
Contact [book.department@intechopen.com](mailto:book.department@intechopen.com)

Numbers displayed above are based on latest data collected.  
For more information visit [www.intechopen.com](http://www.intechopen.com)



# Do Socio-Demographic Characteristics Modify the Association Between Air Pollution and Mortality & Morbidity?

Sabit Cakmak<sup>1</sup>, Sara L. Martin<sup>4</sup>, Claudia Blanco Vidal<sup>2</sup>, Timur Gultekin<sup>3</sup>, Vladislav Brion<sup>4</sup> and Maria Angelica Rubio<sup>5</sup>

<sup>1</sup>Health Canada, Ottawa, Department of Statistics,

<sup>2</sup>Area Descontaminacion Atmosferica, Comisi3n Nacional del Medio Ambiente (CONAMA),

<sup>3</sup>Department of Anthropology, Ankara University,

<sup>4</sup>Health Canada, Ottawa, Department of Statistics

<sup>5</sup>Departamento de Ciencias del Ambiente, Facultad de Qu3mica y Biolog3a, Universidad de Santiago de Chile

Centro de Desarrollo para la Nanociencia y Nanotecnolog3a (CEDENNA),  
<sup>1,4</sup>Canada

<sup>2,5</sup>Chile

<sup>3</sup>Turkey

## 1. Introduction

Historical extreme air pollution events such as those experienced in London in the 1950s and 60s clearly demonstrated the potential of ambient air pollution to cause exacerbation of cardio-respiratory disease, manifested as pre-mature mortality and admission to hospital. In the intervening years, considerable efforts have been made to reduce pollution from the combustion of fossil fuels and industrial activity. Although these pollution mitigation strategies have been largely viewed as successful, evidence from population health studies in North America, Europe, South America, Mexico, Asia, Australia and New Zealand continues to identify ambient air quality as a population health concern (Table 1).

Reference	Data	Location	Focus	Outcomes	Subpop.
Alberdi et al., 1998	1986-1992	Madrid, Spain	TSP, SO <sub>2</sub>	NA, R, CV	Sex, Age >65
Alberdi et al., 1998b	1986-1992	Madrid, Spain	TSP, SO <sub>2</sub>	NA, R, CV	Sex, Age >65
Anderson et al., 1996	1987-1992	London, England	BS, SO <sub>2</sub> , NO <sub>2</sub> , O <sub>3</sub>	NA, R, CV	
Bachárová et al., 1996	1987-1991	Slovak Republic	TSP, SO <sub>2</sub>	A, NA, R, CV	
Ballester et al., 1996	1991-1993	Valencia, Spain	TSP, SO <sub>2</sub>	NA, R, CV	Age >70

Borja-Aburto et al., 1997	1990-1992	Mexico City, Mexico	O <sub>3</sub> , SO <sub>2</sub> , TSP	A, NA, R, CV	Age
Borja-Aburto et al., 1998	1993-1995	Mexico City, Mexico	PM <sub>2.5</sub>	NA, R, CV	Age >65
Boucher et al., 1996	1985-1993	Salt Lake and Utah counties, U.S.	PM <sub>10</sub>	NA	
Burnett et al., 1998	1980-1994	Toronto, Canada	CO, NO <sub>2</sub> , SO <sub>2</sub> , TSP, PM <sub>2.5</sub> , PM <sub>10</sub>	NA, R, CV, O	Age
Burnett et al., 1998b	1980-1991	11 Canadian Cities	CO, NO <sub>2</sub> , SO <sub>2</sub> , O <sub>3</sub>	NA	
Burnett et al., 2000	1986-1996	8 Canadian Cities	PM <sub>2.5</sub> , PM <sub>10</sub> , PM <sub>10-2.5</sub>	NA	
Castillejos et al., 2000	1992-1995	Mexico City, Mexico	PM <sub>2.5</sub> , PM <sub>10</sub> , PM <sub>10-2.5</sub>	NA	
Chock et al., 2000	1989-1991	Pittsburg, Pennsylvania, U.S.	PM <sub>2.5</sub> , PM <sub>10</sub>	NA, R, CV	Age
Cifuenties et al, 2000	1988-1996	Santiago, Chile	PM <sub>2.5</sub> , PM <sub>10</sub> , CO, NO <sub>2</sub> , SO <sub>2</sub> , O <sub>3</sub>	NA	
Dab et al., 1996	1987-1992	Paris, France	BS, SO <sub>2</sub> , O <sub>3</sub> , NO <sub>2</sub> , PM <sub>13</sub>	R morbidity and mortality	
Daniels et al., 2000	1987-1994	20 cities, U.S.	PM <sub>10</sub>	A, CV	
Díaz et al., 1999	1990-1996	Madrid, Spain	TSP, SO <sub>2</sub> , NO <sub>2</sub> , NO <sub>x</sub> , O <sub>3</sub>	R, CV, Emergency hospital admission (94-96)	
Dockery et al., 1992	1985-1986	St. Louis, Illinois and Missouri; Roanne county, Tennessee	PM <sub>2.5</sub> , PM <sub>10</sub> , Aerosols	Mortality	
Fairley, 1990	1980-1986	Santa Clara County, California	PM <sub>10</sub>	NA	
Fairley, 1999	1989-1996	Santa Clara County, California	PM <sub>2.5</sub>	NA, R, CV	
Reference	Data	Location	Focus	Outcomes	Subpop.
Goldberg et al., 2000	1995-1999	Montreal, Quebec	predicted PM <sub>2.5</sub>	NA, CV, C	Age

Gouveia& Fletcher, 2000	1991-1993	São Paulo, Brazil	PM <sub>2.5</sub> , PM <sub>10</sub> , CO, NO <sub>2</sub> , SO <sub>2</sub> , O <sub>3</sub>	NA, R, CV	Socioeco. sex, age
Hales et al., 2000	1988-1993	Christchurch, New Zealand	PM <sub>10</sub> , NO <sub>x</sub> , SO <sub>2</sub> , O <sub>3</sub> , CO	NA, R, CV	Age >65
Gwynn et al., 2000	1988-1990	Buffalo, New York, U.S.	H <sup>+</sup> and SO <sub>4</sub> <sup>2-</sup> PM	R, CV and A mortality and morbidity	
Hatzakis et al., 1986	1975-1982	Athens, Greece	BS, SO <sub>2</sub>	Mortality	
Hoek et al., 1997	1983-1991	Rotterdam, Netherlands	TSP, BS, SO <sub>2</sub> , O <sub>3</sub> , CO	Mortality	
Hoek et al., 2000	1986-1994	The Netherlands	PM <sub>10</sub> , BS	NA, R, CV	
Hong et al., 1999	1995-1996	Inchon, South Korea	PM <sub>10</sub> , SO <sub>2</sub> ,CO, O <sub>3</sub>	NA, R, CV	
Ito et al., 1993	1965-1972	London, England	BS, SO <sub>2</sub> , Acidic Aerosols	Mortality	
Ito et al., 1995	1985-1990	Cook County, Illinois and Los Angeles County, California, U.S.	PM <sub>10</sub>	Mortality	
Ito et al., 1996	1985-1990	Cook County, Illinois, U.S.	PM <sub>10</sub>	R, CV, C	Age, Sex, Race
Kelsall et al., 1997	1974-1988	Philadelphia, Pennsylvania, U.S.	TSP	NA, R, CV	Age
Kinney & Özkaynak, 1991	1970 - 1979	Los Angeles, County, California, U.S.	O <sub>x</sub> , SO <sub>4</sub> , NO <sub>2</sub> , CO	NA, R, CV	
Kinney et al., 1995	1985-1990	Los Angeles, County, California, U.S.	PM <sub>10</sub>	NA	
Klemm & Mason, 2000	1998-1999	Atlanta, Georgia, U.S.	PM <sub>2.5</sub>	NA	Age (>65)
Kotesovec et al., 2000	1982-1994	Northern Bohemia, Czech Republic	TSP, SO <sub>2</sub>	A, CV and C	Sex, Age
Krzyzanowski & Wojtyniak 1991/92	1977-1989	Cracow, Poland	SO <sub>2</sub> , PM <sub>20</sub>	NA, R, CV	Sex, Age
Le Tertre et al., 1998	1987-1990	Paris, France	SO <sub>2</sub>	Mortality	

Lee et al., 1999	1991-1995	Seoul and Ulsan, South Korea	TSP, SO <sub>2</sub> , O <sub>3</sub>	NA	
Lee et al., 2000	1991-1997	7 South Korean cities	TSP, SO <sub>2</sub> , O <sub>3</sub>	NA	
Reference	Data	Location	Focus	Outcomes	Subpop.
Lipfert et al., 2000	1992-1995	Philadelphia, Pennsylvania	TSP	Mortality	
Lippmann et al., 2000	Various	Detroit, Michigan, U.S.	H <sup>+</sup> and SO <sub>4</sub> <sup>2-</sup> PM	Mortality and elderly morbidity	Age
Lyon et al., 1995	1985-1992	Utah County, U.S.	PM <sub>10</sub>	NA, R, CV, O	Age
Machenbach et al., 1993	1979-1987	The Netherlands	SO <sub>2</sub>	Mortality	
Mar et al., 2000	1995-1997	Phoenix, Arizona, U.S.	PM <sub>2.5</sub> , PM <sub>10</sub> , PM <sub>10-2.5</sub>	NA, R, CV	
Michelozzi et al., 1998	1992-1995	Rome, Italy	PM <sub>10</sub> , SO <sub>2</sub> , O <sub>3</sub> , NO <sub>2</sub> , CO	Mortality	
Moolgavak, 2000	1987-1995	3 U.S. Counties	PM <sub>10</sub> , CO, O <sub>3</sub>	NA, R, CV	
Moolgavakar et al., 1995	1974-1984	Steubenville, Ohio, U.S.	TSP, SO <sub>2</sub>	NA	
Moolgavakar et al., 1995b	1973-1988	Philadelphia, Pennsylvania, U.S.	TSP, SO <sub>2</sub> , O <sub>3</sub>	NA	
Morgan et al., 1998	1989-1993	Sydney, Australia	PM, NO <sub>2</sub> , O <sub>3</sub>	NA, R, CV	
Ostro, 1995	1980-1986	California, U.S.	PM <sub>2.5</sub>	NA, R, CV	
Ostro et al., 1996	1989-1991	Santiago, Chile	PM <sub>10</sub>	NA, R, CV	Age >65
Ostro et al., 1999	1989-1992	Coachella Valley, California, U.S.	PM <sub>10</sub>	NA, R, CV	
Ostro et al., 1999b	1992-1995	Bangkok, Thailand	PM <sub>10</sub>	R, CV	Age
Ostro et al., 2000	1989-1998	Coachella Valley, California, U.S.	PM <sub>10</sub>	NA, R, CV	
Peters et al., 2000	1982-1991	Coal districts, Czech Republic; Bavaria n districts, Germany	TSP, SO <sub>2</sub> , PM <sub>2.5</sub> , PM <sub>10</sub>	NA, R(Czech Republic only), CV	
Pope et al., 1992	1985-1989	Utah County	PM <sub>10</sub>	NA, R, CV, O	
Pope et al., 1996	1985-1989	Utah County	PM <sub>10</sub>	NA, R, CV, O	

Pope et al., 1999	1985-1995	Wasatch Front, Utah	PM <sub>10</sub>	NA, R, CV, O	
Rahlenbeck & Kahl, 1996	1981-1989	East Berlin, Germany	TSP, SO <sub>2</sub>	NA	
Rossi et al., 1999	1980-1989	Milan, Italy	TSP, SO <sub>2</sub> , NO <sub>2</sub> , PM <sub>13</sub>	NA, R, CV	
Saldiva et al., 1995	1990-1991	São Paulo, Brazil	PM <sub>10</sub>	NA	
Samet et al., 1998	1973-1980	Philadelphia, Pennsylvania	TSP, SO <sub>2</sub>	NA	
Reference	Data	Location	Focus	Outcomes	Subpop.
Samet et al., 2000	1987-1994	20 cities, U.S.	PM <sub>10</sub>	NA, R, CV	
Schwartz & Dockery, 1992	1974-1984	Steubenville, Ohio	TSP, SO <sub>2</sub>	NA	
Schwartz & Dockery, 1996	1973-1980	Philadelphia, Pennsylvania	TSP, SO <sub>2</sub>	NA, R, CV	Age
Schwartz et al., 1990	1958-1972	London, England	BS, SO <sub>2</sub>	Non-traumatic	
Schwartz, 1991	1973-1982	Detroit, Michigan, U.S.	TSP	NA	
Schwartz, 1993	1985-1988	Birmingham, Alabama, U.S.	PM <sub>10</sub>	NA	
Schwartz, 1994	1977 - 1982	Cincinnati, Ohio, U.S.	TSP	A, R, CV	Age >65
Schwartz, 2000	1986-1993	10 cities, U.S.	PM <sub>10</sub>	Mortality	Socioeco.
Schwartz, 2000b	1986-1993	10 cities, U.S.	PM <sub>10</sub>	Mortality	Age >65
Schwartz, 2000c	1979-1986	Boston, Massachusetts, U.S.	PM <sub>2.5</sub>	A, R, CV	
Schwartz, 2000d	1974-1988	Philadelphia, Pennsylvania	TSP and SO <sub>2</sub>	NA	
Simpson et al., 1997	1987-1993	Brisbane, Australia	PM <sub>10</sub> , SO <sub>2</sub> , O <sub>3</sub>	NA, R, CV	Age >65
Simpson et al., 2000	1991-1996	Melbourne, Australia	PM <sub>2.5</sub> , PM <sub>10</sub>	NA, R, CV	Age
Smith et al., 1999	Various	Alabama & Illinois, U.S.	PM <sub>10</sub>	Mortality	Age >65
Spix & Wichmann, 1996	1976-1985	Köln, Germany	TSP, SO <sub>2</sub> , NO <sub>2</sub>	Mortality	
Spix et al., 1993	1980-1989	Erfurt, Germany	TSP, SO <sub>2</sub>	Mortality	

Styer et al., 1995	1985-1990	Utah, U.S.	PM <sub>10</sub>	NA	Age, Sex, Race
Sunyer et al., 1996	1985-1991	Barcelona, Spain	BS, NO <sub>2</sub> , SO <sub>2</sub> , O <sub>3</sub>	NA, R, CV	Age >70
Szafraniec et al., 1997	1993-1996	Kraków, Poland	SO <sub>2</sub> , PM <sub>10</sub>	NA, CV	Sex
Tobias & Campbell, 1999	1991-1995	Barcelona, Spain	BS	Mortality	
Touloumi et al., 1994	1984-1988	Athens, Greece	BS, CO, SO <sub>2</sub>	Mortality	
Touloumi et al., 1996	1987-1991	Athens, Greece	BS, CO, SO <sub>2</sub>	Mortality	
Touloumi et al., 1997	Various	6 European Cities	NO <sub>2</sub> , O <sub>3</sub>	Mortality	
Vigotti et al., 1996	1980-1989	Milan, Italy	TSP, SO <sub>2</sub>	Respiratory morbidity	
Reference	Data	Location	Focus	Outcomes	Subpop.
Wichmann et al., 2000	1991-2002	Erfurt, Germany	CO, NO <sub>2</sub> , SO <sub>2</sub> , O <sub>3</sub> , PM <sub>10</sub>	Mortality	
Wietlishbach et al., 1996	1984-1989	Zurich, Basle and Geneva, Switerland	TSP, CO, NO <sub>2</sub> , SO <sub>2</sub> , O <sub>3</sub>	NA, R, CV	Age >65
Wojtyniak & Piekarski, 1996	Various	Cracow, Lodz, Poznan and Wroclaw, Poland	SO <sub>2</sub> , BS	NA, R, CV, D	
Wordley et al., 1997	1992-1994	Brimingham, U.K.	PM <sub>10</sub>	R,CV morbidity	
X. Xu et al., 1994	1989	Dongchen and Xichen, Beijing, China	TSP, SO <sub>2</sub>	NA, R, CV, C	
Z. Y. Xu et al., 2000	1992	Shenyang, China	TSP, SO <sub>2</sub>	NA, R, CV, C, O	
Zanobetti & Schwartz, 2000	1986-1993	4 U.S. cities	PM <sub>10</sub>	NA	Sex, Race, Edu.
Zmirou et al., 1996	1985-1990	Lyon, France	S0 <sub>2</sub> , NO <sub>2</sub> , O <sub>3</sub> , PM <sub>13</sub>	NA, R, CV, D	

Table 1. Selected references examining air quality and health outcomes around the world with information on the years in which data was collected, as well as the location, the compounds, the health outcomes and subpopulations studied. In the compound column BS indicates black smoke and TSP indicates total suspended particulates. Cause of death categories studied in each paper were coded as A (accidental), NA (Non-accidental), R (respiratory including lung and chronic obstructive pulmonary disease), CV (cardiovascular or circulatory diseases), C (cancer), D (digestive system) and O (other).



Previous work that has found increases in morbidity and mortality are associated with both ambient air pollution and low socio-economic status (Dockery & Pop, 2002; Brunekreef & Holgate, 2002; Bascom et al., 1996; Hahn et al., 1996; Carr et al., 1992, Chen et al., 2001). However, the literature regarding the effect of age, gender, and social status is conflicting with some studies documenting increased susceptibility studies (Cifuentes et al., 1999; Wojtyniak & Wysocki, 1989; Health Effects Institute[HEI], 2000; Pope, 2000 ) and others finding little or no effect (Gouveia & Fletcher, 2000; Samet et al., 2000; Zanobetti et al., 2000). A variety of factors have been implicated in the increased susceptibility to air pollution among the socially disadvantaged including, higher pollutant levels in living or working areas, increased cigarette smoking, fewer dietary fruits and vegetables, and reduced access to medical care (O'Neill et al., 2003, Sexton et al. 1993). However, identification of subgroups which are more susceptible to the effects of air pollution is important for three reasons: 1) developing targeted intervention programs; 2) determining whether the air pollution-health effects found in one region can be extrapolated to other geographic regions; 3) setting effective air pollution policies that reduce risk for the entire population. This study investigates whether age, gender and an indicator of social status – educational attainment – modify the effect of particulate air pollution on mortality.

## 2. Methods

### 2.1 Air pollution data

Daily air pollution data for the nine communities (communas) that make up the Concepcion Region (Fig. 1.), Tomé, Penco, Talcahuano, Hualpén, Concepción, San Pedro de la Paz, Chiguayante, Lota, and Coronel, were obtained from monitoring stations located within each of the centers (Fig. 2.). We obtained information for the period from 1 January 2000 to 31 December 2009, although some stations had information for only a subset of these dates. The information collected was the average concentration of particulate matter with mass median aerodynamic diameter less than 10 microns ( $PM_{10}$ ) over 24 hour periods

### 2.2 Mortality and sociodemographic data

The daily number of non-accidental deaths (ICD-9 <800) in the study areas were obtained from the Instituto Nacional de Estadísticas, the official source of statistical data in Chile from 1 January 2000 to 31 December 2009 for all nine areas. The daily number of hospitalizations were obtained for five of the areas under study: Tomé, Talcahuano, Concepción, , Lota, and Coronel for the period of January 1 2006 to December 31 2007. Age, gender, and individual educational attainment data were obtained from the Departamento de Estadísticas e Información en Salud (DEIS).

### 2.3 Statistical methods

We used time series analyses and assumed both a Poisson distribution and that there was a linear association between ambient air pollution and mortality or morbidity on a logarithmic scale (Rupprecht et al., 1995).

Natural splines were created for air pollution concentrations on the day of study with one knot for each of 15, 30, 60, 90, 120, 180, and 365 days of observation. We then selected the model with the number of knots that either minimized the Akaike Information Criteria





Fig. 1. Map of Chile. The red area on the map has been declared a non-attainment zone because of failure to maintain daily  $PM_{10}$  concentrations below a standard threshold. The area includes nine communities with a population of 1 million inhabitants.

(AIC), a measure of model prediction, or maximized the evidence that the model residuals did not display any type of structure, including serial correlation, using Bartlett's test (Lindstrom & Bates, 1990; Priestly, 1981). We plotted model residuals against time and found neither a pattern nor a significant correlation between air pollution and time. Once we had selected the optimal model for time, we assessed the value of including terms for the twenty-four hour means of temperature, humidity, and barometric pressure. The best meteorological predictors of death were temperature and humidity while humidex (Meteorological Service of Canada, 2000), a composite measure of temperature and humidity, was the best meteorological predictor of morbidity. We considered temperature and humidex readings on the day of death and the day prior to death and accounted for non-linear associations with death by using natural spline functions. Indicator functions for the day-of-the-week were also included. The association between air pollutants and death was tested at lags of zero to seven days and results were presented for the lags which maximized the effect size. Results from each urban center were pooled using a random effects model.



Fig. 2. Detailed Map of Chile. The locations of ten metropolitan areas highlighted in circles

Here we present the increase in relative risk (RR) of mortality or morbidity with 95% confidence intervals for an increase in PM<sub>10</sub> concentration equal to the interquartile range of the pollutant's concentration over the period of study. The interquartile range includes the middle fifty-percent of the exposure data and provides a realistic estimate of the day-to-day changes in the pollutant's concentration. The interquartile range is a nonparametric measure of the data's spread and, as such, is not influenced by skewed data, extreme values or outliers which are unstable and infrequently seen. A random effects model was used to pool the estimates of relative risk following a DerSimonian- Laird test for homogeneity among estimates.

3. Results

Regional population sizes varied by over fourfold from 49,923 in Penco to 224,212 in Concepción (Table 2). The number of daily deaths varied by four to fivefold between

Concepcion and Penco. In the population of about one million people, there was an average of 15 deaths per day. The twenty-four hour mean concentrations of particulate matter varied by about 50% - 60% between regions with Chiguayante and San Pedro de la Paz reaching the greatest concentrations of PM<sub>10</sub> (Table 2).

Risk of mortality from cardiac disease appeared to be particularly sensitive to increases in air pollution with an estimated increase of 26% (7% to 49%). The point estimate for mortality relative risk was somewhat greater in the oldest compared to the youngest age group, however, the effect was not significantly greater for those at least eighty-five years old compared to less than sixty-five ( $p > 0.05$ ). The point estimates for mortality risk from PM<sub>10</sub> were similar for males and females ( $p > 0.05$ ) indicating a lack of effect modification by sex. The effect of PM<sub>10</sub> on mortality was greatest among those with the lower level of educational attainment. An interquartile increase in pollutants among those who did not complete a college or university degree was associated with a 16.8% (3% to 33%) increase in mortality whereas among college and university graduates there was 13% (-1% to 28%) increase, which was not statistically significant. The risk of death associated with air pollution was particularly high among the elderly with low educational attainment with an increase of 19% (3% to 35%).

	Population 100,000s	Total Mortality	Cardiac Mortality	Respiratory Mortality	PM <sub>10</sub> ( $\mu\text{g}/\text{m}^3$ )	Total Hospitaliza tion	Temperature
Tomé	5.47	0.935 (0.964)	0.081 (0.288)	0.285 (0.539)	47.613 (45.975)	12.441	12.592 (4.388)
Penco	4.99	0.681 (0.819)	0.067 (0.262)	0.175 (0.423)	56.118 (49.525)	NA	12.592 (4.388)
Talcahuano	17.13	3.010 (1.832)	0.264 (0.526)	0.791 (0.921)	50.030 (28.060)	61.760	13.355 (5.871)
Hualpén	8.8	1.272 (1.131)	0.113 (0.343)	0.346 (0.578)	34.645 (19.025)	NA	13.355 (5.871)
Concepción	22.42	3.487 (1.939)	0.321 (0.570)	1.006 (1.002)	41.734 (23.350)	101.414	13.355 (5.871)
San Pedro	8.92	0.975 (1.006)	0.070 (0.265)	0.269 (0.524)	56.118 (49.525)	NA	13.355 (5.871)
Chiguayante	9.98	1.042 (1.022)	0.091 (0.298)	0.302 (0.544)	56.118 (49.525)	NA	12.852 (3.101)
Lota	4.89	1.126 (1.077)	0.100 (0.320)	0.293 (0.544)	49.778 (31.325)	15.047	12.852 (3.101)
Coronel	10.31	1.344 (1.173)	0.129 (0.365)	0.358 (0.610)	52.148 (29.500)	22.280	12.852 (3.101)

Table 2. Population size, mean daily total mortality, 24-hour mean daily air pollution levels and 24-hour mean weather for nine urban centers in Chile from January 2000 to December 2009. Mean daily total mortality rates and 24 hour mean weather variables are accompanied by their standard deviation, while the interquartile range is reported for the concentration of PM<sub>10</sub>.

When regions were pooled, an interquartile increase in concentration of PM<sub>10</sub> was associated with a 5.5% (0.3% to 11%) increased risk of death from all causes (Table 3).

		Relative Risk
Mortality	All Causes	1.055 (1.003, 1.109)
	Cardiac	1.260 (1.065, 1.490)
	Respiratory	1.041 (1.024, 1.076)
Sex	Male	1.043 (1.020, 1.085)
	Female	1.061 (1.024, 1.099)
Age	< 64	1.053 (1.013, 1.096)
	65 - 74	1.048 (1.007, 1.089)
	85 +	1.061 (1.016, 1.107)
Education	< College	1.168 (1.029, 1.325)
	> College	1.130 (0.998, 1.280)
Ages > 85 & lowest educational strata		1.190 (1.031, 1.349)

Table 3. Increase in relative risk of mortality by age group, sex, educational attainment, associated with an interquartile increase in PM<sub>10</sub> adjusted for long-term trends, day-of-the-week, and temperature and humidity for nine urban centers in Chile from January 2000 to December 2009.

The risk of hospitalization from all causes and from respiratory disease showed no evidence of effect modification by age or sex with an increase in air pollution (Table 4). However, risk of hospitalization from cardiac disease was greatest among those 85 years old and greater, with an increase of 23% (6% to 44%) among the elderly versus 3% (-3% to 10%) among those less than 64 years of age; but, similar to risk of hospitalization from all cause and respiratory disease, cardiac disease showed no effect modification by sex.

		All Cause RR	Cardiac RR	Respiratory RR
Age	All	1.032 (1.011 to 1.053)	1.029 (0.983 to 1.077)	1.056 (1.005 to 1.111)
	< 64	1.037 (1.017 to 1.059)	1.033 (0.974 to 1.097)	1.067 (1.014 to 1.123)
	65 to 74	1.034 (0.996 to 1.074)	1.089 (1.006 to 1.178)	1.071 (0.945 to 1.214)
	75 to 84	1.036 (0.991 to 1.084)	1.050 (0.969 to 1.137)	1.119 (1.003 to 1.249)
	> 85	1.048 (0.977 to 1.124)	1.232 (1.058 to 1.435)	1.081 (0.946 to 1.235)
Sex	All	1.032 (1.011 to 1.053)	1.029 (0.983 to 1.077)	1.056 (1.005 to 1.111)
	Females	1.031(1.010 to 1.054)	0.998 (0.940 to 1.059)	1.046 (0.986 to 1.109)
	Males	1.034 (1.007 to 1.062)	1.055 (0.998 to 1.116)	1.073 (1.006 to 1.144)

Table 4. Relative Risk (RR) of hospitalization (morbidity) associated with an interquartile increase in concentrations of PM<sub>10</sub> adjusted for long-term trends, day-of-the-week, and humidex for the five urban centers in Concepcion from January 1 2006 to December 31 2007.

4. Discussion

Although progress has been made steadily over time at reducing ambient concentrations of particulate matter (PM<sub>10</sub>) (Fig.3), the results of this work suggests that there remains a risk to human health from exposure to this pollutant.The burden of mortality and morbidity due

to increases particulate matter (PM<sub>10</sub>) in the short-terms has the greatest influence on the health of those who are elderly with low educational attainment and those with cardiac disease. In general, effect modification was observed by age and by education but not by sex and effect modification was less pronounced for morbidity data than for mortality data. Air quality guidelines that seek to protect the entire population, including high risk subgroups, should consider the greater sensitivity of those who are elderly, have lower educational attainment or suffer from cardiac disease.

4.1 Effect modification by age and educational attainment

Age significantly modified the effect of cardiac morbidity for the five Chilean communities studied here. Modification by age was less pronounced for all cause mortality, all cause morbidity and respiratory morbidity. Similarly, we observed little modification by educational attainment for total mortality. However, we did find that the combination of old age and low educational attainment resulted in elevated risk from air borne particulate matter.




Fig. 3. Air pollution levels over time for all nine centres combined.

Previous work has reported modification of the effect of air pollutants by age (Bell et al., 2005; Pope, 2000; Pope et al., 2002; Spix et al., 1998; Zanobetti et al., 2000). For example, previous work indicated that compared to those under sixty-five years of age, Chileans eighty five years and older were observed to be more than twice as likely to die from acute increases in PM<sub>10</sub>, and over 50% more likely to die from increases in ozone and SO<sub>2</sub> (Camak



et al., 2009). Similarly, Bell et al. (2008) reported increased mortality effects in the elderly from ozone in The National Morbidity Mortality and Air Pollution Study of 98 U.S. cities (Bell & Dominici, 2008) and Filleul et al. (2004) reported a greater effect of air pollution mortality in those over sixty-five years old in France, though these effects were not statistically significant (Filleul et al., 2004).

Previous work has also reported effect modification by educational attainment and other indicators of social status (Bell et al., 2008; Forastiere et al., 2009; Dales, 2002; O'Neil et al., 2008; Ou et al., 2008; Prescott & Vestbo, 1999; Zanobetti et al., 2000). For example, in the Harvard Six-Cities and American Cancer Society cohort studies, there was an increased risk of mortality from long-term exposure to particulate matter among those with lower educational attainment (Health Effects Institute, 2000; Pope et al., 2002; Villeneuve et al., 2002). Similarly, in Hamilton, Canada, the non-accidental mortality risk estimates associated with sulphur dioxide and coefficient of haze were greater in areas of the city with lower educational attainment as well as greater employment in manufacturing (Jerrett et al. 2004). However, this finding is far from consistent: no relation to level of education was found in a study of mortality risk estimates from gaseous and particulate air pollution in Hong Kong (Ou et al., 2008); no effect modification by education was found among urban Americans from 98 communities for ozone levels (Bell & Dominici, 2008); and neither a time-series study of 20 U.S. (Samet et al., 2000) cities nor one focusing on Vancouver, Canada found social status modified the effect of air pollution on mortality (Villeneuve et al., 2003). Furthermore, a study of São Paulo, Brazil the authors reported that a monotonically increasing effects of air pollution with increasing education (Gouvenia & Fletcher, 2000). This type of conflicting results lead the authors of a systematic review of the Medline database up to May 2006 to state that because of inconsistent findings in both long-term and short-term exposure studies "Current evidence does not yet justify a definitive conclusion that socioeconomic characteristics modify the effects of air pollution on mortality" (Laurent et al., 2009). Nevertheless, here we report that in combination with old age, risk increases with lower educational attainment.

#### 4.2 The influence of social status

There are many possible reasons why one might expect lower socioeconomic position to increase susceptibility to the deleterious effects of air pollution including: increased exposure to the air pollutants of interest, increased exposure to co-pollutants from occupational dusts and fumes and cigarette smoke, fewer dietary fruits and vegetables, and reduced access to medical care and medicines (O'Neill et al., 2003; Sexton et al., 1993; Spix et al., 1998). Unfortunately, information on these variables was not available. However, because the overall effect size is based on the association between daily changes in air pollution and daily changes in mortality or morbidity, these other variables would only confound the overall pollution-illness association if they change day-to-day which is unlikely (Bell et al., 2005). It is possible that these variables differ between the educational groups and may partly account for the between-group estimates of effect found here.

#### 5. Conclusion

We found that the burden of all cause mortality and cardiac morbidity due to increased particulate air pollution is disproportionately experienced by the elderly who have low

educational attainment. These findings suggest that the determination of air quality guidelines designed to protect the general population may be insufficient to protect this vulnerable subgroup.

## 6. Acknowledgments

The authors would like to thank Dra. Danuta Rajs Grzebien of the Departamento de Estadísticas e Información de Salud, Ministerio de Salud for providing morbidity data, and Roberto Martínez González and Joyce Vera Bascour from the Secretaría Regional Ministerial del Medio Ambiente Region Metropolitana for providing data and comments.

## 7. References

- Dockery DW, Pope CA. Outdoor Particulates. *Environmental Epidemiology Oxford University Press*, New York. 2002;119-166.
- Brunekreef B, Holgate ST. Air pollution and health. *Lancet* 2002;360:1233-1242.
- Bascom R., Bromberg PA, Costa DA, Devlin R, Dockery DW, Frampton MW, Lambert W, Samet JM, Speizer FE, Utell M. Health effects of outdoor air pollution. *Am J Respir Crit Care Med* 1996;153:3-50.
- Hahn RA, Eaker ED, Barker ND, Teutsch SM, Sosniak WK, Krieger N. Poverty and death in the United States. *Int J Health Serv* 1996; 26:673-690
- Carr W, Zeitel L, Weiss K. Variations in asthma hospitalizations and deaths in New York City. *Am J Public Health* 1992; 82(59):65
- Chen Y, Dales R, Krewski D. Asthma and the risk of hospitalization in Canada : the role of socioeconomic and demographic factors. *Chest* 2001; 119:708-713.
- Cifuentes L, Vega J, Lave L. Daily mortality by cause and socio-economic status in Santiago, Chile 1988-1996. *Epidemiology* 1999; 10:S45
- Wojtyniak B, Wysocki M. Chronic airways disease in Poland. Recent trends. *Chest* 1989; 96(3):324S-328S
- Health Effects Institute. Reanalysis of the Harvard Six Cities Study and the American Cancer Society Study of Particulate Air Pollution and Mortality. 2000;1-285.
- Pope CA, III. Epidemiology of fine particulate air pollution and human health: biologic mechanisms and who's at risk? *Environ Health Perspect* 2000;108(4):713-723.
- Gouveia N, Fletcher T. Time Series analysis of air pollution and mortality: effects by cause, age and socioeconomic status. *J Epidemiol Community Health* 2000;54:750-755.
- Samet JM, Dominici F, Currier FC, Coursac I, Zeger SL. Fine Particulate Air Pollution and Mortality in 20 U.S. Cities 1987-1994. *N Eng J Med* 2000;343:1742-1749.
- Zanobetti A, Schwartz J, Gold D. Are there sensitive subgroups for the effects of airborne particles? *Environ Health Perspect* 2000;108(9):841-845.
- O'Neill, MS, Jerrett M, Kawachi I, Levy JI, Cohen AJ, Gouveia N, Wilkinson P, Fletcher T, Cifuentes L, Schwartz, J, Workshop on Air Pollution and Socioeconomic Conditions. Health, Wealth, and Air Pollution: Advancing Theory and Methods. *Environ Health Perspect* 2003;111:1861-1870.



- Sexton K, Gong Jr. H, Bailar JC III, Ford JG, Gold DR, Lambert WE, Utell MJ. Air pollution health risks: do class and race matter? *Toxicology and Industrial Health* 1993;9:843-878.
- Rupprecht G, Patashnick H, Beeson DE, Green RN, Meyer MD. A new automated monitor for the measurement of particulate carbon in the atmosphere. In: Particulate Matter: Health and Regulatory Issues. Proceedings of an International Specialty Conference, April 1995, Pittsburgh, PA, *Air and Waste Management Publications* Pittsburgh, PA, 1995;309-319
- Lindstrom ML, Bates DM. Nonlinear mixed effects models for repeated measures data. *Biometrics* 1990;46(3):673-687).
- Priestly MB. *Spectral Analysis of Time Series Academic Press, San Diego* 1981.
- Meteorological Service of Canada. 2002.  
[http://www.msccmc.ec.gc.ca/cd/brochures/humidity\\_e.cfm](http://www.msccmc.ec.gc.ca/cd/brochures/humidity_e.cfm)
- Pope CA, III, Burnett RT, Thun MJ, Calle EE, Krewski D, Ito K, Thurston GD, Lung Cancer, Cardiopulmonary Mortality, and Long-term Exposure to Fine Particulate Air Pollution. *JAMA* 2002;287:1132-1141.
- Spix C, Anderson HR, Schwartz J, Vigotti MA, Letertre A, Vonk JM, Touloumi F, Balducci T, Piekarski L, Bacharova A, Tobias A, Ponka, Katsouyanni K. Short-term effects of air pollution on hospital admissions of respiratory diseases in Europe: a quantitative summary of APHEA study results. *Arch Environ Health* 1998;53:54-64.
- Bell ML, Dominici F, Samet JM. A meta-analysis of time-series studies of ozone and mortality with comparison to the national morbidity, mortality, and air pollution study. *Epidemiology* 2005;16:425-426.
- Cakmak S, Dales RE, Blanco Vidal C. Air Pollution and Mortality in Chile: susceptibility among the elderly. *Environ Health Perspect* 2009;115:524-527.
- Bell ML, Dominici F. Effect modification by community characteristics on the short-term effects of ozone exposure and mortality in 98 US communities. *Am J Epidemiol* 2008;167:986-997.
- Filleul L, Le Tertre A, Baldi I, and Tessier JF. Difference in the relation between daily mortality and air pollution among elderly and all-ages populations in southwest France. *Environ Res* 2004;94:249-253.
- Ou CQ, Hedley AJ, Chung RY, Thach TQ, Chau YK, Chan KP, Yang L, Ho SY, Wong CM, Lam TH. Socioeconomic disparities in air pollution-associated mortality. *Environmental Research* 2008 Jan;107[2], 237-244.
- Forastiere F, Stafoggia M., Tasco C, Picciotto S, Agabati N, Cesaroni G, and Perucci CA Socioeconomic status, particulate air pollution, and daily mortality: differential exposure or differential susceptibility. 2009;50[3], 208-216.
- Prescott E, Vestbo J. Socioeconomic status and chronic obstructive pulmonary disease. *Thorax* 1999;54:737-741.
- Dales RE, Choi B, Chen Y, Tang M. Influence of family income on hospital visits for asthma among Canadian school children. *Thorax* 2002;57:513-517.
- O'Neill MS, Bell ML, Ranjit N, C LA, Loomis D, Gouveia N, Borja-Aburto VH, Air Pollution and Mortality in Latin America: The Role of Education. *Epidemiology*. 2008; 19(6): 810-819).

- Villeneuve PJ, Goldberg M.S., Krewski D, Burnett RT, Chen Y. Fine particulate air pollution and all-cause mortality within the Harvard Six-Cities Study: variations in risk by period of exposure. *Ann Epidemiol* 2002;12:568-576.
- Jerrett M. Burnett RT. Brook J. Kanaroglou P. Giovis C. Finkelstein N. Hutchison B. Do socioeconomic characteristics modify the short term association between air pollution and mortality? Evidence from a zonal time series in Hamilton, Canada. *J Epidemiol Community Health* 2004;58:31-40.
- Villeneuve PG, Burnett RT, Shid Y, Krewski D,, Goldberg MS, Hertzman C, Chen Y, Brook J. A time-series study of air pollution, socioeconomic status, and mortality in Vancouver, Canada. *Journal of Exposure Analysis and Environmental Epidemiology* 2003; 13, 427-435. doi:10.1038/sj.jea.7500292
- Laurent O, Bard D, Filleil L, Segala C. Effect of socioeconomic status on the relationship between atmospheric pollution and mortality. *Journal of Epidemiology & Community Health* 2009;61:665-675.
- Klemm, R.J.; Mason, R.M. Jr. Aerosol Research and Inhalation Epidemiological Study (ARIES): Air Quality and Daily Mortality Statistical Modeling - Interim Results . *J. Air Waste Manage. Assoc.* 2000, 50, 1433-1439.
- Daniels, M.J.; Dominici, F.; Samet, J.M.; Zeger, S.L. Estimating Particulate Matter-Mortality Dose-Response Curves and Threshold Levels: An Analysis of Daily Time-Series for the 20 Largest US Cities. *Am. J. Epidemiol.* 2000, 152, 397-406.
- Samet, J.M.; Dominici, F.; Currier, I.; Coursac, I.; Zeger, S.L. Fine Particulate Air Pollution and Mortality in 20 U.S. Cities, 1987-1994. *N. Engl. J. Med.* 2000, 343, 1742-1749.
- Schwartz, J. Assessing Confounding, Effect Modification, and Thresholds in the Association between Ambient Particles and Daily Deaths. *Environ. Health Perspect.* 2000, 108, 563-568.
- Schwartz, J. The Distributed Lag between Air Pollution and Daily Deaths. *Epidemiol.* 2000, 11, 320-326.
- Smith, R.L.; Davis, J.M.; Speckman, P. Assessing the human health risk of atmospheric particles *Novartis. Found. Symp.* 1999, 220, 59-72.
- Schwartz, J. Air Pollution and Daily Mortality in Birmingham, Alabama. *Am. J. Epidemiol.* 1993, 137, 1136-1147.
- Schwartz, J. Harvesting and Long Term Exposure Effects in the Relation between Air Pollution and Mortality. *Am. J. Epidemiol.* 2000, 151, 440-448.
- Gwynn, R.C.; Burnett, R.T.; Thurston, G.D. A Time-Series Analysis of Acidic Particulate Matter and Daily Mortality and Morbidity in the Buffalo, New York, Region. *Environ. Health Perspect.* 2000, 108, 125-133.
- Schwartz, J. Total Suspended Particulate Matter and Daily Mortality in Cincinnati, Ohio. *Environ. Health Perspect.* 1994, 102, 186-189.
- Ostro, B.D.; Hurley, S.; Lipsett, M.J. Air Pollution and Daily Mortality in the Coachella Valley, California: A Study of PM<sub>10</sub> Dominated by Coarse Particles. *Environ. Res.* 1999, 81, 231-238.

- Ostro, B.D.; Broadwin, R.; Lipsett, M.J. Coarse and fine particles and daily mortality in the Coachella Valley, California: a follow-up study. *J. Expo. Anal. Environ. Epidemiol.* 2000, 10, 412-419.
- Ito, K.; Kinney, P.L.; Thurston, G.D. Variations in PM-10 Concentrations within two Metropolitan Areas and their Implications for Health Effects Analyses. *Inhal. Toxicol.* 1995, 7, 735-745.
- Ito, K.; Thurston, G.D. Daily PM10/Mortality Associations: An Investigation of At-Risk Subpopulations. *J. Expo. Anal. Environ. Epidemiol.* 1996, 6, 79-95.
- Styer, P.; McMillan, N.; Gao, F.; Davis, J.; Sacks, J. Effect of Outdoor Airborne Particulate Matter on Daily Death Counts. *Environ. Health Perspect.* 1995, 103, 490-497.
- Moolgavkar, S.H. Pollution and Daily Mortality in Three U.S. Counties. *Environ. Health Perspect.* 2000 108, 777-784.
- Zanobetti, A.; Schwartz, J. Race, Gender, and Social Status as Modifiers of the Effects of PM10 on Mortality. *J. Occup. Environ. Med.* 2000, 42, 469-474.
- Schwartz, J. Particulate Air Pollution and Daily Mortality in Detroit. *Environ. Res.* 1991, 56, 204-213.
- Lippmann, M.; Ito, K.; Nadas, A.; Burnett, R.T. Association of Particulate Matter Components with Daily Mortality and Morbidity in Urban Populations. *Health Effects Institute Research Report Number 95*; Health Effects Institute: Cambridge, MA, 2000.
- Dockery, D.W.; Schwartz, J.; Spengler, J.D. Air Pollution and Daily Mortality: Associations with Particulates and Acid Aerosols. *Environ. Res.* 1992, 59, 362-373.
- Schwartz, J.; Dockery, D.; Neas, L.M. Is Daily Mortality Associated Specifically with Fine Particles? *J. Air Waste Manage. Assoc.* 1996, 46, 927-939.
- Kinney, P.L.; Özkaynak, H. Associations of Daily Mortality and Air Pollution in Los Angeles County. *Environ. Res.* 1991, 54, 99-120.
- Kinney, P.L.; Ito, K.; Thurston, G.D. A Sensitivity Analysis of Mortality/PM-10 Associations in Los Angeles. *Inhal. Toxicol.* 1995, 7, 59-69.
- Goldberg, M.S.; Bailer, J.C. III; Burnett, R.T.; Brook, J.R.; Tamblyn, R.; Bonvalot, Y.; Ernst, P.; Flegel, K.M.; Singh, R.K.; Valois, M.F. *Identifying Subgroups of the General Population That May Be Susceptible to Short-Term Increases in Particulate Air Pollution: A Time-Series Study in Montreal, Quebec.* Health Effects Institute Research Report Number 97; Health Effects Institute, Cambridge, MA, 2000.
- Schwartz, J.; Dockery, D.W. Increased Mortality in Philadelphia Associated with Daily Air Pollution Concentrations. *Am. Rev. Respir. Dis.* 1992, 145, 600-604.
- Moolgavkar, S.H.; Luebeck, E.G.; Hall, T.A.; Anderson, E.L. Air Pollution and Daily Mortality in Philadelphia. *Epidemiol.* 1995, 6, 476-484.
- Schwartz, J. Daily deaths are associated with combustion particles rather than SO<sub>2</sub> in Philadelphia. *Occup. Environ. Med.* 2000, 57, 692-697.
- Samet, J.; Zeger, S.; Kelsall, J.; Xu, J.; Kalkstein, L. Does Weather Confound or Modify the Association of Particulate Air Pollution with Mortality? An analysis of the Philadelphia Data, 1973-1980. *Environ. Res.* 1998, 77, 9-19.
- Kelsall, J.E.; Samet, J.M.; Zeger, S.L.; Xu, J. Air Pollution and Mortality in Philadelphia, 1974-1988. *Am. J. Epidemiol.* 1997, 46, 750-762.

- Wyzga, R.E.; Lipfert, F.W. Temperature - Pollution Interactions with Daily Mortality in Philadelphia. In *Particulate Matter: Health and Regulatory Issues, VIP-49, Proceedings of an International Specialty Conference, Pittsburgh, Pennsylvania, April 4-6, 1995*, 3-42.
- Lipfert, F.W.; Morris, S.C.; Wyzga, R.E. Daily Mortality in the Philadelphia Metropolitan Area and Size-Classified Particulate Matter. *J. Air. Waste Manage. Assoc.* 2000, 50, 1501-1513.
- Mar, T.F.; Norris, G.A.; Koenig, J.Q.; Larson, T.V. Associations between Air Pollution and Mortality in Phoenix, 1995-1997. *Environ. Health Perspect.* 2000, 108, 347-353.
- Chock, D.P.; Winkler, S.L.; Chen, C. A Study of the Association between Daily Mortality and Ambient Air Pollutant Concentrations in Pittsburgh, Pennsylvania. *J. Air Waste Manage. Assoc.* 2000, 50, 1481-1500.
- Pope, C.A. III; Hill, R.W.; Villegas, G.M. Particulate Air Pollution and Daily Mortality on Utah's Wasatch Front. *Environ. Health Perspect.* 1999, 107, 567-573.
- Boucher, K.M.; Lyon, J.L.; Lillquist, D.R. Daily Mortality and Exposure to PM<sub>10</sub> Air Pollution: A Comparison of Salt Lake and Utah Counties, 1985-1993. In *Proceedings of the Second Colloquium on Particulate Air Pollution and Human Health*, Park City, Utah, May 1-3, 1996; pp 4-36-4-54.
- Ostro, B. Fine Particulate Air Pollution and Mortality in Two Southern California Counties. *Environ. Res.* 1995, 70, 98-104.
- Fairley, D. The Relationship of Daily Mortality to Suspended Particulates in Santa Clara County, 1980-1986. *Environ. Health Perspect.* 1990, 89, 159-168.
- Fairley, D. Daily Mortality and Air Pollution in Santa Clara County, California: 1989-1996. *Environ Health Perspect.* 1999, 107, 637-641
- Moolgavkar, S.H.; Luebeck, E.G.; Hall, T.A.; Anderson, E.L. Particulate air pollution, sulfur dioxide and daily mortality: A reanalysis of the Steubenville data. *Inhal. Toxicol.* 1995, 7, 35-44.
- Schwartz, J.; Dockery, D.W. Particulate Air Pollution and Daily Mortality in Steubenville, Ohio. *Am. J. Epidemiol.* 1992, 135, 12-19.
- Burnett, R.T.; Brook, J.R.; Cakmak, S.; Raizenne, M.; Stieb, D.; Vincent, R.; Krewski, D.; Brook, J.R.; Philips, O.; Özkaynak, H. The Association Between Ambient Carbon Monoxide Levels and Daily Mortality in Toronto, Canada. *J. Air Waste Manage. Assoc.* 1998, 48, 689-700.
- Burnett, R.T. In *National Ambient Air Quality Objectives for Ground-Level Ozone, Science Assessment Document*; Canadian Environmental Protection Act Federal - Provincial Advisory Committee Working Group on Air Quality Objectives and Guidelines; Health Canada and Environment Canada: Ottawa, 1999; pp A1-A8.
- Burnett, R.T.; Cakmak, S.; Brook, J.R. The Effect of the Urban Ambient Air Pollution Mix on Daily Mortality Rates in 11 Canadian Cities. *Can. J. Public Health.* 1998, 89, 152-156
- Burnett, R.T.; Brook, J.; Dann, T.; Delocla, C.; Philips, O.; Cakmak, S.; Vincent, R.; Goldberg, M.S.; Krewski, D. Association Between Particulate- and Gas-Phase Components Of Urban Air Pollution and Daily Mortality in Eight Canadian Cities. *Inhal. Toxicol.* 2000, 12 (Supplement 4), 15-39.
- Pope, C.A. III; Schwartz, J.; Ransom, M.R. Daily Mortality and PM<sub>10</sub> Pollution in Utah Valley. *Arch. Environ. Health.* 1992, 47, 211-217.



- Lyon, J.L.; Mori, M.; Gao, R. Is there a causal association between excess mortality and exposure to PM10 air pollution? Additional analyses by location, year, season and cause of death. *Inhal. Toxicol.* 1995, 7, 603-614.
- Pope, C.A. III; Kalkstein, L.S. Synoptic Weather Modeling and Estimates of the Exposure-Response Relationship between Daily Mortality and Particulate Air Pollution. *Environ. Health Perspect.* 1996, 104, 414-420.
- Borja-Aburto, V.H.; Loomis, D.P.; Bangdiwala, S.I.; Shy, C.M.; Rascon-Pacheco, R.A. Ozone, Suspended Particulates, and Daily Mortality in Mexico City. *Am. J. Epidemiol.* 1997, 258-268.
- Borja-Aburto, V.H.; Castillejos, M.; Gold, D.R.; Bierzwinski, S.; Loomis, D. Mortality and Ambient Fine Particles in Southwest Mexico City, 1993-1995. *Environ. Health Perspect.* 1998, 106, 849-855.
- Castillejos, M.; Borja-Aburto, V.H.; Dockery, D.W.; Gold, D.R.; Loomis, D. Airborne coarse particles and mortality. *Inhal. Toxicol.* 2000, 12(S1), 61-72.
- Ostro, B.D.; Sanchez, J.M.; Aranda, C.; Eskeland, G.S. Air Pollution and Mortality: Results from a Study of Santiago, Chile. *J. Expo. Anal. Environ. Epidemiol.* 1996, 6, 97-114.
- Saldiva, P.H.; Pope, C.A. III; Schwartz, J.; Dockery, D.W.; Lichtenfels, A.J.; Salge, J.M.; Barone, I.; Bohm, G.M. Air pollution and mortality in elderly people: A time-series study in Sao Paulo, Brazil. *Arch. Environ. Health.* 1996, 50, 159-163.
- Touloumi, G.; Pocock, S.J.; Katsouyanni, K.; Trichopolous, D. Short-term effects of air pollution on daily mortality in Athens: A time-series analysis. *Int. J. Epidemiol.* 1994, 23, 957-967.
- Hatzakis, A.; Katsouyanni, K.; Kalandidi, A.; Day, N.; Trichopoulos, D. Short-term effects of air pollution on mortality in Athens. *Int. J. Epidemiol.* 1986, 15:73-81.
- Touloumi, G.; Samoli, E.; Katsouyanni, K. Daily mortality and "winter type" air pollution in Athens, Greece - a time series analysis within the APHEA project. *J. Epidemiol. Community Health.* 1996, 50(suppl 1), S47-S51.
- Touloumi, G.; Katsouyanni, K.; Zmirou, D.; et al. Short-term Effects of Ambient Oxidant Exposure on Mortality: A Combined Analysis within the APHEA Project. *Am. J. Epidemiol.* 1997, 146, 177-185.
- Zmirou, D.; Schwartz, J.; Saez, M.; et al. Time-Series Analysis of Air Pollution and Cause Specific Mortality. *Epidemiol.* 1998, 9, 495-503.
- Sunyer, J.; Castellsagué, J.; Sáez, M.; Tobias, A.; Antó, J.M. Air pollution and mortality in Barcelona. *J. Epidemiol. Community Health.* 1996, 50(Suppl 1), S76-S80.
- Tobias, A.; Campbell, M.J. Modelling influenza epidemics in the relation between black smoke and total mortality. A sensitivity analysis. *J. Epidemiol. Community Health.* 1999, 53, 583-584.
- Wietlisbach, V.; Pope, C.A. III; Ackermann-Liebrich, U. Air pollution and daily mortality in three Swiss urban areas. *Soz. Präventivmed.* 1996, 41, 107-115.
- Wordley, J.; Walters, S.; Ayres, J.G. Short term variations in hospital admissions and mortality and particulate air pollution. *Occup. Environ. Med.* 1997, 54, 108-116.
- Bachárová, L.; Fandáková, K.; Bratinka, J.; Budinská, M.; Bachár, J.; Gudába, M. The association between air pollution and the daily number of deaths: findings from the

- Slovak Republic contribution to the APHEA project. *J. Epidemiol. Community Health.* 1996, 50(Suppl 1), S19-S21.
- Krzyzanowski, M.; Wojtyniak, B. Air pollution and daily mortality in Cracow. *Public. Health Rev.* 1991/92, 19, 73-81.
- Wojtyniak, B.; Piekarski, T. Short term effect of air pollution on mortality in Polish urban populations - what is different? *J. Epidemiol. Community Health.* 1996, 50(Suppl 1), S36-S41.
- Szafraniec, K.; Tęcza, W.; Jedrychowski, W. Long-term and Short-term Changes in Air Pollutants and Mortality: Experience among Inhabitants in Krakow, Poland. In *Health Effects of Particulate Matter in Ambient Air: Proceedings of an International Conference*, Prague, Czech Republic, April 23-25, 1997; 196-202.
- Kotesovec, F.; Skorkovsky, J.; Brynda, J.; Peters, A.; Heinrich J. Daily Mortality and Air Pollution in Northern Bohemia: Different Effects for Men and Women. *Cent. Eur. J. Public Health* 2000, 8, 120-127.
- Peters, A.; Skorkovsky, J.; Kotesovec, F.; Brynda, J.; Spix, C.; Wichmann, H.E.; Heinrich, J. Associations between Mortality and Air Pollution in Central Europe. *Environ. Health Perspect.* 2000, 108, 283-287.
- Rahlenbeck, S.I.; Kahl, H. Air pollution and mortality in East Berlin during the winters of 1981-1989. *Int. J. Epidemiol.* 1996, 25, 1220-1226.
- Spix, C.; Heinrich, J.; Dockery, D.; Schwartz, J.; Völksch, G.; Schwinkowski, K.; Cöllen, C.; Wichmann, H.E. Air pollution and daily mortality in Erfurt, East Germany, 1980-89. *Environ. Health Perspect.* 1993, 101, 518-526.
- Wichmann, H.-E.; Spix, C.; Tuch, T.; Wolke, G.; Peters, A.; Heinrich, J.; Kreyling, W.G.; Heyder, J. *Daily Mortality and Fine and Ultrafine Particles in Erfurt, Germany. Part I: Role of Particle Number and Particle Mass, Health Effects Institute Research Report Number 98*; Health Effects Institute: Cambridge, MA, 2000.
- Spix, C.; Wichmann, H.E. Daily mortality and air pollutants: findings from Köln, Germany. *J. Epidemiol. Community Health.* 1996, 50(suppl 1), 52-58.
- Schwartz, J.; Marcus, A. Mortality and Air Pollution in London: A Time Series Analysis. *Am. J. Epidemiol.* 1990 131, 185-194.
- Ito, K.; Thurston, G.D.; Hayes, C.; Lippmann, M. Associations of London, England, Daily Mortality with Particulate Matter, Sulfur Dioxide, and Acidic Aerosol Pollution. *Arch. Environ. Health.* 1993, 48, 213-220.
- Anderson, H.R.; Ponce de Leon, A.; Bland, J.M.; Bower, J.S.; Strachan, D.P. Air pollution and daily mortality in London: 1987-92. *B.M.J.* 1996, 312, 665-669.
- Zmirou, D.; Barumandzadeh, T.; Balducci, F.; Ritter, P.; Laham, G.; Ghilardi, J.-P. Short term effects of air pollution on mortality in the city of Lyon, France, 1985-90. *J. Epidemiol. Community Health.* 1996, 50(suppl 1), S30-S35.
- Alberdi JC, Díaz J, Montero JC, Mirón I. Daily mortality in Madrid community 1986-1992: Relationship with meteorological variables. *Eur J Epidemiol.* 1998;14:571-578.
- Alberdi Odriozola, J.C.; Díaz Jiménez, J.; Montero Rubio, J.C.; Mirón Pérez, I.J.; Pajares Ortiz, M.S.; Ribera Rodrigues P. Air pollution and mortality in Madrid, Spain: a time-series analysis. *Int. Arch. Occup. Environ. Health.* 1998, 71, 543-549.

- Díaz, J.; García, R.; Ribera, P.; Alberdi, J.C.; Hernández, E.; Pajares, M.S.; Otero, A. Modeling of air pollution and its relationship with mortality and morbidity in Madrid, Spain. *Int. Arch. Occup. Environ. Health.* 1999, 72, 366-376.
- Morris, W.T. *Regression analysis of daily mortality and air pollution using real and simulated data*; Department of Trade and Industry: London, 1996.
- Vigotti, M.A.; Rossi, G.; Bisanti, L.; Zanobetti, A.; Schwartz, J. Short term effects of urban air pollution on respiratory health in Milan, Italy, 1980-89. *J. Epidemiol. Community Health.* 1996, 50(suppl 1), S71-S75.
- Rossi, G.; Vigotti, M.A.; Zanobetti, A.; Repetto, F.; Gianelle, V.; Schwartz, J. Air Pollution and Cause-Specific Mortality in Milan, Italy, 1980-1989. *Arch. Environ. Health.* 1999, 54, 158-164.
- Mackenbach, J.P.; Looman, C.W.N.; Kunst, A.E. Air pollution, lagged effects of temperature, and mortality: The Netherlands 1979-87. *J. Epidemiol. Community Health.* 1993, 47, 121-126.
- Buringh, E.; Fischer, P.; Hoek, G. Is SO<sub>2</sub> a causative factor for the PM-associated mortality risks in the Netherlands? *Inhal. Toxicol.* 2000, 12(S1), 55-60.
- Hoek, G.; Brunekreef, B.; Verhoeff, A.; van Wijnen, J.; Fischer, P. Daily Mortality and Air Pollution in the Netherlands. *J. Air Waste Manage. Assoc.* 2000, 50, 1380-1389.
- Le Tertre, A.; Quénel, P.; Medina, S.; Le Moullec, Y.; Festy, B.; Ferry, R.; Dab, W. Modélisation des liens court terme entre la pollution atmosphérique et la santé. Un exemple: SO<sub>2</sub> et mortalité totale, Paris, 1987-1990. *Rev. Épidémiol. Santé Publique.* 1998, 46, 316-328.
- Dab, W.; Medina, S.; Quénel, P.; et al. Short term respiratory health effects of ambient air pollution: results of the APHEA project in Paris. *J. Epidemiol. Community Health.* 1996, 50(suppl 1), S42-S46.
- Michelozzi, P.; Forastiere, F.; Fusco, D.; Perucci, C.A.; Ostro, B.; Ancona, C.; Pallotti, G. Air pollution and daily mortality in Rome, Italy. *Occup. Environ. Med.* 1998, 55, 605-610.
- Hoek, G.; Groot, B.; Schwartz, J.D.; Eilers, P. Effects of ambient particulate matter and ozone on daily mortality in Rotterdam, the Netherlands. *Arch. Environ. Health.* 1997, 52, 455-463.
- Ballester, F.; Corella, D.; Pérez-Hoyos, S.; Hervás, A. Air pollution and mortality in Valencia, Spain: a study using the APHEA methodology. *J. Epidemiol. Community Health.* 1996, 50, 527-533.
- Ostro, B.; Chestnut, L.; Vichit-Vadakan, N.; Laixuthai, A. The Impact of Particulate Matter on Daily Mortality in Bangkok, Thailand. *J. Air Waste Manage. Assoc.* 1996, 49, PM100-PM107.
- Xu, X.; Dockery, D.W.; Gao, J.; Chen, Y. Air Pollution and Daily Mortality in Residential Areas of Beijing, China. *Arch. Environ. Health.* 1994, 49, 216-222.
- Hong, Y.-C.; Leem, J.-H.; Ha, E.-H.; Christiani, D.C. PM<sub>10</sub> Exposure, Gaseous Pollutants, and Daily Mortality in Incheon, South Korea. *Environ. Health. Perspect.* 1999, 107, 873-878.
- Lee, J.-T.; Shin, D.; Chung, Y. Air Pollution and Daily Mortality in Seoul and Ulsan, Korea. *Environ. Health Perspect.* 1999, 107, 149-154.



Lee, J.T.; Kim, H.; Hong, Y.C.; Kwan, H.J.; Schwartz, J.; Christiani, D.C. Air Pollution and Daily Mortality in Seven Major Cities of Korea, 1991-1997. *Environ. Res.* 2000, 84, 247-254.

Xu, Z.Y.; Yu, D.G.; Jing, L.B.; Xu, X.P. Air Pollution and Daily Mortality in Shenyang, China. *Arch. Environ. Health* 2000, 55, 115-120.

IntechOpen

IntechOpen



## **Advanced Topics in Environmental Health and Air Pollution Case Studies**

Edited by Prof. Anca Moldoveanu

ISBN 978-953-307-525-9

Hard cover, 470 pages

**Publisher** InTech

**Published online** 29, August, 2011

**Published in print edition** August, 2011

The book describes the effects of air pollutants, from the indoor and outdoor spaces, on the human physiology. Air pollutants can influence inflammation biomarkers, can influence the pathogenesis of chronic cough, can influence reactive oxygen species (ROS) and can induce autonomic nervous system interactions that modulate cardiac oxidative stress and cardiac electrophysiological changes, can participate in the onset and exacerbation of upper respiratory and cardio-vascular diseases, can lead to the exacerbation of asthma and allergic diseases. The book also presents how the urban environment can influence and modify the impact of various pollutants on human health.

### **How to reference**

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Sabit Cakmak, Sara L. Martin, Claudia Blanco Vidal, Timur Gultekin, Vladislav Brion and Maria Angelica Rubio (2011). Do Socio-Demographic Characteristics Modify the Association Between Air Pollution and Mortality & Morbidity?, Advanced Topics in Environmental Health and Air Pollution Case Studies, Prof. Anca Moldoveanu (Ed.), ISBN: 978-953-307-525-9, InTech, Available from: <http://www.intechopen.com/books/advanced-topics-in-environmental-health-and-air-pollution-case-studies/do-socio-demographic-characteristics-modify-the-association-between-air-pollution-and-mortality-morb>

**INTECH**  
open science | open minds

### **InTech Europe**

University Campus STeP Ri  
Slavka Krautzeka 83/A  
51000 Rijeka, Croatia  
Phone: +385 (51) 770 447  
Fax: +385 (51) 686 166  
[www.intechopen.com](http://www.intechopen.com)

### **InTech China**

Unit 405, Office Block, Hotel Equatorial Shanghai  
No.65, Yan An Road (West), Shanghai, 200040, China  
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元  
Phone: +86-21-62489820  
Fax: +86-21-62489821

© 2011 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike-3.0 License](https://creativecommons.org/licenses/by-nc-sa/3.0/), which permits use, distribution and reproduction for non-commercial purposes, provided the original is properly cited and derivative works building on this content are distributed under the same license.

IntechOpen

IntechOpen