

# We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

186,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index  
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?  
Contact [book.department@intechopen.com](mailto:book.department@intechopen.com)

Numbers displayed above are based on latest data collected.  
For more information visit [www.intechopen.com](http://www.intechopen.com)



# Angiogenesis and Vascularity for Tissue Engineering Applications

Remo A. Largo<sup>1</sup>, Venkat M. Ramakrishnan<sup>1</sup>, Martin Ehrbar<sup>2</sup>, Algirdas Ziogas<sup>2</sup>, Jan A. Plock<sup>3</sup> and Daniel Eberli<sup>1</sup>

<sup>1</sup>*Division of Urology, University Hospital Zurich*

<sup>2</sup>*Division of Obstetrics, University Hospital Zurich*

<sup>3</sup>*Division of Plastic and Hand Surgery, University Hospital Zurich  
Switzerland*

## 1. Introduction

Tissue engineering is a field of medicine that has experienced significant growth in prominence over the past three decades. Though traditional interventions exist for many medical maladies, tissue engineering aims to combat such disorders through the synthesis of body tissues and organs, resulting in functional implants. Tissue engineering takes an innovative approach, often utilizing autologous stem cells for tissue construction or materials that are biocompatible while avoiding immune rejection (Nomi *et al.*, 2002).

Despite its immense successes, a major hurdle still faces tissue engineering. Large volumes of implanted tissue are unable to stimulate the formation of necessary blood vessels required for their survival. In the body, naturally occurring, equivalent vascular networks serve vital functions in gas and nutrient exchange, metabolic processes, and waste expulsion. Though individual, large vessels have been successfully engineered for implant, it is still exceptionally difficult to fashion a stable and sustainable network of vessels for large volumes of tissue (Nomi *et al.*, 2002). Neovascularization after tissue damage requires a level of positive and negative control that has not been successfully replicated in a laboratory environment to date. As such, rapid *de novo* synthesis of a controlled, established vascular network remains a challenge today.

Angiogenesis is the morphogenic process of forming new blood vessels from pre-existing ones (Laschke *et al.*, 2006; Dai and Rabie, 2007; Li and Rabie, 2007). This event plays an important, normal physiological role in wound healing, tissue repair, pregnancy, and exercise (Ferrara and Davis-Smyth, 1997), and exists in contrast to vasculogenesis (the formation of the very first blood vessels in the body, and especially predominant in embryological development). Yet, the abuse of angiogenesis, leading to an uncontrolled vascular formation as a consequence of epigenetic influence, nucleotide polymorphisms, or endocrine irregularities can also result in tumor formation (Verbridge *et al.*, 2010). However, angiogenesis is clearly an activity that is central to development and tissue maintenance. Successful modulation of angiogenesis can have profound therapeutic outcomes for organs and tissues deprived of an adequate, stable vasculature. Studies from the last two decades have shown that the manipulation of various factors directly influences angiogenic outcome.

Angiogenesis requires the activity of soluble factors such as Vascular Endothelial Growth Factor (VEGF; outlined in the next section and in Tables 2 & 3), basic Fibroblast Growth Factor (bFGF), Platelet-derived Growth Factor (PDGF), Transforming Growth Factor- $\beta$  (TGF- $\beta$ ) (Nomi *et al.*, 2002; Dai and Rabie, 2007; Li and Rabie, 2007; Kanczler and Oreffo, 2008; Bates, 2010), Keratinocyte Growth Factor (KGF) (Elia *et al.*, 2010), Hepatocyte Growth Factor (HGF) (Hoot *et al.*, 2010), Ephrin-B2 (Herbert *et al.*, 2009), and Angiopoietin (Han *et al.*, 2010) (Table 1). Morphologically, angiogenesis can be attributed to endothelial cell migration and proliferation as well as pericyte recruitment, migration and differentiation (Egginton, 2010).

Molecule	Known Properties	Citations
Basic Fibroblast Growth Factor (bFGF)	Stimulates activity of fibroblasts, neurons, smooth muscle cells, and endothelial cells; acts via tyrosine kinase receptors; induces production of VEGF during angiogenic stimulation.	(Pepper <i>et al.</i> , 1992; Lee <i>et al.</i> , 2003; Arkudas <i>et al.</i> , 2007; Jung <i>et al.</i> , 2010; Wu <i>et al.</i> , 2010)
Platelet-derived Growth Factor (PDGF)	Certain isoforms possess survival and mitogenic functions; implicated in tumor angiogenesis; compared to VEGF, PDGF: (a) has comparable angiogenic activity, (b) produces blood vessels with decreased permeability and leakage, and (c) is thought to produce functionally different blood vessels.	(Li <i>et al.</i> , 2010; Wu <i>et al.</i> , 2010)
Transforming Growth Factor $\beta$ (TGF- $\beta$ )	Promotes and inhibits angiogenesis and tumor invasion via stimulation of Hepatocyte Growth Factor (HGF) expression.	(Hoot <i>et al.</i> , 2010)
Keratinocyte Growth Factor (KGF)	Member of FGF family; certain isoforms have been implicated in wound healing as well as the inhibition of neovascularization.	(Wang <i>et al.</i> , 2010)
Hepatocyte Growth Factor (HGF)	Stimulates endothelial cell growth, migration, scatter, and elongation independently of VEGF.	(Hoot <i>et al.</i> , 2010)
Ephrin-B2	Implicated in arterial/ venous differentiation (see section: Modulation of Notch Signaling).	(Herbert <i>et al.</i> , 2009)
Angiopoietin	Signals through Tie receptors; certain isoforms are angiogenic inhibitors, while others promote inhibition of apoptosis; Stabilize blood vessels and reduce leakage.	(Han <i>et al.</i> , 2010)

Table 1. Notable Ancillary Growth Factors and Their Properties

2. VEGF and its receptors: a brief overview

In studying the development of cells, tissues, and organs, Vascular Endothelial Growth Factor (VEGF) has been identified as a key, though not sole, proponent of angiogenesis (Takahashi and Shibuya, 2005; Arkudas *et al.*, 2007; Dai and Rabie, 2007; Yla-Herttuala, 2009;

Bates, 2010; Elia *et al.*, 2010). VEGF exists in six classes (Table 2), lettered A through F, though the vast amount of literature concerning VEGF-A indicates that it is by far the most widely understood of all. VEGF-A exists in at least eight homodimeric isoforms, all conceived by alternative mRNA splicing(Ferrara, 2009) and differing by their amino acid number. Of these, VEGF<sub>121</sub> and VEGF<sub>165</sub> are predominant and show great promise in tissue engineering applications.

VEGF Class	Known Properties	Citations
A	Induces proliferation of arterial, venous, and lymphatic vascular endothelial cells; stimulates monocyte chemotaxis; hematopoietic effects; eight isoforms in humans.	(Ferrara, 2009)
B	May play role in atrial conduction, but not required for cardiovascular development.	(Olofsson <i>et al.</i> , 1996a; Olofsson <i>et al.</i> , 1996b; Aase <i>et al.</i> , 2001)
C	Involved in embryonic angiogenesis, lymphangiogenesis, & lymphatic vessel maintenance; mitogenic for cultured endothelial cells.	(Kukk <i>et al.</i> , 1996; Orlandini <i>et al.</i> , 1996; Jeltsch <i>et al.</i> , 1997; Yamada <i>et al.</i> , 1997; Dumont <i>et al.</i> , 1998)
D	Thought to be involved in pulmonary development, endothelial cell mitogen; Function(s) still generally unclear.	(Orlandini <i>et al.</i> , 1996; Yamada <i>et al.</i> , 1997; Achen <i>et al.</i> , 1998; Farnebo <i>et al.</i> , 1999)
E	Proteins are encoded by Orf-viruses and predominantly expressed in sheep, goats, and rarely in humans.	(Lyttle <i>et al.</i> , 1994; Meyer <i>et al.</i> , 1999)
F	Derived from snake venom; enhances the formation of vascular fenestrations in guinea pigs.	(Klein and Catargi, 2007; Matsunaga <i>et al.</i> , 2009; Yamazaki <i>et al.</i> , 2009)
Placental Growth Factor (PlGF)	Induces vascular permeability; supplements VEGF activity during wound-healing; may enhance VEGF-driven angiogenesis.	(Carmeliet <i>et al.</i> , 2001; Adini <i>et al.</i> , 2002; Hattori <i>et al.</i> , 2002; Luttun <i>et al.</i> , 2002; Odorisio <i>et al.</i> , 2002)

Table 2. Six General VEGF Classes

The binding of VEGF to any of its many receptors, such as VEGF-R1 (Flt-1) (Shibuya *et al.*, 1990; de Vries *et al.*, 1992), VEGF-R2 (KDR) (Terman *et al.*, 1991), VEGF-R3 (Flt-4) (Fitz *et al.*, 1997), and Neuropilin (NRP-1 and -2) (Soker *et al.*, 1996; Soker *et al.*, 1998; Dallas *et al.*, 2008), has been shown to trigger a signaling cascade that results in the activation of angiogenesis (Stefanini *et al.*, 2009). Not surprisingly, VEGF receptor and ligand placement and density have been implicated in the successful transmission of the signaling cascades required for angiogenesis (Stefanini *et al.*, 2009). Modulating receptors and their densities appear to be a logical avenue for further efforts in modulating neovascularization. Further, Heparin Sulfate Proteoglycan (HSPG) has been implicated in the binding of VEGF to their receptors, but its exact function remains unclear (Ferrara *et al.*, 2003; Lee *et al.*, 2010). Preliminary findings suggest that HSPGs inhibit VEGF binding to its receptors, and consequently, angiogenic activity (Lee *et al.*, 2010).

### 3. Modulations to VEGF and its delivery

Though VEGF has a significant influence on cell migration, proliferation, and vasodilation, the uncontrolled or sole use of VEGF *in vivo* has been shown to result in the disordered growth of blood vessels into a dense mass (hemangioma), malignant tumor angiogenesis, and the assembly of leaky vessels (Takahashi and Shibuya, 2005; Bates, 2010). As such, the fundamental function of VEGF in physiological development and maintenance, contrasted by its role in tumorigenesis and vessel instability, is paradoxical. In instances where VEGF is heterogeneously present in a microenvironment, areas of high VEGF expression resulted in abnormal angiogenesis. However, implanting VEGF-transfected myoblasts, with each cell equally producing VEGF over time, led to the formation of a stable, normal vascular network (Misteli *et al.*, 2010). The latter provides a compelling case for carefully-controlled VEGF release, distribution, and kinetics in tissue engineering applications. As discussed below, several current tissue-engineering efforts are positioned to resolve this dilemma.

It should not be assumed that blind administration of VEGF is solely responsible for a successful angiogenic effort. Many studies have illustrated that the promotion of successful angiogenesis depends on a prolonged exposure to a low dose of VEGF (Wernike *et al.*, 2010), while other studies claim that micro-environmental conditions must be taken into account (Ferrara and Davis-Smyth, 1997; Misteli *et al.*, 2010). With regards to the latter, techniques like FACS purification (Misteli *et al.*, 2010) and microdialysis (Hoier *et al.*, 2010; Marcus *et al.*, 2010) are becoming increasingly prevalent in closely monitoring VEGF expression in the microenvironment. Further, many attempts at eliciting controlled angiogenesis also focus on coupling the properties of VEGF with certain other growth factors – most notably bFGF (Arkudas *et al.*, 2007) or KGF (Elia *et al.*, 2010) (see Table 1); stable vessels were formed when VEGF was combined with either of these two factors.

There exist contradictions regarding the parameters of tissue exposure to VEGF. In one comparison, a study highlighting growth factor implementation in orthopedic applications suggested that VEGF delivery for over 14 days may have interfered in the vascularization during bone healing and restoration (Wernike *et al.*, 2010). This suggestion challenges an argument that a longer VEGF exposure (of approximately one month) was necessary for the production of stable, but leaky, vessels as brief exposure to VEGF (less than 15 days) resulted in the formation of unstable vessels. Further, these vessels actually degenerated after VEGF delivery cessation (Tafuro *et al.*, 2009). Disparities such as this are common in the



literature and demonstrate the sheer complexity of VEGF activity in various tissue environments.

The view that successful VEGF delivery and angiogenic response are strictly dose-dependent is challenged by factors such as delivery kinetics, which appear to be critical for proper vasculature (Borselli *et al.*, 2010; Wernike *et al.*, 2010). For instance, a characteristic problem with the bolus delivery of VEGF is the outcome of variable systemic effects (Matoka and Cheng, 2009) such as haemorrhage, hypotension, or flu-like symptoms (Benjamin *et al.*, 1999). This method of delivery also fails to achieve the prolonged supply of physiological low dose of VEGF necessary to produce a mature, lasting vascular network. The lack of lasting vessel formation observed in many therapeutic trials is probably due to difficulties in the delivery of VEGF, a growth factor with an apparently narrow therapeutic window (Hariawala *et al.*, 1996; Lee *et al.*, 2000; Dor *et al.*, 2003; Ozawa *et al.*, 2004), in the optimal time and dose for maintaining sufficient vascularity.

A need exists for biomaterials or stable scaffolds that enable slow, sustained VEGF release (Rocha *et al.*, 2008) for a predictable and functional outcome. Studies in different laboratories have indicated that incorporation of growth factor into slow-release polymer formulations could present a means for better control of dose, location, and duration of active signals in tissue (Edelman *et al.*, 1991; Lee *et al.*, 2000; Sheridan *et al.*, 2000; Ehrbar *et al.*, 2004). The kinetics of VEGF delivery today depends on the physical properties (such as cross-linking and porosity, for instance) of biocompatible conduits such as fibrin-gels (Ehrbar *et al.*, 2004; Arkudas *et al.*, 2007), gelatin microparticles (Patel *et al.*, 2008), collagen / fibronectin hydrogels (Glotzbach *et al.*, 2010), and PLG(A) scaffolds (Murphy *et al.*, 2000) (Rocha *et al.*, 2008; Matoka and Cheng, 2009; Borselli *et al.*, 2010; Golub *et al.*, 2010). Moreover, transfecting or transducing developing cells (muscle precursor cells or myoblasts, for instance) with VEGF and subsequently injecting them into a site can also provide a steady, longer-term delivery of the growth factor (Misteli *et al.*, 2010).

Once VEGF is coupled with a delivery conduit, a more intricate approach to controlling VEGF delivery involves modifying the factor itself. Extensive delivery-mechanics research was conducted with the isoform VEGF<sub>121</sub>. Though VEGF<sub>121</sub> is initially confined within a biomaterial (such as a fibrin-gel matrix) upon implant, plasmin and metalloproteinase degradation of the implant over time allows for the rapid, free diffusion of VEGF<sub>121</sub> into the whole body environment (Ehrbar *et al.*, 2004), potentially resulting in angiogenesis that may not be localized. Additionally, the short biological half-life of VEGF impedes its use in long-term applications. A synthetic variant of VEGF<sub>121</sub>, known as TG-VEGF<sub>121</sub>, cross-links to fibrinogen by the transglutaminating activity of factor XIII during fibrin-gel polymerization. This covalently tethered TG-VEGF<sub>121</sub> is protected from rapid diffusion. Gradual degradation of the fibrin-gel by local fibrinolytic activities results in a local liberation of low levels of TG-VEGF<sub>121</sub> into tissue. Experimental animal models have shown that fibrin-conjugated TG-VEGF<sub>121</sub> produced more structurally stable vessels than VEGF<sub>121</sub> while avoiding vascular leakage (Ehrbar *et al.*, 2004; Ehrbar *et al.*, 2008).

#### 4. Hypoxia-mediated control and modulation of notch signaling

Hypoxia presents another means of employing direct control on VEGF and ancillary angiogenic factors. In the low-oxygen environments of normal muscle, VEGF mRNA experiences decreased degradation and increased expression (Ikeda *et al.*, 1995; Levy *et al.*, 1995; Levy *et al.*, 1996; Tang *et al.*, 2004), though severely hypoxic settings will actually

impede VEGF up-regulation (Milkiewicz *et al.*, 2004). VEGF mRNA stability is made possible by its interaction with HuR, a complex that binds and stabilizes RNAs, consequently regulating gene expression (Levy *et al.*, 1998). The increase in VEGF production can be attributed to the Internal Ribosome Entry Site (IRES), which is accountable for efficient factor synthesis under hypoxic conditions (Stein *et al.*, 1998). Other proteins can also impact VEGF efficiency and activity. Control of expressed VEGF lies with ORP150 (oxygen regulated protein), a chaperone that transports VEGF from the cell's endoplasmic reticulum to the Golgi apparatus (Kuwabara *et al.*, 1996; Ozawa *et al.*, 2001). An increase in ORP150 levels correlates with an increased production of VEGF during hypoxia (Ozawa *et al.*, 2001). Further, Hypoxia-inducing factor (HIF) is responsible for activating the transcription of genes associated with neovascularization (Covello and Simon, 2004; Ramirez-Bergeron *et al.*, 2006). These proteins represent points of control for VEGF modulation.

Because angiogenesis is the synthesis of vasculature from existing vessels, constant remodeling and modification of vessels takes place *in vivo*. Existing endothelial cells, such as stalk and tip cells, have receptors that respond to environmental conditions like hypoxia, in which Notch signaling (driven by VEGF-A presence) is most prevalent. The presence of VEGF-A leads to an increase in the presence of Delta-like Ligand 4 (DLL4), a major Notch ligand. It is believed that Notch signaling modulates the ratio of VEGF to its receptors through the inhibition of VEGF-R2.

Manipulation of the Notch signaling pathway presents another means of controlling angiogenesis. Cao, et al., discussed the role of Notch signaling in modulating VEGF activity, resulting in effective pruning and branching of vascular vessels (Cao *et al.*, 2009; Cao *et al.*, 2010). Activation of the Notch signaling pathway inhibits VEGF signaling by down-regulating the VEGF receptor synthesis. With fewer available receptors for free VEGF to bind, endothelial cell proliferation is effectually curbed. This process occurs at the cellular level of the lining endothelium, a major component of vascular vessels, and vascular growth in any particular direction is controlled.

## 5. Applications in regenerative medicine: notable case studies

Recent advances in understanding the angiogenic process and isolating potent and specific angiogenic growth factors prompted their therapeutic usage. Evidence that VEGF is a specific endothelial cell growth factor suggested its potential in therapeutic angiogenesis. Injection of the VEGF<sub>165</sub> protein enhanced revascularization of rabbit ischemic hindlimbs (Takeshita *et al.*, 1994). VEGF treatment induced collateral vessel formation, endothelium-dependent blood flow and tissue perfusion. VEGF has been tested for potential beneficial effects on wound repair in diabetic animal models (Greenhalgh *et al.*, 1990; Tsuboi and Rifkin, 1990; Frank *et al.*, 1995). The growth factor treatment regimens accelerated granulation, tissue formation, and wound closure. However, due to the high clearance and/or degradation of the proteins from the administration site, topically administered growth factors would require high dosages and frequent delivery.

An alternative strategy for therapeutic angiogenesis is gene therapy using recombinant angiogenic growth factors. The first study using VEGF<sub>165</sub> cDNA was performed by gene transfer into the iliac artery of an ischemic hindlimb of a rabbit (Bauters *et al.*, 1994; Takeshita *et al.*, 1996). VEGF protein was expressed at the site of injection, augmenting the formation of collateral vessels. Subsequently, intramuscular gene transfer of VEGF cDNA

was used in a similar model with similar results (Asahara *et al.*, 1996). This technique was further employed in patients with peripheral vascular disease and critical limb ischemia (Isner *et al.*, 1996). Clinical trials showed significant improvement in collateral blood flow, healed ischemic ulcers, and most importantly, salvage of limbs in patients in whom amputation was imminent (Isner, 1998; Isner and Takayuki, 1998). VEGF protein and cDNA have been used for coronary revascularization, resulting in improved myocardial perfusion and increased collateral density (Isner and Losordo, 1999).

As previously discussed, a successful angiogenic outcome is not solely attributed to VEGF implementation alone (discussed in Section 3 and Table 1). Wilcke, *et al.*, showcased the coupling of VEGF with bFGF in a fibrin dermal substitute (Wilcke *et al.*, 2007). This led to a marked improvement in factor delivery, with a notable, prolonged release and resulted in a higher density of newly developed vessels in *in vivo* murine models. Concurrently, an *in vivo* experiment performed by Zacchigna, *et al.*, studied the effect of a VEGF and Angiopoietin 1 (Ang 1) combination (delivered via an adeno-associated viral vector, for long-term protein production and release) on the muscle blood flow (MBF) and vascular permeability of rat skeletal muscle (Zacchigna *et al.*, 2007). The grouping elicited a marked increase in both resting MBF and perfusion post exercise stimulation. To contrast, VEGF expression alone did not enhance the resting MBF and actually reduced tissue perfusion after exercise.

Elcin, *et al.*, researched *in vitro* release kinetics and *in vivo* angiogenic effects of human VEGF-loaded PLGA sponges in rats (Elcin and Elcin, 2006). When compared with control sponges (containing no factor) and bolus injections of VEGF, the use of VEGF-loaded PLGA sponges led to the establishment of neovascularized sites suitable for tissue engineering purposes. Patel, *et al.*, who utilized VEGF-loaded gelatin microparticles infused in biodegradable composite scaffolds, outlined another notable attempt in VEGF delivery modulation in the field of orthopedics (Patel *et al.*, 2008). Findings suggested that modulating the degree of gelatin cross-linking could affect VEGF release into the microenvironment and, consequently, angiogenic outcome. Ennett, *et al.*, studied the temporally regulated delivery of VEGF *in vivo*, hypothesizing that the means of VEGF-loading into a delivery scaffold would have an impact on the factor release kinetics (Ennett *et al.*, 2006). They compared (a) VEGF loaded directly into a PLG scaffold with (b) VEGF pre-encapsulated in PLG microspheres that were later used to fabricate a PLG scaffold. Though pre-encapsulated VEGF microspheres further delayed the factor's release into the surrounding environment, this approach produced a desirable angiogenic outcome, with significant local angiogenesis and negligible systemic effects.

Pre-encapsulation of VEGF by nanoparticles or microspheres has also been investigated in conjunction with Matrigel hydrogels, PLGA-, and collagen-scaffolds. While it is clear that encapsulation protects the factor(s) within the nanoparticles and microspheres, it also offers a mechanism of controlled release – especially ideal for a potent angiogenic factor like VEGF. Even though pre-encapsulated VEGF-loaded PLGA scaffolds performed better than similarly loaded Matrigel hydrogels (with respect to release profiles in a saline solution; also in comparison with free VEGF) *in vitro*, both of the loaded delivery conduits improved angiogenesis *in vivo*. An increase in both endothelial cell counts and red blood cells was noted at the sites of implantation (des Rieux *et al.*, 2010). Further, VEGF-loaded PLGA microspheres combined with collagen elicited strong enhancements to vascular sprouting and activation of endothelial cells *in vivo* and *in vitro*, respectively (des Rieux *et al.*, 2010).



The idea of optimizing delivery agents combined with the notion that co-factor usage is ideal for neovascularization resulted in an experiment that analyzed the co-implementation of VEGF with FGF-2 via an acellular collagen scaffold implant (Nillesen *et al.*, 2007). An implant containing both factors resulted in the highest vessel density and the most mature blood vessels (characteristics of an enhanced, stable vasculature) in a rat model when compared with scaffolds containing either of the factors alone or no factor at all.

## 6. Conclusion

A significant challenge facing tissue engineering lies in eliciting controlled neovascularization. Controlled neovascularization can have a profound impact in vessel and organ synthesis, as well as in the treatment of damaged tissues. Many current attempts for such control entail the use of VEGF. Attempts at fabricating an ordered vascular network include changes to VEGF through structural modification, the employment of strategies controlling its release, and coupling VEGF to other growth factors.

Yet, despite the promises shown by VEGF, significant obstacles remain. Many researchers propose that use of growth factors alone will not ensure stable angiogenesis. Rather, a combination of growth factors (VEGF and bFGF, for example), delivery methods, and modulation of inflammatory responses (via fibroblast, macrophage, cell-adhesion molecule and cytokine manipulation, for instance) and pathways (such as hypoxia and Notch) are thought to facilitate adequate vessel fabrication and stability.

Additionally, investigation into the mechanisms of cellular crosstalk is necessary to better understand angiogenesis in general. Moreover, the refinement and implementation of microenvironment monitoring technologies are vital in ensuring proper vascular development. Such technologies would enable researchers to closely study the impact of gross- or modulated-release of growth factors on the delicate balance required for vascular formation and branching.

## 7. Acknowledgements

Remo A. Largo and Venkat M. Ramakrishnan contributed equally to the production of this chapter.

## 8. References

- Aase, K., von Euler, G., Li, X., Ponten, A., Thoren, P., Cao, R., Cao, Y., Olofsson, B., Gebre-Medhin, S., Pekny, M., Alitalo, K., Betsholtz, C., and Eriksson, U. (2001). Vascular endothelial growth factor-B-deficient mice display an atrial conduction defect. *Circulation* 104, 358-364.
- Achen, M.G., Jeltsch, M., Kukk, E., Makinen, T., Vitali, A., Wilks, A.F., Alitalo, K., and Stacker, S.A. (1998). Vascular endothelial growth factor D (VEGF-D) is a ligand for the tyrosine kinases VEGF receptor 2 (Flk1) and VEGF receptor 3 (Flt4). *Proc Natl Acad Sci U S A* 95, 548-553.
- Adini, A., Kornaga, T., Firoozbakht, F., and Benjamin, L.E. (2002). Placental growth factor is a survival factor for tumor endothelial cells and macrophages. *Cancer Res* 62, 2749-2752.

- Arkudas, A., Tjiawi, J., Bleiziffer, O., Grabinger, L., Polykandriotis, E., Beier, J.P., Sturzl, M., Horch, R.E., and Kneser, U. (2007). Fibrin gel-immobilized VEGF and bFGF efficiently stimulate angiogenesis in the AV loop model. *Mol Med* 13, 480-487.
- Asahara, T., Tsurumi, Y., Takeshita, S., and Isner, J.M. (1996). Naked cDNA encoding secreted proteins for intra-arterial and intramuscular gene transfer. *Seminars in interventional cardiology : SIIC* 1, 225-232.
- Bates, D.O. (2010). Vascular endothelial growth factors and vascular permeability. *Cardiovasc Res* 87, 262-271.
- Bauters, C., Asahara, T., Zheng, L.P., Takeshita, S., Bunting, S., Ferrara, N., Symes, J.F., and Isner, J.M. (1994). Physiological assessment of augmented vascularity induced by VEGF in ischemic rabbit hindlimb. *The American journal of physiology* 267, H1263-1271.
- Benjamin, L.E., Golijanin, D., Itin, A., Pode, D., and Keshet, E. (1999). Selective ablation of immature blood vessels in established human tumors follows vascular endothelial growth factor withdrawal. *J Clin Invest* 103, 159-165.
- Borselli, C., Ungaro, F., Oliviero, O., d'Angelo, I., Quaglia, F., La Rotonda, M.I., and Netti, P.A. (2010). Bioactivation of collagen matrices through sustained VEGF release from PLGA microspheres. *J Biomed Mater Res A* 92, 94-102.
- Cao, L., Arany, P.R., Kim, J., Rivera-Feliciano, J., Wang, Y.S., He, Z., Rask-Madsen, C., King, G.L., and Mooney, D.J. (2010). Modulating Notch signaling to enhance neovascularization and reperfusion in diabetic mice. *Biomaterials* 31, 9048-9056.
- Cao, L., Arany, P.R., Wang, Y.S., and Mooney, D.J. (2009). Promoting angiogenesis via manipulation of VEGF responsiveness with notch signaling. *Biomaterials* 30, 4085-4093.
- Carmeliet, P., Moons, L., Luttun, A., Vincenti, V., Compernelle, V., De Mol, M., Wu, Y., Bono, F., Devy, L., Beck, H., Scholz, D., Acker, T., DiPalma, T., Dewerchin, M., Noel, A., Stalmans, I., Barra, A., Blacher, S., Vandendriessche, T., Ponten, A., Eriksson, U., Plate, K.H., Foidart, J.M., Schaper, W., Charnock-Jones, D.S., Hicklin, D.J., Herbert, J.M., Collen, D., and Persico, M.G. (2001). Synergism between vascular endothelial growth factor and placental growth factor contributes to angiogenesis and plasma extravasation in pathological conditions. *Nat Med* 7, 575-583.
- Covello, K.L., and Simon, M.C. (2004). HIFs, hypoxia, and vascular development. *Current topics in developmental biology* 62, 37-54.
- Dai, J., and Rabie, A.B. (2007). VEGF: an essential mediator of both angiogenesis and endochondral ossification. *Journal of dental research* 86, 937-950.
- Dallas, N.A., Gray, M.J., Xia, L., Fan, F., van Buren, G., 2nd, Gaur, P., Samuel, S., Lim, S.J., Arumugam, T., Ramachandran, V., Wang, H., and Ellis, L.M. (2008). Neuropilin-2-mediated tumor growth and angiogenesis in pancreatic adenocarcinoma. *Clin Cancer Res* 14, 8052-8060.
- de Vries, C., Escobedo, J.A., Ueno, H., Houck, K., Ferrara, N., and Williams, L.T. (1992). The fms-like tyrosine kinase, a receptor for vascular endothelial growth factor. *Science* 255, 989-991.
- des Rieux, A., Ucakar, B., Mupendwa, B.P., Colau, D., Feron, O., Carmeliet, P., and Preat, V. (2010). 3D systems delivering VEGF to promote angiogenesis for tissue engineering. *J Control Release*.

- Dor, Y., Djonov, V., and Keshet, E. (2003). Induction of vascular networks in adult organs: implications to proangiogenic therapy. *Annals of the New York Academy of Sciences* 995, 208-216.
- Dumont, D.J., Jussila, L., Taipale, J., Lymboussaki, A., Mustonen, T., Pajusola, K., Breitman, M., and Alitalo, K. (1998). Cardiovascular failure in mouse embryos deficient in VEGF receptor-3. *Science* 282, 946-949.
- Edelman, E.R., Mathiowitz, E., Langer, R., and Klagsbrun, M. (1991). Controlled and modulated release of basic fibroblast growth factor. *Biomaterials* 12, 619-626.
- Egginton, S. (2010). Physiological factors influencing capillary growth. *Acta physiologica*.
- Ehrbar, M., Djonov, V.G., Schnell, C., Tschanz, S.A., Martiny-Baron, G., Schenk, U., Wood, J., Burri, P.H., Hubbell, J.A., and Zisch, A.H. (2004). Cell-demand liberation of VEGF121 from fibrin implants induces local and controlled blood vessel growth. *Circulation research* 94, 1124-1132.
- Ehrbar, M., Zeisberger, S.M., Raeber, G.P., Hubbell, J.A., Schnell, C., and Zisch, A.H. (2008). The role of actively released fibrin-conjugated VEGF for VEGF receptor 2 gene activation and the enhancement of angiogenesis. *Biomaterials* 29, 1720-1729.
- Elcin, A.E., and Elcin, Y.M. (2006). Localized angiogenesis induced by human vascular endothelial growth factor-activated PLGA sponge. *Tissue engineering* 12, 959-968.
- Elia, R., Fuegy, P.W., VanDelden, A., Firpo, M.A., Prestwich, G.D., and Peattie, R.A. (2010). Stimulation of in vivo angiogenesis by in situ crosslinked, dual growth factor-loaded, glycosaminoglycan hydrogels. *Biomaterials* 31, 4630-4638.
- Ennett, A.B., Kaigler, D., and Mooney, D.J. (2006). Temporally regulated delivery of VEGF in vitro and in vivo. *J Biomed Mater Res A* 79, 176-184.
- Farnebo, F., Piehl, F., and Lagercrantz, J. (1999). Restricted expression pattern of vegf-d in the adult and fetal mouse: high expression in the embryonic lung. *Biochem Biophys Res Commun* 257, 891-894.
- Ferrara, N. (2009). VEGF-A: a critical regulator of blood vessel growth. *Eur Cytokine Netw* 20, 158-163.
- Ferrara, N., and Davis-Smyth, T. (1997). The biology of vascular endothelial growth factor. *Endocr Rev* 18, 4-25.
- Ferrara, N., Gerber, H.P., and LeCouter, J. (2003). The biology of VEGF and its receptors. *Nat Med* 9, 669-676.
- Fitz, L.J., Morris, J.C., Towler, P., Long, A., Burgess, P., Greco, R., Wang, J., Gassaway, R., Nickbarg, E., Kovacic, S., Ciarletta, A., Giannotti, J., Finnerty, H., Zollner, R., Beier, D.R., Leak, L.V., Turner, K.J., and Wood, C.R. (1997). Characterization of murine Flt4 ligand/VEGF-C. *Oncogene* 15, 613-618.
- Frank, S., Hubner, G., Breier, G., Longaker, M.T., Greenhalgh, D.G., and Werner, S. (1995). Regulation of vascular endothelial growth factor expression in cultured keratinocytes. Implications for normal and impaired wound healing. *J Biol Chem* 270, 12607-12613.
- Glotzbach, J.P., Levi, B., Wong, V.W., Longaker, M.T., and Gurtner, G.C. (2010). The basic science of vascular biology: implications for the practicing surgeon. *Plast Reconstr Surg* 126, 1528-1538.

- Golub, J.S., Kim, Y.T., Duvall, C.L., Bellamkonda, R.V., Gupta, D., Lin, A.S., Weiss, D., Robert Taylor, W., and Guldberg, R.E. (2010). Sustained VEGF delivery via PLGA nanoparticles promotes vascular growth. *Am J Physiol Heart Circ Physiol* 298, H1959-1965.
- Greenhalgh, D.G., Sprugel, K.H., Murray, M.J., and Ross, R. (1990). PDGF and FGF stimulate wound healing in the genetically diabetic mouse. *Am J Pathol* 136, 1235-1246.
- Han, S., Arnold, S.A., Sithu, S.D., Mahoney, E.T., Gerald, J.T., Tran, P., Benton, R.L., Maddie, M.A., D'Souza, S.E., Whittemore, S.R., and Hagg, T. (2010). Rescuing vasculature with intravenous angiopoietin-1 and alpha v beta 3 integrin peptide is protective after spinal cord injury. *Brain : a journal of neurology* 133, 1026-1042.
- Hariawala, M.D., Horowitz, J.R., Esakof, D., Sheriff, D.D., Walter, D.H., Keyt, B., Isner, J.M., and Symes, J.F. (1996). VEGF improves myocardial blood flow but produces EDRF-mediated hypotension in porcine hearts. *J Surg Res* 63, 77-82.
- Hattori, K., Heissig, B., Wu, Y., Dias, S., Tejada, R., Ferris, B., Hicklin, D.J., Zhu, Z., Bohlen, P., Witte, L., Hendrikx, J., Hackett, N.R., Crystal, R.G., Moore, M.A., Werb, Z., Lyden, D., and Rafii, S. (2002). Placental growth factor reconstitutes hematopoiesis by recruiting VEGFR1(+) stem cells from bone-marrow microenvironment. *Nat Med* 8, 841-849.
- Herbert, S.P., Huisken, J., Kim, T.N., Feldman, M.E., Houseman, B.T., Wang, R.A., Shokat, K.M., and Stainier, D.Y. (2009). Arterial-venous segregation by selective cell sprouting: an alternative mode of blood vessel formation. *Science* 326, 294-298.
- Hoier, B., Rufener, N., Bojsen-Moller, J., Bangsbo, J., and Hellsten, Y. (2010). The effect of passive movement training on angiogenic factors and capillary growth in human skeletal muscle. *The Journal of physiology* 588, 3833-3845.
- Hoot, K.E., Oka, M., Han, G., Bottinger, E., Zhang, Q., and Wang, X.J. (2010). HGF upregulation contributes to angiogenesis in mice with keratinocyte-specific Smad2 deletion. *J Clin Invest* 120, 3606-3616.
- Ikeda, E., Achen, M.G., Breier, G., and Risau, W. (1995). Hypoxia-induced transcriptional activation and increased mRNA stability of vascular endothelial growth factor in C6 glioma cells. *J Biol Chem* 270, 19761-19766.
- Isner, J.M. (1998). Vascular endothelial growth factor: gene therapy and therapeutic angiogenesis. *The American journal of cardiology* 82, 63S-64S.
- Isner, J.M., and Losordo, D.W. (1999). Therapeutic angiogenesis for heart failure. *Nat Med* 5, 491-492.
- Isner, J.M., and Takayuki, A. (1998). Therapeutic angiogenesis. *Frontiers in bioscience : a journal and virtual library* 3, e49-69.
- Isner, J.M., Walsh, K., Symes, J., Pieczek, A., Takeshita, S., Lowry, J., Rosenfield, K., Weir, L., Brogi, E., and Jurayj, D. (1996). Arterial gene transfer for therapeutic angiogenesis in patients with peripheral artery disease. *Human gene therapy* 7, 959-988.
- Jeltsch, M., Kaipainen, A., Joukov, V., Meng, X., Lakso, M., Rauvala, H., Swartz, M., Fukumura, D., Jain, R.K., and Alitalo, K. (1997). Hyperplasia of lymphatic vessels in VEGF-C transgenic mice. *Science* 276, 1423-1425.
- Jung, S., Wermker, K., Poetschik, H., Ziebur, T., and Kleinheinz, J. (2010). The impact of hyperbaric oxygen therapy on serological values of vascular endothelial growth



- factor (VEGF) and basic fibroblast growth factor (bFGF). *Head & face medicine* 6, 29.
- Kanczler, J.M., and Oreffo, R.O. (2008). Osteogenesis and angiogenesis: the potential for engineering bone. *Eur Cell Mater* 15, 100-114.
- Klein, M., and Catargi, B. (2007). VEGF in physiological process and thyroid disease. *Annales d'endocrinologie* 68, 438-448.
- Kukk, E., Lymboussaki, A., Taira, S., Kaipainen, A., Jeltsch, M., Joukov, V., and Alitalo, K. (1996). VEGF-C receptor binding and pattern of expression with VEGFR-3 suggests a role in lymphatic vascular development. *Development* 122, 3829-3837.
- Kuwabara, K., Matsumoto, M., Ikeda, J., Hori, O., Ogawa, S., Maeda, Y., Kitagawa, K., Imuta, N., Kinoshita, T., Stern, D.M., Yanagi, H., and Kamada, T. (1996). Purification and characterization of a novel stress protein, the 150-kDa oxygen-regulated protein (ORP150), from cultured rat astrocytes and its expression in ischemic mouse brain. *J Biol Chem* 271, 5025-5032.
- Laschke, M.W., Harder, Y., Amon, M., Martin, I., Farhadi, J., Ring, A., Torio-Padron, N., Schramm, R., Rucker, M., Junker, D., Haufel, J.M., Carvalho, C., Heberer, M., Germann, G., Vollmar, B., and Menger, M.D. (2006). Angiogenesis in tissue engineering: breathing life into constructed tissue substitutes. *Tissue engineering* 12, 2093-2104.
- Lee, K.Y., Peters, M.C., Anderson, K.W., and Mooney, D.J. (2000). Controlled growth factor release from synthetic extracellular matrices. *Nature* 408, 998-1000.
- Lee, K.Y., Peters, M.C., and Mooney, D.J. (2003). Comparison of vascular endothelial growth factor and basic fibroblast growth factor on angiogenesis in SCID mice. *J Control Release* 87, 49-56.
- Lee, T.Y., Folkman, J., and Javaherian, K. (2010). HSPG-binding peptide corresponding to the exon 6a-encoded domain of VEGF inhibits tumor growth by blocking angiogenesis in murine model. *PLoS One* 5, e9945.
- Levy, A.P., Levy, N.S., and Goldberg, M.A. (1996). Post-transcriptional regulation of vascular endothelial growth factor by hypoxia. *J Biol Chem* 271, 2746-2753.
- Levy, A.P., Levy, N.S., Wegner, S., and Goldberg, M.A. (1995). Transcriptional regulation of the rat vascular endothelial growth factor gene by hypoxia. *J Biol Chem* 270, 13333-13340.
- Levy, N.S., Chung, S., Furneaux, H., and Levy, A.P. (1998). Hypoxic stabilization of vascular endothelial growth factor mRNA by the RNA-binding protein HuR. *J Biol Chem* 273, 6417-6423.
- Li, Q.F., and Rabie, A.B. (2007). A new approach to control condylar growth by regulating angiogenesis. *Archives of oral biology* 52, 1009-1017.
- Li, X., Kumar, A., Zhang, F., Lee, C., Li, Y., Tang, Z., and Arjuna, P. (2010). VEGF-independent angiogenic pathways induced by PDGF-C. *Oncotarget* 1, 309-314.
- Luttun, A., Tjwa, M., Moons, L., Wu, Y., Angelillo-Scherrer, A., Liao, F., Nagy, J.A., Hooper, A., Priller, J., De Klerck, B., Compennolle, V., Daci, E., Bohlen, P., Dewerchin, M., Herbert, J.M., Fava, R., Matthys, P., Carmeliet, G., Collen, D., Dvorak, H.F., Hicklin, D.J., and Carmeliet, P. (2002). Revascularization of ischemic tissues by PlGF treatment, and inhibition of tumor angiogenesis, arthritis and atherosclerosis by anti-Flt1. *Nat Med* 8, 831-840.



- Lyttle, D.J., Fraser, K.M., Fleming, S.B., Mercer, A.A., and Robinson, A.J. (1994). Homologs of vascular endothelial growth factor are encoded by the poxvirus orf virus. *Journal of virology* 68, 84-92.
- Marcus, H.J., Carpenter, K.L., Price, S.J., and Hutchinson, P.J. (2010). In vivo assessment of high-grade glioma biochemistry using microdialysis: a study of energy-related molecules, growth factors and cytokines. *J Neurooncol* 97, 11-23.
- Matoka, D.J., and Cheng, E.Y. (2009). Tissue engineering in urology. *Canadian Urological Association journal = Journal de l'Association des urologues du Canada* 3, 403-408.
- Matsunaga, Y., Yamazaki, Y., Suzuki, H., and Morita, T. (2009). VEGF-A and VEGF-F evoke distinct changes in vascular ultrastructure. *Biochem Biophys Res Commun* 379, 872-875.
- Meyer, M., Clauss, M., Lepple-Wienhues, A., Waltenberger, J., Augustin, H.G., Ziche, M., Lanz, C., Buttner, M., Rziha, H.J., and Dehio, C. (1999). A novel vascular endothelial growth factor encoded by Orf virus, VEGF-E, mediates angiogenesis via signalling through VEGFR-2 (KDR) but not VEGFR-1 (Flt-1) receptor tyrosine kinases. *Embo J* 18, 363-374.
- Milkiewicz, M., Pugh, C.W., and Egginton, S. (2004). Inhibition of endogenous HIF inactivation induces angiogenesis in ischaemic skeletal muscles of mice. *The Journal of physiology* 560, 21-26.
- Misteli, H., Wolff, T., Fuglistaler, P., Gianni-Barrera, R., Gurke, L., Heberer, M., and Banfi, A. (2010). High-throughput flow cytometry purification of transduced progenitors expressing defined levels of vascular endothelial growth factor induces controlled angiogenesis in vivo. *Stem Cells* 28, 611-619.
- Murphy, W.L., Peters, M.C., Kohn, D.H., and Mooney, D.J. (2000). Sustained release of vascular endothelial growth factor from mineralized poly(lactide-co-glycolide) scaffolds for tissue engineering. *Biomaterials* 21, 2521-2527.
- Nillesen, S.T., Geutjes, P.J., Wismans, R., Schalkwijk, J., Daamen, W.F., and van Kuppevelt, T.H. (2007). Increased angiogenesis and blood vessel maturation in acellular collagen-heparin scaffolds containing both FGF2 and VEGF. *Biomaterials* 28, 1123-1131.
- Nomi, M., Atala, A., Coppi, P.D., and Soker, S. (2002). Principles of neovascularization for tissue engineering. *Molecular aspects of medicine* 23, 463-483.
- Odorisio, T., Schietroma, C., Zaccaria, M.L., Cianfarani, F., Tiveron, C., Tatangelo, L., Failla, C.M., and Zambruno, G. (2002). Mice overexpressing placenta growth factor exhibit increased vascularization and vessel permeability. *J Cell Sci* 115, 2559-2567.
- Olofsson, B., Pajusola, K., Kaipainen, A., von Euler, G., Joukov, V., Saksela, O., Orpana, A., Pettersson, R.F., Alitalo, K., and Eriksson, U. (1996a). Vascular endothelial growth factor B, a novel growth factor for endothelial cells. *Proc Natl Acad Sci U S A* 93, 2576-2581.
- Olofsson, B., Pajusola, K., von Euler, G., Chilov, D., Alitalo, K., and Eriksson, U. (1996b). Genomic organization of the mouse and human genes for vascular endothelial growth factor B (VEGF-B) and characterization of a second splice isoform. *J Biol Chem* 271, 19310-19317.

- Orlandini, M., Marconcini, L., Ferruzzi, R., and Oliviero, S. (1996). Identification of a c-fos-induced gene that is related to the platelet-derived growth factor/vascular endothelial growth factor family. *Proc Natl Acad Sci U S A* 93, 11675-11680.
- Ozawa, C.R., Banfi, A., Glazer, N.L., Thurston, G., Springer, M.L., Kraft, P.E., McDonald, D.M., and Blau, H.M. (2004). Microenvironmental VEGF concentration, not total dose, determines a threshold between normal and aberrant angiogenesis. *J Clin Invest* 113, 516-527.
- Ozawa, K., Kondo, T., Hori, O., Kitao, Y., Stern, D.M., Eisenmenger, W., Ogawa, S., and Ohshima, T. (2001). Expression of the oxygen-regulated protein ORP150 accelerates wound healing by modulating intracellular VEGF transport. *J Clin Invest* 108, 41-50.
- Patel, Z.S., Ueda, H., Yamamoto, M., Tabata, Y., and Mikos, A.G. (2008). In vitro and in vivo release of vascular endothelial growth factor from gelatin microparticles and biodegradable composite scaffolds. *Pharm Res* 25, 2370-2378.
- Pepper, M.S., Ferrara, N., Orci, L., and Montesano, R. (1992). Potent synergism between vascular endothelial growth factor and basic fibroblast growth factor in the induction of angiogenesis in vitro. *Biochem Biophys Res Commun* 189, 824-831.
- Ramirez-Bergeron, D.L., Runge, A., Adelman, D.M., Gohil, M., and Simon, M.C. (2006). HIF-dependent hematopoietic factors regulate the development of the embryonic vasculature. *Developmental cell* 11, 81-92.
- Rocha, F.G., Sundback, C.A., Krebs, N.J., Leach, J.K., Mooney, D.J., Ashley, S.W., Vacanti, J.P., and Whang, E.E. (2008). The effect of sustained delivery of vascular endothelial growth factor on angiogenesis in tissue-engineered intestine. *Biomaterials* 29, 2884-2890.
- Sheridan, M.H., Shea, L.D., Peters, M.C., and Mooney, D.J. (2000). Bioabsorbable polymer scaffolds for tissue engineering capable of sustained growth factor delivery. *J Control Release* 64, 91-102.
- Shibuya, M., Yamaguchi, S., Yamane, A., Ikeda, T., Tojo, A., Matsushime, H., and Sato, M. (1990). Nucleotide sequence and expression of a novel human receptor-type tyrosine kinase gene (flt) closely related to the fms family. *Oncogene* 5, 519-524.
- Soker, S., Fidler, H., Neufeld, G., and Klagsbrun, M. (1996). Characterization of novel vascular endothelial growth factor (VEGF) receptors on tumor cells that bind VEGF165 via its exon 7-encoded domain. *J Biol Chem* 271, 5761-5767.
- Soker, S., Takashima, S., Miao, H.Q., Neufeld, G., and Klagsbrun, M. (1998). Neuropilin-1 is expressed by endothelial and tumor cells as an isoform-specific receptor for vascular endothelial growth factor. *Cell* 92, 735-745.
- Stefanini, M.O., Wu, F.T., Mac Gabhann, F., and Popel, A.S. (2009). The presence of VEGF receptors on the luminal surface of endothelial cells affects VEGF distribution and VEGF signaling. *PLoS computational biology* 5, e1000622.
- Stein, I., Itin, A., Einat, P., Skaliter, R., Grossman, Z., and Keshet, E. (1998). Translation of vascular endothelial growth factor mRNA by internal ribosome entry: implications for translation under hypoxia. *Mol Cell Biol* 18, 3112-3119.
- Tafuro, S., Ayuso, E., Zacchigna, S., Zentilin, L., Moimas, S., Dore, F., and Giacca, M. (2009). Inducible adeno-associated virus vectors promote functional angiogenesis in adult organisms via regulated vascular endothelial growth factor expression. *Cardiovasc Res* 83, 663-671.

- Takahashi, H., and Shibuya, M. (2005). The vascular endothelial growth factor (VEGF)/VEGF receptor system and its role under physiological and pathological conditions. *Clinical science* 109, 227-241.
- Takeshita, S., Weir, L., Chen, D., Zheng, L.P., Riessen, R., Bauters, C., Symes, J.F., Ferrara, N., and Isner, J.M. (1996). Therapeutic angiogenesis following arterial gene transfer of vascular endothelial growth factor in a rabbit model of hindlimb ischemia. *Biochem Biophys Res Commun* 227, 628-635.
- Takeshita, S., Zheng, L.P., Brogi, E., Kearney, M., Pu, L.Q., Bunting, S., Ferrara, N., Symes, J.F., and Isner, J.M. (1994). Therapeutic angiogenesis. A single intraarterial bolus of vascular endothelial growth factor augments revascularization in a rabbit ischemic hind limb model. *J Clin Invest* 93, 662-670.
- Tang, N., Wang, L., Esko, J., Giordano, F.J., Huang, Y., Gerber, H.P., Ferrara, N., and Johnson, R.S. (2004). Loss of HIF-1 $\alpha$  in endothelial cells disrupts a hypoxia-driven VEGF autocrine loop necessary for tumorigenesis. *Cancer Cell* 6, 485-495.
- Terman, B.I., Carrion, M.E., Kovacs, E., Rasmussen, B.A., Eddy, R.L., and Shows, T.B. (1991). Identification of a new endothelial cell growth factor receptor tyrosine kinase. *Oncogene* 6, 1677-1683.
- Tsuboi, R., and Rifkin, D.B. (1990). Recombinant basic fibroblast growth factor stimulates wound healing in healing-impaired db/db mice. *The Journal of experimental medicine* 172, 245-251.
- Verbridge, S.S., Choi, N.W., Zheng, Y., Brooks, D.J., Stroock, A.D., and Fischbach, C. (2010). Oxygen-controlled three-dimensional cultures to analyze tumor angiogenesis. *Tissue Eng Part A* 16, 2133-2141.
- Wang, X., Zhou, X., Ma, J., Tian, H., Jiao, Y., Zhang, R., Huang, Z., Xiao, J., Zhao, B., Qian, H., and Li, X. (2010). Effects of keratinocyte growth factor-2 on corneal epithelial wound healing in a rabbit model of carbon dioxide laser injury. *Biological & pharmaceutical bulletin* 33, 971-976.
- Wernike, E., Montjovent, M.O., Liu, Y., Wismeijer, D., Hunziker, E.B., Siebenrock, K.A., Hofstetter, W., and Klenke, F.M. (2010). VEGF incorporated into calcium phosphate ceramics promotes vascularisation and bone formation in vivo. *Eur Cell Mater* 19, 30-40.
- Wilcke, I., Lohmeyer, J.A., Liu, S., Condurache, A., Kruger, S., Mailander, P., and Machens, H.G. (2007). VEGF(165) and bFGF protein-based therapy in a slow release system to improve angiogenesis in a bioartificial dermal substitute in vitro and in vivo. *Langenbeck's archives of surgery / Deutsche Gesellschaft fur Chirurgie* 392, 305-314.
- Wu, S., Wu, X., Zhu, W., Cai, W.J., Schaper, J., and Schaper, W. (2010). Immunohistochemical study of the growth factors, aFGF, bFGF, PDGF-AB, VEGF-A and its receptor (Flk-1) during arteriogenesis. *Mol Cell Biochem* 343, 223-229.
- Yamada, Y., Nezu, J., Shimane, M., and Hirata, Y. (1997). Molecular cloning of a novel vascular endothelial growth factor, VEGF-D. *Genomics* 42, 483-488.
- Yamazaki, Y., Matsunaga, Y., Tokunaga, Y., Obayashi, S., Saito, M., and Morita, T. (2009). Snake venom Vascular Endothelial Growth Factors (VEGF-Fs) exclusively vary their structures and functions among species. *J Biol Chem* 284, 9885-9891.
- Yla-Herttuala, S. (2009). Gene therapy with vascular endothelial growth factors. *Biochem Soc Trans* 37, 1198-1200.

Zacchigna, S., Tasciotti, E., Kusmic, C., Arsic, N., Sorace, O., Marini, C., Marzullo, P., Pardini, S., Petroni, D., Pattarini, L., Moimas, S., Giacca, M., and Sambuceti, G. (2007). In vivo imaging shows abnormal function of vascular endothelial growth factor-induced vasculature. *Human gene therapy* 18, 515-524.

IntechOpen

IntechOpen



## **Regenerative Medicine and Tissue Engineering - Cells and Biomaterials**

Edited by Prof. Daniel Eberli

ISBN 978-953-307-663-8

Hard cover, 588 pages

**Publisher** InTech

**Published online** 29, August, 2011

**Published in print edition** August, 2011

Tissue Engineering may offer new treatment alternatives for organ replacement or repair deteriorated organs. Among the clinical applications of Tissue Engineering are the production of artificial skin for burn patients, tissue engineered trachea, cartilage for knee-replacement procedures, urinary bladder replacement, urethra substitutes and cellular therapies for the treatment of urinary incontinence. The Tissue Engineering approach has major advantages over traditional organ transplantation and circumvents the problem of organ shortage. Tissues reconstructed from readily available biopsy material induce only minimal or no immunogenicity when reimplanted in the patient. This book is aimed at anyone interested in the application of Tissue Engineering in different organ systems. It offers insights into a wide variety of strategies applying the principles of Tissue Engineering to tissue and organ regeneration.

### **How to reference**

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Remo A. Largo, Venkat M. Ramakrishnan, Martin Ehrbar, Algirdas Ziogas, Jan A. Plock and Daniel Eberli (2011). Angiogenesis and Vascularity for Tissue Engineering Applications, Regenerative Medicine and Tissue Engineering - Cells and Biomaterials, Prof. Daniel Eberli (Ed.), ISBN: 978-953-307-663-8, InTech, Available from: <http://www.intechopen.com/books/regenerative-medicine-and-tissue-engineering-cells-and-biomaterials/angiogenesis-and-vascularity-for-tissue-engineering-applications>

**INTECH**  
open science | open minds

### **InTech Europe**

University Campus STeP Ri  
Slavka Krautzeka 83/A  
51000 Rijeka, Croatia  
Phone: +385 (51) 770 447  
Fax: +385 (51) 686 166  
[www.intechopen.com](http://www.intechopen.com)

### **InTech China**

Unit 405, Office Block, Hotel Equatorial Shanghai  
No.65, Yan An Road (West), Shanghai, 200040, China  
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元  
Phone: +86-21-62489820  
Fax: +86-21-62489821



© 2011 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike-3.0 License](https://creativecommons.org/licenses/by-nc-sa/3.0/), which permits use, distribution and reproduction for non-commercial purposes, provided the original is properly cited and derivative works building on this content are distributed under the same license.

IntechOpen

IntechOpen