

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

186,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Biochemical Properties of MutL, a DNA Mismatch Repair Endonuclease

Kenji Fukui¹, Atsuhiko Shimada², Hitoshi Iino¹,
Ryoji Masui^{1,2} and Seiki Kuramitsu^{1,2}

¹RIKEN SPring-8 Center and

²Graduate School of Science,
Osaka University

Japan

1. Introduction

DNA mismatch repair (MMR) is one of the most widely conserved DNA repair systems, which repairs mismatched bases generated mainly by the error of DNA polymerases during replication (Friedberg, et al., 2006, Iyer, et al., 2006, Kunkel, et al., 2005, Morita, et al., 2010). MMR increases the replication fidelity by 20 to 400-fold (Schaaper, 1993). Mutations and epigenetic silencing in MMR genes cause human hereditary nonpolyposis colon cancers as well as sporadic tumors (Fishel, et al., 1995, Fishel, et al., 1994, Kane, et al., 1997, Leach, et al., 1993, Modrich, et al., 1996, Suter, et al., 2004), indicating the significance of this repair system.

To date, two types of MMR mechanisms have been clarified: one is employed by eukaryotes and most bacteria (Fig. 1A and B) (Modrich, 2006) and the other is specific to *Escherichia coli* and other γ -proteobacteria (Fig. 1C) (Modrich, et al., 1996). The fundamental mechanism and the required proteins in the two types of MMRs are relatively similar to each other. A mismatch is recognized by the bacterial MutS homodimer, eukaryotic MutS α (MSH2-MSH6 heterodimer), or MutS β (MSH2-MSH3 heterodimer) (Acharya, et al., 2003, Drotschmann, et al., 2002, Gradia, et al., 1997, Gradia, et al., 1999, Lamers, et al., 2000, McCulloch, et al., 2003, Obmolova, et al., 2000, Tachiki, et al., 2000). Subsequently, the bacterial MutL homodimer or eukaryotic MutL α (MLH1-PMS2 and MLH1-PMS1 heterodimers in humans and yeast, respectively) is recruited to the mismatched DNA to stimulate downstream events (Acharya, et al., 2003, Kadyrov, et al., 2006). The largest difference between the two types of MMR mechanisms is in the “strand discrimination” system. Although both bases constituting the mismatch are canonical, MMR needs to identify which base is to be repaired. In eukaryotes and most bacteria, MMR directs the repair to the error-containing strand of the mismatched duplex by recognizing the strand discontinuities in the newly synthesized strand (Kadyrov, et al., 2006, Kadyrov, et al., 2007, Larrea, et al., 2010, Modrich, 2006). The termini of leading and lagging strands are thought to serve as discrimination signals. On the other hand, *E. coli* MMR reads the absence of methylation at the restriction site in the newly synthesized strand (Iyer, et al., 2006, Kunkel, et al., 2005, Larrea, et al., 2010). Before the site-specific DNA methylase (e.g., *E. coli* Dam methylase (Schlagman, et




Fig. 1. A schematic representation of MMR pathway models. (A) Eukaryotic MMR. A mismatch is recognized by MutS α , and MutL α nicks the 3'- or 5'-side of the mismatched base on the discontinuous strand. The effective incision by MutL α requires MutS α , replication factor C, proliferating cell nuclear antigen (PCNA), and ATP. The resulting DNA segment is excised by a 5'-3' exonuclease, EXO1, in cooperation with a single-stranded DNA-binding protein, replication protein A (RPA). The DNA strand is resynthesized by DNA polymerase δ and DNA ligase 1. No DNA helicase has been identified to participate in eukaryotic MMR. This mechanism is elucidated by using an *in vitro* reconstituted system.

The pre-existing strand discontinuity can be located on both 5'- and 3'-sides of the mismatch; therefore, there should be 5'- and 3'-nick directed MMR mechanisms. The detail has been described elsewhere (Constantin, et al., 2005, Dzantiev, et al., 2004, Fukui, 2010, Genschel, et al., 2002, Kadyrov, et al., 2009). (B) A speculative model for MMR in *mutH*-less bacteria. The mismatch is recognized by a MutS homodimer. After incision of the discontinuous strand by MutL, the error-containing DNA strand is removed by the cooperative functions of DNA helicases, such as UvrD, the exonucleases RecJ and ExoI, and the single-stranded DNA-binding protein (SSB). DNA polymerase III and DNA ligase fill the gap to complete the repair. Although, no studies have reported the *in vitro* reconstituted system of bacterial nick-directed MMR, it has been elucidated that the endonuclease activity of MutL is required for *in vivo* MMR activity (Fukui, et al., 2008). The involvement of RecJ and ExoI in this MMR system has been implicated experimentally (Shimada et al., 2010). (C) *E. coli* MMR. MutS recognizes the mismatch, and MutL interacts with MutS. Subsequently, the MutH endonuclease is activated to incise the unmethylated strand at the GATC site to create an entry point for the excision reaction. DNA helicase, SSB, and several exonucleases are involved in the excision reaction. At least three models have been proposed for the mechanism by which a MutS homologue stimulates downstream events. They are the "Molecular switch", "Stationary", and "Translocation" models. The major difference between these models is whether a MutS homologue dissociates from the mismatch after recognizing it. Details have been provided in other publications (Kunkel, et al., 2005, Li, 2008).

al., 1986)) completes the modification of the newly synthesized strand, hemi-methylated sites exist and serve as strand-discrimination signals. In both MMR systems, a nicking endonuclease plays a central role in the strand discrimination mechanism. In eukaryotes and most bacteria, MutL homologues are thought to incise the discontinuous strand to introduce the entry or termination point of the excision reaction. In *E. coli*, MutH nicks the unmethylated strand of the duplex to generate the excision entry point. After incision by MutL homologues or MutH, the error-containing strand is removed by helicases and exonucleases. DNA polymerases then resynthesize the strand, and DNA ligases seal the nick to complete the repair reaction.

Although *in vivo* MMR achieves mismatch- and daughter strand-specific incision, eukaryotic MutL α and bacterial MutL show an apparently non-specific endonuclease activity against lesion-less DNA *in vitro* (Duppatla, et al., 2009, Fukui, et al., 2008, Kadyrov, et al., 2006, Kadyrov, et al., 2007, Mauris, et al., 2009), indicating that MMR requires a sequence- or structure-non-specific endonuclease activity to introduce an excision entry point wherever it is needed. The regulatory mechanism underlying this apparently non-specific endonuclease activity has been argued (Yang, 2007). Both eukaryotic and bacterial MutL consist of N-terminal ATPase and C-terminal dimerization (also endonuclease) domains (Fig. 2) (Fukui, et al., 2008, Kadyrov, et al., 2006). The two domains are usually separated by a long, flexible linker region. This domain organization is characteristic of the GHKL ATPase/kinase superfamily that undergoes a large conformational change upon ATP binding and/or hydrolysis (Ban, et al., 1999, Dutta, et al., 2000). Generally, ATP binding and/or hydrolysis control the molecular functions of these superfamily proteins. ATP binding- and/or hydrolysis-induced conformational change is expected to be involved in the regulation of MutL endonuclease activity. Recent biochemical characterizations have demonstrated the effects of ATP binding and hydrolysis on the function of MutL (Duppatla, et al., 2009, Fukui,

et al., 2008, Kim, et al., 2009, Mauris, et al., 2009, Sacho, et al., 2008). In addition to ensuring mismatch-specific incision, cells also need to direct the MutL-dependent nicking reaction to the newly synthesized strand of the mismatched duplex. Interactions of MutL with other MMR proteins have been reported to participate in this regulatory mechanism. In this chapter, we review the biochemical properties of MutL endonucleases that are related to those regulatory mechanisms.




Fig. 2. A schematic representation of the domain structure of MutL homologues. ATPase, endonuclease, and dimerization domains are represented by red, blue, and yellow boxes, respectively. Numbers in parentheses indicate the length of each protein. The interdomain linker regions are shown as gray bars. The crystal structures of the human PMS2 N-terminal ATPase domain (PDB ID: 1EA6) (Guarné, et al., 2001), *E. coli* MutL ATPase domain (PDB ID: 1B63) (Ban, et al., 1999), *Bacillus subtilis* MutL C-terminal endonuclease domain (PDB ID: 3KDK) (Pillon, et al., 2010), and *E. coli* MutL C-terminal dimerization domain (PDB ID: 1X9Z) (Guarné, et al., 2004) are shown.

2. Structure of the C-terminal endonuclease and N-terminal ATPase domains of MutL

The C-terminal domain of MutL endonucleases contains two highly conserved sequence motifs (Fig. 3). One of them is the DQHA(x)₂E(x)₄E motif, which is essential for the nicking endonuclease activity (Fukui, et al., 2008, Kadyrov, et al., 2006). Aspartic acid and histidine residues in this motif are expected to coordinate one or two metal ions to catalyze the nicking reaction (Kosinski, et al., 2008, Pillon, et al., 2010, Yang, 2008). The other is the zinc-binding motif CPHGRP (Kosinski, et al., 2008), which is not essential for the nicking endonuclease activity but is required for the *in vivo* MMR activity (Fukui, et al., 2008, Kosinski, et al., 2008). Crystal structures of *Bacillus subtilis* and *Neisseria gonorrhoeae* MutL C-terminal domains (Namadurai, et al., 2010, Pillon, et al., 2010) revealed that their overall




Fig. 3. Amino acid sequence alignment of the C-terminal regions of MutL homologues. Red and blue boxes indicate the conserved sequence motifs, DQHA(x)₂E(x)₄E and CPHGRP, respectively. The numbers on the left and right show the distances from the N-termini for each protein. The numbers in parentheses show the number of residues that are omitted for the sake of clarity. The NCBI Entrez GI numbers of the sequences are as follows: 4505913 (*Homo sapiens* PMS2), 121583910 (*Mus musculus* PMS2), 18411951 (*Arabidopsis thaliana* PMS1), 46562124 (*Saccharomyces cerevisiae* PMS1), 17562796 (*Caenorhabditis elegans* PMS2), 16078768 (*Bacillus subtilis* MutL), 59801161 (*Neisseria gonorrhoeae* MutL), 15926879 (*Staphylococcus aureus* MutL), 55981292 (*Thermus thermophilus* MutL), 15606703 (*Aquifex aeolicus* MutL), 16131992 (*Escherichia coli* MutL), and 16767605 (*Salmonella typhimurium* MutL).

structures, which are dimeric molecules, resemble that of the *E. coli* MutL C-terminal domain (Fig. 4A) (Guarné, et al., 2004, Kosinski, et al., 2005), although the *E. coli* MutL C-terminal domain lacks the DQHA(x)₂E(x)₄E and CPHGRP motifs. In those crystal structures, the CPHGRP motif is located adjacent to the DQHA(x)₂E(x)₄E motif to form a catalytic site in the subunit (Fig. 4B). In the crystal structure of the *B. subtilis* MutL C-terminal domain, two zinc ions are coordinated near the catalytic site by residues including the histidine and cysteine of the DQHA(x)₂E(x)₄E and CPHGRP motifs, respectively (Fig. 4C) (Pillon, et al., 2010). Although the function of these zinc ions has not been precisely explained, the difference between the zinc-bound and zinc-unbound forms of the *B. subtilis* MutL C-terminal domain demonstrated that binding of zinc ions brings about a local structural rearrangement in the catalytic site (Fig. 4C) (Pillon, et al., 2010). Since the addition of zinc ions to the reaction mixture slightly stimulates the nicking endonuclease activity of MutL (Pillon, et al., 2010), the local structural change would be a prerequisite for the formation of the active form of the catalytic site.

The N-terminal ATPase domain of MutL contains a single ATP-binding motif per subunit just like other GHKL superfamily proteins (Ban, et al., 1998, Guarné, et al., 2001). Unlike the C-terminal domain, the amino acid sequence of the N-terminal ATPase domain of the MutL endonuclease is highly homologous to that of *E. coli* MutL (Iino, et al., 2010). Therefore, the crystal structure of the *E. coli* MutL N-terminal domain can be utilized when considering the structure and function of the N-terminal domain of the MutL endonuclease. Ban and Yang described the apo and AMPPNP-bound forms of the *E. coli* MutL N-terminal domain (Ban, et al., 1999, Ban, et al., 1998), which clearly demonstrated the ATP binding-induced conformational change of this domain (Fig. 5A). Upon AMPPNP binding, the disordered region found in the apo structure formed ordered structures, which led to the dimerization




Fig. 4. Crystal structure of the nicking endonuclease domain of *B. subtilis* MutL (One subunit of the dimer is shown). (A) The overall structure of the endonuclease domain of *B. subtilis* MutL (blue) (zinc-bound form, PDB ID: 3KDK) (Pillon, et al., 2010) is superposed onto the dimerization domain of *E. coli* MutL (yellow) (PDB ID: 1X9Z) (Guarné, et al., 2004). The endonuclease domain is comprised of regulatory and dimerization sub-domains. The DQHA(x)₂E(x)₄E and CPHGRP motifs are included in the dimerization sub-domain. (B) The DQHA(x)₂E(x)₄E motif (green) is located near the CPHGRP motif (cyan). Two zinc ions (pink spheres) are coordinated by several residues including the histidine (a green stick) of the DQHA(x)₂E(x)₄E motif and the cysteine and histidine (cyan sticks) of the CPHGRP motif. (C) The zinc ion binding induces a structural rearrangement of the catalytic site in the endonuclease domain. The zinc-bound form (colored) is superposed onto the unbound crystal forms I (white) (PDB ID: 3GAB) and II (gray) (PDB ID: 3KDG) (Pillon, et al., 2010). All structures are shown in a stereo view.

of the N-terminal domain. As with the MutL endonuclease, the crystal structure of the N-terminal domain of human PMS2 has been reported (Fig. 5B) (Guarné, et al., 2001). Intriguingly, the N-terminal domain of PMS2 bound to ATP γ S even in the absence of the N-terminal domain of MLH1, which is the only report concerning ATP binding by a monomeric GHKL superfamily protein. However, it is expected that in the presence of the MLH1 subunit, ATP binding induces dimerization of the N-terminal domains. In line with this notion, a direct observation using atomic force microscopy suggested that ATP binding causes dimerization of the N-terminal domain in yeast MutL α (Sacho, et al., 2008).




Fig. 5. Crystal structures of the N-terminal ATPase domains of *E. coli* MutL and human PMS2. (A) Stereo view of the *E. coli* MutL N-terminal ATPase domain in the apo form (gray) (PDB ID: 1BKN) and AMPPNP-bound form (red) (PDB ID: 1B63) (Ban, et al., 1999, Ban, et al., 1998). AMPPNP and a magnesium ion are shown as a pink stick and sphere, respectively. (B) Stereo view of human PMS2 N-terminal ATPase domain in the apo form (gray) (PDB ID: 1H7S) and ATP γ S-bound form (red) (PDB ID: 1H7U) (Guarné, et al., 2001). ATP γ S and a magnesium ion are shown as a pink stick and sphere, respectively.

3. ATP modulates the nicking endonuclease activity of MutL

The effect of ATP on the biochemical properties of the MutL endonuclease has been examined using the bacterial MutL endonuclease as a model molecule. *Thermus thermophilus* MutL stably bound one ATP molecule per subunit at a physiological concentration (2 mM) of ATP without any detectable hydrolysis activity in the absence of MutS and mismatch (Fukui, et al., 2008). Limited proteolysis indicated the ATP- or AMPPNP-dependent conformational change of *T. thermophilus* MutL (Fukui, et al., 2008).

In order to detect a nicking endonuclease activity, the covalently closed circular form of plasmid DNA is often used as a substrate (Fukui, et al., 2007). A nicking endonuclease activity converts the closed circular form into an open circular form of the plasmid DNA that can be easily separated from the closed circular form and the linearized form by agarose gel electrophoresis. Mn²⁺ facilitates the mismatch-, MutS-, clamp-, and clamp loader-independent incision of the closed circular form by non-sequence-specific MutL endonuclease activity (Duppatla, et al., 2009, Fukui, et al., 2008, Kadyrov, et al., 2006, Mauris, et al., 2009).

When *T. thermophilus* MutL was preincubated with physiological concentrations of ATP or AMPPNP before to the addition of substrate DNA, the initial rate of the nicking activity was significantly reduced (Fukui, et al., 2008). This was also supported by the result of a gel electrophoretic mobility shift assay, which indicates that ATP or AMPPNP prevents the non-specific DNA binding of *T. thermophilus* MutL (Fukui, et al., 2008). The endonuclease activities of *Aquifex aeolicus* and *N. gonorrhoeae* MutL were also suppressed by the addition of ATP (Duppatla, et al., 2009, Fukui, et al., 2008). One may speculate that the observed suppressing effect is due to the chelating ability of ATP to deprive the manganese ion from MutL. However, this possibility is ruled out by the following two experimental evidences: ATP has no inhibitory effect on the endonuclease activity of the C-terminal domain of MutL (Duppatla, et al., 2009); alteration of the cysteine residue in the CPHGRP motif to an alanine results in perturbation of the suppressing effect of ATP (Fukui, et al., 2008). These results suggest that ATP-dependent suppression requires the binding of ATP to the N-terminal domain and that the zinc ion in the C-terminal domain is required for sensing ATP binding. Interestingly, AMPPNP and a mismatch facilitated the stable interaction between *T. thermophilus* MutL and MutS (Fukui, et al., 2008). The ATP-bound form of MutL would specifically interact with the MutS-DNA complex in the presence of a mismatch. Because the ATPase activity of MutL is expected to be stimulated by its interaction with MutS, the formation of the MutS-MutL complex may promote the endonuclease activity of MutL by unlocking the ATP binding-dependent suppression (Fig. 6).




Fig. 6. A speculative model of the regulatory mechanism for the mismatch-specific enhancement of MutL nicking endonuclease activity. *NTD* and *CTD* represent the N- and C-terminal domains, respectively. ATP binding induces the dimerization of NTD and the approach of NTD to CTD. DNA-unbound MutL exists as an ATP-bound form whose endonuclease activity is inactive, but preferably binds to the MutS-DNA complex. The interaction with the MutS-DNA complex and other MMR proteins induces the ATP hydrolysis of MutL. This ATP hydrolysis induces the tight contact between NTD and CTD, resulting in the stimulation of endonuclease activity.

Mauris and Evans reported the detailed biochemical experiment on *A. aeolicus* MutL, in which they demonstrated that ATP stimulates the nicking endonuclease activity of relatively high concentrations of *A. aeolicus* MutL in the absence of MutS and mismatch (Mauris, et al., 2009). This result suggests that the effect of ATP on the MutL endonuclease activity depends

on the concentration of MutL, which is consistent with the previous report describing the promoting effect of ATP on the nicking endonuclease activity of eukaryotic MutL α (Kadyrov et al., 2006, Kadyrov et al., 2007). These evidences clearly indicate that ATP is utilized not only to suppress the non-specific endonuclease activity of MutL but also to actively enhance its activity. It would therefore be necessary to clarify whether ATP hydrolysis is required for enhancing the endonuclease activity. On one hand, it was reported that AMPPNP can stimulate the endonuclease activity of relatively high concentrations of *A. aeolicus* MutL (Mauris, et al., 2009). On the other hand, the endonuclease activity of *B. subtilis* MutL was not stimulated by AMPPNP even under conditions where ATP could stimulate the activity (Pillon, et al., 2010).

4. The N-terminal ATPase domain stimulates the endonuclease activity of the C-terminal domain

As described in the previous section, the endonuclease activity of MutL is modulated by ATP binding and/or hydrolysis. Because the ATP binding and endonuclease active sites are located in the N- and C-terminal domains, respectively, the interdomain interaction between them had been expected. This prediction was verified by the recent experiment using recombinant N- and C-terminal domains from *A. aeolicus* MutL. The N-terminal domain stimulated the endonuclease activity of the C-terminal domain by at least a 4-fold magnitude in the absence of ATP (Iino, et al., 2010). Interestingly, this promoting effect was abolished by the depletion of zinc ions from the reaction mixture or by the substitution of cysteine in the CPHGRP motif by alanine (Iino, et al., 2010). These results indicate that zinc ions are required for the N-terminal domain-dependent stimulation of the C-terminal domain. It remains to be investigated whether the zinc ions are directly involved in the interdomain interaction or whether they indirectly influence the interaction through rearrangement of the local structure.

It is expected that this interdomain interaction is involved in the ATPase cycle-dependent regulatory mechanism of MutL. Direct observation using atomic force microscopy has suggested the possible ATP binding-induced association of the N-terminal domain to the C-terminal domain (Fig. 6, middle) (Sacho, et al., 2008). Such an approach may reflect the interdomain interaction that is required for stimulating the nicking endonuclease activity. However, as mentioned above, ATP binding suppresses and ATP hydrolysis promotes the nicking endonuclease activity (Duppatla, et al., 2009, Fukui, et al., 2008, Pillon, et al., 2010). Therefore, ATP hydrolysis may create a tighter contact of the N-terminal domain with the C-terminal domain than that created by ATP binding (Fig. 6, right). Such a tight contact may stimulate the nicking endonuclease activity. Further studies are necessary to clarify whether and how ATP hydrolysis affects the structure and function of MutL endonuclease.

5. Interaction with a sliding clamp directs the MutL-dependent incision to the discontinuous strand

In the above sections, we reviewed the possible regulatory mechanism that assures the mismatch-specific nicking endonuclease activity of MutL. We also have to consider a regulatory mechanism that directs the nicking endonuclease activity of MutL to the error-containing strand of the mismatched duplex. Mismatch itself has no signal to discriminate which base is incorrect (Friedberg, et al., 2006). *In vitro* characterization of MMR activity in the

Interestingly, *B. subtilis* MutL endonuclease also interacts with a β -sliding clamp (Pillon, et al., 2011), a bacterial counterpart to eukaryotic PCNA, which also has two distinguishable faces (Kong, et al., 1992). Most bacterial MutL endonucleases have the β clamp-interacting motif QLxLF at the regulatory sub-domain of the C-terminal domain (Fig. 7B) (Pillon, et al., 2010, Pillon, et al., 2011). Mutation of this sequence motif results in defects in the *in vivo* MMR activity (Pillon, et al., 2011), implying that the β -clamp-dependent activation of MutL is necessary in the cell and that bacterial MMR also adopts a strand discrimination mechanism similar to that of eukaryotic MMR. However, MutL endonucleases from the Aquificae phylum lack the regulatory sub-domain (Iino, et al., 2010). In addition, MutL endonucleases from the Thermus-Deinococcus phylum have no obvious β -clamp-interacting motifs (Fig. 7B), although they retain the regulatory sub-domain. Therefore, it should be carefully investigated whether this discrimination mechanism is universally present among all nick-directed MMRs.

6. Bacterial MutL is a homodimeric nicking endonuclease

Crystal structures of *B. subtilis* and *N. gonorrhoeae* MutL C-terminal domains, and other biochemical studies, have revealed that bacterial MutL C-terminal domains are homodimeric (Duppatla, et al., 2009, Iino, et al., 2010, Namadurai, et al., 2010, Pillon, et al., 2010). Generally, linear double-stranded DNA-specific dimeric endonucleases incise both strands of the duplex. Type II (and Type IIs) restriction endonucleases and Type II DNA topoisomerases are representative of double-strand incising dimeric endonucleases. On the other hand, double-stranded DNA-specific nicking endonucleases are usually monomeric, with the exception of several structure-specific nicking endonucleases (Fukui, et al., 2008, Komori, et al., 2002). For example, the following linear double-stranded DNA-specific nicking endonucleases are all monomeric proteins (Table 1): N-type nicking endonucleases (e.g., *N. BspQI*), sequence-specific nicking endonucleases naturally or artificially created by mutating restriction enzymes to lose their dimerization ability (Higgins, et al., 2001, Roberts, et al., 2003, Xu, et al., 2001, Yunusova, et al., 2006, Zheleznyaya, et al., 2009); V-type nicking endonucleases (e.g., *E. coli* Vsr), a short patch MMR nicking endonuclease (Tsutakawa, et al., 1999, Tsutakawa, et al., 1999); Type I DNA topoisomerases (e.g., *E. coli* Topo I), an enzyme with a supercoil-relaxing activity (Kirkegaard, et al., 1978); retrotransposon-targeting endonucleases (e.g., L1 endonuclease), a site-specific nicking endonuclease that directs the invasion of the retrotransposon (Feng, et al., 1996, Feng, et al., 1998, Maita, et al., 2007, Weichenrieder, et al., 2004); bovine DNase I, a non-specific nicking endonuclease that functions in the host defense (Suck, et al., 1988); *E. coli* MutH (Ban, et al., 1998), the MMR nicking endonuclease; bacterial UvrC (Nazimiec, et al., 2001), a nucleotide excision repair nicking endonuclease; bacterial endonuclease V (Dalhus, et al., 2009), a deaminated DNA-specific nicking endonuclease; and bacterial and eukaryotic AP endonucleases (Hosfield, et al., 1999, Mol, et al., 2000), an abasic site-specific nicking endonuclease. Known DNA repair systems other than MMR all adopt a monomeric nicking endonuclease to introduce the entry point for the excision reaction. Therefore, the dimerization ability of the MutL C-terminal domain might be related to the strand-discrimination mechanism of bacterial MMR.

Enzyme	Cellular function	Substrate	Biological unit	References
N-type nicking endonucleases (e.g., N. BspQI)	Host defense (Artificial)	Asymmetric sequence	Monomer	(Higgins, et al., 2001, Roberts, et al., 2003, Xu, et al., 2001, Yunusova, et al., 2006, Zheleznaya, et al., 2009)
V-type nicking endonucleases (e.g., E. coli Vsr)	DNA repair ⁴ and other	Methylated DNA	Monomer	(Tsutakawa, et al., 1999, Tsutakawa, et al., 1999)
Type I DNA topoisomerases (e.g., E. coli Topo I)	Various DNA transactions	Supercoiled DNA	Monomer	(Kirkegaard, et al., 1978)
Retrotransposon-targeting endonucleases ¹ (e.g., L1 endonuclease)	Targeting of retrotransposon	Target sequence	Monomer	(Feng, et al., 1996, Feng, et al., 1998, Maita, et al., 2007, Weichenrieder, et al., 2004)
Bovine DNase I ¹	Host defense	Non-specific	Monomer	(Suck, et al., 1988)
E. coli MutH	DNA repair ⁴	GATC site	Monomer	(Ban, et al., 1998)
E. coli UvrC	DNA repair ⁴	DNA strand with bulky adducts	Monomer	(Nazimiec, et al., 2001)
E. coli endonuclease V	DNA repair ⁴	Deaminated DNA	Monomer	(Dalhus, et al., 2009)
AP endonucleases ¹ (e.g., human APE1)	DNA repair ⁴	DNA with abasic sites	Monomer	(Hosfield, et al., 1999, Mol, et al., 2000)
Serratia nuclease ²	Host defense	Non-specific	Dimer	(Franke, et al., 1998, Franke, et al., 1999)
Bacterial MutL ³ (e.g., B. subtilis MutL)	DNA repair ⁴	DNA strand with mismatched bases	Dimer	(Namadurai, et al., 2010, Pillon, et al., 2010)

Table 1. Linear double-stranded DNA-specific nicking endonucleases. ¹Structural analyses have revealed that AP endonucleases, retrotransposon-targeting endonucleases, and DNase I are closely related to each other. ²*Serratia* nuclease can convert the covalently closed circular form of plasmid DNA not only to the linear form but also to the open circular form; however, the major product of this nuclease is the double-strand break. *Serratia* nuclease also incises single-stranded DNA, and dimerization is not essential for the nuclease activity. ³MutL shows no structural similarity to other known endonucleases. ⁴These nicking endonucleases introduce the starting point for the excision reaction in damaged or error-containing single-stranded DNA. Among these DNA repair nicking endonucleases, only MutL forms a dimer.

The homodimeric structure of bacterial MutL prompts the question of how the symmetric homodimer generates asymmetric nicking products. As with eukaryotic MutL α , the asymmetry would be derived from the nature of the heterodimer. Eukaryotic MutL α has a single catalytic site for the endonuclease activity. On the other hand, bacterial MutL contains two catalytic sites that are apparently equivalent to each other. It may be possible that bacterial MutL dissociates from the substrate DNA before the catalysis of the second strand incision because of its low velocity, or that the binding of the product to the one subunit induces a non-productive binding mode of the substrate to the other subunit. Alternatively, as proposed by Namadurai *et al.*, the inverted arrangement of the bacterial MutL C-terminal domain dimer may enable interactions with other MMR proteins to interfere with one of the two active sites during the reaction (Namadurai, et al., 2010).

7. Conclusion

In this chapter, the biochemical properties of MutL endonucleases are reviewed, with an emphasis on their regulatory mechanisms. The regulatory mechanism needs to ensure both mismatch- and daughter-strand-specific incisions. The ATPase cycle-dependent conformational and functional changes of the MutL endonucleases are expected to play a central role in these mechanisms. Since the ATPase cycle-dependent conformational change would involve the rearrangement of the interaction between the N- and C-terminal domains, the structural analysis of full-length MutL is urgently required. For the structural analysis, MutL homologues from some thermophilic bacterium may be suitable because of the lack of flexible interdomain linker region as well as their extreme thermostability. However, the interdomain linker region plays a significant role in the *in vitro* function of eukaryotic MutL α (Gorman, et al., 2010). Therefore, it is necessary to carefully judge whether the obtained information is universal among all MutL endonucleases.

8. Acknowledgement

This work was partly supported by Grant-in-Aid for Scientific Research 20570131 (to R. M.) from the Ministry of Education, Science, Sports and Culture of Japan.

9. References

- Acharya, S., Foster, P. L., Brooks, P. & Fishel, R., The coordinated functions of the *E. coli* MutS and MutL proteins in mismatch repair, *Molecular Cell*, Vol. 12, No. 1, (July 2003), pp. 233-246, ISSN 1097-2765
- Ban, C., Junop, M. & Yang, W., Transformation of MutL by ATP binding and hydrolysis: a switch in DNA mismatch repair, *Cell*, Vol. 97, No. 1, (April 1999), pp. 85-97, ISSN 0092-8674
- Ban, C. & Yang, W., Crystal structure and ATPase activity of MutL: implications for DNA repair and mutagenesis, *Cell*, Vol. 95, No. 4, (November 1998), pp. 541-552, ISSN 0092-8674
- Ban, C. & Yang, W., Structural basis for MutH activation in *E.coli* mismatch repair and relationship of MutH to restriction endonucleases, *EMBO Journal*, Vol. 17, No. 5, (March 1998), pp. 1526-1534, ISSN 0261-4189

- Bowman, G. D., O'Donnell, M. & Kuriyan, J., Structural analysis of a eukaryotic sliding DNA clamp-clamp loader complex, *Nature*, Vol. 429, No. 6993, (June 2004), pp. 724-730, ISSN 1476-4687
- Constantin, N., Dzantiev, L., Kadyrov, F. A. & Modrich, P., Human mismatch repair: reconstitution of a nick-directed bidirectional reaction, *Journal of Biological Chemistry*, Vol. 280, No. 48, (December 2005), pp. 39752-39761, ISSN 0021-9258
- Dalhus, B., Arvai, A. S., Rosnes, I., Olsen, O. E., Backe, P. H., Alseth, I., Gao, H., Cao, W., Tainer, J. A. & Bjoras, M., Structures of endonuclease V with DNA reveal initiation of deaminated adenine repair, *Nature Structural and Molecular Biology*, Vol. 16, No. 2, (February 2009), pp. 138-143, ISSN 1545-9985
- Drotschmann, K., Hall, M. C., Shcherbakova, P. V., Wang, H., Erie, D. A., Brownnewell, F. R., Kool, E. T. & Kunkel, T. A., DNA binding properties of the yeast Msh2-Msh6 and Mlh1-Pms1 heterodimers, *Biological Chemistry*, Vol. 383, No. 6, (June 2002), pp. 969-975, ISSN 1431-6730
- Duppatla, V., Bodda, C., Urbanke, C., Friedhoff, P. & Rao, D. N., The C-terminal domain is sufficient for endonuclease activity of *Neisseria gonorrhoeae* MutL, *Biochemical Journal*, Vol. 423, No. 2, (September 2009), pp. 265-277, ISSN 1470-8728
- Dutta, R. & Inouye, M., GHKL, an emergent ATPase/kinase superfamily, *Trends in Biochemical Sciences*, Vol. 25, No. 1, (January 2000), pp. 24-28, ISSN 0968-0004
- Dzantiev, L., Constantin, N., Genschel, J., Iyer, R. R., Burgers, P. M. & Modrich, P., A defined human system that supports bidirectional mismatch-provoked excision, *Molecular Cell*, Vol. 15, No. 1, (July 2004), pp. 31-41, ISSN 1097-2765
- Feng, Q., Moran, J. V., Kazazian, H. H., Jr. & Boeke, J. D., Human L1 retrotransposon encodes a conserved endonuclease required for retrotransposition, *Cell*, Vol. 87, No. 5, (November 1996), pp. 905-916, ISSN 0092-8674
- Feng, Q., Schumann, G. & Boeke, J. D., Retrotransposon R1Bm endonuclease cleaves the target sequence, *Proceedings of the National Academy of Sciences of the United States of America*, Vol. 95, No. 5, (March 1998), pp. 2083-2088, ISSN 0027-8424
- Fishel, R. & Kolodner, R. D., Identification of mismatch repair genes and their role in the development of cancer, *Current Opinion in Genetics & Development*, Vol. 5, No. 3, (June 1995), pp. 382-395, ISSN 0959-437X
- Fishel, R., Lescoe, M. K., Rao, M. R., Copeland, N. G., Jenkins, N. A., Garber, J., Kane, M. & Kolodner, R., The human mutator gene homolog MSH2 and its association with hereditary nonpolyposis colon cancer, *Cell*, Vol. 77, No. 1, (April 1994), p. 1 p following 166, ISSN 0092-8674
- Franke, I., Meiss, G., Blecher, D., Gimadutdinow, O., Urbanke, C. & Pingoud, A., Genetic engineering, production and characterisation of monomeric variants of the dimeric *Serratia marcescens* endonuclease, *FEBS Letters*, Vol. 425, No. 3, (April 1998), pp. 517-522, ISSN 0014-5793
- Franke, I., Meiss, G. & Pingoud, A., On the advantage of being a dimer, a case study using the dimeric *Serratia* nuclease and the monomeric nuclease from *Anabaena* sp. strain PCC 7120, *Journal of Biological Chemistry*, Vol. 274, No. 2, (January 1999), pp. 825-832, ISSN 0021-9258

- Friedberg, E. C., Walker, G. C., Siede, W., Wood, R. D., Schultz, R. A., & Ellenberger, T. (2006) DNA repair and mutagenesis, 2nd edition, Washington, DC: American Society for Microbiology, ISBN 978-1555813194
- Fukui, K., DNA mismatch repair in eukaryotes and bacteria, *Journal of Nucleic Acids*, Vol. 2010, (July 2010), p. 260512, ISSN 2090-021X
- Fukui, K., Kosaka, H., Kuramitsu, S. & Masui, R., Nuclease activity of the MutS homologue MutS2 from *Thermus thermophilus* is confined to the Smr domain, *Nucleic Acids Research*, Vol. 35, No. 3, (January 2007), pp. 850-860, ISSN 1362-4962
- Fukui, K., Nakagawa, N., Kitamura, Y., Nishida, Y., Masui, R. & Kuramitsu, S., Crystal structure of MutS2 endonuclease domain and the mechanism of homologous recombination suppression, *Journal of Biological Chemistry*, Vol. 283, No. 48, (November 2008), pp. 33417-33427, ISSN 0021-9258
- Fukui, K., Nishida, M., Nakagawa, N., Masui, R. & Kuramitsu, S., Bound nucleotide controls the endonuclease activity of mismatch repair enzyme MutL, *Journal of Biological Chemistry*, Vol. 283, No. 18, (May 2008), pp. 12136-12145, ISSN 0021-9258
- Genschel, J., Bazemore, L. R. & Modrich, P., Human exonuclease I is required for 5' and 3' mismatch repair, *Journal of Biological Chemistry*, Vol. 277, No. 15, (April 2002), pp. 13302-13311, ISSN 0021-9258
- Georgescu, R. E., Kim, S. S., Yurieva, O., Kuriyan, J., Kong, X. P. & O'Donnell, M., Structure of a sliding clamp on DNA, *Cell*, Vol. 132, No. 1, (January 2008), pp. 43-54, ISSN 0092-8674
- Gorman, J., Plys, A. J., Visnapuu, M. L., Alani, E. & Greene, E. C., Visualizing one-dimensional diffusion of eukaryotic DNA repair factors along a chromatin lattice, *Nature Structural and Molecular Biology*, Vol. 17, No. 8, (August 2010), pp. 932-938, ISSN 1545-9985
- Gradia, S., Acharya, S. & Fishel, R., The human mismatch recognition complex hMSH2-hMSH6 functions as a novel molecular switch, *Cell*, Vol. 91, No. 7, (December 1997), pp. 995-1005, ISSN 0092-8674
- Gradia, S., Subramanian, D., Wilson, T., Acharya, S., Makhov, A., Griffith, J. & Fishel, R., hMSH2-hMSH6 forms a hydrolysis-independent sliding clamp on mismatched DNA, *Molecular Cell*, Vol. 3, No. 2, (February 1999), pp. 255-261, ISSN 1097-2765
- Guarné, A., Junop, M. S. & Yang, W., Structure and function of the N-terminal 40 kDa fragment of human PMS2: a monomeric GHL ATPase, *EMBO Journal*, Vol. 20, No. 19, (October 2001), pp. 5521-5531, ISSN 0261-4189
- Guarné, A., Ramon-Maiques, S., Wolff, E. M., Ghirlando, R., Hu, X., Miller, J. H. & Yang, W., Structure of the MutL C-terminal domain: a model of intact MutL and its roles in mismatch repair, *EMBO Journal*, Vol. 23, No. 21, (October 2004), pp. 4134-4145, ISSN 0261-4189
- Gulbis, J. M., Kelman, Z., Hurwitz, J., O'Donnell, M. & Kuriyan, J., Structure of the C-terminal region of p21(WAF1/CIP1) complexed with human PCNA, *Cell*, Vol. 87, No. 2, (October 1996), pp. 297-306, ISSN 0092-8674

- Higgins, L. S., Besnier, C. & Kong, H., The nicking endonuclease N.BstNBI is closely related to type IIs restriction endonucleases *MlyI* and *PleI*, *Nucleic Acids Research*, Vol. 29, No. 12, (June 2001), pp. 2492-2501, ISSN 1362-4962
- Hosfield, D. J., Guan, Y., Haas, B. J., Cunningham, R. P. & Tainer, J. A., Structure of the DNA repair enzyme endonuclease IV and its DNA complex: double-nucleotide flipping at abasic sites and three-metal-ion catalysis, *Cell*, Vol. 98, No. 3, (August 1999), pp. 397-408, ISSN 0092-8674
- Iino, H., Kim, K., Shimada, A., Masui, R., Kuramitsu, S. & Fukui, K., Characterization of C- and N-terminal domains of *Aquifex aeolicus* MutL endonuclease: N-terminal domain stimulates the endonuclease activity of C-terminal domain in a zinc-dependent manner, *Bioscience Reports*, Vol. 31, No. 5, (October 2011), pp. 309-322, ISSN 0144-8463
- Iyer, R. R., Pluciennik, A., Burdett, V. & Modrich, P. L., DNA mismatch repair: functions and mechanisms, *Chemical Review*, Vol. 106, No. 2, (February 2006), pp. 302-323, ISSN 0009-2665
- Kadyrov, F. A., Dzantiev, L., Constantin, N. & Modrich, P., Endonucleolytic function of MutL α in human mismatch repair, *Cell*, Vol. 126, No. 2, (July 2006), pp. 297-308, ISSN 0092-8674
- Kadyrov, F. A., Genschel, J., Fang, Y., Penland, E., Edelman, W. & Modrich, P., A possible mechanism for exonuclease 1-independent eukaryotic mismatch repair, *Proceedings of the National Academy of Sciences of the United States of America*, Vol. 106, No. 21, (May 2009), pp. 8495-8500, ISSN 1091-6490
- Kadyrov, F. A., Holmes, S. F., Arana, M. E., Lukianova, O. A., O'Donnell, M., Kunkel, T. A. & Modrich, P., *Saccharomyces cerevisiae* MutL α is a mismatch repair endonuclease, *Journal of Biological Chemistry*, Vol. 282, No. 51, (December 2007), pp. 37181-37190, ISSN 0021-9258
- Kane, M. F., Loda, M., Gaida, G. M., Lipman, J., Mishra, R., Goldman, H., Jessup, J. M. & Kolodner, R., Methylation of the hMLH1 promoter correlates with lack of expression of hMLH1 in sporadic colon tumors and mismatch repair-defective human tumor cell lines, *Cancer Research*, Vol. 57, No. 5, (March 1997), pp. 808-811, ISSN 0008-5472
- Kim, T. G., Cha, H. J., Lee, H. J., Heo, S. D., Choi, K. Y., Ku, J. K. & Ban, C., Structural insights of the nucleotide-dependent conformational changes of *Thermotoga maritima* MutL using small-angle X-ray scattering analysis, *Journal of Biochemistry*, Vol. 145, No. 2, (February 2009), pp. 199-206, ISSN 1756-2651
- Kirkegaard, K. & Wang, J. C., *Escherichia coli* DNA topoisomerase I catalyzed linking of single-stranded rings of complementary base sequences, *Nucleic Acids Research*, Vol. 5, No. 10, (October 1978), pp. 3811-3820, ISSN 0305-1048
- Komori, K., Fujikane, R., Shinagawa, H. & Ishino, Y., Novel endonuclease in Archaea cleaving DNA with various branched structure, *Genes & Genetic Systems*, Vol. 77, No. 4, (August 2002), pp. 227-241, ISSN 1341-7568
- Kong, X. P., Onrust, R., O'Donnell, M. & Kuriyan, J., Three-dimensional structure of the β subunit of *E. coli* DNA polymerase III holoenzyme: a sliding DNA clamp, *Cell*, Vol. 69, No. 3, (May 1992), pp. 425-437, ISSN 0092-8674

- Kosinski, J., Plotz, G., Guarné, A., Bujnicki, J. M. & Friedhoff, P., The PMS2 subunit of human MutL α contains a metal ion binding domain of the iron-dependent repressor protein family, *Journal of Molecular Biology*, Vol. 382, No. 3, (Month 2008), pp. 610-627, ISSN 1089-8638
- Kosinski, J., Steindorf, I., Bujnicki, J. M., Giron-Monzon, L. & Friedhoff, P., Analysis of the quaternary structure of the MutL C-terminal domain, *Journal of Molecular Biology*, Vol. 351, No. 4, (October 2005), pp. 895-909, ISSN 0022-2836
- Kunkel, T. A. & Erie, D. A., DNA mismatch repair, *Annual Review of Biochemistry*, Vol. 74, March 2005), pp. 681-710, ISSN 0066-4154
- Lamers, M. H., Perrakis, A., Enzlin, J. H., Winterwerp, H. H., de Wind, N. & Sixma, T. K., The crystal structure of DNA mismatch repair protein MutS binding to a G x T mismatch, *Nature*, Vol. 407, No. 6805, (October 2000), pp. 711-717, ISSN 0028-0836
- Larrea, A. A., Lujan, S. A. & Kunkel, T. A., SnapShot: DNA mismatch repair, *Cell*, Vol. 141, No. 4, (May 2010), p. 730 e731, ISSN 1097-4172
- Leach, F. S., Nicolaides, N. C., Papadopoulos, N., Liu, B., Jen, J., Parsons, R., Peltomaki, P., Sistonen, P., Aaltonen, L. A., Nystrom-Lahti, M., Guan, E. Y., Zhang, J., Meltzer, P. S., Yu, J. W., Kao, F. T., Chen, D. J., Cerosaletti, K. M., Fournier, R. E. K., Todd, S., Lewis, T., Leach, R. J., Naylor, S. L., Weissenbach, J., Mecklin, J. P., Jarvinen, H., Petersen, G. M., Hamilton, S. R., Green, J., Jass, J., Watson, P., Lynch, H. T., Trent, J. M., Chapelle, A., Kinzler, K. W. & Vogelstein, B., Mutations of a *mutS* homolog in hereditary nonpolyposis colorectal cancer, *Cell*, Vol. 75, No. 6, (December 1993), pp. 1215-1225, ISSN 0092-8674
- Lee, S. D. & Alani, E., Analysis of interactions between mismatch repair initiation factors and the replication processivity factor PCNA, *Journal of Molecular Biology*, Vol. 355, No. 2, (January 2006), pp. 175-184, ISSN 0022-2836
- Li, G. M., Mechanisms and functions of DNA mismatch repair, *Cell Research*, Vol. 18, No. 1, (January 2008), pp. 85-98, ISSN 1748-7838
- Maita, N., Aoyagi, H., Osanai, M., Shirakawa, M. & Fujiwara, H., Characterization of the sequence specificity of the R1Bm endonuclease domain by structural and biochemical studies, *Nucleic Acids Research*, Vol. 35, No. 12, (May 2007), pp. 3918-3927, ISSN 1362-4962
- Mauris, J. & Evans, T. C., Adenosine triphosphate stimulates *Aquifex aeolicus* MutL endonuclease activity, *PLoS One*, Vol. 4, No. 9, (September 2009), p. e7175, ISSN 1932-6203
- McCulloch, S. D., Gu, L. & Li, G. M., Nick-dependent and -independent processing of large DNA loops in human cells, *Journal of Biological Chemistry*, Vol. 278, No. 50, (December 2003), pp. 50803-50809, ISSN 0021-9258
- Modrich, P., Mechanisms in eukaryotic mismatch repair, *Journal of Biological Chemistry*, Vol. 281, No. 41, (October 2006), pp. 30305-30309, ISSN 0021-9258
- Modrich, P. & Lahue, R., Mismatch repair in replication fidelity, genetic recombination, and cancer biology, *Annual Review of Biochemistry*, Vol. 65, (March 1996), pp. 101-133, 0066-4154


- Mol, C. D., Izumi, T., Mitra, S. & Tainer, J. A., DNA-bound structures and mutants reveal abasic DNA binding by APE1 and DNA repair coordination [corrected], *Nature*, Vol. 403, No. 6768, (January 2000), pp. 451-456, ISSN 0028-0836
- Morita, R., Nakane, S., Shimada, A., Inoue, M., Iino, H., Wakamatsu, T., Fukui, K., Nakagawa, N., Masui, R. & Kuramitsu, S., Molecular mechanisms of the whole DNA repair system: a comparison of bacterial and eukaryotic systems, *Journal of Nucleic Acids*, Vol. 2010, (October 2010), p. 179594, ISSN 2090-021X
- Namadurai, S., Jain, D., Kulkarni, D. S., Tabib, C. R., Friedhoff, P., Rao, D. N. & Nair, D. T., The C-Terminal Domain of the MutL Homolog from *Neisseria gonorrhoeae* Forms an Inverted Homodimer, *PLoS One*, Vol. 5, No. 10, (October 2010), p. e13726, ISSN 1932-6203
- Nazimiec, M., Lee, C. S., Tang, Y. L., Ye, X., Case, R. & Tang, M., Sequence-dependent interactions of two forms of UvrC with DNA helix-stabilizing CC-1065-N3-adenine adducts, *Biochemistry*, Vol. 40, No. 37, (September 2001), pp. 11073-11081, ISSN 0006-2960
- Obmolova, G., Ban, C., Hsieh, P. & Yang, W., Crystal structures of mismatch repair protein MutS and its complex with a substrate DNA, *Nature*, Vol. 407, No. 6805, (October 2000), pp. 703-710, ISSN 0028-0836
- Pillon, M. C., Lorenowicz, J. J., Uckelmann, M., Klocko, A. D., Mitchell, R. R., Chung, Y. S., Modrich, P., Walker, G. C., Simmons, L. A., Friedhoff, P. & Guarné, A., Structure of the endonuclease domain of MutL: unlicensed to cut, *Molecular Cell*, Vol. 39, No. 1, (July 2010), pp. 145-151, ISSN 1097-4164
- Pillon, M. C., Miller, J. H. & Guarné, A., The endonuclease domain of MutL interacts with the β sliding clamp, *DNA Repair (Amst)*, Vol. 10, No. 1, (January 2011), pp. 87-93, ISSN 1568-7856
- Pluciennik, A., Dzantiev, L., Iyer, R. R., Constantin, N., Kadyrov, F. A. & Modrich, P., PCNA function in the activation and strand direction of MutL α endonuclease in mismatch repair, *Proceedings of the National Academy of Sciences of United States of America*, Vol. 107, No. 37, (September 2010), pp. 16066-16071, ISSN 1091-6490
- Roberts, R. J., Belfort, M., Bestor, T., Bhagwat, A. S., Bickle, T. A., Bitinaite, J., Blumenthal, R. M., Degtyarev, S. K., Dryden, D. T., Dybvig, K., Firman, K., Gromova, E. S., Gumport, R. I., Halford, S. E., Hattman, S., Heitman, J., Hornby, D. P., Janulaitis, A., Jeltsch, A., Josephsen, J., Kiss, A., Klaenhammer, T. R., Kobayashi, I., Kong, H., Kruger, D. H., Lacks, S., Marinus, M. G., Miyahara, M., Morgan, R. D., Murray, N. E., Nagaraja, V., Piekarowicz, A., Pingoud, A., Raleigh, E., Rao, D. N., Reich, N., Repin, V. E., Selker, E. U., Shaw, P. C., Stein, D. C., Stoddard, B. L., Szybalski, W., Trautner, T. A., Van Etten, J. L., Vitor, J. M., Wilson, G. G. & Xu, S. Y., A nomenclature for restriction enzymes, DNA methyltransferases, homing endonucleases and their genes, *Nucleic Acids Research*, Vol. 31, No. 7, (April 2003), pp. 1805-1812, ISSN 1362-4962
- Sacho, E. J., Kadyrov, F. A., Modrich, P., Kunkel, T. A. & Erie, D. A., Direct visualization of asymmetric adenine-nucleotide-induced conformational changes in MutL α , *Molecular Cell*, Vol. 29, No. 1, (January 2008), pp. 112-121, ISSN 1097-2765

- Schaaper, R. M., Base selection, proofreading, and mismatch repair during DNA replication in *Escherichia coli*, *Journal of Biological Chemistry*, Vol. 268, No. 32, (November 1993), pp. 23762-23765, ISSN 0021-9258
- Schlagman, S. L., Hattman, S. & Marinus, M. G., Direct role of the *Escherichia coli* Dam DNA methyltransferase in methylation-directed mismatch repair, *Journal of Bacteriology*, Vol. 165, No. 3, (March 1986), pp. 896-900, ISSN 0021-9193
- Shimada, A., Masui, R., Nakagawa, N., Takahata, Y., Kim, K., Kuramitsu, S. & Fukui, K., A novel single-stranded DNA-specific 3'-5' exonuclease, *Thermus thermophilus* exonuclease I, is involved in several DNA repair pathways, *Nucleic Acids Research*, Vol. 38, No. 17, (September 2010), pp. 5692-5705, ISSN 1362-4962
- Suck, D., Lahm, A. & Oefner, C., Structure refined to 2Å of a nicked DNA octanucleotide complex with DNase I, *Nature*, Vol. 332, No. 6163, (March 1988), pp. 464-468, ISSN 0028-0836
- Suter, C. M., Martin, D. I. & Ward, R. L., Germline epimutation of MLH1 in individuals with multiple cancers, *Nature Genetics*, Vol. 36, No. 5, (May 2004), pp. 497-501, ISSN 1061-4036
- Tachiki, H., Kato, R. & Kuramitsu, S., DNA binding and protein-protein interaction sites in MutS, a mismatched DNA recognition protein from *Thermus thermophilus* HB8, *Journal of Biological Chemistry*, Vol. 275, No. 52, (December 2000), pp. 40703-40709, ISSN 0021-9258
- Tsutakawa, S. E., Jingami, H. & Morikawa, K., Recognition of a TG mismatch: the crystal structure of very short patch repair endonuclease in complex with a DNA duplex, *Cell*, Vol. 99, No. 6, (December 1999), pp. 615-623, ISSN 0092-867
- Tsutakawa, S. E., Muto, T., Kawate, T., Jingami, H., Kunishima, N., Ariyoshi, M., Kohda, D., Nakagawa, M. & Morikawa, K., Crystallographic and functional studies of very short patch repair endonuclease, *Molecular Cell*, Vol. 3, No. 5, (May 1999), pp. 621-628, ISSN 1097-2765
- Weichenrieder, O., Repanas, K. & Perrakis, A., Crystal structure of the targeting endonuclease of the human LINE-1 retrotransposon, *Structure*, Vol. 12, No. 6, (June 2004), pp. 975-986, ISSN 0969-2126
- Xu, Y., Lunnen, K. D. & Kong, H., Engineering a nicking endonuclease N.AlwI by domain swapping, *Proceedings of the National Academy of Sciences of United States of America*, Vol. 98, No. 23, (November 2001), pp. 12990-12995, ISSN 0027-8424
- Yang, W., Human MutL α : the jack of all trades in MMR is also an endonuclease, *DNA Repair (Amst)*, Vol. 6, No. 1, (January 2007), pp. 135-139, ISSN 1568-7864
- Yang, W., An equivalent metal ion in one- and two-metal-ion catalysis, *Nature Structural & Molecular Biology*, Vol. 15, No. 11, (November 2008), pp. 1228-1231, ISSN 1545-9985
- Yao, N., Hurwitz, J. & O'Donnell, M., Dynamics of beta and proliferating cell nuclear antigen sliding clamps in traversing DNA secondary structure, *Journal of Biological Chemistry*, Vol. 275, No. 2, (January 2000), pp. 1421-1432, ISSN 0021-9258
- Yunusova, A. K., Rogulin, E. A., Artyukh, R. I., Zheleznaya, L. A. & Matvienko, N. I., Nickase and a protein encoded by an open reading frame downstream from the nickase *BspD6I* gene form a restriction endonuclease complex, *Biochemistry (Mosc)*, Vol. 71, No. 7, (July 2006), pp. 815-820, ISSN 0006-2979

Zheleznaya, L. A., Kachalova, G. S., Artyukh, R. I., Yunusova, A. K., Perevyazova, T. A. & Matvienko, N. I., Nicking endonucleases, *Biochemistry (Mosc)*, Vol. 74, No. 13, (December 2009), pp. 1457-1466, ISSN 1608-3040

IntechOpen

IntechOpen



DNA Repair - On the Pathways to Fixing DNA Damage and Errors

Edited by Dr. Francesca Storici

ISBN 978-953-307-649-2

Hard cover, 380 pages

Publisher InTech

Published online 09, September, 2011

Published in print edition September, 2011

DNA repair is fundamental to all cell types to maintain genomic stability. A collection of cutting-edge reviews, DNA Repair - On the pathways to fixing DNA damage and errors covers major aspects of the DNA repair processes in a large variety of organisms, emphasizing foremost developments, questions to be solved and new directions in this rapidly evolving area of modern biology. Written by researchers at the vanguard of the DNA repair field, the chapters highlight the importance of the DNA repair mechanisms and their linkage to DNA replication, cell-cycle progression and DNA recombination. Major topics include: base excision repair, nucleotide excision repair, mismatch repair, double-strand break repair, with focus on specific inhibitors and key players of DNA repair such as nucleases, ubiquitin-proteasome enzymes, poly ADP-ribose polymerase and factors relevant for DNA repair in mitochondria and embryonic stem cells. This book is a journey into the cosmos of DNA repair and its frontiers.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Kenji Fukui, Atsuhiko Shimada, Hitoshi Iino, Ryoji Masui and Seiki Kuramitsu (2011). Biochemical Properties of MutL, a DNA Mismatch Repair Endonuclease, DNA Repair - On the Pathways to Fixing DNA Damage and Errors, Dr. Francesca Storici (Ed.), ISBN: 978-953-307-649-2, InTech, Available from: <http://www.intechopen.com/books/dna-repair-on-the-pathways-to-fixing-dna-damage-and-errors/biochemical-properties-of-mutl-a-dna-mismatch-repair-endonuclease>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2011 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike-3.0 License](https://creativecommons.org/licenses/by-nc-sa/3.0/), which permits use, distribution and reproduction for non-commercial purposes, provided the original is properly cited and derivative works building on this content are distributed under the same license.

IntechOpen

IntechOpen