

# We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

185,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index  
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?  
Contact [book.department@intechopen.com](mailto:book.department@intechopen.com)

Numbers displayed above are based on latest data collected.  
For more information visit [www.intechopen.com](http://www.intechopen.com)



# Routine CT- Chest in Primary Evaluation of the Major Blunt Trauma Patients; Pros and Cons

Abdel-Mohsen M. Hamad

*Department of Cardiothoracic Surgery, Tanta University, Tanta, Egypt*

## 1. Introduction

Trauma is a major worldwide public health problem and it is one of the leading causes of death in both industrialized and developing countries. Injuries of the thorax are a major cause of morbidity and mortality in blunt trauma patients. Approximately 20% of trauma-related deaths are attributable to chest injuries (LoCicero and Mattox, 1989).

In trauma patients, a clear history is rarely available as most patients are confused, unconscious or even anesthetized and the clinical findings have been shown to be equivocal or misleading in 20–50% of victims of blunt polytrauma (Poletti et al., 2002).

Consequently, radiology plays a major role in evaluation of the trauma patient.

The Advanced Trauma Life Support (ATLS 2004) course recommended performing the plain film radiography of the chest, abdomen, and cervical spine in all the blunt trauma patients. Nowadays, Chest computed tomography (CCT) is being used with increasing frequency in the evaluation of blunt chest trauma. CCT frequently detects injuries not seen on routine initial chest x-ray (CxR) (occult findings). However, in the vast majority of patients the impact of these findings on patient management is debatable (Blostein et al., 1997, Hamad and regal, 2010).

CT is used primarily to assess for traumatic aortic injuries but also has been shown to be useful in the evaluation of skeletal, pulmonary, airway, and diaphragmatic injuries.

## 2. Chest wall injuries

*Rib fractures* are the most common finding after blunt chest trauma with an incidence reported up to 40%. Chest radiography is routinely used to assist in the diagnosis of rib fractures, even though it has limited sensitivity. It is even more insensitive in showing costochondral fractures. CT is the most sensitive technique for imaging rib fractures, since it can help to determine the site and number of fractures and, more importantly, provide information regarding any associated injuries (Primak and Collins, 2002).

*Sternal fractures* are found in 8–10% of blunt chest traumas and it is a marker of a high-energy trauma. The most common site of the sternal fractures is approximately 2 cm down from the manubrio-sternal joint. Sternal fracture usually cannot be diagnosed on frontal chest radiographs, whereas the lateral projections can detect it with high sensitivity. Spiral

CT, with sagittal and coronal reformations, should be the examination of choice in the suspicion of sternal fracture, as it identifies with high accuracy both the fracture, especially that with minimal dislocation, and the associated lesions.

*Thoracic spine fractures* account for 16% to 30% of all spine fractures. These fractures are usually difficult to detect on routine chest radiographs, especially those located in the upper portion. CT is much more sensitive for diagnosing thoracic spine fractures and is the imaging modality of choice. In addition to soft tissue and lung windows, chest CT scans in the trauma patient also should be viewed in bone windows for skeletal injuries. The most common fractures of the thoracic spine are anterior wedge compression fractures and burst fractures, most of which occur near the thoracolumbar junction (Meyer, 1992). If a thoracic spine fracture is suspected on either plain radiographs or chest CT, a dedicated thoracic spine CT at the level in question should be obtained with sagittal and coronal reconstructions to determine the type of fracture and to assess its stability.

### 3. Pleural space injuries

*Pneumothorax*, an air collection in the pleural space occurs in approximately 30% to 40% of cases of blunt trauma (Sariego et al., 1993). The diagnosis of pneumothorax is important whatever it is small because they may enlarge and progress to tension pneumothorax particularly if the patient undergoes mechanical ventilation or general anesthesia (Enderson et al., 1993). In supine position, pleural air will rise to the most nondependent portion of the thorax, which is the anterior, caudal aspect of the pleural space in the supine patient. Radiographic signs of pneumothorax in the supine trauma patient include 1) the deep sulcus sign, which is a deep, lucent costophrenic sulcus, 2) a relative increase in lucency at the affected lung base, and 3) the double diaphragm sign, which is created by the interfaces between the ventral and dorsal portions of the pneumothorax with the anterior and posterior aspects of the hemidiaphragm. CT is more sensitive for detecting pneumothorax than radiography, particularly in the supine patient. Pneumothoraces that are not apparent on the supine chest radiograph have been shown on CT in 10% to 50% of patients with head and blunt abdominal trauma (Wolfman et al., 1993).

It is generally safe to observe a stable patient with an occult pneumothorax, the situation is more controversial when the patient undergoes PPV. In a randomized study of treatment of occult pneumothorax with or without tube thoracostomy regardless of presence or absence of PPV, there was no difference in the incidence of respiratory distress or the need for emergent tube thoracostomy in either group suggesting that size progression of occult pneumothorax is unrelated to PPV (Brazel et al., 1999).

*Pneumomediastinum* in blunt trauma, the most common cause of pneumomediastinum is from rupture of the alveoli caused by a sudden increase in intra-alveolar pressure. The air then tracks centripetally through the pulmonary interstitium into the mediastinum ("Macklin effect"). On plain radiographs and CT, pneumomediastinum is diagnosed when air is seen outlining mediastinal soft tissue structures and the parietal pleura. The continuous diaphragm sign may be seen when air is present between the pericardium and the diaphragm.

### 4. Lung parenchymal injuries

*Pulmonary contusion* is defined as focal parenchymal injury with edema and alveolar and interstitial hemorrhage. It is reported in 17%–70% of blunt traumas cases, it is usually seen

adjacent to solid structures like vertebrae, ribs, the liver, and the heart (Gavelli et al., 2002). In chest radiographs, lung contusion appears within the first 6-8 hours following trauma as non-segmental, non-lobar, peripheral, and in the form of increased density (Kerns et al., 1990). It is more likely to detect contusions with thoracic CT than with chest radiography (Schild et al., 1989).

*Pulmonary laceration* is defined as the disruption of alveolar spaces with formation of a cavity filled with blood or air, and may occur from penetrating trauma or from shearing forces associated with blunt trauma. It is difficult to detect lacerations with chest radiography as they usually overlap accompanying contusion areas. On CT, pulmonary laceration is characterized by air collections within an area of consolidation. CT detects far more lacerations; in one study of 85 patients with pulmonary laceration, 99 lacerations were present on CT, but only 5 were seen radiographically (Wagner et al., 1988). Pulmonary hematoma results from complete filling of a laceration cavity with blood. Hematomas are seen as well-defined, spherical or oval, homogenous increased densities, both with thoracic CT and chest radiography. A traumatic pneumatocele is a completely air-filled cystic space after either acute laceration or complete resolution of a pulmonary hematoma. The radiographic sequelae of pulmonary laceration may persist for months or years (Mirvis et al., 1992).

## 5. Tracheobronchial injury

Tracheobronchial ruptures due to thoracic trauma are relatively rare, reported in 0.4–1.5% of patients in clinical series of major blunt thoracic trauma. More than 80% of bronchial injuries occur in the main bronchi within 2.5 cm of the carina, with the right side more commonly than the left side (Euathrongchit et al., 2006). Common but nonspecific radiographic findings of tracheobronchial injury are pneumothorax, pneumomediastinum, and subcutaneous emphysema. More specific findings of tracheobronchial injury include persistent pneumothorax after adequate chest tube placement, collapse of the lung away from the hilum ("fallen lung sign"), and overdistension or herniation of the endotracheal balloon. Helical CT with sagittal and coronal reconstructions is more sensitive and specific than radiography (Unger et al., 1989; Wintermark et al., 2001).

## 6. Esophageal injury

Esophageal rupture is extremely rare as a complication of blunt trauma (1/1000 cases of blunt chest trauma). However it must be excluded in any case of mediastinal penetrating trauma. Radiographic signs are not specific and include persistent cervical and mediastinal emphysema, pleural fluid, and abnormal mediastinal contour caused by leakage of fluids, hematoma, or mediastinitis (Rivas et al., 2003). CT findings include the radiographic findings, as well as mediastinal fluid and extraluminal enteric contrast.

## 7. Diaphragmatic injury

Diaphragmatic injuries occur in approximately 1% to 8% of blunt trauma cases (Boulanger et al., 1993). Most injuries occur in the posterolateral portion of the left hemidiaphragm (Worthy et al., 1995). Chest radiographic findings of diaphragmatic injury depend mainly on herniation of hollow viscus into the thorax, and abnormal course of the nasogastric tube. The most common finding of diaphragmatic injury on CT is abrupt discontinuity of the

diaphragm. Other CT findings include herniation of abdominal contents into the thorax and constriction of bowel at the herniation site ("collar sign"). The sensitivity of axial CT for diaphragmatic tears ranges between 70% and 90%, and the specificity is approximately 90% (Murray et al., 1996). Pitfalls in diagnosing diaphragmatic tears on CT occur when intraabdominal blood or hemothorax obscures the diaphragm. Also, small incidental diaphragmatic defects are often identified, particularly in older patients.

## 8. Aortic injury

Diagnosis of acute aortic injury is critical. The frontal chest radiograph is the initial study for the evaluation of aortic injury. The greatest utility of chest radiography is not in diagnosing aortic injury, however, but in excluding it. Normal chest radiograph has a 98% negative predictive value (Pretre et al., 1997). Radiographic findings that may indicate mediastinal hematoma include mediastinal widening, abnormal contour or indistinctness of the aortic knob, apical pleural cap, rightward deviation of the nasogastric tube within the esophagus, rightward deviation of the trachea, downward displacement of the left main stem bronchus, and thickening of the right paratracheal stripe. However, mediastinal widening due to mediastinal hematoma is not specific for aortic injury and can be caused by other injuries such as sternal fractures, thoracic spine fractures, venous injury, or great vessel injury.

According to initial chest radiograph findings, and level of clinical suspicion, further workup for aortic injury may be necessary. For a long time aortography has been considered the gold standard for diagnosing aortic injury. More recently several studies have evaluated the use of contrast-enhanced helical CT as a screening and diagnostic modality and recommended CT for routine evaluation (Mirvis and Shanmuganathan 2007). The chest CT scan performed for possible aortic injury is evaluated for direct evidence of a tear, including abnormal aortic contour, pseudoaneurysm, intimal flap, active extravasation of contrast, or abrupt tapering of the descending aorta relative to the ascending aorta ("pseudocoarctation"). Prospective studies with helical CCT for the evaluation of blunt aortic injury (BAI) shows comparable sensitivity (100% vs. 92%) and negative predictive value (NPV) (100% vs. 97%) when compared to aortography, but performed poorly by comparison in regards to specificity (83% vs. 99%) (Fabian et al., 1998).

## 9. Comment

While conventional X-rays still play an important role as the primary screening method, CT imaging has become an integral part of the trauma screening and resuscitation phase.

In general, The CCT is superior to routine screening CxR with regard to sternal, rib and spine fractures, lung lacerations and contusions, pneumothorax, hemothorax, heart, pericardium, aorta and diaphragm (Guerrero-Lopez et al., 2000, Rivas et al., 2003). However, it is not clear whether CT chest should be used in the general blunt trauma population routinely or on selective basis.

Several authors reported the detection of clinically significant findings more frequently on CCT in 'high risk' blunt trauma admissions, These patients included but were not limited to high-speed motor vehicle collisions, >15 foot falls, pedestrian versus motor vehicle collisions, patients with any sign of thoracic trauma on physical examination or any mediastinal abnormalities on CxR (Demetriades et al., 1998; Exadaktylos et al., 2001). A



significant number of these patients (14–65%) can have a completely normal CxR (Trupka et al., 1997; Demetriades et al., 1998; Exadaktylos et al., 2001; Plurad et al., 2007).

The lethality of the occult injury defines the urgency and pertinence of subsequent clinical action. For example, the finding of an aortic or great vessel injury or spine fracture would mandate critical management or diagnostic manoeuvres, while the diagnosis of an occult pneumothorax may not (Brazel et al., 1999).

Therefore, use of CCT in Selected patients can lead to significant changes in patient management (18– 41%) (Trupka et al., 1997; Guerrero-Lopez et al., 2000; Renton, 2003; Salim et al., 2006; Deunk et al., 2007) while application of CCT more liberally results in little consequential intervention overall (Blostein and Hodgeman, 1997; Plurad et al., 2007; Wisbach et al., 2007) based on these occult diagnoses.

On the other hand, others argue that the higher discovery of injuries with CT is of questionable clinical significance at great costs. For example, it has previously been shown that occult pneumothoraces can be treated expectantly, even with positive pressure ventilation (Guerrero-Lopez et al., 2000; Brazel et al., 1999). Also, The significance of occult rib fractures, hemothorax, or contusion is also questionable because hemothoraces may resolve without intervention (Poole et al., 1993) and there is no treatment for the vast majority of rib fractures other than pain management. The increased use of CCT for blunt trauma, presumably for the diagnosis BAI, has not resulted in a significant increase in detection overall. It can also be questioned as to whether all occult BAIs needs to be treated, because this probably was the case before the evolution of the CCT (Plurad et al., 2007).

The question is, when should CT be used in the general blunt trauma population? Should we scan selectively if clinical examination or plain radiography is abnormal or should we use a lower threshold and scan on a routine basis?

On trying to answer this question, Brink et al., tested a number of risk factors ( age  $\geq 55$  years, abnormal chest PE, altered sensorium, abnormal thoracic spine PE, abnormal chest and thoracic spine CR, abnormal abdominal US or pelvic CR, Hb $<6$  and BE  $< -3$  mmol/l.) as independent predictors of positive findings on chest CT in high-energy blunt trauma patients. Presence of any of these criteria can predict presence of chest injuries on CT with a sensitivity of 95%. However, they reported that if these positive predictors were implemented as scanning indications, 5% of all patients with chest injuries on CT would not be identified. This implies that the chance of missing injuries of the chest remains 13% in the low-risk patients if these patients do not undergo chest CT (Brink et al., 2010).

The liberal use of CT scanning is not without concerns; in addition to the cost issue, radiation exposure, time loss and isolation of patient from medical care are important drawbacks.

There is a general consensus that the current levels of CT radiation may be associated with an increased risk of cancer. The effective radiation dose to all organs from a single full-body CT examination is 12 to 16 milli-Sieverts (mSv). Considering that studies of Japanese atomic bomb survivors who received only 5–150 mSv dose have increased risk of cancer, it is reasonable to assume that the risk of cancer from CCT exposure is small but real. Even the lowest dose in the exposed atomic-bomb survivor population (range, 5–50 mSv; mean, 20 mSv) is associated with an increased cancer mortality risk. This risk is more important in children (Kalra et al., 2004; Brenner et al., 2007; Huda 2007).

In most trauma centers, the CT scanner is located outside of the trauma resuscitation bay, even in a different department of the hospital. This requires patient transportation and isolation from the ER team; this potentially influences the decision to perform CT imaging on a patient-to-patient basis.

In Amsterdam, the Academic Medical Center recently introduced a moveable CT scanner in the trauma room itself (Fung Kon Jin et al., 2008). The aim of this concept is the elimination of patient transfer for CT imaging with subsequent shorter time (79 min. in comparison to 105 min before) until completion of diagnostic imaging in patients requiring CT imaging. Moreover patients in Amsterdam can be treated during CT imaging this progress can encourage ER team to incorporate CT imaging during the initial trauma evaluation thus favoring the decision to go for diagnostic accuracy over immediately performing emergency surgery with Less than optimal insights in the potential injuries of the trauma patient.

On the extreme side, some institutions in Europe have reported their experience with full body CT imaging prior to physical examination by the trauma team in their effort to reduce time (Weninger et al., 2007).

Multiple trauma presentations at the same moment undoubtedly influence the overall time needed for complete radiological evaluation, especially when CT imaging is required. Fung Kon Jin et al in a high-volume level-1 trauma center, reported that complete radiological workup, including CT scanning, of a stable trauma patient is completed in a median of 114 min. Patients that are more severely injured (ISS > 15) were transported faster to CT, resulting in faster diagnostic imaging (Fung Kon Jin et al., 2008).


## 10. References

- American College of Surgeons Committee on Trauma. (2004). ATLS advanced trauma life support program for doctors. American College of Surgeons: Chicago, IL.
- Blostein PA, Hodgman CG. (1997). Computed tomography of the chest in blunt thoracic trauma: results of a prospective study. *J Trauma* 43:13–8.
- Boulanger BR, Milzman DP, Rosati C, Rodriquez A. (1993). A comparison of right and left blunt traumatic diaphragmatic rupture. *J Trauma* 35:255–60.
- Brazel KJ, Stafford RE, Weigelt JA, et al. (1999). Treatment of occult pneumothoraces from blunt trauma. *J Trauma* 46:987–991.
- Brenner DJ, Elliston CD. (2004). Estimated radiation risks potentially associated with full-body CT screening. *Radiology* 232:735–738.
- Brink M, Deunk J, Dekker HM, Edwards MJ, Kool DR, van Vugt AB, van Kuijk C, Blickman JG. (2010). Criteria for the selective use of chest computed tomography in blunt trauma patients. *Eur Radiol* 20(4):818–28.
- Brink M, Deunk J, Dekker HM, et al. (2008). Added value of routine chest MDCT after blunt trauma: evaluation of additional findings and impact on patient management. *Am J Roentgenol* 190:1591–8.
- Demetriades D, Gomez H, Velmahos G, et al. (1998). Routine helical computed tomographic evaluation of the mediastinum in high risk blunt trauma patients. *Arch Surg* 133:1084 –1088.
- Deunk J, Dekker HM, Brink M, van Vugt R, Edwards MJ, van Vugt AB. (2007). The value of indicated computed tomography scan of the chest and abdomen in addition to the conventional radiologic work-up for blunt trauma patients. *J Trauma* 63(4):757–63.
- Enderson BL, Abdalla R, Frame SB, et al. (1993). Tube thoracostomy for occult pneumothorax: a prospective randomized study of its use. *J Trauma* 35:726–9.
- Euathrongchit J, Thoongsuwan N, Stern EJ. (2006). Nonvascular mediastinal trauma. *Radiol Clin North Am* 44:251–258.

- Exadaktylos AK, Sclabas G, Schmid SW, et al. (2001). Do we really need routine computed tomographic scanning in the primary evaluation of blunt chest trauma in patients with "normal" chest radiograph? *J Trauma* 51:1173–1176.
- Fabian TC, David KA, Gavant ML, et al. (1998). Prospective study of blunt aortic injury. *Ann Surg* 227: 666–77.
- Fung Kon Jin PH, van Geene AR, Linnau KF, Jurkovich GJ, Ponsen KJ, Goslings JC. (2009). Time factors associated with CT scan usage in trauma patients. *Eur J Radiol* 72(1):134–8.
- Fung Kon Jin PHP, Goslings JC, Ponsen KJ, et al. (2008). Assessment of a new Trauma Workflow Concept Implementing a Sliding CT Scanner in the Trauma Room: the Effect on Workup Times. *J Trauma* 64(5):1320–6.
- Gavelli G, Canini R, Bertaccini P, et al. (2002). Traumatic injuries: imaging of thoracic injuries. *Eur Radiol* 12:1273–1294.
- Guerrero-Lopez F, Vazquez-Mata G, Cazar- Romero PP, Fernandez-Mondejar E, Guayo-Hoyos E, Linde-Valverde CM. (2000). Evaluation of the utility of computed tomography in the initial assessment of the critical care patient with chest trauma. *Crit Care Med* 28:1370–1375.
- Hamad AM, Regal M. (2009). Is Routine Spiral CT-Chest Justified in Evaluation of the Major Blunt Trauma Patients? *Eur J Trauma* 35:31–4.
- Huda W. (2007). Radiation doses and risks in chest computed tomography examinations. *Proc Am Thorac Soc* 4: 316–20.
- Kalra MK, Maher MM, Rizzo S, et al. (2004). Radiation exposure from chest CT: Issues and strategies. *J Korean Med Sci* 19:159–166.
- Kerns SR, Gay SB. (1990). CT of blunt chest trauma. *Am J Roentgenol* 154:55–60.
- LoCicero III J and Mattox KL. (1989). Epidemiology of chest trauma. *Surg Clin North Am* 69:15–9.
- Meyer S. (1992). Thoracic spine trauma. *Semin Roentgenol* 27:254–61.
- Mirvis SE, Shanmuganathan K. (2007). Diagnosis of blunt traumatic aortic injury: still a nemesis. *Eur J Radiol* 64:27–40.
- Mirvis SE, Templeton P. (1992). Imaging in acute thoracic trauma. *Semin Roentgenol* 27:184–210.
- Murray JG, Gaoili E, Gruden JF, et al. (1996). Acute rupture of the diaphragm due to blunt trauma: diagnostic sensitivity and specificity of CT. *Am J Roentgenol* 166:1035–9.
- Plurad D, Green D, Demetriades D, et al. (2007). The increasing use of chest computed tomography for trauma: is it being overutilized? *J Trauma* 62:631–5.
- Poletti PA, Wintermark M, Schnyder P, Becker CD. (2002). Traumatic injuries: the role of imaging in the management of the polytrauma victim (conservative expectation). *Eur Radiol* 12:969–78.
- Poole GV, Morgan DB, Crenston PE, et al. (1993). Computed tomography in the management of blunt thoracic trauma. *J Trauma* 35:296 –302.
- Prete R and Chilcott M. (1997). Blunt trauma to the heart and great vessels. *N Engl J Med* 336:626–632.
- Primak SL and Collins J. (2002). Blunt nonaortic chest trauma: radiographic and CT findings. *Emergency Radiology* 9:5–12.



- Renton J. (2003). Should helical CT scanning of the thoracic cavity replace the conventional chest x-ray as a primary assessment tool in pediatric trauma? An efficacy and cost analysis. *J Pediatr Surg* 8:793–797.
- Rivas LA, Fishman JE, Munera F, et al. (2003). Multislice CT in thoracic trauma. *Radiol Clin North Am* 41:599–616.
- Salim A, Sangthong B, Martin M, et al. (2006). Whole body imaging in blunt multisystem trauma patients without obvious signs of injury. *Arch Surg* 141:468–475.
- Sariego J, Borwn JL, Matsumoto T, Kerstein MD. (1993). Predictors of pulmonary complications in blunt chest trauma. *Int Surg* 78:320–3.
- Schild HH, Strunk H, Weber W, et al. (1989). Pulmonary contusion CT vs plain radiograms. *J Comput Assist Tomogr* 13:417–420.
- Trupka A, Waydhas C, Hallfeldt K, et al. (1997). Value of thoracic computed tomography in the first assessment of severely injured patients with blunt chest trauma: results of a prospective study. *J Trauma* 43:405–412.
- Unger JM, Schuchmann GG, Grossmann JE, et al. (1989). Tears of the trachea and main bronchi caused by blunt trauma: Radiologic findings. *AJR* 153:1175–80.
- Wagner RB, Crawford Jr. WO Schimpf PP. (1988). Classification of parenchymal injuries of the lung. *Radiology* 167:77–82.
- Weninger P, Mauritz W, Fridrich P, et al. (2007). Emergency room management of patients with blunt major trauma: evaluation of the multislice computed tomography protocol exemplified by an urban trauma center. *J Trauma* 62(3):584–91.
- Wintermark M, Schnyder P, Wicky S. (2001). Blunt traumatic rupture of a mainstem bronchus: spiral CT demonstration of the “fallen lung” sign. *Eur Radiol* 11:409–411.
- Wisbach GG, Sise MJ, Sack DI, et al. (2007). What is the role of chest x-ray in the initial assessment of stable trauma patients? *J Trauma* 62:74–8.
- Wolfman NT, Gilpin JW, Bechtold RE, et al. (1993). Occult pneumothorax in patients with abdominal trauma: CT studies. *J Comput Assist Tomogr* 17:56–9.
- Worthy SA, Kang EY, Hartman TE, Kwong JS, Mayo JR, Müller NL. (1995). Diaphragmatic rupture: CT findings in 11 patients. *Radiology* 194:885–8.



## **CT Scanning - Techniques and Applications**

Edited by Dr. Karupppasamy Subburaj

ISBN 978-953-307-943-1

Hard cover, 348 pages

**Publisher** InTech

**Published online** 30, September, 2011

**Published in print edition** September, 2011

Since its introduction in 1972, X-ray computed tomography (CT) has evolved into an essential diagnostic imaging tool for a continually increasing variety of clinical applications. The goal of this book was not simply to summarize currently available CT imaging techniques but also to provide clinical perspectives, advances in hybrid technologies, new applications other than medicine and an outlook on future developments. Major experts in this growing field contributed to this book, which is geared to radiologists, orthopedic surgeons, engineers, and clinical and basic researchers. We believe that CT scanning is an effective and essential tools in treatment planning, basic understanding of physiology, and and tackling the ever-increasing challenge of diagnosis in our society.

### **How to reference**

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Abdel-Mohsen M. Hamad (2011). Routine CT- Chest in Primary Evaluation of the Major Blunt Trauma Patients; Pros and Cons, CT Scanning - Techniques and Applications, Dr. Karupppasamy Subburaj (Ed.), ISBN: 978-953-307-943-1, InTech, Available from: <http://www.intechopen.com/books/ct-scanning-techniques-and-applications/routine-ct-chest-in-primary-evaluation-of-the-major-blunt-trauma-patients-pros-and-cons>

**INTech**  
open science | open minds

### **InTech Europe**

University Campus STeP Ri  
Slavka Krautzeka 83/A  
51000 Rijeka, Croatia  
Phone: +385 (51) 770 447  
Fax: +385 (51) 686 166  
[www.intechopen.com](http://www.intechopen.com)

### **InTech China**

Unit 405, Office Block, Hotel Equatorial Shanghai  
No.65, Yan An Road (West), Shanghai, 200040, China  
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元  
Phone: +86-21-62489820  
Fax: +86-21-62489821

© 2011 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike-3.0 License](https://creativecommons.org/licenses/by-nc-sa/3.0/), which permits use, distribution and reproduction for non-commercial purposes, provided the original is properly cited and derivative works building on this content are distributed under the same license.

IntechOpen

IntechOpen