We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Angular Dependence of Fluorescence X-Rays and Alignment of Vacancy State Induced by Radioisotopes

İbrahim Han
Ağrı İbrahim Çeçen University
Turkey

1. Introduction

This chapter concerns angular distribution measurements for fluorescence X-ray and the alignments of atoms with inner-shells vacancy resulting from ionization by radioisotope sources. The discussion on this topic is done by evaluating measurements of X-ray fluorescence parameters (such as cross-section, alignment parameter, polarization degree) from sample in various emission angles.

When an atom is ionized in one of its inner shells, the electrons rearrange themselves to fill the vacancy, with the transition energy released as a photon or transferred to another electron. The following X-ray or Auger electron may have an isotropic or non-isotropic angular distribution. The study of alignment of the inner-shell vacancy in ions can provide information about ionization process and the wave functions of inner-shell electrons, and calculations showed that the alignment was a sensitive testing parameter for theoretical models. For the last five decades there have been both theoretically and experimentally renewed efforts towards better understanding of the physics concerned with alignment of atoms with inner-shells vacancy and/or angular dependence of fluorescent X-rays emitted atoms induced photons or charged particle (electrons, protons, heavy ions). Generally, the alignments of atoms with inner-shells vacancy resulting from ionization by photons are investigated by measuring the anisotropic emission of X-ray lines using a detector (such as Si(Li) or Ge(Li)) and radioisotope photon source in various emission angles.

2. Historical background and current status of topic

The aim of paper interested in this topic is to determine the relationship between the angular distributions of X-rays with respect to total angular momentum values (J) of vacancy states. It is well-known that when radioisotope source, X-ray tube or charged particles produce vacancies in atoms at energy levels with $J > 1/2$, the resulting ions will be aligned. The signature of this alignment is the anisotropic angular distribution of the emitted characteristic X-ray radiation, or the degree of polarization of the X-ray radiation. Total angular momentum (J) of vacancy states after photoionization is greater than $1/2$, the population of its magnetic sub-states is non-statistical by the ionized atoms and this is reason of this anisotropic behavior. A lot of theoretical studies have been reported so far

along this topic (Mehlhorn 1968; Mc Farlane, 1972; Berezhko and Kabachnik 1977; Sizov and Kabachnik, 1980, 1983) and the predictions of these researchers have been experimentally supported by some researchers (Schöler and Bell, 1978; Pálincás, 1979, 1982; Wigger et al., 1984; Jesus et al., 1989; Mitra et al., 1996). The experimental study of alignment generally involves measurements of the angular distribution or polarization of the induced X-rays (Hardy et al., 1970; Döbelin et al., 1974; Jamison and Richard, 1977; Jistchin et al., 1979, 1983; Pálincás, et al., 1981; Stachura et al., 1984; Bhalla, 1990; Mehlhorn, 1994; Papp 1999). In 1969, Cooper and Zare, (1969) first suggested a theoretical model relevant to aligned photon induced atoms. According to calculation by Cooper and Zare, (1969), after photoionization the inner-shell vacancy states have statistical population of magnetic substates. The vacancies produced after photoionization in sub-shells are not be aligned at all and so the angular distribution of the fluorescent X-rays subsequent to photoionization will be isotropic. In 1972, 3 years after Cooper and Zare, the predictions of Flügge et al., (1972) showed that when vacancies are created in states with $J > 1/2$, the population of its magnetic sub-states are non-statistical and therefore the resulting ions will be aligned. Mc Farlane (1972) calculated the polarization of X-rays from the decay of a vacancy in the $2p_{2/3}$ sub-shell using hydrogenic wave- functions in the Bethe approximation and the first Born approximation. After Caldwell and Zare (1977) first made an experimental investigation of the photon-induced alignment of Cd and they measured the degree of polarization of the emitted radiation from Cd. Since then, many experiments and calculations have been done to study the alignment of atoms and angular dependence characteristic X-rays by measuring either the angular distribution or the degree of polarization of the emitted X-rays. All these studies confirmed either alignment or not-alignment of the atoms after photoionization. The angular correlation between ionizing and fluorescent X-rays has been calculated relativistically, including all the radiation multipoles using single particle wavefunctions calculated in the Hartree-Slater model, by (Scofield, 1976). More recently, Scofield, (1989) used a relativistic model to study the angular distribution of the photoelectrons produced from photo- ionization by linear polarize photons and its inverse process (radiative recombination) in the energy region of 1-100keV. Scofield, (1989) found that the cross-section has a maximum at 90° compared to the direction of the incoming photons in the x-z plane (polarization plane) while the cross-section is independent of the angle between the incoming photon and the ejected electron in the y-z plane (normal to the polarization plane). Kamiya et al., (1979) measured L X-rays of Ho and Sm produced by protons and ^3He impacts with Si(Li) detector over the incident energy ranges $E_p = 0.75\text{--}4.75\text{MeV}$ and $E_{^3\text{He}} = 1.5\text{--}9.4\text{MeV}$ in the direction of 90° to the projectile. Kamiya, et al., (1979) reported that the ratios of X-ray production cross-sections for the $L\alpha$ and $L\beta$ lines depend clearly on projectile energy, but are independent of the projectile charge. Theoretical values of the alignment parameter for different states of various atoms calculated using the Herman-Skillman wave functions, have been reported by Berezhko and Kabachnik, (1977). The very strong anisotropy was reported for the emission of L lines for various elements by several scientists (Kahlon, et al., 1990a,b, 1991a,b; Ertuğrul, et al., 1995, 1996; Ertuğrul, 1996, 2002; Kumar, et al., 1999 Sharma and Allawadhi, 1999; Seven and Koçak, 2001,2002; Seven, 2004; Demir, et al., 2003). However, in all these investigations, the observed anisotropy is much higher than the predicted theoretical values of Scofield, (1976) and Berezhko and Kabachnik, (1977). On the other hand, anisotropic emission for L X-rays of Pb, Th and U was reported by some scientists (Mehta, et al., 1999; Kumar, et al 1999, 2001). Recently, Yamaoka et al., (2002, 2003) performed experiments using synchrotron radiation to determine the angular distribution of

L X-ray photons of Pb and Au. Although they found an isotropic distribution of the Pb L_3 lines within the experimental errors, non-isotropic angular distribution of the Au L_3 lines have been obtained. Papp and Campbell, (1992) reported the magnitude of the anisotropy and the alignment parameter for the L lines of Er. The alignment parameter of the ions of Xe was obtained by Küst, et al., (2003).

Kahlon et al., (1990a) reported experimental investigation of the alignment of the L_3 subshell vacancy state produced after photoionization in lead by 59.57 keV photons. The values of differential cross sections for the emission of the Ll , $L\alpha$, $L\beta$ and $L\gamma$ X-ray lines were determined at different emission angles varying from 40° to 120° . It was seen from the results that the Ll , and $L\alpha$ peaks show anisotropic emission, while the $L\beta$ and $L\gamma$ peaks are emitted isotropically. The angular dependence of emission intensity of L shell X rays induced by 59.57 keV photons in Pb and U was investigated by Kahlon et al., (1990b) measuring the normalized intensities of the resolved L X-ray peaks at different angles varying from 40° to 140° . It was observed that while the Ll and $L\alpha$ peaks (originating from $J=3/2$ state) show some anisotropic angular distribution, the emission of the $L\beta$ and $L\gamma$ peaks are emitted isotropically. Kahlon et al., (1991a) measured the angular distribution and polarization of the L shell fluorescent X-rays excited by 59.54 keV photons in Th and U. It was found that the $L\gamma$ group of L X-rays is isotropic in spatial distribution and unpolarized but, the Ll and $L\alpha$ groups are anisotropically distributed and polarized. Although no anisotropy of the $L\beta$ group is detected, it was slightly polarized. Kahlon et al., (1991b) investigated the differential cross sections for emission of Ll , $L\alpha_2$, $L\alpha$, $L\beta$ and $L\gamma$ groups of L X-ray lines induced in Au by 59.54 keV photons at different angles varying from 40° to 120° . The L X-rays represented by Ll , $L\alpha_2$ and $L\alpha$ peaks were found to be anisotropic in the spatial distribution while those in $L\beta$ and $L\gamma$ peaks were isotropic. Papp and Campbell, (1992) measured angular distributions of the L_l , $L\alpha_{1,2}$ and $L\beta_{2,15}$ transitions of erbium in the angular range of 70° – 150° following photoionization by 8.904 keV photons. A Johansson-type monochromatic was used to select the $Cu K\beta_1$ line for ionization. Anisotropy parameters for Ll , $L\alpha_{1,2}$ and $L\beta_{2,15}$ were found as 0.052 ± 0.016 , 0.16 ± 0.022 and 0.012 ± 0.015 , respectively. Ertugrul et al., (1995, 1996a, 1996b) measured differential cross-sections for the emission of Ll , $L\alpha$, $L\beta$ and $L\gamma$ X-rays of Au, Hg, Tl, Pb, Bi, Tb and U at different emission angles varying from 45° to 135° . They found that Ll and $L\alpha$ peaks are emitted isotropically, while $L\beta$ and $L\gamma$ peaks show anisotropic emission. Sharma and Allawadhi, (1999) measured values of Ll , $L\alpha$ and $L\beta$ differential X-ray production cross sections in Th and U at 16.896 and 17.781 keV at emission angles 60° , 70° , 80° and 90° . From the results of the measurements it was evident that, in the present case, all the three Ll , $L\alpha$ and $L\beta$ differential X-ray production cross sections depend on the emission angle and thus, the emission is anisotropic. Demir et al., (2000) indicated differential cross-sections for the emission of M shell fluorescence X-rays from Pt, Au and Hg by 5.96 keV photons at seven angles ranging from 50° to 110° at. The differential cross-sections were found to decrease with increase in the emission angle, showing an anisotropic spatial distribution of M shell fluorescence X-rays. Seven and Koçak (2001, 2002) measured the Ll , $L\alpha$, $L\beta$ and $L\gamma$ X-ray production cross-sections in U, Th, Bi, Pb, Tl, Hg, Au, Pt, Re, W, Ta, Hf, Lu, and Yb using 59.5 keV incident photon energies in the angular range 40° – 130° . Although differential cross sections for $L\beta$ and $L\gamma$ X-rays were found to be angle independent within experimental error, those for the Ll and $L\alpha$ X-rays were found to be angle dependent. Ertugrul et al., (2002) measured the alignment parameter the $I_{L\alpha}/I_{Ll}$ intensity ratio. The Ll and $L\alpha$ X-rays of the elements were measured with a Si (Li) detector at a direction of 90° to the projectile. The L_3 edges of Nd,

Gd, Tb, Dy, Ho, Er, Yb, Hf, Ta, W, Au, Hg, Tl, Pb, Bi, Th and U the elements were excited with the K X-ray energy of 17.781(MoK $_{\alpha,\beta}$), 16.896(NbK $_{\alpha,\beta}$), 14.980(RbK $_{\beta}$), 13.300(BrK $_{\beta}$), 12.503(SeK $_{\beta}$), 12.158(BrK $_{\alpha,\beta}$), 10.983(GeK $_{\beta}$), 10.073(GeK $_{\alpha_{1,\beta}}$), 9.572(ZnK $_{\beta}$), 8.976(CuK $_{\beta_2}$), 8.907(CuK $_{\beta}$), 8.265(NiK $_{\beta}$), 7.649(CoK $_{\beta_1}$), 6.490(MnK $_{\beta_1}$) keV from the selected elements, respectively. They noticed that the L $_3$ X-rays show large anisotropy, the measured alignment parameter varying from -0.115 to +0.355. Demir et al., (2003) reported Ll , $L\alpha$, $L\beta$ and $L\gamma$ X-ray differential cross-sections, fluorescence cross-sections and σ_{L1} , σ_{L2} and σ_{L3} subshell fluorescence cross-sections for Er, Ta, W, Au, Hg and Tl at an excitation energy of 59.6 keV. The differential cross-sections for these elements have been measured at different angles varying from 54° to 153°. The Ll and $L\alpha$ groups in the L X-ray lines were found to be spatially anisotropic, while those in the $L\beta$ and $L\gamma$ peaks are isotropic. The Ll , $L\alpha$, $L\beta_{2,4}$, $L\beta_{1,3}$ and $L\gamma$ X-ray production cross-sections and L-subshell fluorescence yields ω_1 and ω_2 in Th and U have been determined by Seven (2004) at an incident photon energy of 59.54 keV by measuring differential cross-sections with angles changing from 40° to 130°. The Ll , $L\alpha$ and $L\beta_{2,4}$ X-rays have an anisotropic spatial distribution while $L\beta_{1,3}$ and $L\gamma$ X-rays have isotropic spatial distributions. Özdemir et al., (2005) measured the angular dependence of L $_3$ subshell to M-shell vacancy transfer probabilities for the elements Lu, Hf, Ta, W, Os and Pt at the excitation energies of 5.96 keV and K X-rays of Zn, Ga, Ge, and As, respectively, at seven angles varying from 120° to 150°. It was observed that angular dependence from L $_3$ subshell to M-shell vacancy transfer probabilities increase with increasing $\cos\theta$. The angular dependence of M X-ray production differential cross-sections for selected heavy elements between Lu and Pt have been measured by Durak (2006) at 5.59 keV of incident photon energy and at seven emission angles in the range of 120°-150°. Angular dependence of M X-ray production differential cross sections has been derived, using the M-shell fluorescence yields, experimental total M X-ray production cross sections and theoretical M-shell photoionization cross sections. M X-ray production differential cross-sections were found to decrease with increase in the emission angle, showing an anisotropic spatial distribution of M X-rays. Angular dependence from L $_3$ subshell to M-shell vacancy transfer probabilities for selected heavy elements from Au to U were measured by Özdemir and Durak (2008) at different angles varying from 120° to 150°. It was observed that angular dependence from L $_3$ -subshell to M-shell vacancy transfer probabilities increase with increasing $\cos\theta$. Apaydın et al., (2008) measured Mi ($i = \alpha + \beta$) X-ray production differential cross sections for Re, Bi and U elements at the 5.96 keV incident photon energy in an angular range 135°-155°. They found that the angular dependence M X-rays production cross sections decrease with increase in the emission angle, showing anisotropic spatial distribution. Kumar et al., (1999) investigated the angular dependence of emission of L x-rays following photoionization at 22.6 and 59.5 keV in 82Pb by measuring the intensity ratios $I_{Ll}/I_{L\gamma}$, $I_{L\alpha}/I_{L\gamma}$ and $I_{L\beta}/I_{L\gamma}$ at different angles varying from 50° to 140°. The measured intensity ratios for various L x-rays were found to be angle independent within experimental error. Mehta et al., (1999) measured the L_l , $L\alpha$, $L\eta$, $L\beta_6$, $L\beta_{2,4}$, $L\beta_{1,3}$, $L\beta_{9,10}$ and $L\gamma$ x-ray production differential cross sections in 92U using the 22.6- and 59.5-keV incident photon energies in an angular range 43°-140°. Differential cross sections for various L x rays were found to be angle independent within experimental error. Puri et al., (1999) measured The Ll , $L\alpha$, $L\beta_{2,4}$, $L\beta_{1,3}$ and $L\gamma_{1,5}$ X-ray production differential cross sections in 90Th have at 22.6 keV incident photon energy in an angular range 50° -130° The measured differential cross sections for various L X-rays were found to be angle-independent within experimental error. Kumar et al., (2001a) measured the the Ll , $L\alpha$ and $L\beta_{2,5,6,7,15}$ X-ray fluorescence (XRF)

differential cross-sections in Pb at the 13.6 keV incident photon energy ($E_{L3} < E_{inc} < E_{L2}$, E_{Li} being the Li sub-shell binding energy) and in the angular range 90-160°. At this incident photon energy, the L_3 sub-shell vacancies ($J = 3/2$) are produced only due to the direct ionization and the reduction in the observed anisotropy in the emission of the Ll , $L\alpha$ and $L\beta_{2,5,6,7,15}$ X-rays due to the transfer of unaligned L_1 and L_2 subshell vacancies ($J = 1/2$) to the L_3 sub-shell through Coster-Kronig transitions was eliminated. The differential cross-sections for various x-rays were found to be angle-independent within experimental error. The L X-ray production (XRP) differential cross sections in Th and U have been measured by Kumar et al., (2001b) at the 17.8 keV incident photon energy ($E_{L3} < E_{inc} < E_{L2}$, E_{Li} is the Li subshell ionization threshold) in an angular range 90°-160° and at the 25.8 and 46.9 keV incident photon energies ($E_{L1} < E_{inc} < E_K$) at an angle of 130°. The present measurements rule out the possibility of a strong angular dependence of differential cross sections for various L_3 subshell X-rays following selective photoionization of the L_3 subshell. Tartari et al., (2003) investigated the anisotropy of L X-ray fluorescence induced by 59.54 keV unpolarized photons by means of an experimental procedure which allows the relative L X-ray production cross section to be evaluated without taking account of the angular set-up and the instrumental efficiency. Thick targets of Yb, Hf, Ta, W and Pb are considered, and the angular trend of the relative experimental ratios, $I_{L\alpha}/I_{L\beta}$, is calculated by simple evaluations of the peak area alone. Within the experimental uncertainties, which were found to be of the order of 1.6% in the worst cases, the results do not show any significant angular dependence of the $L\alpha$ emission lines. Santra et al., (2007) measured the angular distribution of the L X-ray fluorescent lines from Au and U induced by 22.6-keV X-rays in the angular range of 70°-150°. No strong anisotropy was observed as mentioned by some groups. In the case of Au, a maximum anisotropy of 5% was observed while for U it was within experimental errors 2%. From the angular distribution of the L_i line of Au, the alignment parameter was obtained and its value was found to be 0.10 ± 0.14 . Kumar et al., (2008) investigated alignment of the $M_3(J=3/2)$, $M_4(J=3/2)$ and $M_5(J=5/2)$ subshell vacancy states produced following photoionization in the M_i ($i=1-5$) subshells of Au, Bi, Th and U through angular distribution of the subsequently emitted M X-rays. The unpolarized Mn K X rays ($E_{KX}=5.97$ keV) from the ^{55}Fe radioisotope were used to ionize the M_i subshells in an angular range 90°-160° and the emitted M X-rays were measured under vacuum using a low energy Ge detector. The M X-ray spectra taken at different emission angles were normalized using the isotropically emitted K shell ($J=1/2$) X-rays measured simultaneously from a ^{23}V thin target placed adjoining the M X-ray target. The present precision measurements infer that anisotropy in the $M_{\alpha\beta\gamma}$ X-ray emission shows trends and order of magnitude predicted by theoretical calculations, i.e., anisotropy parameter (β_2) ~ 0.01 .

In the recent experimental study (Han et al., 2008), the angular distribution of characteristic K and L X-rays, emitted from Sm, Eu, Gd Tb, Dy, Ho, and Er as a result of K and L shell vacancies produced by 59.54 keV photon impact was investigated. Thus, K and L X-rays emitted from these elements were simultaneously measured in the same experimental geometry. In this study, Sm, Eu, Gd, Tb, Dy, Ho, and Er lanthanides were chosen since both K shell and L shell electrons of these elements can be excited simultaneously by an Am-241 point source. Also, K and L peaks of the chosen elements are well resolved. Earlier experimental investigations have been only performed on the K X-ray cross sections or on the angular distribution of L X-rays. This is the first report of the angular distributions of L_i X-ray and K_i X-ray ($i = \alpha, \beta$) cross sections for Sm, Eu, Gd, Tb, Dy, Ho, and Er at different angles. It is well known that K X-ray cross sections have no angular dependency ($J = 1/2$). The experimental investigation on K X-ray cross sections at different angles was made to

check the validity of the angular dependency of experimental L X-ray cross sections. The experimental K X-ray cross sections were compared with theoretically calculated values and fairly good correspondence was observed. This means that the present measurements regarding angular dependency of L X-rays are reliable.

In following the work of us (Han et al., 2009) experimental results of the angular distribution of characteristic X-rays were introduced. We preferred to use of $I_{La} / I_{Li}(\theta)$ intensity ratios to obtain the values of alignment parameters (A_2). In that case, the background subtraction problem is considerably reduced and statistical errors are significantly less. It was observed from measured intensities that La and Ll X-ray intensities for the L_3 sub-state depended on the emission angle, meaning that La and Ll X-rays had an anisotropic spatial distribution. Thus, the La to Ll intensity ratios for a set of elements was determined and alignment parameters for each element were obtained using these ratios. In this study, three L sub-shells electrons were excited. Therefore, alignment parameter values are influenced by Coster-Kronig transitions from vacancies induced in the L_1 or L_2 sub-shells. L_1 and L_2 sub-shells have the same $J = 1/2$ value therefore the transferred vacancies are not-aligned and the observed anisotropy of the X-rays is attenuated. For this reason, corrected value of the alignment parameter was calculated using attenuation factor F . If photon energies exciting only L_3 sub-shell electrons are chosen, the alignment parameter will be independent from Coster-Kronig transitions

In more recently study (Han and Demir, 2011a), we investigated the angular distribution of characteristic L X-rays emitted from heavy elements (Pt, Au, Pb, Bi, Th and U) as a result of L shell vacancy production by 59.54 keV photon impact and angular distribution of Compton scattering photons from the same elements. Thus, emitted fluorescent L X-rays and Compton scattering photons from elements were simultaneously measured in the same experimental geometry. Earlier experimental investigations have been only performed on the angular distribution of L X-rays or Compton scattering photons. This is the first report of the angular distribution of Li ($i = l, a, \beta$ and γ) X-rays fluorescent and Compton scattering differential cross sections for Pt, Au, Pb, Bi, Th and U at different angles in the same experimental geometry. It is well known that Compton scattering differential cross sections have angular distribution. The experimental investigation on Compton scattering differential cross sections at different angles was made to check the validity of angular distribution of experimental L X-rays fluorescent differential cross sections. The experimental Compton scattering differential cross sections were compared with theoretically calculated values and fairly good correspondence was observed. This means that the present measurements regarding angular distribution of L X-rays are reliable. In the meantime, L_3 -subshell alignment of Th and U ionized by 59.5 keV photons has been investigated by evaluating the angular dependence of Li ($i = l, a, \eta, \beta$ and γ) X-ray lines. The angular dependence measurements were performed by measuring the fluorescence cross section, σ_{Li} ($i = l, a, \eta, \beta$ and γ) and $\sigma_{Li} / \sigma_{L\gamma}$, $\sigma_{L\eta} / \sigma_{L\gamma}$, $\sigma_{La} / \sigma_{L\gamma}$ and $\sigma_{L\beta} / \sigma_{L\gamma}$ ratios at different angles. It was observed from the measurements that Li ($i = l$ and a) X rays for the L_3 -subshell depended on the emission angle and had an anisotropic spatial distribution. On the other hand, there was no dependence of emission angle and any significant anisotropy for other L X rays. The both Ll and La X-rays originate from the filling of vacancies in states L_3 -subshell with $J = 3/2$. The results of measurements indicate that the L_3 -subshell vacancy states with $J = 3/2$ are aligned, whereas L_1 , and L_2 vacancy states with $J = 1/2$ are non-aligned. Integral cross-sections for the Li ($i = l, a, \eta, \beta$ and γ) X-rays and L subshell fluorescence yields ω_i ($i = 1, 2$ and 3) were also determined and results were compared with theoretically calculated

values and results of others and fairly good correspondence was observed. The $L\gamma$ X-rays, originating purely from the L_1 and L_2 subshells, having isotropic emission were used to normalize the intensities of the anisotropic Ll and the La X-rays originating from the L_3 subshell. It was observed from measurements that Ll and La X-ray for the L_3 sub-state depended on the emission angle, meaning that Ll and La X-rays had an anisotropic spatial distribution. On the other hand, the $L\beta$ and $L\gamma$ X-rays don't show any significant anisotropy. The fluorescence cross sections for Ll and La X-rays are decreased with increased emission angles (Han and Demir, 2011b).

3. Conclusion

In the light of all these, above; data from different researchers show contradictory and the existing results on the angular dependence of fluorescence X-ray and the alignment of atoms with inner-shells vacancy following ionization are still controversial and quite confusing. Therefore, more experimental and theoretical investigations should be required to settle the present discrepancies

4. Acknowledgment

I thank to M.R. Kacal for his help and advice during the preparation of this chapter.

5. References

- Apaydin, G., Tirasoglu, E., Sogut, O., 2008. Measurement of angular dependence of M X-ray production cross-sections in Re, Bi and U at 5.96 keV Eur. Phys. J. D 46, 487-492
- Berezhko, E.G., Kabachnik, N.M., 1977. Theoretical study of inner-shell alignment of atoms in electron impact ionisation: angular distribution and polarization of X-rays and Auger electrons. J. Phys. B At. Mol. Opt. Phys. 10, 2467-2477.
- Bhalla, C.P., 1990. Angular distribution of Auger electrons and photons in resonant transfer and excitation in collisions of ions with light targets. Phys. Rev. Lett. 64, 1103-1106.
- Caldwell, C.D., Zare, R.N., 1977. Alignment of Cd atoms by photoionization. Phys. Rev. A 16, 255-262.
- Cooper, J., Zare, N., 1969. Photoelectron angular distributions. In: Geltman, S., Mahanthappa, K.T., Brittin, W.E. (Eds.), Lectures in Theoretical Physics: Atomic Collision Processes, vol. XI C. Gordon and Breach, New York, pp. 317-337.
- Demir, L., Şahin, M., Kurucu, Y., Karabulut, A., Şahin, Y., 2000. Measurement of angular dependence of photon-induced differential cross-sections of M X-rays from Pt, Au and Hg at 5.96 keV. Radiat. Phys. Chem. 59 355-359
- Demir, L., Şahin, M., Kurucu, Y., Karabulut, A., Şahin, Y., 2003. Angular dependence of Ll , La , $L\beta$ and $L\gamma$ X-ray differential and fluorescence cross-sections for Er, Ta, W, Au, Hg and Tl Radiat. Phys. Chem. 67, 605-612
- Döbelin, E., Sandner, W., Mehlhorn, W., 1974. Experimental study of inner shell alignment of atoms in electron impact ionization. Phys. Lett. A 49, 7-8.
- Durak, R., 2006. Measurement of angular dependence of M X-ray production differential cross-sections in heavy elements at 5.96 keV. Can. J. Anal. Sci. Spectrosc. 51, No. 2.

- Ertuğrul, M., Büyükkasap, E., Erdoğan, H., 1996a. Experimental investigation of the angular dependence of photon-induced differential cross-sections of L X-rays from U, Th and Bi at 59.5 keV. *Il Nuovo Cimento D*, 18, 671–676.
- Ertuğrul, M., Büyükkasap, E., Küçükönder, A., Kopya, A.İ., Erdoğan, H., 1995. Anisotropy of L -shell X-rays in Au and Hg excited by 59.5 keV photons. *Il Nuovo Cimento D*, 17, 993–998.
- Ertuğrul, M., Öz, E., Şahin, Y., 2002. Measurement of alignment parameters for photon induced L_3 vacancies in the elements $59 \leq Z \leq 92$. *Physica Scripta* 66, 289–292.
- Ertuğrul, M., 1996b. Measurement of cross-sections and Coster–Kronig transition effect on L sub-shell X-rays of some heavy elements in the atomic range $79 \leq Z \leq 92$ at 59.5 keV. *Nucl. Instrum. Methods Phys. Res. B* 119, 345–351.
- Flügge, S., Mehlhorn, W., Schmidt, V., 1972. Angular distribution of auger electrons following photoionization. *Phys. Rev. Lett.* 29, 7–9 Erratum: *Phys. Rev. Lett.* 29, 1288.
- Han, I., Demir, L., 2011a. Angular distribution of fluorescent L X-rays and Compton scattering photons *Spectrosc. Lett.* 44, 95–102.
- Han, I., Demir, L., 2011b. Angular dependence of L_3 -subshell X-ray emission following photoionisation. *J X-Ray Sci. Technol.* 19, 13–21.
- Han, I., Sahin, M., Demir, L., 2008. Angular variations of K and L X-ray fluorescence cross sections for some lanthanides *Can. J. Phys.* 86, 361–367.
- Han, I., Sahin, M., Demir, L., 2009. The polarization of X-rays and magnetic photoionization cross-sections for L_3 sub-shell. *Appl. Radiat. Isot.* 67, 1027–1032.
- Hardy, J., Henins, A., Bearden, J.A., 1970. Polarization of the L_{a1} X-rays of mercury. *Phys. Rev. A* 2, 1708–1710.
- Jamison, K.A., Richard, P., 1977. Polarization of target K X-rays. *Phys. Rev. Lett.* 38, 484–487.
- Jesus, A.P., Ribeiro, J.P., Niza, I.B., Lopes, J.S., 1989. L_3 -subshell alignment of Au induced by proton, deuteron and alpha-particle impact. *J. Phys. B At. Mol. Opt. Phys.* 22, 65.
- Jitschin, W., Hippler, R., Shanker, R., Kleinpoppen, H., Schuch, R., Lutz, H.O., 1983. L X-ray anisotropy and L_3 -sub-shell alignment of heavy atoms induced by ion impact. *J. Phys. B At. Mol. Opt. Phys.* 16, 1417–1431.
- Jitschin, W., Kleinpoppen, H., Hippler, R., Lutz, H.O., 1979. L-shell alignment of heavy atoms induced by proton impact ionization. *J. Phys. B At. Mol. Opt. Phys.* 12, 4077–4084.
- Kahlon, K.S., Aulakh, H.S., Singh, N., Mittal, R., Allawadhi, K.L., Sood, B.S., 1990a. Experimental investigation of alignment of the L_3 sub-shell vacancy state produced after photoionization in lead by 59.57 keV photons. *J. Phys. B At. Mol. Opt. Phys.* 23, 2733–2743.
- Kahlon, K.S., Aulakh, H.S., Singh, N., Mittal, R., Allawadhi, K.L., Sood, B.S., 1991a. Measurement of angular distribution and polarization of photon-induced fluorescent x rays in thorium and uranium. *Phys. Rev. A* 43, 1455–1460.
- Kahlon, K.S., Shatendra, K., Allawadhi, K.L., Sood, B.S., 1990b. Experimental investigation of angular dependence of photon induced L shell X-ray emission intensity. *Pramana* 35, 105–114.
- Kahlon, K.S., Singh, N., Mittal, R., Allawadhi, K.L., Sood, B.S., 1991b. L_3 -sub-shell vacancy state alignment in photon–atom collisions. *Phys. Rev. A* 44, 4379–4385.

- Kamiya, M., Kinefuchi, Y., Endo, H., Kuwako, A., Ishii, K., Morita, S., 1979. Projectile- energy dependence of intensity ratio of La to Ll X-rays produced by proton and ^3He impacts on Ho and Sm. *Phys. Rev. A* 20, 1820–1827.
- Kumar, A., Garg, M.L., Puri, S., Mehta, D., Singh, N., 2001a. Angular dependence of L_3 x-ray emission following L_3 sub-shell photoionization in Pb. *X-Ray Spectrom.* 30, 287–291
- Kumar, A., Puri, S., Mehta, D., Garg, M.L., Singh, N., 1999. Angular dependence of L x-ray emission in Pb following photoionization at 22.6 and 59.5 keV. *J. Phys. B At. Mol. Opt. Phys.* 32, 3701–3709.
- Kumar, A., Puri, S., Shahi, J.S., Garg, M.L., Mehta, D., Singh, N., 2001b. L X-ray production cross-sections in Th and U at 17.8, 25.8 and 46.9 keV photon energies. *J. Phys. B At. Mol. Opt. Phys.* 34, 613–623.
- Kumar, S., Sharma, V., Mehta, D., Singh, N., 2008. Alignment of M_i ($i=3-5$) subshell vacancy states in ^{79}Au , ^{83}Bi , ^{90}Th , and ^{92}U following photoionization by unpolarized Mn K x rays. *Phys. Rev. A* 77, 032510
- Küst, H., Kleiman, U., Mehlhorn, W., 2003. Alignment after Xe L_3 photoionization by synchrotron radiation. *J. Phys. B At. Mol. Opt. Phys.* 36, 2073.
- Mc Farlane, S.C., 1972. The polarization of characteristic X radiation excited by electron impact. *J. Phys. B At. Mol. Opt. Phys.* 5, 1906–1915.
- Mehlhorn, W., 1968. On the polarization of characteristic X radiation. *Phys. Lett. A* 26, 166–167.
- Mehlhorn, W., 1994. Alignment after inner-shell ionization by electron impact near and at threshold. *Nucl. Instrum. Methods Phys. Res. B* 8, 7227–7233.
- Mehta, D., Puri, S., Singh, N., Garg, M.L., Trehan, P.N., 1999. Angular dependence of L X-ray production cross-sections in U at 22.6 and 59.5 keV photon energies. *Phys. Rev. A* 59, 2723–2731.
- Mitra, D., Sarkar, M., Bhattacharya, D., Chatterjee, M.B., Sen, P., Kuri, G., Mahapatra, D.P., Lapicki, G., 1996. L_3 -subshell alignment in gold and bismuth induced by low-velocity carbon ions. *Phys. Rev. A* 53, 2309–2313.
- Ozdemir, Y., Durak R., 2008. Angular dependence from L_3 -subshell to M-shell vacancy transfer probabilities for heavy elements using EDXRF technique. *Annals of Nuclear Energy* 35 1335–1339.
- Ozdemir, Y., Durak R., Esmer, K., Ertugrul, M., 2005. Measurement of angular dependence from L_3 -subshell to M-shell vacancy transfer probabilities for the elements in the atomic region $71 \leq Z \leq 78$. *J Quant. Spectrosc. Radiat. Transf.* 90 161–168.
- Pálincás, J., Sarkadi, L., Schlenk, B., Török, I., Kálmán, Gy., 1982. L_3 -subshell alignment of gold by C^+ and N^+ impact ionisation. *J. Phys. B At. Mol. Opt. Phys.* 15, L451.
- Pálincás, J., Schlenk, B., Valek, A., 1979. Experimental investigation of the angular distribution of characteristic X-radiation following electron impact ionisation. *J. Phys. B At. Mol. Opt. Phys.* 12, 3273.
- Pálincás, J., Schlenk, B., Valek, A., 1981. The Coulomb deflection effect on the L_3 -sub-shell alignment in low-velocity proton impact ionization. *J. Phys. B At. Mol. Opt. Phys.* 14, 1157–1159.
- Papp, T., 1999. On the angular distribution of X-rays of multiply ionized atoms. *Nucl. Instrum. Methods Phys. Res. B* 154, 300–306.
- Papp, T., Campbell, J.L., 1992. Non-statistical population of magnetic substates of the erbium L_3 subshell in photoionization. *J. Phys. B At. Mol. Opt. Phys.* 25, 3765.

- Puri, S., Mehta, D., Shahid, J.S., Garg, M.L., Singh, N., Trehan, P.N., 1999. Photon-induced L X-ray production differential cross sections in thorium at 22.6 keV. *Nucl. Instrum. Methods Phys. Res. B* 152 19-26
- Santra, S., Mitra, D., Sarkar, M., Bhattacharya, D., 2007. Angular distribution of Au and U L x rays induced by 22.6-keV photons. *Phys. Rev. A* 75, 022901.
- Schöler, A., Bell, F., 1978. Angular distribution and polarization fraction of characteristic X-radiation after proton impact. *Z. Phys. A* 286, 163–168.
- Scofield, J.H., 1976. Angular dependence of fluorescent X-rays. *Phys. Rev. A* 14, 1418–1420.
- Scofield, J.H., 1989. Angular and polarization correlations in photoionization and radiative recombination. *Phys. Rev. A* 40, 3054–3060.
- Seven, S., 2004. Measurement of angular distribution of fluorescent X-rays and L subshell fluorescence yields in thorium and uranium. *Radiat. Phys. Chem.* 69, 451–460
- Seven, S., Koçak, K., 2001. Angular dependence of L x-ray production cross sections in seven elements from Yb to Pt at a photon energy of 59.5 keV. *J. Phys. B At. Mol. Opt. Phys.* 34, 202.
- Seven, S., Koçak, K., 2002. Angular dependence of L x-ray production cross-section in seven elements from Au to U at 59.5 keV photon energy. *X-Ray Spectrom.* 31, 75–83.
- Sharma, J.K., Allawadhi, K.L., 1999. Angular distribution of $L\beta$ X-rays from decay of L_3 subshell vacancies in uranium and thorium following photoionization. *J. Phys. B At. Mol. Opt. Phys.* 32, 2343–2349.
- Sizov, V.V., and Kabachnik, N.M., 1980. Inner-shell alignment of atoms in ion-atom collisions. I. Impact ionisation. *J. Phys. B At. Mol. Opt. Phys.* 13, 1601.
- Sizov, V.V., and Kabachnik, N.M., 1983. Inner-shell alignment of atoms in ion-atom collisions. III. Light target atoms. *J. Phys. B At. Mol. Opt. Phys.* 16, 1565.
- Stachura, Z., Bosch, F., Hambsch, F.J., Liu, B., Maor, D., Mokler, P.H., Schonfeldt, W.A., Wahl, H., Cleff, B., Brüssermann, M., Wigger, J., 1984. Anisotropy of $L\alpha$ X-ray transition observed in 1.4 MeV N^{-1} heavy ion-atom collisions. *J. Phys. B At. Mol. Opt. Phys.* 17, 835–847.
- Tartari, A., Baraldi, C., Casnati, E., Re, A.D., Fernandez, J.E., Simone, T., 2003. On the angular dependence of L x-ray production cross sections following photoionization at an energy of 59.54 keV. *J. Phys. B: At. Mol. Opt. Phys.* 36 843–851.
- Wigger, J., Altevogt, H., Brüssermann, M., Richter, G., Cleff, B., 1984. M_3 , M_4 and M_5 alignment of thorium by proton impact ionisation. *J. Phys. B At. Mol. Opt. Phys.* 17, 4721.

Radioisotopes - Applications in Physical Sciences

Edited by Prof. Nirmal Singh

ISBN 978-953-307-510-5

Hard cover, 496 pages

Publisher InTech

Published online 19, October, 2011

Published in print edition October, 2011

The book Radioisotopes - Applications in Physical Sciences is divided into three sections namely: Radioisotopes and Some Physical Aspects, Radioisotopes in Environment and Radioisotopes in Power System Space Applications. Section I contains nine chapters on radioisotopes and production and their various applications in some physical and chemical processes. In Section II, ten chapters on the applications of radioisotopes in environment have been added. The interesting articles related to soil, water, environmental dosimetry/tracer and composition analyzer etc. are worth reading. Section III has three chapters on the use of radioisotopes in power systems which generate electrical power by converting heat released from the nuclear decay of radioactive isotopes. The system has to be flown in space for space exploration and radioisotopes can be a good alternative for heat-to-electrical energy conversion. The reader will very much benefit from the chapters presented in this section.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Ibrahim Han (2011). Angular Dependence of Fluorescence X-Rays and Alignment of Vacancy State Induced by Radioisotopes, Radioisotopes - Applications in Physical Sciences, Prof. Nirmal Singh (Ed.), ISBN: 978-953-307-510-5, InTech, Available from: <http://www.intechopen.com/books/radioisotopes-applications-in-physical-sciences/angular-dependence-of-fluorescence-x-rays-and-alignment-of-vacancy-state-induced-by-radioisotopes>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2011 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](https://creativecommons.org/licenses/by/3.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen