

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

185,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



UV Damaged DNA Repair & Tolerance in Plants

Ashwin L. Ganpudi and Dana F. Schroeder
*University of Manitoba,
 Canada*

1. Introduction

In both natural and agricultural conditions, plants are frequently subjected to unfavourable environments, resulting in varying degrees of stress. UV irradiation, drought, heat shock, chilling/freezing, salinity and oxygen deficiency are a few of the major abiotic factors restricting plant growth and development. An important consequence of stratospheric ozone depletion is increased transmission of solar Ultra Violet (UV) radiation through the earth's atmosphere. This increased incidence of UV irradiation causes detrimental effects to all life forms on earth.

1.1 UV irradiation

The spectrum of solar electromagnetic radiation striking the earth's atmosphere ranges from 100 nm to 1 mm. This includes the UV spectrum (100-400 nm), visible spectrum (380-780 nm) and infrared spectrum (700 nm-1 mm). The UV spectrum is further subdivided into three categories: UV-C (100-280 nm), UV-B (280-315 nm), and UV-A (315-400 nm) (Ballaré, 2003). The shortest of these wavelengths, UV-C, is blocked completely by the ozone layer and atmospheric oxygen. In contrast, UV-A is weakly absorbed and directly transmitted to the earth's surface. Wavelengths in the UV-B range are absorbed efficiently though not completely by ozone, as a very small percentage may pass through holes in the ozone. UV-C is extremely harmful, followed by UV-B, while UV-A has milder effects (Batschauer et al., 1999).

1.2 Plants and UV radiation

Plants, due to their non-motile nature, generally have a higher rate of UV tolerance than animals. Plant secondary metabolites aid in defence against both abiotic and biotic stress factors. Plants are capable of reflecting or absorbing harmful UV rays via thick layers of waxy cutin or submerin on the cell walls and intracellular accumulation of chemical substances such as flavanols or phenolics. The biological effects of UV radiation on plants include altered growth responses, reproductive deformities, epigenetic variations, plant susceptibility to biotic factors, premature senescence, damage to the photosynthetic apparatus, and altered conformation of membrane structures. A wide array of genes were found to be induced upon prolonged exposure to low doses of UV-B in the model plant *Arabidopsis thaliana* (Frohnmeier & Staiger, 2003; Mackerness, 2000; Ries et al., 2000). Upregulated transcripts include: antioxidant/free radical scavenging enzymes, proteins involved in: DNA repair, translation, E3 ligase system, cell cycle, signal transduction, and secondary

metabolites, as well as several other genes with unknown function (Brosché et al., 2002; Jansen et al., 2008). UV-B also results in numerous changes in plant morphology. This signalling cascade is well reviewed elsewhere (Jenkins, 2009). Here we focus on plant responses to UV-induced DNA damage.

1.3 UV induced DNA damage

UV-C/B radiation is directly absorbed by DNA, inducing lesions which inhibit vital cellular functions such as transcription and DNA replication. UV-A is comparatively less efficient in lesion induction but triggers the production of reactive oxygen species (ROS) (Kunz et al., 2006). The primary UV induced DNA lesions include cyclobutane pyrimidine dimers (adjacent pyrimidines covalently linked between C-5 and C-6 carbon atoms) and secondary lesions 6-4 pyrimidine-pyrimidone photoproducts (6-4 PP) (covalent linkage between the C-4 position of a pyrimidine to the C-6 position of an adjacent pyrimidine) (Fig. 1). In order to respond to this damage, plants employ specific mechanisms (Britt, 1999). In light conditions, photoreactivation catalyses dimer monomerizations while during dark conditions, Nucleotide Excision Repair (NER) excises these helix-distorting lesions. Finally, residual lesions are bypassed via translesion synthesis (TLS).




Fig. 1. A) Normal adjacent pyrimidine residues. B) UV-induced Cyclobutane Pyrimidine Dimer (CPD) and C) 6-4 Pyrimidine-Pyrimidone photoproduct (6-4 PP).

2. Photoreactivation

Photoreactivation is a blue light dependant DNA repair mechanism catalysed by the photolyase (E.C 4.1.99.3) class of enzymes. Pyrimidine dimers are split by the action of two photoactive damage specific DNA repair enzymes – CPD photolyase and 6-4 PP photolyase. Both classes of photolyase require two co-factors, one being the two electron reduced form of Flavin Adenine Dinucleotide (FAD) and the second chromophore, a blue light harvesting photoantenna, being either 5,10- methenyltetrahydrofolate (MTHF) or 8-hydroxy-7,8-didemethyl-5-deazariboflavin (8-HDF). FAD is an essential co-factor for regulating DNA binding and repair. In contrast, the second chromophore has a higher extinction co-efficient and an absorption maximum at longer wavelengths, hence regulates the rate of repair depending on the external light intensity. MTHF or 8-HDF absorbs the photoreactivating blue light photons and transfers this excitation energy to the reduced FAD. The FADH⁻ in

turn transfers electrons to the lesions, catalyzing the cleavage of the cyclobutane rings and dimer monomerization (Deisenhofer, 2000; Sancar, 2003, 2008). Multiple sequence alignment reveals that conserved homology between prokaryotic and eukaryotic CPD photolyases is limited to the C-terminal FAD binding site. It has been suggested that a common ancestor gave rise to both type I and type II photolyases but diverged at an early evolutionary stage (Yasui et al., 1994). CPD photolyases have been classified into Class I (microbial) and Class II (higher eukaryotes excluding placental mammals) groups, respectively. The 6-4 photolyases from *Drosophila* and *Arabidopsis* have strong sequence similarity to class I CPD photolyases (Nakajima et al., 1998; Todo et al., 1996). Similarly cryptochromes, the plant blue light photoreceptors, are 30% similar to the class I microbial photolyases, but demonstrate no photolyase activity (Ahmad & Cashmore, 1993). Thus, microbial Class I CPD photolyases, eukaryotic 6-4 photolyases, and blue light photoreceptors constitute the class I photolyase/photoreceptor family.

Genes encoding CPD photolyases and 6-4 PP photolyases have been identified and characterized in a range of prokaryotic and eukaryotic systems (Sancar, 2003). In plants genes encoding CPD photolyases have been identified in *Arabidopsis thaliana* (Ahmed et al., 1997), cucumber (Takahashi et al., 2002), rice (Hirouchi et al., 2003), spinach (Yoshihara et al., 2005), and soybean (Yamamoto et al., 2008). Genes encoding 6-4 PP photolyases have been identified in *Arabidopsis* and rice (Chen et al., 1994; Singh et al., 2010). In *Arabidopsis*, the highest levels of both photolyases are associated with floral tissues, which may presumably serve to minimize lesions in germline cells. While expression of the CPD photolyase is light/UV-B regulated, 6-4 PP photolyase is constitutively expressed (Takahashi et al., 2002; Waterworth et al., 2002). The *Arabidopsis* CPD photolyase gene (*AtPHR1*) encodes a class II CPD photolyase. An *Arabidopsis* mutant (*uvr2*) lacking this gene is hypersensitive to UV. *AtPHR1* is efficient in CPD photoreactivation but deficient in 6-4 photoproduct repair (Ahmed et al., 1997; Landry et al., 1997). *AtPHR1* is upregulated several fold in a UV insensitive mutant of *Arabidopsis* (*uvi1*) irrespective of light conditions, conferring constitutive protection (Tanaka et al., 2002). Overexpression of CPD photolyase in *Arabidopsis* and rice resulted in enhanced CPD removal (Hidema et al., 2007; Kaiser et al., 2009; Ueda et al., 2005). Genetic complementation of photolyase deficient *E.coli* strains with soybean, rice, spinach and *Arabidopsis* CPD photolyase genes restored photoreactivation activity (Yamamoto et al., 2007, 2008; Yoshihara et al., 2005). CPD photolyase activity in *Arabidopsis* (Pang & Hays, 1991; Waterworth et al., 2002), soybean (Yamamoto et al., 2008) and rice (Hidema et al., 2000) has been reported to be temperature sensitive. Rice CPD photolyase, encoded by a single copy gene in the nuclear genome, translocates to chloroplasts, mitochondria and nuclei to repair UV-induced CPDs in all three genomes (Takahashi et al., 2011), a phenomenon not observed in spinach chloroplasts (Hada et al., 2000) or young *Arabidopsis* seedlings (Chen et al., 1996). However, upon exposure to photoreactivating blue light, *Arabidopsis* seedlings did exhibit efficient repair of CPDs in the extracellular organelles (Draper & Hays, 2000). The *Arabidopsis* 6-4 PP photolyase, encoded by the *UVR3* gene, encodes a 62 kDa protein with 45% identity to *Drosophila* 6-4 PP photolyase and 17% identity to the Class II CPD photolyases. *AtUVR3* is proficient in 6-4 photoproduct removal but deficient in CPD repair. Both *uvr2* and *uvr3* are nonsense mutations, and the double mutants are extremely sensitive to UV relative to the single mutants (Nakajima et al., 1998). Photolyases appear to be the sole repair mechanism active in non-proliferating plant tissues (Kimura et al., 2004). Hence, photolyases play an important role in plant repair of UV damaged DNA.

3. Nucleotide excision repair

Nucleotide excision repair (NER) is a light independent repair process involving a series of reactions: initial damaged DNA recognition, DNA unwinding, dual incision bracketing the lesion, repair synthesis and final ligation to seal the repaired site. NER initiates with specific sub-pathways for transcriptionally active (Transcription Coupled Repair (TC-NER)) or silent (Global Genomic Repair (GG-NER)) DNA. TC-NER and GG-NER exhibit different damage recognition strategies followed by a common repair pathway (Gillet & Scharer, 2006) (Fig. 2). Defects in human NER genes result in the photosensitive syndromes Xeroderma pigmentosum (XP) or Cockayne syndrome (CS). Eight genetic complementation groups for XP have been identified (XPA-G, V) as well as two for CS (CSA and CSB). While XP-V mutation uniquely results in defects in translesion synthesis, XP -A, -B, -D, -F, and -G mutation results in both TC-NER and GG-NER defects, while XP -C and -E mutation results in GG-NER defects only. CSA and CSB mutation results exclusively in TC-NER defects (Hoeijmakers, 2001; Svejstrup, 2002). Bioinformatic analysis of the plant NER protein machinery suggests the molecular mechanisms are largely but not entirely conserved with that of the extensively studied yeast *S. cerevisiae* and mammalian cells (Kimura & Sakaguchi, 2006; Kunz et al., 2005, 2006). NER in plants has been studied primarily in rice and *Arabidopsis* (Singh et al., 2010). Many *Arabidopsis* NER related genes were initially isolated by analysis of UV hypersensitive (*uvh*) and UV repair defective (*uvr*) mutants which were subsequently mapped to homologues of the human XP genes (Table 1).

3.1 Global genomic repair

3.1.1 Damage recognition

3.1.1.1 DDB1 & DDB2

In mammalian systems, damage detection in the GG-NER pathway involves UV-Damaged DNA Binding protein 1 and 2 (DDB1 and DDB2) followed by the XPC-HR23B-CEN2 complex. In humans DDB2 complements the XPE mutation and plays a role in recognition of the UV-induced lesion, while DDB1 is required for high affinity interaction of the DDB1-DDB2 complex (Groisman et al., 2003; Luijsterburg et al., 2007; Rapic-Otrin et al., 2002). *S. pombe* Ddb1 knockouts result in chromosomal segregation defects, UV sensitivity and slow S phase progression leading to defects in meiosis (Holmberg et al., 2005). DDB1 and DDB2 homologues have been identified in rice, where they are UV-induced in proliferating tissues (Ishibashi et al., 2003). *Arabidopsis thaliana* encodes two homologs of DDB1 – DDB1A and DDB1B. These proteins are 91% identical with redundant function. Although *ddb1b* null alleles appear lethal, a viable partial loss of function allele exhibits no developmental or UV sensitive phenotypes (Bernhardt et al., 2010; Schroeder et al., 2002). Overexpression of DDB1A in *Arabidopsis* confers enhanced UV resistance and *ddb1a* knockouts exhibit mild UV sensitivity (Al Khateeb & Schroeder, 2009; Molinier et al., 2008). *AtDDB2* encodes a 48 kDa nuclear localized protein with upregulated expression upon UV-induction. *AtDDB2* loss of function results in UV sensitivity while overexpression increases UV tolerance (Biedermann & Hellmann, 2010; Koga et al., 2006; Molinier et al., 2008).

3.1.1.2 Cullin based E3 ligases

The 127 kDa DDB1 homologs function as substrate adaptors for CULLIN4 based E3 ubiquitin (Ub) ligases (Groisman et al., 2003). E3 Ub ligases are multimeric complexes that add ubiquitin moieties to target proteins and contain CULLIN proteins as scaffolding




Fig. 2. Overview of mammalian nucleotide excision repair. In GG-NER, DDB2-CUL4 mediated histone (H) and XPC ubiquitination facilitates lesion binding. In TC-NER, stalled RNA POL II recruits CSB and the CSA-CUL4-CSN complex, followed by recruitment of other TC-NER specific factors. In both cases, NER core players follow suit: XPB and XPD helicases of the TFIIH complex, XPF-ERCC1 and XPG endonucleases, and the ssDNA binding XPA-RPA complex. The fragment encompassing the lesion is excised, followed by repair synthesis and ligation. Repair synthesis requires DNA POL δ/ϵ in concert with accessory proteins PCNA, RFC and RPA. See text for details.

subunits (Hua & Vierstra, 2011). CUL4 based E3 ubiquitin ligases consist of three core subunits: CULLIN4 (CUL4), RING finger protein REGULATOR OF CULLINS1 (ROC1)/RING-BOX1 (RBX1), and DDB1. The CUL4 – RBX1 – DDB1 complex interacts with a large number of proteins containing WD40 motifs referred to as DCAF proteins (DDB1-CUL4 Associated Factor) or DWD proteins (DDB1 binding WD40 proteins) (Lee & Zhou, 2007). DDB2 is an example of a WD40 domain containing DCAF protein. WD40 motifs are characterized by 40 amino acid repeats initiated by a glycine-histidine dipeptide and terminated by a tryptophan-aspartate (WD) dipeptide facilitating protein-protein interactions. DDB1 is composed of three β propeller domains (BPA, BPB and BPC) and DDB2, in addition to the WD40 domain, contains a helix loop helix (HLX) segment in the N terminal. While the clam shaped BPA-BPC of DDB1 mediates interaction with the HLX motif of DDB2 and other DCAF substrates, BPB exhibits exclusive interactions with CUL4 (Scriama et al., 2008).

AtCUL4 is a 91 kDa protein with a conserved CH motif and an extended N terminal region of 65 amino acids that shares close sequence similarity to its human/mouse orthologs. *AtCUL4* loss of function results in abnormal plant development (Bernhardt et al., 2006; Chen et al., 2006) and UV sensitivity (Molinier et al., 2008). Examples of DCAF proteins interacting with the *Arabidopsis* CUL4-DDB1A/B complex include AtDDB2 (Bernhardt et al., 2006), AtCSA-1&2 (Biedermann & Hellmann, 2010; Zhang et al., 2010), as well as the negative regulator of photomorphogenesis DET1 (De-etiolated1) (Schroeder et al., 2002), and many other DWD proteins (Lee et al., 2008). Recent results have shed light on the cross talk between photomorphogenesis regulation and repair of UV damaged DNA. HY5, a positive regulator of photomorphogenesis, has been shown to regulate gene sets connected to UV tolerance, such as the *UVR3* and *PHR1* photolyases, as well as secondary metabolite transcriptional regulators (Oravec et al., 2006; Ulm et al., 2004). DET1, initially identified as a nuclear localized negative regulator of photomorphogenesis, exhibits a constitutively light grown phenotype in addition to increased levels of flavanoids (Pepper et al., 1994). Recent papers show that *det1* mutants exhibit enhanced UV tolerance through increased levels of secondary metabolites reflecting/absorbing UV irradiation as well as by upregulation of photolyase genes. Further it appears that DET1 protein dosage influences UV sensitivity of plants as DET1 overexpressing lines exhibit increased UV sensitivity (Castells et al., 2010, 2011).

3.1.1.3 Histone ubiquitination facilitates NER machinery entry

In mammals, in the absence of UV irradiation, DDB2-DDB1-CUL4-RBX1 (DDB2 complex) was found to be associated with the COP9 Signalosome complex (CSN). CSN shares significant structural homology with the 19S lid of 26S proteasome. The CSN deconjugates neddylation (Nedd8) from CULLINs, thereby regulating the activation, stability or the disassembly of CULLIN based E3 ligase activity (Parry & Estelle, 2004; Schwechheimer & Deng, 2001). The DDB2 - CSN complex show no ubiquitin ligase activity, but upon UV irradiation, these complexes disassociate in parallel with increased neddylation and activation of CUL4 (Groisman et al., 2003). Core histone proteins have been identified as potential targets for DDB2-DDB1-CUL4-RBX1 mediated proteasomal degradation. Kapetanaki et al. (2006) and Wang et al. (2006) describe the ubiquitination of H2A, H3 and H4 histone proteins. Reduction of histone H3 and H4 ubiquitination by knockdown of *cul4* impairs recruitment of the repair protein XPC to the damaged foci and inhibits the repair process. Thus biochemical studies indicate that DDB-CUL4-RBX1-mediated histone

Human	Yeast	Function	ATG no.	Arabidopsis
Photoreactivation:				
ND	ND	6-4 PP Photolyase	At3g15620	UVR3
ND	PHR1	Class II CPD Photolyase	At1g12370	PHR1/UVR2
Nucleotide Excision Repair:				
DDB1	ND	Substrate adaptor for CUL4. Interacts with DCAF proteins	At4g05420 At4g21100	AtDDB1a AtDDB1b
DDB2/XPE	ND	Damaged DNA binding (DCAF)	At5g58760	AtDDB2
CUL4	CUL4	Scaffolding subunit of E3 Ub ligase	At5g46210	AtCUL4
XPC	RAD4	GG-NER damage recognition	At5g16630	AtXPC
HR23B	RAD23	Binds to XPC	At1g79650 At1g16190 At3g02540 At5g38470	AtRAD23A AtRAD23B AtRAD23C AtRAD23D
CEN2	CEN2	Stabilizes XPC-HR23B complex	At3g50360	AtCEN2
XPB	RAD25	Subunit of TFIIH. 3'->5' helicase	At5g41370 At5g41360	AtXPB1 AtXPB2
XPD	RAD3	Subunit of TFIIH. 5'->3' helicase	At1g03190	AtXPD/UVH6
GTF2H1	TFB1	Core TFIIH subunits	At1g55750	AtTFB1-1
GTF2H2	SSL1		At3g61420	AtTFB1-2
GTF2H3	TFB4		At1g05055	Atp44
GTF2H4	TFB2		At1g18340	AtTFB4
GTF2H5	TFB5		At4g17020	AtTFB2
XPA	RAD14	ssDNA binding	ND	ND
XPG	RAD2	3' endonuclease	At3g28030	AtXPG/UVH3
ERCC1	RAD10	5' endonuclease with XPF	At3g05210	AtERCC1/UVR7
XPF	RAD1	5' endonuclease with ERCC1	At5g41150	AtXPF/UVH1
PCNA	PCNA	RFC dependant sliding clamp	At1g07370 At2g29570	AtPCNA1 AtPCNA2
RFC1	RFC1	DNA-dependent ATPase required for DNA replication and repair	At5g22010	AtRFC1
RFC2	RFC2		At1g21690	AtRFC2
RFC3	RFC3		At1g77470	AtRFC3
RFC4	RFC4		At1g63160	AtRFC4
RFC5	RFC5		At5g27740	AtRFC5
RPA70	RFA1	ssDNA binding protein required for architectural role in dual lesion incision and repair synthesis	At2g06510	AtRPA70A
			At4g19130	AtRPA70B
			At5g08020	AtRPA70C
			At5g45400	AtRPA70D
			At5g61000	AtRPA70E
RPA32	RFA2		At2g24490	AtRPA32A
			At3g02920	AtRPA32B
RPA14	RFA3		At3g52630	AtRPA14A
			At4g18590	AtRPA14B
CSA	RAD28	TC-NER specific DCAF protein	At1g27840 At1g19750	AtCSA1A AtCSA1B

Human	Yeast	Function	ATG no.	Arabidopsis
CSB	RAD26	SWI/SNF2 like ATPase	At2g18760	AtCSB
XAB2	SYF1	Complex stabilization	At5g28740	AtXAB2
TFIIS	TFIIS	TC-NER elongation factor	At2g38560	AtTFIIS
HMGN1	ND	Nucleosome binding	ND	ND
EP300/CBP	ND	Histone acetyl transferase	At1g79000 At1g67220 At1g55970 At3g12980 At1g16710	HAC1 HAC2 HAC4 HAC5 HAC12
Translesion Synthesis				
POLH/XPV	RAD30	Y-family DNA polymerase	At5g44740	AtPOLH
REV1	REV1	Y-family DNA polymerase	At5g44750	AtREV1
POLZ	REV3	B-family DNA polymerase	At1g67500	AtREV3
REV7	REV7	REV3 accessory subunit	At1g16590	AtREV7
POLK	ND	Y-family DNA polymerase	At1g49980	AtPOLK

Table 1. Genes involved in UV damaged DNA repair and tolerance. ND=not detected.

ubiquitination weakens the interaction between histones and DNA to further facilitate the recruitment of repair proteins to damaged DNA (Guerrero-Santoro et al., 2008; Higa et al., 2006). The activated DDB2 complex recruits XPC via protein-protein interactions, followed by ubiquitination of XPC and DDB2. Polyubiquitinated DDB2 exhibits reduced affinity for damaged DNA and is subsequently displaced from the damaged foci, whereas polyubiquitinated XPC enhances its binding to DNA (Sugasawa et al., 2005). In *Arabidopsis*, DDB2 turnover is abrogated in *cul4*, *ddb1a*, *atr* and *det1* mutants suggesting that ATR and DET1 may co-operate with the CUL4-DDB1 E3 ligase complex in regulating NER (Castells et al., 2011; Molinier et al., 2008).

3.1.1.4 XPC-HR23B-CEN2

The next step in GG-NER involves the homologous heterodimers hXPC-hHR23B (in Humans) and RAD4-RAD23 (in yeast). In addition to hXPC-hHR23B, Araki et al. (2001) identified hCEN2, a Ca²⁺ binding protein contributing to the stability of the hXPC-hHR23B complex. Hence in mammalian systems the identified protein recognition complex is hXPC-hHR23B-hCEN2, however neither hHR23B nor hCEN2 bind to damaged DNA but enhance both the affinity and specificity of hXPC binding to damaged DNA (Fitch et al., 2003; Nishi et al., 2005). AtCEN2 shares 49% identity with hCEN2, *Atcen2* mutants are UV sensitive, and AtCEN2 overexpression resulted in enhanced repair. Upon UV irradiation, AtCEN2 level increases and it rapidly translocates to the nucleus. AtCEN2-AtXPC interaction in *Arabidopsis thaliana* has also been demonstrated (Liang et al., 2006; Molinier et al., 2004). Potential homologs of HR23B/RAD23 have been identified in *Arabidopsis thaliana*, *Oryza sativa* and *Daucus carota* (Schultz & Quatrano, 1997; Sturm & Leinhard, 1998). The *Arabidopsis* genome has 4 loci encoding RAD23 homologs (RAD23a, RAD23b, RAD23c, RAD23d), and although mutants exhibit multiple pleiotrophic developmental defects (Farmer et al., 2010), UV sensitivity or complex interactions with the *Arabidopsis* NER machinery have not yet been reported.

3.1.2 DNA unwinding complex assembly

Following recognition, the damaged region is unwound by the TFIIH transcription factor which joins the XPC-CEN2-HR23B complex. TFIIH is a complex of 10 proteins, seven of which are players in the NER pathway (helicases XPB and XPD, p62, p44, p34, p52, and p8). The last five proteins are encoded by GTF2H1, GTF2H2, GTF2H3, GTF2H4, GTF2H5 (Frit et al., 1999). TFIIH not only participates in NER of transcriptionally active and inactive DNA, but also in RNA POL II dependant transcription, cell cycle control and regulation of nuclear receptor activity (Chen & Suter, 2003). ATP dependant 5'→3' and 3'→5' helicase activities associated with XPD/RAD3 and XPB/RAD25 respectively unwind the DNA encompassing the lesion with the concomitant release of the recognition complex. Human XPB and the corresponding yeast RAD25 knockouts are lethal. *Arabidopsis thaliana*, unlike the sugarcane, rice or mammalian genomes, encodes two homologs of XPB – AtXPB1 and AtXPB2. These proteins are 95% identical with redundant functions and are expressed constitutively throughout plant development (Morgante et al., 2005; Ribeiro et al., 1998). Despite this redundancy, *xpb1* mutants exhibit delayed germination and flowering phenotypes but are not UV sensitive (Costa et al., 2001). Phenotypes of *Arabidopsis xpb2* or double mutants have not yet been reported. *Arabidopsis* XPD is 56% and 50% identical to human and yeast sequences. *Arabidopsis* XPD/RAD3 null mutations are lethal, however viable point mutation alleles are UV sensitive and were identified as *uvh6* (*uv hypersensitive 6*) mutants. (Jenkins et al., 1997; Liu et al., 2003). Another component of the of TFIIH complex, p44, was identified in *Arabidopsis* as ATGTF2H2 and was found to interact in vivo with AtXPD (Vonarx et al., 2006).

3.1.3 Endonuclease recruitment following unwinding

TFIIH further recruits additional factors such as XPA and heterotrimeric Replication Protein A (RPA), composed of 70, 32 and 14 kDa subunits, to promote and stabilize the formation of an open intermediate essential for the dual incision by XPG and XPF-ERCC1 (Excision Repair Complementing defective repair in Chinese hamster 1) (RAD1/RAD10) endonucleases at the 3' and 5' sites respectively (Tapais et al., 2004). The RPA-XPA complex exhibits interactions with both endonucleases (He et al., 1995; Wold, 1997), however the specific function of XPA is still not evident. Initially it was thought to function as a lesion recognition complex in concert with XPC, but was later determined to be recruited after TFIIH entry and facilitate XPC complex departure (Hey et al., 2002; Volker et al., 2001). In addition, XPA homologues do not exist in plants (Kunz et al., 2005). The dual incisions catalyzed by the endonucleases excise oligonucleotides of about 20-30 bases containing the lesion (Reidl et al., 2003).

Potential homologs of ERCC1, XPF, XPG and RPA have been identified in plants. Although ERCC1 was first cloned from male germ line cells of *Lilium longiflorum*, southern blots confirmed the presence of ERCC1 across diverse plant species such as *A. thaliana*, *B. napus*, *Z. mays*, *L. esculentum*, *N. tabacum*, and *O. sativa* (Xu et al., 1998). Hefner et al. (2003) mapped the *Arabidopsis uvr7* mutant to *AtERCC1*. *Atercc1* knockouts are phenotypically normal in contrast to the lethal mammalian counterparts (Weeda et al., 1997). *Atercc1* mutants are extremely sensitive to DNA damaging chemicals such as mitomycin and ionizing agents such as UV and γ – radiation (Hefner et al., 2003). More recent studies in *Arabidopsis* indicate significant roles of *AtERCC1* in concert with *AtXPF* in homologous recombination and chromosomal stability (Dubest et al., 2002, 2004; Vannier et al., 2009). Gallego et al. (2000) and Liu et al. (2000) characterized the single copy AtXPF which is 37% and 29% identical to

human XPF and *S. cerevisiae* RAD1 respectively. *AtXPF* is homogenously expressed, was mapped to the *uvh1* mutant in *Arabidopsis*, and partially complements the yeast *rad1* mutant (Gallego et al., 2000). *AtXPF* point mutations result in sensitivity to ionizing radiation and mitomycin C, and impaired removal of photoproducts (Fidanstef et al., 2000; Vonarx et al., 2002). *AtXPG* was mapped to the *UVH3* locus and knockouts result in UV sensitivity as well as premature senescence and reduced seed production (Liu et al., 2001). The XPG rice homolog, *OsSEND-1*, exhibits upregulated mRNA levels in response to UV and DNA damaging agents (Furukawa et al., 2003a).

ssDNA binding RPA proteins in plants were first identified in rice (Ishibashi et al., 2001). Unlike most eukaryotic organisms, *Arabidopsis* and rice possess multiple copies of the RPA homologs. In addition to participating in DNA repair, RPA proteins play a role in homologous recombination and DNA replication in humans and yeast (Sakaguchi et al., 2009). The rice genome encodes three OsRPA70 (OsRPA70A, OsRPA70B, OsRPA70C), three OsRPA32 (OsRPA32-1, OsRPA32-2, OsRPA32-3) and one OsRPA14. *In vivo* interactions in rice confirms three different complex formations: OsRPA70A-OsRPA32-2-OsRPA14 (Type1); OsRPA70B-OsRPA32-1-OsRPA14 (Type2); and OsRPA70C-OsRPA32-3-OsRPA14 (Type3). Subcellular localization of all three OsRPA32 was detected in both the nucleus and chloroplasts. OsRPA70A was only in the chloroplast whereas OsRPA70B and OsRPA70C were exclusively to the nucleus. This data suggest that while the Type1 complex may participate in chloroplast DNA repair, Type2 and Type3 complexes concentrate on nuclear DNA repair (Ishibashi et al., 2006). OsRPA70A and OsRPA70B share only 33% sequence homology and exhibit differences in expression pattern (Ishibashi et al., 2001). A T-DNA insertion in *OsRPA70A* resulted in partial male sterility, complete female sterility and hypersensitivity to DNA mutagens (Chang et al., 2009). OsRPA32 protein abundance is regulated by both UV irradiation and cell cycle phase (Marwedel et al., 2002). *Arabidopsis*, on the other hand, encodes five putative RPA70 genes and two copies each of RPA32 and RPA14 (Shultz et al., 2007). *Arabidopsis* RPA70A interacts preferentially with AtRPA32A rather than AtRPA32B. Knockouts of both AtRPA70A and AtRPA70B exhibited UV sensitivity when irradiated, but exhibited wildtype characteristics under normal conditions (Ishibashi et al., 2005; Takashi et al., 2009).

3.1.4 Repair synthesis and ligation

In mammals, the gap formed by the excision is filled via PCNA (Proliferating Cell Nuclear Antigen) and RFC dependant DNA synthesis by DNA POL δ/ϵ . These components are likely recruited by XPG and RPA as RFC exhibits interaction with RPA (Yuzhakov et al., 1999). RFC catalyzes the ATP dependant loading of PCNA to DNA at the 3' OH. PCNA is a homotrimeric protein which forms a sliding clamp-like complex (Gulbis et al., 1996) and interacts with the DNA POL to ensure replication occurs processively (Mocquet et al., 2008). The final phase of NER is completed by phosphodiester backbone rejoining of the repaired DNA strand by DNA Ligase I.

Although PCNA and RFC homologues have been identified in plants, their specific role in nucleotide excision repair has not yet been elucidated (Furukawa et al., 2003b; Strzalka & Ziemienowicz, 2011). Recently, Roy et al. (2011) cloned and characterized a homolog of mammalian DNA POL λ in *Arabidopsis*. AtPOL λ was upregulated upon UV induction under dark conditions, and *Atpol λ* mutants exhibited UV sensitivity and decreased DNA repair. Thus, this report suggests a role for DNA POL λ in plant NER.

3.2 Transcription coupled repair

The emerging picture of mammalian TC-NER is of a complex mechanism requiring two essential assembly factors (CSA and CSB), certain TC-NER specific proteins (P300, HMGN1, XAB2 and TFIIS), as well as core NER proteins. UV induced DNA damage is initially recognised by RNA POL II, which stalls when it encounters helix-distorting lesions on the template strand during transcription. Stalled RNA POL II backtracks a few nucleotides and is recognised by the CSB protein which in turn co-ordinates the recruitment of the repair factors required to accomplish Transcription Coupled NER (Mellon, 2005).

Cloning and characterization of the mammalian CSB gene revealed a nuclear protein of 168 kDa with a region of homology to the SWI2/SNF2 family of helicases. CSB has been shown to interact with RNA POL II and this interaction is enhanced and prolonged by UV exposure (van den Boom et al., 2004). Studies propose that functional CSB in the absence of UV could also serve as a component of the transcriptional machinery promoting elongation (Fousteri & Mullenders, 2008; Hanawalt & Spivak, 2008). Further, CSB facilitates the entry of the core NER factors XPA, XPG and TFIIH through complex interactions (Laine & Egly, 2006; Saxowsky & Doetsch, 2006; Svejstrup, 2002). Mammalian CSA on the other hand is a 46 kDa DWD protein containing seven WD40 repeats that associates with DDB1-CUL4 type E3 ligases and is recruited to the damaged site after CSB. CSA physically interacts with the CSB-RNA POL II complex in a UV dependant manner (Groisman et al., 2003; Fousteri et al., 2006). Interestingly, unlike the DDB2 complex, the CSA-CUL4 Ub ligase complex is active under normal conditions but is rapidly inactivated upon UV irradiation by CSN. Hence CSN plays a differential and dynamic role in regulating both pathways of Nucleotide Excision Repair. The stable CSN-CSA-CSB complex is required for the recruitment of the other NER factors. Following repair, CSN dissociates, reactivating CSA Ub ligase activity and resulting in CSB degradation. Clearance of CSB is required for the reinitiation of transcription by RNA POL II (Groisman et al., 2003, 2006).

Several papers over the years propose the fate of RNA POL II during the coupling process: either ubiquitinated and degraded, translocated or bypassed from the lesion site, or simply stalled during the entire repair process, is still a matter of debate (Reviewed in Tornaletti, 2009). XAB2 (XPA binding protein 2) is a RNA-binding protein with 15 tetratricopeptide repeats. In addition to interacting with XPA, XAB2 is capable of interacting with CSA, CSB and RNA POL II (Nakatsu et al., 2000). XAB2 is thought to stabilize protein assemblies by functioning as a scaffolding subunit. XAB2 knockouts in mammalian cells exhibit hypersensitivity and decreased recovery of mRNA synthesis post UV irradiation (Kuraoka et al., 2008). Increased amounts of histone acetyl transferase p300 and High Mobility Group Nucleosome binding domain containing protein 1 (HMGN1) interact with RNA POL II in a CSB-dependant manner upon UV irradiation but exhibit weak interactions under normal conditions (Fousteri et al., 2006). Both HMGN1 and p300 are nucleosome interacting proteins which remodel the chromatin structure behind the arrested polymerase and facilitating the backward translocation of RNA POL II (Hanawalt & Spivak, 2008). TFIIS, functioning as a transcription elongation factor, stimulates the arrested RNA POL II to restart elongation. This TFIIS-RNA POL II interaction is significantly increased upon UV irradiation (Fousteri et al., 2006). Hence it is proposed that TFIIS facilitates resumption of transcription post DNA lesion removal in a CSA/B-dependent manner.

Elucidation of the TC-NER mechanism in plants is still at its infancy. While there is no plant homologue for HMGN1, the *Arabidopsis* genome encodes homologues of XAB2,

TFIIIS, and five p300/CBP homologues, however the role of these genes in DNA repair has not been assessed (Grasser et al., 2009; Kunz et al., 2005; Pandey et al., 2002). Only recently was the homolog of human CSA cloned and characterized in *Arabidopsis*. In contrast to mammalian systems, the *Arabidopsis* genome encodes two homologs of CSA – AtCSA1A and AtCSA1B, 92% identical DWD proteins with overlapping subcellular localization and expression patterns. These proteins exist as heterotetramers in planta and are capable of interacting with the DDB1-CUL4 E3 complex. Knockouts of either gene result in UV sensitivity and decreased photoproduct removal (Zhang et al., 2010). Concurrently, another group overexpressed AtCSA1A, which surprisingly also resulted in increased UV sensitivity. This result is hypothesised to be due to competition between CSA and with other DWD proteins to interact with the DDB1-CUL4 complex. Interestingly AtCSA1A levels remained constant upon UV induction (Biedermann & Hellmann, 2010). RNAi of a putative *Arabidopsis* CSB homolog resulted in a UV sensitive phenotype (Shaked et al., 2006). Hence, taken as a whole, these results confirm the role of the CUL4-DDB1-CSA and CSB pathway in plants.

4. Translesion synthesis

Despite the available repair mechanisms, there are times when UV damage persists and DNA replication proceeds past UV lesions via translesion synthesis (TLS). This synthesis proceeds via one of several DNA polymerases: Pol η , Rev1, Pol ζ , Polk or Pol ι . Pol η (eta) is encoded by the *POLH* locus in humans and is homologous to yeast RAD30. Pol η performs error-free bypass of CPDs, and loss of function results in increased mutation and xeroderma pigmentosa (XP-V) in humans. REV1 exhibits dCMP transferase activity but is required for 6-4 PP bypass independent of this activity, suggesting its role may be in recruitment of other polymerases. Pol ζ is composed of REV3 and REV7 and exhibits error prone repair. REV3 is a B-family polymerase (unlike the other TLS polymerases which are all Y-family polymerases) and REV7 an accessory subunit that enhances REV3 activity (Waters et al., 2009). Polk activity is specific to N2-dG lesions in both animals and plants, while plants do not have a Pol ι homologue (Garcia-Diaz & Bebenek, 2007; Garcia-Ortiz et al., 2007).

Homologues of REV1, REV3 and REV7 as well as Pol η have been identified in *Arabidopsis* (Sakamoto et al., 2003; Santiago et al., 2006; Takahashi et al., 2005). Mutant alleles of *REV3*, *REV1* and *REV7* all result in UV sensitivity, although *rev1* only weakly and *rev7* only to long-term UV exposure (Sakamoto et al., 2003; Takahashi et al., 2005). Interestingly, AtREV1 cannot insert nucleotides across from UV-induced DNA lesions, suggesting that its role is primarily in the recruitment of other TLS polymerases (Takahashi et al., 2007). The UV sensitivity of the *rev1 rev3* double mutant is similar to that of *rev3*, additional evidence that the role of REV1 may be in REV3 recruitment (Takahashi et al., 2005). Analysis of mutation frequency in *rev1* and *rev3* mutants shows a reduction relative to wildtype rate, evidence that these genes contribute to error-prone repair (Nakagawa et al., 2011).

Pol η is encoded by the *Arabidopsis* *POLH* gene. The *POLH* transcript is ubiquitously expressed and alternatively spliced, and AtPol η exhibits similar levels of activity *in vitro* as the human protein (Hoffman et al., 2008; Santiago et al., 2008a, 2009). At*POLH* overexpression results in increased UV tolerance, while At*POLH* loss of function results in weak UV sensitivity, but enhances the sensitivity of the *rev3* mutant, evidence that Pol η is acting independently of Pol ζ (Curtis & Hays, 2007; Santiago et al., 2008b). *POLH* mutants

exhibit an increased mutation frequency, indicating a role in error-free repair. Interestingly, this increased mutation rate is suppressed in the *rev3 polh* double mutant, indicating that REV3 is required for the increased mutations in *polh* (Nakagawa et al., 2011). AtPol η interacts with the *Arabidopsis* PCNA homologues PCNA1 and PCNA2. In yeast the interaction with PCNA2 was found to be functionally important and require the PCNA-interacting protein (PIP) box and Ubiquitin-binding motif (UBM) of AtPol η (Anderson et al., 2008).

In other systems, PCNA monoubiquitination by RAD6/RAD18 is implicated in polymerase switching during translesion synthesis (Waters et al., 2009). While the *Arabidopsis* genome contains RAD6 homologues (*AtUBC1-3*), no obvious RAD18 homologue exists (Kraft et al., 2005). Interestingly, in mammalian cells, the CUL4-DDB1-CDT2 E3 ubiquitin ligase was recently shown to also monoubiquitinate PCNA and promote translesion synthesis (Terai et al., 2010). The CUL4-DDB1-CDT2 complex also ubiquitinates Pol η (Kim & Mischeal, 2008; Soria & Gottifredi, 2010), suggesting that this complex may be a key regulator of translesion synthesis (Abbas & Dutta, 2011). All components of the CUL4-DDB1-CDT2 complex exist in plants (Lee et al., 2008), so it will be interesting to examine the role of this complex in translesion synthesis.

5. Other responses

In addition to DNA repair and TLS, UV-damaged DNA also triggers other cellular responses such as cell cycle arrest and cell death (Batista et al., 2009). In general, relatively little is known about the molecular basis of these responses in plants. As in mammalian systems, in plants the initial damage sensors include ATM (double strand breaks), ATR (ssDNA) and ATR-interacting protein ATRIP (Culligan et al., 2004; Furukawa et al., 2010; Sakamoto et al., 2009). Plants do not have a p53 homologue, in contrast damage signalling pathways converge on the SOG1 transcription factor, resulting in cell cycle arrest and cell death (Furukawa et al., 2010; Preuss & Britt, 2003; Yoshiyama et al., 2009). UV-induced cell death in plants also involves topoisomerase I (Balestrazzi et al., 2010), metacaspase-8 and caspase-3-like activities (Danon et al., 2004; He et al., 2008; Zhang et al., 2009), as well as changes in reactive oxygen species and mitochondria (Gao et al., 2008).

6. Conclusions

Thus, despite the barrage of damage resulting from solar UV exposure plants face every day, they have a variety of mechanisms which allow them to survive. UV induced DNA damage is repaired by direct photoreactivation via photolyases, or by dark repair (NER) in both transcribed (TC-NER) or non-transcribed regions (GG-NER). Finally, rather than repair UV damage, plants can tolerate its presence and proceed with DNA replication via translesion synthesis, although there can be mutagenic consequences of this activity. The continued study of these pathways and the interplay between them in plants is sure to bring additional insight.

7. Acknowledgements

Research in our laboratory is supported by a grant from the Natural Sciences & Engineering Research Council of Canada (NSERC).

8. References

- Abbas, T. & Dutta, A. (2011). CRL4Cdt2: master coordinator of cell cycle progression and genome stability. *Cell Cycle*, 10(2):241-9
- Ahmad, M. & Cashmore, A.R. (1993). *HY4* gene of *A. thaliana* encodes a protein with characteristics of a blue-light photoreceptor. *Nature*, 366 (6451):162-6
- Ahmad, M., Jarillo, J.A., Klimczak, L.J., Landry, L.G., Peng, T., Last, R.L. & Cashmore, A.R. (1997). An enzyme similar to animal type II photolyases mediates photoreactivation in *Arabidopsis*. *Plant Cell*, 9(2):199-207
- Al Khateeb, W.M. & Schroeder, D.F. (2009). Overexpression of *Arabidopsis* damaged DNA binding protein 1A (DDB1A) enhances UV tolerance. *Plant Mol Biol.*, 70(4):371-83
- Anderson, H.J., Vonarx, E.J., Pastushok, L., Nakagawa, M., Katafuchi, A., Gruz, P., Di Rubbo, A., Grice, D.M., Osmond, M.J., Sakamoto, A.N., Nohmi, T., Xiao, W. & Kunz, B.A. (2008). *Arabidopsis thaliana* Y-family DNA polymerase eta catalyses translesion synthesis and interacts functionally with PCNA2. *Plant J.*, 55(6):895-908
- Araki, M., Masutani, C., Takemura, M., Uchida, A., Sugawara, K., Kondoh, J., Ohkuma, Y. & Hanaoka, F. (2001). Centrosome protein centrin 2/caltractin 1 is part of the xeroderma pigmentosum group C complex that initiates global genome nucleotide excision repair. *J Biol Chem.*, 276(22):18665-72
- Balestrazzi, A., Locato, V., Bottone, M.G., De Gara, L., Biggiogera, M., Pellicciari, C., Botti, S., Di Gesù, D., Donà, M. & Carbonera, D. (2010). Response to UV-C radiation in topo I-deficient carrot cells with low ascorbate levels. *J Exp Bot.*, 61(2):575-85
- Ballaré, C.L. (2003). Stress under the sun: spotlight on ultraviolet-B responses. *Plant Physiol.*, 132:1725-1727
- Batista, L.F., Kaina, B., Meneghini, R. & Menck, C.F. (2009). How DNA lesions are turned into powerful killing structures: insights from UV-induced apoptosis. *Mutat Res.*, 681(2-3):197-208
- Batschauer, A. (1999). Light perception in higher plants. *Cell Mol Life Sci.*, 55:153-165
- Bernhardt, A., Lechner, E., Hano, P., Schade, V., Dieterle, M., Anders, M., Dubin, M.J., Benvenuto, G., Bowler, C., Genschik, P. & Hellmann, H. (2006). CUL4 associates with DDB1 and DET1 and its downregulation affects diverse aspects of development in *Arabidopsis thaliana*. *Plant J.*, 47(4):591-603
- Bernhardt, A., Mooney, S. & Hellmann, H. (2010). *Arabidopsis* DDB1a and DDB1b are critical for embryo development. *Planta*, 232(3):555-566
- Biedermann, S. & Hellmann, H. (2010). The DDB1a interacting proteins ATCSA-1 and DDB2 are critical factors for UV-B tolerance and genomic integrity in *Arabidopsis thaliana*. *Plant J.*, 62(3):404-15
- Biedermann, S. & Hellmann, H. (2011). WD40 and CUL4-based E3 ligases: lubricating all aspects of life. *Trends Plant Sci.*, 16(1):38-46
- Britt, A.B. (1999). Molecular genetics of DNA repair in higher plants. *Trends Plant Sci.*, 4(1):20-25
- Brosché, M., Schuler, M.A., Kalbina, I., Connor, L. & Strid, A. (2002). Gene regulation by low level UV-B radiation: identification by DNA array analysis. *Photochem Photobiol Sci.*, 1(9):656-64
- Castells, E., Molinier, J., Drevensek, S., Genschik, P., Barneche, F. & Bowler, C. (2010). *det1-1*-induced UV-C hyposensitivity through *UVR3* and *PHR1* photolyase gene over-expression. *Plant J.*, 63:392-404

- Castells, E., Molinier, J., Benvenuto, G., Bourbousse, C., Zabulon, G., Zalc, A., Cazzaniga, S., Genschik, P., Barneche, F. & Bowler, C. (2011). The conserved factor DE-ETIOLATED 1 cooperates with CUL4-DDB1(DDB2) to maintain genome integrity upon UV stress. *EMBO J.*, 30(6):1162-72
- Chang, Y., Gong, L., Yuan, W., Li, X., Chen, G., Li, X., Zhang, Q. & Wu, C. (2009). Replication protein A (RPA1a) is required for meiotic and somatic DNA repair but is dispensable for DNA replication and homologous recombination in rice. *Plant Physiol.*, 151(4):2162-73
- Chen, H., Shen, Y., Tang, X., Yu, L., Wang, J., Guo, L., Zhang, Y., Zhang, H., Feng, S., Strickland, E., Zheng, N. & Deng, X. W. (2006). *Arabidopsis* CULLIN4 Forms an E3 Ubiquitin Ligase with RBX1 and the CDD Complex in Mediating Light Control of Development. *Plant Cell*, 18(8):1991-2004
- Chen, J.J., Mitchell, D.L. & Britt, A.B. (1994). A Light-Dependent Pathway for the Elimination of UV-Induced Pyrimidine (6-4) Pyrimidinone Photoproducts in *Arabidopsis*. *Plant Cell*, 6(9):1311-1317
- Chen, J.J., Jiang, C.Z. & Britt, A.B. (1996). Little or no repair of cyclobutyl pyrimidine dimers is observed in the organellar genomes of the young *Arabidopsis* seedlings. *Plant Physiol.*, 111(1):19-25
- Chen, J. & Suter, B. (2003). Xpd, a structural bridge and a functional link. *Cell Cycle*, 2(6):503-6
- Costa, R.M., Morgante, P.G., Berra, C.M., Nakabashi, M., Bruneau, D., Bouchez, D., Sweder, K.S., Van Sluys, M.A. & Menck, C.F. (2001). The participation of AtXPB1, the XPB/RAD25 homologue gene from *Arabidopsis thaliana*, in DNA repair and plant development. *Plant J.*, 28(4):385-95
- Culligan, K., Tissier, A. & Britt, A. (2004). ATR regulates a G2-phase cell-cycle checkpoint in *Arabidopsis thaliana*. *Plant Cell*, 16(5):1091-104
- Curtis, M.J. & Hays, J.B. (2007). Tolerance of dividing cells to replication stress in UVB-irradiated *Arabidopsis* roots: requirements for DNA translesion polymerases eta and zeta. *DNA Repair (Amst.)*, 6(9):1341-58
- Danon, A., Rotari, V.I., Gordon, A., Mailhac, N. & Gallois, P. (2004). Ultraviolet-C overexposure induces programmed cell death in *Arabidopsis*, which is mediated by caspase-like activities and which can be suppressed by caspase inhibitors, p35 and Defender against Apoptotic Death. *J Biol Chem.*, 279(1):779-87
- Deisenhofer, J. (2000). DNA photolyases and cryptochromes. *Mutat Res.*, 460(3-4):143-9
- Draper, C.K. & Hays, J.B. (2000). Replication of chloroplast, mitochondrial and nuclear DNA during growth of unirradiated and UVB-irradiated *Arabidopsis* leaves. *Plant J.*, 23(2):255-65
- Dubest, S., Gallego, M.E. & White, C.I. (2002). Role of the AtRad1p endonuclease in homologous recombination in plants. *EMBO Rep.*, 3(11):1049-54
- Dubest, S., Gallego, M.E. & White, C.I. (2004). Roles of the AtErcc1 protein in recombination. *Plant J.*, 39(3):334-42
- Farmer, L.M., Book, A.J., Lee, K.H., Lin, Y.L., Fu, H. & Vierstra, R.D. (2010). The RAD23 family provides an essential connection between the 26S proteasome and ubiquitylated proteins in *Arabidopsis*. *Plant Cell*, 22(1):124-42
- Fidantsef, A.L., Mitchell, D.L. & Britt, A.B. (2000). The *Arabidopsis* UVH1 gene is a homolog of the yeast repair endonuclease RAD1. *Plant Physiol.*, 124(2):579-86

- Fitch, M.E., Nakajima, S., Yasui, A. & Ford, J.M. (2003). In vivo recruitment of XPC to UV-induced cyclobutane pyrimidine dimers by the DDB2 gene product. *J Biol Chem.*, 278(47):46906-10
- Fousteri, M., Vermeulen, W., van Zeeland, A.A. & Mullenders, L.H. (2006). Cockayne syndrome A and B proteins differentially regulate recruitment of chromatin remodeling and repair factors to stalled RNA polymerase II in vivo. *Mol Cell*, 23(4):471-82
- Fousteri, M. & Mullenders, L.H. (2008). Transcription-coupled nucleotide excision repair in mammalian cells: molecular mechanisms and biological effects. *Cell Res.*, 18(1):73-84
- Frit, P., Bergmann, E. & Egly, J.M. (1999). Transcription factor IIIH: a key player in the cellular response to DNA damage. *Biochimie*, 81(1-2):27-38
- Frohnmeier, H. & Staiger, D. (2003). Ultraviolet-B Radiation-Mediated Responses in Plants. Balancing Damage and Protection. *Plant Physiol.*, 133(4):1420-1428
- Furukawa, T., Kimura, S., Ishibashi, T., Mori, Y., Hashimoto, J. & Sakaguchi, K. (2003a). OsSEND-1: a new RAD2 nuclease family member in higher plants. *Plant Mol Biol.*, 51(1):59-70
- Furukawa, T., Ishibashi, T., Kimura, S., Tanaka, H., Hashimoto, J. & Sakaguchi, K. (2003b). Characterization of all the subunits of replication factor C from a higher plant, rice (*Oryza sativa* L.), and their relation to development. *Plant Mol Biol.*, 53(1-2):15-25
- Furukawa, T., Curtis, M.J., Tominey, C.M., Duong, Y.H., Wilcox, B.W., Aggoune, D., Hays, J.B. & Britt, A.B. (2010). A shared DNA-damage-response pathway for induction of stem-cell death by UVB and by gamma irradiation. *DNA Repair (Amst.)*, 9(9):940-8
- Gallego, F., Fleck, O., Li, A., Wyrzykowska, J. & Tinland, B. (2000). AtRAD1, a plant homologue of human and yeast nucleotide excision repair endonucleases, is involved in dark repair of UV damages and recombination. *Plant J.*, 21(6):507-18
- Gao, C., Xing, D., Li, L. & Zhang, L. (2008). Implication of reactive oxygen species and mitochondrial dysfunction in the early stages of plant programmed cell death induced by ultraviolet-C overexposure. *Planta*, 227(4):755-67
- Garcia-Diaz, M. & Bebenek, K. (2007). Multiple functions of DNA polymerases. *CRC Crit Rev Plant Sci.*, 26(2):105-122
- García-Ortiz, M.V., Roldán-Arjona, T. & Ariza, R.R. (2007). The noncatalytic C-terminus of AtPOLK Y-family DNA polymerase affects synthesis fidelity, mismatch extension and translesion replication. *FEBS J.*, 274(13):3340-50
- Gillet, L.C.J. & Scharer, O.D. (2006). Molecular mechanisms of mammalian global genome nucleotide excision repair. *Chem. Rev.*, 106(2):253-276
- Grasser, M., Kane, C.M., Merkle, T., Melzer, M., Emmersen, J. & Grasser, K.D. (2009). Transcript elongation factor TFIIS is involved in *Arabidopsis* seed dormancy. *J Mol Biol.*, 386(3):598-611
- Groisman, R., Polanowska, J., Kuraoka, I., Sawada, J., Saijo, M., Drapkin, R., Kissilev, A.F., Tanaka, K. & Nakatani, Y. (2003). The ubiquitin ligase activity in the DDB2 and CSA complexes is differentially regulated by the COP9 signalosome in response to DNA damage. *Cell*, 113(3): 357-367
- Groisman, R., Kuraoka, I., Chevallier, O., Gaye, N., Magnaldo, T., Tanaka, K., Kisselev, A.F., Harel-Bellan, A. & Nakatani, Y. (2006). CSA-dependent degradation of CSB by the ubiquitin-proteasome pathway establishes a link between complementation factors of the Cockayne syndrome. *Genes Dev.*, 20(11):1429-34

- Guerrero-Santoro, J., Kapetanaki, M.G., Hsieh, C.L., Gorbachinsky, I., Levine, A.S. & Rapić-Otrin, V. (2008). The cullin 4B-based UV-damaged DNA-binding protein ligase binds to UV-damaged chromatin and ubiquitinates histone H2A. *Cancer Res.*, 68(13):5014-22
- Gulbis, J.M., Kelman, Z., Hurwitz, J., O'Donnell, M. & Kuriyan, J. (1996). Structure of the C-terminal region of p21(WAF1/CIP1) complexed with human PCNA. *Cell*, 87(2):297-306
- Hada, M., Hino, K., Buchholz, G., Goss, J., Wellmann, E. & Shin, M. (2000). Assay of DNA photolyase activity in spinach leaves in relation to cell compartmentation-evidence for lack of DNA photolyase in chloroplasts. *Biosci Biotechnol Biochem.*, 64:1288-1291
- Hanawalt, P.C. & Spivak, G. (2008). Transcription-coupled DNA repair: two decades of progress and surprises. *Nat Rev Mol Cell Biol.*, 9(12):958-70
- He, R., Drury, G.E., Rotari, V.I., Gordon, A., Willer, M., Farzaneh, T., Woltering, E.J. & Gallois, P. (2008). Metacaspase-8 modulates programmed cell death induced by ultraviolet light and H₂O₂ in *Arabidopsis*. *J Biol Chem.*, 283(2):774-83
- He, Z., Henricksen, L.A., Wold, M.S. & Ingles, C.J. (1995). RPA involvement in the damage-recognition and incision steps of nucleotide excision repair. *Nature*, 374: 566-569
- Hefner, E., Preuss, S.B. & Britt, A.B. (2003). *Arabidopsis* mutants sensitive to gamma radiation include the homologue of the human repair gene ERCC1. *J Exp Bot.*, 54(383):669-80
- Hey, T., Lipps, G., Sugawara, K., Iwai, S., Hanaoka, F. & Krauss, G. (2002). The XPC-HR23B complex displays high affinity and specificity for damaged DNA in a true-equilibrium fluorescence assay. *Biochemistry*, 41(21): 6583-6587
- Hidema, J., Kumagai, T. & Sutherland, B.M. (2000). UV radiation-sensitive norin 1 rice contains defective cyclobutane pyrimidine dimer photolyase. *Plant Cell*, 12(9):1569-78
- Hidema, J., Taguchi, T., Ono, T., Teranishi, M., Yamamoto, K. & Kumagai, T. (2007). Increase in CPD photolyase activity functions effectively to prevent growth inhibition caused by UVB radiation. *Plant J.*, 50(1):70-9
- Higa, L.A., Wu, M., Ye, T., Kobayashi, R., Sun, H. & Zhang, H. (2006). CUL4-DDB1 ubiquitin ligase interacts with multiple WD40-repeat proteins and regulates histone methylation. *Nat Cell Biol.*, 8(11):1277-83
- Hirouchi, T., Nakajima, S., Najrana, T., Tanaka, M., Matsunaga, T., Hidema, J., Teranishi, M., Fujino, T., Kumagai, T. & Yamamoto, K. (2003). A gene for a Class II DNA photolyase from *Oryza sativa*: cloning of the cDNA by dilution-amplification. *Mol Genet Genomics*, 269(4):508-16
- Hoeijmakers, J.H. (2001). Genome maintenance mechanisms for preventing cancer. *Nature*, 411: 135-141
- Hoffman, P.D., Curtis, M.J., Iwai, S. & Hays, J.B. (2008). Biochemical evolution of DNA polymerase eta: properties of plant, human, and yeast proteins. *Biochemistry*, 47(16):4583-96
- Holmberg, C., Fleck, O., Hansen, H.A., Liu, C., Slaaby, R., Carr, A.M. & Nielsen, O. (2005). Ddb1 controls genome stability and meiosis in fission yeast. *Genes Dev.*, 19:853-862
- Hua, Z. & Vierstra, R.D. (2011). The Cullin-RING Ubiquitin-Protein Ligases. *Annu Rev Plant Biol.*, 62:299-334

- Ishibashi, T., Kimura, S., Furukawa, T., Hatanaka, M., Hashimoto, J. & Sakaguchi, K. (2001). Two types of replication protein A 70 kDa subunit in rice, *Oryza sativa*: molecular cloning, characterization, and cellular & tissue distribution. *Gene*, 272(1-2):335-43
- Ishibashi, T., Kimura, S., Yamamoto, T., Furukawa, T., Takata, K., Uchiyama, Y., Hashimoto, J. & Sakaguchi, K. (2003). Rice UV-damaged DNA binding protein homologues are most abundant in proliferating tissues. *Gene*, 308:79-87
- Ishibashi, T., Koga, A., Yamamoto, T., Uchiyama, Y., Mori, Y., Hashimoto, J., Kimura, S. & Sakaguchi, K. (2005). Two types of replication protein A in seed plants. *FEBS J.*, 272(13):3270-81
- Ishibashi, T., Kimura, S. & Sakaguchi, K. (2006). A higher plant has three different types of RPA heterotrimeric complex. *J Biochem.*, 139(1):99-104
- Jansan, M.A.K., Hectors, K., O'Brien, N.M., Guisez, Y. & Potters, G. (2008). Plant stress and human health: Do human consumers benefit from UV-B acclimated crops? *Plant Science*, 175:449-458
- Jenkins, G.I. (2009). Signal transduction in responses to UV-B radiation. *Annu Rev Plant Biol.*, 60:407-31
- Jenkins, M.E., Suzuki, T.C. & Mount, D.W. (1997). Evidence that heat and ultraviolet radiation activate a common stress-response program in plants that is altered in the *uvh6* mutant of *Arabidopsis thaliana*. *Plant Physiol.*, 115(4):1351-8
- Kaiser, G., Kleiner, O., Beisswenger, C. & Batschauer, A. (2009). Increased DNA repair in *Arabidopsis* plants overexpressing CPD photolyase. *Planta*, 230(3):505-15
- Kapetanaki, M.G., Guerrero-Santoro, J., Bisi, D.C., Hsieh, C.L., Rapić-Otrin, V. & Levine, A.S. (2006). The DDB1-CUL4ADDDB2 ubiquitin ligase is deficient in xeroderma pigmentosum group E and targets histone H2A at UV-damaged DNA sites. *Proc Natl Acad Sci U S A.*, 103(8):2588-93
- Kim, S.H. & Michael, W.M. (2008). Regulated proteolysis of DNA polymerase eta during the DNA-damage response in *C. elegans*. *Mol Cell*, 32(6):757-66
- Kimura, S., Tahira, Y., Ishibashi, T., Mori, Y., Mori, T., Hashimoto, J. & Sakaguchi, K. (2004). DNA repair in higher plants; photoreactivation is the major DNA repair pathway in non-proliferating cells while excision repair (nucleotide excision repair and base excision repair) is active in proliferating cells. *Nucleic Acids Res.*, 32(9):2760-7
- Kimura, S. & Sakaguchi, K. (2006). DNA repair in plants. *Chem Rev.*, 106(2):753-66
- Koga, A., Ishibashi, T., Kimura, S., Uchiyama, Y. & Sakaguchi, K. (2006). Characterization of T-DNA insertion mutants and RNAi silenced plants of *Arabidopsis thaliana* UV-damaged DNA binding protein 2 (AtUV-DDB2). *Plant Mol Biol.*, 61(1-2):227-40
- Kraft, E., Stone, S.L., Ma, L., Su, N., Gao, Y., Lau, O.S., Deng, X.W. & Callis, J. (2005). Genome analysis and functional characterization of the E2 and RING-type E3 ligase ubiquitination enzymes of *Arabidopsis*. *Plant Physiol.*, 139(4):1597-611
- Kunz, B.A., Anderson, H.J., Osmond, M.J. & Vonarx, E.J. (2005). Components of nucleotide excision repair and DNA damage tolerance in *Arabidopsis thaliana*. *Environ Mol Mutagen.*, 45(2-3):115-27
- Kunz, B.A., Cahill, D.M., Mohr, P.G., Osmond, M.J. & Vonarx, E.J. (2006). Plant responses to UV radiation and links to pathogen resistance. *Int Rev Cytol.*, 255:1-40
- Kuraoka, I., Ito, S., Wada, T., Hayashida, M., Lee, L., Saijo, M., Nakatsu, Y., Matsumoto, M., Matsunaga, T., Handa, H., Qin, J., Nakatani, Y. & Tanaka, K. (2008). Isolation of

- XAB2 complex involved in pre-mRNA splicing, transcription, and transcription-coupled repair. *J Biol Chem.*, 283(2):940-50
- Lainé, J.P. & Egly, J.M. (2006). Initiation of DNA repair mediated by a stalled RNA polymerase II. *EMBO J.*, 25(2):387-97
- Landry, L.G., Stapleton, A.E., Lim, J., Hoffman, P., Hays, J.B., Walbot, V. & Last, R.L. (1997). An *Arabidopsis* photolyase mutant is hypersensitive to ultraviolet-B radiation. *Proc Natl Acad Sci U S A.*, 94(1):328-32
- Lee, J. & Zhou, P. (2007). DCAFs, the missing link of the CUL4-DDB1 ubiquitin ligase. *Mol Cell*, 26(6):775-80
- Lee, J.H., Terzaghi, W., Gusmaroli, G., Charron, J.B., Yoon, H.J., Chen, H., He, Y.J., Xiong, Y. & Deng, X.W. (2008). Characterization of *Arabidopsis* and rice DWD proteins and their roles as substrate receptors for CUL4-RING E3 ubiquitin ligases. *Plant Cell*, 20(1):152-67
- Liang, L., Flury, S., Kalck, V., Hohn, B. & Molinier, J. (2006). CENTRIN2 interacts with the *Arabidopsis* homolog of the human XPC protein (AtRAD4) and contributes to efficient synthesis-dependent repair of bulky DNA lesions. *Plant Mol Biol.*, 61(1-2):345-56
- Liu, Z., Hossain, G.S., Islas-Osuna, M.A., Mitchell, D.L. & Mount, D.W. (2000). Repair of UV damage in plants by nucleotide excision repair: *Arabidopsis* UVH1 DNA repair gene is a homolog of *Saccharomyces cerevisiae* Rad1. *Plant J.*, 21(6):519-28
- Liu, Z., Hall, J.D. & Mount, D.W. (2001). *Arabidopsis* UVH3 gene is a homolog of the *Saccharomyces cerevisiae* RAD2 and human XPG DNA repair genes. *Plant J.*, 26(3):329-38
- Liu, Z., Hong, S.W., Escobar, M., Vierling, E., Mitchell, D.L., Mount, D.W. & Hall, J.D. (2003). *Arabidopsis* UVH6, a homolog of human XPD and yeast RAD3 DNA repair genes, functions in DNA repair and is essential for plant growth. *Plant Physiol.*, 132(3):1405-14
- Luijsterburg, M.S., Goedhart, J., Moser, J., Kool, H., Geverts, B., Houtsmuller, A.B., Mullenders, L.H., Vermeulen, W. & van Driel, R. (2007). Dynamic in vivo interaction of DDB2 E3 ubiquitin ligase with UV-damaged DNA is independent of damage-recognition protein XPC. *J Cell Sci.*, 120(15):2706-16
- Mackerness, S. (2000). Plant responses to ultraviolet-B (UV-B:280–320 nm) stress: What are the key regulators? *Plant Growth Regul.*, 37:27-39
- Marwedel, T., Ishibashi, T., Lorbiecke, R., Jacob, S., Sakaguchi, K. & Sauter, M. (2003). Plant-specific regulation of replication protein A2 (OsRPA2) from rice during the cell cycle and in response to ultraviolet light exposure. *Planta.*, 217(3):457-65
- Mellon, I. (2005). Transcription-coupled repair: a complex affair. *Mutat Res.*, 577(1-2):155-61
- Mocquet, V., Lainé, J.P., Riedl, T., Yajin, Z., Lee, M.Y. & Egly, J.M. (2008). Sequential recruitment of the repair factors during NER: the role of XPG in initiating the resynthesis step. *EMBO J.*, 27(1):155-67
- Molinier, J., Ramos, C., Fritsch, O. & Hohn, B. (2004). CENTRIN2 modulates homologous recombination and nucleotide excision repair in *Arabidopsis*. *Plant Cell*, 16(6):1633-43
- Molinier, J., Lechner, E., Dumbliuskas, E. & Genschik, P. (2008). Regulation and role of *Arabidopsis* CUL4-DDB1A-DDB2 in maintaining genome integrity upon UV stress. *PloS Genetics*, 4(6):e10000

- Morgante, P.G., Berra, C.M., Nakabashi, M., Costa, R.M., Menck, C.F. & Van Sluys, M.A. (2005). Functional XPB/RAD25 redundancy in *Arabidopsis* genome: characterization of AtXPB2 and expression analysis. *Gene*, 344:93-103
- Nakagawa, M., Takahashi, S., Tanaka, A., Narumi, I. & Sakamoto, A.N. (2011). Role of AtPol{zeta}, AtRev1, and AtPol{eta} in UV light-induced mutagenesis in *Arabidopsis*. *Plant Physiol.*, 155(1):414-20
- Nakajima, S., Sugiyama, M., Iwai, S., Hitomi, K., Otsoshi, E., Kim, S.T., Jiang, C.Z., Todo, T., Britt, A.B. & Yamamoto, K. (1998). Cloning and characterization of a gene (*UVR3*) required for photorepair of 6-4 photoproducts in *Arabidopsis thaliana*. *Nucleic Acids Res.*, 26(2):638-44
- Nakatsu, Y., Asahina, H., Citterio, E., Rademakers, S., Vermeulen, W., Kamiuchi, S., Yeo, J.P., Khaw, M.C., Saijo, M., Kodo, N., Matsuda, T., Hoeijmakers, J.H. & Tanaka, K. (2000). XAB2, a novel tetratricopeptide repeat protein involved in transcription-coupled DNA repair and transcription. *J Biol Chem.*, 275(45):34931-7
- Nishi, R., Okuda, Y., Watanabe, E., Mori, T., Iwai, S., Masutani, C., Sugawara, K. & Hanaoka, F. (2005). Centrin2 stimulates nucleotide excision repair by interacting with xeroderma pigmentosum group C protein. *Mol Cell Biol.*, 25(13):5664-74
- Oravec, A., Baumann, A., Máté, Z., Brzezinska, A., Molinier, J., Oakeley, E.J., Adám, E., Schäfer, E., Nagy, F. & Ulm, R. (2006). CONSTITUTIVELY PHOTOMORPHOGENIC1 is required for the UV-B response in *Arabidopsis*. *Plant Cell*, 18(8):1975-90
- Pandey, R., Müller, A., Napoli, C.A., Selinger, D.A., Pikaard, C.S., Richards, E.J., Bender, J., Mount, D.W. & Jorgensen, R.A. (2002). Analysis of histone acetyltransferase and histone deacetylase families of *Arabidopsis thaliana* suggests functional diversification of chromatin modification among multicellular eukaryotes. *Nucleic Acids Res.*, 30(23):5036-55
- Pang, Q. & Hays, J.B. (1991). UV-B-Inducible and Temperature-Sensitive Photoreactivation of Cyclobutane Pyrimidine Dimers in *Arabidopsis thaliana*. *Plant Physiol.*, 95(2):536-43
- Parry, G. & Estelle, M. (2004). Regulation of Cullin based ubiquitin ligases by the Nedd8/RUB ubiquitin like proteins. *Semin Cell Dev. Biol.*, 15: 221-229
- Pepper, A., Delaney, T., Washburn, T., Poole, D. & Chory, J. (1994). DET1, a negative regulator of light-mediated development and gene expression in *Arabidopsis*, encodes a novel nuclear-localized protein. *Cell*, 78(1):109-16
- Preuss, S.B. & Britt, A.B. (2003). A DNA-damage-induced cell cycle checkpoint in *Arabidopsis*. *Genetics*, 164(1):323-34
- Rapić-Otrin, V., McLenigan, M.P., Bisi, D.C., Gonzalez, M. & Levine, A.S. (2002). Sequential binding of UV DNA damage binding factor and degradation of the p48 subunit as early events after UV irradiation. *Nucleic Acids Res.*, 30(11):2588-98
- Ribeiro, D.T., Machado, C.R., Costa, R.M., Praekelt, U.M., Van Sluys, M.A. & Menck, C.F. (1998). Cloning of a cDNA from *Arabidopsis thaliana* homologous to the human XPB gene. *Gene*, 208(2):207-13
- Riedl, T., Hanaoka, F. & Egly, J.M. (2003). The comings and goings of nucleotide excision repair factors on damaged DNA. *EMBO J.*, 22(19):5293-303
- Ries, G., Heller, W., Puchta, H., Sandermann, H., Seidlitz, H.K. & Hohn, B. (2000). Elevated UV-B radiation reduces genome stability in plants. *Nature*, 406(6791):98-101


- Roy, S., Choudhury, S.R., Singh, S.K. & Das, K.P. (2011). AtPol λ , A Homolog of Mammalian DNA Polymerase λ in *Arabidopsis thaliana*, is Involved in the Repair of UV-B Induced DNA Damage through the Dark Repair Pathway. *Plant Cell Physiol.*, 52(2): 448-467
- Sakaguchi, K., Ishibashi, T., Uchiyama, Y. & Iwabata, K. (2009). The multi-replication protein A (RPA) system--a new perspective. *FEBS J.*, 276(4):943-63
- Sakamoto, A., Lan, V.T., Hase, Y., Shikazono, N., Matsunaga, T. & Tanaka, A. (2003). Disruption of the *AtREV3* gene causes hypersensitivity to ultraviolet B light and gamma-rays in *Arabidopsis*: implication of the presence of a translesion synthesis mechanism in plants. *Plant Cell*, 15(9):2042-57
- Sakamoto, A.N., Lan, V.T., Puripunyanich, V., Hase, Y., Yokota, Y., Shikazono, N., Nakagawa, M., Narumi, I. & Tanaka, A. (2009). A UVB-hypersensitive mutant in *Arabidopsis thaliana* is defective in the DNA damage response. *Plant J.*, 60(3):509-17
- Sancar, A. (2003). Structure and function of DNA photolyase and cryptochrome blue-light photoreceptors. *Chem Rev.*, 103(6):2203-37
- Sancar, A. (2008). Structure and function of photolyase and in vivo enzymology: 50th anniversary. *J Biol Chem.*, 283(47):32153-7
- Santiago, M.J., Alejandre-Durán, E. & Ruiz-Rubio, M. (2006). Analysis of UV-induced mutation spectra in *Escherichia coli* by DNA polymerase eta from *Arabidopsis thaliana*. *Mutat Res.*, 601(1-2):51-60
- Santiago, M.J., Ruiz-Rubio, M., Dio, L.D., González-Reyes, J.A. & Alejandre-Durán, E. (2008a). Ubiquitous expression of two translesion synthesis DNA polymerase genes in *Arabidopsis*. *Planta*, 227(6):1269-77
- Santiago, M.J., Alejandre-Durán, E., Muñoz-Serrano, A. & Ruiz-Rubio, M. (2008b). Two translesion synthesis DNA polymerase genes, *AtPOLH* and *AtREV1*, are involved in development and UV light resistance in *Arabidopsis*. *J Plant Physiol.*, 165(15):1582-91
- Santiago, M.J., Alejandre-Durán, E. & Ruiz-Rubio, M. (2009). Alternative splicing of two translesion synthesis DNA polymerases from *Arabidopsis thaliana*. *Plant Science*, 176(4):591-596
- Saxowsky, T.T. & Doetsch, P.W. (2006). RNA polymerase encounters with DNA damage: transcription-coupled repair or transcriptional mutagenesis? *Chem Rev.*, 106(2):474-88
- Schroeder, D.F., Gahrtz, M., Maxwell, B.B., Cook, R.K., Kan, J.M., Alonso, J.M., Ecker, J.R. & Chory, J. (2002). De-etiolated1 and Damaged DNA Binding Protein 1 interact to regulate *Arabidopsis* photomorphogenesis. *Curr Biol.*, 12(17):1462-72
- Schultz, T.F. & Quatrano R.S. (1997). Characterization and expression of a rice *RAD23* gene. *Plant Mol Biol.*, 34: 557-562
- Schwechheimer, C. & Deng, X.W. (2001). COP9 signalosome revisited: a novel mediator of protein degradation. *Trends Cell Biol.*, 11(10):420-426
- Scrima, A., Konícková, R., Czyzewski, B.K., Kawasaki, Y., Jeffrey, P.D., Groisman, R., Nakatani, Y., Iwai, S., Pavletich, N.P. & Thomä, N.H. (2008). Structural basis of UV DNA-damage recognition by the DDB1-DDB2 complex. *Cell*, 135(7):1213-23
- Shaked, H., Avivi-Ragolsky, N. & Levy, A.A. (2006). Involvement of the *Arabidopsis* SWI2/SNF2 chromatin remodeling gene family in DNA damage response and recombination. *Genetics*, 173(2):985-94

- Shultz, R.W., Tatineni, V.M., Hanley-Bowdoin, L. & Thompson, W.F. (2007). Genome-wide analysis of the core DNA replication machinery in the higher plants *Arabidopsis* and rice. *Plant Physiol.*, 144(4):1697-714
- Singh, S.K., Roy, S., Choudhury, S.R. & Sengupta, D.N. (2010). DNA repair and recombination in higher plants: insights from comparative genomics of *Arabidopsis* and rice. *BMC Genomics*, 11:443
- Soria, G. & Gottifredi, V. (2010). PCNA-coupled p21 degradation after DNA damage: The exception that confirms the rule? *DNA Repair (Amst)*, 9(4):358-64
- Strzalka, W. & Ziemienowicz, A. (2011). Proliferating cell nuclear antigen (PCNA): a key factor in DNA replication and cell cycle regulation. *Ann Bot.* 107(7):1127-40
- Sturm, A. & Leinhard, S. (1998). Two isoforms of plant RAD 23 complement a UV sensitive rad23 mutant in yeast. *Plant J.*, 13(6): 815-821
- Sugasawa, K., Okuda, Y., Saijo, M., Nishi, R., Matsuda, N., Chu, G., Mori, T., Iwai, S., Tanaka, K., Tanaka, K. & Hanaoka, F. (2005). UV-induced ubiquitylation of XPC protein mediated by UV-DDB-ubiquitin ligase complex. *Cell*, 121(3):387-400
- Svejstrup, J.Q. (2002). Mechanisms of Transcription coupled repair. *Nat Rev Mol Cell Biol.*, 3(1):21-29
- Takahashi, M., Teranishi, M., Ishida, H., Kawasaki, J., Takeuchi, A., Yamaya, T., Watanabe, M., Makino, A. & Hidema, J. (2011). Cyclobutane pyrimidine dimer (CPD) photolyase repairs ultraviolet-B-induced CPDs in rice chloroplast and mitochondrial DNA. *Plant J.*, 66(3):433-42
- Takahashi, S., Nakajima, N., Saji, H. & Kondo, N. (2002). Diurnal change of cucumber CPD photolyase gene (*CsPHR*) expression and its physiological role in growth under UV-B irradiation. *Plant Cell Physiol.*, 43(3):342-9
- Takahashi, S., Sakamoto, A., Sato, S., Kato, T., Tabata, S. & Tanaka, A. (2005). Roles of *Arabidopsis* AtREV1 and AtREV7 in translesion synthesis. *Plant Physiol.*, 138(2):870-81
- Takahashi, S., Sakamoto, A.N., Tanaka, A. & Shimizu, K. (2007). AtREV1, a Y-family DNA polymerase in *Arabidopsis*, has deoxynucleotidyl transferase activity in vitro. *Plant Physiol.*, 145(3):1052-60
- Takashi, Y., Kobayashi, Y., Tanaka, K. & Tamura, K. (2009). *Arabidopsis* replication protein A 70a is required for DNA damage response and telomere length homeostasis. *Plant Cell Physiol.*, 50(11):1965-76
- Tanaka, A., Sakamoto, A., Ishigaki, Y., Nikaido, O., Sun, G., Hase, Y., Shikazono, N., Tano, S. & Watanabe, H. (2002). An ultraviolet-B resistant mutant with enhanced DNA repair in *Arabidopsis*. *Plant Physiol.*, 129(1):64-71
- Tapias, A., Auriol, J., Forget, D., Enzlin, J.H., Schärer, O.D., Coin, F., Coulombe, B. & Egly, J.M. (2004). Ordered conformational changes in damaged DNA induced by nucleotide excision repair factors. *J Biol Chem.*, 279(18):19074-83
- Terai, K., Abbas, T., Jazaeri, A.A. & Dutta, A. (2010). CRL4(Cdt2) E3 ubiquitin ligase monoubiquitinates PCNA to promote translesion DNA synthesis. *Mol Cell*, 37(1):143-9
- Todo, T., Ryo, H., Yamamoto, K., Toh, H., Inui, T., Ayaki, H., Nomura, T. & Ikenaga, M. (1996). Similarity among the *Drosophila* (6-4)photolyase, a human photolyase homolog, and the DNA photolyase-blue-light photoreceptor family. *Science*, 272(5258):109-12

- Tornaletti, S. (2009). DNA repair in mammalian cells: Transcription-coupled DNA repair: directing your effort where it's most needed. *Cell Mol Life Sci.*, 66(6):1010-20
- Ueda, T., Sato, T., Hidema, J., Hirouchi, T., Yamamoto, K., Kumagai, T. & Yano, M. (2005). qUVR-10, a major quantitative trait locus for ultraviolet-B resistance in rice, encodes cyclobutane pyrimidine dimer photolyase. *Genetics*, 171(4):1941-50
- Ulm, R., Baumann, A., Oravecz, A., Máté, Z., Adám, E., Oakeley, E.J., Schäfer, E. & Nagy, F. (2004). Genome-wide analysis of gene expression reveals function of the bZIP transcription factor HY5 in the UV-B response of *Arabidopsis*. *Proc Natl Acad Sci U S A.*, 101(5):1397-402
- Vannier, J.B., Depeiges, A., White, C. & Gallego, M.E. (2009). ERCC1/XPF protects short telomeres from homologous recombination in *Arabidopsis thaliana*. *PLoS Genet.*, 5(2):e1000380
- van den Boom, V., Citterio, E., Hoogstraten, D., Zotter, A., Egly, J.M., van Cappellen, W.A., Hoeijmakers, J.H., Houtsmuller, A.B. & Vermeulen, W. (2004). DNA damage stabilizes interaction of CSB with the transcription elongation machinery. *J Cell Biol.*, 166(1):27-36
- Volker, M., Mone, M.J., Karmakar, P., van Hoffen, A., Schul, W., Vermeulen, W., Hoeijmakers, J.H., van Driel, R., van Zeeland, A.A. & Mullenders, L.H. (2001). Sequential assembly of the nucleotide excision repair factors in vivo. *Mol Cell*, 8(1):213-224
- Vonarx, E.J., Howlett, N.G., Schiestl, R.H. & Kunz, B.A. (2002). Detection of *Arabidopsis thaliana* AtRAD1 cDNA variants and assessment of function by expression in a yeast rad1 mutant. *Gene*, 296(1-2):1-9
- Vonarx, E.J., Tabone, E.K., Osmond, M.J., Anderson, H.J. & Kunz, B.A. (2006). *Arabidopsis* homologue of human transcription factor IIH/nucleotide excision repair factor p44 can function in transcription and DNA repair and interacts with AtXPD. *Plant J.*, 46(3):512-21
- Wang, H., Zhai, L., Xu, J., Joo, H.Y., Jackson, S., Erdjument-Bromage, H., Tempst, P., Xiong, Y. & Zhang, Y. (2006). Histone H3 and H4 ubiquitylation by the CUL4-DDB-ROC1 ubiquitin ligase facilitates cellular response to DNA damage. *Mol Cell*, 22(3):383-94
- Waters, L.S., Minesinger, B.K., Wilttrout, M.E., D'Souza, S., Woodruff, R.V. & Walker, G.C. (2009). Eukaryotic translesion polymerases and their roles and regulation in DNA damage tolerance. *Microbiol Mol Biol Rev.*, 73(1):134-54
- Waterworth, W.M., Jiang, Q., West, C.E., Nikaido, M. & Bray, C.M. (2002). Characterization of *Arabidopsis* photolyase enzymes and analysis of their role in protection from ultraviolet-B radiation. *J Exp Bot.*, 53(371):1005-15
- Weeda, G., Donker, I., De Wit, J., Morreau, H., Janssens, R., Vissers, C.J., Nigg, A., van Steeg, H., Bootsma, D. & Hoeijmakers, J.H. (1997). Disruption of mouse ERCC1 results in a novel repair syndrome with growth failure, nuclear abnormalities and senescence. *Curr Biol.*, 7(6): 427-39
- Wold, M.S. (1997). Replication protein A: a heterotrimeric, single-stranded DNA-binding protein required for eukaryotic DNA metabolism. *Annu Rev Biochem.*, 66:61-92
- Xu, H., Swoboda, I., Bhalla, P.L., Sijbers, A.M., Zhao, C., Ong, E.K., Hoeijmakers, J.H. & Singh, M.B. (1998). Plant homologue of human excision repair gene ERCC1 points to conservation of DNA repair mechanisms. *Plant J.*, 13(6):823-9

- Yamamoto, A., Hirouchi, T., Mori, T., Teranishi, M., Hidema, J., Morioka, H., Kumagai, T. & Yamamoto, K. (2007). Biochemical and biological properties of DNA photolyases derived from ultraviolet-sensitive rice cultivars. *Genes Genet Syst.*, 82(4):311-9
- Yamamoto, A., Tanbir, N., Hirouchi, T., Teranishi, M., Hidema, J., Morioka, H. & Yamamoto, K. (2008). Temperature-sensitive photoreactivation of cyclobutane thymine dimer in soybean. *J Radiat Res (Tokyo)*, 49(2):189-96
- Yasui, A., Eker, A.P., Yasuhira, S., Yajima, H., Kobayashi, T., Takao, M. & Oikawa, A. (1994). A new class of DNA photolyases present in various organisms including aplacental mammals. *EMBO J.*, 13(24):6143-51
- Yoshihara, R., Imaki, T., Hori, M., Watanabe, C., Yamamoto, K. & Takimoto, K. (2005). CPD photolyase gene from *Spinacia oleracea*: repair of UV-damaged DNA and expression in plant organs. *J Radiat Res (Tokyo)*, 46(2):157-64
- Yoshiyama, K., Conklin, P.A., Huefner, N.D. & Britt, A.B. (2009). Suppressor of gamma response 1 (SOG1) encodes a putative transcription factor governing multiple responses to DNA damage. *Proc Natl Acad Sci USA*, 106(31):12843-8
- Yuzhakov, A., Kelman, Z., Hurwitz, J. & O'Donnell, M. (1999). Multiple competition reactions for RPA order the assembly of the DNA polymerase delta holoenzyme. *EMBO J.*, 18(21):6189-99
- Zhang, C., Guo, H., Zhang, J., Guo, G., Schumaker, K.S. & Guo, Y. (2010). *Arabidopsis* cockayne syndrome A-like proteins 1A and 1B form a complex with CULLIN4 and damage DNA binding protein 1A and regulate the response to UV irradiation. *Plant Cell*, 22(7):2353-69
- Zhang, L., Xu, Q., Xing, D., Gao, C. & Xiong, H. (2009). Real-time detection of caspase-3-like protease activation in vivo using fluorescence resonance energy transfer during plant programmed cell death induced by ultraviolet C overexposure. *Plant Physiol.*, 150(4):1773-83

IntechOpen



Selected Topics in DNA Repair

Edited by Prof. Clark Chen

ISBN 978-953-307-606-5

Hard cover, 572 pages

Publisher InTech

Published online 26, October, 2011

Published in print edition October, 2011

This book is intended for students and scientists working in the field of DNA repair, focusing on a number of topics ranging from DNA damaging agents and mechanistic insights to methods in DNA repair and insights into therapeutic strategies. These topics demonstrate how scientific ideas are developed, tested, dialogued, and matured as it is meant to discuss key concepts in DNA repair. The book should serve as a supplementary text in courses and seminars as well as a general reference for biologists with an interest in DNA repair.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Ashwin L. Ganpudi and Dana F. Schroeder (2011). UV Damaged DNA Repair & Tolerance in Plants, Selected Topics in DNA Repair, Prof. Clark Chen (Ed.), ISBN: 978-953-307-606-5, InTech, Available from: <http://www.intechopen.com/books/selected-topics-in-dna-repair/uv-damaged-dna-repair-tolerance-in-plants>

INTeCH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2011 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](https://creativecommons.org/licenses/by/3.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen