We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Childhood Acute Leukemias in Hispanic Population: Differences by Age Peak and Immunophenotype

Juan Manuel Mejía-Aranguré et al.*

Unidad de Investigación en Epidemiología Clínica, Enfermedades Oncológicas, Enfermedades Infecciosas y Parasitarias/Hospital de Pediatría & División de Laboratorios de Vigilancia e Investigación Epidemiológica, Instituto Mexicano del Seguro Social México

1. Introduction

Childhood cancer represents 0.5–5.7% of all cancers (Birch & Blair, 1992; Smith & Gloecker, 2002). The most important kinds of cancer during childhood differ from those most frequently found in adulthood (Schellong, 1985). During infancy, the principal cancers are not epithelial, in contrast to those in the later stages of life (Miller, 1983). Therefore, it is possible that the risk factors for cancer are different during infancy than during adulthood; for children, risk factors may be present *in utero* or during the firsts months of the life (Draper, 1994).

The age peak of cancer during infancy, especially those for leukemias and lymphomas, varies among countries. Although the peak age for leukemias worldwide is principally between 2–5 years of age, a peak as late as 7–13 years of age was reported for Niger (William, 1975). In developed countries such as Germany or the United States of America (USA), the age peak for lymphomas is between 10–14 years of age, whereas in less developed countries, Mexico for example, the age peak is between 5–9 years of age (Fajardo-Gutiérrez et al., 1995; Kaatsch et al., 1995; Nully-Brown et al., 1989; Young et al., 1986).

Leukemia is the most common cancer in children under 15 years old, representing between 25–35% of all childhood cancers in most populations (Parkin et al., 1988, 1998). Leukemia is a heterogeneous group of hematopoietic malignancies, with several biologically distinct subgroups. The main subtypes of leukemia described by most cancer registries include acute lymphoblastic leukemia (ALL), representing about 80% of all leukemias; AML, with a frequency of 15%; and chronic myeloid leukemia (CML) with a frequency of 3–5% (Bathia, 2003). ALL is the most frequent leukemia in infancy (Mejía Arangure et al., 2005a). The age

* María Luisa Pérez-Saldivar, Rosana Pelayo-Camacho, Ezequiel Fuentes-Pananá, Carolina Bekker-Mendez, Abigail Morales-Sánchez, David Aldebarán Duarte-Rodríguez and Arturo Fajardo-Gutiérrez

Unidad de Investigación en Epidemiología Clínica, Enfermedades Oncológicas, Enfermedades Infecciosas y Parasitarias/Hospital de Pediatría & División de Laboratorios de Vigilancia e Investigación Epidemiológica, Instituto Mexicano del Seguro Social, México

peak at 2–5 years of age was reported for England after 1920, for the USA in 1940, and for Japan in 1960; during this time, this same peak was demonstrated for African-American with ALL in the USA (Greaves et al., 1993; Pratt et al., 1988; Ramot & Magrath, 1982).

We analyzed the immunophenotype of all cases of ALL registered in eight of the nine public hospitals, located in Mexico City, which attended children with ALL from January 2010 to May 2011. We showed that, for the 320 cases registered, the frequency of childhood ALL in Mexico City had two age peaks (Fig. 1), in agreement with data reported previously (Bernáldez et al., 2008; Mejía Arangure et al., 2005a). In Brazil, two peaks in ALL have also been reported (de Souza Reis et al., 2011).

Fig. 1. Immunophenotypes in children from Mexico City during 2010-2011. The children with acute lymphoblastic leukemia have two age peaks, specially the children with Pre B cell immunophenotype.

The aim of this chapter is to present different hypotheses related to the age peak of the ALL in children. We briefly describe the ontology of B and T lymphocytes and discuss an apparent association between the genetic rearrangements involved and the age peak in leukemias. We also review data on the age peak of ALL in different countries. At the end of the chapter, we propose an hypothesis with respect to the age peak in Latin American populations and in general.

2. B- and T-cell development

After birth and throughout life, development and replenishment of lymphoid cells is a highly ordered process that starts in bone marrow (BM) with the differentiation of multipotential hematopoietic stem cells (HSC). In this multi-step process, multiple alternate lineage potentials are gradually lost and lineage commitment is coincident with gain of specialized functions. Over the last few years, remarkable advances have been made in characterizing the primitive progenitors that initiate the lymphoid program; the patterns of transcriptional activity that control decisions about the fate of early lineages and late maturation process; and the environmental signals that influence the differentiation pathway during normal hematopoiesis. Malignant early lymphoid development is currently under intense investigation, and the existence of leukemic stem cells is still a matter of debate. However, identification of cytogenetic abnormalities in cells lacking lineage markers

and the unsuspected genetic diversity within individuals strongly suggests the participation of primitive cells in this disease.

2.1 From stem cells to committed B-cell progenitors

Because mature blood cells have finite life spans, they are constantly replaced from a unique cell population of HSC that resides in specialized niches in the BM (Baba et al., 2004). Within the hematopoietic system, HSC are the only cells with the ability to extensively give rise to identical daughter stem cells (self-renewal) and to differentiate into all functional blood cells (multipotency) (Seita & Weissman, 2010). The hematopoietic system is organized as a hierarchy of cell types with differing capacities for self-renewal, proliferation, and differentiation. In the pathways of this system, developing lymphoid or myeloid cells progress through critical stages of differentiation of multipotential and self-renewing HSC to multipotential early progenitors, which give rise to oligopotent progenitors. Downstream, the production of lineage-committed precursors is crucial for maturation of blood cells (Fig. 2). The lymphoid lineage consists of B, T, and natural killer (NK) cells; the myeloid lineage includes granulocytes, monocytes, macrophages, erythrocytes, megakaryocytes, and mast cells. Dendritic cells (DCs) are generated starting from the pathways of lymphoid or myeloid differentiation.

Fig. 2. Hematopoietic development in humans. The self-renewing hematopoietic stem cell (HSC) gives rise to multipotent progenitors (MPP), which have the ability to differentiate into common myeloid progenitors (CMP), and into multi-lymphoid progenitors (MLP). Mature blood cells are produced from lineage-committed precursors. The hierarchical structure of the hematopoietic system is shown. GMP, granulocyte and monocyte progenitors; MEP, megakaryocyte and erythrocyte progenitors; MDP, monocyte and dendritic cell progenitors; NK, natural killer cells; DC, dendritic cell; Mac, macrophage; Gran, granulocyte; Plat, platelets; Eryt, erythrocyte.

Current knowledge about the development of the lymphoid system is based, in great part, on the work done in mouse models, which has demonstrated that this differentiation program begins in BM in the fractions of lymphoid-primed multipotent progenitors (LMPP) and early lymphoid progenitors (ELP) capable of differentiating into T, B, NK, and conventional dendritic cells (Pelayo et al., 2005a, 2006a; Welner et al., 2008b). ELP also produce plasmacytoid dendritic cells (pDC), and interferon-producing killer dendritic cells (IKDC), both of which are key components of the innate immune response to infections (Pelayo et al., 2005b; Welner et al., 2007). The more differentiated common lymphoid progenitors (CLP) have substantially lost the possibility of differentiating into multiple lineages and efficiently generate B- and NK-cell precursors. Interestingly, HSC and early progenitors proliferate in response to systemic infection and replenish innate immune cells by using mechanisms that apparently involve interferons, tumor necrosis factor- α , and Toll-like receptors (TLR)—these last recognize viral/bacterial components. Thus, plasticity in primitive progenitor cells is sensitive to extrinsic agents that can modify differentiation fates during infection, thus supporting the idea that the stages of restriction of hematopoietic lineages are less abrupt than previously thought (Baldridge et al., 2011; Welner et al., 2008a). In humans, the early hematopoietic progenitors are confined in the BM to a cellular compartment that expresses CD34 (Bloom & Spits, 2006). This fraction of multipotent stem cells is characterized by the phenotype Lin-CD34⁺CD38^{-/lo}CD10⁺CD45RA⁻, whereas that of the earliest lymphoid progenitors is characterized by the phenotype Lin-CD34⁺CD38^{-/lo}CD45RA⁺CD10⁺ and has been recently designated as multi-lymphoid progenitor (MLP) (Doulatov et al., 2010). While a description that fully matches the definition of mouse ELP is still missing, Lin-CD34⁺CD38⁺CD45RA⁺CD10⁺ cells, which differentiate principally into B and NK cells, are considered the counterparts of mouse CLP (Fig. 2). Downstream, the sequential differentiation of lineage-restricted precursors generates Pro-B, Pre-B, and immature-B cells that eventually are exported to the peripheral lymphoid tissues. As in mice, human lymphopoiesis can undergo adjustments in cell-fate decisions under inflammatory conditions or during infections (Baldridge et al., 2011; Kim et al., 2005).

2.1.1 Biological differences between early hematopoiesis in neonatal and in adult

Some properties in hematopoietic development, including cell-cycle progression, transcription-factor networks, and growth-factor production, show substantial differences between newborns and adults (Mayani, 2010; Nguyen et al., 2010; Pelayo et al., 2006b).

During fetal and neonatal development, the hematopoietic system faces a complex set of demands, including rapid population turnover, protection against infection, and avoidance of harmful inflammatory immune responses (Levy, 2007; Pelayo et al., 2006b). After birth, there is an age-dependent maturation of the immune response; apparently, prenatal and postnatal exposure to microbial components or products may accelerate this maturation process (Levy, 2007). In response to TLR agonists, production of cytokines and chemokines is poor in early life but increases with age, though it is still limited in one-year-old infants, suggesting that the first year of life represents a critical period for the acquisition of competence by developing hematopoietic cells (Nguyen et al., 2010).

Moreover, hematopoietic stem and progenitor cells must divide extensively to generate the billions of new blood cells needed each day. Yet, little is known about the period when the expansion takes place or about the mechanisms responsible for the biological differences between fetal and adult primitive cells. However, it is widely recognized that fetal- and

neonatal-derived HSC and lymphoid progenitors possess higher proliferation and expansion potentials. Furthermore, during rebound from chemotherapy, adult progenitors acquire some, but not all, of the characteristics of fetal cells (Mayani, 2010; Pelayo et al., 2006b). Telomere dynamics, key cell-cycle mediators, and differential gene expression may account for the fetal/adult disparity. Whether the potential for high proliferation during early life makes hematopoietic precursor cells more vulnerable to gene mutations is not as yet clear, but its implication in the onset of neoplastic hematological diseases in neonates might be decisive.

2.1.2 Early steps in malignant lymphoid development

ALL is characterized by the monoclonal/oligoclonal proliferation of hematopoietic precursor cells of the lymphoid series within the BM. Although in recent years studies have reported important advances in the investigation of genetic, molecular, karyotypic, and phenotypic abnormalities that are prevalent in this disorder, the understanding of the mechanisms that damage the earliest program of lymphoid development remains poor. Results from cell cultures for early hematopoiesis, detection of specific leukemic karyotypes in primitive CD34⁺ cells, and data showing that cells with immature phenotypes are capable of engrafting and reconstituting leukemia in immunodeficient mice suggest that infant B cell-leukemia initiating cells have primitive characteristics (Cobaleda et al., 2000; Cox et al., 2004, 2009; Espinoza-Hernández et al., 2001) and are subject to intrinsic and extrinsic stimuli that could trigger lineage instability. On the other hand, leukemic blasts can also completely re-establish leukemic phenotypes *in vivo*, conferring them with stem-cell properties (Heidenreich & Vormoor, 2009). These conflicting results reveal that key questions regarding leukemic stem cells and lymphoid development in ALL remain unsolved. Recently, the combination of clonal studies, alterations in genetic copies, and xenotransplantation models have shown unsuspected genetic diversity, supporting the idea of multi-clonal evolution of leukemogenesis, rather than that of lineal succession (Dick, 2008). Future progress in this area will ultimately lead to an understanding of the biology of leukemia, as well as behavior and prognosis among individual groups.

2.2 Murine B-cell maturation in bone marrow

The first cells that exhibit commitment to the B-cell lineage in mice (the model in which this process is currently better understood) are B220⁺, CD19⁺, and CD43⁺ pro-B cells (Allman et al., 1999). During B-cell development, the main goal is to generate functional cell populations that are capable of expressing a diverse repertoire of B-cell antigen receptors (BCR), with each clone having specificity to recognize and counteract a new or recurrent pathogen. Antigen recognition is mediated by a heterodimer of immunoglobulin (Ig) heavy chains and light chains and signaling for an antigen-induced B-cell response is mediated by the molecules Igα (CD79a) and Igβ (CD79b). Igα and Igβ exist as a disulfide-coupled heterodimer in non-covalent association with the Ig antigen-recognition element. Igα/Igβ signaling is dependent upon distinct motifs localized in the cytoplasmic tails of these proteins, the immunoreceptor tyrosine-based activation motifs (ITAM) (Cambier, 1995; Fuentes-Pananá et al., 2004b). The BCR can generate signals that lead to a variety of outcomes, depending upon the developmental stage of the B cell and the degree and persistence of receptor aggregation. Although the mature form of the BCR is present only in immature and mature B cells, genetic and biochemical studies have shown that various

forms of the BCR are expressed at defined stages of B-cell development that are required for progression of B cells through several defined developmental checkpoints (Fuentes-Pananá & Monroe 2001; Fuentes-Pananá et al., 2004a).

In the pro-B stage in which the heavy chain is in the process of recombination, the signaling proteins Ig α and Ig β are expressed on the cell surface and are associated with the protein calnexin, and perhaps other as yet unknown proteins, forming the so-called "pro-BCR" (Nagata et al., 1997). Pro-B cells that successfully produce an Ig heavy-chain protein transport it to the surface in association with Ig α /Ig β and with the surrogate light-chain complex that is composed of $\lambda 5$ and V pre-B (the pre-BCR receptor) (Karasuyama et al., 1996; Melchers et al., 1993). Surface expression of this receptor marks the transition to the pre-B stage, a step that is marked by loss of CD43 and gain of CD22 expression, by allelic exclusion of the un-recombined heavy-chain allele, and by the opening and initiation of recombination at the λ Ig light chain loci (Rolink & Melchers, 1993). A similar mechanism operates in the pre-B stage in which the successful recombination of light chain and its pairing with the Ig heavy chain and Ig α and Ig β allows the assembly of the mature form of the BCR and marks the transition to the immature stage. Intimate contact between the immature-B cell and the stromal cells of the BM allows those receptors capable of recognizing self-antigens to be identified and eliminated through a variety of mechanisms collectively termed "tolerance". Non-self-reactive B cells exit to the periphery and reach the spleen where they are again tested for reactivity against self-antigens before they transition to the mature stage (von Boehmer & Melcher, 2010).

2.2.1 Heavy and light chain VDJ recombination and positive selection in BM

The main goal during the pro-B and pre-B stages is to generate a signaling-competent antigen receptor, with the progression of development conditioned by the signaling capacity of the receptor complex. The BCR Ig heavy and light chain genes are composed of constant and variable domains, the recombination of which makes BCR clonal diversity possible. The variable domain is formed by a series of segments termed variable (V), diversity (D), and joining (J) (Fig. 3A), which are brought together by a site-specific recombination process termed VDJ recombination. This is a highly ordered process during which the D and J fragments are rearranged and the V segment is then joined to the DJ fragment; these steps occur first in the heavy and then in the light chain loci (Fig. 3B) (Fuentes-Pananá et al., 2004b; Thomas et al., 2009). This process is essential for the adaptive immune function of the lymphocyte. As mentioned above, the pro-B stage is characterized by rearrangement of the Ig heavy chain and it is further divided according to the status of the rearrangement; in mice, Marshall named these sub-stages proB-A (during which the heavy chain is in the germ-line state), proB-B (during which D and J are recombined), and proB-C (during which V-DJ is recombined) (Hardy et al., 1991). In the pre-B stage, the light chain V and J fragments are recombined, first in the λ and then in the κ loci (Fig. 3A).

Unsuccessful recombination (e.g., heavy chain VDJ fragments that are not in a proper reading frame) or unsuccessful pairing of the pre-BCR components results in pro-B cells that are unable to proceed to the pre-B stage, thus leading the developing pro-B cell to apoptosis. This observation supports the concept of an active signaling role for the pre-BCR in generating the permissive signal that allows differentiation through the pre-B stage (Bannish et al., 2001; Mandal et al., 2009; Yasuda et al., 2008). A similar mechanism is thought to operate for the transition from pre-B to immature cells, which requires expression of the

mature form of the BCR (Bannish et al., 2001; Mandal et al., 2009; Yasuda et al., 2008). In addition to their VDJ recombination status, all pro-B and pre-B stages can be recognized by their patterns of surface-marker expression (Hardy et al., 1991).

2.3 T-cell development

T-cell development occurs in the thymus through equivalent processes of VDJ recombination, selection against self-reactive clones, and maturation into fully functional clones (Figure 3B). Expression of the T-cell antigen receptor (TCR) on the surface of the T cell marks the transition through all developmental stages. The TCR is a complex of proteins that may be functionally divided into the antigen-binding unit formed by the β and α chains (also in a smaller fraction of T cells, the γ and δ chains) and the signaling unit formed by the CD3 chains. The β and α (and γ/δ) chains consist of constant and variable regions, with VDJ recombination taking place in the latter. There are different types of CD3 chains and each mature TCR is associated with six of them: two CD3 ϵ , two CD3 ζ , one CD3 γ , and one CD3 δ . The stages of T-cell development are known as the following: pro-T (CD3^{pos}CD44^{pos}CD4^{neg}CD8^{neg}; the TCR β chain is rearranged); pre-T (CD3^{pos}CD44^{pos}CD25^{pos}CD4^{neg}CD8^{neg}; the TCR α chain is rearranged); CD4 and CD8 double-positive T cells [CD3^{pos}CD4^{pos}CD8^{pos}; equivalent to immature/transitional B cells (the stage in which tolerance mechanisms are applied to the developing T cell)]; and CD4 or CD8 single-positive cells (mature helper or cytotoxic T cells). Pro-T and pre-T cells are further subdivided according to the status of the β and α chains rearrangements: DN1 (β chain is in germ-line configuration); DN2 (D and J rearrangement of the β chain); DN3 (V joins to DJ of the β chain); and DN4 (V and J rearrangement of the α chain). DN1, DN2, and DN3 are pro-T stage; DN4 is a pre-T stage. The progression of recombination-associated development is illustrated in Fig. 3B. Expression of pro-TCR, pre-TCR, and mature-TCR receptors mark the progression through T-cell developmental checkpoints, homologous to the positive- and negative-selection mechanisms for B cells (Love et al., 2010; Wiest, 1994, 1995).

2.4 Human B- and T-cell development

Selection processes operating on developing B and T cells of all mammals are the same; thus, B- and T-cell development in human is also mainly divided by the process of VDJ recombination and expression of antigen-recognition complexes on the cell surface and by the proliferative expansion of clones that have successfully completed the rearrangement of their receptors (Blom & Spits, 2006). However, for the human, all these processes are less well understood than are their counterparts in the murine model. Importantly in human lymphocyte development, the pro-B and pre-B sub-stages are the ones generally found to be compromised in human pediatric B-cell ALL. B-cell ALL is characterized by cells unable to progress along the differentiation pathway. The stages more often compromised are: proB-A (before heavy-chain recombination), proB-C (after heavy-chain recombination), and large pre-B stage (before light-chain rearrangement) (Fig. 3B) (Hardy et al., 1991; Rolink et al., 1991). These stages are better known in humans as early proB or pre-proB (A); proB (B); preB1 (C) (proB stage); and preB2 (large preB stage) (Fig. 3B). All these stages in human B and T cells can also be recognized by the expression of specific surface markers, a characteristic that has helped to classify the different types of pediatric ALL. In humans, B cells are recognized by the expression of CD19 and CD10; T cells by CD2, CD3, CD5, and

CD7; and, for the proliferating immature-B cells, pre-proB cells by CD34⁺CD19⁻CD10⁺; preB1 by CD34⁺CD19⁻CD10⁺; and preB2 by CD34⁺CD19⁺CD10⁺. Immature T cells are mainly recognized by the lack of expression of CD4 and CD8 and expression of T-cell markers.

Fig. 3. B- and T-cell development. A) The Ig heavy and light chain genes are comprised of constant and variable domains; the variable domain is formed by an n number of segments termed variable (V), diversity (D), and joining (J) in the heavy chain and by segments V and J in the light chain. These segments are brought together by a site-specific recombination process, termed VDJ recombination, which is responsible for the extensive repertoire of BCR specificities. There are two loci for light chain, κ and λ . Here, all the loci are shown in germ-line configuration, prior to the process of VDJ recombination. B) All B- and T-cell stages can be divided according to the main processes guiding development: receptor assembly, tolerance, and activation. Receptor assembly stages (light blue box) in B and T cells are differentiated by the process of VDJ recombination in the heavy (IGH) and light (IGL) chains, which are recombined in the pro-B and pre-B stages, respectively (β and α rearrangement in pro-T and pre-T cells). The nomenclature of each sub-stage in the mouse model is shown in black letters, A-D (Marshall's) for B cells and DN1–4 for T cells; the most common human nomenclature is shown in red letters. The dashed lines separating all stages indicate checkpoints at which signaling from the preBCR and BCR is required for positive selection and progression along the B-cell maturation pathway. The proBCR, proTCR, preBCR, preTCR, and mature receptors are also illustrated in their respective stages.

3. Incidence peak of acute leukemia with specific cytogenetic aberrations in childhood

ALL is a clonal disease driven by mutations. The incidence of leukemia among children varies considerably with age. Age at diagnosis has played a relevant role in the epidemiology of these diseases and in the determination of risk groups and treatment stratification (Smith et al., 1997). The age-related incidence pattern of ALL has been established by the demonstration of the clone-specific 11q23-, t(12;21)-, as determined by blood testing (Guthrie cards), and hyperdiploidy (Gale et al., 1997; Wiemels et al., 2002).

The majority of childhood ALL cases show the emergence *in utero* of pre-leukemic cells and, postnatally, rare interactions between the immune system and childhood infections (Greaves, 2002).

The incidence rates of childhood ALL vary from country to country; in Europe and the USA, 85–90% of the cases have a B-cell-precursor phenotype (pre-B) with an incidence peak in the 2–7 years age group. For childhood T-cell ALL, lower incidence peaks have been reported (Hjalgrim et al., 2003). Most cases of pre-B ALL carry either a high hyperdiploid karyotype or the chromosomal translocation t(12;21) (p13;q22). A high hyperdiploid karyotype, which is present in 30% of pediatric patients, shows a modal chromosome number above 50, which involve trisomies of chromosomes X, 4, 6, 10, 14, 17, 18, and 21 and which, in most cases, arise by simultaneous chromosomal gain in a single abnormal mitosis (Paulsson et al., 2005). The chromosomal translocation t(12;21) (p13;q22) is found in 20–25% of cases and involves the gene *RUNX1* in chromosome region 21q22 and the gene *ETV6* in chromosome region 12p13. Both encode transcriptional factors that are essential for normal fetal hematopoiesis (Romana et al., 1995). High-hyperdiploid leukemic clones are sometimes missed by standard G-band karyotyping due to poor metaphase quality. The detection rates can be raised by extended fluorescence in situ hybridization (FISH) analyses, flow cytometry, DNA-index analyses, high-resolution comparative genomic hybridization (CGH), and comparative genomic array CHG (Kristensen et al., 2003; Nyggard et al., 2006). Translocation t(12;21) is cryptic by standard G-band karyotyping; its diagnosis is based on FISH analyses and reverse transcriptase-polymerase chain reaction (RT-PCR) (Nordgren, 2003).

Recent studies from twins with concordant leukemia suggest that a genetic event is involved in these acute leukemias: chromosomal translocations can have a prenatal origin (Ford, 1993, 1997, 1998; Gill et al., 1994; Wiemels et al., 1999a, 1999b). There is also indirect support that the prenatal origin of leukemic clones is derived from the presence of clonotypic rearrangements at the IGH and TCR loci in Guthrie spots (Fasching et al., 2000; Yagi et al., 2000). Because most of the studies have included a limited subset of leukemias, primarily those of lymphoid phenotypes, it remains an open question as to whether the prenatal origin of childhood leukemia is a possible, or a common, occurrence in other subgroups.

The most frequent translocation both in children (1–20 years of age) and in adults (older than 20 years) is t(8;21) AML1-ETO, which results in a fusion protein that disrupts the normal function of the transcription factors AML1 and ETO (Finnette et al., 1996). The incidence of leukemia with the t(8;21) AML1-ETO translocation increases slowly during childhood and is constant thereafter (Downing, 1999).

MLL/11q23 translocations are known to be involved in the leukemia of infancy, but they can also be found in older children and sometimes in the elderly (Johansson et al., 1998). MLL/11q23 translocations, of which t(11;19)(q23;p13) and t(4;11)(q21;q23) are the most

common, have an incident peak during first year of life; 11q23-aberrant leukemias are the second most common malignancy. When these leukemias appear in very early infancy, their prenatal origin is proven; in monozygous twins (with identical MLL-rearrangements), the concordance rate is almost 100%. This suggests that, when these mutations are present at birth, they invariably lead to overt leukemia (Greaves et al., 2003). It is not known when MLL/11q23 translocations occur in adults.

Childhood acute leukemias are clinically related, but cytogenetically different, diseases; therefore, cytogenetic subgrouping is important to understand the diversity in their etiology, natural history, and epidemiology. An understanding of the epidemiology of childhood acute leukemia is relevant in order to clarify the cytogenetic backbone of etiologic research.

4. Infectious etiology of childhood leukemia

The hypothesis concerning an infectious etiology of childhood leukemia was first raised several decades ago (Cooke, 1942). Although, to date, it has not been possible to demonstrate the involvement of one or more particular infectious agent(s), different hypotheses are still in force based on epidemiological and demographic studies that have yielded evidence about the possible participation of infectious agents in childhood leukemogenesis. Independently, Greaves (1988), Kinlen (1988), and Smith (1997) have suggested different mechanisms by which certain events of infection may explain at least some cases of childhood leukemia.

4.1 Greaves' hypothesis

More than 20 years ago, Greaves (1988) proposed the involvement of infectious agents in the etiology of childhood leukemia. In the most recent version, his hypothesis includes a model of the natural history of ALL with a minimum of two oncogenic hits (Greaves, 2002), whereby a first hit occurs *in utero* and forms a pre-leukemic clone that requires at least one secondary, postnatal carcinogenic hit to unleash the malignant transformation. According to this hypothesis of "delayed infection", the second oncogenic event could be indirectly promoted by delayed exposure to an infectious agent that causes an uncontrolled immune response and promotes the malignant, abnormal proliferation of the pre-leukemic clone.

A number of studies have shown that first-born children have a higher risk of developing leukemia. Only children and those children who do not attend daycare have been assumed to be evidence of the hypothesis, in that, by having less contact with others children and potentially less frequency of exposure to infectious agents, the probability of developing the disease increased, when compared to children outside these groups (Dockerty et al., 2001; Infante-Rivard et al., 2000; Jourdan-Da Silva et al., 2004; Perrillat, et al., 2002; Petridou et al., 1993),

Nevertheless, some research has not provided absolute support for Greaves' hypothesis: data from the Northern California study group (Ma et al., 2005) indicated that daycare attendance and ear infections during infancy are associated with a lower risk of ALL; however, this was true only for non-Hispanic children.

It should be noted that Greaves' hypothesis concerns the common form of ALL (B-cell precursor). This form comprises most of the ALL cases that peak at 2–5 years of age and that are seen in developed countries or in affluent communities that have improved their living

standards and have become "more hygienic". Therefore, any relevant infection should occur in the first two years of life. Through comparison of international reports, variations in the peaks of childhood ALL have been identified. The aforementioned peak at 2–5 years of age is reduced, or even absent, for Black Africans and for other developing communities (Court & Doll, 1961; Hewitt, 1955; Ramot & Magrath, 1982, as cited in Chan et al., 2002). In Mexico, for example, two peaks have been reported; the first occurring at 2–3 years of age and the second at 6–9 (Bernaldez-Rios et al., 2008). A recent epidemiological study done in Mexico City showed that severe infections, occurring in the first year of life and requiring hospitalization, were associated with a higher risk of developing leukemia for children with Down syndrome (Flores-Lujano et al., 2009).

A possible interpretation of these results showing that infections in the first year of life represent risk factors (Cardwell et al., 2008; Roman et al., 2007) is that such infections could play a role as promoters, not as protectors, in the genesis of leukemia. In principle, this could mean that, for these groups, the infectious etiology of leukemia might be different than that proposed by Greaves.

4.2 Kinlen's hypothesis

When clusters of childhood leukemia and non-Hodgkin lymphoma were observed in the early 1980's near nuclear plants in Cumbria, England (Black, 1984), and at Dounreay, Scotland (Heasman et al., 1986), it was thought that such increase could have been a consequence of radioactive contamination by the nuclear plants, which might have caused somatic or germinal line mutations that produce cancer. However, there was no evidence of radioactive leaks (Committee on Medical Aspects of Radiation in the Environment [COMARE], 1986) or of epidemiological associations that showed greater occupational exposures of the parents (Gardner et al., 1990).

Kinlen proposed that the observed clusters of leukemia could have resulted from the unusual population mixing that occurred due to migration of workers and their families, who had travelled to work in the nuclear plants; he also hypothesized that a common, but unidentified, infectious agent could have been involved (Kinlen, 1988, 1995a). According to his proposal, cases of childhood leukemia would be a rare response from the isolated, uninfected individuals who were, therefore, susceptible to a putative infectious agent carried by these newcomers.

By extrapolating from animal leukemias caused by virus, a subtype of leukemia in adults, and by considering that childhood leukemia is not a contagious disease, Kinlen proposed that the agent involved could be a common virus causing a non-common response (Kinlen, 2011).

To date, Kinlen's working group has directed several studies, mainly in England and Scotland, and has observed a significant increase in cases of childhood leukemia where large-scale mixing between rural and urban populations occurred with unusual patterns of contact in the different communities (Kinlen et al., 1990, 1993, 1995b; Kinlen & Balkwill, 2001; Kinlen & Hudson, 1991; Kinlen & John, 1994).

Kinlen first proved his population-mixing hypothesis in the infectious etiology of childhood leukemia in Scotland, in a rural area that had received large influxes of people: the results showed a significant increase in the leukemia cases in children under five years of age (Kinlen, 1988). Another of his studies showed that excess of leukemia was higher in the 0–4 years age range; it was predominantly higher for children younger than 1 year, suggesting

an infection during pregnancy (Kinlen & Hudson, 1991). From his observations, Kinlen proposed that, during population influx, adults are the main transmitters of an infectious agent and that a rare response was more likely to be made by a naive immune system; thus, in the foregoing case, population mixing could be responsible for the leukemia cases seen, even in the first year of life. Nonetheless, the author did not associate the leukemia clusters with a particular subtype of disease and interpreted that any type of childhood and infant leukemia might have a common cause.

4.3 Smith's hypothesis

A third hypothesis regarding the infectious etiology of childhood leukemia was proposed by Smith (1997). He hypothesized that some cases of ALL observed in the 2–5 years age group (B-cell precursor) could be the result of an infection that occurred during pregnancy and that was transmitted from mother to fetus.

Unlike Greaves, who postulated that infections are secondary etiological factors acting indirectly, Smith suggested that infection is one of the first hits in leukemogenesis and that a related infectious agent acts directly, i.e., the agent is able to infect the immature-B cell and to promote genomic instability, thereby leading to the transforming process. According to his model, the agent involved must be able to cross the placenta and infect the fetus without causing serious abnormalities, have limited oncogenic capacity, and produce minimal primary symptoms (Smith, 1997).

Studies that have shown that maternal infections are associated with an increased risk of ALL support this model. Lehtinen et al. (2003) showed that maternal infection with Epstein-Barr Virus (EBV) was associated with a significantly increased risk of ALL. Naumberg's work group found a similar association when the mother had lower genital tract infections (Naumberg, 2002). However, other studies have shown little or no association between infections, either by varicella or influenza during pregnancy (Little, 1999, as cited in McNally 2004), by unspecified infection in pregnancy (McKinney et al., 1999), or by recurrent maternal infections (Infante-Rivard et al., 2000), and subsequent childhood leukemia in their offspring.

Molecular screening for viruses in leukemic cells has been used as a distinct approach to explore possible mechanism(s) of direct transformation occurring *in utero* or during infancy. Smith postulated that JC polyomavirus (polyomavirus are named from the initials of the patients from whom the first isolates were made) would be a good candidate, because it met all the conditions for his model. However, to date, viral sequences from the polyomaviruses, JC or BK (a closely related virus), have not been found in leukemic cells (MacKenzie, 1999). A number of viruses have been analyzed by different researchers. When blood or BM samples from patients diagnosed with childhood ALL were screened for members of the human herpesvirus family (HHV4, HHV6, HHV7, and HHV8), no evidence of the presence of these viruses in leukemic cells was found (MacKenzie, et al., 2001). Screenings for bovine leukemia virus (BLV) and transfusion-transmitted virus (TTV) have also been negative (Bender et al., 1988; Shiramizu et al., 2002). However, as the list of candidates has not been exhausted, massive sequencing to analyze non-human genomic components from leukemic cells, even those not previously identified, is an attractive approach.

There is evidence supporting each of the hypotheses presented here. Although the postulated mechanisms differ from each other, such hypotheses are not mutually exclusive. Because the notion of a unique carcinogenic factor is a biologically absurd assumption, the

various cases of leukemia in Mexico and around the world must be triggered by a variety of agents. Given the evidence of an infectious etiology involved in childhood leukemia, the identification of this agent, or agents, remains a scientific challenge.

5. Age peak of ALL in the world: epidemiologic aspects

Industrialization, urbanization, economic growth, and technological development have brought benefits to people, making everyday tasks easier and offering accessibility to services. However, modernization and industrialization have also led to changes in the environment and in lifestyle, which have changed the pattern of diseases in adults and children over the last century. Children now face predominantly chronic and disabling diseases such as obesity, diabetes, asthma, learning disabilities, birth defects, and cancer (Suk et al., 2003).

Internationally, published studies have reported lower incidence rates of ALL in Africa, Asia, and Vietnam, with the high rates found for Hong Kong, United Kingdom (UK), USA, and Japan, thus suggesting a correlation between the reported variations in incidence of ALL and socio-economic status (Parkin, et al., 1988, 1998; Stiller, 2004). However, the highest incidence rates of ALL have been reported for Costa Rica, Mexico City, and the Hispanic populations of Los Angeles, California and of Florida in the USA. (American Cancer Society, Surveillance Research [ACSSR], 2009; Glazer et al., 1999; Mejía-Aranguré et al., 2005b; Monge et al., 2002; Surveillance Epidemiology and End Results [SEER], 2008; Wilkinson et al., 2001). Of all cases diagnosed with ALL in the 2–7 years age group in Europe and the USA, approximately 85–90% had a precursor-B (pre-B) cell immunophenotype; the remainder were classified as having either T-cell or B-cell immunophenotype (14% or 2%, respectively (Pui, 1997). Reports from the USA indicate that African-American children have half the risk of developing ALL than do White children, with a peak age of presentation of ALL between 2–6 years old (median 4 years old) and with ALL predominating in males (male:female ratio of 1.2: 1) (Eden, 2010).

Reports regarding the incidence and trend for ALL from the Surveillance Epidemiology and End Results (SEER) in the USA and the Manchester Children's Tumor Registry (MCTR) in the UK indicate that the rates have increased annually by 0.6% and 1.1%, respectively. However, recent reports indicate a modest increase of 0.4% annually for the USA (Linabery & Ross 2008) and 1–4% for Europe (McNally et al., 2001) in the group of 1–4 years old.

Of particular note in reports of cancer registries are the high incidences of ALL in Costa Rica, Mexico City, and the Hispanic populations of the USA (Table 1). For the last case, it has been suggested that, because this population tends to live in more crowded conditions and poverty that does the White population, thereby exposing this population to infectious agents that have not as yet been identified but which may promote the development of acute leukemia (Yaris et al., 2004). There are marked variations in the incidence of ALL among populations worldwide; these variations can provide valuable clues to help understand the etiology of ALL.

A review of the latest articles published about the incidence of ALL in countries around the world (Table 2), some studies reported only the incidence of acute leukemia; nevertheless we included these studies because, as 80% of leukemias are ALL, the incidence rates reported provide very important information. We considered only those works that reported incidence rate, age group, and when done, immunophenotype. In these studies, the

results were generally reported by age groups: <1 year; 1–4 years; 0–4 years; 5–9 years; 10+ years; or 0–14 years. Some studies did not estimate the overall incidence rate and reported only by sex. Few studies had a graphic concerning the incidence of ALL and age, which would have better illustrated the peak incidence. Most published studies are from the Americas or Europe; few studies determined the immunophenotype of acute leukemia.

Report	Period	Age group (years)	Incidence rate per 10 ⁶ children
United States SEER ¹ (Howlader et al., 2010)	2007	0–14	35.0
CDC (NPCR) ² (USCS ³ Working Group, 2010)	2003–2007	0–19	46.0
ACSSR ⁴ (American Cancer Society, 2009)	2002–2006	0–14	46.7
Florida (Wilkinson et al., 2001)	1985–1997	0–14	49.7
California (Glazer et al., 1999)	1988–1994	0–14	44.0
Mexico City IMSS ⁵ (Mejía-Aranguré et al., 2005b)	2006–2007	0–14	57.6
Costa Rica (Monge et al., 2002)	1981–1996	0–14	43.1

¹ Surveillance, Epidemiology and End Results Program in United States of America.
² Center for Disease Control, Division of Cancer Prevention and Control, National Program of Cancer Registries (NPCR).
³ United States Cancer Statistics
⁴ American Cancer Society, Surveillance Research.
⁵ Instituto Mexicano del Seguro Social

Table 1. Incidence rates of lymphoid leukemias per million Mexican and Costa Rican children from cancer registries.

For the Americas, a highest incidence rate of ALL, 73.2 per 10⁶ children in the 1–4 years age group, was reported for the USA for the period 1992–2004 (Linabery & Ross, 2008); the lowest incidence rate of ALL (35.5 per 10⁶ children) was in Uruguay (Castillo et al., 2001), this latter report covered a different period and did not report incidence rates for different age groups. It is very important to note that, for both Brazil (de Souza Reis R et al., 2011) and Mexico City (Bernáldez-Ríos et al., 2008), the highest peak incidence of ALL was reported for the same age groups, with high (albeit distinct) incidence rates in both countries. From these data, one can speculate that these two countries share certain genetic or environmental characteristics. Although in a report from Puerto Rico (Pérez-Perdomo & Rodríguez-Figueroa, 2000), there are data only for individuals <20 years, these data were included here to provide information, not for comparative purposes. For countries in Europe, the highest incidence rate was reported for Italy (95.6 per 10⁶) (Magnani et al., 2003) and Greece (82.5 per 10⁶) (Petridou et al., 2008), for different periods and immunophenotypes, but for the same age group. The lowest incidence rate (35.7 per 10⁶) was reported for Ireland (Stack et al., 2007).

The differences in trends across demographic subgroups provide starting points for the etiological investigation of ALL. Such research should generate testable hypotheses that include factors such as the child's birth characteristics and environmental exposures and topics such as the creation of cancer registries.

Location	Period	Type of cancer	Immuno-phenotype	Age group	Incidence rate ¹
The Americas					
North America					
USA					
SEER (Linabery & Ross, 2008)	1992–2004	ALL ²	ND ³	1–4	73.2
Hispanic children (Glazer et al., 1999)	1988–1994	ALL	ND	0–4	66.0
Hawaii, USA (Goodman et al., 1989)	1960–1984	ALL	ND	0–4	49.6 (M) 44.8 (F)
Mexico City (Bernáldez-Ríos et al.,	1996–2006	ALL	B-cell Precursor	3	65.0
				6	50.0
Mexico City (Mejía-Aranguré et al., 2005b)	1996–2000	ALL	ND	1–4	64.6
Ontario, Canada (Agha et al., 2006)	1986	AL ⁴	ND	0–14	52.8
	2001				44.4
Puerto Rico (Pérez-Perdomo & Rodríguez-Figueroa et al., 2000)	1980–1991	AL	ND	<20	35.7
Central America					
Costa Rica (Monge et al., 2002)	1981–1996	ALL	ND	1–4	68.5
El Salvador (Mejía-Aranguré et al., 2005b)	1996–2000	ALL	ND	1–4	48.4
South America					
Brazil (de Souza Reis et al., 2011)	1997–2004	ALL	ND	3	72.8
				6	40.0
Trinidad and Tobago (Bodkyn & Lalchandani, 2010)	2001–2006	AL	ND	0–4	49.0 (M) 27.0 (F)
Uruguay (Castillo et al., 2001)	1992–1994	ALL	ND	0–14	35.5
Asia					
Iran (Mousavi et al., 2010)	2003–2007	AL	ND	0–14	15.9 (M)
					14.1 (F)

Europe					
North Europe					
Nordic Countries (Hjalgrim et al., 2003)	1998–2001	ALL	B-cell precursor	0–14	35.9
West Europe					
Ireland (Stack et al., 2007)	1994–2000	ALL	ND	0–14	35.7
Northwest England (McNally et al., 2001)	1954–1998	ALL	ND	1–4	51.2
Northwest England (McNally et al., 2000)	1995–1998	ALL	B-cell precursor	1–4	54.0
Southwest England (McKinney et al., 1993)	1984–1988	ALL	Pre-B	1–4	50.0 (M) 54.0 (F)
Yorkshire, UK (Feltbower et al., 2001)	1995–1997	ALL	B-cell precursor	1–4	46.6
Central Europe					
Eastern Germany (Spix et al., 2008)	1991–2004	ALL	ND	0–14	44.0
Germany (Kaatsch, 2010)	1988–1997	AL	ND	0–14	44.0
South Europe					
Castilla and León, Spain (González García et al., 2010)	2003–2007	AL	ND	0–4	47.6
Greece (Petridou et al., 2008)	1996–2006	ALL	B-cell precursor	1–4	82.5
Northwest Italy (Magnani et al., 2003)	1995–1998	ALL	Common Pre-B	1–4	95.6
Spain (Peris-Bonet et al., 2010)	1983–2002	AL	ND	0–14	45.9
Oceania					
Australia (Baade et al., 2010)	1997–2006	ALL	ND	0–4	84.6

¹per 10⁶ million children
²Acute lymphoblastic leukemia
³No data
⁴Acute leukemia

Table 2. The incidence of AL and ALL and the peak age in different continents expressed in cases by million.

6. Conclusions

The ontology of B and T lymphocytes demonstrates that different factors may be involved at the moment when the cell is most vulnerable to undergo an alteration that promotes the development of ALL. An important factor may be an infectious agent(s). For children with

ALL, both the presence of MLL/AF4, principally during the first year of life, and the presence of ETV6/RUNX1 in 2–6 year olds are considered causes of ALL. MLL/AF4 is sufficient to promote the development of ALL and that children born with this genetic rearrangement will always develop ALL. However, in the case of children born with the rearrangement ETV6/RUNX1, one or more additional risk factors (e.g., an exaggerated response to late common infections) would be needed to initiate leukemogenesis. It is possible that the age peak of ALL reflects a higher degree of susceptibility with which a child is born, such as is the case with both bilateral retinoblastoma and bilateral Wilm's tumor: an earlier age peak is found in the bilateral forms of these diseases than when the retinoblastoma or Wilm's tumor is unilateral (National Wilm's Tumor Study Committee, 1999; Pastore et al., 1988; Sanders et al., 1988; Teppo et al., 1975). Thus, children with the highest susceptibility at birth should show an early age peak for ALL (say, during the firsts year of life), whereas children born with the lowest susceptibility will evidence an age peak of ALL in later years. For those children for whom it is not possible to determine their susceptibility to ALL at birth, it may be that many environmental factors are involved when the child develops ALL during the firsts year of life. That is to say, in order for a child to develop ALL, both risk factors (susceptibility at birth or degree of exposure) must be present and at least one of the two risks factors must be high. Thus, to develop ALL, a child born with a high susceptibility for developing the disease may need only a low degree of exposure to environmental factor(s), whereas a child born with low susceptibility for developing ALL will need a higher degree of exposure to environmental factor(s) (Figure 4).

Fig. 4. Interaction between a gradient of susceptibility and a gradient of exposure to carcinogenic environmental factors. From this, to develop acute leukemia, an individual with a higher susceptibility, as determined by the interplay of genetic factors, would need a lower exposure, as determined by the interplay (unknown, possibly synergistic) of the characteristics of the exposure. Conversely, the higher the exposure, the lower the susceptibility needed to result in developing the disease.

It is possible that the two age peaks for ALL reported for children from Brazil and Mexico City reflect different etiologies, or are a result of each child's having a different etiology. It is probable that there is no one genetic rearrangement that determines susceptibility to ALL, nor one environmental factor associated with ALL. However, if the effect(s) of exposure to

environmental factor(s) is(are) cumulative and if the degree of exposure to environmental factor(s) perfectly complements the degree of susceptibility with which the child was born, then the child develops ALL. The age peak of ALL permits the prediction of the degree of susceptibility with which the child was born and/or the degree of exposure to environmental factor(s) experienced by the child during the first years of life.

7. Acknowledgments

This work was supported by a grant (FIS/IMSS/PROT/056) from the Fondo de Investigacion en Salud, Mexico, and of projects financed by grants (2007-1-18 71223/FIS/IMSS/PROT/592 & CB-2007-1-83949/FIS/IMSS/PROT/616) from the Consejo Nacional de Ciencia y Tecnologia, Mexico, through its division for the dissemination of the results of scientific investigation. We thank Veronica Yakoleff for reviewing of the original manuscript, editorial revision, and for helpful comments. Janet Flores-Lujano did the figure 1.

8. References

- Agha, M., DiMonte, B., Greenberg, M., Greenberg, C., Barr, R. & McLaughlin JR. (2006). Incidence trends and projections for childhood cancer in Ontario. *International Journal of Cancer*, Vol. 118, No. 11, (June 2006), pp. 2809-2815, ISSN 1097-0215.
- Allman, D., Li, J. & Hardy, RR. (1999). Commitment to the B lymphoid lineage occurs before DH-JH recombination. *Journal of Experimental Medicine*, Vol.189, No.4, (February 1999), pp. 735-740, ISSN 0022-100.
- American Cancer Society (2009). Cancer in children and adolescents, In: Cancer Facts & Figures for Hispanics/Latinos 2009-2011, Accessed on Jun 8th 2011, Available from: <<http://www.cancer.org/acs/groups/content/@nho/documents/document/ffhispanicslatinos20092011.pdf>>
- Baade, PD., Youlten, DR., Valery, PC., Hassall, T., Ward, L., Green, AC. & Aitken, JF. (2010). Trends in incidence of childhood cancer in Australia, 1983-2006. *British Journal of Cancer*, Vol.102, No.3, (February 2010), pp. 620-626, ISSN 1532-1827.
- Baba, Y., Pelayo, R. & Kincade, PW. (2004). Relationships between hematopoietic stem cells and lymphocyte progenitors. *Trends in Immunology*, Vol.25, No.12, (December 2004), pp. 645-649, ISSN 1471-4906.
- Baldrige, MT., King, KY. & Goodell, MA. (2011). Inflammatory signals regulate hematopoietic stem cells. *Trends in Immunology*, Vol.32, No.2, (February 2011), pp.57-65, ISSN 1471-4906.
- Bannish, G., Fuentes-Pananá, EM., Cambier, JC., Pear, WS. & Monroe, JG. (2001). Ligand-independent signaling functions for the B lymphocyte antigen receptor and their role in positive selection during B lymphopoiesis. *Journal of Experimental Medicine*, Vol.194, No.11 (December 2001), pp. 1583-1596, ISSN 0022-100.
- Bathia, S. & Robison, L. (2003). Epidemiology of leukemia in childhood, In: *Hematology of infancy and childhood*, Nathan, DG., Oski, FA. Eds., pp. (1081-1100), Saunders, ISBN 978-1-4160-9977-2, Philadelphia.
- Bender, AP., Robison, LL., Kashmiri, SV., McClain, KL., Woods, WG., Smithson, WA., Heyn, R., Finlay, J., Schuman, LM., Renier, C. & Gibson, R. (1988). No involvement of bovine leukemia virus in childhood acute lymphoblastic leukemia and non-

- Hodgkin's lymphoma. *Cancer Research*. No.48, Vol.10, (May 1988), pp. 2919-2922, ISSN 0008-5472.
- Bernaldez-Rios, R., Ortega-Alvarez, MC., Perez-Saldivar, ML., Alatoma-Medina, NE., Del Campo-Martinez, Mde L., Rodriguez-Zepeda, Mdel C., Montero-Ponce, I., Franco-Ornelas, S., Fernandez-Castillo, G., Nuñez-Villegas, NN., Taboada-Flores, MA., Flores-Lujano, J., Argüelles-Sanchez, ME., Juarez-Ocaña, S., Fajardo-Gutierrez, A. & Mejia-Arangure, JM. (2008). The age incidence of childhood B-cell precursor acute lymphoblastic leukemia in Mexico City. *Journal of Pediatric Hematology/ Oncology*, Vol. 30, No. 3, (March 2008), pp. 199-203, ISSN 1536-3678.
- Birch, J. & Blair, V. (1992). The epidemiology of infant cancers. *British Journal of Cancer*; Vol. 66, No. Suppl. 18, (August 1992), pp. 2-4, ISSN 0007-0920.
- Black, D. (1984). Investigation of the Possible Increased Incidence of Cancer in West Cumbria, London. *Proceedings of Report of the Independent Advisory Group*. HMSO. London.
- Blom, B. & Spits, H. (2006). Development of human lymphoid cells. *Annual Review of Immunology*, Vol.24 (April 2006), pp. 287-320, ISSN 0732-0582.
- Bodkyn, C. & Lalchandani, S. (2010). Incidence of childhood cancer in Trinidad and Tobago. *West Indian Medical Journal*, Vol.59, No.5, (October 2010), pp. 465-468, ISSN 0043-3144.
- Cambier, J. C. (1995). Antigen and Fc receptor signaling. The awesome power of the immunoreceptor tyrosine-based activation motif (ITAM). *Journal of Immunology*, Vol.155, No.7, (October 1995) pp. 3281-3285, ISSN 0022-1767.
- Cardwell, CR., McKinney, PA., Patterson, CC., & Murray, LJ., (2008). Infections in early life and childhood leukaemia risk: a UK case-control study of general practitioner records. *British Journal of Cancer*, Vol.99, No.9, (November 2008), pp. 1529-1533, ISSN 0007-0920.
- Castillo, L., Fluchel, M., Dabezies, A., Pieri, D., Brockhorst, N. & Barr, R. (2001). Childhood cancer in Uruguay: 1992-1994. Incidence and mortality. *Medical and Pediatric Oncology*, Vol.37, No.4, (October 2001), pp. 400-404, ISSN 0098-1532.
- Chan, LC., Lam, TH., Li, CK., Lau, YL., Li, CK., Yuen, HL., Lee, CW., Ha, SY., Yuen, PM., Leung, NK., Patheal, SL., Greaves, MF. & Alexander, FE. (2002). Is the timing of exposure to infection a major determinant of acute lymphoblastic leukaemia in Hong Kong? *Paediatric Perinatal Epidemiology*, Vol.16, No.2, (April 2002), pp. 154-165, ISSN 0269-5022.
- Cobaleda, C., Gutiérrez-Cianca, N., Pérez-Losada, J., Flores, T., García-Sanz, R., González, M. & Sánchez-García, I. (2000). A primitive hematopoietic cell is the target for the leukemic transformation in human philadelphia-positive acute lymphoblastic leukemia. *Blood*, Vol.95, No.3, (February 2000) pp. 1007-1013, ISSN 0006-4971.
- Committee on Medical Aspects of Radiation in the Environment (COMARE). First Report Proceedings of The Implications of the New Data on the Releases From Sellafield in the 1950s for the Conclusions of the Report on the Investigation of the Possible Increased Incidence of Cancer in West Cumbria (Department of Health, 1986)
- Cooke, V. (1942). The incidence of acute leukemia in children. *The journal of the American Medical Association*, Vol.119, No.7, (June 1942), pp. 547-550, ISSN 00987484.

- Court, WM. & Doll, R. (1961). Leukaemia in childhood and young adult life. Trends in mortality in relation to aetiology. *British Medical Journal*, Vol.1, No.5231, (April 1961), pp. 981-988, ISSN 09598138.
- Cox, CV., Diamanti, P., Evely, RS., Kearns, PR. & Blair, A. (2009). Expression of CD133 on leukemia-initiating cells in childhood ALL. *Blood*, Vol.113, No.14, (April 2009), pp.3287-3295, ISSN 0006-4971.
- Cox, CV., Evely, RS., Oakhill, A., Pamphilon, DH., Goulden, NJ. & Blair, A. (2004). Characterization of acute lymphoblastic leukemia progenitor cells. *Blood*, Vol.104, No.9, (November 2004), pp. 2919-2925, ISSN 0006-4971
- de Souza Reis R, Sr., de Camargo, B., de Oliveira Santos, M., de Oliveira, JM., Azevedo Silva, F. & Pombo-de-Oliveira, MS. (2011). Childhood leukemia incidence in Brazil according to different geographical regions. *Pediatric blood & Cancer*, Vol.56, No.1, (January 2011), pp. 58-64, ISSN 1545-5017.
- Dick, J.E. (2008). Stem cell concepts renew cancer research. *Blood*, Vol.112, No.13, (December 2008), pp. 4793-4807, ISSN 0006-4971.
- Dockerty, JD., Draper, G., Vincent, T., Rowan, SD. & Bunch, KJ. (2001). Case-control study of parental age, parity and socioeconomic level in relation to childhood cancers. *International Journal of Epidemiology*, Vol.6, No.30, (December 2001), pp. 1428-1437, ISSN 0300-5771.
- Doulatov, S., Notta, F., Eppert, K., Nguyen, LT., Ohashi, PS. & Dick, JE. (2010). Revised map of the human progenitor hierarchy shows the origin of macrophages and dendritic cells in early lymphoid development. *Nature Immunology*, Vol.11, No.7, (July 2010), pp. 585-593, ISSN 1529-2908.
- Downing, JR. (1999). The AML1-ETO chimeric transcription factor in acute myeloid leukaemia. Biology and clinical significance. *British Journal of Haematology*, Vol.106, No.2, (August 1999), pp. 296-308, ISSN 1365-2141.
- Draper, G., Kroll, M. & Stiller, C. (1994). Childhood cancer. *Cancer Surveys*; Vol.19-20, (April 1994), pp. 493-517, ISSN 0261-2429.
- Eden, T. (2010). Aetiology of childhood leukaemia. *Cancer Treatment Reviews*, Vol.36, No.4, (June 2010), pp. 286-297, ISSN 0305-7372.
- Espinoza-Hernández, L., Cruz-Rico, J., Benítez-Aranda, H., Martínez-Jaramillo, G., Rodríguez-Zepeda, MC., Vélez-Ruelas, MA. & Mayani, H. (2001). In vitro characterization of the hematopoietic system in pediatric patients with acute lymphoblastic leukemia. *Leukemia Research*, Vol.25, No.4, (April 2001), pp. 295-303, ISSN 0145-2126.
- Fajardo-Gutiérrez, A., Mejía-Aranguré, M., Gómez-Delgado, A., Mendoza-Sánchez, H., Garduño-Espinosa, J. & Martínez-García, MC. (1995). Epidemiología de las neoplasias malignas en niños residentes del Distrito Federal (1982-1991). *Boletín Médico del Hospital Infantil de México*, Vol.52, No.9, (September 2005), pp. 513-522, ISSN 1665-1146.
- Fasching, K., Panzer, S., Haas, OA., Marschalek, R., Gadner, H. & Panzer-Grumayer, ER. (2000). Presence of clone-specific antigen receptor gene rearrangement at birth indicates an in utero origin of diverse types of early childhood acute lymphoblastic leukemia. *Blood*, Vol.95, No.8, (April 2000), pp. 2722-2724, ISSN 1528-0020.

- Feltbower, RG., Moorman, AV., Dovey, G., Kinsey, SE. & McKinney, PA. (2001). Incidence of childhood acute lymphoblastic leukaemia in Yorkshire, UK. *The Lancet*, Vol.358, No.9279, (August 2001), pp. 385-387, ISSN 0140-6736.
- Finnette, BA., Posseno, T. & Albertini, RJ. (1996). V(D)J recombinase-mediated HPRT mutations in peripheral blood lymphocytes of normal children. *Cancer Research*, Vol.56, No.6, (March 1996), pp. 1405-1412, ISSN 1538-7445.
- Flores-Lujano, J., Perez-Saldivar, ML., Fuentes-Pananá, EM., Gorodezky, C., Bernaldez-Rios, R., Del Campo-Martinez, MA., Martinez-Avalos, A., Medina-Sanson, A., Paredes-Aguilera, R., De Diego-Flores Chapa, J., Bolea-Murga, V., Rodriguez-Zepeda, MC., Rivera-Luna, R., Palomo-Colli, MA., Romero-Guzman, L., Perez-Vera, P., Alvarado-Ibarra, M., Salamanca-Gómez, F., Fajardo-Gutierrez, A. & Mejía-Aranguré, JM. (2009). Breastfeeding and early infection in the aetiology of childhood leukaemia in Down syndrome. *British Journal of Cancer*, Vol.101, No.5, (September 2009), pp. 860-864, ISSN 0007-0920.
- Ford, AM., Bennett, CA., Price, CM., Bruin, MC., Van Wering, ER. & Greaves, M. (1998). Fetal origins of the TEL-AML1 fusion gene in identical twins with leukemia. *Proceedings of the National Academy of Sciences of United States of America*, Vol.95, No.8, (April 1998), pp. 4584-4588, ISSN 1091-6490.
- Ford, AM., Pombo-de-Oliveira, MS., McCarthy, KP., MacLean, JM., Carrico, KC., Vincent, RF. & Greaves, M. (1997). Monoclonal origin of concordant T-cell malignancy in identical twins. *Blood*, Vol.89, No.1, (January 1997), pp. 281-285, ISSN 1528-0020.
- Ford, AM., Ridge, SA., Cabrera, ME., Mahmoud, H., Steel, CM., Chan, LC. & Greaves, M. (1993). In utero rearrangements in the trithorax-related oncogene in infant leukemias. *Nature*, Vol.363, No.6427, (May 1993), pp. 358-360, ISSN 1476-5551.
- Fuentes-Pananá, EM., & Monroe, J. G. (2001). Ligand-dependent and -independent processes in B-cell-receptor-mediated signaling. *Springer Seminars in immunopathology*, Vol.23, No.4, (December 2001), pp. 333-350, ISSN 1863-2297.
- Fuentes-Pananá, EM., Bannish, G. & Monroe, J. G. (2004a). Basal B-cell receptor signaling in B lymphocytes: mechanisms of regulation and role in positive selection, differentiation, and peripheral survival. *Immunological Reviews*, Vol.197, No.1, (January 2004), pp. 26-40, ISSN 0105-2896.
- Fuentes-Pananá, EM., Bannish, G. & Monroe, J. G. (2004b). Basal Igα/Igβ signals trigger the coordinated initiation of preBCR-dependent processes. *Journal of Immunology*, Vol.173, No.2, (July 2004), pp. 1000-1011, ISSN 0022-1767.
- Gale, KB., Ford, AM., Repp, R., Borkhardt, A., Keller, C., Eden, OB. & Greaves, MF. (1997). Backtracking leukemia to birth: identification of clonotypic gene fusion sequences in neonatal blood spots. *Proceedings of the National Academy of Sciences of United States of America*, Vol.94, No.25, (December 1997), pp. 13950-13954, ISSN 1091-6490.
- Gardner, MJ., Snee, MP., Hall, AJ., Powell, CA., Downes, S. & Terrell, JD. (1990). Results of case-control study of leukaemia and lymphoma among young people near Sellafield nuclear plant in West Cumbria. *British Medical Journal*, Vol.300, No.6722, (February 1990), pp. 423-429, ISSN 0959-8138.
- Gill Super, HJ., Rothberg, PG., Kobayashi, H., Freeman, AL., Diaz, MO. & Rowly, JD. (1994). Clonal Nonconstitutional rearrangements of the MLL gene in infant twins with acute lymphoblastic leukemia: in utero chromosome rearrangement of 11q23. *Blood*, Vol.83, No.3, (May 1994), pp. 641-644, ISSN 1528-0020.

- Glazer, ER., Perkins, CI., Young, JL. Jr., Schlag, RD., Campleman, SL. & Wright, WE. (1999). Cancer among Hispanic children in California, 1988-1994: comparison with non-Hispanic white children. *Cancer*, Vol.86, No.6, (September 1999), pp. 1070-1079, ISSN 1097-0142.
- González García, H., Quirós, AB., Crespo, CV., Meléndez, PG., Gómez, ST. & Sanz, JC. (2010). Incidence of childhood cancer in the autonomous region of Castilla y León in Spain (2003-2007). *Anales de Pediatría (Barcelona, Spain)*, Vol.73, No.4, (October 2010), pp. 169-179, ISSN 1695-9531.
- Goodman, MT., Yoshizawa, CN. & Kolonel, LN. (1989) Incidence trends and ethnic patterns for childhood leukaemia in Hawaii: 1960-1984. *British Journal of Cancer*, Vol.60, No.1, (July 1989), pp. 93-97, ISSN 1532-1827.
- Greaves, M. (2002). Childhood leukemia. *British Medical Journal*, Vol.324, No.7332, (February 2002), pp. 283-287, ISSN 1468-5833.
- Greaves, M., Maia, AT., Wiemels, JL. & Ford, AM. (2003). Leukemia in twins; lessons in natural history. *Blood*, Vol.102, No.7, (October 2003), pp. 2321-2333, ISSN 1528-0020.
- Greaves, MF. (1988). Speculations on the cause of childhood acute lymphoblastic leukemia. *Leukemia*, Vol.2, No.2, (February 1988), pp. 120-125, ISSN 1476-5551.
- Greaves, MF., Colman, SM., Beard, MEJ., Bradstock, K., Cabrera, ME., Chen, PM., Jacobs, P., Lam-Po-Tang, PR., MacDougall, LG. & Williams, CK. (1993). Geographical distribution of acute lymphoblastic leukemia subtypes: Second report of the collaborative group study. *Leukemia*, Vol.7, No.1, (January 1993), pp. 27-34, ISSN 1476-5551.
- Hardy, RR., Carmack, CE., Shinton, SA., Kemp, JD. & Hayakawa, K. (1991). Resolution and characterization of pro-B and pre-pro-B cell stages in normal mouse bone marrow. *Journal of Experimental Medicine*, Vol.173, No.5, (May 1991), pp. 1213-1225, ISSN 0022-1007.
- Heasman, MA., Kemp, IW., Urquhart, JD. & Black, R. (1986). Childhood leukaemia in northern Scotland. *Lancet*, Vol.327, No.8475, (February 1986), pp. 266, ISSN 0140-6736.
- Heidenreich, O. & Vormoor, J. (2009). Malignant stem cells in childhood ALL: the debate continues! *Blood*, Vol.113, No.18, (April 2009), pp. 4476-4477, ISSN 0006-4971.
- Hewitt, D. (1955). Some features of leukaemia mortality. *British Journal of Preventive and Social Medicine*, Vol.9, No.2, (April 1955), pp. 81-88, ISSN 0007-1242.
- Hjalgrim, LL., Rostgaard, K., Schmiegelow, K., Söderhäll, S., Kolmannskog, S., Vettenranta, K., Kristinsson, J., Clausen, N., Melbye, M., Hjalgrim, H. & Gustafsson, G. (2003). Age-and sex-specific incidence of childhood leukemia by immunophenotype in the Nordic countries. *Journal of the National Cancer Institute*, Vol.95, No.20, (October 2003), pp. 1539-1544, ISSN 1460-2105.
- Howlader, N., Noone, AM., Krapcho, M., Neyman, N., Aminou, R., Waldron, W., Altekruse, SF., Kosary, CL., Ruhl, J., Tatalovich, Z., Cho, H., Mariotto, A., Eisner, MP., Lewis, DR., Chen, HS., Feuer, EJ., Cronin, KA. & Edwards, BK. (2010). *SEER Cancer Statistics Review, 1975-2008*, National Cancer Institute. Bethesda, MD. Retrieved from <http://seer.cancer.gov/csr/1975_2008/>
- Infante-Rivard, C., Fortier, I. & Olsen, E. (2000). Markers of infection, breast-feeding and childhood acute lymphoblastic leukaemia. *British Journal of Cancer*, Vol.83, No.11, (December 2000), pp. 559-1564, ISSN 0007-0920.

- Johansson, B., Moorman, AV., Haas, OA., Watmore, AE., Cheung, KL., Swanton, S. & Secker-Walker, LM. (1998). Hematologic malignancies with t(4;11)(q21;q23)-a cytogenetic, morphologic, immunophenotypic and clinical study of 183 cases. European 11q23 Workshop participants. *Leukemia*, Vol.12, No.5, (May 1998), pp. 779-787, ISSN 1476-5551.
- Jourdan-Da Silva, N., Perel, Y., Méchinaud, F., Plouvier, E., Gandemer, V., Lutz, P., Vannier, JP., Lamagnère, JL., Margueritte, G., Boutard, P., Robert, A., Armari, C., Munzer, M., Millot, F., De Lumley, L., Berthou, C., Rialland, X., Pautard, B., Hémon, D. & Clavel, J. (2004). Infectious diseases in the first year of life, perinatal characteristics and childhood acute leukaemia. *British Journal of Cancer*, Vol.90, No.1, (January 2004), pp. 139-145, ISSN 0007-0920.
- Kaatsch, P. (2010). Epidemiology of childhood cancer. *Cancer Treatment Reviews*, Vol.36, No.4, (June 2010), pp. 277-285, ISSN 0305-7372.
- Kaatsch, P., Haaf, G. & Michaelis, J. (1995). Childhood malignancies in Germany. Methods and results of a nationwide registry. *European Journal of Cancer*, Vol.31A, No.6, (June 1995), pp. 993-999, ISSN 0959-8049.
- Karasuyama, H., Rolink, A. & Melchers, F. (1996). Surrogate light chain in B cell development. *Advances in Immunology*, Vol.63, No.2, (1996), pp. 1-41, ISSN 0065-2776.
- Kim, JM., Kim, NI., Oh, YK., Kim, YJ., Youn, J. & Ahn, MJ. (2005). CpG oligodeoxynucleotides induce IL-8 expression in CD34+ cells via mitogen-activated protein kinase-dependent and NF- κ B-independent pathways. *International Immunology*, Vol.17, No.12, (December 2005), pp. 1525-1531, ISSN 0953-8178.
- Kinlen, L. (2011). Childhood leukaemia, nuclear sites, and population mixing. *British Journal of Cancer*, Vol.104, No.1, (January 2011), pp. 12-18, ISSN 0007-0920.
- Kinlen, LJ., O'Brien, F., Clarke, K., Balkwill, A. & Matthews, F. (1993). Rural population mixing and childhood leukaemia: effects of the North Sea oil industry in Scotland, including the area near Dounreay nuclear site. *British Journal of Cancer*, Vol.306, No.6880, (March 1993), pp. 743-748, ISSN 0007-0920.
- Kinlen, L. (1988). Evidence for an infective cause of childhood leukaemia: comparison of a Scottish New Town with nuclear reprocessing sites in Britain. *The Lancet*, Vol.2, No.8624, (December 1988), pp. 1323-1327, ISSN 0140-6736.
- Kinlen, L. (1995a). Epidemiological evidence for an infective basis in childhood leukaemia. *British Journal of Cancer*, Vol.71, No.1, (January 1995), pp. 1-5, ISSN 0007-0920.
- Kinlen, LJ. & Balkwill, A. (2001). Infective cause of childhood leukaemia and wartime population mixing in Orkney and Shetland, UK. *Lancet*, Vol.357, No.9259, (March 2001), pp. 357-358, ISSN 0140-6736.
- Kinlen, LJ. & Hudson, C. (1991). Childhood leukaemia and poliomyelitis in relation to military encampments in England and Wales in the period of national military service, 1950 -63. *British Medical Journal*, Vol.303, No.6814, (November 1991), pp. 1357-1362, ISSN 0959-8138.
- Kinlen, LJ. & John, SM. (1994). Wartime evacuation and mortality from childhood leukaemia in England and Wales in 1945-9. *British Medical Journal*, Vol.309, No.6963, (November 1994), pp. 1197-1202, ISSN 0959-8138.

- Kinlen, L.J., Clarke, K. & Hudson, C. (1990). Evidence from population mixing in British New Towns of an infective basis for childhood leukaemia. *Lancet*, Vol.3368, No.715, (September 1990), pp. 577-582, ISSN 0140-6736.
- Kinlen, L.J., Dickson, M. & Stiller, CA. (1995b). Childhood leukaemia and non-Hodgkin's lymphoma near large rural construction sites, with a comparison with Sellafield nuclear site. *British Medical Journal*, Vol.310, No.6982, (March 1995), pp. 763-768, ISSN 0959-8138.
- Kristensen, T., Wesenberg, F., Jonsoon, OG., Carlsen, NT., Forestier, E., Kirchhoff, M., Lundsteen, C. & Schmiegelow, K. (2003). High resolution comparative genomic hybridisation yields at high detection rate of chromosomal aberrations in childhood acute lymphoblastic leukemia. *European Journal of Haematology*, Vol.70, No.6, (June 2003), pp. 363-372, ISSN 1600-0609.
- Lehtinen, M., Koskela, P., Ogmundsdottir, H., Bloigu, A., Dillner, J., Gudnadottir, M., Hakulinen, T., Kjartansdottir, A., Kvarnung, M., Pukkala, E., Tulinius, H & Lehtinen, T. (2003). Maternal herpesvirus infections and risk of acute lymphoblastic leukemia in the offspring. *American Journal Epidemiology*, Vol.58, No.3, (August 2003), pp. 207-213, ISSN 0002-9262.
- Levy, O. (2007). Innate immunity of the newborn: basic mechanisms and clinical correlates. *Nature Reviews Immunology*, Vol.7, No.5, (May 2007), pp. 379-390, ISSN 1474-1733.
- Linabery, AM. & Ross, JA. (2008). Trends in childhood cancer incidence in the U.S. (1992-2004). *Cancer*, Vol. 112, No.2, (January 2008), pp. 416-432, ISSN 1097-0142.
- Little, J. (August 15, 1999). *Epidemiology of Childhood Cancer*. (1st edition), International Agency for Research on Cancer Scientific Publications, ISBN-10 9283221494, ISBN-13 978-9283221494, Lyon, France
- Love, P. E., & Hayes, S. M. (2010). ITAM-mediated signaling by the T-cell antigen receptor. *Cold Spring Harbor Perspectives in Biology*, Vol.2, No.6, (June 2010), pp. 1-11, ISSN 1943-0264.
- Ma, X., Buffler, PA., Wiemels, JL., Selvin, S., Metayer, C., Loh, M., Does, MB. & Wiencke, JK. (2005). Ethnic difference in daycare attendance, early infections, and risk of childhood acute lymphoblastic leukemia. *Cancer Epidemiology, Biomarkers & Prevention*, Vol.14, No.8. (August 2005), pp. 1928-1934, ISSN 1055-9965.
- MacKenzie, J., Gallagher, A., Clayton, RA., Perry, J., Eden, OB., Ford, AM., Greaves, MF. & Jarrett, RF. (2001). Screening for herpesvirus genomes in common acute lymphoblastic leukemia. *Leukemia*, Vol.15, No.3, (March 2001), pp. 415-421, ISSN 0887-6924.
- MacKenzie, J., Perry, J., Ford, AM., Jarrett, RF. & Greaves, M. (1999). JC and BK virus sequences are not detectable in leukaemic samples from children with common acute lymphoblastic leukaemia. *British Journal of Cancer*, Vol.81, No.5, (November 1999), pp. 898-899, ISSN 0007-0920.
- Magnani, C., Dalmaso, P., Pastore, G., Terracini, B., Martuzzi, M., Mosso, ML. & Merletti, F. (2003). Increasing incidence of childhood leukemia in Northwest Italy, 1975-98. *International Journal of Cancer*, Vol. 105, No. 4, (July 2003), pp. 552-557, ISSN 1097-0215.
- Mandal, M., Powers, S. E., Ochiai, K., Georgopoulos, K., Kee, B. L., Singh, H. & Clark, M. R. (2009). Ras orchestrates exit from the cell cycle and light-chain recombination

- during early B cell development. *Nature Immunology*, Vol.10, No.10, (October 2009), pp. 1110-1117, ISSN 1529-2908.
- Mayani, H. (2010). Biological differences between neonatal and adult human hematopoietic stem/progenitor cells. *Stem Cells and Development*, Vol.19, No.3, (March 2010), pp. 285-298, ISSN 1547-3287.
- McKinney, PA., Alexander, FE., Cartwright, RA., Scott, CS. & Staines, A. (1993). Acute lymphoblastic leukaemia incidence in the UK by immunophenotype. *Leukemia*, Vol. 7, No. 10, (October 1993), pp. 1630-1634, ISSN 1476-5551.
- McKinney, PA., Juszczak, E., Findlay, E., Smith, K. & Thomson, CS. (1999). Pre- and perinatal risk factors for childhood leukaemia and other malignancies: a Scottish case control study. *British Journal of Cancer*, Vol.80, No.11, (August 1999), pp. 1844-1851, ISSN 0007-0920.
- McNally, RJ. & Eden, TO. (2004). An infectious aetiology for childhood acute leukaemia: a review of the evidence. *British Journal of Haematology*, Vol.127, No.3, (November 2004), pp. 243-263, ISSN 0007-1048.
- McNally, RJ., Birch, JM., Taylor, GM. & Eden, OB. (2000). Incidence of childhood precursor B-cell acute lymphoblastic leukaemia in north-west England. *The Lancet*, Vol.356, No.9228, (August 2000), pp. 485-486, ISSN 0140-6736.
- McNally, RJ., Cairns, DP., Eden, OB., Kelsey, AM., Taylor, GM. & Birch, JM. (2001). Examination of temporal trends in the incidence of childhood leukaemias and lymphomas provides aetiological clues. *Leukemia*, Vol.15, No.10, (October 2001), pp. 1612-1618, ISSN 1476-5551.
- Mejía-Aranguré, JM., Flores Aguilar, H., Juárez Muñoz, I., Vázquez Langle, J., Games Eternod, J., Pérez Saldívar, ML., Ortega Alvarez, MC., Rendón Macías, ME. & Gutiérrez, AF. (2005a). Age of onset of different malignant tumors in childhood. *Revista Médica del Instituto Mexicano del Seguro Social*, Vol.43, No.1, (January-February 2005), pp. 25-37, ISSN 0443-5117.
- Mejía-Aranguré, JM., Bonilla, M., Lorenzana, R., Juárez-Ocaña, S., de Reyes, G., Pérez-Saldivar, ML., González-Miranda, G., Bernáldez-Ríos, R., Ortiz-Fernández, A., Ortega-Alvarez, M., Martínez-García, Mdel C. & Fajardo-Gutiérrez, A. (2005b). Incidence of leukemias in children from El Salvador and Mexico City between 1996 and 2000: population-based data. *Bio Medical Central Cancer*, Vol.4, No.5, (April 2005), pp. 33, ISSN 1471-2407.
- Melchers, F., Karasuyama, H., Haasner, D., Bauer, S., Kudo, A., Sakaguchi N., Jameson B. & Rolink A. (1993). The surrogate light chain in B-cell development. *Immunology Today*, Vol.14, No.2, (February 1993), pp. 60-68, ISSN 0167-5699.
- Miller, RW. & Myers, MH. (1983). Age distribution of epithelial and non-epithelial cancers. *Lancet*, Vol.2, No.8361, (November 1983), pp. 1250, ISSN 0140-6736.
- Monge, P., Wesseling, C., Rodríguez, AC., Cantor, KP., Weiderpass, E., Reutfors, J., Ahlbom, A. & Partanen, T. (2002). Childhood leukaemia in Costa Rica, 1981-96. *Paediatric and Perinatal Epidemiology*, Vol.16, No.3, (July 2002), pp. 210-218, ISSN 1365-3016.
- Mousavi, SM., Pourfeizi, A. & Dastgiri, S. (2010). Childhood cancer in Iran. *Journal of Pediatric Hematology/Oncology*, Vol.32, No.5, (July 2010), pp. 376-382, ISSN 1536-3678.
- Nagata, K., Nakamura, T., Kitamura, F., Kuramochi, S., Taki, S., Campbell, K. S. & Karasuyama, H. (1997). The Ig alpha/Igbeta heterodimer on mu-negative proB cells

- is competent for transducing signals to induce early B cell differentiation. *Immunity*, Vol.7, No.4, (October 1997), pp. 559-570, ISSN 1074-7613.
- National Wilm's Tumor Study Committee (1991). Wilm's tumor: status report, 1990. *Journal of Clinical Oncology*, Vol.9, No.5, (May 1991), pp. 877-887, ISSN 1527-7755.
- Naumberg, E., Bellocco, R., Cnattingius, S., Jonzon, A. & Ekbom, A. (2002). Perinatal exposure to infection and risk of childhood leukemia. *Medical and Pediatric Oncology*, Vol.38, No.6, (June 2002), pp. 391-397, ISSN 0098-1532.
- Nguyen, M., Leuridan, E., Zhang, T., De Wit, D., Willems, F., Van Damme, P., Goldman, M. & Goriely, S. (2010). Acquisition of adult-like TLR4 and TLR9 responses during the first year of life. *Plos One*, Vol.5, No.4, (April 2010), pp. e10407, ISSN 1932-6203.
- Nordgren, A. (2003). Hidden aberrations diagnosed by interphase fluorescence in situ hybridization and spectral karyotyping in childhood acute lymphoblastic leukemia. *Leukemia and Lymphoma*, Vol.44, No.12, (March 2003), pp. 2039-2053, ISSN 1029-2403.
- Nully-Brown, P., Hertz, H., Olsen, JH., Yssing, M., Scheibel, E. & Jensen, OM. (1989). Incidence of childhood cancer in Denmark 1943-1984. *International Journal of Epidemiology*, Vol.18, No.3, (September 1989), pp. 546-555, ISSN 1464-3685.
- Nyggard, U., Larsen, J., Kristensen, TD., Wesenberg, F., Jonsson, OG., Carlsen, NT., Forestier, E., Kirchhoff, M., Larsen, JK., Schmiegelow, K. & Christensen, IJ. (2006). The sensitivity of G-band karyotyping, flow cytometric DNA index and comparative genomic hybridization in detection of high hiperdiploidy in childhood acute lymphoblastic leukemia. *Journal of Pediatric Hematology/Oncology*, Vol.28, No.3, (March 2006), pp. 134-140, ISSN 1536-3678.
- Parkin, D.M., Kramárová, E., Draper, G.J., Masuyer, E., Michaelis, J., Neglia J., Qureshi, S. & Stiller, CA (Eds.). (1998) *International Incidence of Childhood Cancer: Volume 2*, IARC Scientific Publications No. 144). International Agency for Research on Cancer, ISBN 9789283221449 , Lyon France.
- Parkin, DM., Stiller, CA., Draper, GJ. & Bieber, CA. (1988). The international incidence of childhood cancer. *International Journal of Cancer*, Vol.42, No.4, (October 1988), pp. 511-520, ISSN 1097-0215.
- Pastore, G., Carli, M., Lemerle, J., Tournade, MF., Voute, PA. Rey, A., Burgers, JMV., Zucker, JM., Burger, D., Kraker, JD. & Delemarre, JFM. (1988). Epidemiological features of Wilms' tumor: Results of studies by the International Society of Pediatric Oncology (SIOP). *Medical and Pediatric Oncology*, Vol.16, No.1, (1988), pp. 7-11, ISSN 1096-911X.
- Paulsson, K., Mörse, H., Fioretos, T., Behrendtz, M., Strömbeck, B. & Johansson, B. (2005). Evidence for a single-step mechanism in the origin of hyperdiploid childhood acute lymphoblastic leukemia. *Genes Chromosomes and Cancer*, Vol.44, No.2, (October 2005), pp. 113-122, ISSN 1098-2264.
- Pelayo, R., Hirose, J., Huang, J., Garrett, KP., Delogu, A., Busslinger, M. & Kincade, PW. (2005b). Derivation of 2 categories of plasmacytoid dendritic cells in murine bone marrow. *Blood*, Vol.105, No.11, (June 2005), pp. 4407-4415, ISSN 0006-4971.
- Pelayo, R., Miyazaki, K., Huang, J., Garrett, KP., Osmond, DG. & Kincade, PW. (2006b). Cell cycle quiescence of early lymphoid progenitors in adult bone marrow. *Stem Cells*, Vol.24, No.12, (December 2006), pp. 2703-2713, ISSN 1549-4918.

- Pelayo, R., Welner, R., Perry, SS., Huang, J., Baba, Y., Yokota, T. & Kincade, PW. (2005a). Lymphoid progenitors and primary routes to becoming cells of the immune system. *Current Opinion in Immunology*, Vol.17, No.2, (April 2005), pp. 100-107, ISSN 0952-7915.
- Pelayo, R., Welner, RS., Nagai, Y. & Kincade, PW. (2006a). Life before the pre-B cell receptor checkpoint: specification and commitment of primitive lymphoid progenitors in adult bone marrow. *Seminars in Immunology*, Vol.18, No.1, (February 2006), pp. 2-11, ISSN 1044-5323.
- Pérez-Perdomo, RV. & Rodríguez-Figueroa, L. (2000). Characteristics of cancer patients under age 20 at a population-based registry, Puerto Rico, 1980-1991. *Puerto Rico Health Sciences Journal*, Vol.19, No.2, (June 2000), pp. 123-129, ISSN 0738-0658.
- Peris-Bonet, R., Salmerón, D., Martínez-Beneito, MA., Galceran, J., Marcos-Gragera, R., Felipe, S., González, V. & Sánchez de Toledo Codina, J. (2010). Spanish Childhood Cancer Epidemiology Working Group. Childhood cancer incidence and survival in Spain. *Annals of Oncology*, Vol.21, No.Suppl 3, (May 2010), pp. 103-110, ISSN 1569-8041.
- Perrillat, F., Clavel, J., Auclerc, MF., Baruchel, A., Leverger, G., Nelken, B., Philippe, N., Schaison, G., Sommelet, D., Vilmer, E. & Hémon, D. (2002). Day-care, early common infections and childhood acute leukaemia: a multicentre French case-control study. *British Journal of Cancer*, Vol.86, No.7, (April 2002), pp. 1064-1069, ISSN 0007-0920.
- Petridou, E., Kassimos, D., Kalmanti, M., Kosmidis, H., Haidas, S., Flytzani, V., Tong, D. & Trichopoulos, D. (1993). Age of exposure to infections and risk of childhood leukaemia. *British Medical Journal*, Vol.307, No.6907, (September 1993), pp. 774, ISSN 09598138.
- Petridou, ET., Pourtsidis, A., Dessypris, N., Katsiardanis, K., Baka, M., Moschovi, M., Polychronopoulou, S., Kolioukas, D., Sidi, V., Athanasiadou-Piperopoulou, F., Kalmanti, M., Belechri, M., La Vecchia, C., Curado, MP. & Skalkidis, I. (2008). Childhood leukaemias and lymphomas in Greece (1996-2006): a nationwide registration study. *Archives of disease in childhood*, Vol.93, No.12, (December 2008), pp. 1027-1032, ISSN 1468-2044.
- Pratt, JA., Velez, R., Brender, JD. & Manton, KG. (1988). Racial differences in acute lymphoblastic leukemia mortality and incidence trends. *Journal of Clinical Epidemiology*, Vol.41, No.4, (August 1988), pp. 367-371, ISSN 0895-4356.
- Pui, CH. (1997). Acute lymphoblastic leukemia. *Pediatric Clinics of North America*, Vol.44, No.4, (August 1997), pp. 831-846, ISSN 0031-3955.
- Ramot, B. & Magrath, I. (1982). Hypothesis: the environment is a major determinant of the immunological sub-type of lymphoma and acute lymphoblastic leukaemia in children. *British Journal of Haematology*, Vol.50 No.2, (February 1982), pp. 183-189, ISSN 0007-1048.
- Rolink, A., Kudo, A., Karasuyama, H., Kikuchi, Y. & Melchers, F. (1991). Long-term proliferating early pre B cell lines and clones with the potential to develop to surface Ig-positive, mitogen reactive B cells in vitro and in vivo. *The EMBO Journal*, Vol.10, No.2, (February 1991), pp. 327-336, ISSN 0261-4189.
- Rolink, A. & Melchers, F. (1993). B lymphopoiesis in the mouse. *Advances in Immunology*, Vol.53, (1993), pp. 123-156, ISSN 0065-2776.

- Roman, E., Simpson, J., Ansell, P., Kinsey, S., Mitchell, CD., McKinney, PA., Birch, JM., Greaves, M. & Eden, T. (2007). Childhood acute lymphoblastic leukemia and infections in the first year of life: a report from the United Kingdom Childhood Cancer Study. *American Journal of Epidemiology*, Vol.165, No.5, (March 2007), pp. 496-504, ISSN 0002-9262.
- Romana, SP., Poiriet, H., Leconiat, M., Flexor, MA., Mauchauffé, M., Jonveaux, P., Macintyre, EA., Berger, R. & Bernard OA. (1995). High frequency of t(12:21) in childhood B-lineage acute lymphoblastic leukemia. *Blood*, Vol.86, No.11, (December 1995), pp. 4263-4269, ISSN 1528-0020.
- Sanders, BM., Draper, GJ. & Kingston, JE (1988). Retinoblastoma in Great Britain 1969-80: incidence, treatment, and survival. *The British Journal of Ophthalmology*, Vol.72, No.8, (August, 1988), pp. 282-286, ISSN 1468-2079.
- Schellong, G. (1985). What's new in pediatric oncology? Epidemiology, treatment principles and prognosis in childhood malignancies. *Pathology, Research and Practice*, Vol.179, No.3, (January 1985), pp. 425-428, ISSN 0344-0338.
- Seita, J. & Weissman, I.L. (2010). Hematopoietic stem cell: self-renewal versus differentiation. *Wires Systems Biology and Medicine*, Vol.2, No.6, (November/December 2010), pp. 640-653, ISSN 1939-005X.
- Shiramizu, B., Yu, Q., Hu, N., Yanagihara, R. & Nerurkar, VR. (2002). Investigation of TT virus in the etiology of pediatric acute lymphoblastic leukemia. *Pediatric Hematology and Oncology*, Vol.19, No.8, (December 2002), pp. 543-551, ISSN 1077-4114.
- Smith, MA., Chen, T. & Simon, R. (1997). Age-specific incidence of acute lymphoblastic leukemia in U.S. children: in utero initiation model. *Journal of the National Cancer Institute*, Vol.89, No.20, (October 1997), pp. 1542-1544, ISSN 1460-2105.
- Smith, MA. & Gloecker RL. (2002). Childhood cancer: incidence, survival, and mortality. In: *Principles and practice of pediatric oncology*, Pizzo PA, Poplack DG., pp. 1-12, Lippincott Williams & Wilkins Publishers, ISBN 0-7817-2658-1, Philadelphia.
- Smith, M. (1997). Considerations on a possible viral etiology for B-precursor acute lymphoblastic leukemia of childhood. *Journal of Immunotherapy*, Vol.20, No.2, (March 1997), pp. 89-100, ISSN 1053-8550.
- Spix, C., Eletr, D., Blettner, M. & Kaatsch, P. (2008). Temporal trends in the incidence rate of childhood cancer in Germany 1987-2004. *International Journal of Cancer*, Vol.122, No.8, (April 2008), pp. 1859-1867, ISSN 1097-0215.
- Stack, M., Walsh, PM., Comber, H., Ryan, CA. & O'Lorcain, P. (2007). Childhood cancer in Ireland: a population-based study. *Archives of disease in childhood*, Vol.92, No.10, (October 2010), pp. 890-897, ISSN 1468-2044.
- Stiller, CA. (2004). Epidemiology and genetics of childhood cancer. *Oncogene*, Vol.23, No.38, (August 2004), pp. 6429-6444, ISSN 1476-5594.
- Suk, WA., Murray, K. & Avakian, MD. (2003). Environmental hazards to children's health in the modern world. *Mutation Research*, Vol.544, No.2-3, (November 2003), pp. 235-242, ISSN 0027-5107.
- Teppo, L., Salonen, T. & Hakulinen, T. (1975). Incidence of childhood cancer in Finland. *Journal of the National Cancer Institute*, Vol.55, No.5, (November 1975), pp. 1065-1067, ISSN 1460-2105.
- Thomas, L. R., Cobb, R. M. & Oltz, E. M. (2009). Dynamic regulation of antigen receptor gene assembly. *Advances in Experimental Medicine and Biology*. Vol.650, (June 2009),

- pp. 103-115, ISSN 0065-2598. U.S. Cancer Statistics Working Group. (2010). *United States Cancer Statistics: 1999–2007 Incidence and Mortality Web-based Report*. Atlanta: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention and National Cancer Institute. Retrieved from < www.cdc.gov/uscs >
- von Boehmer, H. & Melcher, F. (2010). Checkpoints in lymphocyte development and autoimmune disease. *Nature Immunology*, Vol.11, No.1, (January 2010), pp. 14-20, ISSN: 1529-2908.
- Welner, RS., Pelayo, R. & Kincade, PW. (2008a). Evolving views on the genealogy of B cells. *Nature Reviews Immunology*, Vol.8, No.2, (February 2008), pp. 95-106, ISSN 1474-1733.
- Welner, RS., Pelayo, R., Garrett, KP., Chen, X., Perry, SS., Sun, XH., Kee, BL. & Kincade, PW. (2007). Interferon-producing dendritic cells (IKDC) arise via a unique differentiation pathway from primitive c-kit^{hi}CD62L⁺ lymphoid progenitors. *Blood*, Vol.109, No.11, (June 2007), pp. 4825-4831, ISSN 0006-4971.
- Welner, RS., Pelayo, R., Nagai, Y., Garrett, KP., Wuest, TR., Carr, DJ., Borghesi, LA., Farrar, MA. & Kincade, PW. (2008b). Lymphoid precursors are directed to produce dendritic cells as a result of TLR9 ligation during Herpes infection. *Blood*, Vol.112, No.9, (November 2008), pp. 3753-3761, ISSN 0006-4971.
- Wiemels, JL., Ford, AM., Van Weiring, ER., Postma, A. & Greaves, M. (1999b). Protracted and variable latency of acute lymphoblastic leukemia after TEL-AML1 gene fusion in utero. *Blood*, Vol.94, No. 3, (August 1999), pp. 1057-1062, ISSN 1528-0020.
- Wiemels, J., Cazzaniga, G., Daniotti, M., Eden, OB., Addison, GM., Masera, G., Saha, V., Biondi, A. & Greaves, M. (1999a). Prenatal origin of acute lymphoblastic leukaemia in children. *Lancet*, Vol.354, No.9189, (October 1999), pp. 1499-1503, ISSN 0140-6736.
- Wiest, D. L., Kears, K. P., Shores, E. W. & Singer, A. (1994). Developmentally regulated expression of CD3 components independent of clonotypic T cell antigen receptor complexes on immature thymocytes. *Journal of Experimental Medicine*, Vol.180, No.4, (October 1994) pp. 1375-1382, ISSN 0022-100.
- Wiest, D. L., Burgess, W. H., McKean, D., Kears, K. P. & Singer, A. (1995). The molecular chaperone calnexin is expressed on the surface of immature thymocytes in association with clonotype-independent CD3 complexes. *The EMBO Journal*, Vol.14, No.14, (July 1995), pp. 3425-3433, ISSN 0261-4189.
- Wilkinson, JD., Fleming, LE., MacKinnon, J., Voti, L., Wohler-Torres, B., Peace, S. & Trapido, E. (2001). Lymphoma and lymphoid leukemia incidence in Florida children: ethnic and racial distribution. *Cancer*, Vol.91, No.7, (April 2001), pp. 1402-1408, ISSN 1097-0142.
- William, AO. (1975). Tumors of childhood in Ibadan, Nigeria. *Cancer*, Vol.36, No.2, (August 1975), pp. 370-378, ISSN 1097-0142.
- Yagi, T., Hibi, S., Tabata, Y., Kuriyama, K., Teramura, T., Hashida, T., Shimizu, Y., Takimoto, T., Todo, S., Sawada, T. & Imashuku, S. (2000). Detection of clonotypic IGH and TCR rearrangements in the neonatal blood spots of infants and children with the B-cell precursors acute lymphoblastic leukemia. *Blood*, Vol.96, No.1, (July 2000), pp. 264-268, ISSN 1528-0020.
- Yaris, N., Mandiracioglu, A. & Büyükpamukcu, M. (2004). Childhood cancer in developing countries. *Journal of Pediatric Hematology/ Oncology*, Vol.21, No.3, (April-May 2004), pp. 237-253, ISSN 1536-3678.

- Yasuda, T., Sanjo, H., Pagès, G., Kawano, Y., Karasuyama, H., Pouysségur, J., Ogata, M. & Kurosaki, T. (2008). Erk kinases link pre-B cell receptor signaling to transcriptional events required for early B cell expansion. *Immunity*, Vol.28, No.4, (April 2008), pp. 499-508, ISSN 1074-7613.
- Young, JI., Gloeckler, RL., Silverberg, E., Horm, JW. & Miller R. (1986). Cancer incidence, survival, and mortality for children younger than age 15 years. *Cancer*, Vol.58, No.2Suppl, (July 1986), pp. 598-602, ISSN 1097-0142.

Novel Aspects in Acute Lymphoblastic Leukemia

Edited by Dr. Stefan Faderl

ISBN 978-953-307-753-6

Hard cover, 258 pages

Publisher InTech

Published online 16, November, 2011

Published in print edition November, 2011

Acute lymphoblastic leukemia (ALL) has turned from a universally fatal to a highly curable disease in little more than four decades. Even though differences in outcome continue to exist between children and adults, intense efforts are under way to overcome this discrepancy and improve the prognosis of adult patients as well. This exemplary progress in ALL therapy has been possible by the combination of an increasingly better understanding of the biology of the disease, availability of a range of effective drugs, and astute designs and relentless executions of many clinical trials. ALL is a complex disease requiring complex therapy. Whereas this book cannot provide a comprehensive review of every one of its many facets, the chapters from many investigators from around the world nevertheless cover a number of relevant topics: aspects of the epidemiology of ALL in Hispanics, ophthalmologic manifestations of ALL, overviews of current therapy and drug-resistance mechanisms, novel biological pathways and targets, new drugs in development, and long-term consequences of CNS prophylaxis and therapy. The publishers and editor therefore hope that the prospective readers will find enough insight and information for their own endeavors.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Juan Manuel Mejía-Arangur, Mara Luisa Prez-Saldivar, Rosana Pelayo-Camacho, Ezequiel Fuentes-Panan, Carolina Bekker-Mendez, Abigail Morales-Snchez, David Aldebarn Duarte-Rodrguez and Arturo Fajardo-Gutirrez (2011). Childhood Acute Leukemias in Hispanic Population: Differences by Age Peak and Immunophenotype, *Novel Aspects in Acute Lymphoblastic Leukemia*, Dr. Stefan Faderl (Ed.), ISBN: 978-953-307-753-6, InTech, Available from: <http://www.intechopen.com/books/novel-aspects-in-acute-lymphoblastic-leukemia/childhood-acute-leukemias-in-hispanic-population-differences-by-age-peak-and-immunophenotype>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2011 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](https://creativecommons.org/licenses/by/3.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen