

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

185,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Genetics of Type 1 Diabetes

Tatijana Zemunik and Vesna Boraska
*University of Split School of Medicine,
 Split,
 Croatia*

1. Introduction

Type 1 diabetes (T1D) is an autoimmune disease characterized by immune destruction of insulin-producing pancreatic β cells. This leads to dysfunctional regulation of blood glucose levels in T1D patients. The destruction of β cells of Langerhans islets is caused by infiltration of dendritic cells, macrophages and T lymphocytes. The destruction of β cells starts with an autoimmune process that is followed by massive destruction of β cells later on. Autoantibodies against T1D-specific antigens are present in serum and can be detected in the early stage of the disease (Ounis-Benkalha & Polychronakos, 2008). There are several main types of T1D autoantibodies: islet antibodies, antibodies to insulin (IAA), glutamic acid decarboxylase (GADA) and tyrosine phosphatase IA-2. In the last few years antibodies to zinc transporter (ZnT8) have been added to this group (Mehers & Gillespie, 2008). It is generally accepted that T1D occurs as a result of genetic and environmental factors when presence of many alleles combined with effects of numerous environmental factors lead to disease development (Pociot et al., 2010). Research of T1D genetic basis and environmental factors has increased dramatically in the last two decades. Today it is considered that beside *HLA* region on chromosome 6q21 that contributes approximately with 40% to T1D development, more than 50 non-*HLA* genes significantly increase the risk of T1D occurrence (MacFarlane et al., 2009; Ziegler et al., 2010; Concannon et al., 2010). The final aim of genetic research is integration with clinical practice, which is expected once the main understanding of genetic etiology of T1D is achieved. Translation to clinics includes development of genetic-based diagnostic tests, population screening methods and prevention strategies, and finally, development of new treatments and therapies (Manolio et al., 2009).

2. Genetic studies

There are two main approaches in dissecting T1D genetic background: linkage analysis and association analysis (Figure 1). Linkage analysis is based on simple Mendelian inheritance and it uses affected relatives (typically siblings) to identify regions on chromosomes that are shared more frequently than expected by chance. Since affected siblings are relatively rare in T1D, linkage studies have been performed in somewhat unique subgroup of families with T1D (Concannon et al., 2009). In general, samples are genotyped for a modestly dense panel of markers, typically microsatellites, to search for linked alleles i.e. alleles that are inherited together. Regions in the genome with accumulating evidence of linkage are further fine mapped, which means that additional markers are typed in the same chromosomal region,




Fig. 1. Linkage analysis tests for the co-segregation of alleles within family members whereas association analysis searches for the difference in allele frequency between unrelated groups of affected and unaffected individuals or within families. Adapted from Concannon et al., 2009.




Fig. 2. Relation of risk allele frequencies, effect sizes (odds ratios) and feasibility of identifying risk variants by common genetic tests. Generally, linkage studies are better in indentifying low-frequency alleles with larger effect sizes, whereas association studies are more effective in identifying common variants with small to moderate effects. Adapted from McCarthy et al., 2008 & Manolio et. al., 2009.

in order to narrow down regions associated with disease. Linkage analyses are most effective in identifying rare alleles with large effect sizes (Figure 2) (Concannon et al., 2009). On the other hand, common alleles with modest and small effect sizes can be identified through association analysis. Association studies test for association of genotyped marker (typically single nucleotide polymorphism, SNP) with the disease of interest in a case-control or family-based sample. These studies rely on the assumption that investigated allele is associated with disease if it differs in frequency between two investigated groups of individuals. Tested polymorphism is usually not the causative one but it will show an association if it is in linkage disequilibrium (LD) with an unknown causative, risk or protective, variant. Human genome is divided into regions of high and low LD, and if allele resides in the region of high LD that means that many SNPs from the same region will be inherited together and therefore reflect one another. This means that genotyping of only limited, but carefully selected, set of SNPs can actually capture a majority of information within tested gene region (Lander, 2011).

For the last decade the most common design of association analysis used to be candidate gene approach that searches for differences in allele frequencies in specifically selected genes between affected and healthy groups of individuals or affected subjects and their parents. There are few general limits of candidate gene studies that include modest sample sizes, limited number of investigated variants, the fact that selection of genes/variants is often based on inadequate understanding of biological pathways and, most importantly, observed associations are usually difficult to replicate (Manolio et al., 2009). However, just few years ago a complete dominance in association analyses design was taken by genome-wide association studies (GWAS) approach. These are hypothesis free studies that usually test between 300,000 up to 1 million directly genotyped SNPs that capture substantial proportion of common genetic variation of the genome (McCarthy et al. 2008). The methodology behind the GWAS is the same as in any association study and that is to map susceptibility variants through identification of associations between allele (genotype) frequency and disease status (McCarthy et al. 2008). The development of both, high-throughput genotyping platforms and a catalogue of human variation by International HapMap Project (<http://hapmap.ncbi.nlm.nih.gov/>) and The 1000 Genomes project (<http://www.1000genomes.org/>) have made possible high utilization of GWAS. In addition, development of imputation methods that infer missing genetic variants enabled inclusion and comparison of different GWAS in large-scale meta-analysis framework (Marchini & Howie, 2010). Also, huge collaborative international projects such as The Type 1 Diabetes Genetics Consortium (T1DGC) (<https://www.t1dgc.org/home.cfm>) have put efforts to collect and systemise data from several thousand T1D affected and healthy individuals worldwide in order to identify genes contributing to an individual's risk for T1D susceptibility. Overall, the combination of linkage, association and large-scale GWAS approaches provided evidence of genetic contribution of many common and rare alleles with wide range of effect sizes to the T1D development.

2.1 Linkage analysis approach

Genetic linkage studies have shown the biggest success in discovering genetic loci underlying monogenic disorders where risk factors, even rare in frequency, have large effects and often lead to change in amino acid sequence (Smith & Newton-Cheh 2009). In complex diseases, such as T1D, situation is less straightforward since there are loci of small, modest and large effect sizes contributing to the disease (Concannon et al., 2009; Kere, 2010).

Several linkage analyses of T1D provided evidence for linkage between the *HLA* region on chromosome 6p21 and T1D. There were also other non-*HLA* loci findings but these were not consistently replicated among studies, mostly because of limited sample sizes. T1DGC conducted linkage meta-analysis of most of T1D genome-wide linkage studies and showed strong evidence of linkage with *HLA* class II genes encoding *HLA-DR* and *HLA-DQ*. In addition, this study demonstrated supporting evidence of linkage of additional genes within *HLA* region and small number of other regions in the genome (Concannon et al., 2009).

2.2 Association analysis approach

2.2.1 Case-control design

Case-control design is one of the most common association study designs. Case-control study compares two groups of individuals, one with the disease (cases) and the other without disease (controls). It is assumed that cases have higher prevalence of susceptibility alleles for disease of interest than controls and that susceptibility alleles can be detected through direct comparison of allele frequencies between two groups (McCarthy et al. 2008). A lot of attention is given to case ascertainment to minimize phenotypic heterogeneity. In addition, study power can be improved by selection of cases, for example to those with the extremes of phenotypes (McCarthy et al. 2008). Since it is observed that incident rates of T1D highly increase in the very young group of children, an early age of disease onset could be an example of extreme T1D phenotype and enrichment of those cases in the sample set is likely to improve power (McCarthy et al. 2008; Maahs et al. 2010).

In genetic epidemiological studies a lot of attention is given to control selection. Controls are matched with cases by ethnicity to avoid problems of population stratification that may result with spurious associations (false positives). This means that controls are selected from the same population, preferably from the same region, as cases (Zondervan & Cardon 2007). Nowadays, with the genome-wide data, it is possible to estimate the level of relatedness among individuals and, also, the matching of cases and controls by ancestry (Anderson et al. 2010). Principal component analysis is one of the most common methods that enables clustering of individuals by ancestry (Figure 3). To further reduce stratification within the sample set, controls can also be matched to cases by age, sex and environmental factors. Usually, association analyses are adjusted for covariates with strong impact on phenotype to reduce non-genetic contribution to phenotype variation (Smith & Newton-Cheh 2009).

2.2.2 Family-based design

Family-based association studies, most commonly in the form of parent-offspring trios, use another analytical approach to test for association. To make assumptions on association with the disease these studies examine the transmission of alleles from heterozygous parents to affected offspring that is observed more frequently than expected by chance (Smith & Newton-Cheh 2009). Since these studies are conducted within families they offer a protection from population stratification but they also rely on informative parent-offspring trios which usually reduce the effective sample size, thus power as well (McCarthy et al. 2008; Smith & Newton-Cheh 2009). Family-based studies are particularly useful in finding variants underlying relatively rare phenotypes that segregate within families. Also, these studies have advantages when age of disease onset is low, as in the case of T1D, because it enables easier collection of many family members (Smith & Newton-Cheh 2009).




Fig. 3. Principal component analysis of samples deriving from population of interest (POP) and 11 HapMap phase II populations (CEU - Utah residents with Northern and Western European ancestry from the CEPH collection; TSI - Tuscans in Italy; MEX - Mexican ancestry in Los Angeles; GIH - Gujarati Indians in Houston; CHB - Han Chinese in Beijing, China; CHD - Chinese in Metropolitan Denver; JPT - Japanese in Tokyo, YRI - Yoruba in Ibadan, Nigeria; LWK - Luhya in Webuye, Kenya; ASW - African ancestry in Southwest USA; MKK - Maasai in Kinyawa, Kenya)

2.2.3 Genome-wide association studies (GWAS) and meta-analyses

Rationale underlying GWAS is the ‘common disease, common variant’ hypothesis. It is believed that both, common and rare variants, contribute to complex disease risk. However, GWAS are generally powered only to detect association of common variants (allelic variants present in more than 5% of the population) with modest to large effect sizes (Manolio et al., 2009). GWAS are not designed to identify multiple rare mutations within a gene (Kere, 2010).

Because of the modest sample sizes individual GWAS have limited power to detect all associations underlying complex diseases. Increase in sample size achieved by combination of statistical evidence of individual studies through meta-analysis approach can improve study power and raise a discovery of susceptibility loci. Nowadays, a majority of new genetic findings underlying complex traits are found through meta-analysis approach. Since different studies differ in design, sample collection, genotyping platform and analysis

methodology, one of the most important prerequisites for meta-analysis is capability to uniform study results. Most genotyping platforms have different representation of genetic markers and harmonisation of studies through expansion of SNP coverage can be achieved by imputation processes. Imputation infers and fills in missing genotypes on the basis of HapMap, the 1000 Genomes or other reference panels to allow different studies to analyse the same set of common SNPs (de Bakker et al., 2010). The biggest meta-analysis for T1D combined results from two studies and included total of 7,514 cases and 9,045 reference samples. This study identified another 18 regions associated with T1D that suggested novel candidate genes such as *IL10*, *IL19*, *IL20*, *GLIS3*, *CD69* and *IL27* (Barrett et al., 2009).

2.3 Other types of genetic research

There are many other types of genetic research that contribute to understanding of complex disease mechanisms. Many of these studies use knowledge on susceptible genetic variation accumulated through linkage and association studies. Identified variants are usually additionally analysed for gene-gene and gene-environment interactions. Also, it is well known that genes interact through complex molecular networks and integrating the prior knowledge of biological pathways of genes of interest may increase a chance to find genes involved in disease development. These pathway-based analyses use different software packages that search through variety of web-based databases and take into account the existing data on biological pathways of investigated genes (Wang et al., 2010). Cross-disorder overlap is another search that looks for evidence of potential overlapping regions of the genome affecting various different diseases, such as T1D and other autoimmune traits (Eyre et al. 2010). All these supplementary analyses help in elucidating genetic contribution in complex disease development.

There are functional studies that also use data derived from genetic analyses. Most commonly performed ones are gene expression analyses that may investigate susceptible genes/gene variants in different tissues or investigate them under different environmental stimuli. Combining the information on gene expression profiles and alternate splicing sites across a range of human tissues together with genetic mapping for the same samples will be valuable in deciphering the roles of genetic variants (McCarthy et al., 2008). Genetical genomics analysis also offers new means in understanding the genetic architecture of gene expression (Cui et al., 2010)

2.4 Finding the missing heritability

GWAS have identified more than 50 genetic variants associated with T1D. However, just like in most other complex traits, associated variants explain only a small proportion of heritability of T1D ($\lambda_s \sim 5$, whereas it is estimated to be 15) and have rather small effect on disease risk (Clayton, 2009). The remaining missing heritability can be explained in several different ways such as an influence of much larger number of common variants of smaller effect sizes that still need to be identified, an influence of rare variants of modest and small effect sizes that have not yet been discovered because of their underrepresentation in the current genotyping platforms and because of underpowered sample sizes, an influence of structural variants that are also poorly captured by existing platforms and generally low power to detect gene-gene and gene-environment interactions (Manolio et al., 2009). Sample size is generally one of the major limiting factors for discovery of common alleles with small effect sizes. Augmenting the number of investigated individuals through meta-analysis

approach to more than tens of thousands of individuals is another way for discovery of new genetic loci (Lander, 2011). On the other hand sequencing is the best way for discovering rare and structural variants such as copy number variants, inversions, translocations, microsatellite repeat expansions, insertions of new sequence and complex rearrangements (Manolio et al., 2009). Because of immense decrease in price, sequencing is becoming a common practice and next generation sequencing (exon or whole-genome sequencing) might provide many clues for missing heritability. The 1000 Genomes Project (<http://www.1000genomes.org/>) aims to provide a complete catalogue of human genome sequence variation and the pilot phase of the project already identified around 15 million SNPs, 1 million short insertions and deletions and 20,000 structural variants (The 1000 Genomes Project Consortium et al., 2010). Most of these variants were previously unknown and will provide a foundation for future genetic research of human diseases, including T1D. Also, sequencing of individuals with extreme phenotypes, for example individuals with the extreme age of T1D diagnosis, might provide important findings because it is thought that they carry more deleterious, loss-of-function mutations (Romeo et al., 2007).

It is also thought that some of the missing heritability might be discovered by conducting studies in populations of non European ancestry. Most genetic studies have been limited to European populations even it is known that genetic variation is highest in the populations of recent African ancestry. These studies might prove useful in detecting rare variants associations and narrowing down associated regions due to smaller LD windows (International HapMap Consortium et al., 2007). Family studies and isolated populations are another sample sets that might help in identifying missing heritability due to their enrichment of unique genetic variants (Sabatti et al., 2009).

2.5 Prevention, diagnostics and clinical application of genetic findings

Genetic research of complex diseases aims to improve understanding of biological and physiological pathways involved in disease etiology. The main goal is integration of newly accumulated knowledge with clinical practice by development of more effective means of diagnosis, prevention, treatment and response to therapies. Identification of predictive variants for considerable proportion of disease, even with identification of many other risk variants with smaller effect sizes, is very challenging (Manolio et al., 2009). The biggest influence for T1D development is carried by *HLA* loci and it is shown that only few *HLA* SNPs capture most of the heritability of T1D risk that is attributable to *HLA* associations (Clayton, 2009). There is substantial genetic risk to T1D that can be attributable to yet undiscovered loci and it is assumed that the majority of these loci will have smaller effects than those loci which have already been discovered (Clayton, 2009). So far, genetic prediction for T1D is modest and it is still not reaching criteria required for a targeted disease prevention strategy (Clayton, 2009). However, it is believed that a small proportion of population at the highest risk will be identifiable and the development of diagnostic and targeted prevention strategies for those individuals will be feasible. These diagnostic and interventional strategies will require more accurate genetic prediction and it is necessary that they are developed through ethical, safe, effective and individualized approach (Manolio et al., 2009; Clayton, 2009). One of suggested ways to discover individuals at risk is through population screening which means genotyping specific genetic variants, whole-genome typing and whole-genome sequencing of entire population (Manolio et al., 2009; Pharoah et al., 2008). Figure 4 shows the ways of translation of genetic findings into clinical practice and disease management.




Fig. 4. Translation of genetic information into clinical practice. Adapted from McCarthy et al. 2008.

3. Genetic background of T1D

3.1 Rare monogenetic forms of T1D

Very rare form of autoimmune diabetes is monogenetic diabetes, which means that it is caused by mutation of a single gene. In such cases, diabetes occurs as part of multiple set of autoimmune diseases. One of them is known as the immune dysregulation, polyendocrinopathy, enteropathy, X-linked (IPEX) syndrome in which a function of regulatory T cells is impaired (van Belle et al., 2011). It occurs as a result of mutations of *FOXP3*, the master gene for normal functioning of regulatory T cells (Wang et al., 2010). About 80% of children with IPEX syndrome develop autoimmune diabetes at an early age (van Belle et al., 2011). Another example of multiple autoimmune disease is autoimmune polyendocrine syndrome type 1 (APS-1) or autoimmune polyendocrinopathy-candidiasis-ectodermal dystrophy (APECED) (van Belle et al., 2011). The main cause of APECED development is mutations in the autoimmune regulator (AIRE) protein (Villaseñor et al. 2005). Lack of AIRE protein decreases the expression of insulin and other peripheral molecules in the thymus. This allows effector T cells escape to the periphery and prevent their negative selection in the thymus through apoptosis (Liston et al., 2004). About 20% of APS1 or APECED patients develop T1D (van Belle et al., 2011). Although the number of patients with this monogenic form of T1D is almost negligible, these findings indicate the high impact of immune status on the disease occurrence (van Belle et al., 2011).

3.2 Family history of T1D

Over 85% of T1D patients do not have positive family history of T1D, however there is a 6% of disease clustering among siblings. Siblings have 15 times greater chance of developing T1D in comparison with the general population which gives strong evidence of the genetic background of this disease. The pattern of inheritance seems very complicated, and disease development further depends on the triggers from the environment. Long-term monitoring showed that the concordance rate of inheritance is greater than 50% in monozygous twins, while it is 6-10% in dizygous twins, which is similar to that of siblings. Interestingly, the

siblings who share both identical haplotypes of *DR3/DR4* *HLA* class II region, which shows highest susceptibility to T1D, have a higher risk of disease development than those who share only one or no haplotypes of that region. (Steck & Rewers 2011).

3.3 HLA

First and consistent evidence of *HLA* gene contribution to the disease was provided by linkage analyses and further confirmed by association analyses. *HLA* genes contribute with approximately 40% of genetic risk to T1D development (MacFarlane et al., 2009). The proteins encoded by *HLA* genes are cell-surface proteins grouped in class I (A, B and C) and class II (DP, DQ and DR) of *HLA* region on chromosome 6p21 (Figure 5). Both groups of proteins are essential in self and non-self immune recognition. The proteins encoded by the *HLA* class I genes are single chain proteins that present intracellular antigen to CD8⁺ T killer cells while proteins encoded by *HLA* class II are built of two chains and present extracellular antigen to CD4⁺ T helper cells (Ounisis-Benkalha & Polychronakos, 2008). It is considered today that the *HLA* region class II has the strongest input in the development of T1D, which has been deeply investigated in the past several decades (van Belle et al., 2011). *HLA DR4* and *DR3* class II haplotypes are of particular importance in the T1D development. Even greater risk of T1D development have individuals with genetic combinations of two susceptible alleles, *DR3/DR4*. *HLA* region on chromosome 6p21.3 comprises more than 200




Fig. 5. *HLA* region on hromosome 6p21.3. Adapted from Mehers & Gillespie, 2008.

genes that are in high LD. Therefore, it is difficult to determine which gene gives the observed effect. It is considered that haplotypes of high risk for T1D are *DRB1*0401-DQA1*0301-DQB1*0302* and *DRB1*0301-DQA1*0501-DQB1*0201* (Mehers & Gillespie, 2008; Skrodeniene et al., 2010). Antigen-presenting cells (APC) carry HLA class II molecules that bind key T1D autoantigens such as preproinsulin, insulinoma associated antigen 2, glutamic acid decarboxylase (GAD) and zinc transporter (ZnT8) and present them to thymocytes in the thymus. Strongly self-reactive thymocytes die by apoptosis, negative selection that eliminates 98% of thymocytes. Only 2% of thymocytes that have low affinity migrate as mature T cells in the periphery where they develop into CD4⁺ and CD8⁺ T cells. CD4⁺ T cells are helper cells to CD8⁺ T killer cells in the processes of destruction of pancreatic islet β cells (Figure 6). It is believed that the most important autoantigen in the onset of T1D is preproinsulin, whose N-terminal signal peptide and the peptidase cleavage site is recognised by CD8⁺ T killer cells. The proteins encoded by *HLA* class I, *HLA-A* and *HLA-B* exert smaller effects in the pathogenesis of T1D. Proteins encoded by class I *HLA* genes are expressed on nucleated cells, often in the pancreatic insulin-producing β cells. These proteins present antigens directly to the CD8⁺ T killer cells (Figure 6) (Ounissi-Benkalha & Polychronakos, 2008; Todd, 2010; Pociot et al., 2010; Blueston et al., 2010; van Belle et al., 2011). However, it was observed that some *HLA* haplotype combinations have protective association with T1D. It is believed that haplotype *DR15-DQ6*, which was found in about 20% of the general population and even less than 1% of patients with T1D has protective effect (Ounissi-Benkalha & Polychronakos, 2008).

3.4 Other non-*HLA* pathways

Linkage analyses additionally pointed to linkage of some non-*HLA* loci but without consistent replication. Candidate-gene association studies have further confirmed *HLA* loci as the major T1D genetic factors but also identified four other T1D susceptibility loci: gene for insulin (*INS*), cytotoxic T-lymphocyte-associated protein 4 (*CTLA4*), protein tyrosine phosphatase, non-receptor type 22 (lymphoid) (*PTPN22*) and interleukin 2 receptor, alpha (*IL2RA*) (Bell et al., 1984; Nisticò et al., 1996; Bottini et al., 2004; Lowe et al., 2007).

The first strong non-*HLA* association with T1D was shown with polymorphisms within the *INS* gene (Bell et al., 1984). Variable number of tandem repeats (VNTR) composed of 14-15 bp tandem repeat sequence (ACAGGGGTGTGGGG) is located 596 bp upstream of the *INS* gene on chromosome 11p15 and regulates gene transcription (Ounissi-Benkalha & Polychronakos, 2008). Alleles of this region are divided into three classes with respect to the number of consecutive VNTRs: class I VNTR alleles (short, 26-63 repeats), class II VNTR alleles (63-140 repeats) and class III VNTR alleles (long, 141-209) (Durinovic-Bello et al., 2010). They correlate with *INS* mRNA production in pancreas and thymus. Class I alleles of the *INS* VNTR increase the risk of T1D and have been associated with high mRNA levels in pancreas and low levels in thymus. Class III alleles are associated with 20% lower mRNA than class I in pancreas but two to three times higher mRNA levels in thymus and therefore are considered to be protective. It seems that high levels of proinsulin in the thymus may stimulate negative selection (dying by apoptosis) of insulin specific T-lymphocytes crucial in the pathogenesis of T1D. On the other hand it is believed that lower levels of proinsulin in thymus affect the positive selection of T cells in the thymus, migration of CD4⁺ proinsulin specific T cells in the periphery, that increase risk to T1D. In that way, the genetically regulated selection mechanisms affect the selection of autoreactive cells in the immune response to autoantigens (Todd, 2010; Pociot et al., 2010; Ounissi-Benkalha & Polychronakos, 2008).




Fig. 6. The development of an autoimmune reaction in T1D. Adapted from Ounissi-Benkalha & Polychronakos, 2008.

In 1996 *CTLA4* gene located on chromosome 2q33 was confirmed as another T1D susceptibility gene (Nistico L et al., 1996.). *CTLA4* protein is a co-stimulatory receptor on the cell surface of $CD4^+$ T cells. It binds B7 ligands of APCs that activate main component of the co-receptor, CD28. In the same time intracellular part of the *CTLA4* interact with intracellular domain of CD3 receptor and start phosphorylation of the several downstream target molecules. Consequently, this leads to activation of T cells after their binding to HLA molecules on APCs (Figure 6) (Ounissi-Benkalha & Polychronakos, 2008). Reduction of the *CTLA4* protein in $CD4^+$ T cells increases susceptibility to T1D (Todd, 2010; Pociot et al., 2010). The A49G polymorphism in exon 1 causes substitution of an alanine with a threonine in the signal sequence of *CTLA4* protein that leads to incorrect glycosylation of mutant protein and reduction of its expression on T-cell surface. Conversely, C318T

polymorphism of the *CTLA4* promoter gene region causes higher promoter activity and increases the amount of CTL4 protein on the T cell surface. Therefore, it can be considered as protective polymorphism of this gene (Ounissi-Benkhalha & Polychronakos, 2008). More recently in 2004, *PTPN22* gene located on chromosome 1q13 that encodes lymphoid tyrosine phosphatase was recognized as further T1D susceptibility gene (Bottini et al., 2004). Lymphoid tyrosine phosphatase “Lyp” inhibits T cell receptor (TCR) signal transduction and causes inhibition of CD4⁺ T-cell activation (Figure 6). The Lyp inhibitory function is enhanced by its interaction with C-terminal Src tyrosine kinase (Csk). T1D is associated with a polymorphism at position 1858 by replacing the C to T, which leads to the substitution of arginine to tryptophan at position 620 of Lyp protein (R620W) (Ounissi-Benkhalha & Polychronakos, 2008; Mehers & Gillespie, 2008; Todd, 2010; Pociot et al., 2010). The Lyp interaction with tyrosine kinase Csk occurs just in the 620 region. A 620W allele interacts less strongly with Csk than 620R allele. A 620W homozygous shows increased inhibition of TCR signalling, reduces CD4⁺ T cell activation thus resulting in increased autoimmunity (Todd, 2010; Pociot et al., 2010; Ziegler et al., 2010).

Another strongly associated gene, *IL2RA* located on chromosome 10p15 that encodes alpha chain (CD25) molecules of IL-2 receptor has been reported as T1D associated gene (Vella et al. 2005). The CD25 is responsible for binding of IL-2 and proliferation of regulatory T cell and consequently, it affects their function in inhibition of effector T cells and autoimmune disease (Figure 6). T1D-predisposing alleles of *IL2RA* gene correlate with lower amounts of CD25 on the surface of the regulatory T cells which are suppressors of autoreactivity and that can be important in regulation of T-cell proliferation by an immunogenic stimulus (Ounissi-Benkhalha & Polychronakos, 2008; Mehers & Gillespie, 2008; Todd, 2010; Pociot et al., 2010).

Since 2001 a significant number of GWA studies have been reported. Data from The International Type 1 Diabetes Genetics Consortium (T1DGC) collected through multiple genome-wide association studies are available to the scientific community by request. Recently, GWAS and large scale meta-analyses identified more than 40 loci that affect the risk of developing T1D (Table 1.) (Barrett et al., 2009, Pociot et al., 2010). The analysis included 7.514 cases and 9.045 control samples. Fifteen of these regions have been previously reported as regions associated with T1D susceptibility. Eighteen additional regions showed significant association with T1D and several of them contain new candidate genes of possible relevance to T1D (*IL10*, *IL19*, *IL20*, *GLIS3*, *CD69* and *IL27*) (Pociot et al., 2010). Most of the listed genes mediate the immune response, some exert their functions in the process of destruction of pancreatic β cells and some have a dual role (Pociot et al., 2010). A list of SNPs and genes associated with T1D is presented in Table 1.

| SNP | Chromosome | OR minor allele | Gene of interest |
|-----------|------------|-----------------|------------------|
| rs2476601 | 1p13.2 | 2.05 | PTPN22 |
| rs2269241 | 1p31.3 | 1.10 | PGM1 |
| rs2816316 | 1q31.2 | 0.89 | RGS1 |
| rs3024505 | 1q32.1 | 0.84 | IL10 (CNTN2) |
| rs1534422 | 2p25.1 | 1.08 | (gene desert) |

| SNP | Chromosome | OR minor allele | Gene of interest |
|------------|------------|-----------------|-----------------------------|
| rs917997 | 2q12.1 | 0.83 | IL18RAP |
| rs1990760 | 2q24.2 | 0.86 | IFIH1 |
| rs3087243 | 2q33.2 | 0.88 | CTLA4 |
| rs11711054 | 3p21.31 | 0.85 | CCR5 |
| rs10517086 | 4p15.2 | 1.09 | (gene desert) |
| rs4505848 | 4q27 | 1.13 | IL2 |
| rs6897932 | 5q13.2 | 0.89 | IL7R |
| rs9268645 | 6p21.32 | 6.8 | MHC |
| rs11755527 | 6q15 | 1.13 | BACH2 |
| rs9388489 | 6q22.32 | 1.17 | C6orf173 |
| rs2327832 | 6q23.3 | 0.90 | TNFAIP3 |
| rs1738074 | 6q25.3 | 0.92 | TAGAP |
| rs7804356 | 7p15.2 | 0.88 | SKAP2 |
| rs4948088 | 7p12.1 | 0.77 | COBL |
| rs7020673 | 9p24.2 | 0.88 | GLIS3 |
| rs12251307 | 10p15.1 | 1.61 | IL2RA |
| rs11258747 | 10p15.1 | 0.84 | PRKCQ |
| rs10509540 | 10q23.31 | 0.75 | RNLS |
| rs7111341 | 11p15.5 | 2.38 | INS (TH) |
| rs4763879 | 12p13 | 1.09 | CD69 |
| rs2292239 | 12q13.2 | 1.31 | ERBB3 |
| rs1678536 | 12q13.3. | | Multiple (MMP19-LOCx-GSTPP) |
| rs3184504 | 12q24.12 | 1.28 | SH2B3 |
| rs1465788 | 14q24.1 | 0.86 | C14orf181 |
| rs4900384 | 14q32.2 | 1.09 | (0; gene desert) |
| rs3825932 | 15q25.1 | 0.86 | CTSH |
| rs12708716 | 16p13.13 | 0.81 | CLEC16A |
| rs12444268 | 16p12.3 | 1.10 | UMOD |
| rs4788084 | 16p11.2 | 0.86 | IL27 (NUPR1) |
| rs7202877 | 16q23.1 | 1.28 | CTRB1 |
| rs16956936 | 17p13.1 | 0.92 | DNAH2 |
| rs2290400 | 17q12 | 0.87 | ORMDL3 (GSDML3) |
| rs7221109 | 17q21.2 | 0.95 | SMARCE1 |
| rs1893217 | 18p11.21 | 1.28 | PTPN2 |
| rs763361 | 18q22.2 | 1.16 | CD226 |
| rs425105 | 19q13.32 | 0.86 | PRKD2 |
| rs2281808 | 20p13 | 0.90 | SIRPG |
| rs11203203 | 21q22.3 | 1.13 | UBASH3A |
| rs5753037 | 22q12.2 | 1.10 | LOC729980/HORMAD2 |
| rs229541 | 22q13.1 | 1.12 | C1QTNF6 |
| rs2664170 | Xq28 | 1.16 | GAB3 |

Table 1. SNPs associated with T1D according to Pociot et al. 2010 & Barret et al 2008.

Additional functional studies provided evidence of causality of several genes within established loci, such as several cytokines and their receptors (*IL10*, *IL2*, *IL27*, *IL7R*, *CCR5*, *SH2B3*, *IL18RAP*), immunomodulatory molecules (*IFIH1*, *TLR7-TLR8*, *TAGAP*) and other types of proteins (*PTPN2*, *GLIS3*). However, for the majority of associated regions the most likely causal gene still needs to be identified (Todd, 2010). *IFIH1* gene on chromosome 2q24.2 encodes intracellular pathogen receptor, MDA5, responsible for binding of viral RNA. This binding stimulates the production of type 1 interferon that could enhance cytotoxic activity of T cells on pancreatic β cells. *TLR7-TLR8* genes on chromosome Xp22.2 also encode intracellular receptors for viral RNA. This may explain induction of autoimmune destruction of β cells by numerous viral infections that may raise type 1 interferon levels (Todd, 2010; Pociot et al., 2010; Concannon et al., 2010). *GLIS3* acts as both transcriptional activator and repressor and is specifically involved in the development of pancreatic β cells. Mutations in this gene have been associated with a rare syndrome of neonatal diabetes and congenital hypothyroidism (Grant SF et al., 2009). It is the only gene that shows overlap between T1D and T2D that may be due to its function in the development and/or function of β cells (Todd, 2010).

There are more than 300 candidate genes that are in LD with T1D associated genetic regions. Also, it has been shown that at least 10 T1D associated regions do not contain a functional candidate gene which suggests that distant, long-range gene regulation might underly some of the observed associations. The main focus of current research is to identify causal risk genes and to understand how they influence the disease (Todd, 2010; Pociot et al., 2010). T1DGC is involved in the research of many autoimmune diseases since it is believed that many of them share common genetic background. A genotyping assay called ImmunoChip, that includes ~200 000 SNPs that are expected to be involved or were previously associated with immune reactions, was developed in order to disentangle the genetic background of various autoimmune diseases including T1D (Pociot et al., 2010).

3.5 Genetic markers in prediction and prevention of T1D

Recently, several population studies attempted to stratify children at birth according to their predisposition for T1D development by examining their *HLA* genotypes and insulin gene polymorphisms. Denver, Germany and Finland studies showed that children with the high risk *HLA* genotypes or polymorphisms within the insulin gene have about 50% higher risk of developing T1D-specific antibodies by the age of 5 (Barker et al., 2004, Walter et al., 2003, Hummel et al., 2004, Steck & Rewers, 2011). Assessment of risk can be extended by including other polymorphisms of susceptible genes for T1D development as well as family history of diabetes (Steck & Rewers, 2011). The application of preventive therapy would be focused on those individuals who have the highest genetic predisposition for the T1D development. Animal model of mucosal administration of insulin in the prevention of autoimmune diabetes has proven to be safe. Mucosal administration of insulin as a self-antigen stimulates a protective immune response that has the potential to affect the destructive immune response that would otherwise follow. Low doses of autoantigenes induce regulatory T cells that release inhibitory cytokines, while high doses destroy autoreactive effector T cells. Pre-POINT (Primary Oral Insulin) Trial in siblings with the high genetic risk for T1D, involving several European populations, Canada and USA, plans to investigate such treatment. Children aged 18 months and 7 years with a family history and genetic predisposition to T1D will be enrolled in the study. Conduction of study is planned in two steps: -step one -

genetic screening and determination of serum antibodies: step two - recruitment of selected children and treatment (test of four doses of insulin). As the effect depends on the dose and route at which insulin is administered in primary mucosal insulin therapy, goal of this study is to find a safe dose (applied orally or intranasal) that will induce an immune response to insulin. Pre-POINT studies should continue with POINT study with the determinate set dose. The main objective of this study is to determine whether the use of oral insulin can prevent or delay the onset of T1D in individuals with high genetic risk (Steck & Rewers, 2011, Achenbach et al, 2008).

4. Epigenetics

There is a significant increase in the incidence of T1D in the last 50 years that is mainly explained by changes in the environment. It is believed that environmental factors can affect epigenetic mechanisms of candidate genes expression and development of T1D. Epigenetic mechanisms encouraged with environmental factors can cause identical genotypes to exhibit different phenotypes. The proposed environmental factors that can trigger an autoimmune process involve nutrition and viruses. Nutrients that may trigger epigenetic mechanisms are considered to be substances that provide a methyl group (methionine, choline) or cofactors (folic acid, vitamin B12 and pyridoxal phosphate) required for DNA and histone methylation (Hewagama & Richardson, 2009). Actually, there are three ways in which phenotype can be altered by epigenetic modifications of gene expression: methylation of DNA, histone modification or activation of micro-RNA. It is well known that silencing of gene expression can be achieved by methylation of cytosine in CpG dinucleotides. Acetylation, methylation, phosphorylation and ubiquitination of histones modify the chromatin conformation, which can stimulate or silent gene expression. MicroRNAs bind to mRNA causing degradation before the translation in protein. These mechanisms that alter gene expression may influence development and function of immune system, as well as development, function and recovery of pancreatic β cells. Differentiation of T-helper cells is regulated by a complex epigenetic control. Critical epigenetic process in T-helper cells differentiation is DNA methylation, which can affect the expression of specific cytokines (interferons, interleukins) and encourage autoreactivity. Development, function and regeneration of pancreatic β cells largely depend on the genetic profile that will be expressed. Progressive decline of pancreatic β cells in type 1 and type 2 diabetes is strongly associated with the expression of genes responsible for the development and function of β cells. It is shown that the activity of the insulin gene is dependent on mechanisms of histone acetylation and methylation. It was also shown that the blood glucose concentration can affect the activity of enzymes that regulate the methylation process, but this seems to be associated with type 2 diabetes (MacFarlane et al., 2009).

5. Environment

Numerous environmental factors are implicated in T1D disease development in genetically susceptible individuals. Many of these factors act in uterine life, infancy and early childhood and are namely associated with viral infections and diet (Norris, 2010; Roivainen & Klingel, 2010).

Viral infections are considered to be the major environmental factors predisposing to T1D. Rotaviruses, adenoviruses, retroviruses, reoviruses, cytomegalovirus, Epstein-Barr virus,

mumps virus and rubella virus are the ones that have been implicated in T1D pathogenesis but the most risk ones are human enteroviral (HEV) intestinal infections. Coxsackievirus and echovirus serotypes of HEV infections, are highly cytotoxic and can cause β cell cytolysis and activate innate and adaptive immune system but can also activate autoreactive T cells (Roivainen & Klingel, 2010). A recent study examined autoimmune microbiome for T1D and came out with conclusion that microbiomes of healthy children differ to those of children that develop T1D later in their lives. This means that microbiome could be used as bacterial marker for the early T1D diagnosis. Also, the “healthy” microbiome could be used in the prevention of T1D development in children at genetically high risk of developing disease (Giongo et al. 2011).

The maternal diet during pregnancy, such as vegetable and vitamin D consumption, and wheat, cow’s milk and omega-3 fatty acids early exposures in life are speculated to have a role in the aetiology of the disease. Introduction of cow’s milk and meat prior 6 months of age have shown to have risk effects. Likewise, cereal, gluten or wheat antigens may cause an aberrant response in developing immune system. Some other factors may have protective effects against T1D development, such as early introduction of vegetable oil and high omega-3 fatty acid intake (Norris, 2010). Huge international collaborative effort, The Environmental Determinants of Diabetes in the Young (TEDDY) (<http://teddy.epi.usf.edu/>), was developed with the aim of identifying environmental factors that modify risk for T1D (TEDDY, 2008).

It is generally considered that environmental and behavioural factors have stronger effects on disease development than genetic loci itself, but it is very hard to accurately identify and measure them. The largest effects are expected from gene and environment interaction in individuals that are genetically at high-risk for disease development (Clayton, 2009).

6. Conclusion

The main genetic predisposition for developing T1D comes from the HLA region. There are currently identified additional ~50 non-HLA loci that predispose to risk of T1D. It is expected that the remaining T1D susceptible loci will be explained by additional common and rare genetic variants, structural polymorphisms, gene-gene and gene-environment interactions and epigenetic events. A role of associated genes and their protein products in disease aetiology is under intense investigation but the candidacy of many loci implicate to the combined effect of adaptive and innate immune action in insulin-producing β cell destruction. Further genetic studies performed on much bigger datasets comprising tens of thousands of individuals, detailed genetic mapping, genotype-phenotype correlation studies and other functional studies will be crucial in deciphering a complete genetic architecture of T1D and understanding the disease mechanisms. The main goal of genetic research is to link research findings with advances in therapy such as screening of individuals and implementation of preventive measures to those with high genetic predisposition to T1D and development of new, more efficient treatments and therapies. Detection of major pathways in the development of T1D opens up new therapeutic targets, development of more efficient treatments and individual approaches to patients.

7. Acknowledgement

We would like to thank Marina Pehlić, MD and Ante Kokan for their help in preparation of figures.


8. References

- 1000 Genomes Project Consortium, Durbin, R.M., Abecasis, G.R., Altshuler, D.L., Auton, A., Brooks, L.D., Durbin, R.M., Gibbs, R.A., Hurles, M.E. & McVean, G.A. (2010). A map of human genome variation from population-scale sequencing. *Nature*, Vol.467, No.7319, (Oct 2010), pp.1061-1073, ISSN 0028-0836
- Achenbach P, Barker J, Bonifacio E; Pre-POINT Study Group. (2008). Modulating the natural history of type 1 diabetes in children at high genetic risk by mucosal insulin immunization. *Current diabetes reports*, Vol.8, No.2, (Apr 2008), pp.87-93, ISSN 1534-4827
- Anderson, C.A., Pettersson, F.H., Clarke, G.M., Cardon, L.R., Morris, A.P., Zondervan, K.T. (2010). Data quality control in genetic case-control association studies. *Nature Protocols*, Vol.5, No.9, (Sep 2010), pp.1564-1573, ISSN 1754-2189
- de Bakker, P.I., Neale, B.M., Daly, M.J. (2010). Meta-analysis of genome-wide association studies. *Cold Spring Harbor Protocols*, Vol.6, (Jun 2010), pdb.top81, ISSN 1559-6095
- Barker, J.M., Goehrig, S.H., Barriga, K., Hoffman, M., Slover, R., Eisenbarth, G.S., Norris, J.M., Klingensmith, G.J., Rewers, M, DAISY study. (2004). Clinical characteristics of children diagnosed with type 1 diabetes through intensive screening and follow-up. *Diabetes Care*, Vol.27, No.6, (Jun 2004), pp.1399-404, ISSN 0149-5992
- Barrett, J.C., Clayton, D.G., Concannon, P., Akolkar, B., Cooper, J.D., Erlich, H.A., Julier, C., Morahan, G., Nerup, J., Nierras, C., Plagnol, V., Pociot, F., Schuilenburg, H., Smyth, D.J., Stevens, H., Todd, J.A., Walker, N.M. & Rich, S.S. (2009). Type 1 Diabetes Genetics Consortium. Genome-wide association study and meta-analysis find that over 40 loci affect risk of type 1 diabetes. *Nature Genetics*, Vol.41, No.6, (Jan 2009), pp.703-707, ISSN 1061-4036
- Bell, G.I., Horita, S. & Karam, J.H. (1984). A polymorphic locus near the human insulin gene is associated with insulin-dependent diabetes mellitus. *Diabetes*, Vol.33, No.2, (Feb 1984), pp.176-183, ISSN 0012-1797
- van Belle, T.L., Coppieters, K.T., von Herrath, M.G. (2011). Type 1 diabetes: etiology, immunology, and therapeutic strategies. *Physiological reviews*, Vol. 91, No.1, (Jan 2011), pp. 79-118. ISSN 0031-9333
- Bluestone, J.A., Herold, K. & Eisenbarth, G. (2010). Genetics, pathogenesis and clinical interventions in type 1 diabetes. *Nature*, Vol.464, No.7293, (Apr 2010), pp.1293-1300, ISSN 0028-0836
- Bottini, N., Musumeci, L., Alonso, A., Rahmouni, S., Nika, K., Rostamkhani, M., MacMurray, J., Meloni, G.F., Lucarelli, P., Pillecchia, M., Eisenbarth, G.S., Comings, D. & Mustelin, T. (2004). A functional variant of lymphoid tyrosine phosphatase is associated with type I diabetes. *Nature Genetics*, Vol.36, No.4, (Apr 2004), pp.337-338, ISSN 1061-4036
- Clayton, D.G. (2009). Prediction and interaction in complex disease genetics: experience in type 1 diabetes. *PLoS Genetics*, Vol.5, No.7, (Jul 2009), e1000540, ISSN 1553-7390
- Concannon, P., Rich, S.S. & Nepom, G.T. (2009). Genetics of type 1A diabetes. *The New England Journal of Medicine*, Vol.360, No.16, (Apr 2009), pp.1646-1654, ISSN 0028-4793
- Cui, Y., Li, G., Li, S. & Wu, R. (2010). Designs for linkage analysis and association studies of complex diseases. *Methods in Molecular Biology*, Vol.620, Part.2, pp.219-242, ISSN 1064-3745

- Durinovic-Belló, I., Wu, R.P., Gersuk, V.H., Sanda, S., Shilling, H.G., Nepom, G.T. (2010). Insulin gene VNTR genotype associates with frequency and phenotype of the autoimmune response to proinsulin. *Genes and immunity*, Vol.11, No.2, (Mar 2010), pp.188-93, ISSN 1466-4879
- Eyre, S., Hinks, A., Bowes, J., Flynn, E., Martin, P., Wilson, A.G., Morgan, A.W., Emery, P., Steer, S., Hocking, L.J., Reid, D.M., Harrison, P., Wordsworth, P., Yorkshire Early Arthritis Consortium, Biologics in RA Control Consortium, Thomson, W., Worthington, J., Barton, A. (2010). Overlapping genetic susceptibility variants between three autoimmune disorders: rheumatoid arthritis, type 1 diabetes and coeliac disease. *Arthritis Research & Therapy*, Vol.12, No.5, (2010), R175, ISSN 1478-6354
- Giongo, A., Gano, K.A., Crabb, D.B., Mukherjee, N., Novelo, L.L., Casella, G., Drew, J.C., Ilonen, J., Knip, M., Hyöty, H., Veijola, R., Simell, T., Simell, O., Neu, J., Wasserfall, C.H., Schatz, D., Atkinson, M.A., Triplett, E.W. (2011). Toward defining the autoimmune microbiome for type 1 diabetes. *Multidisciplinary Journal of Microbial Ecology*, Vol.5, No.1, (Jan 2011), pp. 82-91, ISSN 1751-7362
- Grant, S.F., Qu, H.Q., Bradfield, J.P., Marchand, L., Kim, C.E., Glessner, J.T., Grabs, R., Taback, S.P., Frackelton, E.C., Eckert, A.W., Annaiah, K., Lawson, M.L., Otieno, F.G., Santa, E., Shaner, J.L., Smith, R.M., Skraban, R., Imielinski, M., Chiavacci, R.M., Grundmeier, R.W., Stanley, C.A., Kirsch, S.E., Waggott, D., Paterson, A.D., Monos, D.S., DCCT/EDIC Research Group, Polychronakos, C., Hakonarson, H. (2009). Follow-up analysis of genome-wide association data identifies novel loci for type 1 diabetes. *Diabetes*, Vol.58, No.1, (Jan 2009), pp.:290-5, 0012-1797
- Hewagama, A., Richardson, B. (2009). The genetics and epigenetics of autoimmune diseases. *Journal of autoimmunity*. Vol. 3, No.1, (Aug 2009), pp.3-11, ISSN 0896-8411
- Hummel, M., Bonifacio, E., Schmid, S., Walter, M., Knopff, A., Ziegler, A.G. (2004). Brief communication: early appearance of islet autoantibodies predicts childhood type 1 diabetes in offspring of diabetic parents. *Annals of internal medicine*, Vol.140, No.11, (Jun 2004), pp.82-6, ISSN 0003-4819
- International HapMap Consortium, Frazer, K.A., Ballinger, D.G. et al. (2007). A second generation human haplotype map of over 3.1 million SNPs. *Nature*, Vol.18, No.449, (Oct 2007), pp.851-861, 0028-0836
- Kere, J. (2010). Genetics of complex disorders. *Biochemical and Biophysical Research Communications*. Vol.21, No.396, (May 2010), pp.143-6, ISSN 0006-291X
- Lander, E.S. (2011). Initial impact of the sequencing of the human genome. *Nature*, Vol.470, No.7333, (Feb 2011), pp.187-197, ISSN 0028-0836
- Liston, A., Gray, D.H., Lesage, S., Fletcher, A.L., Wilson, J., Webster, K.E., Scott, H.S., Boyd, R.L., Peltonen, L., Goodnow, C.C. (2004). Gene dosage-limiting role of Aire in thymic expression, clonal deletion, and organ-specific autoimmunity. *The Journal of experimental medicine*. Vol.18, No.8, (Oct. 2004), pp.1015-26, ISSN 0022-1007
- Lowe, C.E., Cooper, J.D., Brusko, T., Walker, N.M., Smyth, D.J., Bailey, R., Bourget, K., Plagnol, V., Field, S., Atkinson, M., Clayton, D.G., Wicker, L.S. & Todd, J.A. (2007). Large-scale genetic fine mapping and genotype-phenotype associations implicate polymorphism in the IL2RA region in type 1 diabetes. *Nature Genetics*, Vol.39, No.9, (Sep 2007), pp.1074-1082, ISSN 1061-4036

- Maahs, D.M., West, N.A., Lawrence, J.M., Mayer-Davis, E.J. (2010). Epidemiology of type 1 diabetes. *Endocrinology and Metabolism Clinics of North America*, Vol.39, No.3, (Sep 2010), pp.481-497, ISSN 0889-8529
- MacFarlane, A.J., Strom, A. & Scott, F.W. (2009). Epigenetics: deciphering how environmental factors may modify autoimmune type 1 diabetes. *Mammalian Genome*, Vol.20, No.9-10, (Sep-Oct 2009), pp.624-632, ISSN 0938-8990
- Manolio, T.A., Collins, F.S., Cox, N.J., Goldstein, D.B., Hindorff, L.A., Hunter, D.J., McCarthy, M.I., Ramos, E.M., Cardon, L.R., Chakravarti, A., Cho, J.H., Guttmacher, A.E., Kong, A., Kruglyak, L., Mardis, E., Rotimi, C.N., Slatkin, M., Valle, D., Whittemore, A.S., Boehnke, M., Clark, A.G., Eichler, E.E., Gibson, G., Haines, J.L., Mackay, T.F., McCarroll, S.A. & Visscher, P.M. (2009). Finding the missing heritability of complex diseases. *Nature*, Vol.461, No.7265, (Oct 2009), pp.747-753, ISSN 0028-0836
- Marchini, J. & Howie, B. (2010). Genotype imputation for genome-wide association studies. *Nature Reviews Genetics*, Vol.11, No.7, (Jun 2010), pp.499-511, ISSN 1471-0056
- McCarthy, M.I., Abecasis, G.R., Cardon, L.R., Goldstein, D.B., Little, J., Ioannidis, J.P. & Hirschhorn, J.N. (2008). Genome-wide association studies for complex traits: consensus, uncertainty and challenges. *Nature Reviews Genetics*, Vol.9, No.5, (May 2008), pp.356-369, ISSN 1471-0056
- Mehers, K.L., Gillespie, K.M. (2008). The genetic basis for type 1 diabetes. *British medical bulletin*, Vol.88, No.1, (2008), 115-29, ISSN 0007-1420
- Nisticò, L., Buzzetti, R., Pritchard, L.E., Van der Auwera, B., Giovannini, C., Bosi, E., Larrad, M.T., Rios, M.S., Chow, C.C., Cockram, C.S., Jacobs, K., Mijovic, C., Bain, S.C., Barnett, A.H., Vandewalle, C.L., Schuit, F., Gorus, F.K., Tosi, R., Pozzilli, P. & Todd, J.A. (1996). The CTLA-4 gene region of chromosome 2q33 is linked to, and associated with, type 1 diabetes. Belgian Diabetes Registry. *Human Molecular Genetics*, Vol.5, No.7, (Jul 1996), pp.1075-1080, ISSN 0964-6906
- Norris, J.M. (2010). Infant and childhood diet and type 1 diabetes risk: recent advances and prospects. *Current diabetes reports*, Vol.10, No.5, (Oct 2010), pp.345-349, ISSN 1534-4827
- Ounissi-Benkhalha, H., Polychronakos, C. (2008). The molecular genetics of type 1 diabetes: new genes and emerging mechanisms. *Trends in molecular medicine*, Vol.14, No.6, (Jun 2008), pp.268-75, ISSN 1471-4914
- Pharoah, P.D., Antoniou, A.C., Easton, D.F., Ponder, B.A. (2008). Polygenes, risk prediction, and targeted prevention of breast cancer. *New England Journal of Medicine*, Vol.358, No.26, (Jun 2008), pp.2796-2803, ISSN 0028-4793
- Pociot, F., Akolkar, B., Concannon, P., Erlich, H.A., Julier, C., Morahan, G., Nierras, C.R., Todd, J.A., Rich, S.S. & Nerup, J. (2010). Genetics of type 1 diabetes: what's next? *Journal of Diabetes*, Vol.59, No.7, (Jul 2010), pp.1561-1571, ISSN 1753-0407
- Roivainen, M. & Klingel K. (2010). Virus infections and type 1 diabetes risk. *Current diabetes reports*, Vol.10, No.5, (Oct 2010), pp.350-356, ISSN 1534-4827
- Romeo, S., Pennacchio, L.A., Fu, Y., Boerwinkle, E., Tybjaerg-Hansen, A., Hobbs, H.H. & Cohen, J.C. (2007). Population-based resequencing of ANGPTL4 uncovers variations that reduce triglycerides and increase HDL. *Nature Genetics*, Vol. 9, No.4, (Apr 2007), pp.513-516, ISSN 1061-4036
- Sabatti, C., Service, S.K., Hartikainen, A.L., Pouta, A., Ripatti, S., Brodsky, J., Jones, C.G., Zaitlen, N.A., Varilo, T., Kaakinen, M., Sovio, U., Ruokonen, A., Laitinen, J., Jakkula, E., Coin,

- L., Hoggart, C., Collins, A., Turunen, H., Gabriel, S., Elliot, P., McCarthy, M.I., Daly, M.J., Järvelin, M.R., Freimer, N.B. & Peltonen, L. (2009). Genome-wide association analysis of metabolic traits in a birth cohort from a founder population. *Nature Genetics*, Vol.41, No.1, (Jan 2009), pp. 35-46, ISSN 1061-4036
- Skrodenienė, E., Marčiulionytė, D., Padaiga, Z., Jašinskienė, E., Sadauskaitė-Kuehne, V., Sanjeevi, C.B., Ludvigsson, J. (2010). HLA class II alleles and haplotypes in Lithuanian children with type 1 diabetes and healthy children (HLA and type 1 diabetes). *Medicina (Kaunas)*, Vol.46, No.8, (2010), pp. 505-10, ISSN 1010-660X
- Smith, J.G., Newton-Cheh, C. (2009). Genome-wide association study in humans. *Methods in Molecular Biology*, Vol.573, (2009), pp.231-258, ISSN 1064-3745
- Smyth, D.J., Cooper, J.D., Bailey, R., Field, S., Burren, O., Smink, L.J., Guja, C., Ionescu-Tirgoviste, C., Widmer, B., Dunger, D.B., Savage, D.A., Walker, N.M., Clayton, D.G. & Todd, J.A. (2006). A genome-wide association study of nonsynonymous SNPs identifies a type 1 diabetes locus in the interferon-induced helicase (IFIH1) region. *Nature Genetics*, Vol.38, No.6, (Jun 2006), pp.617-619, ISSN 1061-4036
- Steck AK, Rewers MJ. (2011). Genetics of type 1 diabetes. *Clinical chemistry*, Vol.57, No.2, (Feb 2011), pp.176-85, ISSN 0009-9147
- TEDDY Study Group. (2008). The Environmental Determinants of Diabetes in the Young (TEDDY) Study. *Annals of the New York Academy of Sciences*, Vol.150, No.1150, (Dec 2008), 1-13, ISSN 0077-8923
- Todd, J.A. Etiology of type 1 diabetes. (2010). *Immunity*, Vol.32, No.4, (Apr 2010), pp457-467, ISSN 1074-7613
- Vella, A., Cooper, J.D., Lowe, C.E., Walker, N., Nutland, S., Widmer, B., Jones, R., Ring, S.M., McArdle, W., Pembrey, M.E., Strachan, D.P., Dunger, D.B., Twells, R.C., Clayton, D.G., Todd, J.A. (2005). Localization of a type 1 diabetes locus in the IL2RA/CD25 region by use of tag single-nucleotide polymorphisms. *The American Journal of Human Genetics*, Vol.76, No.5, (Maj 2005), pp.773-779, ISSN 0002-9297
- Villaseñor, J., Benoist, C., Mathis, D. (2005). AIRE and APECED: molecular insights into an autoimmune disease. *Immunological reviews*. Vol.204, (Apr 2005), pp.156-64, ISSN 0105-2896.
- Ziegler, A.G. & Nepom, G.T. (2010). Prediction and pathogenesis in type 1 diabetes. *Immunity*, Vol.32, No.4, (Apr 2010), pp.468-478, ISSN 1074-7613
- Zondervan, K.T., Cardon, L.R. (2007). Designing candidate gene and genome-wide case-control association studies. *Nature Protocols*, Vol.2, No.10, (Oct 2007), pp.92-2501, ISSN 1754-2189
- Walter, M., Albert, E., Conrad, M., Keller, E., Hummel, M., Ferber, K., Barratt, B.J., Todd, J.A., Ziegler, A.G., Bonifacio, E. (2003). IDDM2/insulin VNTR modifies risk conferred by IDDM1/HLA for development of Type 1 diabetes and associated autoimmunity. *Diabetologia*, Vol.46, No.5, (May 2003), pp.712-20, ISSN 0012-186X
- Wang, J., Li, X., Jia, Z., Tian, Y., Yu, J., Bao, L., Wu, Y., Ni, B. (2010). Reduced FOXP3 expression causes IPEX syndrome onset: An implication from an IPEX patient and his disease-free twin brother. *Clinical Immunology*, Vol.137, No.1, (Oct 2010), pp. 178-80, ISSN 1744-666X
- Wang, K., Li, M., Hakonarson, H. (2010). Analysing biological pathways in genome-wide association studies. *Nature Reviews Genetics*, Vol.11, No.12, (Dec 2010), pp.843-854, ISSN 1471-0056



Type 1 Diabetes - Pathogenesis, Genetics and Immunotherapy

Edited by Prof. David Wagner

ISBN 978-953-307-362-0

Hard cover, 660 pages

Publisher InTech

Published online 25, November, 2011

Published in print edition November, 2011

This book is a compilation of reviews about the pathogenesis of Type 1 Diabetes. T1D is a classic autoimmune disease. Genetic factors are clearly determinant but cannot explain the rapid, even overwhelming expanse of this disease. Understanding etiology and pathogenesis of this disease is essential. A number of experts in the field have covered a range of topics for consideration that are applicable to researcher and clinician alike. This book provides apt descriptions of cutting edge technologies and applications in the ever going search for treatments and cure for diabetes. Areas including T cell development, innate immune responses, imaging of pancreata, potential viral initiators, etc. are considered.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Tatijana Zemunik and Vesna Boraska (2011). Genetics of Type 1 Diabetes, Type 1 Diabetes - Pathogenesis, Genetics and Immunotherapy, Prof. David Wagner (Ed.), ISBN: 978-953-307-362-0, InTech, Available from: <http://www.intechopen.com/books/type-1-diabetes-pathogenesis-genetics-and-immunotherapy/genetics-of-type-1-diabetes>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2011 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](https://creativecommons.org/licenses/by/3.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen