

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

186,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Defining the Deletion Size in Williams-Beuren Syndrome by Fluorescent *In Situ* Hybridization with Bacterial Artificial Chromosomes

Marc De Braekeleer^{1,2,3} et al.*

¹Faculté de Médecine et des Sciences de la Santé, Université de Brest, Brest

²Institut National de la Santé et de la Recherche Médicale (INSERM), U613, Brest

³Service de Cytogénétique, Cytologie et Biologie de la Reproduction
Hôpital Morvan, CHRU Brest, Brest
France

1. Introduction

Williams-Beuren syndrome (WBS, MIM No. 194050) is a contiguous gene deletion syndrome that was described independently by Williams et al. (1961) in patients with supra-aortic stenosis, growth retardation and an unusual facial appearance (Williams et al., 1961) and by Beuren et al. (1962) in patients having the same features as well as dental anomalies and friendly personality (Beuren et al., 1962).

The clinical picture of WBS includes a characteristic craniofacial dysmorphism with broad forehead, periorbital fullness, flat nasal bridge, broad nasal tip, long philtrum, full lips and lower cheeks, micrognathia, wide mouth and stellate irides, growth retardation, cardiovascular anomalies (mostly supra-aortic stenosis and pulmonary artery stenosis), mild to moderate mental retardation and unique behavioral and neurocognitive profile (relative preservation of linguistic abilities and gross visual-spatial processing dysfunction) (Morris et al., 1988; Pober, 2010). Its incidence is estimated at 1/7,500-1/20,000 (Grimm & Wesselhoeft, 1980; Morris et al., 1988; Stromme et al., 2002).

Williams-Beuren syndrome results from the hemizygous deletion of several genes, including *ELN* (elastin), encompassing 1.55 to 1.84 Mb on chromosome 7q11.23 (Ewart et al., 1993; Meng et al., 1998; Pober, 2010; Schubert & Laccone, 2006; Wang et al., 1999). WBS usually results from de novo deletion. Twenty-six to 28 genes have been identified within the WBS deletion region (Merla et al., 2010; Pober, 2010; Schubert, 2009).

*Audrey Basinko^{1,2,3}, Nathalie Douet-Guilbert^{1,2,3}, Séverine Audebert-Bellanger⁴, Philippe Parent⁴, Clémence Chabay-Vichot^{1,4}, Clément Bovo^{1,2}, Nadia Guéganic^{1,2}, Marie-Josée Le Bris³, Frédéric Morel^{1,2,3}

¹Faculté de Médecine et des Sciences de la Santé, Université de Brest, Brest, France

²Institut National de la Santé et de la Recherche Médicale (INSERM), U613, Brest, France

³Service de Cytogénétique, Cytologie et Biologie de la Reproduction, Hôpital Morvan, CHRU Brest, Brest, France

⁴Département de Pédiatrie et de Génétique Médicale, Hôpital Morvan, CHRU Brest, Brest, France

The WBS deletion region is flanked by highly homologous clusters of genes and pseudogenes organized into low-copy-repeat (LCR) blocks. Unequal meiotic recombination during meiosis can lead to deletion of the WBS region. The unique genetic architecture of the region explains why the size of WBS deletion is almost the same in most of the patients (Baumer et al., 1998; Dutly & Schinzel, 1996; Pober, 2010; Schubert, 2009; Valero et al., 2000).

Several studies have investigated the size of the WBS deletion using multiplex PCR with several microsatellite markers (Brondum-Nielsen et al., 1997; Perez Jurado et al., 1996; Wang et al., 1999; Wu et al., 1998). Most of the patients were found to carry the 1.5 or 1.8 Mb deletion. In the present study, we performed fluorescent *in situ* hybridization (FISH) with Bacterial Artificial Chromosome (BAC) clones in an attempt to map the WBS deletion in 14 WBS patients.

2. Patients and methods

2.1 Patients

Fifteen patients were referred to the cytogenetic laboratory of the Brest University Hospital by medical geneticists, pediatricians and pediatric cardiologist for suspicion of WBS between 2002 and March 2011. The reasons for referral included cardiovascular anomalies, dysmorphism or a combination of signs suspecting WBS (Table 1). However, because of limited cell pellet available for one patient, the study could be performed only on 14 patients.

2.2 Conventional cytogenetics

Metaphase chromosomes were prepared from peripheral blood lymphocytes of the 14 patients after having obtained their parents' informed consent. Standard R banding chromosomal analyses were performed according to the standard procedures and the karyotypes described according to the International System for Cytogenetic Nomenclature (ISCN 2005).

2.3 FISH analyses with commercially available probe

A FISH study using the Vysis Williams Region Probe - LSI ELN SpectrumOrange/LSI D7S486, D7S522 SpectrumGreen was carried out on the metaphase preparations from all 14 patients, as recommended by the manufacturer (Abbott, Rungis, France).

The Williams Region Probe consists of a probe of approximately 180 kb in size for *ELN*, *LIMK1* and the D7S613 locus located in band 7q11.23, labeled in Spectrum Orange, and a control probe for the region containing loci D7S486 and D7S522 located in band 7q31, labeled in Spectrum Green.

2.4 FISH analyses with BAC clones

To delineate the extent of the deletion on chromosome 7, FISH analyses were carried out using BAC clones mapping to the long arm of chromosome 7 (7q11.23).

We identified the BAC clones of interest through the Human Genome Browser Database of the Genome Bioinformatics Group at the University of California at Santa Cruz (<http://genome.ucsc.edu/>). They were then ordered by Internet on the site of the Children's Hospital Oakland Research Institute in Oakland, California (<http://bacpac.chori.org/>).

Patient Nr.	1	2	3	4	5	6	7	8	9	10	11	12	13	14
Sex	F	M	F	M	F	M	M	F	M	M	F	M	F	F
Age at diagnosis	3m	2m	7m	1y4m	3m	1y7m	1y9m	9y11m	8m	2y9m	9m	2m	3m	2m
Reason for referral	Heart anomaly	Heart anomaly	Pulmonary artery stenosis	WBS suspicion	Aortic stenosis	Pulmonary artery stenosis	Heart anomaly	WBS suspicion	Dysmorphism	Dysmorphism	Heart anomaly	Dysmorphism	Heart anomaly	Heart anomaly
Heart														
Ventricular septal defect	-	+	+	-	-	-	+	+	-	-	-	-	-	-
Supra valvular aortic stenosis	+	+	-	+	+	-	-	-	+	-	-	+	-	+
Pulmonary artery stenosis	-	+	+	-	+	+	-	-	-	-	+	+	+	+
Growth & Development														
IUGR	+	+	+	-	+	NA	-	-	+	+	-	-	-	-
Growth retardation	+	+	+	+	+	+	+	+	+	+	+	+	+	-
Psychomotor retardation	+	+	+	+	+	+	+	+	+	+	+	+	+	?
Digestive system														
Neonatal feeding difficulties	+	+	-	+	+	NA	+	-	+	+	+	+	+	-
Inguinal hernia	+	-	-	+	-	NA	-	-	-	+	-	+	-	-
Gastroesophageal reflux	+	-	-	+	+	NA	-	+	-	+	NA	+	+	+
Dysmorphology														
Periorbital oedema	-	+	-	+	+	+	+	+	-	-	+	+	-	+

Table 1. Clinical characteristics of the 14 patients with Williams-Beuren syndrome.

Patient Nr.	1	2	3	4	5	6	7	8	9	10
Sex	F	M	F	M	F	M	M	F	M	M
Large mouth	-	-	+	+	-	-	+	+	+	+
Fine upper lip	-	+	-	-	+	-	+	+	+	+
Flat philtrum	-	+	-	-	-	-	-	+	+	+
Bulbous nose	-	+	+	+	+	+	+	+	+	+
Anteverted nares	-	-	-	-	+	-	+	+	-	-
Low-set dysplastic ears	-	+	-	-	-	-	-	-	+	+
Prominent forehead	-	+	-	+	-	-	-	-	-	-
ORL										
High arched palate or cleft palate	-	+	-	-	NA	NA	+	+	-	+
Ophthalmology										
Stellate irides	+	-	+	-	NA	NA	+	-	-	+
Strabismus ± astigmatism	NA	-	-	-	NA	NA	-	-	+	+

M: month; y: year

WBS suspicion: combination of signs (facial dysmorphology and cardiovascular anomaly) making the diagnosis

IUGR: intra uterine growth retardation

NA: not available

Patient # 14 - ?: patient still too young (4 months old)

When received, bacterial cultures were prepared from a single colony picked from a selective plate in the presence of chloramphenicol. Plasmids were obtained from bacterial cultures grown in the presence of chloramphenicol (10 mg/L). After having lysed bacteria using SDS1%/NaOH 0.2 N, DNA was purified from RNA, proteins and other cellular contaminants. Probes were then labeled by nick translation in Spectrum Orange (Nick Translation Kit, Abbott, Rungis, France) or in FITC (Prime-it Fluor Fluorescence Labeling Kit, Stratagene, Amsterdam, Netherlands). All BAC clones were applied to normal lymphocyte metaphases to confirm their chromosomal location.

After hybridization, the slides were counterstained with 4-6-diamino-2-phenyl-indole-dihydrochloride (DAPI). The preparations were examined using a Zeiss Axio Plan Microscope (Zeiss, Le Pecq, France). Images acquisition was performed using a CCD camera and analyzed using the ISIS program (In Situ Imaging System) (MetaSystems, Altussheim, Germany).

Twenty-two overlapping BACs covering the WBS deletion region and beyond (about 1.9 Mb) were applied on the 14 patients (Table 2).

BAC name	Centromeric start (Mb from telomere)	Telomeric start (Mb from telomere)	BAC length (bp)
RP11-48D17	72,557,570	72,705,204	147,635
RP11-483G21	72,688,731	72,875,946	187,216
RP11-614D7	72,737,634	72,926,932	189,299
RP11-101D2	72,814,548	72,991,974	177,427
RP11-598B14	72,850,682	73,042,752	192,071
RP11-622P13	73,002,073	73,181,256	179,184
RP11-73G23	73,035,513	73,184,825	149,313
RP11-148M21	73,153,473	73,327,919	174,447
RP11-1011-F11	73,166,799	73,342,468	175,670
RP11-1056I4	73,437,240	73,639,567	202,328
RP11-7M12	73,611,329	73,793,367	182,039
RP11-351B3	73,705,662	73,899,619	193,958
RP11-247L6	73,872,593	74,038,131	165,539
RP11-196F10	73,872,610	74,038,131	165,522
RP11-137E8	73,944,720	74,129,587	184,868
RP11-728M8	73,959,533	74,148,428	188,896
RP11-926D5	73,959,536	74,145,706	186,171
RP11-19F19	73,994,434	74,173,462	179,029
RP11-813J7	74,091,733	74,261,309	169,577
RP11-1105J19	74,128,975	74,321,491	192,517
RP11-1094P22	74,129,067	74,321,019	191,953
RP11-379L10	74,307,663	74,480,665	173,003

The base pairs position (bp) are predicted on Build 39 National Center for Biotechnology Information (<http://www.ncbi.nlm.nih.gov>) and assembly February 2009 by The UCSC Genome Browser Database (<http://genome.ucsc.edu/index.html>),

Table 2. BAC library used to delimitate the deletion size in the 14 WBS patients.

3. Results

Fourteen patients were included in the study (Table 1). There were 7 boys and 7 girls. The median age at the time of cytogenetic exam was 7.5 months. Except for a patient who was almost 10 years old, all the other patients were less than 3 years old at the time of cytogenetic analyses, the majority of them being less than 1 year old (9 of 14 patients).

Cardiovascular anomalies and facial dysmorphism were present in all patients, but for one without cardiovascular anomaly and another without dysmorphism (Table 1). Growth and psychomotor retardation was noted in all patients. Stellate irides and/or strabismus was also present in 7 of the 12 patients for whom the data was available. No cognitive nor personality profile was available for the patients, mainly due to their young age.

R-banding conventional cytogenetics showed a normal 46,XY or 46,XX karyotype whereas FISH using the Vysis Williams Region Probe revealed a deletion of the *ELN*, *LIMK1* and D7S613 locus in all 14 patients (Figure 1-A).




Fig. 1. Example of FISH results in a patient with Williams-Beuren syndrome. (A) FISH using the commercially available probe (*ELN* in SpectrumOrange and D7S486, D7S522 in SpectrumGreen). (B) FISH using two BAC clones (RP11-614D7 in green and RP11-1105J19 in red).

Sequential FISH analyses with BAC clones were applied on metaphases of all 14 patients having a deletion of the Vysis Williams Region Probe (Figure 1-B). They showed the centromeric deletion boundary to be located in a 145 kb interval, between RP11-598B14, deleted in 13 patients (P1 to P13), and RP11-48D17, always present (Figure 2). The telomeric deletion boundary was found to be located in a 229 kb interval, between RP11-351B3, deleted in 13 patients (P1 to P13), and RP11-1105J19, always present. Therefore, the minimum and maximum estimated deletion sizes for these 13 patients (P1 to P13) were 1,048,937 bp and 1,423,771 bp, respectively. Using overlapping BAC clones, both centromeric and telomeric boundaries could be refined. FISH signals with BAC clones RP11-614D7 and RP11-101D2, located at the centromeric deletion boundary, and those RP11-

247L6 and RP11-137E8, located at the telomeric deletion boundary, showed decreased intensities. The centromeric and telomeric intervals were reduced to 36 kb and 93 kb, respectively, giving an estimated WBS deletion size of 1.17Mb.




Fig. 2. Size of the deletions in the 14 WBS patients estimated by BAC clones (ELN: elastin gene)

The fourteenth patient (P14) was found to have a centromeric deletion boundary to be located in a 109 kb interval, between RP11-101D2 (deleted) and RP11-48D17 (present). Using overlapping BAC clones RP11-483G21 and RP11-614D7, the centromeric boundary was refined to a 77 kb interval. The telomeric deletion boundary was found to be located at the junction of two overlapping BAC clones (RP11-926D5 and RP11-1105J19). Therefore, the estimated WBS deletion size for this patient was 1.41Mb.

4. Discussion

Molecular genetic studies have shown that hemizyosity of the elastin (*ELN*) gene accounts for the cardiovascular abnormalities observed in WBS patients. The other signs (facial dysmorphism, growth and mental retardation, etc.) observed in WBS are likely to be due to hemizyosity of other genes flanking the *ELN* locus. Although the unique combination of signs and anomalies is highly evocative, WBS patients usually present phenotypic variability, which could be explained by different deletion size and, therefore, by different sets of genes showing hemizyosity.

Several studies tried to define the WBS critical deletion region. Using sequence tagged site (STS) markers flanking the *ELN* gene, Perez-Jurado et al. (1996) and Wu et al. (1998) looked for the minimal deletion region responsible for the WBS phenotype in 123 patients. Markers from D7S489B (also named D7S489U) to D7S1870 were shown to be consistently removed, defining a 2 cM deletion in all informative patients (Perez Jurado et al., 1996; Wu et al., 1998).

Using BAC and PAC (P1-derived artificial chromosome) clones, Meng et al. (1998) constructed a physical map of the common deletion region. They localized the centromeric and telomeric deletion breakpoints to two genomic clones (containing D7S489B and D7S1870) flanking a 1.5 Mb deletion interval (Meng et al., 1998).

Based on the work by Valero et al. (2000) who found that 3 large region-specific segmental duplication or low copy repeat (LCR) elements flanked the common deletion region (Valero et al., 2000), Bayes et al. (2003) defined two common deletion regions in a set of 74 patients with WBS. Most of the patients (95%) had a 1.55 Mb deletion whereas the remaining 5% exhibited a larger deletion of approximately 1.84 Mb (Bayes et al., 2003).

Botta et al. (1999) reported a deletion of about 850 kb in two patients showing the full spectrum of the WBS phenotype (Botta et al., 1999). Furthermore, the analysis of the deletion size in WBS patients with "incomplete" phenotype revealed even more heterogeneity (Antonell et al., 2010). Using quantitative real-time PCR to scan 2.5 Mb of the WBS deletion region at a resolution of 100-300 kb among 65 patients with strong clinical indication of having WBS, Schubert and Laccone (2006) found that 21 patients had a deletion in the WBS region. Nineteen patients had a deletion ranging from 1.4 to 1.8 Mb in size whereas one had a 200 kb deletion and the remaining one a 2.5 Mb deletion (Schubert & Laccone, 2006).

FISH using BAC clones has been extensively used to construct physical maps of the 7q region deleted in WBS and to define the commonly deleted region (Meng et al., 1998; Peoples et al., 2000; Perez Jurado et al., 1996; Wu et al., 1998). However, to our knowledge, no study has been conducted to determine the deletion size in WBS individuals using overlapping BACs. Indeed, the level of resolution is a limit to this technique. A BAC is considered as partially deleted by FISH analysis when the fluorescent signal of this clone is much stronger on one chromosome 7 than on the other. Furthermore, a BAC could be considered as present or absent (not partially deleted) if only a small part of the DNA sequence is removed or kept (usually less than 50 kb). Having resource to overlapping BAC clones can increase the level of confidence but some uncertainty will remain.


5. Conclusion

Thirteen patients had a deletion estimated at 1.17 Mb in size and the remaining patient a deletion size estimated at 1.41 Mb. All the patients had the “classical” WBS phenotype, but for the cognitive profile which was not evaluated. The margins of uncertainty of the centromeric and telomeric boundaries of the deletion were less than 100 kb in all 14 patients, a level obtained by microsatellite analysis or quantitative real-time PCR. Therefore, this approach is efficient to define the deletion size among WBS patients.

6. References

- Antonell, A., Del Campo, M., Magano, L.F., Kaufmann, L., de la Iglesia, J.M., Gallastegui, F., Flores, R., Schweigmann, U., Fauth, C., Kotzot, D., & Perez-Jurado, L.A. (2010) Partial 7q11.23 deletions further implicate GTF2I and GTF2IRD1 as the main genes responsible for the Williams-Beuren syndrome neurocognitive profile. *Journal of Medical Genetics*, Vol. 47, pp. 312-320
- Baumer, A., Dutly, F., Balmer, D., Riegel, M., Tukel, T., Krajewska-Walasek, M., & Schinzel, A.A. (1998) High level of unequal meiotic crossovers at the origin of the 22q11. 2 and 7q11.23 deletions. *Human Molecular Genetics*, Vol. 7, pp. 887-894
- Bayes, M., Magano, L.F., Rivera, N., Flores, R., & Perez Jurado, L.A. (2003) Mutational mechanisms of Williams-Beuren syndrome deletions. *American Journal of Human Genetics*, Vol. 73, pp. 131-151
- Beuren, A.J., Apitz, J., & Harmjanz, D. (1962) Supravalvular aortic stenosis in association with mental retardation and a certain facial appearance. *Circulation*, Vol. 26, pp. 1235-1240
- Botta, A., Novelli, G., Mari, A., Novelli, A., Sabani, M., Korenberg, J., Osborne, L.R., Digilio, M.C., Giannotti, A., & Dallapiccola, B. (1999) Detection of an atypical 7q11.23 deletion in Williams syndrome patients which does not include the STX1A and FZD3 genes. *Journal of Medical Genetics*, Vol. 36, pp. 478-480
- Brondum-Nielsen, K., Beck, B., Gyftodimou, J., Horlyk, H., Liljenberg, U., Petersen, M.B., Pedersen, W., Petersen, M.B., Sand, A., Skovby, F., Stafanger, G., Zetterqvist, P., & Tommerup, N. (1997) Investigation of deletions at 7q11.23 in 44 patients referred for Williams-Beuren syndrome, using FISH and four DNA polymorphisms. *Human Genetics*, Vol. 99, pp. 56-61
- Dutly, F. & Schinzel, A. (1996) Unequal interchromosomal rearrangements may result in elastin gene deletions causing the Williams-Beuren syndrome. *Human Molecular Genetics*, Vol. 5, pp. 1893-1898
- Ewart, A.K., Morris, C.A., Atkinson, D., Jin, W., Sternes, K., Spallone, P., Stock, A.D., Leppert, M., & Keating, M.T. (1993) Hemizygoty at the elastin locus in a developmental disorder, Williams syndrome. *Nature Genetics*, Vol. 5, pp. 11-16
- Grimm, T. & Wesselhoeft, H. (1980) Zur Genetik des Williams-Beuren-Syndroms und der isolierten Form der supravalvulaeren Aortenstenose (Untersuchungen von 128 Familien). *Zeitschrift für Kardiologie*, Vol. 69, pp. 168-172
- ISCN (2005). An International System for Human Cytogenetic Nomenclature. Basel: S. Karger.
- Meng, X., Lu, X., Li, Z., Green, E.D., Massa, H., Trask, B.J., Morris, C.A., & Keating, M.T. (1998) Complete physical map of the common deletion region in Williams

- syndrome and identification and characterization of three novel genes. *Human Genetics*, Vol. 103, pp. 590-599
- Merla, G., Brunetti-Pierri, N., Micale, L., & Fusco, C. (2010) Copy number variants at Williams-Beuren syndrome 7q11.23 region. *Human Genetics*, Vol. 128, pp. 3-26
- Morris, C.A., Demsey, S.A., Leonard, C.O., Dilts, C., & Blackburn, B.L. (1988) Natural history of Williams syndrome: physical characteristics. *Journal of Pediatrics*, Vol. 113, pp. 318-326
- Peoples, R., Franke, Y., Wang, Y.K., Perez-Jurado, L., Paperna, T., Cisco, M., & Francke, U. (2000) A physical map, including a BAC/PAC clone contig, of the Williams-Beuren syndrome--deletion region at 7q11.23. *American Journal of Human Genetics*, Vol. 66, pp. 47-68
- Perez Jurado, L.A., Peoples, R., Kaplan, P., Hmal, B.C.J., & Francke, U. (1996) Molecular definition of the chromosome 7 deletion in Williams syndrome and parent-of-origin effects on growth. *American Journal of Human Genetics*, Vol. 59, pp. 781-792
- Pober, B.R. (2010) Williams-Beuren syndrome. *New England Journal of Medicine*, Vol. 362, pp. 239-252
- Schubert, C. (2009) The genomic basis of the Williams-Beuren syndrome. *Cellular and Molecular Life Sciences*, Vol. 66, pp. 1178-1197
- Schubert, C. & Laccone, F. (2006) Williams-Beuren syndrome: determination of deletion size using quantitative real-time PCR. *International Journal of Molecular Medicine*, Vol. 18, pp. 799-806
- Stromme, P., Bjornstad, P.G., & Ramstad, K. (2002) Prevalence estimation of Williams syndrome. *Journal of Child Neurology*, Vol. 17, pp. 269-271
- Valero, M.C., de Luis, O., Cruces, J., & Perez Jurado, L.A. (2000) Fine-scale comparative mapping of the human 7q11.23 region and the orthologous region on mouse chromosome 5G: the low-copy repeats that flank the Williams-Beuren syndrome deletion arose at breakpoint sites of an evolutionary inversion(s). *Genomics*, Vol. 69, pp. 1-13
- Wang, M.S., Schinzel, A., Kotzot, D., Balmer, D., Casey, R., Chodirker, B.N., Gyftodimou, J., Petersen, M.B., Lopez-Rangel, E., & Robinson, W.P. (1999) Molecular and clinical correlation study of Williams-Beuren syndrome: No evidence of molecular factors in the deletion region or imprinting affecting clinical outcome. *American Journal of Medical Genetics*, Vol. 86, pp. 34-43
- Williams, J.C., Barratt-Boyes, B.G., & Lowe, J.B. (1961) Supravalvular aortic stenosis. *Circulation*, Vol. 24, pp. 1311-1318
- Wu, Y.Q., Sutton, V.R., Nickerson, E., Lupski, J.R., Potocki, L., Korenberg, J.R., Greenberg, F., Tassabehji, M., & Shaffer, L.G. (1998) Delineation of the common critical region in Williams syndrome and clinical correlation of growth, heart defects, ethnicity, and parental origin. *American Journal of Medical Genetics*, Vol. 78, pp. 82-89



Bacterial Artificial Chromosomes

Edited by Dr Pradeep Chatterjee

ISBN 978-953-307-725-3

Hard cover, 148 pages

Publisher InTech

Published online 25, November, 2011

Published in print edition November, 2011

This book focuses on the numerous applications of Bacterial Artificial Chromosomes (BACs) in a variety of studies. The topics reviewed range from using BAC libraries as resources for marsupial and monotreme gene mapping and comparative genomic studies, to using BACs as vehicles for maintaining the large infectious DNA genomes of viruses. The large size of the insert DNA in BACs and the ease of engineering mutations in that DNA within the bacterial host, allowed manipulating the BAC-viral DNA of Varicella-Zoster Virus. Other reviews include the maintenance and suitable expression of foreign genes from a Baculovirus genome, including protein complexes, from the BAC-viral DNA and generating vaccines from BAC-viral DNA genomes of Marek's disease virus. Production of multi-purpose BAC clones in the novel *Bacillus subtilis* host is described, along with chapters that illustrate the use of BAC transgenic animals to address important issues of gene regulation in vertebrates, such as functionally identifying novel cis-acting distal gene regulatory sequences.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Audrey Basinko, Nathalie Douet-Guilbert, Séverine Audebert-Bellanger, Philippe Parent, Clémence Chabay-Vichot, Clément Bovo, Nadia Guéganic, Marie-Josée Le Bris, Frédéric Morel and Marc De Braekeleer (2011). Defining the Deletion Size in Williams-Beuren Syndrome by Fluorescent In Situ Hybridization with Bacterial Artificial Chromosomes, *Bacterial Artificial Chromosomes*, Dr Pradeep Chatterjee (Ed.), ISBN: 978-953-307-725-3, InTech, Available from: <http://www.intechopen.com/books/bacterial-artificial-chromosomes/defining-the-deletion-size-in-williams-beuren-syndrome-by-fluorescent-in-situ-hybridization-with-bac>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2011 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen