

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

186,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Marine Macrophytic Algae of the Western Sector of North Pacific (Russia)

Olga N. Selivanova

*Kamchatka Branch of Pacific Institute of Geography,
Far Eastern Division of the Russian Academy of Sciences,
Russia*

1. Introduction

Marine algal flora of the western coasts of Bering Sea is studied non-uniformly, i.e. to a variable degree in different areas. Perhaps, algae of the Commander Islands are studied more thoroughly as compared to the other areas of the Russian Pacific. The data on marine algae of the Islands were presented in many published works by the Russian authors. The comprehensive survey of the literature on the Islands' marine flora was given in our papers (Selivanova & Zhigadlova, 1997 a, b, c). Thereafter we continued our floristic and taxonomic studies on the Commander Islands and published many new papers: Selivanova, 2001 a, b; 2008 a, b, c; 2009; Selivanova & Zhigadlova, 2000; 2003; 2010; Zhigadlova, 2009.

In contrast to Commander Islands algae of the continental part of the Bering Sea are studied rather poorly. Though floristic investigations began there above 200 years ago information on algal flora and structure of benthic communities of this area is still scanty. Remoteness and inaccessibility, severe climate and ice conditions, and a short navigation season make this area very inconvenient for natural studies, so they were episodic and uncoordinated. Practically no seasonal field observations have been conducted there, no marine biological stations have ever existed, and scientific expeditions have been infrequent and sporadic. Therefore information on benthic algae of the area is limited. Special publications on this subject are rare (e.g. Vinogradova, 1973a; 1978; Perestenko, 1988; Zhigadlova & Selivanova, 2004), although data on the marine algae of this area may be found in some general taxonomic, floristic and hydrobiological studies (e.g. Kongisser, 1933; Petrov, 1972, 1973; Vinogradova, 1973b, 1974, 1979; Vinogradova et al., 1978; Vinogradova & Perestenko, 1978; Kussakin & Ivanova, 1978; Perestenko, 1994).

The most detailed and contemporary information on marine algae of the Russian part of Bering Sea is given in our recent publication (Selivanova & Zhigadlova, 2010). This work however was published in Russian in a book with a small number of copies so it is scarcely available to phycologists outside Russia. Partially data on algae of western part of Bering Sea (from Ozernoi Gulf to Dezhnev Cape) was presented earlier in my work (Selivanova, 2002), its electronic version is still available at the address: <http://ucjeps.berkeley.edu/constancea/83>

Data from this publication are cited in the species list of Bering Sea algae presented in a current paper. However some part of this information has already become out-of date due to considerable and fast changes in the algal systematics caused by the application of

molecular-genetic studies in phycology in the latest decade. So our main purpose was to make the inventory of the flora of the western Bering Sea on the basis of our personal collections in conformity with new world data in algal taxonomy and nomenclature.

We studied benthic algae of the western part of Bering Sea that belongs to the administrative region – Kamchatskii Krai, including Koryak Autonomous District (from Ozernoi Gulf to Dezhnev Bay) (Fig. 1). Water areas northwards of it located on the territory of Chukchi Autonomous District (Anadyrskii Gulf, north of Dezhnev Bay to Bering Strait) were unstudied by us. However data on the algae of this area may be found in the papers of Vinogradova (1973a, b), Tolstikova (1974), Kussakin & Ivanova (1978); Perestenko (1988; 1994) but the inventory of marine algae of this region is uncomplete, and additional floristic and taxonomic studies are necessary.

It should be noted that in spite of better knowledge on algae of the Commander Islands as compared to other areas of western coasts of Bering Sea inventory of their flora is still not finished either because of the difficulties of collection of algae, especially subtidal ones, in this remote hard-to-reach and little-inhabited area. Besides that the general taxonomic base is insufficient and there are still unsolved nomenclatural problems.

That is why the species list of marine benthic algae of the Russian sector of Bering Sea presented in the current work in tables (Table 1-3) should be still treated as preliminary. It is not only due to the permanently renovated information on algal systematics but also due to more careful examination and re-identification of our material and finding of new species especially having sub-microscopic size.

Moreover collections of algae on the studied area were carried out in a short period of time (mostly in August-September) that also makes our list insufficient and obviously its completion needs additional field expeditions in other seasons. Necessity to continue studies on marine algae of the western coasts of Bering Sea is also caused by the threat of extermination of algal species as a result of uncontrolled harvest of marine bioresources. The fact is that rich and diverse vegetation on the shelf of this area attracts increasing attention of the commercial sea fishery organizations in the recent decade of years.

2. Materials and methods

The list of algae presented in the paper is based on phycological material collected by the author during expedition of the Laboratory of Hydrobiology of KBPIG in the Bering Sea in 1988. The material was collected from August through October on the littoral fringe during low tides, with the help of a long hook called “kanza” from the depths of 1 to 3 m, and with usage of SCUBA technique from the depths of 1 to 30 m and with a dredge from deeper waters (up to 120 m). The material from Commander Islands was collected by the author during expeditions of the same laboratory annually in 1986-1992 and incidentally by individual collectors in 1997, 2007-2010. Algae were collected from April through September on the littoral fringe during low tides, and with use of SCUBA from the depths of 1-30 m. Algae cast ashore were also picked up. Material was sectioned freehand with a razor blade, placed in a drop of fresh water on the slides and examined using the light microscope. The sections were studied uncolored or stained with iodine Lugol solution. The processing of collections was conducted at Kamchatka Branch of Pacific Institute of Geography (Petropavlovsk-Kamchatskii, Russia). Material is stored in the herbarium of the above mentioned institute.




Fig. 1. A schematic map of the part of Bering Sea including Commander Islands. Arrows show the direction of the Eastern Kamchatka and Alaskan currents. Numbers correspond to the specially protected nature areas (SPNA-s) within the studied area: 1 – Commander state biosphere reserve; 2 - Reserve “Karaginskii Island”; 3 - State nature reserve of federal significance “Koryakskii”; 4 - Nature park “Beringia”. Abbreviations correspond to the names of major water areas of Bering Sea where algae were collected: NR- Navariniskii Region, O – Olyutorskii Gulf, K- Karaginskii Gulf, OZ – Ozernoi Gulf.

3. Species composition and distribution of marine macroalgae of the upper part of the western North Pacific (continental coast of Bering Sea and Commander Islands)

Up to the present day the general list of marine benthic algae of the western coasts of Bering Sea including Commander Islands based on our own materials contains 193 species, but taking into consideration data from the literature their number totals to 209 ones: 36 species of Chlorophyta, 129 Rhodophyta and 44 Ochrophyta (class Phaeophyceae). The list is considerably enlarged as compared to the previous publications (Selivanova and Zhigadlova, 1997a, b, c; Selivanova, 2002) due to new finds of algal samples and description of new taxa.

The author’s own data on the species composition of poorly studied areas of Bering Sea are supplemented with the generalized information from separate literature sources only if this information seems to be reliable. In quoting the data of the other authors I do not always share their opinion on taxonomic status of the species and their nomenclature because quite often the data cited are based on rather old literature or archieve materials, so they are outdated, not taking into account changes in the nomenclature and systematics of algae which have occurred recently. Nevertheless, they are included in the table with corrections, where it is possible, in conformity with modern taxonomic data. The original information of the other authors in the case I do not accept their interpretation of the taxa or doubt in correctness of the species identification is not included in the present list.

As it was already mentioned marine macrophytic algae of the Commander Islands are studied more thoroughly as compared to the other areas of the Russian Pacific. By the present day the species list of algae of the Islands based on our collections includes 164 species but it is still being enlarged in the process of the material processing. Recently we found there 9 species new for whole Russian Pacific coastal area (Selivanova and Zhigadlova, in press) and described a new taxon of red algae: *Gloiocladia guiryi* (Selivanova) Selivanova (2009). It belongs to the order Rhodymeniales, family Faucheaceae and appears to be the first representative of the genus and the family Faucheaceae to be reported from the Far Eastern seas of Russia though members of the genera *Faucheia* Bory et Montagne in Montagne and *Gloiocladia* J. Agardh were well known from the other areas. They were recorded from the American Pacific coasts (e.g. O'Clair, Lindstrom, 2000; Gabrielson et al., 2000, 2006), from Japan (Yoshida, 1998), Korea (Oak et al., 2005) and China (Xia, Wang, 2000). But our species still remains the only representative of the family Faucheaceae in the Russian Pacific sector. Originally it was described as *Faucheia guiryi* Selivanova (2008c). However a taxonomic revision of the genus *Faucheia* based on morphologic and molecular-genetic data resulted in the transfer of the overwhelming majority of the species of the genus including the generitype *F. repens* (C. Agardh) Montagne et Bory de Saint-Vincent to the genus *Gloiocladia* (Rodríguez-Prieto et al., 2007). Consequently it caused the necessity of the same transfer of *F. guiryi* to *Gloiocladia*. So a new nomenclatural combination for this species was proposed by me: *Gloiocladia guiryi* (Selivanova) Selivanova (2009).

Earlier one more taxon was described from western Bering Sea (Olytorskii Gulf, Lavrov Bay) and Glubokaya Bay (Selivanova & Zhigadlova, 2003). It represented the genus *Phycodryis* (Delesseriaceae, Rhodophyta) and was named *P. valentinae* Selivanova et Zhigadlova in honour of the well-known Russian phycologist Valentina F. Przhemenetskaya (Makienko) who has been studying marine algae of the Far Eastern seas of Russia for about 30 years. Later this species was also found by us in Avacha Gulf (Starichkov Island, south-eastern Kamchatka) (Selivanova and Zhigadlova, 2009).

The list of marine macroalgae of the Russian continental part of Bering Sea (without Commander Islands) totals to 153 species. Some of them were recorded by as as new for the Far Eastern seas of Russia (e.g. *Palmaria mollis* (Setchell et Gardner) Van der Meer et Bird, *Opuntella californica* (Farlow) Kylin, *Membranoptera setchellii* Gardner (Selivanova, 2002).

The general list of algae of the western Bering Sea is presented below in the table format (Tables 1-3) for convenience of information retrieval and with the purposes of shortening printed space. Separate tables contain data on algae of 3 major groups of the colour spectrum corresponding to higher taxa: green (Division Chlorophyta), red (Division Rhodophyta) and brown (Division Ochrophyta, class Phaeophyceae).

It should be noted that taxonomic viewpoints of Russian and western phycologists may differ, and that is the case with interpretation of brown algae. According to the first viewpoint this taxon is treated as a class Phaeophyceae within the phylum Heterokontophyta, kingdom Chromista, empire Eukaryota (Guiry and Guiry, 2011). But we prefer to follow Russian concept of Belyakova et al. (2006) and to attribute the class Phaeophyceae to the division Ochrophyta, kingdom Straminopila, empire Chromalveolata. Higher taxa (orders within divisions) are arranged in the tables according to their systematic position, while families within the orders, genera within the families and species within the genera are given in alphabetic order. The list also contains information on the distribution of algae. Abbreviations in the column "Distribution within the studied area" correspond to the names of the areas shown on schematic map (Fig. 1). In addition to the abbreviations adopted in the legend to the figure Commander Islands are abbreviated in the tables as CI.

3.1 Species composition and distribution of brown algae in the studied area

On the total there are 44 species of brown algae found up to now on the shelf of the western part of Bering Sea and Commander Islands. The species list is given below in the table format (Table 1).

Record number	Taxon	Distribution within the studied area
Order Desmarestiales		
Family Desmarestiaceae		
1	<i>Desmarestia aculeata</i> (Linnaeus) Lamouroux	CI, K, O
2	<i>Desmarestia ligulata</i> (Lightfoot) Lamouroux	CI
3	<i>Desmarestia viridis</i> (O.F. Müller) Lamouroux	CI
Order Ectocarpales		
Family Pylaiellaceae		
4	<i>Pylaiella littoralis</i> (Linnaeus) Kjellman	CI, K,O, NR
Family Chordariaceae		
5	<i>Chordaria chordaeformis</i> (Kjellman) Kawai et Kim	O, NR
6	<i>Chordaria flagelliformis</i> (O.F. Müller) C. Agardh	in all places
7	<i>Coilodesme bulligera</i> Strömfelt	CI
8	<i>Coilodesme californica</i> (Ruprecht) Kjellman	CI
9	<i>Delamarea attenuata</i> (Kjellm.) Rosenv	CI
10	<i>Dictyosiphon foeniculaceus</i> (Hudson) Greville	CI, K, O, NR
11	<i>Elachista tenuis</i> Yamada	CI
12	<i>Halothrix lumbricalis</i> (Kützinger) Reinke	K, O
13	<i>Leathesia marina</i> (Lyngbye) Decaisne	CI
14	<i>Leptonematella fasciculata</i> (Reinke) Silva	CI, O
15	<i>Melanosiphon intestinalis</i> (Saund.) Wynne	K, O
16	<i>Punctaria plantaginea</i> (Roth) Greville	K, O
17	<i>Saundersella simplex</i> (Saunders) Kylin	OZ, K, O
18	<i>Stictyosiphon tortilis</i> (Ruprecht) Reinke	O
19	<i>Soranthera ulvoidea</i> Postels et Ruprecht	CI
20	<i>Streblonema scabiosum</i> Setchell et Gardner	CI
Family Ectocarpaceae		
21	<i>Ectocarpus siliculosus</i> (Dillwyn) Lyngbye	CI, K, O
Order Scytosiphonales		
Family Scytosiphonaceae		
22	<i>Analipus filiformis</i> (Ruprecht) Papenfuss	K, O
23	<i>Analipus japonicus</i> (Harvey) Wynne	CI, K

24	<i>Petalonia fascia</i> (O.F. Müller) Kuntze	CI, K, O, NR
25	<i>Scytosiphon dotyi</i> Wynne	CI, O
26	<i>Scytosiphon lomentaria</i> (Lyngb.) Link	CI, K, O
Order Laminariales		
Family Alariaceae		
27	<i>Alaria angusta</i> Kjellman <i>emend.</i> Petrov	CI, OZ, K, NR
28	<i>Alaria marginata</i> Postels et Ruprecht	CI, K, O
29	<i>Eualaria fistulosa</i> (Postels et Ruprecht) Wynne	CI, K
Family Arthrothamnaceae		
30	<i>Arthrothamnus bifidus</i> (Gmelin) Ruprecht	OZ, K
Family Chordaceae		
31	<i>Chorda asiatica</i> Sasaki et Kawai	K, O
Family Costariaceae		
32	<i>Agarum clathratum</i> Dumortier	in all places
33	<i>Thalassiophyllum clathrus</i> (Gmelin) Postels et Ruprecht	CI, K
Family Laminariaceae		
34	<i>Cymathaere triplicata</i> (Postels et Ruprecht) J.Agardh	CI
35	<i>Laminaria longipes</i> Bory	CI, OZ, K
36	<i>Laminaria yezoensis</i> Miyabe	CI, OZ, K, O
37	<i>Saccharina bongardiana</i> (Postels et Ruprecht) Selivanova, Zhigadlova et G.I. Hansen	In all places
38	<i>Saccharina dentigera</i> (Kjellman) Lane, Mayes, Druehl et Saunders	CI, K
39	<i>Saccharina gurjanovae</i> (A. Zinova) Selivanova, Zhigadlova et G.I. Hansen	OZ, K
Order Fucales		
Family Fucaceae		
40	<i>Fucus evanescens</i> C. Agardh	In all places
Order Ralfsiales		
Family Ralfsiaceae		
41	<i>Ralfsia fungiformis</i> (Gunnerus) Setchell et Gardner	CI, K, O
42	<i>Ralfsia verrucosa</i> (Areschoug) Areschoug	K
Order Sphacelariales		
Family Sphacelariaceae		
43	<i>Battersia arctica</i> (Harvey) Draisma, Prud’homme et Kawai	K, O
44	<i>Chaetopteris plumosa</i> (Lyngbye) Kützing	K

Table 1. Empire Chromalveolata Kingdom Straminopila Division Ochrophyta Class Phaeophyceae

3.2 Species composition and distribution of green algae in the studied area

Up now there are 36 species of green algae found on the shelf of the western part of Bering Sea and Commander Islands. The species list is given below in the table format (Table 2).

Record number	Taxon	Distribution within the studied area
Class Bryopsidophyceae		
Order Bryopsidales		
Family Derbesiaceae		
1	<i>Derbesia marina</i> (Lyngbye) Solier	CI
Family Codiaceae		
2	<i>Codium ritteri</i> Setchell et Gardner	CI
Class Chlorophyceae		
Order Chlorococcales		
Family Chlorochytriaceae		
3	<i>Chlorochytrium inclusum</i> Kjellman	In all places
Family Endosphaeraceae		
4	<i>Codiolum gregarium</i> Braun	CI
Class Ulvophyceae		
Order Cladophorales		
Family Cladophoraceae		
5	<i>Chaetomorpha ligustica</i> (Kützing) Kützing	CI, K, O
6	<i>Chaetomorpha linum</i> (O.F. Müller) Kützing	CI
7	<i>Chaetomorpha melagonium</i> (Weber et Mohr) Kützing	OZ, K
8	<i>Cladophora speciosa</i> Sakai	CI, K
9	<i>Rhizoclonium riparium</i> (Roth) Harvey	CI, K, O
Order Ulotrichales		
Family Gomontiaceae		
10	<i>Monostroma crassidermum</i> Tokida	K, O
11	<i>Monostroma grevillei</i> (Thuret) Wittrock	CI, K, NR
Family Ulotrichaceae		
12	<i>Acrosiphonia duriuscula</i> (Ruprecht) Yendo	CI, K, O, NR
13	<i>Psedothrix groenlandica</i> (J.Agardh) Hanic et Lindstrom	CI, O
14	<i>Spongomorpha arcta</i> (Dillwyn) Kützing	CI
15	<i>Spongomorpha mertensii</i> (Yendo) Setchell et Gardner	CI
16	<i>Ulothrix flacca</i> (Dillwyn) Thuret in Le Jolis	CI, K, O
17	<i>Ulothrix implexa</i> (Kützing) Kützing	O
18	<i>Urospora penicilliformis</i> (Roth) Areschoug	CI, K, O
19	<i>Urospora wormskjoldii</i> (Mertens ex Horneman) Rosenvinge	CI, O
Order Ulvales		
Family Kornmanniaceae		
20	<i>Blidingia chadefaudii</i> (Feldman) Bliding	K
21	<i>Blidingia minima</i> (Nägeli ex Kützing) Kylin	CI, K, O
22	<i>Blidingia subsalsa</i> (Kjellman) Kornman et Sahling	CI, O

23	<i>Kornmannia leptoderma</i> (Kjellman) Bliding	CI
Family Ulvaceae		
24	<i>Percursaria percursa</i> (C. Agardh) Rosenvinge	CI, O
25	<i>Ulva clathrata</i> (Roth) C. Agardh	K, O
26	<i>Ulva fenestrata</i> Postels et Ruprecht	In all places
27	<i>Ulva flexuosa</i> Wulfen	CI, K
28	<i>Ulva linza</i> Linnaeus	CI, K, O
29	<i>Ulva procera</i> (Ahlner) Hayden, Blomster, Maggs, Silva, Stanhope et Waaland	CI, O
30	<i>Ulva prolifera</i> O.F. Müller	CI, K, O
31	<i>Ulvaria splendens</i> Ruprecht	In all places
Family Ulvellaceae		
32	<i>Acrochaete flustrae</i> (Reinke) O’Kelly	CI, O
33	<i>Acrochaete geniculata</i> (Gardner) O’Kelly	CI
34	<i>Acrochaete ramosa</i> (Gardner) O’Kelly	K
35	<i>Acrochaete viridis</i> (Reinke) R. Nielsen	CI, K
Class Trebouxiophyceae		
Order Prasiolales		
Family Prasiolaceae		
36	<i>Prasiola borealis</i> Reed	CI

Table 2. Empire Eukaryota kingdom Plantae subkingdom Viridiplantae division Chlorophyta

3.3 Species composition and distribution of red algae in the studied area

On the total there are 129 species of red algae found up to now on the shelf of the western part of Bering Sea and Commander Islands. The species list is given below in the table format (Table 3).

Record number	Taxon	Distribution within the studied area
Class Compsopogonophyceae		
Order Erythropeltidales		
Family Erythrotrichiaceae		
1	<i>Erythrocladia irregularis</i> Rosenvinge	CI, K
Class Bangiophyceae		
Order Bangiales		
Family Bangiaceae		
2	<i>Bangia fuscopurpurea</i> (Dillwyn) Lyngbye	CI
3	<i>Porphyra abbottiae</i> Krishnamurthy	CI
4	<i>Porphyra brumalis</i> Mumford	CI, O, NR
5	<i>Porphyra gardneri</i> (Smith et Hollenberg) Hawkes	CI
6	<i>Porphyra kurogii</i> Lindstrom in Lindstrom et Cole	CI, K, O, NR
7	<i>Porphyra miniata</i> (C. Agardh) C. Agardh	CI, K, O, NR
8	<i>Porphyra ochotensis</i> Nagai	CI

9	<i>Porphyra pseudolinearis</i> Ueda	CI, K, O
10	<i>Porphyra purpurea</i> (Roth) C. Agardh	CI, O, NR
11	<i>Porphyra tasa</i> (Yendo) Ueda	O
12	<i>Porphyra torta</i> Krishnamurthy	CI, NR
13	<i>Porphyra variegata</i> (Kjellman) Kjellman in Hus	CI, K, O, NR
Class Florideophyceae		
Order Hildenbrandiales		
Family Hildenbrandiaceae		
14	<i>Hildenbrandia rubra</i> (Sommerfelt) Meneghini	CI, K, O
Order Corallinales		
Family Corallinaceae		
15	<i>Bossiella frondescens</i> (Postels et Ruprecht) Dawson	CI
16	<i>Corallina pilulifera</i> Postels et Ruprecht	CI, K,O
17	<i>Pachyarthron cretaceum</i> (Postels et Ruprecht) Manza	CI, K
Family Hapalidiaceae		
18	<i>Clathromorphum circumscriptum</i> (Strömfelt) Foslie	CI, K,O, NR
19	<i>Clathromorphum compactum</i> (Kjellman) Foslie	CI, K, O
20	<i>Clathromorphum loculosum</i> (Kjellman) Foslie	CI, K, O
21	<i>Clathromorphum nereostratum</i> Lebednik	CI, O
22	<i>Clathromorphum reclinatum</i> (Foslie) Adey	CI
23	<i>Phymatolithon lenormandii</i> (Areschoug) Adey	CI, OZ, K
24	<i>Lithothamnion phymatodeum</i> Foslie	CI
25	<i>Lithothamnion sonderi</i> Hauck	CI, OZ, K
Colaconematales		
Colaconemataceae		
26	<i>Colaconema savianum</i> (Meneghini) R. Nielsen	O
Order Acrochaetiales		
Family Acrochaetiaceae		
27	<i>Acrochaetium alariae</i> (Jónsson) Bornet	O
28	<i>Acrochaetium arcuatum</i> (Drew) Tseng	OZ, K
29	<i>Acrochaetium microscopicum</i> (Nägeli ex Kützing) Nägeli	K
30	<i>Acrochaetium parvulum</i> (Kylin) Hoyt	K
31	<i>Rhodochorton purpureum</i> (Lightfoot) Rosenvinge	CI, K
Order Palmariales		
Family Meiodiscaceae		
32	<i>Meiodiscus spetsbergensis</i> (Kjellman) Saunders et McLachlan	O
33	<i>Rubrointrusa membranacea</i> (Magnus) Clayden et Saunders	CI
Family Palmariaceae		
34	<i>Devaleraea compressa</i> (Ruprecht) Selivanova et Kloczcova	K, O, NR
35	<i>Devaleraea microspora</i> (Ruprecht) Selivanova et Kloczcova	K, NR

36	<i>Halosaccion firmum</i> (Postels et Ruprecht) Kützing	CI, K, O
37	<i>Halosaccion glandiforme</i> (Gmelin) Ruprecht	CI, K, O, NR
38	<i>Halosaccion minjaiti</i> I.K. Lee	CI
39	<i>Palmaria callophyllloides</i> Hawkes et Scagel	CI, O
40	<i>Palmaria hecatensis</i> Hawkes	CI, O
41	<i>Palmaria marginicrassa</i> I.K. Lee	CI
42	<i>Palmaria mollis</i> (Setchell et Gardner) van der Meer et Bird	CI, O, K
43	<i>Palmaria stenogona</i> (Perestenko) Perestenko	CI, O, K
Order Ahnfeltiales		
Family Ahnfeltiaceae		
44	<i>Ahnfeltia fastigiata</i> (Endlicher) Makienko	CI
45	<i>Ahnfeltia plicata</i> (Hudson) Fries	CI, K
Order Bonnemaisoniales		
Family Bonnemaisoniaceae		
46	<i>Pleuroblepharidella japonica</i> (Okamura) Wynne	CI
Order Cryptonemiales		
Family Crossocarpaceae		
47	<i>Beringia castanea</i> Perestenko	CI
48	<i>Cirrulicarpus gmelinii</i> (Grunow) Tokida et Masaki	CI
49	<i>Cirrulicarpus ruprechtianum</i> (Sinova) Perestenko	CI, K, O, NR
50	<i>Crossocarpus lamuticus</i> Ruprecht	CI, OZ, K
51	<i>Hommersandia palmatifolia</i> (Tokida) Perestenko	CI, K, O
52	<i>Kallymeniopsis lacera</i> (Ruprecht) Perestenko	In all places
53	<i>Velatocarpus kurilensis</i> Perestenko	K, O
54	<i>Velatocarpus pustulosus</i> (Postels et Ruprecht) Perestenko	CI, OZ, K, O
Family Dumontiaceae		
55	<i>Constantinea rosa-marina</i> (Gmelin) Postels et Ruprecht	CI
56	<i>Constantinea subulifera</i> Setchell	CI
57	<i>Dilsea socialis</i> (Postels et Ruprecht.) Perestenko	OZ, K, O
58	<i>Dumontia contorta</i> (Gmelin) Ruprecht	CI, K, O
59	<i>Neodilsea natashae</i> Lindstrom	CI, K
60	<i>Neodilsea yendoana</i> Tokida	CI, K, O
Family Kallymeniaceae		
61	<i>Callophyllis radula</i> Perestenko	CI
62	<i>Callophyllis rhynchocarpa</i> Ruprecht	CI, K, O
63	<i>Euthora cristata</i> (C. Agardh) J. Agardh	In all places
Order Halymeniales		
Family Halymeniaceae		
64	<i>Neoabbottiella araneosa</i> (Perestenko) Lindstrom	CI, OZ, K
Order Gigartinales		
Family Cystocloniaceae		
65	<i>Fimbrifolium dichotomum</i> (Lepechin) Hansen	In all places

66	<i>Rhodophyllis capillaris</i> Tokida	CI, K
Family Endocladiaceae		
67	<i>Gloiopeltis furcata</i> (Postels et Ruprecht) J. Agardh	CI, K, O
Family Furcellariaceae		
68	<i>Opuntiella californica</i> (Farlow) Kylin	OZ, K
69	<i>Opuntiella ornata</i> (Postels et Ruprecht) A. Zinova	CI, OZ, K, O
70	<i>Turnerella mertensiana</i> (Postels et Ruprecht) Schmitz	In all places
71	<i>Turnerella pennyi</i> (Harvey) Schmitz in Rosenvinge	K
Family Gigartinaceae		
72	<i>Mazzaella parksii</i> (Setchell et Gardner) Hughey, Silva et Hommersand	CI, K
73	<i>Mazzaella phyllocarpa</i> (Postels et Ruprecht) Perestenko	CI, K, O
Family Phyllophoraceae		
74	<i>Coccotylus truncatus</i> (Pallas) Wynne et Heine	K, O
75	<i>Lukinia dissecta</i> Perestenko	CI
76	<i>Mastocarpus pacificus</i> (Kjellman) Perestenko	CI, K, O, NR
77	<i>Mastocarpus papilatus</i> (C. Agardh) Kützing	CI
Order Nemastomatales		
Family Schizymeniaceae		
78	<i>Schizymenia pacifica</i> (Kylin) Kylin	K, NR
Order Rhodymeniales		
Family Faucheaceae		
79	<i>Gloiocladia guiryi</i> (Selivanova) Selivanova	CI
Family Rhodymeniaceae		
80	<i>Sparlingia pertusa</i> (Postels et Ruprecht) Saunders, Strachan et Kraft	CI, K
Order Ceramiales		
Family Ceramiaceae		
81	<i>Ceramium cimbricum</i> H. Petersen in Rosenvinge	K
82	<i>Ceramium deslongchampsii</i> Chauvin ex Duby	K
83	<i>Ceramium kondoi</i> Yendo	CI, K, O
84	<i>Microcladia borealis</i> Ruprecht	CI
85	<i>Scagelia breviarticulata</i> Perestenko	CI
86	<i>Scagelia pylaisaei</i> (Montagne) Wynne	CI, K, O
Family Delesseriaceae		
87	<i>Congregatocarpus kurilensis</i> (Ruprecht) Wynne	CI
88	<i>Hideophyllum yezoense</i> (Yamada et Tokida) A. Zinova	CI, OZ, K, O
89	<i>Hymenena ruthenica</i> (Postels et Ruprecht) A. Zinova	CI, OZ, K, O
90	<i>Laingia aleutica</i> Wynne	CI
91	<i>Membranoptera spinulosa</i> (Ruprecht) Kuntze	In all places
92	<i>Membranoptera serrata</i> (Postels et Ruprecht) A. Zinova	K, NR
93	<i>Membranoptera setchellii</i> Gardner	OZ, K
94	<i>Mikamiella ruprechtiana</i> (A. Zinova) Wynne	CI

95	<i>Nienburgia prolifera</i> Wynne	CI
96	<i>Pantoneura fabriciana</i> (Lyngbye) Wynne	K
97	<i>Pantoneura juergensii</i> (J. Agardh) Kylin	CI, OZ, K, NR
98	<i>Phycodrys fimbriata</i> (Kuntze) Kylin	CI, OZ, O, NR
99	<i>Phycodrys serratiloba</i> (Ruprecht) A. Zinova	CI, OZ, K, O
100	<i>Phycodrys valentinae</i> Selivanova et Zhigadlova	O, NR
101	<i>Phycodrys vinogradovae</i> Perestenko et Gussarova in Perestenko	CI
102	<i>Tokidadendron bullata</i> (Gardner) Wynne	CI, O
103	<i>Yendonia crassifolia</i> (Ruprecht) Kylin	CI, OZ, K, O
Family Rhodomelaceae		
104	<i>Beringiella labiosa</i> Wynne	CI
105	<i>Harveyella mirabilis</i> (Reinsch) Schmitz et Reinke	K, O
106	<i>Neorhodomela larix</i> (Turner) Masuda	CI, K, O
107	<i>Neorhodomela oregona</i> (Doty) Masuda	CI
118	<i>Neosiphonia japonica</i> (Harvey) M.S.Kim et I.K.Lee	K
109	<i>Odonthalia annae</i> Perestenko	CI
110	<i>Odonthalia corymbifera</i> (Gmelin) Greville	CI
111	<i>Odonthalia dentata</i> (Linnaeus) Lyngbye	CI, K, O, NR
112	<i>Odonthalia floccosa</i> (Esper) Falkenberg	CI
113	<i>Odonthalia kamtschatica</i> (Ruprecht) J. Agardh	CI, OZ, K, NR
114	<i>Odonthalia ochotensis</i> (Ruprecht) J. Agardh	CI, K, O
115	<i>Odonthalia setacea</i> (Ruprecht) Perestenko	CI, K, O
116	<i>Polysiphonia morrowii</i> Harvey	CI
117	<i>Polysiphonia stricta</i> (Dillwyn) Greville	CI, K, O
118	<i>Pterosiphonia bipinnata</i> (Postels et Ruprecht) Falkenberg	In all places
119	<i>Pterosiphonia hamata</i> Sinova	CI
120	<i>Rhodomela pinnata</i> Perestenko	K, O
121	<i>Rhodomela sibirica</i> A. Zinova et Vinogradova in Vinogradova	K, O
122	<i>Rhodomela tenuissima</i> (Ruprecht) Kjellman	K, O
123	<i>Tayloriella abyssalis</i> Wynne	CI
Family Wrangeliaceae		
124	<i>Neoptilota asplenioides</i> (Esper) Kylin	CI, NR
125	<i>Pleonosporium kobayashii</i> Okamura	CI
126	<i>Pleonosporium vancouverianum</i> (J. Agardh) Setchell et Gardner	CI
127	<i>Ptilota filicina</i> J. Agardh	CI, K, O, NR
128	<i>Ptilota serrata</i> Kützing	In all places
129	<i>Tokidaea serrata</i> (Wynne) Lindstrom et Wynne	CI

Table 3. Empire Eukaryota Kingdom Plantae subKingdom Biliphyta Division Rhodophyta

In conclusion of this part I would like to point out that the studies on the species composition of any group of organisms are the basis for further scientific studies in

biocenology, biogeography, ecology etc. As far as marine macrophytes are among the leading components of benthic communities of the shelf of Bering Sea their inventory is very important. In spite of the incompleteness of inventory works in this region main counters of the marine algal flora are already outlined and might serve as background of development of different scientific conceptions.

For instance knowledge on contemporary algal flora may be helpful for clarification of the problems of historical phytogeography, processes of the species formation and migration. It is also important for understanding of the regularities of forming and functioning of the ecosystems and the role of seaweeds in benthic communities. Shallow water zones of the shelf of the western part of Bering Sea and Commander Islands are also of interest for economic activity. Competent exploitation of their resources is possible only on the basis of reliable scientific data on the species composition, structure and distribution of benthic communities. These problems are discussed to some extent in the sections below.

4. Brief characteristics of the algal flora of the western part of Bering Sea and Commander Islands

In Russian phycological literature the whole water area of the Far Eastern Seas is considered to be divided into 7 large floristic regions: 1) the Sea of Japan, 2) small Kurile Islands region; 3) southern Kurile Islands region; 4) northern Kurile Islands region; 5) the Sea of Okhotsk; 6) Bering Sea; 7) Commander Islands (Perestenko, 1994). We carried out our studies in the last 2 regions and came to a conclusion that the Bering Sea region that according to Perestenko (1994) includes the whole Eastern Kamchatka coasts should be subdivided into 2 subregions practically equal in geographic extension and floristic significance: a) proper Bering Sea area in its geographic borders (from Bering Strait to Ozernoi Gulf) and b) southeastern Kamchatka (from Kamchatskii Gulf to Lopatka Cape). The latter is out of the limits of the present study. Here we discuss marine algal flora of 2 major floristic complexes of the regions located in the upper part of North Pacific, those of Commander Islands and western continental coast of Bering Sea.

From the point of view of phytogeography both floristic complexes judging from the elements forming them may be attributed to boreal (cold-temperate) type (Table 4). The number of arctic-boreal and high-boreal species in the floristic complex of the Commander Islands totals to 41 %, in the floristic complex of western continental part of Bering Sea their number comes to 44 %. If so called wide-boreal elements are taken into account these figures increase correspondingly to 92% and 90 %.

Systematic analysis of the species list revealed that the largest orders in the floristic complex of the Commander Islands are: 1) within the division Ochrophyta, class Phaeophyceae: Ectocarpales (contains 3 families, 11 genera and 12 species) and Laminariales (5 families, 7 genera and 10 species); 2) within the division Rhodophyta: Ceramiales (4 families, 24 genera and 38 species) and Cryptonemiales (in its traditional interpretation) (5 families, 13 genera and 17 species); 3) within the division Chlorophyta: Ulvales (3 families, 5 genera and 14 species). These orders include above 55 % of the total species number of the flora.

The situation on the western continental coast of Bering Sea is much the same, the largest orders in this floristic complex are: 1) within the division Ochrophyta, class Phaeophyceae: Ectocarpales (contains 3 families, 10 genera and 11 species) and Laminariales (5 families, 8 genera and 12 species); 2) within the division Rhodophyta: Ceramiales (4 families, 18 genera and 31 species) and Cryptonemiales (in its traditional interpretation) (5 families, 12 genera

Phytogeographic group	Commander Islands		Continental part of Bering Sea	
	Species number	%	Species number	%
Arctic-boreal	17	10	24	16
High boreal	50	31	43	28
Wide boreal	83	51	70	46
Boreal-tropical	8	4	10	6
Cosmopolitan	6	4	6	4
Total	164		153	

Table 4. Phytogeographic composition of the macrophytic algae floras of Bering Sea and Commander Islands

and 14 species); and 3) within the division Chlorophyta: Ulvales (3 families, 4 genera and 14 species). These orders include about 54 % of the total species number of the flora.

The largest higher taxon of the Commander Islands' flora is the division Rhodophyta consisting of 13 orders, 25 families, 63 genera and 103 species and exceeding the number of green and brown algae in sum (Phaeophyceae + Chlorophyta) in 1.7 times. Thus, the nucleus of the Islands' flora is represented by red algae. The largest genera within Rhodophyta, consisting of 5 and more species are as follows: *Porphyra* (10 species), *Odonthalia* (7 species), *Palmaria* (5 species) and *Clathromorphum* (5 species). The largest genus among Chlorophyta is *Ulva* (6 species), however Phaeophyceae has no genera with more than 3 species, though it contains the complex of genera *Laminaria*+ *Saccharina*, but even in this case the number of the species totals to only 4.

The division Rhodophyta is also the largest one in the flora of the western continental coasts of Bering Sea consisting of 14 orders, 25 families, 56 genera and 91 species and exceeding the total number of green and brown algae in 1.5 times. So the nucleus of the Bering Sea flora is also formed by red algae though their number is less than that of Commander Islands (91 species versus 103 accordingly). The largest genus within Rhodophyta in the Bering Sea flora is *Porphyra* (8 species), other large genera containing 4 species each are: *Clathromorphum*, *Acrochaetium*, *Palmaria* and *Odonthalia*. The largest taxa among division Chlorophyta and class Phaeophyceae are correspondingly the genus *Ulva* (7 species) and the complex of genera *Laminaria*+ *Saccharina* (5 species).

The figures given above make it possible to outline main characteristics of the floras. Particularly, in accordance with data of systematic analysis in general marine algal flora of both regions can be characterized as variegated in the species composition and allochthonous by origin. It contains big number of monotypic genera and families. I mean those taxa of higher rank that are represented in the studied area by only one taxon of lower rank, i.e. a family represented by one genus and a genus represented by one species. On Commanders there are: 17 genera and 5 families in Phaeophyceae; 12 genera and 7 families in Chlorophyta; 42 genera and 9 families in Rhodophyta; in Bering Sea: 18 genera and 7 families in Phaeophyceae; 7 genera and 5 families in Chlorophyta; 35 genera and 11 families in Rhodophyta. Allochthonous type of the flora is also confirmed by relatively low ratios: species/genus (1.6 for both regions); species/family (3.6 for Commander Islands and 3.4 for Bering Sea) and genus/family (2.3 for Commander Islands and 2.1 for Bering Sea). No endemic species were found. Variegated species composition shows that both floristic complexes are under strong influence of adjacent areas. Main ecologic factors of the existence of contemporary floras of algae-macrophytes in the area are formed by ocean currents.

The Bering Sea marine algal flora presumably (on the assumption of paleontologically verified data on the other groups of organisms) was formed mostly by low-boreal elements that adapted to fall of temperature in Cainozoic period. At present this flora is being enriched due to the penetration of high-boreal and arctic species (e.g. *Dilsea socialis*, *Rhodomela sibirica*) via cold-water Eastern Kamchatka current. This current goes further to the south and in such a way cold-water species are spread along the south-eastern Kamchatka coasts (up to Lopatka Cape) (Fig. 1). Thus the marine algal flora of Bering Sea has a well-defined migratory character and is under strong influence of the Arctic area.

On the other hand Commander Islands are greatly influenced by American continent due to the branch of Alaska current providing penetration of American elements of flora. I suppose this process to be main tendency of the Islands' flora development. As a result of invasion of American elements marine algal flora of the Commander Islands is more rich and diverse as compared to other areas of Bering Sea. For example, elementary analysis of the data from our Tables 1-3 show that there are 57 species of marine macrophytes (9 species of green, 10 – of brown and 38 - of red algae) that are met only on Commander Islands and are not found on the continental coast of Bering Sea. Many of these species were described from the American Pacific coasts (e.g. *Streblonema scabiosum*, *Acrochaete geniculata*, *Erythrocladia irregularis*, *Palmaria callophyloides*, *P. hecatensis*, *Laingia aleutica*, *Nienburgia prolifera*, *Beringiella labiosa*, *Tayloriella abyssalis*). Of course, absence of records of some species from the continental part of Bering Sea does not in all cases mean that these algae do not grow there, possibly it is due to less careful studies in this area in contrast to Commander Islands.

It should be noted that in spite of the fact that all above-mentioned species were described from the American continent they are not necessarily American in their origin. It is quite possible that marine algal flora of the American continent is just more comprehensively studied and new species are found and described there more often in comparison with the Russian Pacific sector. And it is no wonder, because big and well-equipped groups of phycologists from Canada and USA effectively work in their sector of the Pacific Ocean while in Russia in fact only small isolated groups of enthusiasts try to study marine algae of the western Pacific coasts. In any case the presence of the species on Commander Islands that are not met in other areas of the Russian Pacific but are common with American coast (*Coilodesme californica*, *Pleonosporium vancouverianum*, *Microcladia borealis*, *Tokidaea serrata*, *Nienburgia prolifera*, *Laingia aleutica*, *Beringiella labiosa*, *Odonthalia floccosa* etc.) brings together their flora with the floras of Aleutian Islands and Alaska. So Commander Islands serve as a peculiar bridge uniting American and Asian algal floras.

Comparison of the species composition of the macrophytes of the Commander Islands' shelf and that of the south-eastern Kamchatka reveals certain similarity of the floras of these two regions (Jaccard's coefficient of community $K_j = 0.68$) and testify to belonging of the Commander Islands flora to Asian type.¹ According to my calculations similarity between

¹ The geological data are also in favour of hypothesis of Asian origin of marine algal flora of the Commander Islands. In accordance with the theory of continental drift the Pacific islands (together with their underwater foundations) are considered to be the edge chains separated from the continental blocks. During general movement of the Earth crust along the mantle oriented mostly to the west they lagged behind and remained in the east. So their original position was closer to the Asian continent in comparison with the present day (Vegener, 1984). In some sense hypothesis of Asian origin of the Commander Islands marine algal flora is confirmed by data on terrestrial flora of some well studied

the floras of south-eastern Kamchatka and western part of Bering Sea is a little less ($K_j = 0.63$), and it is the least between the floras of the Commander Islands and western continental coast of Bering Sea ($K_j = 0.51$). I explain phenomenon of such low similarity (in other words, considerable difference) between floristic complexes of geographically close water areas by peculiarities of the system of ocean currents in this sector of the North Pacific, existence of already mentioned Eastern Kamchatka and Alaska currents (Fig. 1) providing favourable conditions for penetration of invasive species of different origin from the adjacent areas, namely from the Arctic into northern part of Bering Sea and from the American continent to Commander Islands. To tell the truth, such deviations in comparative floristic coefficients may happen due to purely statistical reasons (Kafanov et al., 2004). Nevertheless my conclusions are in agreement with the data of K.L. Vinogradova & L.P. Perestenko (1978) who also showed considerable difference of marine flora of the western Bering Sea from those of the south-eastern Kamchatka and Commander Islands.

5. The problem of diversity conservation of algae in the western Bering Sea including Commander Islands

Conservation of biodiversity is a primary condition for stability of biosphere. Studies on biodiversity became a component part of National strategy of Russia more than 10 years ago. Creation of the system of the specially protected nature areas (SPNA-s) serves as the most important instrument of biodiversity conservation in our country. At present 9 SPNA-s with marine coastal area aimed at conservation of hydrobionts diversity have been created in the studied sector of the North Pacific. The northernmost of these areas – Nature Park “Beringia” is located within administrative borders of Chukchi Autonomous District and is just mentioned but not included in our discussion.

5.1 Commander State Biosphere Reserve

Commander State Biosphere Reserve is the most important among the rest 8 SPNA-s. It embraces practically the whole archipelago Commander Islands (Fig. 1). Necessity of protection of the unique biodiversity of Commander Islands was acknowledged in Russia many years ago.

In 1958 the Soviet government prohibited fishery and other economic activity in 30-miles sea area around the Islands with the purpose of conservation of marine mammals (sea otters, fur seals). In 1993 in accordance with Decision of the government of the Russian Federation the state biosphere “Komandorskii” (Commander) reserve with the area of 3648679 hectares was established, its coastal area covering 3463300 hectares. This is one of the largest reserves in the world and taking into account the size of its coastal zone it could be considered as a marine reserve. Main objectives of this state biosphere reserve are: studies and protection of the unique natural complexes of the archipelago, genofond of plants and animals, protection of large rookeries of marine mammals, population of blue fox, nesting spots of rare birds and also conservation of historical traces of Vitus Bering’s expedition. Besides that Commander Islands are the place of compact residence

Pacific islands and archipelagos. For instance, aboriginal flora of Hawaiian Islands (without agricultural and invasive species) is more allied with the flora of the Old World than with that of their nearest neighbour – North America (Drude, 1890).

of indigenous small people of the North – Aleutians with their traditional mode of life and culture that also need protection. Based on these grounds the Commander Reserve was included in the worldwide net of the biosphere reservations in the frames of the UNESCO program “The man and the biosphere” (MAB).

Organization of the state reserve helps to take under protection all marine organisms inhabiting the Islands, including marine macrophytic algae. And this fact inspires optimism because the majority of rare species included in the Red Data Books of Kamchatka and Russia grow on Commander Islands. In most cases the Islands are the only area of their inhabitation within the Russian coasts (Selivanova, 2007). Conservation of these species in the nature has not only scientific importance but is an integral part of conservation of biodiversity of the coastal zone in its broad sense (i.e. not only of the species diversity but also diversity and stability of marine ecosystems).

It is expedient to note that all species included in the Red Data Book of Kamchatka are rare because of natural reasons, their number is not reduced by human activity and their conservation needs rather non-interference in natural habitats than any active measures of protection. However it is necessary to exclude anthropogenic pollution of the coastal area and poaching in the Reserve’s water area. Of course, rare algae themselves cannot be the objects of harvesting but irrational catch of other hydrobionts may cause nonreversible changes in marine ecosystems and as a result – extermination of rare species of algae. In fact there are groups of algae growing on Commander Islands that should be attributed to the category of so called vulnerable species and these are commercial laminarian species (*Laminaria*, *Saccharina*). At present they are abundant on the Islands and there are no real threats for decrease in their diversity. In my opinion remoteness of the Islands, small number of their human population and relatively low demand for the plant marine products will prevent local community from poach harvest of seaweeds as it happened in the vicinity of Petropavlovsk-Kamchatskii. However there is a risk of their overharvest in case of large-scale uncontrolled commercial catch. I think that this perspective should be treated as a possible threat for the biodiversity conservation on the shelf of the Commander Islands.

5.2 Reserve “Karaginskii Island”

Reserve “Karaginskii Island” is another important SPNA in the Bering Sea that covers Karaginskii Island located in Karaginskii Gulf and separated from Kamchatka Peninsula by Litke Strait (Fig. 1(2)). At organization of the Reserve “Karaginskii Island” in 1996 all types of commercial catch were prohibited in the water areas around the Island. It was supposed to organize here marine biosphere reserve in succeeding years but the program did not receive financial support. At present Reserve “Karaginskii Island” does not exist any more however the Island is included in the list of wetlands in accordance with Ramsar Convention. There still remains possibility of getting support for the program of organization of marine biosphere reserve on Karaginskii Island, we consider it very expedient because of high biodiversity of marine flora and fauna of the area that needs protection.

Besides the above-mentioned SPNA-s there are areas within Bering Sea with different nature protective status aimed at conservation of marine ecosystems:

5.3 State nature reserve of federal significance “Koryakskii”

State nature reserve of federal significance “Koryakskii” was founded in 1995 by the Decision of the government of the Russian Federation. The reserve covers a part of Goven

Peninsula and adjacent water area of Lavrov Bay (Olyutorskii Gulf) (Fig. 1 (3)). The protected water area totals to 83000 hectares. The purpose of creation of the “Koryakskii” Reserve was to provide protection of the whole ecosystem complex of the northern Kamchatka including marine and coastal ecosystems of Bering Sea with large colonies of sea birds.

5.4 Nature park “Beringia”

Nature park “Beringia”, founded in 1992, is located in Chukchi Autonomous District and includes a part of Bering Sea water area (Fig. 1 (4)).

In addition there are five **Nature Monuments of regional significance** that have protected marine area within their borders: «**Verkhoturov Island**», «**Witgenstein Cape**», «**Witgenstein Rock**», «**Yuzhnaya Glubokaya Bay**» and «**Anastasia Bay**». It is planned to create some new marine nature parks and state nature reserves of federal significance. However all already existing SPNA-s with the exception of Commander State Biosphere Reserve are just nominal organizations without real system of nature protection service. As far as the level of knowledge on their biodiversity is concerned it is extremely low and scientific studies seem to us to be the major task for future.

However it should not preclude the possibility of seaweed harvest on the territory of SPNA-s even with high nature protective rank. For instance, Commander Reserve is thought to be perspective for plant resources exploitation in the areas with permitted economic activity (Selivanova, 2008b). It is necessary to elaborate well-grounded recommendations on permissible level of harvest that in its turn should be based on reliable estimate of algal resources. Correct estimation of resources is important for proper decisions on biodiversity conservation not destroying ecosystem integrity and providing restoration of natural seaweed resources.

6. Preliminary data on seaweed resources of the Russian sector of North Pacific

The largest and most productive algal communities in the Russian sector of North Pacific (western continental part of Bering Sea and Commander Islands) are formed by seaweeds of the order Laminariales. These algae are main objects of commercial harvest. Unfortunately there are no exact present-day quantitative data on seaweed resources in the studied area. The only information available is just preliminary, fragmentary and rather old, dated to the first decades of the last century (Kongisser, 1933; Gail, 1936). Naturally we need more updated information concerning this matter but practical estimation works require considerable expeditionary expenses, equipment and efforts of many specialists. At present it is unreal to expect that these works will be carried out. So the only way out is to make calculations using so called expert method. It is a theoretical estimation based on remote sensing data on the bottom area suitable for the growth of seaweeds and their projective cover in communities. In accordance with this expert estimation general stock of seaweeds resources in the Russian sector of North Pacific comes to 10 million tons.

In spite of such rich stock there is a risk of overharvest in case of poor organization of catch. Exceeding press on commercial algae may cause abrupt decrease of their number and further replacement by competitive species, for example, by crustose coralline algae. This may cause irreversible changes in the structure of plant benthic communities and as a result total loss of commercial species. That is why scientific recommendations on permissible level of harvest are very important.

In my opinion the most perspective region for sustainable use of seaweed resources is a shallow water area of Commander Islands in spite of the fact that their major part is included in the state biosphere reserve. The reasons for such statement are as follows: 1) this region is one of the most pristine in the Far Eastern seas of Russia, seaweeds growing there are suitable for food and pharmaceutical use not only by the local population but also for export; 2) analysis of the benthic communities of the Islands reveal that their state is good enough and the number of laminarian algae is actually excessive; 3) it will be expedient and even necessary to organize harvesting of commercial algae in the zones of permitted economic activity; 4) in the latest few years the scheme of initial zonation of the Reserve was reconsidered and the limits of economic zones were appreciably extended.

However as it was already said, competent exploitation of seaweed resources should be based on reliable research data on the algal species composition, number and distribution within the area. This information is obviously insufficient and additional studies are necessary.

We have at our disposal only relatively old data on the stock and distribution of the laminarian algae on Commander Islands obtained during expeditions of Kamchatka Branch of the Pacific Institute of Geography in 1986, 1989-1991. In accordance with these data *Saccharina bongardiana* forms relatively small areas (20-30 hectares) covered by kelp at the depths of 1-2 meters. In sheltered bays *S. bongardiana* is met up to the depth of 12 m, forming mixed communities with other brown algae. Mean biomass of *S. bongardiana* varied in different years. The area covered by the kelp with the dominance of this species did not exceed 10 km², total stock came to a 1-1.5 thousand tons. In exposed habitats *S. bongardiana* was replaced by a competitive species *S. dentigera*.

S. dentigera formed a distinct belt around the coast of Bering Island at the depths from 3 to 10 m. Some patches are met from the depth of 0 (in exposed to waves places) to 25 m. Plants often settled on coralline alga *Clathromorphum nereostratum* and obtaining big size were cast ashore by storms together with coralline crusts on haptera. The portion of *S. dentigera* in cast ashore mass reached 70%. The biomass of *S. dentigera* varied in different years in 3-4 times. The area occupied by the kelp of this species came to 200 km², and the stock averaged to 2 million tons.

At present only algae of the genera *Saccharina* and *Laminaria* are the objects of harvest in Kamchatka Region though other laminarian algae (*Alaria*, *Eualaria*) are also suitable for food, forage and technological use. In particular a giant alga *Eualaria fistulosa* that forms dense population around Bering Island at the depths 5-15 m may be treated as a perspective commercial species. Sometimes it is met in deeper places (up to 25 m) and vice versa rises to 1-2 m in sheltered habitats. The density of populations of *E. fistulosa* varied up to 10 times from a year to year (from 0.6 to 7 kg/m²). Kelp formed by this alga occupied the area of about 100 km². Floating plants of *E. fistulosa* and their parts formed rather thick fields with projective cover up to 60%. In autumn-winter period these floating fields of *E. fistulosa* were destroyed by storms but in April-May they appeared again and reached their maximal size (10-12 m) by June-July due to the fast growth of attached algae. The stock of this alga on Bering Island was estimated at its minimal amount in 1992 to be 82 thousand tons and at its maximal amount in 1990-1991 – up to 700 thousand tons. The total amount of *E. fistulosa* at both Bering and Medny Island in 1990-es averaged to 925 thousand tons.

As is well known abundance of brown algae is subjected to unpredictable changes as a result of harvesting, so their natural fluctuations should be taken into account because harvesting in unfavourable for seaweeds years may lead to abrupt decrease in their stock.

Three species we discuss (*S. bongardiana*, *S. dentigera* and *E. fistulosa*) have different growth peculiarities, reproduction and demographic structure of the populations, so the strategy of their harvest should be different.

For instance, *S. bongardiana* may be harvested in the volumes of 50 % of its stock in each area (for Bering Island 0.5-0.75 thousand tons). The most expedient way of this species harvesting is cutting of algae in the intertidal zone during low tides using special knife (pruner) in 20 cm above the meristem zone at basal part of the blade. This provides fast re-growth of the blade and makes it possible to get aftercrop in 1.5-2 months, i.e. to carry out multiple harvesting during a vegetation period.

On the other hand the total harvest volume of *S. dentigera* should not exceed 30 % of the total stock because algae of this species grow much slower than *S. bongardiana* but have longer lifetime (presumably, 4-5 years of *S. dentigera* versus 2-3 years of *S. bongardiana*). Inasmuch as *S. dentigera* grows at considerable depths and constitutes a major part in cast ashore algae, collection of fresh samples of *S. dentigera* from the stormy beached seaweed mass seems to be the most rational method of its harvesting.

Harvesting of *E. fistulosa* is possible from the boat by wrapping floating parts of the plants round long hook (kanza). For the sake of fast restoration of the stock it is expedient to cut them at the depth less than 1 m from water surface. Owing to high growth rate of the fronds of *E. fistulosa* it is possible to harvest phytomass several times during one vegetation season. Plants are able to reproduce successfully because the lower parts with sporophylles are not damaged. Annual estimated yield of *E. fistulosa* for Bering Island comes to 120 thousand tons.

It should be accentuated that the use of dragrope is inadmissible in seaweed harvest as it causes considerable loss of raw material, damage of substratum, death of juvenile plants and as a result – elimination of thickets of algae. Moreover torn off plants sink and form dead bottom sediments, in the process of decay they reduce hydrogen contents in the near-bottom water layers.

Recommended by us methods of brown algae harvesting at the coasts of Commander Islands satisfy the international requirements of exploitation of seaweed resources and do not conflict with the nature protection regime in the region. They are ecologically justified because produce no negative effects on the benthic marine ecosystems and populations of marine mammals. It is necessary to take into account the depths of algal vegetation and water temperature in order to fix optimal period of harvest in each specific area. According to our data the most favourable time of seaweed harvest on Bering Island is May-June. Plants growing in shallow water areas develop more rapidly and start reproduction earlier owing to faster water warming-up than algae growing in deeper areas. That is why first of all seaweeds from shallow water zones should be harvested. According to my calculations in the intertidal zone of Commander Islands it is possible to get 3-5 kg from m² and in the upper sublittoral zone – up to 10-12 kg of raw seaweeds.

As far as the continental part of Bering Sea is concerned the present-day information on its seaweed resources is practically absent. That is why I would like to withhold recommendations on organization of harvest in this area. Commercial catch of seaweeds without scientific grounds at contemporary technological approach may cause irreplaceable losses of seaweeds and even destruction of the benthic communities. I think that studies on biodiversity of the western part of Bering Sea and ensuring of effective work of SPNA-s should precede organization of seaweed harvesting in the area.

7. Conclusion

Inventory of marine flora of the Russian sector of North Pacific is necessary not only for clarification of scientific problems of phycology but also for solving of some important practical problems. In particular, there is a threat of decrease of biodiversity of the region and loss of the still unknown species. Already known rare species included in the Red Data Book of Kamchatka (Selivanova, 2007) are also under the threat of loss. So I consider it very important to continue studies on biodiversity of marine coastal ecosystems of Bering Sea paying special attention to their plant components that form basic environments for valuable commercial objects (herring, rock trout, sea urchins etc.). In order to prevent their loss or decrease of number it is necessary to organize valid protection of resources of the shelf of the area. I am confident that sustainable harvest will make it possible to avoid destruction of marine ecosystem integrity and provide restoration of natural seaweed resources and their long-term exploitation. This will help to supply population of Kamchatka Region with this valuable food and technological natural raw material and organize its export to other areas of Russia and abroad.

8. Acknowledgment

I am grateful to my colleague Galina G. Zhigadlova for participation in algal collection and identification.

9. References

- Belyakova, G.A.; Diakov, Yu. N. & Tarasov, K.L. (2006). *Botany in 4 volumes. V. 2. Algae and fungi*, Center 'Academia' Press, ISBN 5-7695-2750-1, Moscow, Russia, 320 pp. (In Russian)
- Drude, O. (1890). *Handbuch der Pflanzengeographie*. Stuttgart, Germany, 487 S.
- Gabrielson, P.W.; Widdowson, T.B.; Lindstrom, S.C.; Hawkes, M.W. & Scagel, R.F. (2000). *Keys to the benthic marine algae and seagrasses of British Columbia, Southeast Alaska, Washington and Oregon*. Phycological Contributions N 5. University of British Columbia, ISBN 0-88865-466-9, Vancouver, BC, Canada. 189 pp.
- Gabrielson, P.W.; Widdowson, T.B. & Lindstrom, S.C. (2006). *Keys to the seaweeds and seagrasses of Southeast Alaska, British Columbia, Washington and Oregon*. Phycological Contributions N 7. University of British Columbia, ISBN 0-9763817-1-0, BC, Canada. 209 pp.
- Guiry, M.D. & Guiry, G.M. (2011). *AlgaeBase*. World-wide electronic publication, National University of Ireland, Galway. <http://www.algaebase.org> [searched on February, 24, 2011].
- Kafanov, A.I.; Borisovetz, Ye.E. & Volvenko, V.I. (2004). On application of cluster analysis in the biogeographic classifications. *Zhurnal obshchei biologii* [Journal of general biology], Vol. 65, No 3, (May-June, 2004) pp. 255-270, ISSN 0044-4596 (In Russian).
- Kongisser, R.A. (1933). Hydrobiological studies in Bering Sea at the northeastern coasts of Kamchatka (Preliminary information). *Issledovaniya dal'nevostochnykh morei SSSR* [Investigations of the Far Eastern seas of the USSR]. Issue 2, pp. 115-124 (In Russian).
- Kussakin, O.G. & Ivanova, M.B. (1978). Intertidal zone of Chukotka part of Bering Sea, In: *Litoral' Beringova morya i yugo-vostochnoi Kamchatki* [Intertidal zone of Bering Sea and

- south-eastern Kamchatka], O.G. Kussakin, (Ed.), 10–40, Nauka, Moscow, Russia (In Russian)
- Gail, G. (1936). Laminarian algae of the Far Eastern Seas. *Vestnik DV Filiala AN SSSR [Bulletin of the Far Eastern branch of Academy of Sciences of the USSR]*, issue 19, pp. 31–64. Moscow, Russia (In Russian).
- Oak, J.H.; Keum, Y.-S.; Hwang, M.S. & Oh, Y.S. (2005). New records of marine algae from Korea. II. *Algae*, Vol. 20, No 3, (September, 2005), pp. 177–181, ISSN 0273-1177.
- O’Clair, R.M. & Lindstrom, S.C. (2000). *North Pacific Seaweeds*, Plant Press, ISBN 0-9664245-1-4, Auke Bay, Alaska, USA, 162 pp.
- Perestenko, L.P. (1988). Additions to the flora of red algae from Bering Sea, *Novitates systematicae plantarum non vascularum (News of systematics of spore plants)*, Vol. 25, pp. 54–57, Leningrad, Russia, (In Russian).
- Perestenko, L.P. (1994). *Krasnye vodorosli dal'nevostochnykh morei Rossii [Red algae of the Far Eastern seas of Russia]*. “Olga” Publishing House, ISBN 5-86093-007-0, St. Petersburg, Russia, 331 pp. (In Russian).
- Petrov, Yu.E. (1972). Systematics of some species of the genus *Laminaria* Lamour. from the Far East. *Novitates systematicae plantarum non vascularum (News of systematics of spore plants)*. Vol. 9, pp. 47–52, Leningrad, Russia, (In Russian)
- Petrov, Yu.E. (1973). The genus *Alaria* in the seas of the USSR. *Novitates systematicae plantarum non vascularum (News of systematics of spore plants)*. Vol. 10, pp. 49–59, Leningrad, Russia, (In Russian)
- Rodríguez-Prieto, C.; Freshwater, D.W. & Sánchez, N. (2007). Vegetative and reproductive morphology of *Gloiocladia repens* (C. Agardh) Sánchez et Rodríguez-Prieto, comb. nov. (Rhodymeniales, Rhodophyta) with a taxonomic re-assessment of the genera *Fauchea* and *Gloiocladia*. *European Journal of Phycology*, Vol. 42, No 2 (May, 2007), pp. 145–162, ISSN 0967-0262.
- Selivanova, O. N. (2001a). Dynamics of the species diversity of macrophytic algae of the shelf of the Commander Islands, *Proceedings of the 2nd International Conference “Conservation of biodiversity of Kamchatka and coastal waters*, pp. 91–93, ISBN 5-8440-0026-9, Petropavlovsk-Kamchatskii, Russia, April 9–10, 2001 (In Russian).
- Selivanova, O.N. (2001b). Changes in the structure of marine coastal communities of the Commander Islands caused by extermination of Steller sea cow (*Hydrodamalis gigas*), *Proceedings of the International Conference “Man in the coastal zone: Experience of centuries”*, pp. 99–104, ISBN 5-89131-029-5, Petropavlovsk-Kamchatskii, Russia, September 18–20, 2001
- Selivanova, O. N. (2002). Marine benthic algae of the Russian coasts of Bering Sea (from the Ozernoi Gulf to Dezhnev Bay, including Karaginskii Island). In: *Constancea*, University of California electronic publications in Botany, Vol. 83. Available from <http://ucjeps.berkeley.edu/constancea/83/selivanova/Selivanova.html>
- Selivanova, O.N. (2007). Marine algae-macrophytes. In: *Red Data Book of Kamchatka. V. 2. Plants, mushrooms and thermophilic organisms*. O.A. Chernyagina (Ed.), 234–254, Kamchatsky Pechatny Dvor Publishing House, ISBN 5-85857-084-4, Petropavlovsk-Kamchatsky, Russia (In Russian)
- Selivanova, O.N. (2008a.) Are giant kelp algae extincting from the Northern Pacific?, *Proceedings of Symposium “Asia-Pacific Network for Global Change Research: Marine*

- biodiversity and bioresources of the North-Eastern Asia*", pp. 160-164, Cheju, Korea, October 21-22, 2008.
- Selivanova, O.N. (2008b). The present day state of the algal communities of the Commander Islands shelf and perspectives of sustainable nature use in the region. *Rybovodstvo i rybnoye khozyaistvo [Pisciculture and fish industry]*, Vol. 2, No. 7, pp. 34-38, ISSN 2074-5990, Moscow, Russia (In Russian).
- Selivanova, O.N. (2008c). *Faucheia guiryi* Selivanova sp.n.: the first record of a species of the family Faucheaceae (Rhodophyta: Rhodymeniales) in the Far Eastern seas of Russia. *Russian Journal of Marine Biology*, Vol. 34, No 6, (December, 2008), pp. 351-358 (translated into English from Russian), ISSN 1063-0740.
- Selivanova, O.N. (2009). *Gloiocladia guiryi* (Selivanova) comb. nov. – a new name for the first member of the family Faucheaceae (Rhodymeniales, Rhodophyta) from the Russian Pacific. *Phycologia*, Vol. 48, No 5 (September, 2009) pp. 439–440, ISSN 0031-8884.
- Selivanova, O.N. & Zhigadlova, G.G. (1997a). Marine algae of the Commander Islands. Preliminary remarks on the revision of the flora. I. Chlorophyta. *Botanica Marina*. Vol. 40, No pp. 1-8.
- Selivanova, O.N. & Zhigadlova, G.G. (1997b). Marine algae of the Commander Islands. Preliminary remarks on the revision of the flora. II. Phaeophyta. *Botanica Marina*. Vol. 40, No pp. 9-13.
- Selivanova, O.N. & Zhigadlova, G.G. (1997c). Marine algae of the Commander Islands. Preliminary remarks on the revision of the flora. III. Rhodophyta. *Botanica Marina*. Vol. 40, No pp. 15-24.
- Selivanova, O.N. & Zhigadlova, G.G. (2000). Benthic algae of the Russian coasts of Bering Sea (including Commander Islands). I. Medny Island. In: *Proceedings of Kamchatka Institute of Ecology and Nature management*, R.S. Moiseev, (Ed.), 71-108, Kamchatsky Pechatny Dvor Publishing House, ISBN 5-85857-108-5, Petropavlovsk-Kamchatskii, Russia (In Russian)
- Selivanova, O.N. & Zhigadlova, G.G. (2003). Benthic algae of the Russian coasts of Bering Sea (including Commander Islands). II. Bering Island. In: *Proceedings of Kamchatka Institute of Ecology and Nature management*, R.S. Moiseev, (Ed.), 172-208, Kamchatsky Pechatny Dvor Publishing House, ISBN 5-85857-054-02, Petropavlovsk-Kamchatskii, Russia, Issue 4 (In Russian)
- Selivanova, O.N. & Zhigadlova, G.G. (2009). Algofloristic studies on Starichkov Island (Eastern Kamchatka). Species composition and distribution of benthic algae. *Journal of Siberian Federal University, Biology*, Vol. 2, No 3, (September, 2009), ISSN 1997-1389, pp. 271-285 (In Russian).
- Selivanova, O.N. & Zhigadlova, G.G. (2010). Diversity, systematics, distribution and resources of marine algae-macrophytes of the Western Bering Sea. In: *Current state of the ecosystem of the Western Bering Sea*, P.S. Makarevich, (Ed.), 37- 78, Southern Science Center, Russian Academy of Sciences Publishing House, ISBN 978-5-902982-78-4, Rostov-on-Don, Russia (In Russian)
- Tolstikova, N.E. (1974). New data on ecology of subtidal macrophytes of Anadyrskii Gulf, Bering Sea. *Novitates systematicae plantarum non vascularum (News of systematics of spore plants)*. Vol. 11, pp. 147–152, Leningrad, Russia (In Russian).

- Vegener, A. (1984). Die Entstehung der Kontinente und Ozeane [*Origin of the continents and oceans*], P.N. Kropotkin, (Ed.), Nauka, Leningrad, Russia, 285 pp. (translated from German into Russian).
- Vinogradova, K.L. (1973a). Species composition of algae from the intertidal and subtidal zones of the north-western part of Bering Sea. *Novitates systematicae plantarum non vascularum (News of systematics of spore plants)*. Vol. 10, pp. 32–44. Leningrad, Russia (In Russian).
- Vinogradova, K.L. (1973b). On new species of *Rhodomela* Ag. and *Polycerea* J.Ag. from Bering Sea. *Novitates systematicae plantarum non vascularum (News of systematics of spore plants)*. Vol. 10, pp. 22–28. Leningrad, Russia (In Russian).
- Vinogradova, K.L. (1974). *Ulvovye vodorosli (Chlorophyta) morei SSSR [Ulvaceous algae (Chlorophyta) of the seas of the USSR]*. Nauka. Leningrad, Russia, 165 pp. (In Russian).
- Vinogradova, K.L. (1978). Algae of the south-western coasts of Bering Sea. *Novitates systematicae plantarum non vascularum (News of systematics of spore plants)*. Vol. 15, pp. 3–11. Leningrad, Russia. (In Russian).
- Vinogradova, K.L. (1979). *Opredelitel' vodoroslei dal'nevostochnykh morei. Zelenye vodorosli [Guide-book to the algae of the Far Eastern Seas. Green Algae]*. Nauka. Leningrad, Russia. 147 pp. (In Russian).
- Vinogradova, K.L.; Klochko, N.G. & Perestenko, L.P. (1978). List of intertidal algae of the Eastern Kamchatka and western coasts of Bering Sea. In: *Litoral' Beringova morya i yugo-vostochnoi Kamchatki [Intertidal zone of Bering Sea and south-eastern Kamchatka]*. O.G. Kussakin (ed.), 150–155. Nauka. Moscow, Russia (In Russian).
- Vinogradova, K.L. & Perestenko, L.P. (1978). Main regularities of algal distribution in the intertidal zone of the western part of Bering Sea. In: *Zakonomernosti raspredeleniya i ekologiya pribrezhnykh biotsenozov [Regularities of distribution and ecology of the coastal communities]*. pp. 72–75. Nauka. Leningrad. (In Russian).
- Xia, B.-M. & Wang, Y.-Q. (2000). Studies on some new red algae from China (II). *Acta Phytotaxonomica Sinica*, vol. 38, No 1, pp. 86–95, ISSN 05291526.
- Yoshida, T. (1998). *Marine algae of Japan*. Uchida Rokakuho Publishing CO., ISBN 4-7536-4049-3 C3045, Tokyo, Japan, 1222 pp. (In Japanese).
- Zhigadlova, G.G. (2009). Additions to the flora of macroalgae from southeast Kamchatka and Commander Islands. *Proceedings of the 10th International Conference "Conservation of biodiversity of Kamchatka and coastal waters"*, pp. 232–235, ISBN 978-5-9610-0133-4, Petropavlovsk-Kamchatskii, Russia, November 17–18, 2009 (In Russian).
- Zhigadlova, G.G. & Selivanova, O.N. (2004). Benthic algae of the Russian coasts of Bering Sea. III. Karaginskii Gulf. In: *Proceedings of Kamchatka Branch of Pacific Institute of Geography*, R.S. Moiseev, (Ed.), 47–89, Kamchatsky Pechatny Dvor Publishing House, ISBN 5-85857-062-3, Petropavlovsk-Kamchatskii, Russia, Issue 5 (In Russian).



The Dynamical Processes of Biodiversity - Case Studies of Evolution and Spatial Distribution

Edited by PhD. Oscar Grillo

ISBN 978-953-307-772-7

Hard cover, 366 pages

Publisher InTech

Published online 02, December, 2011

Published in print edition December, 2011

Driven by the increasing necessity to define the biological diversity frame of widespread, endemic and threatened species, as well as by the stimulating chance to describe new species, the study of the evolutive and spatial dynamics is in constant execution. Systematic overviews, biogeographic and phylogenetic backgrounds, species composition and distribution in restricted areas are focal topics of the 15 interesting independent chapters collected in this book, chosen to offer to the reader an overall view of the present condition in which our planet is.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Olga N. Selivanova (2011). Marine Macrophytic Algae of the Western Sector of North Pacific (Russia), The Dynamical Processes of Biodiversity - Case Studies of Evolution and Spatial Distribution, PhD. Oscar Grillo (Ed.), ISBN: 978-953-307-772-7, InTech, Available from: <http://www.intechopen.com/books/the-dynamical-processes-of-biodiversity-case-studies-of-evolution-and-spatial-distribution/marine-macrophytic-algae-of-the-western-sector-of-north-pacific-russia->

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2011 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](https://creativecommons.org/licenses/by/3.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen