We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Aquaculture Water Quality for Small-Scale Producers

Oscar Alatorre-Jácome, Fernando García-Trejo,
Enrique Rico-García and Genaro M. Soto-Zarazúa
*C.A. de Ingeniería de Biosistemas, Campus Amazcala,
Facultad de Ingeniería, Universidad Autónoma de Querétaro,
Amazcala, Mpio, El Marqués, C.P. 76260, Querétaro, Qro
México*

1. Introduction

Today we know that solar energy can be stored as edible biomass by photosynthesis, but 10,000 years ago this assumption was unthinkable. However, the Neolithic man could develop agriculture, based on plant germination observations. Nowadays, the systems used for food production are practically the same (Koning et al., 2008). But for neolithic man, the empirical observation of germination processes and biological cycles of certain species ended in the discovery of agriculture.

Even our agriculture is still the same, the climate scenario is changing. Our current carbon-based economy has caused massive greenhouse emissions, and consequences are global warming and the depletion of the ozone layer. The biological, economical and social impact on the environment is significant. Water scarcity and natural catastrophes (tsunamis, earthquakes, floods and erosion) damage cultivable land, reducing local food availability (Franck et al., 2011, IPCC, 2007).

Clearly, it is necessary the innovation of new food production systems. Aquatic food production is an alternative to land-based, 2-dimensional systems (Alagaraja, 2007). Due to physical differences between air and water (thermal conductivity, heat capacity, density, etc), most aquatic organisms do not use energy in thermal regulation. The consequence is a more efficient food-tissue conversion ratio compared with land-based system yield (FAO, 2010).

There are limitations, of course. Because overfishing causes fish stock depletion, aquaculture has become an alternative to provide fish products (Naylor, 2009; Brander, 2007; Pillay et al., 2003; Naylor, 2000).

The notorious growth in aquaculture has been possible by the application of science. Aquacultural and fisheries science produce new and replicable knowledge based on scientific method. The research is usually done in farms or in a laboratory. The mathematical and analytical methods used by researches are generally robust and provided a verifiable knowledge. Nevertheless, there is a gap between the scientific results and its practical use. Lukefar (1999) pointed out that the examples taught on classroom lectures are usually based

under realistic, idealized situations. But there is a necessity to find practical solutions applicable for commercial aquaculture.

The problem of the academic extension is an important issue in aquaculture if we consider that small-scale aquaculture provides almost half of the worldwide inland fish production (FAO, 2010). Kawarazuka (2010) commented “aquaculture and small-scale fisheries can improve the food intake”. Mohanty et al. (2010) mentioned that at least 90 % of the people involved in aquacultural practices works in small fisheries.

Facing the problem of the continuous rising world population, it seems to be clear that the good use of all natural resources (including small freshwater bodies), must be done. So there is an important, unattended practical knowledge field to be considered. Technologies for small-scale fish producers are ready available (Van Gorder, 2003), and the current knowledge on aquatic science can be helpful for the producer with the goal to optimize its own resources (time, land, water and energy).

The objective of the present chapter is to present selected topics on water quality management, especially if they can be carried out by low-cost technologies in small systems. Due to the close relationship among water quality and yields, we hope it can be helpful as a practical guide for fish producers based on scientific principles.

2. Overview of water quality in aquaculture

Water is the physical environment where fish develop, growth and reproduce. The dynamic of the mass and energy involved on an aquaculture system is complex, because bacteria, algae and fish growth together in the pond (Wheaton, 1982). The main energy input comes from sunlight, and nutrients of the system are commonly provided by pelletized fish feed. So the transformation of the elements carried out by autotrophic and heterotrophic organisms change the physical/chemical/biological composition of the water (Hargreaves, 1998).

Temperature and pH are fundamental for the aquatic living organisms, due to the intimate relationship between them and the velocity of its biochemical processes. Oxygen (O_2) and carbon dioxide (CO_2) are important molecules because they are involved in photosynthesis and respiration processes. Nutrients like nitrogen are essential on biological metabolism: when is ingested as protein or amino acid, it can be incorporated by the organism as functional proteins or in structural tissues. But when is excreted by the fish as ammonium (NH_3), in certain circumstances it can be toxic and even lethal in high doses. Another nutrients, like phosphorus, potassium and calcium are also important, and they lack are usually diagnosed by deficiency.

Besides the physical and chemical factors, the biotic component also can change the water composition. For example, algae can consume or produce oxygen and carbon dioxide, depending on light presence or absence. The nitrogen, phosphorus and potassium can be used and assimilated by unicellular organism, incorporating them as biomolecules. Moreover, the nitrogen can be also used to produce energy on chemoautotrophic biochemical cycles (Hargreaves, 1998).

So, the biological impacts of water quality over the cultivated species could be analyzed under physical, chemical or biological perspectives. For practical purposes, a quick

overview of the basic principles is given below. The parameters mentioned have been divided into two main categories: Critical and important parameters.

2.1 Critical water quality parameters

The critical parameters are very important in the aquacultural system. They are temperature, pH, dissolved oxygen (DO) and ammonia. They must be measure daily, or in the case of intensive systems, all day long. They do influence the physical properties and chemical composition of the water, and thereafter its correct management can improve the overall fish performance (health and growth). In the other hand, if they are not properly attended, the consequences can be serious, varying from poor growth rates, stress, and death.

2.1.1 Temperature

Temperature is probably the most important physical variable on aquatic ecology. It affects directly the metabolism of all living organisms. As a consequence, temperature set the growth, development and reproduction rates in biological species in biological species. This fact is very useful in aquaculture: because fish do not expend energy on corporal temperature regulation, they can assimilate almost the food nutrients into muscular tissue (Soto-Zarazúa et al., 2011). As results of an adequate temperature condition, the biomass production and final yield of the fish farm can increase.

It is important to remark the influence of temperature in fish respiration rate. A rise of temperature causes more oxygen consumption in bacteria, algae and fish (Boyd, 1998). Because respiration implies carbon dioxide release and energy consumption, the gas balance can be dangerous for fish. If there are enough inorganic nutrients in the water, the algae biomass can increase to considerable levels. Even some algae species can double its biomass in only 3.5 hours (Brennan & Owende, 2010). Then, the elevated rates of nutrient assimilation will produce significant impacts on the water quality.

In other hand, a higher temperature produces higher metabolic rates. The increment in fish metabolism enhances the protein breakdown. As a consequence, the release of NH_3 by fish will be high, too. The resulting combination of high temperatures with NH_3 high concentration is very a toxic environment (Eshchar et al., 2006).

Finally, if the exposure to high temperatures is very long, the structure of the proteins begins to break, causing fish death.

2.1.2 pH

The pH is a measure of acidity and basicity inside an aqueous solution. It indicates the concentration of hydrogen ions on water. When pH is below 7 the solution is considered acid, and when is above 7 is named basic or alkaline. Distilled water has a pH of 7. pH can be defined as the negative logarithm of the molar concentration of hydrogen ions on water. In mathematical notation, pH is described as

$$pH = \log \frac{1}{[H^+]} = -\log[H^+]$$

Where $[H^+]$ is the concentration of hydrogen ions. Fish and other vertebrates have a pH blood value near to 7.4. The contact between environmental water and fish blood is only separated by one or two cells of the gills. An ideal pH for an aquacultural system must be near to 7. The lethal limits are below 5 and above 10, for most of the fish species.

There is an important relation between fish respiration and pH. In the gills the gaseous interchange of O_2 instead CO_2 occurs. This interchange can be difficult if pH is not optimum. The effects are called Bohr and root. So even if we have enough oxygen in our system, if pH is not adequate fish could not breathe (Wurts & Durborow, 1992).

pH, like temperature, is always changing. For example: In afternoon, the oxygen concentration decrease and the phytoplankton photosynthesis stops by the absence of sunlight. The concentrations of O_2 and CO_2 began to change, and the pH can vary due its intimate relationship with CO_2 equilibrium (Wurts & Durborow, 1992). Another important factor to take into account is that cinematic of certain types of bacteria can change in low pH, so the mass and energy transformations of the pool carried out by unicellular organism can influence the environment (Ebeling et al., 2006).

2.1.3 Dissolved oxygen

According to Rumei et al. (2003), dissolved oxygen is the most important manageable variable in aquaculture. The oxygen is necessary to glucose breakdown and energy release inside fish cells. However, its diffusivity and availability on water is mediated by temperature, elevation and salinity (Boyd, 1998).

Low concentrations of oxygen can produce negative impacts on fish health, like poor growth performance, low feeding rate, and increase risk on potential diseases or even fish death. These impacts are specific for every fish species. The particular oxygen requirements depend principally on the fish biology. For example, trout needs a high quantity of oxygen (about 7 ppm), but catfish (bottom, detritivore fish) can survive with only 0.5 ppm of oxygen (Akinwole & Faturoti, 2007).

The presence of dissolved oxygen on aquaculture water ponds depends on physical factors (salinity, temperature and altitude), algae photosynthesis or artificial supplying (Boyd, 1998). Oxygen consumption depends on the carry capacity of the systems. The fish biomass is an important factor to be considered in a system, because the food supplied to the fish plays an important role in water ponds biogeochemistry (Timmons, 2002; Hargreaves, 1998).

2.1.4 Ammonia

The ammonia is a nitrogen compound excreted by fish through gills and faeces. So the amount of ammonia is in direct relation with the amount of feed input on the pond. Ammonia can also be produced in pond by organic material decomposition driven by bacterial activity (Durborow et al., 1997a).

The ammonia is presented in two forms: the high toxic un-ionized ammonia (NH_3), and the non-toxic ionized ammonia (NH_4^+). They are in chemical equilibrium driven by temperature and pH. The sum of NH_3 and NH_4^+ is called Total Nitrogen Ammonia (TAN). In general,

the fraction of toxic ammonia increases at elevated temperature and high values of pH (Durborow et al., 1997a), as it can be seen in the following equation:

$$\frac{NH_3}{TAN} = \frac{K_{eq}}{10^{-pH} + K_{eq}}$$

Where k_{eq} is the negative logarithm of pK ($pK = -\log(K_{eq})$), and $pK = 0.09018 + 2727.9 - (T + 273.1) + (0.1552 + 0.0003142T)I$, T is temperature ($^{\circ}C$) and I is Ionic strength (M) (Eshchar et al., 2006). For more convenient purposes, computations of ammonia toxic fractions can be made with the tables provided by Durborow et al. (1997a). Measures of TAN, pH and temperature are required.

2.2 Important water quality parameters

The dynamic of the important parameters occurs slower than the reactions implicit on critical ones, so they can be monitored with less frequency. Solids, nitrite (NO_2), nitrate (NO_3), carbon dioxide (CO_2), hardness and alkalinity can be measure once a week. Lethal effects on the fish at high concentrations of this parameters are not common, but their accumulation can affects directly and indirectly the fish growth. Also, if unfavourable conditions are presented, an enhanced risk to infections or diseases can be presented.

2.2.1 Solids

In aquaculture, the solids are placed into a special category because in most of the cases they can be controlled by good management practices. Solids in pond are presented as uneaten food, faeces, fish scales, dead bacteria and algae, dust, and dead fish (Cripps and Bergheim, 2000). The adverse consequences driven by the presence of organic solids in the system are caused mainly by bacterial processes. It implies additional oxygen consumption and carbon dioxide release, among other effects. In some anoxic environments, hydrogen sulfide (H_2S) and ammonia can be release to water by bacterial anaerobic processes.

In general, the solids can be classified in three major divisions. 1) Settleable solids, which can be easily separated by sedimentation or decantation (for example, uneaten feed, fish scales, faeces); 2) Suspended solids, which are very fine solids without the capacity of rapid sedimentation (about 1-10 μm on diameter); and 3) dissolved solids, which are nano-scale elementary forms as molecules or atoms. (Losordo et al., 1999; Malone, 1991).

In general, the impacts of the solids in aquaculture systems are negative. For example, in the surface when the water turbidity rises, the photosynthetic activity in algae decreases. In bottom, accumulation of solid wastes causes anaerobic zones, in which undesirable bacteria can proliferate. In ponds, high clay turbidity usually causes acidity, low nutrient levels, and limited light penetration for photosynthesis (Yi et al., 2003). Dead fish are solids too, and its presence on culture water can be a factor for pathogenic propagation inside the farms (Cripps & Bergheim, 2000).

Is useful to think that solids on aquacultural environments are mainly composed by organic material. If they remain in the water, they become basic nutrients, like nitrogen and phosphorus that could be easily assimilated by microscopic organisms. So the total oxygen budget will rise (Timmons et al., 2002).

2.2.2 Nitrite

The presence of nitrite inside an aquaculture pond is often caused by ammonia biological oxidation (Hargreaves, 1998). The stoichiometric reaction is the following:

As a result, there is a free energy yield (ΔG) of $-65 \text{ kcal mole}^{-1}$ from the ammonia oxidation.

This reaction is carried out by bacteria that naturally grow in the aquacultural environment. Bacteria of the genus *Nitrosomonas* are the main responsible for nitrification in aquaculture, but there are other important genera involved in this process (Hovanek & DeLong, 1996; Ebeling et al., 2006).

The physiological effects of nitrite in fish health are mainly caused by a chemical reaction on haemoglobin. The role of this protein is the oxygen transportation all over the blood stream. In the presence of nitrite, haemoglobin becomes methaemoglobin, a non-efficient oxygen transporter (Jensen, 2003). As a result of oxygen deficit, the fish blood becomes brown, and a gasping behaviour can be observed. If nitrite intoxication remains unattended, massive fish death can occur after a short time caused by hypoxia (Masser et al., 1999).

High nitrogen concentrations in ponds occur more frequently in the fall and spring, when low and fluctuating temperatures cause decay rates on phytoplankton and bacteria metabolisms (Durborow et al., 1997b).

2.2.3 Nitrate

The presence of nitrate in aquaculture water is delivered as a waste product of organic bacterial activity in the pond. The reaction involves the presence of nitrite and oxygen. The following stoichiometric reaction shows the overall process (Wheaton, 1982).

In total, when a nitrite molecule is oxidized, a free energy yield (ΔG) of $-18 \text{ kcal mole}^{-1}$ is released. In aquacultural ponds, the most representative bacteria genus that can perform the nitrite conversion to nitrate is *Nitrobacter*, but other genera of bacteria are commonly presented during the nitrification (Camargo et al., 2005).

Nitrate is commonly controlled in aquaculture systems by dilution. In intensive recirculation systems between 5 and 10% of water are removed and replaced every day. In systems with low technification, the daily water exchange usually is more than 10%.

A natural pathway to remove the nitrite in an aquacultural system is done by denitrification. The reaction is carried out by bacteria in absence of oxygen and in presence of methanol as a carbon compound. The general reaction is done in two steps (Van Rijn et al., 2006):

2.2.4 Carbon dioxide

Carbon dioxide (CO_2) in aquatic systems is very important, because its presence is required in some chemical and biological processes. For example, CO_2 interacts with water to form a

natural buffer system which helps to maintain a constant pH. Is also important for biological primary production, because is necessary in algal photosynthesis to synthesize glucose. Is important to note that biological activity can also release CO₂ to the media, when fish and algae breathe. Some bacteria also need CO₂ to maintain its metabolic cycles in constant function.

Bacteria also release CO₂, both autotrophs and heterotrophs. For example: The free acid produced during nitrification reacts with bicarbonate alkalinity in water to release more CO₂ than autotrophs consume. So for every gram of TAN metabolized, 4.6 ppm of oxygen will be consumed, and 5.9 g of CO₂ will be released. In heterotrophic reactions, for every gram of O₂ consumed, 1.38 g of CO₂ is released (Summerfelt & Sharrer, 2004).

When CO₂ is dissolved in water, part of it combines to form carbonic acid (H₂CO₃). This weak acid tend to react with calcium carbonate (CaCO₃) to form calcium bicarbonate [Ca(HCO₃)], which is often dissociated on hydrogen ions and carbonate ions (Wheaton, 1982):

These four reactions are mediated by hydrogen concentration. They proceed to the right when pH rises and they go to the left when pH decreases. Therefore, the carbon dioxide usually can be found in water under four different forms: As free gas (CO₂), as carbonic acid (H₂CO₃), as carbonate (CO₃²⁻) and as bicarbonate (HCO₃⁻) (Wheaton, 1982). Thus, the chemical importance of CO₂ relies in the fact that is a pH buffer, and it has a lot of relation with other physicochemical parameters, like hardness and alkalinity.

2.2.5 Hardness

The hardness is defined as the total concentration of calcium and magnesium ions, expressed as calcium carbonated. However, if other metallic ions are presented in the water (Al, Fe, Mg, Sr, Zn) they can be also considered in the definition (Wheaton, 1982).

The hardness is an important water quality parameter because a direct relation between water metal content and pH variations exists. When the concentration of Ca and Mg trends to be higher, the buffering capacity of the water becomes higher, too, and is more capable to smooth pH variations. In other words, hard water is more stable than soft water.

In the biological perspective, calcium is important in fish metabolism, because is used on the scale and bone formation, and to keep the adequate balance of Na and K in the blood (Wurst & Durborow, 1992). Calcium is often required in neural synapses, and in physiological ion balance. In the case of magnesium, it is used by photosynthetic organism because is embedded in the center of the chlorophyll molecule, and it is also required as prosthetic group in proteins (Müller-Esterl, 2008)

In the farm, hardness is very important when the organisms are cultivated for reproduction purposes in hatcheries. If invertebrates are cultivated, hardness becomes an important variable to consider, because Ca and Mg are very important in the formation of hard parts (i.e., exoskeleton or shells).

Changes in the water hardness are slowly. They take place typically in weeks or months.

However, is a good management practice to monitoring this variable constantly.

2.2.6 Alkalinity

The alkalinity is the amount of acids (H^+) that water can neutralize before to reach a given pH. It is defined as the stoichiometric sum of the bases in a solution. Common bases found in fish ponds include carbonates, bicarbonates, hydroxides, phosphates and berates. Carbonates and bicarbonates are the most common and the most important components of alkalinity. Because total alkalinity can be expressed as ppm of $CaCO_3$, is common to confuse it with hardness (Wurts & Durborow, 1992).

Like another water quality parameters, alkalinity can be affected by the biological activity of systems. Because photosynthesis of phytoplankton requires CO_2 to synthesize glucose, pH in the water increases due to inorganic carbon adsorption in water (mainly H_2CO_3 and CO_3^{2-}). During long periods of intensive photosynthesis, the release of carbonate can elevate the pH levels over 9. These effects can be observed if water has low alkalinity (20 to 50 ppm) or low bicarbonate (75 to 200 ppm). High photosynthetic activity can be presented when the sodium and potassium carbonates are dissolved in water, because they are more soluble than calcium and magnesium bicarbonates (Wurts & Durborow, 1992).

3. Low-tech water quality management

Currently, the aquacultural engineering gives a considerable number of solutions to control the variables involved in aquaculture water quality. In general, they are available commercial devices and chemical products to control water quality. However, there are alternative techniques used by scientist that considered the pool as a bioreactor. New aquaculture techniques are bio-flocs (Avnimelech, 2006), Integrated Multi-trophic Aquaculture (IMTA) (Chopin, 2003), greenwater systems (Hargreaves, 2006), Zero-Exchange water systems (Panjaitan, 2010; Olvera-Olvera et al., 2009) and aquaponics (Rakocy & Hargreaves, 1993), among others.

3.1 Temperature

The temperature control on aquaculture could be difficult, because the high specific heat capacity of water ($4.18 \text{ kJ kg}^{-1} \text{ }^\circ\text{K}^{-1}$) implies a high amount of energy when water is heated or cooled. In addition, the water volume used on aquacultural facilities is frequently high, so the monetary and environmental cost to raise water temperature could be unaffordable (Seginer & Mozes, 2008).

There are several alternatives to increase thermal stability on aquaculture. The most effective system is the use of greenhouse to cover aquaculture ponds or tanks (Soto-Zarazúa et al, 2011; Fuller, 2007). Some farms use heating pumps, thermosolar systems, fossil fuel heaters or electric resistances, but all of them are expensive ways to heat water, both by initial and operational cost. Besides, some of them imply negative environmental impacts caused by greenhouse gas emissions (Mohanty et al., 2010).

Nevertheless, there are alternatives for the small-scale producers. The most effective strategy is a good planning on farm building. Then, the most important thing is to pick a good geographical location. The climate and the temperature of the make-up water must be adequate for the cultivated fish. An analysis of the local climatologically data, if available, can give us an idea if the selected location is proper to our intentions.

If low air temperatures are presented, covering the tanks with plastic sheets could be useful (Van Gorder, 2003; Crab, 2009). On the other hand, if temperature rises, a packed aeration column can be built as a chiller (Wheaton, 1982). In both cases (low and high water temperatures), to add new make-up water can be useful. It is also important to remember that some fish doesn't grow on winter, like tilapia (Crab et al., 2009).

If fish tanks are used, it could be helpful to cover its walls with heat insulators. Elastomeric foams are a good choice, because they are generally designed to support extreme insulation and hard management. Fiber glass and polystyrene foams could be used, too (Alatorre, 2010).

Another option is to consider solar heating water systems. Despite the high initial cost, they can help to raise the water temperature from 1 to 5 °C. Solar pool blankets, geodesic structures and active/passive solar collectors have been tested with considerable results (Fuller, 2007). A practical guide and cases studies is available in www.retscreen.net. A complete guide to thermosolar processes can be found on Duffie and Beckman (1992).

3.2 pH

A common action to pH control is the addition of chemical substances. When pH is low, it could be useful adding lime (Wurts and Durborow, 1992). If pH rises, small amounts of phosphoric acid or acetic acid can help to neutralize water. Some people uses sulphuric acid, but it can be dangerous if there is no previous experience on dangerous substances management. Another way to pH control is the enrichment of the biological activity in the pool (Hargreaves, 2006).

In our experience, many traditional aquaculture managers do not monitoring pH. This can be a bad choice, because as we discussed previously, a certain combination of temperature, pH and ammonium can kill entire production stocks.

In the case of aquaponics systems, pH management is fundamental. In this kind of systems plants, fish and bacteria are cultivated. In general, the nitrogen cycle is completed inside the culture water, and the final metabolite (nitrate) is assimilated by plants. As a result, the nutrients provided by fish feed are recycled, and the negative environment impacts are diminished (Diver, 2006). But there is a problem. The optimum pH value for plants, fish and bacteria is 6, 7 and 8 respectively. There is necessary reconciling pH in this systems. Tyson (2008a,b) recommended a pH value of 8 to improve nitrogen assimilation by bacteria.

3.3 Dissolved oxygen

Mechanical aeration is by far the most common and effective way to increasing DO concentrations in ponds. In semi-intensive aquaculture, aeration is applied in case of emergency. (Boyd, 1998).

Paddlewheel aerators and propeller-aspirator-pumps are the most common aerators used in aquaculture. Aeration amounts varies from 1-2 kW ha⁻¹ in extensive cultures, to 15-20 kW ha⁻¹ in intensive culture of marine shrimp. For every kW of aeration extra, the gains are estimated over 500 kg fish/crustacean biomass (Boyd, 1998).

All basic types of mechanical aerators have been used in aquaculture, but vertical pumps, pump sprayers, propeller-aspirator-pumps, paddle wheels, and diffused-air systems are the

most common. However, paddle wheels aerators and propeller-aspirator-pumps are the most efficient devices (Boyd, 1998).

There are some useful indications about the use of aerators. In semi-intensive culture, is common to turn on mechanical aeration at night, when is a lack of photosynthesis and the respiration rate is in a maximum value. However, is important to note that if the aerators are operated on sunny afternoons when water is supersaturated, oxygen will be lost (degassed) from the water (Tucker, 2005)

When commercial aerators are not available, a degassing column can be constructed with local available materials. Wheaton (1982) mentioned how to build a simple aeration tower. Tucker (2005) gives photos and a short description about it uses on farm systems. Theoretical background can be found in Boyd (1998), Summerfelt et al. (2000) and Vinatea & Carvalho (2007)

3.4 Nitrogen compounds

Ebeling et al. (2006) pointed out the following: for every gram of ammonia-nitrogen converted to nitrate-nitrogen, 4.18 of dissolved oxygen and 7.05 g of alkalinity (1.69 g of inorganic carbon) are consumed and 0.20 g of microbial biomass (0.105 g organic carbon) and 5.85 g of CO₂ (1.59 g inorganic carbon) are produced. So the nitrogen cycle in pond water affects the physical, chemical and biological components present in the system. To control the accumulation of every kind of components in the pond the most usual method is the addition of make-up water. However, there are alternatives to manage the levels of nitrogen compounds. Some of them are described below.

3.4.1 Ammonia

The most extended method to control ammonia inside aquacultural ponds is to keep a good feeding schedule based on nutrimental tables. If the feeding regimen in the fish farm is intensive, then pH control is also recommended. If pH can be maintained lower than 8, the toxic fraction of TAN will remain in minimal percentage (Hargreaves and Tucker, 2004).

A common method to control ammonia in recirculating aquaculture systems is a biofilter addition in the system. Biofilters design and theoretical foundations are already in literature (Bazil (2006), Drennan II et al. (2006), Eding et al. (2006), Gutierrez-Wing and Malone (2006), Kuo-Feng & Kuo-Ling (2004)). Examples of applications and affordable designs can be found in Soto-Zarazúa et al. (2010), Timmons et al. (2006), Al-Hafedh et al. (2003) and Ridha & Cruz (2001)

Another method to control ammonia is the increment of microbiological activity inside the pond (Hargreaves and Tucker, 2004). The addition of organic carbon in relation with nitrogen concentration is very useful (Crab et al., 2009). Theoretical background of this technique (also named bio-floc technology) is already available in Avnimelech (2006, 2003 and 1999).

3.4.2 Nitrite

A common practice to reduce nitrite toxicity is the elevation of chloride concentration in the culture water (Losordo et al., 1998). For this purpose, common salt (sodium chloride) is

used. Calcium chloride can also be used. A high chloride:nitrate ratio of 10:1 is used to prevent brown blood disease. In catfish, for example, is recommended to maintain at least 100 ppm chloride in pond waters (Durborow et al., 1997).

When brown gill disease is not presented, a good practice is to flush water. In general, the absorption of NO_2 by bacteria is presented in tanks and ponds. To enhance its activity, biofiltration is recommended.

3.4.3 Nitrate

In general, nitrate is a non-toxic form of nitrogen in pond. However, nitrate management is usually carried out by water exchanges. Phytoplankton and bacterial uptake is another method to assimilate nitrate into cellular tissue (Gross et al., 2000). Aquaponics is another way to assimilate NO_3 into fresh, marketable plant biomass, as lettuce, tomato, basil and other crops (Graber & Junge, 2009; Rico-García et al., 2009; Savidov et al., 2005).

Denitrification is also an alternative to eliminate aquacultural nitrate. Theory and practice examples can be found in van Rijn et al., 2006.

3.5 Solids

The management of solids include feed design and management, flow regulation and separation treatment technology (Cripps and Bergheim, 2000). In small scale aquaculture, a good start point is to asses an adequate feed schedule. In our experience, we have noted that overfeeding is usual in new managers, so the detriment of the water quality is faster.

Fish also waste a lot of feed. In the case of tilapia, even 50% of their feed can be wasted (Avnimelech, 2003). So the experience of the manager is fundamental in solids control.

A complementary approach to treat solid wastes is a quickest remotion of them (Timmons et al., 1998). In general, the settleable solids can be easily remove from the culture water. If circular tanks are used, the centrifugal forces and conical bottoms must help to accumulate the solids in the center drain. (Timmons et al., 1998; Summerfelt et al., 1998). If rectangular tanks are used, is recommended to adjust length/width ratios to increased bottom velocities and reduced biosolid accumulation (Oca and Masaló, 2007). In some cases additional components can be used, like settling basins or hydroclones (Wheaton, 1982). For aquaponics, the size reduction of solid wastes from pellets to fine particles or even until basic organic compounds is desirable (Rakocy & Hargreaves, 1993).

If extensive ponds are used, the use of chemicals to enhance flocculation of organic particulate material could be useful. The addition of gypsum helps to diminish the negative electrical charges between particulate material, enhancing the flocculation and sedimentation of suspended solids. The recommended amounts are between 100 to 300 mg per liter (Hargreaves, 1999).

3.6 Carbon dioxide (CO_2)

In general terms, rarely CO_2 concentration cause problems in fish ponds with sufficient alkalinity. However, if the pond is deeper than 5 feet and poor mixing is presented, there is a risk of water stratification (Hargreaves and Brunson, 1996). Vigorous aeration can prevent

stratification, and it would be helpful if CO₂ is saturated on pond water (Wurts and Durborow, 1992).

Other alternative is chemical treatment. The addition of quicklime, hydrated lime or sodium carbonate will increase the alkalinity of the pond, dropping the presence of dissolved CO₂ in water. Treatment calculations can be found on Hargreaves and Brunson, (1996). In the case of intensive systems, CO₂ could be a problem. In this case, the addition of a degassing tower is recommended (Summerfelt and Sharrer, 2004).

4. Conclusions

The processes involved in aquaculture water quality are affected by many variables and they are generally complex. However, the driving forces involved in fish farms can be controlled in order to increase fish productivity. This control involves learned skills to make the right decisions to correct present problems and to prevent new ones.

In this chapter we had discussed extensively the theoretical foundations of fish water quality, with emphasis on small-scale aquaculture correction techniques. As it can be seen, physical, chemical and biological processes involved in the pond dynamics are strongly related, and they can't be considered as isolated phenomena. The understanding of these processes can be useful to improve the final yields on the farm with the minimal waste of time, energy and money. As a result, the sustainability of the protein production can be rise and the negative environmental impact can be diminished.

5. References

- Akinwale, A. O. & Faturoti, E. O. (2007). Biological performance of African catfish (*Clarias gariepinus*) cultured in recirculating system in Ibadan. *Aquacultural engineering* 36:18-23.
- Al-Hafedh Y. S., Alam A. & Alam M. A. (2003). Performance of plastic biofilter media with different configuration in a water recirculation system for the culture of Nile tilapia (*Oreochromis niloticus*). *Aquacultural engineering* 29:139-154.
- Alagaraja, K. (1991). Aquaculture productivity. On *Proceedings of the Symposium on Aquaculture Productivity*. December 1998. Hindustan Lever Research Foundation. Oxford & IBH Publishing Co. PVT. LTD. New Delhi. Sinha, V.R.P and H. C. Srivastava.
- Alatorre-Jácome, O. (2010). Implementación de un sistema termosolar para alevinaje (Nursery) de tilapia (*Oreochromis niloticus* L.). MSc. thesis dissertation. Universidad Autónoma de Querétaro, México.
- Alatorre-Jácome, O., Rico-García, E., Soto-Zarazúa, G. M. García-Trejo, F. & Herrera-Ruiz, G. (2011). A termosolar nursery for tilapia (*Oreochromis niloticus* L.). Scientific Research and Essays. In Press.
- Avnimelech, Y. (2006). Bio-filters: The need for a new comprehensive approach. *Aquacultural engineering* 34:172-178.
- Avnimelech, Y. (2003). Control of microbial activity in aquaculture systems: active suspension ponds. *World Aquaculture* 34(4):19-21.
- Avnimelech, Y. (1999). Carbon/nitrogen ratio as a control element in aquaculture systems. *Aquaculture* 176: 227-235.

- Brazil, B. L. (2006). Performance and operation of a rotating biological contactor in a tilapia recirculating aquaculture system. *Aquacultural engineering* 34:261-274.
- Boyd, C. (1998) Pond water aeration systems. *Aquacultural Engineering* 18:9-40.
- Brander, K. M. (2007). Global fish production and climate change. *Proceedings of the National Academy of Sciences* 104 (50): 19709-19714
- Brenan, L. & Owende, P. (2010). Biofuels from microalgae – A review of technologies for production, processing, and extractions of biofuels and co-products. *Renewable and Sustainable Energy Reviews* 14:557-577.
- Camargo, J. A., Alonso, A. & Salamanca, A. Nitrate toxicity to aquatic animals: a review for freshwater invertebrates. *Chemosphere* 58:1255-1267
- Chen, S. & Malone, R. F. (1991). Suspended solids control in recirculating aquacultural systems. Engineering aspects of intensive aquaculture, pp. 170-186 in Proceedings from the aquaculture symposium, Cornell University, Ithaca, New York, 1991.
- Chopin, T. (2003). Integrated aquaculture. *Canadian Geographic* 123(5):24
- Crab, R., Kochva, M., Verstraete, W. & Avnimelech, Y. (2009). Bio-flocs technology application in over-wintering of tilapia. *Aquacultural Engineering* 40:105-112.
- Cripps, S. J. & Bergheim, A. (2000). Solids management and removal for intensive land-based aquaculture production systems. *Aquacultural Engineering* 22:33-56.
- Diver, S. (2006). Aquaponics - Integration of hydroponics with aquaculture. Publication No. IP163. ATTRA. National Sustainable Agriculture Information Service.
- Duffie, J. A. & Beckman, W. A. (1992). Solar Engineering of thermal processes. 2th edition. John Wiley & Sons, Inc. USA.
- Drennan II, D. G., Hosler, K. C., Francis, M., Weaver, D., Aneshansley, E., Beckman, G., Johnson, C. H. & Cristina, C.M. (2006). Standardized evaluation and rating of biofilters II. Manufacturer's and user's perspective. *Aquacultural engineering* 34:403-416
- Durborow, R. M., Crosby, D. M. & Brunson, M. W. (1997a). Ammonia in fish ponds. SRAC Publication No. 463.
- Durborow, R. M., Crosby, D. F. & Brunson, M. W. (1997b). Nitrite in fish ponds. SRAC Publication No. 462.
- Eding, E. H., Kamstra, A., Verreth J. A.J., Huisman E. A. & Klapwijk, A. (2006). Design and operation of nitrifying trickling filters in recirculating aquaculture: A review. *Aquacultural engineering* 34:234-260.
- Ebeling, J. E., Timmons, M. B. & Bisogni, J. J. (2006) Engineering analysis of the stoichiometry of photoautotrophic, autotrophic, and heterotrophic removal of ammonia-nitrogen in aquaculture systems. *Aquaculture* 257:346-358.
- Eshchar, M., O., Lahav, N. Mozes, A. Peduel & B. Ron (2006). Intensive fish culture at high ammonium and low pH. *Aquaculture* 255:301-313.
- Food and Agriculture Organizations of the United Nations (FAO). (2010). The State of World Fisheries and Aquaculture (SOFIA), 2010. FAO Fisheries Department, Roma.
- Franck, S., von Bloh, W., Müller, C., Bondeau, A. & Sakschewski, B. (2011). Harvesting the sun: New estimations of the maximum population of planet earth. *Ecological Modelling* 222(12):2019-2026.
- Fuller, R. J. (2007). Solar heating systems for recirculation aquaculture. *Aquacultural Engineering* 36:250-260.
- Graber, A. & Junge, R. (2009). Aquaponic systems: Nutrient recycling from fish wastewater by vegetable production. *Desalination* 246:147-156.

- Gross, A., Boyd, C. E, Wood & C. W. (2000). Nitrogen transformations and balance in channel catfish ponds. *Aquacultural Engineering* 24:1-14.
- Gutierrez-Wing, M. T. & Malone, R. F. (2006). Biological filters in aquaculture: Trends and research directions for freshwater and marine applications. *Aquacultural Engineering* 34:163-171.
- Hannesson, R. (2003). Aquaculture and fisheries. *Marine policy* 27:196-178.
- Hargreaves, J. A. (2006). Photosynthetic suspended-growth systems in aquaculture. *Aquacultural engineering* 34:344-363
- Hargreaves, J. A. & Tucker, C. S. (2004). Managing ammonia in fish ponds. SRAC Publication No. 4603. USA.
- Hargreaves, J. A. (1999). Control of clay turbidity in ponds. SRAC Publication No. 460.
- Hargreaves, J. A. (1998). Nitrogen biogeochemistry of aquaculture ponds. *Aquaculture* 166:181-212.
- Hargreaves, J. & Brunson, M.(1996). Carbon dioxide in fish ponds. SRAC Publication No. 468.
- Hovanec, T. A. & DeLong, E. F. (1996). Comparative analysis of nitrifying bacteria associated with freshwater and marine aquaria. *Applied and environmental microbiology* 62(8):2888-2896.
- Jensen, F. B. (2003). Nitrite disrupts multiple physiological functions in aquatic animals. *Comparative Biochemistry and Physiology Part A* 135:9-24.
- Kawarazuka, N. (2010). The contribution of fish intake, aquaculture, and small-scale fisheries to improving nutrition: A literature review. The World Fish Center Working Paper No. 2106. The World Fish Center, Penang, Malaysia. 51 p.
- Koning, N. B. J., Van Ittersum, M. K., Becx, G. A., Van Boekel, M. A. J. S., Brandenburg, W. A., Van Den Broek, J. A., Goudriaan, J., Van Hofwegen, Jongeneel, G., R. A. Schiere, J. B. & Smies, M. (2008). Long-term global availability of food: continued abundance or new scarcity? *NJAS* 55(3):229-292.
- Kuo-Feng, T. & Kuo-Lin W. (2004). The ammonia removal cycle for a submerged biofilter used in a recirculating ell culture system. *Aquacultural Engineering* 31:17-30.
- Losordo, T. M., Masser, M. P. & Rakocy, J. E. (1999). Recirculating aquaculture tank production systems. A review of component options. SRAC publication No. 453. USA.
- Losordo, T. M., Masser, M. P. & Rakocy. (1998). Recirculating aquaculture tank production systems. An overview of critical considerations. SRAC publication No. 451. USA.
- Lukefar, S. (1999). Teaching international animal agriculture. *Journal of Animal Science* 77(11):3106-3113.
- Masser, M. P; Rakocy, J. & Losordo, T. M. (1999). Recirculating aquaculture tank production systems. Management of recirculating systems. Sourthern Regional Aquaculture Center, Publication No. 452. USA.
- Müller-Esterl, W. (2008). Bioquímica. Fundamentos para la medicina y ciencias de la visa. Editorial Reverté, S. A. España.
- Mohanty, B. P., Mohanty, S., Sahoo, J. K. & Sharma, A. P. (2010). Climate change: impacts on fisheries and aquaculture. In Simard, S. (ed.) Climate change and variability InTech Publishing ISBN 978-953-307-144-2
- Naylor, R. L., Hardy, R.W., Bureau, D.P., Chiu, A., Elliot, M., Farrell, A. P., Forster, I., Gatlin, D.M., Goldberg, R.J., Hua, K. & Nichols, P. D. (2009). Feeding aquaculture in an era of finite resources. *Proceedings of the National Academy of Sciences* 106 (36):15103-15110.

- Naylor, R. L., Goldburg, R.J., Primavera, J. H., Kautsky, N., Beveridge, M. C. M., Clay, J., Folke, C., Lubchenco, J., Mooney, H., & Troell, M. (2000). Effect of aquaculture on world fish supplies. *Nature* 405(29):1017-1024.
- Oca, J., & Masaló, I. (2007). Design criteria for rotating flow cells in rectangular aquaculture tanks. *Aquacultural Engineering* 36:36-44.
- Olvera-Olvera, C., Olvera-González, J. E., Mendosa-Jasso, J., Peniche-Vera, R., Castañeda-Miranda, R. & Herrera-Ruiz, G. (2009). Feed dosage and ammonium control device base on C/N ratio for a zero-discharge system. *International Journal of Agriculture & Biology* 11(2):173-177.
- Panjaitan, P. (2010). Shrimp culture of *Penaeus monodon* with zero water exchange model (ZWEM) using molasses. *Journal of Coastal Development* 14(1):35-44
- Rakocy, J. E., & Hargreaves, J. A. (1993). Integration of vegetable hydroponics with fish culture: A review. In Wang, Jaw-Kai. Techniques of modern aquaculture. Proceedings of an Aquacultural Engineering Conference. 21 - 23 June, 1993. Spokane, Washington .pp. 112-136. Published by American Society of Agricultural Engineerings.
- Rico-García, E.; Casanova-Villareal, V. E., Mercado-Luna, A., Soto-Zarazúa, G. M., Guevara-González, R. G., Herrera-Ruiz, G., Torres-Pacheco, I., Velázquez-Ocampo, R. V. (2009). Nitrate content on summer lettuce production using fish culture water. *Trends in Agriculture Economics* 2(1):1-9.
- Ridha, M. T. & Cruz, E. M. (2001). Effect of biofilter media on water quality and biological performance of the Nile tilapia *Oreochromis niloticus* L. reared in a simple recirculating system. *Aquacultural Engineering* 24:157-166
- Rumei, W., Setian, F., Guo, T., Lizhong, F. & Xiaoshuan (2003). Evaluation of the aquaculture pond water quality. ASAE Annual International Meeting. Nevada, USA. Paper ASAE 031298.
- Savidov, N. A., Hutchings, E., Rakocy, J. E. (2005). Fish and plant production in a recirculating aquaponic system: A new approach to sustainable agriculture in Canada. *Acta horticulturae* 742:209-222.
- Seginer, I. & Halachmi, I.(2008). Water heating to enhance fish growth. Is it justified? Proceedings AgEng2008 (CD), Paper OP-630 (1131237).
- Summerfelt, S. T. & Sharrer, M. J. (2004). Design implication of carbon dioxide production within biofilters contained in recirculating salmonid culture systems. *Aquacultural engineering* 32:171-182.
- Summerfelt, S. T., Vinci, B. J. & Piedrahita, R. H. (2000). Oxygenation and carbon dioxide control in water reuse systems. *Aquacultural engineering* 22:87-108.
- Summerfelt, S. T., Timmons, M. B., Watten, B. J. (1998). Culture tank designs to increase profitability. In Libey, G. S. and Timmons, M. B. (1998) Proceedings of the Second International Conference on Recirculating Aquaculture. July 16-19, Roanoke, Virginia.
- Timmons, M. B.; Holder, J. L., Ebeling, J. B. (2006). Application of microbead biological filters. *Aquacultural engineering* 34:332-343.
- Timmons, M. B.; Ebeling, J. M., Wheaton, F. W., Summerfelt, S. T., Vinci, B. J. (2002). Recirculating aquaculture systems. Cayuga Aqua Ventures Inc. 2th. Edition. USA.
- Timmons, M. B.; Summerfelt, S. T., Vinci, B. J. (1998). Review of circular tank technology and management. *Aquacultural Engineering* 18:51-69.

- Soto-Zarazúa, M. G., Rico-García, E. & Toledano-Ayala, M. (2011). International Journal of the Physical Sciences 6(5):1039-1044.
- Soto-Zarazúa, M. G., Herrera-Ruiz, G., Rico-García, E., Toledano-Ayala, M., Peniche-Vera, R., Ocampo-Velázquez, R. & Guevara-González, R. (2010). Development of efficient recirculation system for Tilapia (*Oreochromis niloticus*) culture using low cost materials. *African Journal of Biotechnology* 9(32):5203-5211.
- Tucker, C. (2005). Pond aeration. Southern Regional Aquaculture Center, Publication No. 3700. USA.
- Tyson, R. V., Simonne, E. H., Treadwell, D. D., Davis, M. & White, J. M. (2008a). Effect of water pH on yield and nutritional status of greenhouse cucumber grown in recirculating hydroponics. *Journal of plant nutrition* 31(11):2018-2030.
- Tyson, R. V., Simonne, E. H., Treadwell, D. D., White, J.M. & Simone, A. (2008b). Reconciling pH for ammonia biofiltration and cucumber yield in a recirculating aquaponic system with perlite biofilters. *HortScience* 43(3):719-724.
- Van Gorder, S. (2003). Small-scale aquaculture and aquaponics. The new and the nostalgic. *Aquaponics Journal* 7(3):14-17.
- Van Rijn, J., Tal, J. & Schreier, H. J. (2006). Denitrification in recirculating systems: Theory and applications. *Aquacultural Engineering* 34:364-376.
- Vinatea, L. & Carvalho, J. W. Influence of water salinity on the SOTR of paddlewheel and propeller-aspirator-pump aerators, its relation to the number of aerators per hectare and electricity costs. *Aquacultural engineering* 37 (2): 73-78.
- Wheaton, F. W. (1982). Acuicultura. Diseño y construcción de sistemas. AGT Editor, S. A. México, D. F.
- Wurts, W. A. & Durborow, R. M. (1992). Interactions of pH, carbon dioxide, alkalinity and hardness in fish ponds. Southern Regional Aquaculture Center, Publication No. 464. USA.
- Yi, Y. & Kwei-Lin, C. & Diana, J. S. (2003). Techniques to mitigate clay turbidity problems in fertilized earthen fish ponds. *Aquacultural Engineering* 27:39-51.

IntechOpen

Aquaculture and the Environment - A Shared Destiny

Edited by Dr. Barbara Sladonja

ISBN 978-953-307-749-9

Hard cover, 246 pages

Publisher InTech

Published online 22, December, 2011

Published in print edition December, 2011

Aquaculture is the art, science and business of cultivating aquatic animals and plants in fresh or marine waters. It is the extension of fishing, resulted from the fact that harvests of wild sources of fish and other aquatic species cannot keep up with the increased demand of a growing human population. Expansion of aquaculture can result with less care for the environment. The first pre-requisite to sustainable aquaculture is clean water, but bad management of aquatic species production can alter or even destroy existing wild habitat, increase local pollution levels or negatively impact local species. Aquatic managers are aware of this and together with scientists are looking for modern and more effective solutions to many issues regarding fish farming. This book presents recent research results on the interaction between aquaculture and environment, and includes several case studies all over the world with the aim of improving and performing sustainable aquaculture.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Oscar Alatorre-Jácome, Fernando García-Trejo, Enrique Rico-García and Genaro M. Soto-Zarazúa (2011). Aquaculture Water Quality for Small-Scale Producers, Aquaculture and the Environment - A Shared Destiny, Dr. Barbara Sladonja (Ed.), ISBN: 978-953-307-749-9, InTech, Available from:
<http://www.intechopen.com/books/aquaculture-and-the-environment-a-shared-destiny/aquaculture-water-quality-for-small-scale-producers>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2011 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](https://creativecommons.org/licenses/by/3.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen