

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

186,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Up-to-Date Insight About Membrane Remodeling as a Mechanism of Action for Ethanol-Induced Liver Toxicity

Odile Sergent, Fatiha Djoudi-Aliche and Dominique Lagadic-Gossmann
*EA 4427 SeRAIC/IRSET, Université de Rennes 1, IFR 140, UFR des Sciences
Pharmaceutiques et Biologiques, 2, av Pr Léon Bernard, 35043 Rennes cédex,
France*

1. Introduction

Hepatocellular death is a key mechanism in alcoholic liver diseases. Although ethanol has been described for many years as capable of increasing membrane fluidity, it is only recently that this fluidizing effect has been reported to be involved in ethanol-induced liver toxicity. In addition, in the last decade, a better understanding of plasma membrane has led to suggest that this membrane is not a random association of lipids, but is rather heterogeneous, with various microstructures enriched in specific components depending on their affinity. Special attention has been paid on lipid rafts that are cholesterol- and sphingolipid- rich microstructures, conferring them higher rigidity compared to other plasma membrane microdomains. As lipid rafts can also activate or suppress cell signaling pathways, lipid raft discovery provides new arguments for several researchers to revisit the fluidizing effect of ethanol by studying the possible ethanol-induced physical and biochemical alteration of lipid rafts. Thus, in this chapter, we have considered to review the capacity of ethanol to induce a membrane remodeling, depicted as an increase in membrane fluidity and alterations of physical and biochemical properties of lipid rafts, and its relationship with ethanol liver toxicity.

2. Membrane fluidity

The Singer-Nicolson fluid mosaic model indicates that membranes consist of a phospholipid bilayer, where lipids, in a fluid phase, act as solvent for proteins (Singer & Nicolson, 1972). In this chapter, membrane fluidity means the relative freedom of motion for membrane components, especially phospholipids, and represents the combination of various types of mobility (Figure 1). Membrane fluidity is principally determined by the acyl chain swinging movement and phospholipid rotation. Thus, short chains and double bonds in acyl chains of phospholipids create spaces in the bilayer and promote membrane fluidization. At the opposite, the rigid steroid nucleus of cholesterol, lying next to the first 9 or 10 carbon atoms of the phospholipid acyl chains, prevents the swinging movement of the acyl chains thereby stiffening membranes. For the evaluation of this membrane parameter, most studies have used either electron paramagnetic resonance (EPR) with spin-labeled fatty acids, or

polarization of fluorescence with hydrophobic fluorescence polarization probes. An increased membrane fluidity for EPR is usually assessed by a decrease of order parameter (S), and for fluorescence, by a decrease of polarization (P), anisotropy (A) or microviscosity (η).




Fig. 1. Different types of mobility of phospholipids.

Any alteration of the optimal range for membrane fluidity has influence on many biological functions such as membrane enzyme and receptor activities, or transmembrane transport processes (Ho et al, 1994; Schachter, 1984; Stubbs et al, 1988). Furthermore, more recently, it was also shown its fundamental role in cell signalling responses to xenobiotic stress (polycyclic aromatic hydrocarbons, cisplatin or ethanol), leading to cell death such as apoptosis (Rebillard et al, 2007; Sergent et al, 2005; Tekpli et al, 2011).

2.1 Plasma liver membranes

Since the end of the seventies, many papers have provided strong evidence that ethanol very rapidly induces a fluidization of membranes as reported by several reviews (Goldstein, 1987; Rottenberg, 1992; Wood & Schroeder, 1988).

2.1.1 Tissue type-dependent effect of ethanol

Using electron paramagnetic resonance, Chin and Goldstein (1977a) were the first to demonstrate the ability of ethanol used at low concentrations (from 20 mM - 40 mM) to increase *in vitro* membrane fluidity of erythrocyte and synaptosomal plasma membranes. In addition, they showed that continuous exposure of mice to ethanol provided in the diet for a short period (8 days) (Chin & Goldstein, 1977b) or by inhalation (3 days) (Lyon & Goldstein, 1983) respectively restored a membrane fluidity near controls or even rigidified membranes in the inner hydrophobic regions, testifying an adaptation. Thus, in alcoholic patients, erythrocytes exhibited a decrease in membrane fluidity (Beaugé et al, 1985; Parmahamsa et al, 2004). However, the effect of ethanol on plasma membranes is different for the liver. Indeed, they become more fluid, mainly in the inner hydrophobic regions, for chronically ethanol-intoxicated rats (Schüller et al, 1984; Yamada & Lieber, 1984) and an increase in fluidity was also observed in plasma membranes isolated from Reuber H35 rat hepatoma

cells (Polokoff et al, 1985) or WRL-68 human hepatic cells (Gutierrez-Ruiz et al, 1995) following a long term treatment with ethanol (3 or 4 weeks). Such an effect could contribute to the special sensitivity of liver to ethanol toxicity. At the opposite, other organelles in the liver did not exhibit any membrane fluidification after ethanol intoxication of rats (Table 1). It should be noted that, when primary hepatocytes isolated from chronically ethanol-treated rats were cultured before the evaluation of plasma membrane fluidity by fluorescence polarization, an increased ordering was observed (Benedetti et al, 1991).

Organelles	Methods	Type of Lipid Bilayer Environment	Membrane Fluidity (compared to controls)	References
Mitochondria	Electron Paramagnetic Resonance	Polar region	=	Waring et al, 1981
	Fluorescence Polarization	Polar region	↓	Castro et al, 1991
	Fluorescence Polarization	Polar region	↓	Colell et al, 1997
Microsomes	Electron Paramagnetic Resonance	Polar region	↓	Ponnappa et al, 1982
	Electron Paramagnetic Resonance	Apolar core region	=	Taraschi et al, 1985
	Electron Paramagnetic Resonance	Apolar core region	=	Aloia et al, 1985
Plasma membranes	Fluorescence Polarization	Apolar core region	↑	Schüller et al, 1984
	Fluorescence Polarization	Apolar core region	↑	Yamada and Lieber, 1984


Table 1. Effect of chronic ethanol intoxication on membrane fluidity of various organelles in the liver. (In all experiments, rats were fed a diet containing 36 % of total calories as ethanol for 30 to 40 days.)

Whatever exposure modes (ingestion, inhalation or intraperitoneal injections) (Chin & Golstein, 1977b; Lyon & Golstein, 1983; Johnson et al, 1979), erythrocyte and synaptosomal plasma membranes isolated from ethanol-treated mice did not exhibit an increase in

membrane fluidity after a further *in vitro* ethanol addition in contrast to membranes isolated from untreated mice. Such an *in vitro* resistance was also observed in erythrocyte membranes from alcoholic patients (Beaugé et al, 1985). Even though liver plasma membranes remained more fluid after chronic rat intoxication (Schüller et al, 1984; Yamada & Lieber, 1984) or after long term ethanol treatment of cultured hepatocytes (Gutierrez-Ruiz et al, 1995; Polokoff et al, 1985), these isolated membranes also exhibited an *in vitro* resistance to the disordering effect of a further direct addition of ethanol. Finally, most of the papers quoted in table 1 indicated such a process for microsomes or mitochondria. This phenomenon could be related to several changes in membrane lipid composition (Johnson et al, 1979) *ie* an increase in cholesterol within brain and liver cell membranes in rats (Chin et al, 1978) and in monkeys (Cunningham et al, 1983), an increased ratio of saturated to polyunsaturated fatty acids (Johnson et al, 1979), or reduced concentrations of sialic acid and galactose in the membrane surface of human erythrocytes (Beaugé et al, 1985).

2.1.2 Molecular mechanisms whereby ethanol could increase membrane fluidity

These mechanisms, summarized in figure 2, can occur simultaneously. The first described mechanism was in brain membranes and concerns physical properties of ethanol which allow it to directly interact with the lipid bilayer, thus triggering a direct membrane disorder (Goldstein, 1984; Gurtovenko & Anwar, 2009; Marquês et al, 2011; Rottenberg, 1992). This theory was particularly developed in the field of drug tolerance and physical dependence, but, in the liver, other mechanisms were also described. First, it was proposed that the fluidizing effect of chronic ethanol treatment could be related to changes in membrane lipid composition as acyl chain saturation and cholesterol are well-described to affect membrane fluidity. Thus, Yamada et al (1984) related the increase in membrane fluidity of liver plasma membranes after chronic ethanol feeding to a decrease in cholesterol plasma membrane content by an unknown mechanism. In hepatoma cells chronically exposed to ethanol for 3 weeks, the increase in membrane fluidity of plasma membranes was linked to the elevation of the ratio phosphatidylcholine/sphingomyelin (Polokoff et al, 1985). However, the main distinction of liver is that most of the ethanol metabolism occurs in this organ. Thus, ethanol metabolism appeared to play a key role since blocking ethanol metabolism by methylpyrazole inhibited changes in membrane fluidity both in acute intoxicated primary rat hepatocytes (Sergent et al, 2005), and in chronically treated hepatoma cells (Polokoff et al, 1985). Logically, as ethanol metabolism was involved, our team was interested in looking at the involvement of oxidative stress following an acute ethanol intoxication of primary rat hepatocytes. Using antioxidant such as thiourea (reactive oxygen species (ROS) scavenger) or vitamin E (lipid peroxidation inhibitor), we showed that oxidative stress played a role in the fluidizing effect of ethanol (Sergent et al, 2005). This new mechanism explained how ethanol could very rapidly (30 minutes) increase membrane fluidity since ROS production could be detected as soon as 15 minutes. Several molecular mechanisms can be proposed to explain the influence of oxidative stress on membrane fluidity. First, lipid peroxidation by-products could increase membrane fluidity either by interacting with membrane proteins (Buko et al, 1996; Subramaniam et al, 1997), or more directly by their own rearrangement (Jain et al, 1994; Gabbita et al, 1998). ROS, by oxidizing tubulin could also disrupt microtubule cytoskeleton, thereby increasing membrane fluidity (Yoon et al, 1998; Remy-Kristensen et al, 2000). In our model of primary rat hepatocytes, paclitaxel (a microtubule stabilizer) prevented from the fluidizing effect of ethanol (unpublished data).



(ROS : reactive oxygen species).

Fig. 2. Possible molecular mechanisms for ethanol to increase membrane fluidity.


2.2 Liver mitochondria membranes

As shown above, a great body of evidence indicated that, in inner membranes of mitochondria, ethanol intoxication induced a decrease rather than an increase in fluidity. This was demonstrated for chronically intoxicated rats but also with HepG2 human hepatocytes treated with acetaldehyde, a product of ethanol metabolism (Lluis et al, 2003), providing a further proof of the involvement of ethanol metabolism in membrane fluidity changes. In addition, this decrease was related to an elevation of cholesterol content in mitochondria which concerns both outer and inner membranes. Finally, the acetaldehyde stimulation of cholesterol incorporation into mitochondria membranes was attributed to endoplasmic reticulum stress.

2.3 Membrane pharmacology of ethanol liver toxicity by manipulation of membrane fluidity

Since the eighties, many studies suggested the influence of ethanol fluidizing effect on membrane protein activities (McCall et al, 1989; Mills et al, 1985; Rubin & Rottenberg, 1982). Only recently, researchers became interested in determining the role of membrane fluidity changes in ethanol-induced hepatocellular death. Thus, manipulation of plasma membrane fluidity by exposing primary rat hepatocytes to membrane stabilizing agents (ursodeoxycholic acid (UDCA) or ganglioside GM1 (GM1)) led to the inhibition of ethanol-induced cell death, while fluidizing compounds (tween 20 or A₂C) enhanced it (Sergent et al, 2005). In order to explain how plasma membrane fluidity could affect cell death, oxidative stress was also studied. At the opposite of fluidizing compounds, membrane stabilizing agents were shown to protect from ethanol-induced lipid peroxidation, ROS production and the elevation of another prooxidant factor, namely low-molecular-weight iron. Low-molecular-weight iron consists of iron species that can trigger oxidative stress by catalyzing the formation of a highly reactive free radical, the hydroxyl radical. It should be noted that UDCA and GM1 displayed a protection towards ethanol-induced ROS production only when ROS were evaluated after 1 or 5 hours of incubation with ethanol. At 15 minutes, no protection was afforded by membrane stabilizing agents, unlike the inhibitor

of ethanol metabolism, 4-methyl-pyrazole. This led us to postulate a sequence of events whereby the early ROS formation was mainly due to ethanol metabolism and the late phase to the increase in membrane fluidity (Figure 3). Interestingly, the increased mitochondrial membrane ordering was also associated with the development of oxidative stress. Indeed, stabilizing agents such as S-adenosyl-L methionine (SAME) or taurine conjugate of UDCA (tauroursodeoxycholic acid) protected from glutathione depletion in mitochondria obtained from the liver of rats chronically fed with ethanol (Colell et al, 1997; Colell et al, 2001). Reduced glutathione, the main non protein thiol in cells, plays an important role to detoxify hydrogen peroxide and other organic peroxides in mitochondria. Glutathione depletion in mitochondria made them more sensitive to ROS production and subsequent oxidative stress. Thus, it was demonstrated that the increased mitochondrial membrane microviscosity impaired the glutathione transporter which normally allows the glutathione transport from cytosol to mitochondrial matrix (Coll, 2003; Lluís et al, 2003) (Figure 3).



(GSH : reduced glutathione; ROS : reactive oxygen species).

Fig. 3. Relationship between membrane fluidity and ethanol-induced oxidative stress.

Cholesterol involvement in this process should be pointed out. Indeed, as mitochondrial membrane enrichment in cholesterol was responsible for the decreased mitochondrial membrane fluidity, lovastatin, an inhibitor of hydroxymethylglutaryl coenzyme A involved in cholesterol synthesis, was able to protect hepatocytes from acetaldehyde sensitization to tumor necrosis factor (TNF) α (Lluís et al, 2003). Similarly to membrane stabilizing agents, membrane fluidizer (A₂C) restored the initial glutathione transport rate and mitochondrial content (Coll et al, 2003; Lluís et al, 2003). However, the use of membrane fluidizers should be done with caution since, from our results about the involvement of plasma membrane fluidization in ethanol-induced cell death, it appears that they can be injurious for hepatocytes. At the opposite, UDCA and its conjugates seem to be good candidates for a potential therapeutic use, because, due to their membrane stabilizing properties (Güldütuna et al, 1993), they restore the normality in membrane fluidity for every type of membranes. Thus, in case of ethanol intoxication, they were able to prevent both the increase of plasma membrane fluidity, as we observed in primary rat hepatocytes (Sergent, 2005), and the decrease in mitochondria membranes of hepatocytes from ethanol-fed rats (Colell et al,

2001). In addition, UDCA was also shown to protect rats from the increase in liver plasma membrane fluidity due to chronic ethanol intake and hence from liver lipid peroxidation and necrosis (Oliva et al, 1998). However, although UDCA is a therapeutically relevant bile acid, already used for preventing human primary biliary cirrhosis (Poupon et al, 2003; Corpechot et al, 2011), it did not exhibit any beneficial effect on a 6-month survival of patients with severe alcohol-induced cirrhosis, but possibly because of inappropriate dosage (Pelletier et al, 2003).

3. Lipid rafts

Because of the well-described effect of ethanol on plasma membrane fluidity, it is not surprising that some researchers about alcoholic liver diseases were interested in the possible involvement of lipid rafts in ethanol toxicity. Indeed, plasma membrane is not constituted by a random lipid distribution but rather by a selective lateral lipid segregation due to self-associative properties of sphingolipid and cholesterol, leading to the concept of "lipid rafts" (Simons & Toomre, 2000; Lingwood & Simons, 2010). Thus, lipid rafts are detergent-resistant, sphingolipid- and cholesterol-rich microdomains of the plasma membrane, which form highly ordered spatial nanoscale assemblies separated from other membrane regions composed of more unsaturated and loosely packed fatty acids (Figure 4).




Fig. 4. Schematic representation of a lipid raft (without proteins).


Lipid rafts as nanoscale assemblies are dynamics and after cell stimulation, can coalesced to larger levels to form raft platforms (Harder & Engelhardt, 2004). Concerning proteins, lipid rafts are notably enriched in glycosylphosphatidylinositol (GPI) proteins, receptors such as cell death receptors and Toll-like receptors (TLR), and signaling proteins like mitogen-activating protein kinases, protein kinases C etc. Some proteins are raft residents, whereas others are recruited after cell stimulation with receptor-specific ligands. In addition, based on their mobility, lipid rafts, through their aggregation, can form platforms that assembly many proteins on a same place leading to the formation of a receptor cluster, which can then activate or suppress signaling pathways (Pike, 2003; Schmitz & Orso, 2002). One might suppose that ethanol, through its capacity to increase liver plasma membrane fluidity, can disturb these microdomains and hence, various cell signaling pathways. Consequently, in the last past decade, new investigations were undertaken to possibly link lipid rafts to ethanol toxicity. Researches were conducted in two directions: the main one concerned perturbation of innate immunity *via* TLR4 signaling and the other one, hepatocyte cell death *via* the activation of phospholipase C (PLC) signaling.

3.1 Lipid rafts in the TLR4 signaling dysfunction by ethanol

Several components of innate immunity contribute to the pathogenesis of alcoholic liver disease (Gao et al, 2011). Here, we will mainly focus on lipopolysaccharide (LPS)/TLR4 signaling pathways because of the necessary translocation of TLR4 receptor into lipid rafts for its activation.

3.1.1 Involvement of LPS/TLR4 signaling pathway in alcoholic liver disease

Strong evidence suggest that the immune cells of the liver (phagocytic cells such as neutrophils or resident Küpffer cells, and lymphocytes such as natural killer [NK] cells or T cells) play a crucial role in alcoholic liver disease including steatosis, hepatitis and fibrosis (Suh & Jeong, 2011). Thus, Küpffer cells are main actors in the immune response against endotoxin/lipopolysaccharide (LPS) *via* Toll-Like Receptor type 4 (TLR4) signaling pathway leading to the production of pro-inflammatory mediators such as cytokines (TNF- α , interleukin [IL]-1, IL-6), chemokines (monocyte chemotactic protein-1 [MCP-1]), ROS and profibrogenic factors (transforming growth factor [TGF]- β , platelet-derived growth factor [PDGF]), which subsequently activate hepatic stellate cells for the production of extracellular matrix (Jeong & Gao, 2008) (Gao et al, 2011) (Figure 5). Indeed, it is well established that ethanol intake, by increasing gut permeabilization, allows the uptake of LPS in portal circulation (Parlesak et al, 2000) promoting liver ethanol toxicity (Nanji et al, 1994). In addition, in the liver, TLR4 is also expressed on recruited macrophages, hepatocytes, sinusoidal endothelial cells and hepatic stellate cells (Seki & Brenner, 2008). Consequently, *via* TLR4 signalling, these last cells can also contribute to liver inflammation by releasing proinflammatory cytokines and chemokines. Finally, TLR4 signalling in hepatic stellate cells can also participate to the development of alcoholic fibrosis by enhancing TGF- β signalling (Seki et al, 2007). Therefore, TLR4 receptor appeared crucial in the development of alcoholic liver disease (Gao et al, 2011).



(HEP : hepatocytes; HSC: hepatic stellate cells; KC : Küpffer cells; IL : interleukin; LPS : lipopolysaccharide; MCP-1 : monocyte chemotactic protein-1 ; PDGF : platelet-derived growth factor; TGF: tumor growth factor; TLR4 : Toll-like receptor 4; TNF : tumor necrosis factor)

Fig. 5. Contribution of TLR4 receptor to the pathogenesis of alcoholic liver disease.

3.1.2 Effects of ethanol on the recruitment of TLR4 into lipid rafts

LPS does not bind TLR4 receptor directly, but is rather first bound to cell surface co-receptors, the cluster of differentiation 14 (CD14) and the myeloid differentiation protein 2 (MD-2), without cytoplasmic domains (Fitzgerald, 2004). However, TLR4 is the integrator of cell signalling since it has intracellular signaling domains. Close interactions between these membrane receptors are made possible by their recruitment and assembly within lipid rafts (Schmitz & Orso, 2002; Triantafilou et al, 2002). Thus, CD14 is a glycosyl phosphatidylinositol-linked protein which therefore constitutively resides in lipid rafts, while TLR4 needs translocation into rafts for the complex formation (Dolganiuc et al, 2006). Two features of the ethanol effect on TLR4 and other receptor signaling could be distinguished depending on ethanol concentration. 1) At high concentration (≥ 50 mM), ethanol prevented from LPS-induced redistribution pattern of the co-receptor CD14 within lipid rafts, and from the translocation of TLR4 receptor into rafts (Table 2). This alteration could partly explain why ethanol consumption is recognized as a risk factor for concomitant bacterial or viral infections (Nelson and Kolls, 2002; Szabo, 1999). Dai et al (2005) and Dolganiuc et al (2006) suggested that ethanol, at the concentration of 50 or 86 mM, may disrupt lipid rafts because similar effects were obtained with lipid raft disrupters. However, a protein raft marker, flotillin did not exhibit any alteration and no clear evidence of lipid raft disruption was given, since the cholesterol decrease was detected in culture media instead of lipid rafts. They also attributed changes in partitioning cellular membrane in raft and nonraft structures to the increase in bulk membrane fluidity (Dolganiuc et al, 2006) without checking this influence by the use of membrane stabilizing agents or measuring the increase in membrane fluidity directly in lipid rafts. Their hypothesis would be that ethanol by this way could disrupt lipid protein interactions (Szabo et al, 2007). Only at very high concentrations (200 mM), a lipid raft disruption was really observed in RAW 264.7 macrophages (Fernandez-Lizarbe et al, 2008). However, at 50 mM, in primary rat cortical astrocytes, a partial disruption of lipid raft could be detected suggesting that ethanol at this concentration induced both effects : i) disruption leading to the inhibition of lipid raft - induced cell signalling, and ii) promotion of TLR4 recruitment in lipid rafts (see below)) (Blanco et al, 2008). More recently, it was also demonstrated an ethanol inhibition of lipid raft-mediated T-Cell Receptor (TCR) signalling in human CD4+ T cells and in Jurkat T cells, but no alteration of lipid raft markers was observed suggesting that ethanol had no direct effect on lipid rafts (Ghare et al, 2011). Interestingly, the authors proposed a post-translational modification of proteins to explain the inhibition of protein translocation into lipid rafts. These mechanisms could also be explored for the other models. 2) At lower concentration (≤ 50 mM), mimicking LPS effects both in macrophages and astrocytes, ethanol induced the recruitment of TLR4 into lipid rafts, thus allowing the activation of TLR4 dependent cell signalling (Table 2). A similar process was also observed for IL-1R1 (IL1 receptor 1) (Blanco et al, 2008). Thus, ethanol triggered cytokine and other inflammatory mediator secretion *via* lipid raft-dependent signalling pathway. According to Blanco et al (2008), low ethanol concentrations (10 - 50 mM) may facilitate protein-protein and protein-lipid interactions within the membrane microdomains to promote receptor recruitment into the lipid rafts. Even if this effect has not yet been directly described in the liver, lipid rafts might participate to the mechanisms involved in the enhancement by chronic ethanol treatment of liver inflammation associated with the activation of IL-1R1 receptor in rat liver and hepatocytes (Valles et al, 2003), or TLR4 in immune cells (Szabo & Bala, 2010).

Ethanol Concentration	Cell Type	Ethanol effects on		References
		lipid raft biochemical properties	receptor signalling and inflammation	
86 mM	RAW macrophage like cell line	- Alteration of the redistribution pattern of CD14 in lipid rafts after LPS stimulation	- Decrease in LPS-induced TNF α production - Similar effect with nystatin, a lipid raft disrupter	Dai et al, 2005
50 mM	Chinese hamster ovary cell line (CHO) Primary human monocytes	- Alteration of the redistribution pattern of CD14 in lipid rafts after LPS stimulation - Decrease in LPS-induced TLR4 translocation into lipid rafts - Increase in residual TLR4 in the nonraft fractions - No changes in flotillin, a protein raft marker	- Decrease in LPS-induced TNF α production and NF-kB activation - Similar effects with methyl- β -cyclodextrin, a lipid raft disrupter	Dolganiuc et al, 2006
10 mM 50 mM 200 mM	RAW macrophage like cell line Primary mice peritoneal macrophages	-10 mM : CD14 and TLR4 translocation into lipid rafts -50 mM: CD14 translocation into lipid rafts, but at a lesser extent for TLR4 -10 and 50 mM : translocation of proteins associated with TLR4 response into lipid rafts -200 mM : No translocation of TLR4 into lipid rafts and disruption of lipid rafts	- Direct production of TNF α (10 mM : by 2.4 fold; 50 mM: by 1.8 fold; 100 mM : no effect)	Fernandez-Lizarbe et al, 2007

Table 2. (Continued)

Ethanol Concentration	Cell Type	Ethanol effects on		References
		lipid raft biochemical properties	receptor signalling and inflammation mediators	
10 mM 50 mM	Primary rat cortical astrocytes	- 10 mM: translocation of IL-1R1 and TLR4 into lipid rafts; recruitment of activated signalling proteins into rafts (P-IRAK and P-ERK) - 50 mM: idem, but at a lesser extent, with a slight disruption	- Expression of IL-1R1 and TLR4 receptors in cell lysates was abolished by lipid raft disrupters (saponin, or methyl- β -cyclodextrin) - Internalization of IL-1R1 receptor in caveosomes	Blanco et al, 2008
25 mM 75 mM	Primary human CD4+ T cells Jurkat T cell line	- Inhibition of PHA or anti-CD23/CD28 antibodies induced Lck, LAT or PLC γ recruitment into lipid rafts - No changes in lipid raft protein marker (flotillin)	Decrease in IL2 expression and down-regulation of TCR signaling	Ghare et al, 2011


(ERK : extracellular regulated kinase; IL : interleukin; IRAK : interleukin-1 receptor associated kinase; LAT : linker for activation of T cells; lck : lymphocyte-specific protein tyrosine kinase; LPS : lipopolysaccharide; NF-kB : nuclear factor kappa B; PHA : phytohaemagglutinin; PLC: phospholipase C; TCR : T cell receptor; TNF : tumor necrosis factor).

Table 2. Effects of acute ethanol exposure on lipid raft-mediated receptor activation. (In these studies, rafts were isolated by their *in vitro* property to resist to solubilization in non-ionic detergents at low temperature and to float and concentrate in low-density sucrose (Brown & Rose, 1992), leading to raft and non-raft fractions.)

3.2 Lipid rafts in ethanol-induced hepatocyte damage

Another approach was to consider the role of lipid rafts in ethanol-induced oxidative stress. The occurrence of oxidative stress in alcoholic liver disease and its relationship with ethanol liver damage have been extensively documented (Albano, 2008; Cederbaum et al, 2009; De Minicis & Brenner, 2008; Wu & Cederbaum, 2009), but less is known about the possible role of lipid rafts. Thus, it was shown by our team that lipid raft disrupters were able to protect from ethanol-induced ROS production and lipid peroxidation in primary rat hepatocytes (Nourissat et al, 2008). In addition, we have showed for the first time that oxidative changes within lipid rafts are a prerequisite for the oxidative stress to develop in rat hepatocytes.


Thus, ethanol metabolism, by producing a rapid and mild oxidative stress, was able to induce oxidative damage within lipid rafts leading to their clustering following protein crosslinkages (Figure 6).



(CHO : carbonyl group; ROS : reactive oxygen species; SH : thiol group)

Fig. 6. Ethanol-induced lipid raft clustering *via* oxidative stress and protein crosslinkage.

Protein crosslinkages were obtained by the formation of disulfide bridges from two intermolecular thiol (SH) groups from several rafts, and by the formation of adducts with malondialdehyde, a well-known product of lipid peroxidation in ethanol treated-rat hepatocytes (Nourissat et al, 2008). This aldehyde like 4-hydroxynonenal can react with nucleophile residues in proteins to form carbonyl groups which then may form Schiff base with a lysine of another protein. Such a protein can be included in another raft leading to raft clustering (Figure 6). Interestingly, according to experiments performed on the translocation of TLR4 (see above) which proposed a role for membrane fluidity without fully demonstrating it, we expressly proved the involvement of the fluidizing effect in the ethanol-induced lipid raft clustering by the use of membrane stabilizer or fluidizers. In addition, ethanol was shown to be able to fluidize lipid rafts, but at a lesser extent compared to bulk membranes. These results also confirmed our previous results which showed the pivotal role of the increased membrane fluidity in ethanol-induced cell death of rat hepatocytes (Sergent et al, 2005), thereby emphasizing on the contribution of membrane remodeling in ethanol liver toxicity. Finally, lipid raft clustering also participated to the activation of phospholipase C- γ -dependent signaling pathway. Indeed, this clustering induced translocation of phospholipase C- γ into rafts, which induced elevation of low-molecular-weight-iron, a potent prooxidant factor, and hence, lipid peroxidation. To summarize, ethanol metabolism, by producing a mild oxidative stress can rapidly affect both membrane fluidity and lipid rafts, thus promoting lipid raft aggregation (Figure 7). Then, this lipid raft clustering, by activating phospholipase C- γ dependent signaling pathway, may in turn trigger amplification of oxidative stress and cell death (Figure 7).



(LMW iron: low molecular weight iron; PLC: phospholipase C)

Fig. 7. Amplification of oxidative stress *via* lipid raft clustering during acute ethanol intoxication of rat hepatocytes.

In this context, new therapeutic approach, called membrane lipid therapy (Escriba et al, 2006), could be a very effective strategy to protect hepatocytes from membrane-dependent oxidative damage in alcoholic liver damage, especially as an increasing body of evidence indicated that some dietary compounds such as plant flavonoids (Tarahosky et al, 2008) or fatty fish long-chain polyunsaturated n-3 fatty acids (n-3 PUFAs) (Wassal & Stillwell, 2009) might modify physical and chemical properties of lipid rafts. Thus, n-3 PUFAs have been extensively described as efficient modifiers of lipid and protein composition of lipid rafts in many cell types such as T lymphocytes (Fan et al, 2004; Stulnig, 2001), Caco-2 cells (Duraismy et al, 2007), retinal vascular endothelial cells (Chen et al, 2007) and macrophages (Wong et al, 2009). In this context, the nutritional significance of lipid rafts has been recently pointed out (Yaqoob and Shaikh, 2010).

4. Conclusion

Taken altogether, these studies show that physical alterations of membranes (changes in membrane fluidity and microstructures) can be considered as an additional mechanism involved in ethanol liver toxicity. It is only in the last past decade that membrane remodeling appeared to be linked to ethanol liver toxicity (Figure 8). Therefore, further studies are needed in order to determine the role of lipid rafts in chronic ethanol intoxication, to further explore the downstream cell signaling after lipid raft clustering such as pathways involved in the elevation of low-molecular weight iron cell content, or to understand whether receptor recruitment in lipid raft might participate to alcoholic liver disease. In addition, other investigation should shed light on the possible beneficial effect of the modulation of membrane fluidity and lipid raft. Thus, statins that are already currently used in patients suffering from hypercholesterolemia, have demonstrated their efficiency to protect hepatocytes from acetaldehyde sentization to TNF (Lluis et al, 2003), and might also be proposed to disrupt lipid rafts. Finally, nutritional compounds such as plant flavonoids

or fatty fish long-chain polyunsaturated n-3 fatty acids might represent a new therapeutic approach for patients with alcoholic liver disease based upon modulation of the membrane structures.

1980'	<ul style="list-style-type: none"> ✓ Ethanol induces membrane fluidization ✓ Membrane adaptation in chronically ethanol intoxicated rats, except for plasma membranes.
1990'	<ul style="list-style-type: none"> ✓ Decreased membrane fluidity of liver mitochondria membranes and consequences on ethanol-induced oxidative stress ✓ First demonstration of the protection afforded by membrane stabilizers on <i>in vivo</i> ethanol liver toxicity
2000'	<ul style="list-style-type: none"> ✓ Demonstration of the involvement of membrane fluidization in the <i>in vitro</i> ethanol-induced oxidative damages of hepatocytes ✓ Various effects of ethanol on lipid rafts ✓ Involvement of lipid rafts in ethanol triggering of inflammation and oxidative stress

Fig. 8. Evolution of the "membrane remodelling" concept for alcoholic liver diseases.

5. Acknowledgement

The authors gratefully acknowledge IREB (Institut de Recherches Scientifiques sur les Boissons, Paris, France) for its financial support. They also wish to thank "Région Bretagne", which provided a grant for Fatiha Djoudi. The authors are also very grateful to Martine Chevanne for her skilfull technical assistance.

6. References

- Albano, E. (2008). Oxidative mechanisms in the pathogenesis of alcoholic liver disease. *Molecular Aspects of Medicine*, Vol.29, No.1-2, (February-April 2008), pp.9-16, ISSN 0098-2997.
- Aloia, R.C.; Paxton, J.; Daviau, J.S.; Van Gelb, O.; Mekusch, W.; Truppe; W.; Meyer, J.A. & Brauer, F.S. (1985). Effect of chronic alcohol consumption on rat brain microsome lipid composition membrane fluidity and Na⁺-K⁺-ATPase activity. *Life Sciences*, Vol.36, No. 10, (March 1985), pp.1003-1017, ISSN 0024-3205.
- Beaugé, F.; Stibler, H & Borg S. (1985). Abnormal fluidity and surface carbohydrate content of the erythrocyte membrane in alcoholic patients. *Alcoholism: Clinical and Experimental Research*, Vol.9, No.4, (July-August 1985), pp.322-326, ISSN 0145-6008.
- Benedetti, A.; Tangorra, A.; Baroni, G.S.; Ferretti, G.; Marucci, L.; Jezequel, A.M. & Orlandi, F. (1994). Plasma membrane order parameter in periportal and perivenular hepatocytes isolated from ethanol-treated rats. *American Journal of Physiology*, Vol.266, No.2Pt1, (February 1994), pp.G282-G291, ISSN 0193-1857.
- Blanco, A.M.; Perez-Arago, A.; Fernandez-Lizarbe, S. & Guerri, C. (2008). Ethanol mimics ligand-mediated activation and endocytosis of IL-1RI/TLR4 receptors via lipid rafts caveolae in astroglial cells. *Journal of Neurochemistry*, Vol.106, No.2, (July 2008), pp.625-639, ISSN 0022-3042.

- Brown, D.A. & Rose, J.K. (1992). Sorting of GPI-anchored proteins to glycolipid enriched membrane subdomains during transport to the apical cell surface. *Cell*, Vol.68, No.3, (February 1992), pp.533-544, ISSN 0092-8674.
- Buko, V.; Artsukevich, A.; Zavodnik, I.; Maltsev, A.; Suhko, L.; Zimmermann, T. & Dianzani, M.U. (1996). Interactions of malondialdehyde and 4-hydroxynonenal with rat liver plasma membranes and their effect on binding of prostaglandin E2 by specific receptors. *Free Radical Research*, Vol.25, No. 5, (November 1996), pp.415-420, ISSN 1071-5762.
- Castro, J.; Cortés, J.P. & Guzman, M. (1991). Properties of the mitochondrial membrane and carnitine palmitoyltransferase in the periportal and the perivenous zone of the liver. Effects of chronic ethanol feeding. *Biochemical Pharmacology*, Vol.41, No.12, (Jun 1991), pp.1987-1995, ISSN 0006-2952.
- Cederbaum, A.I.; Lu, Y. & Wu, D. (2009). Role of oxidative stress in alcohol-induced liver injury. *Archives in Toxicology*, Vol.83, No.6, (June 2009), pp.519-548, ISSN 0340-5761.
- Chen, W.; Jump, D.B.; Esselman, W.J. & Busik, J.V. (2007). Inhibition of cytokine signaling in human retinal endothelial cells through modification of caveolae/lipid rafts by docosahexaenoic acid. *Investigative Ophthalmology and Visual Science*, Vol.48, No.1, (January 2007), pp.18-26, ISSN 0146-0404.
- Chin, J.H. & Goldstein, D.B. (1977a). Effects of low concentrations of ethanol on the fluidity of spin-labeled erythrocyte and brain membranes. *Molecular Pharmacology*, Vol.13, No.3, (May 1977), pp.435-441, ISSN 0026-895x.
- Chin, J.H. & Goldstein, D.B. (1977b). Drug tolerance in biomembranes. *Science*, Vol.196, No. 4290, (May 1977), pp.684-685, ISSN 0036-8075.
- Chin, J.H.; Parson, L.M. & Goldstein D.B. (1978). Increased cholesterol content of erythrocyte and brain membranes in ethanol-tolerant mice. *Biochimica et Biophysica Acta*, Vol.513, No.3, (September 1978), pp.358-363, ISSN 0270-9139.
- Coll, O.; Colell, A.; Garcia-Ruiz, C.; Kaplowitz, N. & Fernandez-Checa, J.C. (2003). Sensitivity of the 2-oxoglutarate carrier to alcohol intake contributes to mitochondrial glutathione depletion. *Hepatology*, Vol. 38, No.3, (September 2003), pp.692-702, ISSN 0270-9139.
- Colell, A.; Garcia-Ruiz, C.; Morales, A.; Ballesta, A.; Ookhtens, M.; Rodés, J.; Kaplowitz, N. & Fernandez-Checa, J.C. (1997). Transport of reduced glutathione in hepatic mitochondria and mitoplasts from ethanol-treated rats: Effect of membrane physical properties and S-adenosyl-L-Methionine. *Hepatology*, Vol.26, No.3, (September 1997), pp.699-708, ISSN 0270-9139.
- Colell, A.; Coll.O.; Garcia-Ruiz, C.; Paris, R.; Tiribelli, C.; Kaplowitz, N.; Fernandez-Checa, J.C. Tauroursodeoxycholic acid protects hepatocytes from ethanol-fed rats against tumor necrosis factor-induced cell death by replenishing mitochondrial glutathione. *Hepatology*, Vol.34, No.5, (November 2001), pp.964-971, ISSN 0270-9139.
- Corpechot, C.; Chazouillères, O. & Poupon, R. (2011). Early primary biliary cirrhosis : biochemical response to treatment and prediction of long-term outcome. *Journal of Hepatology*, (2011), in press, ISSN 0168-8278.
- Cunningham, C.C.; Bottenus, R.E.; Spach, P.I & Rudel, L.L. (1983) Ethanol-induced changes in liver microsomes and mitochondria from the monkey, *Macaca fascicularis*. *Alcoholism : Clinical and Experimental Research*, Vol. 7, No. 4, (September 1983), pp.424-430, ISSN 0145-6008.


- Dai, Q.; Zhang, J. & Pruetz, S.B. (2005). Ethanol alters cellular activation and CD14 partitioning in lipid rafts. *Biochemical and Biophysical Research Communications*, Vol.332, No.1, (June 2005), pp. 37-42, ISSN 0006-291x.
- De Minicis, S. & Brenner, D.A. (2008). Oxidative stress in alcoholic liver disease : role of NADPH oxidase complex. *Journal of Gastroenterology and Hepatology*, Vol.23, Suppl.1, (2008), pp.S98-S103, ISSN 0815-9319.
- Dolganiuc, A.; Bakis, G.; Kodys, K.; Mandrekar, P. & Szabo, G. (2006). Acute ethanol treatment modulates toll-like receptor-4 association with lipid rafts. *Alcoholism : Clinical and Experimental Research*, Vol.30, No.1, (January 2006), pp.76-85, ISSN 0145-6008.
- Duraisamy, Y.; Lambert, D.; O'Neill, C.A. & Padfield, P.J. (2007). Differential incorporation of docosahexaenoic acid into distinct cholesterol-rich membrane raft domains. *Biochemical and Biophysical Research Communications*, Vol.360, No.4, (September 2007), pp.885-890, ISSN 0006-291x.
- Escriba, P.V. (2006). Membrane-lipid therapy : a new approach in molecular medicine. *Trends in Molecular Medicine*, Vol.12, No.1, (January 2006), pp.34-43, ISSN 1471-4914.
- Fan, Y.-Y.; Ly, L.H.; Barhoumi, R.; McMurray, D.N.; Chapkin, R.S. (2004). Dietary docosahexaenoic acid suppresses T cell protein kinase C theta lipid raft recruitment and IL-2 production. *Journal of Immunology*, Vol. 173, No.10, (November 2004), pp.6151-6160, ISSN 0022-1767.
- Fernandez-Lizarbe, S.; Pascual, M.; Gascon, M.S.; Blanco, A. & Guerri, C. (2008). Lipid rafts regulate ethanol-induced activation of TLR4 signaling in murine macrophages. *Molecular Immunology*, Vol.45, No.7, (April 2008), pp.2007-2016, ISSN 0161-5890.
- Fitzgerald, K.A.; Rowe, R.C. & Golenbock, D.T. (2004). Endotoxin recognition and signal transduction by the TLR4/MD2-complex. *Microbes and Infection*, Vol.6, No.15, (December 2004), pp.1361-1367, ISSN 1286-4579.
- Gabbita, S.P.; Subramaniam, R.; Allouch, F.; Carney, J.M. & Butterfield, D.A. (1998). Effects of mitochondrial respiratory stimulation on membrane lipids and proteins : an electron paramagnetic resonance investigation. *Biochimica et Biophysica Acta*, Vol.1372, No.2, (July 1998), pp.163-173, ISSN 0006-3002.
- Gao, B.; Deki, E.; Benner, D.A.; Frideman, S.; Cohen, J.I.; Nagy, L.; Szabo, G. & Zakhari, S. (2011). Innate immunity in alcoholic liver disease. *American Journal of Physiology*, Vol. 300, No.4, (April 2011), pp.G516-G525, ISSN 0193-1857.
- Ghare, S.; Patil, M.; Hote, P.; Suttles, J.; McClain, C.; Barve, S. & Joshi-Barve, S. (2011). Ethanol inhibits lipid raft-mediated TCR signaling and IL-2 expression: potential mechanism of alcohol-induced immune suppression. *Alcoholism : Clinical and Experimental Research*, Vol.35, No.8, (August 2011), pp.1-10, ISSN 0145-6008.
- Goldstein, D.B (1984). The effects of drugs on membrane fluidity. *Annual Review of Pharmacology and Toxicology*, Vol.24, (1984), pp.43-64, ISSN 0362-1642.
- Goldstein, D.B. (1987). Ethanol-induced adaptation in biological membranes. *Annals of the New York Academy of Sciences*, Vol.492, (April 1987), pp.103-111, ISSN 0077-8923.
- Güldütuna, S.; Zimmer, G., Imhof, M.; Bhatti, S.; You, T. & Leuschner, U. (1993). Molecular aspects of membrane stabilization by ursodeoxycholate. *Gastroenterology*, Vol.104, No.6, (June 1993), pp. 1736-1744, ISSN 0016-5085.
- Gurtovenko, A.A. & Anwar, J. (2009). Interaction of ethanol with biological membranes : the formation of non-bilayer structures within the membrane interior and their

- significance. *The Journal of Physical Chemistry B*, Vol.113, No. 7, (February 2009), pp.1983-1992, ISSN 1520-6106.
- Gutiérrez-Ruiz, M.C.; Gomez, J.L.; Souza, V. & Bucio, L. (1995). Chronic and acute ethanol treatment modifies fluidity and composition in plasma membranes of a human hepatic cell line (WRL-68). *Cell Biology and Toxicology*, Vol.11, No.2, (April 1995), pp.69-78, ISSN 0742-2091.
- Harder, T. & Engelhardt, K.R. (2004). Membrane microdomains in lymphocytes – from lipid rafts to protein scaffolds. *Traffic*, Vol.5, No.4, (April 2004), pp.265-275, ISSN 1600-0854.
- Ho, C.; Williams, B.W.; Kelly, M.B. & Stubbs, C.D. (1994) Chronic ethanol intoxication induces adaptative changes in the membrane protein/lipid interface. *Biochimica et Biophysica Acta*, Vol.1189, No.7, (January 1994), pp.135-142, ISSN 0006-3002.
- Jain, S.; Thomas, M.; Kumar, P. & Laloraya, M. (1994). Appearance of homogeneous smectic multilamellar microenvironments in biomembranes undergoing superoxide-initiated lipid peroxidation : lipid-dienyl radical accumulation and fluidity management in lipid bilayers. *Biochemistry and Molecular Biology International*, Vol.33, No.5, (August 1994), pp.853-862, ISSN 1039-9712.
- Jeong, W.I & Gao, B. (2008). Innate immunity and alcoholic liver fibrosis. *Journal of Gastroenterology and Hepatology*, Vol.23, No. Suppl.1, (2008), pp.S112-S118, ISSN 0815-9319.
- Johnson, D.A.; Lee, N.M.; Cooke R. & Loh, H.H. (1979). Ethanol-induced fluidization of brain lipid bilayers : required presence of cholesterol in membranes for the expression of tolerance. *Molecular Pharmacology*, Vol. 15, No., (1979), pp.739-746, ISSN 0026-895x.
- Lingwood, D. & Simons, K. (2010). Lipid rafts as a membrane-organizing principle. *Science*, Vol. 237, No., (January 2010), pp.46-50, ISSN 0036-8075.
- Lluis, J.M.; Colell, A.; Garcia-Ruiz, C.; Kaplowitz, N. & Fernandez-Checa, J.C. (2003). Acetaldehyde impairs mitochondrial glutathione transport in HepG2 cells through endoplasmic reticulum stress. *Gastroenterology*, Vol.124, No.3, (March 2003), pp.708-724, ISSN 0016-5085.
- Lyon, R.C. & Goldstein D.B. (1983). Changes in synaptic membrane order associated with chronic ethanol treatment in mice. *Molecular Pharmacology*, Vol.23, No.1, (January 1983), pp.86-91, ISSN 0026-895x.
- Marquês, J.T.; Viana, A.S. & De Almeida, R.F. (2011) Ethanol effects on binary and ternary supported lipid bilayers with gel/fluid domains and lipid rafts. *Biochimica and Biophysica Acta*, Vol.1808, No.1, (January 2011), pp.405-414, ISSN 00036-3002.
- McCall, D.; Henderson, G.I.; Gray, P. & Schenker, S. Ethanol effects on active Na⁺ and K⁺ transport in cultured fetal rat hepatocytes. *Biochemical Pharmacology*, Vol.38, No.16, (August 1989), pp.2593-2600, ISSN
- Mills, P.R.; Meier, P.J.; Boyer, J.L. & Gordon, E.R. The effect of ethanol and calcium on fluid state of plasma membranes of rat hepatocytes. *Alcohol*, Vol.2, No.1, (January-February 1985), pp.153-156, ISSN 0006-2952.
- Nanji, A.A.; Khettry, U. & Sadrzadeh, S.M. (1994). Lactobacillus feeding reduces endotoxemia and severity of experimental alcoholic liver disease. *Proceedings of the Society for Biology and Medicine*, Vol.205; No.3, (March 1994), pp.243-247, ISSN 0037-9727.

- Nelson, S. & Kolls, J.K. (2002). Alcohol, host defence and society. *Nature reviews. Immunology*, Vol.2, No.3, (March 2002), pp.205-209, ISSN 1471-1733.
- Nourissat, P.; Travert, M.; Chevanne M.; Tekpli, X.; Rebillard, A.; Le Moigne-Müller, G.; Rissel, M.; Cillard, J.; Dimanche-Boitrel, M.-T.; Lagadic-Gossmann, D. & Sergent, O. (2008). Ethanol induces oxidative stress in primary rat hepatocytes through the early involvement of lipid raft clustering. *Hepatology*, Vol.47, No.1, (January 2008), pp.59-70, ISSN 0270-9139.
- Oliva, L.; Beaugé, F.; Choquart, A.M.; Guitaoui, M. & Montet, J.C. (1998). Ursodeoxycholate alleviates alcoholic fatty liver damage in rats. *Alcoholism : Clinical and Experimental Research*, Vol.22, No.7, (October 1998), pp.1538-1543, ISSN 0145-6008.
- Parlesak, A.; Schäfer, C.; Schütz, T. Bode, J.C. & Bode C. (2000). Increased intestinal permeability to macromolecules and endotoxemia in patients with chronic alcohol abuse in different stages of alcohol-induced liver disease. *Journal of Hepatology*, Vol.32, No.5, (May 2000), pp.742-747, ISSN 0168-8278.
- Parmahamsa, M.; Reddy, K.R. & Varadacharyulu, N. (2004). Changes in composition and properties of erythrocyte membrane in chronic alcoholics. *Alcohol & Alcoholism*, Vol.39, No.2, (March-April 2004), pp.110-112, ISSN 0735-0414.
- Pelletier, G.; Roulot, D.; Davion, T.; Masliah, C.; Causse, X.; Oberti, F.; Raabe, J.J.; Van Lemmens, C.; Labadie, H. & Serfaty, L. A randomized controlled trial of ursodeoxycholic acid in patients with alcohol-induced cirrhosis and jaundice. *Hepatology*, Vol.37, No.4, (April 2003), pp.887-892, ISSN 0270-9139.
- Pike, L.J. (2003). Lipid rafts : bringing order to chaos. *Journal of Lipid Research*, Vol.44, No.4, (April 2003), pp.655-667, ISSN 0022-2275.
- Polokoff, M.A.; Simon, T.J.; Harris A.; Simon, F.R & Iwahashi. (1985). Chronic ethanol increases liver plasma membrane fluidity. *Biochemistry*, Vol.24, No.13, (June 1985), pp.3114-3120, ISSN 0006-2960.
- Ponnappa, B.C.; Waring, A.J.; Hoeck, J.B.; Rottenberg, H. & Rubin, E. (1982). Chronic ethanol ingestion increases calcium uptake and resistance to molecular disordering by ethanol in liver microsomes. *The Journal of Biological Chemistry*, Vol.257, No.17, (September 1982), pp.10141-10146, ISSN 0021-9258.
- Poupon, R.E.; Lindor, K.D.; Pares, A.; Chazouilleres, O.; Poupon, R. & Heathcote, E.J. (2003). Combined analysis of the effect of treatment with ursodeoxycholic acid on histologic progression in primary biliary cirrhosis. *Journal of Hepatology*, Vol. 39, No.1, (July 2003), pp.12-16, ISSN 0168-8278.
- Rebillard, A.; Tekpli, X.; Meurette, O.; Sergent, O.; LeMoigne-Muller, G.; Vernhet, L.; Gorria, M.; Chevanne, M.; Christmann, M.; Kaina, B.; Counillon, L.; Gulbins, E.; Lagadic-Gossmann, D. & Dimanche-Boitrel, M.-T. (2007). Cisplatin-induced apoptosis involves membrane fluidification via inhibition of NHE1 in human colon cancer cells. *Cancer Research*, Vol.67, No.16, (August 2007), pp.7865-7874, ISSN 0008-5472.
- Remy-Kristensen, A.; Duportail, G.; Coupin, G. & Kuhry, J.G. (2000). The influence of microtubule integrity on plasma membrane fluidity in L929 cells. *Molecular Membrane Biology*, Vol.17, No.2, (April-June 2000), pp.95-100, ISSN 0968-7688.
- Rottenberg, H. (1992). Liver cell membrane adaptation to chronic alcohol consumption, In : *Drug and Alcohol Abuse Reviews, Vol.2 : Liver Pathology and Alcohol*, R.R. Watson (Ed.), 91-115, The humana Press, ISBN 978-0-89603-206-4, Totowa, New Jersey, USA.

- Rubin, E. & Rottenberg, H. (1982). Ethanol-induced injury and adaptation in biological membranes. *Federation Proceedings*, Vol.41, No.8, (June 1982), pp. 2465-2471, ISSN 0014-9446.
- Schachter D. Fluidity and function of hepatocyte plasma membranes. (1984). *Hepatology*, Vol. 4, No.1, (January-February 1984), pp.140-151, ISSN 0270-9139.
- Schmitz, G. & Orso, E. (2002). CD14 signaling in lipid rafts : new ligands and co-receptors, *Current Opinion in Lipidology*, Vol.13, No.5, (October 2002), pp.513-521, ISSN 0957-9672.
- Schüller, A.; Moscat, J.; Diez, E.; Fernandez-Checa, C.; Gavilanes F.G. & Muncio, A.M. (1984). The fluidity of plasma membranes from ethanol-treated rat liver. *Molecular and Cellular Biochemistry*, Vol.64, No.1, (September 1984), pp.89-95, ISSN 0300-8177.
- Seki, E; De Minicis, S.; Osterreicher, C.H.; Kluwe, J.; Osawa, Y.; Brenner D.A. & Schwabe, R.F. (2007).TLR4 enhance TGF-beta signaling and hepatic fibrosis. *Nature Medicine*, Vol.13, No.11, (November 2007), pp.1324-1332, ISSN 1078-8956.
- Seki, E. & Brenner, D.A. (2008). Toll-like receptors and adaptator molecules in liver disease: update. *Hepatology*, Vol.48, No.1, (July 2008), pp.322-335, ISSN 0270-9139.
- Sergent, O.; Pereira, M.; Belhomme, C.; Chevane, M.; Huc, L. & Lagadic-Gossmann, D. (2005). Role for membrane fluidity in ethanol-induced oxidative stress in primary rat hepatocytes. *The Journal of Pharmacology and Experimental Therapeutics*, Vol. 313, No.1, (April 2005), pp.104-111, ISSN 0022-3665.
- Simons, K. & Toomre, D. Lipid rafts and signal transduction. *Nature Reviews. Molecular Cell Biology*, Vol.1, No.1, (October 2000), pp.31-39, ISSN 1471-0072.
- Singer, S.J. & Nicolson, G.L. (1972). The fluid mosaic model of the structure of cell membranes. *Science*, Vol.175, No.23, (February 1972), pp.720-731, ISSN 0036-8075.
- Stubbs, C.D.; Williams, D.B.; Pryor, C.L. & Rubin, E. (1988) Ethanol-induced modifications to membrane lipid structure : effect on phospholipase A₂ membrane interactions. *Archives of Biochemistry and Biophysics*, Vol. 262, No.2, (May 1988), pp.560-573, ISSN 0003-9861.
- Stulnig, T.M.; Huber, J.; Leitinger, N.; Imre, E.-M., Angelisova, P.; Nowotny, P. & Waldhausl W. (2001). Polyunsaturated eicosapentaenoic acid displaces proteins from membrane rafts by altering raft lipid composition. *Journal of Biological Chemistry*, Vol.276, No., pp.37335-37340.
- Suh, Y.-G. & Jeong, W.-I. (2011). Hepatic stellate cells and innate immunity in alcoholic liver disease. *World Journal of Gastroenterology*, Vol.17, No.20, (May 2011), pp.2543-2551, ISSN 1007-9327.
- Subramaniam, R.; Roediger, F.; Jordan, B.; Mattson, M.P.; Keller, J.N.; Waeg, G. & Butterfield, D.A. (1997). The lipid peroxidation product, 4-hydroxy-2-trans-nonenal, alters the conformation of cortical synaptosomal membrane proteins. *Journal of Neurochemistry*, Vol.69, No.3, (September1997), pp.1161-1169, ISSN 0022-3042.
- Szabo, G. (1999). Consequences of alcohol consumption on host defense. *Alcohol and Alcoholism*, Vol.34, No.6, (November-December 1999), pp.830-841, ISSN 0735-0414.
- Szabo, G.; Dolganiuc A.; Dai, Q. & Pruett, S.B. (2007). TLR4, ethanol and lipid rafts : a new mechanism of ethanol action with implications for other receptor-mediated effects. *Journal of Immunology*, Vol.178, No.3, (February 2007), pp.1243-1249, ISSN 0022-1767.

- Szabo, G. & Bala, S. (2010). Alcoholic liver disease and the gut-liver axis. *World Journal of Gastroenterology*, Vol.16, No.11, (March 2010), pp.1321-1329, ISSN 1007-9327.
- Tarahosky, Y.S.; Muzafarov, E.N. & Kim, Y.A. (2008). Rafts making and rafts braking : how plant flavonoids may control membrane heterogeneity. *Molecular and Cellular Biochemistry*, Vol.314, No.1-2, (July 2008), pp.65-71, ISSN 0300-8177.
- Taraschi, T.F.; Wu, A. & Rubin, E. (1985). Phospholipid spin probes measure the effects of ethanol on the molecular order of liver microsomes. *Biochemistry*, Vol.24, No., (1985), pp.7096-7101, ISSN 0021-2960.
- Tekpli, X.; Holme, J.A.; Sergent, O. & Lagadic-Gossmann, D. (2011). Importance of plasma membrane dynamics in chemical-induced carcinogenesis. *Recent Patents on Anti-Cancer Drug Discovery*, Vol.6, (2011), in press, ISSN 1574-8928.
- Triantafilou, M.; Miyake, K.; Golenbock, D.T. & Triantafilou, K. (2002). Mediators of innate immune recognition of bacteria concentrate in lipid rafts and facilitate lipopolysaccharide-induced cell activation. *Journal of Cell Science*, Vol.115, No.Pt12, (June 2002), pp. 2603-2611, ISSN 0021-9533.
- Valles, S.L.; Blanco, A.M.; Azorin, I.; Guasch, R.; Pascual, M.; Gomez-Lechon, M.J.; Renau-Piqueras, J. & Guerri, C. (2003). Chronic ethanol consumption enhances interleukin-1 mediated signal transduction in rat liver and in cultured hepatocytes. *Alcohol: Clinical and Experimental Research*, Vol.27, No.12, (December 2003), pp.1979-1986, ISSN 0145-6008.
- Waring, A.J.; Rottenberg, H.; Ohnishi, T. & Rubin, E. (1981). Membranes and phospholipids of liver mitochondria from chronic alcoholic rats are resistant to membrane disordering by ethanol. *Proceedings of the National Academy of Sciences of the United States of America*, Vol.78, No.4, (April 1981), pp.2582-2586, ISSN 0027-8424.
- Wassal, S.R. & Stillwell, W. (2009). Polyunsaturated fatty-acid-cholesterol interactions : domain formation in membranes. *Biochimica et Biophysica Acta*, Vol. 1788, No.1, (January 2009), pp.24-32, ISSN 0006-3002.
- Wong, S.W.; Kwon, M.-J.; Choi, A.M.K.; Kim, H.-P.; Nakahira, K. & Hwang, D.H. (2009). Fatty acids modulate toll-like receptor 4 activation through regulation of receptor dimerization and recruitment into lipid rafts in a reactive oxygen species-dependent manner. *Journal of Biological Chemistry*, Vol. 284, No.40, (October 2009), pp.27384-27392, ISSN 0021-9258.
- Wood, W.G. & Schroeder, F. (1988). Membrane effects of ethanol : bulk lipid versus lipid domains. *Life Science*, Vol.43, No.6, (1988), pp.467-475, ISSN 0024-3205.
- Wu, D. & Cederbaum, A.I. (2009). Oxidative stress and alcoholic liver disease. *Seminars in Liver Disease*, Vol.29, No.2, (May 2009), pp.141-154, ISSN 0272-8087.
- Yamada, S. & Lieber C.S. (1984). Decrease in microviscosity and cholesterol content of rat liver plasma membranes after chronic ethanol feeding. *The Journal of Clinical Investigation*, Vol.74, No.6, (December 1984), pp.2285-2289, ISSN 0021-9738.
- Yaqoob, P. & Shaikh, S.R. (2010). The nutritional and clinical significance of lipid rafts. *Current Opinion in Clinical Nutrition and Metabolic Care*, Vol.13, No.2, (March 2010), pp.156-166, ISSN 1363-1950.
- Yoon, Y.; Török, N.; Krueger, E.; Oswald, B & McNiven M.A. (1998). Ethanol-induced alterations of the microtubule cytoskeleton in hepatocytes. *The American Journal of Physiology*, Vol.274, No.4Pt1, (April 1988), pp.G757-G766, ISSN 0002-9513.



Trends in Alcoholic Liver Disease Research - Clinical and Scientific Aspects

Edited by Dr. Ichiro Shimizu

ISBN 978-953-307-985-1

Hard cover, 220 pages

Publisher InTech

Published online 11, January, 2012

Published in print edition January, 2012

Alcoholic liver disease occurs after prolonged heavy drinking. Not everyone who drinks alcohol in excess develops serious forms of alcoholic liver disease. It is likely that genetic factors determine this individual susceptibility, and a family history of chronic liver disease may indicate a higher risk. Other factors include being overweight and iron overload. This book presents state-of-the-art information summarizing the current understanding of a range of alcoholic liver diseases. It is hoped that the target readers - hepatologists, clinicians, researchers and academicians - will be afforded new ideas and exposed to subjects well beyond their own scientific disciplines. Additionally, students and those who wish to increase their knowledge will find this book a valuable source of information.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Odile Sergent, Fatiha Djoudi-Aliche and Dominique Lagadic-Gossmann (2012). Up-to-Date Insight About Membrane Remodeling as a Mechanism of Action for Ethanol-Induced Liver Toxicity, Trends in Alcoholic Liver Disease Research - Clinical and Scientific Aspects, Dr. Ichiro Shimizu (Ed.), ISBN: 978-953-307-985-1, InTech, Available from: <http://www.intechopen.com/books/trends-in-alcoholic-liver-disease-research-clinical-and-scientific-aspects/up-to-date-insight-about-membrane-remodeling-as-a-mechanism-of-action-for-ethanol-induced-liver-toxi>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen