

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

186,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



The Human Papilloma Virus – Ion Channel Link in Cancer: *An Alternative Opportunity for Diagnosis and Therapy*

Ana Ramírez and Javier Camacho

Department of Pharmacology,

Centro de Investigación y de Estudios Avanzados

Mexico City

Mexico

1. Introduction

Despite hundreds of clinical trials being conducted for cancer patients, the overall response is below 4% and cancer remains a major health problem worldwide (Roberts, 2007; Kamb et al., 2007). Indeed, early detection of the disease should help improve diagnosis and treatment leading to a reduction in cancer mortality. The arising link between human papilloma virus (HPV) and ion channels presents a very interesting opportunity for the early diagnosis and therapy of different types of cancer, including cervical, head and neck, oral cavity and lung cancer, some of them within those of the highest incidence in the world.

HPV infection has been proposed as the main etiological factor for cervical cancer (Walboomers et al., 1999; zur Hausen, 2002). Nevertheless, HPV infection has been also suggested to be associated with head, neck and oral cavity cancer (Anaya-Saavedra et al., 2008). Interestingly, in a population of taiwanesse women, more than 90% of lung cancer cases were not associated to cigarette smoke (Chen et al., 1990). Analysis of HPV expression and its E6 oncoprotein in lung cancer biopsies from non-smoker taiwanesse women, led to suggest HPV as a lung cancer risk factor in such population (Cheng et al., 2001; 2007). Therefore, HPV presence might be used as an early marker for several types of cancer.

Ion channels play important roles in cell physiology, including excitability, neural transmission, cardiac contraction, pancreatic cell metabolism, apoptosis and cell proliferation. Accordingly, alterations in either channel activity or expression are associated to several diseases (Ashcroft, F. 2006), and cancer is not an exception. Actually, several ion channels are suggested as tumor markers and therapeutic targets for different types of cancer, including those malignancies associated to HPV infection.

Ion channels are integral membrane proteins transferring small ions through the hydrophobic lipid bilayer of the cell membrane, such as potassium (K^+), sodium (Na^+), chloride (Cl^-) and calcium (Ca^{2+}). They are present in the plasma and intracellular membranes of every cell type in the human body. Most ion channels require the presence of a stimulus to be activated (*gated*), this can be accomplished by changes in the membrane

potential (*voltage-gated ion channels*), neurotransmitters or other molecules (*ligand-gated ion channels*), as well as light, temperature, mechanical forces, etc. Ion channels play an important role in a variety of cellular functions regulating every aspect of the cell physiology, the voltage-gated channels provide the ionic currents to generate and spread neuronal activity, calcium channels trigger synaptic transmission, hormonal secretion, and muscle contraction, and some channels participate in the regulation of cell migration, cell cycle progression, apoptosis and gene transcription. An increased number of human diseases has been found to result from defects in ion channel function or expression, including epilepsy, cardiac arrhythmias, skeletal muscle disorders and diabetes (Hübner, C. & Jentsch, T. 2002; Ashcroft, F. 2006). Since channels play an important role in proliferation and growth, and because cancer is a multifactorial disease, these membrane proteins also participate in tumor development. Many types of human cancers show alterations on ion channel expression presumably to help to transform healthy cells into malignant, invasive, and fast growing tissue (Schönherr, R. 2005).

In this Chapter, we begin with a general overview of ion channels in cancer followed by the specific description of the participation of potassium, calcium, sodium and chloride channels in tumor cells. Then we focus on cervical cancer as the best example of a malignancy in which a link between HPV and ion channels can be found. Lastly, we describe in detail the regulation of human oncogenic *ether à-go-go-1* (Eag1) channels by HPV oncogenes and estradiol.

2. Ion channels in cancer

The control of cell proliferation involves diverse signaling pathways, growth factors, and receptors, which have a restrict regulation in order to maintain the cell homeostasis (Vermeulen, et al., 2003). From a general view, the eukaryotic cell cycle consists of four phases (Figure 1) and some checkpoints (Norbury, C. & Nurse, P. 1992; Massagué, J., 2004). Ion channels coordinate the upstream and downstream signals that converge on the cell cycle machinery. Both voltage- and ligand-gated channels have been implicated in the control of different cell cycle checkpoints in normal as well as neoplastic cells. Cell proliferation involves at some point the activation of Cl^- channels, K^+ channels and Ca^{2+} channels; these channels appear to play an active role in the pathways leading to duplication of any given cell (Kunzelmann, K. 2005).

There are more than 400 genes encoding ion channel subunits that regulate the flow of ions across the plasma membrane and the intracellular organelle membrane. In tumor cells, during the change from normal to cancer phenotype, a series of genetic alterations occur in which genes encoding ion channels might be affected. This might lead to changes in either channel expression or activity, which may be responsible, in part, of the pathophysiological features that cause malignant growth.

Many types of ion channels have been described to play a potential role in the development and growth of cancer cells, some of them are listed in Table 1. Hallmarks of cancer have been recently reviewed and include altered cell cycle progression, self sufficiency in growth signals, insensitivity to antigrowth signals, evasion of apoptosis, adaptation to harsh conditions, sustained angiogenesis, tissue invasion and metastasis (Hanahan, D. & Weinberg, R., 2011). Most of the ion channels that contribute to the development of cancer

have the capacity to induce proliferation by regulating the cell cycle at some point. Cell proliferation is a highly regulated process in which ion channels participate as regulators of the cell cycle, but surprisingly, the same ion channel mechanisms that regulate cell proliferation are involved in the control of apoptosis (programmed cell death) (Wang, Z. 2004). Cell proliferation and apoptosis are two counterparts that are responsible for maintaining normal cellular functions. Abnormal enhanced proliferation and/or impaired apoptosis alters the cell homeostasis leading to loss of control of the cellular growth.




Fig. 1. Phases of the eukaryotic cell cycle.

Details of the different types of ion channels involved in proliferation or invasion of tumor cells are described in Table 1.

2.1 Potassium channels

Potassium channels are the most extensive family of ion channels and some of the most studied in cancer. Several potassium channels are overexpressed in tumors as compared to the corresponding healthy tissues. Some types of these channels include Kv's (voltage-gated potassium channels), K_{Ca} (calcium-activated potassium channels), K_{2P} (two-pore domain) and K_{ir} (inward rectifier). All of these classes of potassium channels are further subdivided into subfamilies (Chandy & Gutman, 1993), and they regulate a vast diversity of functions in the cells including maintenance of the cell membrane potential, regulation of the cell volume (Lang, 2007), and cell cycle progression (Blackiston et al., 2009).

Potassium ions are important for the osmotic regulation of cell volume by working in concert with the Na^+H^+ exchanger and the Na^+K^+ ATPase, contributing to the regulation of intracellular pH (pH_i) (Ikuma et al., 1998). Thus, K^+ channels participate in physiological and pathological proliferation by playing a role in membrane potential, cell volume and pH_i change during cell-cycle progression. K^+ channels also provide a hyperpolarizing effect on membrane voltage during cell cycle and therefore stimulate calcium influx. This increase in intracellular Ca^{2+} concentration is needed during progression through G1 phase and the G1/S transition (Wonderlin and Strobl, 1996; Kulzenmann, 2005). Usually, cancer cells have less negative voltage membrane than normal cells (Marino et al., 1994); therefore, it is likely that they would require a higher expression of certain types of K^+ channels to produce the transient hyperpolarization required to proceed with cell cycle progression. Therefore, K^+

Channels name IUPHAR	Type of cancer	Reference
K ⁺ channels		
K _v 1.3	Breast, Colon, Prostate	Abdul, M., et al. (2003) ; Abdul, M. & Hoosein, N. (2002a) ; Abdul, M. & Hoosein, N. (2002c)
K _v 1.5	Glioma	Preussat, K., et al. (2003)
K _v 3.4	Oral Squamous cell carcinoma	Chang, K., et al. (2003)
K _{ca} 1.1	Cervical, Ovarian, Breast, Glioma	Han, et al. (2007); Ouadid, H., et al. (2004a); Liu, X., et al. (2002)
K _{ca} 3.1	Prostate, Breast, Melanoma, Pancreas	Parihar, A., et al. (2003); Ouadid, H., et al. (2004b); Jäger, H., et al. (2004); Gavrilova, O., et al. (2002)
K _v 10.1	Breast, Colon, Cervix, Gastric, Prostate, Sarcoma, Ovarian	Pardo, L., et al. (1999); Hemmerlein, B., et al. (2006); Ousingsawat, J., et al (2007); Farias, L., et al. (2004); Ding, X., et al. (2007); Mello, F., et al. (2006); Asher, V., et al. (2010)
K _v 11.1	Colon, Endometrial, Glioma, Leukemia, Stomach	Lastraioli, E., et al. (2004); Cherubini, A., et al. (2000); Masi, et al., (2008); Smith, G., et al. (2002); Pillozzi, S., et al. (2002); Shao, X., et al. (2008)
K _{2p} 9.1	Breast, Lung	Mu, D., et al. (2003); Pei, L., et al. (2003)
K _{ir} 3.1	Lung	Plummer, H., et al. (2005)
Ca ²⁺ Channels		
Ca _v 1.2	Colon	Wang, X., et al. (2000)
Ca _v 3.2	Prostate	Mariot, P., et al. (2002)
Na ⁺ Channels		
Na _v 1.2	Prostate	Anderson, J., et al. (2003)
Na _v 1.4	Prostate	Bennett, E., et al. (2004)
Na _v 1.5	Breast, Ovarian	Roger, S., et al. (2003); Gao, R., et al. (2010)
Na _v 1.6	Cervical, Glioma	Hernandez, E., et al. (2011); Schrey, M., et al. (2002)
Na _v 1.7	Prostate	Diss, J., et al. (2205)
Other channels		
Cl channels	Breast, Cervical, Hepatoma, Glioma, Prostate	Abdel, M., et al. (2003); Mao, et al. (2009); Roman, R., et al. (2001); Olsen, M., et al. (2003) ; Shuba, Y., et al. (2000).
TRPC1	Prostate	Vanden, F., et al. (2003)
TRPV6	Breast, Colon, Ovarian, Prostate, Thyroid	Bolanz, K., et al. (2008); Vanden, F., et al. (2003); Zhuang, L., et al. (2002); Fixemer, T., et al. (2003)
TRPM8	Breast, Colon, Lung, Melanoma, Prostate	Tsavaler, L., et al. (2001); Yamamura, H., et al. (2008)
P2X7	Cervical, Leukemia, Neuroblastoma,	Feng, Y., et al. (2006); Wiley, J., et al. (2002); Per Larsson, K., et al. (2002)

Table 1. Examples of ion channels involved in cancer.

channel activation is particularly important for the early G1 phase of the cell cycle. During G1/S transition and S phase, cells swell and activate a regulatory volume decrease (RVD) mechanism, which allows shrinkage of the cell after mitosis. RVD requires efflux of water

and solutes, which is due to simultaneous opening of K^+ and Cl^- channels (Kulzenmann, 2005), allowing the cell to proliferate. Another evidence associating K^+ channels with proliferation is that voltage-gated K^+ channels blockers inhibit proliferation in many cell types; this has been observed in normal physiological proliferation (lymphocytes) and in pathological conditions (as in cancer cells). Thus the cells require K^+ channels in order to proceed with cell cycle progression (Wonderlin and Strobl, 1996).

Most studies are devoted to the impact of voltage-gated potassium channels on proliferation of tumor cells, particularly those of epithelial origin (Abdul & Hoosein, 2002; Farias et al., 2004; O'Grady & Lee, 2005; Pardo, 2004; Pardo et al., 2005; Arcangeli et al., 2009). One of the most studied voltage-gated potassium channel in cancer is the Eag1 channel. Eag1 displays oncogenic properties and shows a very restricted distribution in normal tissues (Pardo et al., 1999; Hemmerlein et al., 2006) but it is expressed in many types of tumors including cervical, breast, lung, prostate, liver and colon carcinoma (Farias et al., 2004; Hemmerlein et al., 2006; Ousingsawat et al., 2007). Inhibition of either channel activity or expression leads to decreased tumor cell proliferation both *in vitro* and *in vivo* (Pardo et al., 1999; Gómez-Varela et al., 2007; Ousingsawat et al., 2007). The restricted distribution of Eag1 and its role in proliferation have converted this ion channel in an attractive tool for diagnose and therapy of many cancers.

It has also been shown that several K^+ channels associate with other proteins related to proliferation, for instance Kv11.1 channels associate with 14-3-3 (Kagan et al., 2002), Src (Cayabyab, 2002), or TNF- α receptors (Wang et al., 2002), Kv1.3 associate with integrins (Levite et al., 2000) and p56lck (Hanada et al., 1997), and Kv10.1 associates with calmodulin (Schönherr et al., 2000). Increased expression of K^+ channels in tumors offers an additional tool for cancer diagnose and treatment.

2.2 Calcium channels

Calcium entry via different channels activates intracellular signalling cascades. Calcium channels play pivotal roles in many human diseases, particularly of the cardiac and nervous systems, for example, epilepsy, hypertension and migraine. In addition, these channels have been involved in pain and cancer. Cytosolic Ca^{2+} activity is necessary throughout the cell cycle since it plays a critical role on the regulation of cell proliferation (Whitfield et al., 1995; Parekh & Penner, 1997; Berridge et al., 1998, 2000, 2003). It has been shown that decreased levels of extracellular Ca^{2+} inhibit the progression through the G1 phase, causing cells to remain at the G1/S boundary. Moreover, Ca^{2+} is highly concentrated in intracellular stores and is released from the endoplasmic reticulum or mitochondria upon mitogenic stimulation (Nilius et al., 1993; Lepple et al., 1996). In excitable tissues, Ca^{2+} influx occurs through voltage-gated Ca^{2+} channels depending on the cell type. In non-excitable tissues, hyperpolarization of the membrane voltage is important for the increase of intracellular Ca^{2+} , providing the driving force for Ca^{2+} entry from the extracellular space.

Among voltage-gated Ca^{2+} channels, members of the Ca_v3 subfamily, and in particular $Ca_v3.2$ channels, are implicated in proliferation. It has been observed that voltage-gated calcium channels Ca_v1 (L-type currents), are expressed in non-proliferative phases, while expression of Ca_v3 channels (T-type currents) often increases during the proliferative phases. This is observed in both normal and cancer cells, although the precise physiological significance is uncertain. $Ca_v3.2$ channels are expressed in several cancer-derived cell lines,

and blockage of the channel generates antiproliferative action (Roger et al., 2006, Panner & Wurster, 2006). T-type calcium channels have been found in lung carcinoma cells (Oguro-Okano et al., 1992) and calcium influx through different ion channels has been suggested to participate in migration and invasion of prostate, breast and fibrosarcoma cells (Monet et al., 2009; Yang et al., 2009; Huang et al., 2004).

Other types of calcium channels have also been involved in cancer progression and prognosis. For instance, expression of the Ca^{+2} channel TRPV6 was correlated with prostate cancer grade, in which patients with positive TRPV6 prostate cancer had a poor prognosis (Fixemer et al., 2003).

2.3 Sodium channels

Voltage gated sodium channels (VGSCs) that selectively conduct sodium due to changes in membrane potential, allow sodium entry and thus the propagation of depolarization along the plasma membrane of nerve, muscle and other electrically excitable cells. Non-voltage-gated sodium channels like the degenerin/epithelial sodium channel (ENaC) superfamily, which are permeable also to lithium and potassium, is a group of proteins involved in diverse biological processes, including sodium homeostasis, salt taste, nociception, pain transduction, touch sensation and mechanotransduction (Goldin et al., 2000). Sodium channels have various functional and pharmacological properties in different tissues and species, and VGSCs in particular play an important role in generating action potentials. There are ten genes that encode VGSCs α subunits, from which nine constitute a single family named Nav1 (Nav1.1 to Nav1.9), whose members associate with one or more auxiliary β subunits (Nav β 1 to Nav β 4) to form the whole channel protein complex (Yu et al., 2003). The remaining isoform, Na_x, shows a structure diverging from the Nav1 family and seems to be gated by sodium concentration and not by voltage.

VGSCs have been suggested as participants in the development of cancer. Enhanced metastasis correlates with the appearance of membrane channels and currents that are characteristic of excitable membranes. Metastasis is a process where cells escape from a primary tumor, enter circulation (blood or lymph), migrate and invade other tissues, proliferate and form secondary tumors. In *in vitro* experiments, it has been shown that VGSCs are associated to proliferation, motility, and invasion of breast, lung, ovary and prostate cancer (Roger et al., 2003; Gao et al., 2010; Diss et al. 2005; Chioni et al., 2010; Roger et al., 2007; House, et al., 2010; Bennett et al., 2004). In prostate cancer cells, the main VGSC overexpressed is the Nav1.7 subunit, while in breast, colon and ovary the Nav1.5 subunit is the predominant subunit overexpressed. In addition, functional expression of VGSCs (mainly Nav1.7, Nav1.6 and Nav1.5) is often associated with metastasis and VGSCs have been found in biopsies from prostate and breast metastatic cancer (Roger et al., 2006). The mechanisms responsible for VGSCs upregulation and for their pro-invasive roles are still poorly understood, but diverse hypothesis exists. One hypothesis strongly suggests that there is a regulation of growth factors release and/or activity which is common to all the cancers described, highly malignant cancer cell types that overexpress VGSCs also express growth factors such as epidermal growth factor (EGF) and nerve growth factor (NGF) (Brackenbury & Djamgoz, 2007; Uysal & Djamgoz, 2007), emphasizing that growth factors could play a major role in upregulating VGSCs. Another hypothesis describes that embryonic genes, which are silent in the cells of the mature organ, are re-expressed in cancer

cells (Monk & Holding, 2001). This might be the case for VGSCs, since highly metastatic cancers (prostate and breast) mostly express embryonic isoforms of VGSCs (Diss et al., 2005; Fraser et al., 2005). Hence overexpression of VGSCs contributes to the physiological and pathophysiological invasive processes in several metastatic cancers, representing a potential target to inhibit invasion and metastasis.

2.4 Chloride channels

Cl⁻ channels play a crucial role in controlling the ionic composition of the cytoplasm and the volume of cells. According to their gating mechanisms there are five classes of chloride channels:

- Voltage-gated chloride channels (CLC).
- Volume/swell-regulated/sensitive anion/chloride channels (VRAC).
- Cystic fibrosis transmembrane conductance regulator (CFTR).
- Calcium-activated chloride channels (CLCA).
- Ligand-activated chloride channels, which mainly form synaptic channels.

Cl⁻ channels play a crucial role in controlling the ionic composition of the cytoplasm and the cell volume. Cl⁻ channels are expressed in a variety of tumor cells and participate in cell proliferation, invasion and migration. Proliferation is associated with volume increase along the G1 phase, but non-specific cell swelling can inhibit proliferation. To regulate their volume, cells are endowed with various ions and organic osmolyte transport proteins that become activated upon cell swelling or shrinkage. In the presence of a significant water permeability of the plasma membrane, water follows osmotically, resulting in a regulated change of cell volume. This is called regulatory volume increase (RVI) or regulatory volume decrease (RVD). RVI most often involves the uptake of Na⁺ and Cl⁻, for instance, by the concomitant activation of Na⁺/H⁺ and Cl⁻/HCO₃⁻ exchangers. Na⁺ is replaced by K⁺ through the Na⁺-K⁺-ATPase, resulting in a net intracellular accumulation of KCl. In RVD, intracellular KCl may be extruded by KCl cotransporters or by the concerted activation of swelling-activated Cl⁻ channels and K⁺ channels (Jentsch et al., 2002). Cell proliferation has been shown to correlate with increases in cell volume in fibroblasts, mesangial cells, lymphocytes, human promyelocytic leukemia cells (HL-60 cells), hybridoma cells (GAP A3), smooth muscle cells and cervical carcinoma cells (HeLa cells). The signaling of cell proliferation needs at some stage transient cell shrinkage, which may require the activation of Cl⁻ channels. Usually, intracellular Cl⁻ is above electrochemical equilibrium and activation of Cl⁻ channels leads to Cl⁻ efflux and thus depolarization. As long as K⁺ channels are active, the Cl⁻ outward movement is paralleled by K⁺ efflux. The loss of KCl and water shrinks the cells (Lang et al., 1998).

It has been observed that pharmacological inhibition of Cl⁻ channels impairs the cell ability to migrate and limits tumor progression in experimental tumor models (McFerrin & Sontheimer, 2006). The outer membrane of mitochondria contains a Cl⁻ selective porin, the so-called voltage-dependent anion channel (VDAC). On the other hand, VRAC's modulate progression of nasopharyngeal carcinoma cells through the G1 restriction point and endow nasopharyngeal carcinoma cells with enhanced proliferation ability. In addition, it is reported that VRAC displays cell cycle-dependent expression in tumor cells. Also CLC-3, an important member of the CLC superfamily, plays a crucial role in a variety of cellular

processes, including cell proliferation and cell cycle progression, and it has been demonstrated that inhibition of CLC-3 protein expression down-regulates invasion and migration ability of tumor cells.

3. Ion Channels in apoptosis

Cell homeostasis requires a delicate balance between formation of new cells by cell proliferation and cell elimination by apoptosis (programmed cell death). Apoptosis eliminates abundant and potentially harmful cells (Green & Reed, 1998). Apoptotic pathways involve several proteins with great enzymatic cell degrading potential including Bcl-2, caspases and cytochrome C. This process is initiated by death-promoting molecules such as TNF- α or CD95 Fas ligand, DNA damage, lack of growth factors, or cell exposure to genotoxics like radiation or oxidants. Apoptosis evasion might be one of the first steps for the transformation of normal cells into cancerous cells, and it is a tightly regulated and highly efficient cell death program which requires the interplay of multiple factors. Upon receiving specific signals instructing the cell to enter apoptosis, distinctive changes occur in the cell including shrinkage, nuclear condensation, DNA fragmentation, formation of sub-cellular apoptotic bodies, and mitochondrial depolarization. Ion channels have been involved in the regulation of apoptosis.

Potassium ions must leave the cell as an obligatory step in the apoptotic pathway. Activation of K⁺ channels and loss of intracellular K⁺ has been directly correlated to trigger apoptosis (Wang, 2004). Therefore, inhibition of apoptosis might take place by either increasing extracellular K⁺ concentration (Prehn et al., 1997; Colom et al., 1998) or inhibiting K⁺ channels (Lang et al., 2003). In any case, cellular K⁺ loss seems to be an important trigger of apoptosis in a wide variety of cells. Activation of K⁺ channels leads to hyperpolarization of the cell membrane, thus increasing the electrical driving force for Cl⁻ outflow into the extracellular space. Then, if K⁺ channel activity is paralleled by Cl⁻ channel activity, it leads to cellular loss of KCl with osmotically obliged water and hence to cell shrinkage, a hallmark of apoptosis (Lang et al., 1998). Subsequently, metabolic enzymes are activated, such as caspases and nucleases, which further propagate death signals. Remarkably, all these enzymes are controlled by the intracellular K⁺ concentration and while the concentration of various ions may change during apoptotic cell shrinkage, that of K⁺ plays a necessary and probably pivotal role in the cell death program. Cl⁻ and K⁺ conductances must stay within a certain values in order to support proliferation, otherwise programmed cell death is triggered (Lang et al., 2004).

Sustained increase of cytosolic Ca²⁺ activity has been shown to trigger apoptosis in a variety of cells (Green & Reed, 1998; Spassova et al., 2004; Parekh & Putney, Jr., 2005). Cytosolic Ca²⁺ may trigger mechanisms required for cell proliferation and stimulate enzymes executing apoptosis. Ca²⁺ signal convergence results in activation of intracellular channels that leads to cytochrome C release from mitochondria. During apoptosis, small amounts of mitochondrial cytochrome C translocate to the endoplasmic reticulum and trigger Ca²⁺ release via IP3R channels. This leads to a bursting Ca²⁺ overload, which coordinates massive cytochrome C release from mitochondria, leading to activation of the caspase cascade, essential for the development of apoptosis (Boehning, D. *et al.*, 2003). Both the magnitude, space- and time- occurrence of Ca²⁺ entry is a major determinant to trigger apoptosis.

Both cell proliferation and apoptosis involve at some point activation of Cl^- channels, K^+ channels and Ca^{2+} channels. Due to complex interaction with other signaling pathways, a given ion channel may play a dual role in both cell proliferation and apoptosis.

At this point we focus on cervical cancer, a very well recognized cancer type strongly associated to HPV infection and ion channel expression. A potential link between HPV and ion channel expression has been also proposed for this type of cancer.

4. Overview of ion channels in cervical cancer

Expression of several ion channels has been reported in cervical cancer suggesting these proteins as potential markers and/or therapeutic targets for this malignancy. Large Ca^{2+} -activated K^+ (BK) channels are ion channels activated by changes in membrane electrical potential and/or by increases in the intracellular concentration of Ca^{2+} , contributing to cell proliferation and migration (Yuan et al., 2010). BK channels are expressed in HeLa cells, and it has been observed that they play a significant role in the regulation of proliferation of this cell line. Blockage of BK channels in HeLa cells results in tumor cell apoptosis and cycle arrest at G1 phase. The transduction pathway underlying such anti-proliferative effects is linked to the increased expression of apoptotic protein p53 and the decreased expression of its chaperone heat shock proteins (Hsp). Heat shock proteins function as intra-cellular chaperones (proteins that bind to and stabilize an otherwise unstable conformer of another protein) for other proteins, such as p53 (Zylicz M, King FW, Wawrzynow A., 2001); upon cell stress, the levels of these proteins increase dramatically. Other K^+ channels induce expression of heat shock proteins to exert tissue protective effects (Shinohara et al. 2004), suggesting that the intracellular potassium homeostasis may play a certain role in modulating heat shock proteins. The tumor suppressor gene p53 is involved in a variety of cellular processes including induction of G1 arrest and apoptosis by transactivating a number of downstream genes (Jin S, Levine A., 2001). One of those genes is p21 Cip1 (cyclin-dependent kinase inhibitor 1A), which is a cell-cycle regulatory protein that interacts with cyclin-CDK2 and -CDK4, inhibiting cell cycle progression at G1. The expression of p21 is tightly controlled by p53, through which this protein mediates the p53-dependent cell cycle arrest at G1 phase (Zamzami N, Kroemer G., 2005). Blockage of BK channels increases p53, p21Cip1, and Bax protein levels inducing cell arrest and apoptosis. In addition, inhibition of BK channels decreased the expression levels of some Hsp's, which could be an upstream signaling of the BK channel-mediated p53 change. This would keep p53 under negative control by the activity of BK channels, inhibiting apoptosis in cervical cancer (Han et al., 2007).

Purinergic ATP-gated nonselective ion channels (P2X receptors) have been also suggested to participate in cervical cancer. These channels are permeable to Ca^{2+} , Na^+ and K^+ (Burnstock, 2004) and several members have been identified and termed P2X1 through P2X7. Extracellular ATP is a physiological ligand that activates P2X7 receptor (Surprenant et al., 1996). Under normal conditions, extracellular ATP is present in only low concentrations, but increases significantly under inflammatory conditions and in response to tissue trauma (e.g., ischemia/hypoxia). The ionotropic purinergic P2X7 receptors activate a diverse range of cellular responses that play a role in evading apoptosis of cervical cancer cells. In normal cervical epithelial cells, the activation of P2X7 receptors forms a pore that increases calcium influx and induces apoptosis via the Ca^{2+} -dependent mitochondrial pathway. Human

cervical cancer cells express a P2X splice variant (P2X7j) that do not mediate P2X7-dependent apoptosis, causing downregulation of the functional P2X7 receptor, thereby preventing the Ca^{2+} influx required to trigger apoptosis and leading to defective apoptosis and enhanced growth of cervical cancer cells (Feng, et al., 2006).

Voltage-gated sodium channels (VGSC) have also been detected in cervical carcinoma cells that may be involved in metastasis. Expression of Nav1.4, Nav1.6 and Nav1.7 channels has been detected in biopsies from cervical cancer (Díaz et al., 2007). Whether the activity of the sodium channel participates in the invasiveness of cervical cancer cells remains to be determined, but the pharmacological blockade of VGSCs commonly results in the reduced migration of highly metastatic cell lines, whereas the facilitation of channel opening by agonists enhances migration without impairing cell proliferation or viability (Roger et al., 2007). The abnormal expression of these VGSCs might be used as a tumor marker and a potential therapeutic target for cervical carcinoma.

It has been suggested that the expression of volume-activated current ($I_{\text{Cl,vol}}$) in HeLa cells contributes to the cell cycle dependent regulation of cell migration. The density of $I_{\text{Cl,vol}}$ was positively correlated to the rate of cell migration during cell-cycle progression indicating that volume-activated Cl^- channels are involved in the cell-cycle-dependent migratory behaviour of HeLa cells. Moreover, expression inhibition of the voltage-gated chloride channel ClC-3 arrested HeLa cells in S phase (Mao et al., 2009). The capacity of Cl^- channels to participate in migration could be associated to cell volume regulation, since regulatory volume decrease plays an important role in migration of tumor cells. Cl^- channels help the cell to undergo changes in their shape and volume, facilitating tumor cells to move through the extracellular space. In the following section we describe the very interesting oncogenic Eag channels, which are to our best knowledge the only known channels regulated by HPV oncogenes.

5. Oncogenic Eag1 K^+ channels in cervical cancer and regulation by HPV oncogenes

Ether à go go (Eag1) K^+ channel has gained great interest in cervical cancer diagnosis and therapy (Farias et al., 2004). Eag1 is a voltage gated K^+ channel first described as a cell-cycle regulated channel (Brüggemann et al., 1997; Camacho et al., 2000), and was the first ion channel identified to display oncogenic properties (Pardo et al., 1999). These channels are widely distributed in the central nervous system, but their expression in peripheral tissues is very restricted, finding the channel only in placenta, transiently in myoblasts, testis and adrenal glands (Occhiodoro et al., 1998; Pardo et al., 1999; Hemmerlein et al., 2006). This restricted distribution in normal tissues is one of the most attractive features of Eag1 as a potential tumor marker. Eag1 channels are overexpressed in various cancer cell lines including IGR1, IPC298, and IGR39 from melanoma, SH-SY5Y from neuroblastoma, MCF-7 from breast cancer and HeLa from cervical cancer. Eag1 channels have transforming properties, they confer loss of contact inhibition and sustained growth in the absence of serum. Cells transfected with Eag1 channel and implanted into immunosuppressed mice induce formation of aggressive tumors (Pardo et al. 1999). Eag1 has been found to be overexpressed in many types of tumors including breast, lung, prostate, liver and colon (Hemmerlein et al., 2006). Thus, Eag is a promising tumor

marker. On the other hand, inhibition of either channel activity or expression decreases proliferation of tumor cells both *in vitro* and *in vivo* (Pardo et al., 1999; Ousingsawat et al., 2007; Gómez-Varela et al. 2007; Downie et al., 2008; Díaz et al., 2009). Therefore, Eag is also a promising therapeutic target for many types of cancer (Camacho, 2006; Pardo & Stühmer, 2008).

Eag1 mRNA expression was found in 100% of cervical cancer human biopsies while only in 33% of normal control samples (Farias et al., 2004). Interestingly, in this study it was observed that in one of the patients that was submitted to hysterectomy without any previous evidence of cervical malignancy (negative pap smears), postsurgery pathological studies showed an unexpected endocervical adenocarcinoma expressing Eag1. This case, although unique in such study, emphasizes the potential significance of Eag1 as a tumor marker in cervical cancer (Farias et al., 2004; Camacho, 2006). In addition, one of the normal cervical samples that were positive for Eag1 expression was correlated with HPV infection. This led to suggest that Eag1 expression in normal cervix could be an early sign of tumor development associated to HPV infection.

The primary transforming activity of high-risk HPVs is provided by the E6 and E7 oncoproteins which act cooperatively in the development of HPV-induced cancers. A primary target of E7 is the retinoblastoma (Rb) family of proteins that control the activity of E2F transcription factors, which are key regulators of S phase genes. The efficient abrogation of Rb function by E7 leads to increased levels of p53 and, consequently, the E6 proteins have evolved to target p53 for degradation. Researchers investigated the potential link between HPV infection and Eag oncogenic channels by transfecting normal human keratinocytes with E6 and/or E7 HPV oncogenes (Díaz et al., 2009). They observed that normal human keratinocytes do not express Eag channels. Interestingly, keratinocytes transfected with either E6, E7 or both HPV oncogenes, displayed a strong Eag channel expression. This finding suggests a novel mechanism by which HPV induces tumor formation, namely, up-regulation of the oncogenic Eag1 channel.

Regulation of Eag1 channels by E6/E7 HPV oncogenes suggests that the regulation of Eag1 in cervical cancer might be via p53 and Rb pathways. A novel signaling pathway has been described that might explain how Eag is regulated by p53. It appears that Eag1 is a terminal component in the p53-miR-34-E2F1 pathway. miR-34 is a micro RNA, which silences expression of target genes through the RNA interference pathway and is commonly downregulated in human cancers; one example is miR-34, which is a direct target of p53. miR-34 transcription is activated by p53, and expression of miR-34 inhibits proliferation by inducing cellular senescence and cell cycle arrest at G1. When cellular stress or damage exist, p53 increases, causing the miR-34 transcription to increase, and the increased miR-34 will decrease the transcription factor E2F1. One of the target genes for transcription of E2F1 is the K⁺ channel Eag1. Suppression of E2F1 will repress Eag1, this will diminish Eag1 expression and function, resulting in a shut down of the cell proliferation or a cell cycle arrest, thus upregulation of miR-34 represses E2F1 and Eag1. Therefore, p53 negatively regulates Eag1 expression by a negative feed forward mechanism through the p53-miR-34-E2F1 pathway (Lin et al., 2011). This pathway might also help to explain the effect of the E7 oncogene on Eag regulation. E7 binds to Rb disrupting the Rb-E2F complexes, resulting in the constitutive expression of E2F1-responsive genes including Eag1.

6. Effect of estrogens on HPV and eag channels

HPV infection has been suggested as a necessary but not sufficient factor to induce cervical cancer. Thus other contributing factors have been proposed, especially estrogens. The uterine cervix is highly responsive to steroidal hormones and the use of oral contraceptives and multiple pregnancies have been shown to significantly increase the risk for cervical cancer in HPV-infected women (Moreno et al., 2002; Muñoz et al., 2002). In mice models for HPV-associated cancers, estrogen is required for the development of cervical and vaginal cancers (Brake & Lambert, 2005). The estrogen receptor alpha (ERα) is also required in mice for these cancers to develop and ER antagonists can cause efficient regression of cancer, dysplasia, and atypical squamous metaplasia, preventing malignant progression (Chung & Lambert, 2009). Other studies also suggest the relevance of estrogen in the carcinogenesis of cervical cancer. Growth stimulation of SiHa cervical cancer cells by estrogens appeared to be related to the increased expression of HPV E6/E7 oncogenes, estradiol stimulated both cell growth and transcription of E6/E7 viral oncogenes (Rosembaum, et al., 1989; Kim, et al., 2000). In addition, overexpression of aromatase, the enzyme that transforms testosterone into estrogen, is known to increase estrogen activity in breast tissue, and in cervical cancers it has been reported that 35% of human cervical cancer tested express aromatase, but




Fig. 2. Regulation of Eag1 oncogenic Eag1 channels by HPV oncogenes and estradiol. Eag1 might become extremely up-regulated since HPV oncogenes are in turn regulated by estradiol, leading cervical cells to cancer.

aromatase expression was not detected in precancerous or normal cervical tissue samples. Aromatase overexpression induced the expression of cyclin D1, proliferating cell nuclear antigen and HPV oncogenes (Nair, et al., 2005). It was also observed that women that expressed higher levels of estrogen receptors transcripts were significantly more likely to have cervical HPV infection; it may be that the presence of the receptor allows cellular acquisition of HPV and increased viral transcription (Shew, M., et al. 2005). These data suggest that a mechanism of synergistic cooperation exist between estrogen exposure and viral oncogenes, so it raises the possibility that steroidal hormones, such as estrogen, might affect cancers of the cervix, much like that of other hormonally responsive female organs. In summary, several studies suggest that estrogens play a critical role not only in the genesis of cervical cancer but also in its persistence and continuous development.

Eag1 channel expression is also up-regulated by estrogens. HeLa cells transfected with ER α and treated with 17- β estradiol, induce a strong up-regulation of Eag1 channels. These results suggest ER α activation as one of the mechanisms of the estrogenic regulation of Eag1. Thus the regulation of the Eag1 channel in cervical cancer is mediated by HPV oncogenes and estrogens (Diaz, L., et al., 2009). HPV-infection in cervical cells might lead to a very significant increase in oncogenic Eag channels. Since estrogens might regulate both HPV-oncogens and Eag1 channels, these related pathways might easily drive the cell into a tumor phenotype (Figure 2).

7. Conclusion

Ion channels are emerging as potential tools for diagnosis and treatment of many types of cancer including those where HPV infection plays a major role. Particularly, up-regulation of the oncogenic human Eag channel by HPV oncoogens offers a novel mechanism by which HPV associates to some types of cancer. Since several ion channels are up-regulated by HPV oncogenes and/or factors closely associated to HPV, for instance estrogens, simultaneous detection of HPV infection and ion channel expression should provide an alternative option for early detection of tumors. Besides, HPV detection might also serve to target ion channels involved in tumor progression. This combined approach should help reduce cancer mortality.

8. Acknowledgment

We thank all of our collaborators participating in our publications. The current research project on cervical cancer in the laboratory of JC is partially funded by Conacyt, Mexico (Grant No. 141126).

9. References

- Abdel, M., Cheng, H., Elble, R., Lin, H., DiBiasio, J. & Pauli, B. (2003). The interacting binding domains of the beta (4) integrin and calcium-activated chloride channels (CLCAs) in metastasis. *The Journal of Biological Chemistry*. Vol. 278, pp. 49406-49416.
- Abdul, M., & Hoosein, N. (2002a). Voltage-gated potassium ion channels in colon cancer. *Oncology Reports*, Vol. 9, pp. 961-964.

- Abdul, M., & Hoosein, N. (2002b). Voltage-gated sodium ion channels in prostate cancer: expression and activity. *Anticancer Research*, Vol. 22, pp. 1727–1730.
- Abdul, M. & Hoosein, N. (2002c). Expression and activity of potassium ion channels in human prostate cancer. *Cancer Letters*. Vol. 186, pp. 99-105.
- Abdul, M., Santo, A. & Hoosein, N. (2003). Activity of potassium channel-blockers in breast cancer. *Anticancer Research*. Vol. 23, pp. 3347-3351.
- Anaya-Saavedra, G. Ramírez-Amador V, Irigoyen-Camacho ME, García-Cuellar CM, Guido-Jiménez M, Méndez-Martínez R, & García-Carrancá A. (2008). High association of human papillomavirus infection with oral cancer: a case-control study. *Archives of Medical Research*, Vol. 39, pp. 189-197.
- Anderson, J., Hansen, T., Lenkowski, P., Walls, A., Choudhury, I., Schenck, H., Friehling, M., Höll, G., Patel, M., Sikes, R. & Brown, M. (2003). Voltage-gated sodium channel blockers as cytostatic inhibitors of the androgen-independent prostate cancer cell line PC-3. *Molecular Cancer Therapeutics*. Vol. 11, pp. 1149-1154.
- Arcangeli, A., Crociani, O., Lastraioli, E., Masi, A., Pillozzi, S., & Becchetti A., (2009). Targeting ion channels in cancer: a novel frontier in antineoplastic therapy. *Current Medicinal Chemistry*, Vol. 16, pp. 66–93.
- Ashcroft, F. (2006). From molecule to malady. *Nature*, Vol. 440, pp. 440-447.
- Asher, V., Khan, R., Warren, A., Shaw, R., Schalkwyk, G., Bali, A. & Sowter, H. (2010). The Eag potassium channel as a new prognostic marker in ovarian cancer. *Diagnostic Pathology*. Vol. 7.
- Bennett, E., Smith, B., & Harper, J. (2004). Voltage-gated Na⁺ channels confer invasive properties on human prostate cancer cells. *Pflügers Archiv European Journal of Physiology*. Vol. 447, pp. 908– 914.
- Berridge, M., Bootman, M., & Lipp, P. (1998). Calcium—a life and death signal. *Nature*, Vol. 395, pp. 645– 648.
- Berridge, M., Lipp, P., & Bootman, M. (2000). The versatility and universality of calcium signalling. *Nature Reviews Molecular Cell Biology*, Vol. 1, pp. 11–21.
- Berridge, M., Bootman, M., & Roderick, H. (2003). Calcium signalling: dynamics, homeostasis and remodelling. *Nature Reviews Molecular Cell Biology*, Vol. 4, pp. 517– 529.
- Blackiston, D.J., McLaughlin, K.A., & Levin, M. (2009). Bioelectric controls of cell proliferation: ion channels, membrane voltage and the cell cycle. *Cell Cycle*, Vol. 8, No. 21, pp. 3519-3528.
- Boehning, D., Patterson, R., Sedaghat, L., Glebova, N., Kurosaki, T., & Snyder, S. (2003). Cytochrome c binds to inositol (1,4,5) trisphosphate receptors, amplifying calcium-dependent apoptosis. *Nature Cell Biology*. Vol. 5, No. 12, pp. 1051-1061.
- Bolanz, K., Hediger, M. & Landowski, C. (2008) The role of TRPV6 in breast carcinogenesis. *Molecular Cancer Therapeutics*. Vol. 7, pp. 271–279.
- Brackenbury, W. & Djamgoz, M. (2007) Nerve growth factor enhances voltage-gated Na⁺ channel activity and Transwell migration in Mat-LyLu rat prostate cancer cell line. *Journal of Cell Physiology*, Vol. 210, pp. 602–608.

- Brüggemann A., Stühmer W., & Pardo L. (1997). Mitosis-promoting factor-mediated suppression of a cloned delayed rectifier potassium channel expressed in *Xenopus* oocytes. *Proceedings of the National Academic of Sciences*, Vol. 94, pp. 537-542.
- Brake, T. & Lambert, P. (2005) Estrogen contributes to the onset, persistence, and malignant progression of cervical cancer in a human papillomavirus-transgenic mouse model. *Proceedings of the National Academy of Sciences USA*. Vol. 102, pp. 2490-2495.
- Burnstock, G. (2004). Introduction: P2 receptors. *Current Topics in Medicinal Chemistry*, Vol. 4, pp. 793– 803.
- Camacho, J. (2006). Ether à go-go potassium channels and cancer. *Cancer Letters*, Vol. 233, pp. 1-9.
- Camacho J., Sánchez A., Stühmer W., & Pardo, L., (2000). Cytoskeletal interactions determine the electrophysiological properties of human EAG potassium channels. *Pflügers Archiv European Journal Physiology*, Vol. 441, pp. 167-174.
- Cayabyab, F., & Schlichter, L., (2002). Regulation of an ERG K⁺ current by *Src* tyrosine kinase. *Journal of Biological Chemistry*, Vol. 277, pp. 13673–13681.
- Chandy, K., & Gutman, G. (1993). Nomenclature for mammalian potassium channel genes. *Trends in Pharmacological Sciences*, Vol. 14, No. 12, Dec, 1993, pp. 434.
- Chang, K., Yuan, T., Fang, K., Yang, F., Liu, C., Chang, C. & Lin, S. (2003). The increase of voltage-gated potassium channel Kv3.4 mRNA expression in oral squamous cell carcinoma. *Journal of Oral Pathology and Medicine*. Vol. 32, pp. 606-611.
- Chen CJ, Wu HY, Chuang YC, Chang AS, Luh KT, Chao HH, Chen KY, Chen SG, Lai GM, Huang HH, et al. (1990). Epidemiologic characteristics and multiple risk factors of lung cancer in Taiwan. *Anticancer Research*, Vol. 10, pp. 971-979.
- Cheng, YW, Chiou HL, Sheu GT, Hsieh LL, Chen JT, Chen CY, Su JM, & Lee H. (2001). The association of human papillomavirus 16/18 infection with lung cancer among non-smoking Taiwanese women. *Cancer Research*, Vol. 61, pp. 2799-2803.
- Cheng, YW, Wu MF, Wang J, Yeh KT, Goan YG, Chiou HL, Chen CY, & Lee H. (2007). Human papillomavirus 16/18 E6 oncoprotein is expressed in lung cancer and related with p53 inactivation. *Cancer Research*, Vol. 67, pp. 10686-10693.
- Cherubini, A., Taddei, G., Crociani, O., Paglierani, M., Buccoliero, A., Fontana, L., Noci, I., Borri, P., Borrani, E., Giachi, M., Becchetti, A., Rosati, B., Wanke, E., Olivotto, M. & Arcangeli, A. (2000). HERG potassium channels are more frequently expressed in human endometrial cancer as compared to non-cancerous endometrium. *British Journal of Cancer*. Vol. 83, pp. 1722-1729.
- Chioni, A., Shao, D., Grose, R. & Djamgoz, M. (2010). Protein kinase A and regulation of neonatal Nav1.5 expression in human breast cancer cells: activity-dependent positive feedback and cellular migration. *International Journal of Biochemistry and Cell Biology*, Vol. 42, pp. 346-358.
- Chung, S., & Lambert, P. (2009). Prevention and treatment of cervical cancer in mice using estrogen receptor antagonists. *Proceedings of the National Academic of Sciences* Vol. 106, pp. 19467–19472

- Colom, L., Diaz, M., Beers, D., Neely, A., Xie, W., & Appel, S. (1998). Role of potassium channels in amyloid-induced cell death. *Journal of Neurochemistry*, Vol. 70, pp. 1925–1934.
- Diaz, D., Delgadillo, D., Hernández, E., Ramírez, M., Hinojosa, L., Ortiz, C., Berumen, J., Camacho, J. & Gomora, J. (2007) Functional expression of voltage-gated sodium channels in primary cultures of human cervical cancer. *Journal of Cell Physiology*, Vol. 210, pp. 469–478.
- Díaz, L., Ceja, I., Restrepo, I., Larrea, F., Avila, E., García, R., Borja, E., Barrera, D., Ahumada, E., Gariglio, P., Alvarez, E., Ocadiz, R., Garcia, E., Hernandez, E., Camacho, I., Morales, A., Ordaz, D., Garcia, E., Escamilla, J., Sanchez, L., Saqui, M., Gamboa, A., Vera, E., Uribe, M., Murbartian, J., Ortiz, C., Rivera, C., De Vizcaya, A. & Camacho, J. (2009). Estrogens and human papilloma virus oncogenes regulate Human Ether-à-go-go-1 Potassium Channel Expression. *Cancer Research*, Vol. 69, pp. 3300–3307.
- Ding, X., Luo, H., Jin, X., Yan, J. & Ai, Y. (2007). Aberrant expression of Eag1 potassium channels in gastric cancer patients and cell lines. *Medical Oncology*. Vol. 24, pp. 345–350.
- Diss, J., Stewart, D., Pani, F., Foster, C., Walker, M., Patel, A. & Djamgoz, M. (2005). A potential novel marker for human prostate cancer: voltage-gated sodium channel expression in vivo. *Prostate Cancer and Prostatic Diseases*, Vol. 8, pp. 266–273.
- Downie, B., Sánchez, A., Knötgen, H., Contreras, C., Gymnopoulos, M., Weber, C., Stühmer, W. & Pardo, L., (2008). Eag1 expression interferes with hypoxia homeostasis and induces angiogenesis in tumors. *Journal of Biological Chemistry*, Vol. 283, pp. 36234–36240.
- Farias, L., Ocana, D., Diaz, L., Larrea, F., Avila, E., Cadena, A., Hinojosa, L., Lara, G., Villanueva, L., Vargas, C., Hernandez, E., Camacho, I., Duenas, A., Perez, E., Pardo, L., Morales, A., Taja, L., Escamilla, J., Sanchez, C., & Camacho, J. (2004). Ether à go-go potassium channels as human cervical cancer markers. *Cancer Research*, Vol. 64, pp. 6996–7001.
- Feng, Y., Li, X., Wang, L., Zhou, L. & Gorodeski GI. (2006). A truncated P2X7 receptor variant (P2X7-j) endogenously expressed in cervical cancer cells antagonizes the fulllength P2X7 receptor through hetero-oligomerization. *Journal of Biological Chemistry*, Vol. 281, pp. 17228–17237.
- Fixemer, T., Wissenbach, U., Flockerzi, V. & Bonkhoff, H. (2003) Expression of the Ca²⁺-selective cation channel TRPV6 in human prostate cancer: a novel prognostic marker for tumor progression. *Oncogene*, Vol. 22, pp. 7858–7861
- Fraser, S., Diss, J., Chioni, A., Mycielska, M., Pan, H., Yamaci, R., Pani, F., Siwy, Z., Krasowska, M., Grzywna, Z., Brackenbury, W., Theodorou, D., Koyutürk, M., Kaya, H., Battaloglu, E., De Bella, M., Slade, M., Tolhurst, R., Palmieri, C., Jiang, J., Latchman, D., Coombes, R. & Djamgoz, M. (2005) Voltage-gated sodium channel expression and potentiation of human breast cancer metastasis. *Clinical Cancer Research*, Vol. 11, pp. 5381–5389.

- Gao, R., Shen, Y., Cai, J., Lei, M. & Wang, Z. (2010). Expression of voltage-gated sodium channel alpha subunit in human ovarian cancer. *Oncology Reports*, Vol. 23, pp. 1293-1299.
- Gavrilova, O., Schönherr, K., Gessner, G., Schönherr, R., Klapperstück, T., Wohlrab, W. & Heinemann, S. (2002). Effects of imipramine on ion channels and proliferation of IGR1 melanoma cells. *Journal of Membrane Biology*. Vol. 188, pp. 137-149.
- Goldin, A., Barchi, R., Caldwell, J., Hofmann, F., Howe, J., Hunter, J., Kallen, R., Mandel, G., Meisler, M., Netter, Y., Noda, M., Tamkun, M., Waxman, S., Wood, J. & Catterall, W. (2000). Nomenclature of voltage-gated sodium channels. *Neuron*, Vol. 28, pp. 365-368.
- Gómez-Varela, D., Zwick, E., Knötgen, H., Hettmann, T., Ossipov, D., Weseloh, R., Contreras, C., Rothe, M., Stühmer, W. & Pardo, L. (2007). Monoclonal antibody blockade of the human Eag1 potassium channel function exerts antitumor activity. *Cancer Research*, Vol. 67, pp. 7343-7349.
- Green, D. & Reed, J. (1998). Mitochondria and apoptosis. *Science*, Vol. 281, pp. 1309-1312.
- Han, X., Wang, F., Yao, W., Xing, H., Weng, D., Song, X., Chen, G., Xi, L., Zhu, T., Zhou, J., Xu, G., Wang, S., Meng, L., Ladecola, C., Wang, G., & Ma, D. (2007). Heat shock proteins and p53 play a critical role in K⁺ channel-mediated tumor cell proliferation and apoptosis. *Apoptosis*, Vol. 12, pp. 1837-1846.
- Hanada, T., Lin, L., Chandy, K., Oh, S., & Chishti, A., (1997). Human homologue of the *Drosophila Discs Large* tumor suppressor binds to p56lck tyrosine kinase and Shaker type Kv1.3 potassium channel in T lymphocytes. *Journal of Biological Chemistry*, Vol. 272, pp. 26899-26904.
- Hanahan, D. & Weinberg, RA. (2011). Hallmarks of cancer: the next generation. *Cell*, Vol. 144, pp. 646-674.
- Hemmerlein B., Weseloh R., Mello F., Knötgen H., Sánchez A., Rubio M., Martin S., Schliephacke T., Jenke M., Joachim H., Stühmer W., & Pardo L., (2006). Overexpression of Eag1 potassium channels in clinical tumours. *Molecular Cancer*, Vol. 5.
- Hernández, E., Ortiz, C., Marquina, B., Medina, I., Alfaro, A., Berumen, J., Rivera, M. & Gomora, J. (2011). Over expression of Na (V) 1.6 channels is associated with the invasion capacity of human cervical cancer. *International Journal of Cancer*. DOI: 10.1002/ijc.26210
- House, C., Vaske, C., Schwartz, A., Obias, V., Frank, B., Luu, T., Sarvazyan, N., Irby, R., Strausberg, R., Hales, T., Stuart, J. & Lee, N. (2010). Voltage-Gated Na⁺ Channel SCN5A Is a Key Regulator of a Gene Transcriptional Network That Controls Colon Cancer Invasion. *Cancer Research*, Vol. 70., pp. 6957-6967.
- Huang, J.B., Kindzelskii, A.L., Clark, A.J. and Petty, H.R. 2004. Identification of channels promoting calcium spikes and waves in HT1080 tumor cells: Their apparent roles in cell motility and invasion. *Cancer Research*, Vol. 64, pp. 2482-2489.
- Hübner, C. & Jentsch, T. (2002). Ion channel diseases. *Human Molecular Genetics*, Vol. 11, pp. 2435-2445.

- Ikuma, M., Binder, H.J., & Geibel, J. (1998). Role of apical H-K exchange and basolateral K channel in the regulation of intracellular pH in rat distal colon crypt cells. *Journal of Membrane Biology*, Vol. 166, December, pp. 205–212.
- Jager, H., Dreker, T., Buck, A., Giehl, K., Gress, T., & Grissmer, S. (2004). Blockage of intermediate-conductance Ca^{2+} -activated K^{+} channels inhibit human pancreatic cancer cell growth in vitro. *Molecular Pharmacology*, Vol. 65, pp. 630–638.
- Jentsch, T., Stein, V., Weinreich, F., & Zdebik, A. (2002). Molecular Structure and Physiological Function of Chloride Channels. *Physiological Reviews*, Vol. 82, pp. 503–568.
- Jin, S., & Levine, A. (2001). The p53 functional circuit. *Journal of Cell Science*, Vol. 114, pgs. 4139–4140.
- Kagan, A., Melman, Y., Krumerman A., & McDonald, T., (2002). 14-3-3 amplifies and prolongs adrenergic stimulation of HERG K^{+} channel activity. *EMBO Journal*, Vol. 21, pp. 1889–1898.
- Kamb, A. et al. (2007). Why is cancer drug discovery so difficult? *Nature Reviews Drug Discovery*. 6:115-120.
- Kim, C., Um, S., Kim, T., Kim, E., Park, T., Kim, S., Namkoong, S., & Park, J. (2000). Regulation of cell growth and HPV genes by exogenous estrogen in cervical cancer cells. *International Journal of Gynecological Cancer*, Vol. 10, pp. 157-164.
- Kunzelmann, K., 2005. Ion channels and cancer. *Journal of Membrane Biology*, Vol. 205, pags. 159-173.
- Lang, F., Busch, G., Ritter, M., Volkl, H., Waldegger, S., Gulbins, E., & Haussinger, D. (1998). Functional significance of cell volume regulatory mechanisms. *Physiological Reviews*, Vol. 78, pp. 247–306.
- Lang, F., Gulbins, E., Szabo, I., Lepple, A., Huber, S., Duranton, C., Lang, K., Lang, P., & Wieder, T. (2004). Cell volume and the regulation of apoptotic cell death. *Journal of Molecular Recognition*, Vol. 17, pp. 473–480.
- Lang, F. (2007). Mechanisms and significance of cell volume regulation. *Journal of the American College of Nutrition*, Vol. 26, No. 5, Oct., (5 Suppl):613S-623S.
- Lang, P., Warskulat, U., Heller, B., Huang, D., Grenz, A., Myssina, S., Duszenko, M., Lang, F., Haussinger, D., Vallon, V., & Wieder, T. (2003). Blunted apoptosis of erythrocytes from taurine transporter deficient mice. *Cell Physiology Biochemistry*, Vol. 13, pp. 337–346.
- Lastraioli, E., Guasti, L., Crociani, O., Polvani, S., Hofmann, G., Witchel, H., Bencini, L., Calistri, M., Messerini, L., Scatizzi, M., Moretti, R., Wanke, E., Olivotto, M., Mugnai, G. & Arcangeli, A. (2004). *herg1* gene and HERG1 protein are overexpressed in colorectal cancers and regulate cell invasion of tumor cells. *Cancer Research*. Vol. 15, pp. 606-611.
- Lepple, A., Berweck, S., Bohmig, M., Leo, C., Meyling, B., Garbe, C., & Wiederholt, M. (1996). K^{+} channels and the intracellular calcium signal in human melanoma cell proliferation. *Journal of Membrane Biology*, Vol. 151, pp. 149–157
- Levite, M., Cahalon, L., Peretz, A., Hershkovich, R., Sobko, A., Ariel, A., Desai, R., Attali, B., & Lider, O. (2000). Extracellular K^{+} and opening of voltage-gated potassium channels activate T cell integrin function: physical and functional association

- between Kv1.3 channels and b1 integrins. *The Journal of Experimental Medicine*, Vol. 191, pp. 1167–1176.
- Lin, H., Li, Z., Chen, C., Luo, X., Xiao, J., Dong, D., Lu, Y., Yang, B. & Wang, Z. (2011) Transcriptional and Post-Transcriptional Mechanisms for Oncogenic Overexpression of Ether A` Go-Go K⁺ Channel. *PLoS One*, Vol. 6, pg. e20362
- Liu, X., Chang, Y., Reinhart, P., & Sontheimer, H. (2002). Cloning and Characterization of Glioma BK, a Novel BK Channel Isoform Highly Expressed in Human Glioma Cells. *The Journal of Neuroscience*. Vol. 22, pp. 1840–1849.
- Lu, F., Chen, H., Zhou, C., Liu, S., Guo, M., Chen, P., Zhuang, H., Xie, D., & Wu, S. (2008). T-type Ca²⁺ channel expression in human esophageal carcinomas: A functional role in proliferation. *Cell Calcium*, Vol. 43, pp. 49–58
- Mc Ferrin, M. & Sontheimer, H. (2006) A role for ion channels in glioma cell invasion. *Neuron Glia Biology*, Vol. 2, pp. 39–49.
- Mao, J., Chen, L., Xu, B., Wang, L., Wang, W., Li, M., Zheng, M., Li, H., Guo, J., Li, W., Jacob, T. & Wang L. (2009). Volume-activated chloride channels contribute to cell-cycle-dependent regulation of HeLa cell migration. *Biochemical Pharmacology*, Vol. 15, pp. 159–168.
- Marino, A., Iliev, I., Schwalke, M., Gonzalez, E., Marler, K., & Flanagan, C. (1994). Association between cell membrane potential and breast cancer. *Tumour Biology*, Vol. 15, pp. 82–89.
- Mariot, P., Vanoverberghe, K., Lalevee, N., Rossier, M. & Prevarskaya, N. Overexpression of an alpha 1H (Cav3.2) T-type calcium channel during neuroendocrine differentiation of human prostate cancer cells. *The Journal of Biological Chemistry*. Vol. 277, pp. 10824–10833.
- Masi, A., Becchetti, A., Restano, R., Polvani, S., Hofmann, G., Buccoliero, A., Paglierani, M., Pollo, B., Taddei, G., Gallina, P., Di Lorenzo, N., Franceschetti, S., Wanke, E. & Arcangeli, A. (2008) hERG1 channels are overexpressed in glioblastoma multiform and modulate VEGF secretion in glioblastoma cell lines. *Journal of Biological Chemistry*. Vol. 283, pp. 36234–36240.
- Massagué, J. (2004). G1 cell-cycle control and cancer. *Nature*, Vol. 432, pp. 298–306.
- Mattson, M., & Chan, S. (2003). Calcium orchestrates apoptosis. *Nature Cell Biology*, Vol. 5, No. 12, pp. 1041–1043.
- Mello, F., Suarez, G., Stühmer, W. & Pardo, L. (2006). Ether à go-go potassium channel expression in soft tissue sarcoma patients. *Molecular Cancer*. Vol. 5.
- Monet, M., Gkika, D., Lehen'kyi V., Pourtier, A., Abeela, F.V., Bidaux, G., Juvin, J., Rassendren, F., Humez, S. & Prevarskaya, N. (2009). Lysophospholipids stimulate prostate cancer cell migration via TRPV2 channel activation. *Biochimica et Biophysica Acta-Molecular Cell*, Vol. 1793, pp. 528–539.
- Monk, M. & Holding, C. (2001). Human embryogenic genes re-expressed in cancer cells. *Oncogene*, Vol. 20, pp. 8085–8091.
- Moreno, V., Bosch, F., Muñoz, N., Meijer, C., Shah, K., Walboomers, J., Herrero, R. & Franceschi, S. (2002). Effect of oral contraceptives on risk of cervical cancer in women with human papillomavirus infection: The IARC multicentric case-control study. *Lancet*, Vol. 359, pp. 1085–1092.

- Mu. D., Chen, L., Zhang, X., See, L., Koch, C., Yen, C., Tong, J., Spiegel, L., Nguyen, K., Servoss, A., Peng, Y., Pei, L., Marks, J., Lowe, S., Hoey, T., Jan, L., McCombie, W., Wigler, M. & Powers, S. (2003). Genomic amplification and oncogenic properties of the KCNK9 potassium channel gene. *Cancer Cell*, Vol. 3, pp. 297-302.
- Munoz, N., Franceschi, S., Bosetti, C., Moreno, V., Herrero, R., Smith, J., Shah, K., Meijer, C., & Bosch, F. (2002). Role of parity and human papillomavirus in cervical cancer: The IARC multicentric case-control study. *Lancet*, Vol. 359, pp. 1093-1101.
- Nair, H., Luthra, R., Kirma, N., Flowers, L., Evans, D., & Tekmal, R. (2005). Induction of aromatase expression in cervical carcinomas: effects of endogenous estrogen on cervical cancer cell proliferation. *Cancer Research*, Vol 65, pp. 11164-11173.
- Nilius, B., Schwarz, G., & Droogmans, G. (1993). Control of intracellular calcium by membrane potential in human melanoma cells. *American Journal of Physiology*, Vol. 265, pp. C1501-C1510.
- Norbury, C. & Nurse, P. (1992). Animal cell cycles and their control. *Annual Review of Biochemistry*, Vol. 61, pp. 441-470.
- Occhiodoro TL, Bernheim L, Liu JH, Bijlenga P, Sinnreich M, Bader CR, Fischer-Lougheed J: Cloning of a human ether-a-go-go potassium channel expressed in myoblasts at the onset of fusion. *FEBS Lett* 1998, 434:177-182.
- Oguro-Okano M., Griesmann G.E., Wieben, E.D., Slaymaker, S.J., Snutch, T.P. & Lennon, V.A. (1992). Molecular Diversity of neuronal-type calcium channels identified in small cell lung carcinoma. *Mayo Clinic Proceedings*, Vol. 67, pp. 1150-1159.
- O'Grady, S., & Lee, S. (2005). Molecular diversity and function of voltage-gated (Kv) potassium channels in epithelial cells. *International Journal of Biochemistry and Cell Biology*, Vol. 37, pp. 1578-1594.
- Olsen, M., Schade, S., Lyons, S., Amaral, M. & Sontheimer, H. (2003) Expression of voltage-gated chloride channels in human glioma cells. *Journal of Neuroscience*. Vol. 23, pp. 5572-5582.
- Ouadid, H., Roudbaraki, M., Ahidouch, A., Delcourt, P., Prevarskaya, N. (2004a). Cell-cycle-dependent expression of the large Ca^{2+} -activated K^{+} channels in breast cancer cells. *Biochemical and Biophysical Research Communications*. Vol. 316, pp. 244-251.
- Ouadid, H., Roudbaraki, M., Delcourt, P., Ahidouch, A., Joury, N., & Prevarskaya, N. (2004b). Functional and molecular identification of intermediate-conductance Ca^{2+} -activated K^{+} channels in breast cancer cells: association with cell cycle progression. *American Journal of Physiology, Cell Physiology*, Vol. 287, pp. C125-C134.
- Ousingsawat J, Spitzner M, Puntheeranurak S, Terraciano L, Tornillo L, Bubendorf L, Kunzelmann K and Schreiber R. (2007). Expression of voltage-gated potassium channels in human and mouse colonic carcinoma. *Clinical Cancer Research*, Vol. 13, pp. 824-831.
- Panner, A., & Wurster, R. (2006). T-type calcium channels and tumor proliferation. *Cell Calcium*, Vol. 40, pp. 253-259.
- Pardo, L., Camino D., Sánchez A., Alves F., Brüggemann A., Beckh S., & Stühmer W., (1999). Oncogenic potential of EAG K^{+} channels. *EMBO Journal*, Vol. 18, pp. 5540-5547.
- Pardo, L. (2004). Voltage-gated potassium channels in cell proliferation. *Physiology (Bethesda)*, Vol.19, pp. 85-292.


- Pardo, L., Contreras, C., Zientkowska, M., Alves, F., & Stühmer W. (2005). Role of Voltage-gated Potassium Channels in Cancer. *Journal of Membrane Biology*, Vol. 205, pp. 115–124.
- Pardo, L. & Stühmer, W. (2008) Eag1: an emerging oncological target. *Cancer Research*, Vol. 68, pp. 1611–1613.
- Parekh, A., & Penner, R. (1997). Store depletion and calcium influx. *Physiological Reviews*, Vol. 77, pp. 901–930.
- Parekh, A., & Putney JW Jr. (2005). Store-operated calcium channels. *Physiological Reviews*, Vol. 85, pp. 757–810.
- Parihar, A., Coghlan, M., Gopalakrishnan, M. & Shieh, C. (2003). Effects of intermediate-conductance Ca^{2+} -activated K^{+} channel modulators on human prostate cancer cell proliferation. *The European Journal of Pharmacology*. Vol. 471, pp. 157–164.
- Pei, L., Wiser, O., Slavin, A., Mu, D., Powers, S., Jan, L. & Hoey, T. (2003). Oncogenic potential of TASK3 (Kcnk9) depends on K^{+} channel function. *The Proceedings of the National Academy of Sciences USA*. Vol. 100, pp. 7803–7807.
- Per Larsson, K., Hansen, A. & Dissing, S. (2002). The human SH-SY5Y neuroblastoma cell-line expresses a functional P2X7 purinoceptor that modulates voltage-dependent Ca^{2+} channel function. *Journal of Neurochemistry*. Vol. 83, pp. 285–298.
- Pillozzi, S., Brizzi, M., Balzi, M., Crociani, O., Cherubini, A., Guasti, L., Bartolozzi, B., Becchetti, A., Wanke, E., Bernabei, P., Olivotto, M., Pegoraro, L. & Arcangeli, A. (2002). HERG potassium channels are constitutively expressed in primary human acute myeloid leukemias and regulate cell proliferation of normal and leukemic hemopoietic progenitors. *Leukemia*. Vol. 16, pp. 1791–1798.
- Plummer, H., Dhar, M., Cekanova, M. & Schuller, H. (2005). Expression of G-protein inwardly rectifying potassium channels (GIRKs) in lung cancer cell lines. *BMC Cancer*. Vol. 5.
- Prehn, J., Jordan, J., Ghadge, G., Preis, E., Galindo, M., Roos, R., Kriegstein, J., & Miller, R. (1997). Ca^{2+} and reactive oxygen species in staurosporine-induced neuronal apoptosis. *Journal of Neurochemistry*, Vol. 68, pp. 1679–1685.
- Preussat, K., Beetz, C., Schrey, M., Kraft, R., Wölfl, S., Kalff, R. & Patt, S. (2003). Expression of voltage-gated potassium channels Kv1.3 and Kv1.5 in human gliomas. *Neuroscience Letters*. Vol. 346, pp. 33–36.
- Roberts, T.G., Goulart, B.H., Squitieri, L., Stallings, S.C., Halpern, E.F., Chabner, B.A., Gazelle, G.S., Finkelstein, S.N, & Clark J.W. 2004. Trends in the risk and benefits to patients with cancer participating in phase 1 clinical trials. *The Journal of the American Medical Association*, 292, 2130–2140.
- Roger S, Besson P, & Le Guennec JY. (2003). Involvement of a novel fast inward sodium current in the invasion capacity of a breast cancer cell line. *Biochimica et Biophysica Acta*, Vol. 1616, pp. 107–111.
- Roger, S., Potier, M., Vandier, C., Besson, P., & Le Guennec, J. (2006) Voltage-gated sodium channels: new targets in cancer therapy? *Current Pharmaceutical Design*, Vol. 12, pp. 3681–3695
- Roger, S., Rollin, J., Barascu, A., Besson, P., Raynal, P., Iochmann, S., Lei, M., Bougnoux, P., Gruel, Y. & Le Guennec, J. (2007) Voltage-gated sodium channels potentiate the

- invasive capacities of human non-small-cell lung cancer cell lines. *Int. J. Biochem. Cell Biology*, Vol. 39, pp. 774–786.
- Roman, R., Smith, R., Feranchak, A., Clayton, G., Doctor, R. & Fitz, J. (2001). ClC-2 chloride channels contribute to HTC cell volume homeostasis. *American Journal of Physiology - Gastrointestinal and Liver Physiology*. Vol. 280, pp. G344-G353.
- Rosenbaum, S., Tsvieli, R., & Kaspas, R. (1989). Oestrogen stimulates differential transcription of human papillomavirus type 16 in SiHa cervical carcinoma cells. *Journal of General Virology*, Vol. 70, pp. 2227- 2232.
- Shao, X., Wu, K., Guo, X., Xie, M., Zhang, J. & Fan, D. (2008). Expression and significance of HERG protein in gastric cancer. *Cancer Biology and Therapy*. Vol. 7, pp. 45-50.
- Shew, M., McGlennen, R., Zaidi, N., Westerheim, M., Ireland, M., & Anderson, S. (2002). Oestrogen receptor transcripts associated with cervical human papillomavirus infection. *Sexually Transmitted Infections*, Vol. 78, pp. 210-214.
- Schönherr, R., Lober, K., & Heinemann, S. (2000). Inhibition of human ether à go-go potassium channels by Ca^{2+} /calmodulin. *EMBO Journal*, Vol. 19, pp. 3263–3271.
- Schönherr, R. (2005). Clinical Relevance of Ion Channels for Diagnosis and Therapy of Cancer. *Journal of Membrane Biology*, Vol. 205, pp. 175–184.
- Schrey, M., Codina, C., Kraft, R., Beetz, C., Kalff, R., Wölfl, S. & Patt, S. (2002). Molecular characterization of voltage-gated sodium channels in human gliomas. *Neuroreport*. Vol. 13, pp. 2493-2498.
- Shuba, Y., Prevarskaya, N., Lemonnier, L., Van, F., Kostyuk, P., Mauroy, B. & Skryma, R. (2000). Volume-regulated chloride conductance in the LNCaP human prostate cancer cell line. *American Journal of Physiology - Cell Physiology*. Vol. 279, pp. C1144-C1154.
- Shinohara, T., Takahashi, N., Ooie, T., Ichinose, M., Hara, M., Yonemochi, H., Saikawa, T., & Yoshimatsu, H. (2004). Estrogen inhibits hyperthermia-induced expression of heat-shock protein 72 and cardioprotection against ischemia/reperfusion injury in female rat heart. *Molecular Cell Cardiology*, Vol. 37, pp. 1053–1061.
- Smith, G., Tsui, H., Newell, E., Jiang, X., Zhu, X., Tsui, F., Schlichter, L. (2002). Functional up-regulation of HERG K^{+} channels in neoplastic hematopoietic cells. *The Journal of Biological Chemistry*. Vol. 277, pp. 18528-18534.
- Spasova, M., Soboloff, J., He, L., Hewavitharana, T., Xu, W., Venkatachalam, K., van Rossum, D., Patterson, R., & Gill, D. (2004). Calcium entry mediated by SOCs and TRP channels: variations and enigma. *Biochimica et Biophysica Acta*, Vol. 1742, pp. 9–20.
- Surprenant, A., Rassendren, F., Kawashima, E., North, R. A., & Buell, G. (1996). The cytolytic P2Z receptor for extracellular ATP identified as a P2X receptor (P2X7). *Science*. Vol. 272, pp. 735–738
- Tsavalier, L., Shapero, M., Morkowski, S. & Laus, R. (2001). Trp-p8, a novel prostate-specific gene, is up-regulated in prostate cancer and other malignancies and shares high homology with transient receptor potential calcium channel proteins. *Cancer Research*. Vol. 61, pp. 3760-3769.

- Uysal, P. & Djamgoz, M. (2007) Epidermal growth factor potentiates in vitro metastatic behavior of human prostate cancer PC-3 M cells: involvement of voltage-gated sodium channel. *Molecular Cancer*, Vol. 6, pg., 76.
- Vanden, F., Shuba, Y., Roudbaraki, M., Lemonnier, L., Vanoverberghe, K., Mariot, P., Skryma, R. & Prevarskaya, N. (2003). Store-operated Ca^{2+} channels in prostate cancer epithelial cells: function, regulation, and role in carcinogenesis. *Cell Calcium*. Vol. 33, pp. 357-373.
- Vermeulen, K., Van, D. & Berneman Z. (2003). The cell cycle: a review of regulation, deregulation and therapeutic targets in cancer. *Cell Proliferation*, Vol. 36, pp. 131-149.
- Walboomers JMM, Jacobs MV, Manos MM, Bosch FX, Kummer JA, Shah KV, Snijders PJF, Peto J, Meijer CJLM, & Munoz N (1999). Human papillomavirus is a necessary cause of invasive cervical cancer worldwide *Journal of Pathology*, Vol. 189, pp. 12-19.
- Wang, H., Zhang, Y., Cao, L., Han, H., Wang, J., Yang, B., Nattel, S., & Wang, Z., (2002). HERG K^{+} channel, a regulator of tumor cell apoptosis and proliferation. *Cancer Research*, Vol. 62, pp. 4843-4848.
- Wang, Z. (2004). Roles of K^{+} channels in regulating tumour cell proliferation and apoptosis. *Pflügers Archiv: European Journal of Physiology*, Vol. 448, pp. 274-286.
- Wang, Q., Li, X., Wang, L., Feng, Y. H., Zeng, R., & Gorodeski, G. (2004). Antiapoptotic effects of estrogen in normal and cancer human cervical epithelial cells. *Endocrinology*, Vol. 145, pp. 5568-5579.
- Wang, X., Nagaba, Y., Cross, H., Wrba, F., Zhang, L. & Guggino, S. (2000). The mRNA of L-type calcium channel elevated in colon cancer: protein distribution in normal and cancerous colon. *American Journal of Pathology*. Vol. 157, pp. 1549-1562.
- Whitfield, J., Bird, R., Chakravarthy, B., Isaacs, R., & Morley, P. (1995). Calcium-cell cycle regulator, differentiator, killer, chemopreventor, and maybe, tumor promoter. *Journal of Cellular Biochemistry*, Vol 22, pp. 74-91.
- Wiley, J., Dao, L., Gu, B., Sluyter, R., Shemon, A., Li, C., Taper, J., Gallo, J. & Manoharan, A. (2002). A loss-of-function polymorphic mutation in the cytolytic P2X7 receptor gene and chronic lymphocytic leukaemia: a molecular study. *Lancet*. Vol. 359, pp. 1114-1119.
- Wonderlin, W., & Strobl, J. (1996). Potassium channels, proliferation and G1 progression. *Journal of Membrane Biology*, Vol. 154, pp. 91-107.
- Yamamura, H., Ugawa, S., Ueda, T., Morita, A. & Shimada, S. (2008). TRPM8 activation suppresses cellular viability in human melanoma. *American Journal of Physiology - Cell Physiology*. Vol. 295, pp. C296-C301.
- Yang, S.Y., Zhang, J.J.L. & Huang, X.Y. (2009). Orail and STIM1 are critical for breast tumor cell migration and metastasis. *Cancer Cell*, Vol. 15, pp. 124-134.
- Yu, F. & Catterall, W. (2003). Overview of the voltage-gated sodium channel family. *Genome Biology*, Vol. 4, pp. 207.
- Yuan, P., Leonetti, M., Pico, A., Hsiung, Y., & MacKinnon, R. (2010). Structure of the Human BK Channel Ca^{2+} - Activation Apparatus at 3.0 Å Resolution. *Science*. Vol. 329, pp. 182-186.

- Zamzami, N., & Kroemer, G. (2005). p53 in apoptosis control: an introduction. *Biochemical and Biophysical Research Communications*, Vol. 331, pp. 685–687.
- Zhuang, L., Peng, J., Tou, L., Takanaga, H., Adam, R., Hediger, M. & Freeman, M. (2002). Calcium-selective ion channel, CaT1, is apically localized in gastrointestinal tract epithelia and is aberrantly expressed in human malignancies. *Laboratory Investigations*. Vol. 82, pp. 1755-1764.
- zur Hausen H (2002) Papillomaviruses and cancer: from basic studies to clinical applications *Nature Reviews Cancer*, Vol. 2, pp. 342-350.
- Zylicz, M., King, F.W., & Wawrzynow, A. (2001). Hsp70 interactions with the p53 tumour suppressor protein. *EMBO Journal*, Vol. 20, pp. 4634–4638

IntechOpen



Human Papillomavirus and Related Diseases - From Bench to Bedside - Research aspects

Edited by Dr. Davy Vanden Broeck

ISBN 978-953-307-855-7

Hard cover, 406 pages

Publisher InTech

Published online 25, January, 2012

Published in print edition January, 2012

Cervical cancer is the second most prevalent cancer among women worldwide, and infection with Human Papilloma Virus (HPV) has been identified as the causal agent for this condition. The natural history of cervical cancer is characterized by slow disease progression, rendering the condition, in essence, preventable and even treatable when diagnosed in early stages. Pap smear and the recently introduced prophylactic vaccines are the most prominent prevention options, but despite the availability of these primary and secondary screening tools, the global burden of disease is unfortunately still very high. This book will focus on epidemiological and fundamental research aspects in the area of HPV, and it will update those working in this fast-progressing field with the latest information.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Ana Ramírez and Javier Camacho (2012). The Human Papilloma Virus – Ion Channel Link in Cancer: An Alternative Opportunity for Diagnosis and Therapy, Human Papillomavirus and Related Diseases - From Bench to Bedside - Research aspects, Dr. Davy Vanden Broeck (Ed.), ISBN: 978-953-307-855-7, InTech, Available from: <http://www.intechopen.com/books/human-papillomavirus-and-related-diseases-from-bench-to-bedside-research-aspects/the-human-papilloma-virus-ion-channel-link-in-cancer-an-alternative-opportunity-for-diagnosis-and-th>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](https://creativecommons.org/licenses/by/3.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen