We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Proximal Tibiofibular Joint in Knees with Arthroplasty

Hakan Boya

*Başkent University, Faculty of Medicine,
Department of Orthopaedics and Traumatology,
Zübeyde Hanım Hospital, İzmir
Turkey*

1. Introduction

Because proximal tibiofibular joint (PTFJ) is a diarthrodial joint encased in a synovial-lined articular capsule, it is possible to observe disorders at synovial joints, such as traumatic dislocation, osteoarthritis, inflammatory arthritis, ganglion cysts, pigmented villonodular synovitis, and infection.

Because of its close proximity to the knee joint, PTFJ may be the cause of lateral knee pain. This issue is important, especially in knees with arthroplasty. However, knee arthroplasty may exacerbate PTFJ. Consequently, the joint should be examined in detail before and after knee arthroplasty operations.

2. Embryology and postnatal development

Before 12 weeks of fetal age, PTFJ does not create a cavity (Bozkurt et al., 2003; Resnick et al., 1978). Subsequently, narrow cavities, which may be separated from the lateral femorotibial joint by a small amount of loose fibrous or areolar tissue, are apparent (Resnick et al., 1978). Subsequent development of the PTFJ includes the formation of articular cartilage, synovial tissue, synovial recesses, and a fibrous capsule (Resnick et al., 1978). Ossification usually begins in the proximal tibia within the first three months following birth. The tibiofibular joint morphology has considerable morphologic variation, and the joint may communicate with the knee joint (Ogden, 1984).

3. Anatomy

PTFJ is a diarthrodial joint between the lateral tibial condyle and fibular head; it is located posterolaterally on proximal tibia (Bozkurt et al., 2003). PTFJ has certain characteristics of synovial joints, such as synovial membrane, hyaline cartilage, and a fibrous capsule (Bozkurt et al., 2003; Resnick et al., 1978). The stability of the joint is provided by anterosuperior and posteriosuperior capsular ligaments (Ogden, 1974; Resnick et al., 1978; Gray, 1977). The tendon of the biceps femoris muscle inserts into the anterior part of the fibular head and enforces the anterosuperior ligament of the joint (Bozkurt et al., 2003; Marshall et al., 1972).

Communication between the knee joint and PTFJ is reported to be 10-63% (Bozkurt et al., 2003; De Franca, 1992; Eichenblat et al., 1983, Veth et al., 1984). Communication between the

proximal tibiofibular and the knee joint occurs via the subpopliteal recess-associated defect in the posterior ligament of the fibular head (Dirim et al., 2008). The relation of the defect to trauma or developmental deficiency is unclear (Dirim et al., 2008).

Two types of PTFJ were defined according to joint line inclination; oblique (inclination $> 20^\circ$) and horizontal (inclination $< 20^\circ$) (Ogden, 1974). Moreover, planar, trochoid and double trochoid types of PTFJ have been reported (Espregueira-Mendes & Vieira, 2006).

Fig. 1. Horizontal (a) and oblique (b) type PTFJ

3.1 Functional anatomy

Primary functions of PTFJ include the following:

1. Dissipation of torsional stresses applied at the ankle. With external rotation of the fibula about its longitudinal axis during dorsiflexion of the ankle joint, the proximal fibula rotates a few degrees externally (Barnett & Napier, 1952). The amount of external rotation is much greater in horizontal-type PTFJs (Barnett & Napier, 1952).
2. Dissipation of lateral tibial bending movements. Tensile and torsional forces influence the proximal-middle fibula in contrast to the distal fibula, which is affected by compressive forces (Ogden, 1974).
3. Transmitting axial loads in weight-bearing. Approximately one-sixth of the static load is applied at the ankle being transmitted to the PTFJ (Lambert, 1971).

4. Pathologies of PTFJ

It is possible to observe all disease at the PTFJ, similar to other synovial joints. Pathologies of this joint include primary osteoarthritis (Bozkurt et al., 2004; Öztuna et al., 2003; Özcan et al., 2009), trauma (Ogden, 1974; Resnick et al., 1978), infection, and inflammatory arthritis (Resnick & Niwayama, 1995), synovial osteochondromatosis (Bozkurt et al., 2007; Heybeli et al., 2009; Weiss et al., 1975), neoplasms (Forster et al., 2007), ganglion cysts (Miskovsky et al., 2004; Mortazavi et al., 2006; Ward & Echardt, 1994), and pigmented villonodular synovitis (Ryan et al., 2004).

4.1 PTFJ in knees with severe primary osteoarthritis

PTFJ can be affected by primary osteoarthritis (Öztuna et al., 2003). The degree of osteoarthritis of the proximal tibiofibular joint strongly correlates with the degree of arthritis

in tibiofemoral joints (TFJ) that have severe degenerative joint disease (Boya et al., 2008). Inflammatory enzymes passing between the joint spaces through possible anatomical communication between the TFJ and PTFJ may contribute to the advancement of arthritis in the respective compartments (Boya et al., 2008; Bozkurt et al., 2003). As with other joints, osteophytes, subchondral cysts, subchondral sclerosis, and joint-space narrowing are typical imaging findings (Forster et al., 2007).

Although primary degenerative disease of the PTFJ is commonly associated with primary degenerative disease of the knee joint, radiographic findings of the PTFJ in patients with severe degenerative knee osteoarthritis and varus misalignment do not correlate with clinical findings (Özcan et al., 2009).

4.2 PTFJ in other pathologies

Various neoplasms can affect the proximal tibiofibular joint, including osteochondroma, osteblastoma, osteosarcoma, and nerve sheath tumors (Schwannomas and neurofibromas) (Forster et al., 2007).

Tuberculosis lesions at the fibular head can destroy PTFJ and mimic tumoral lesions (Abdelwahab et al., 2003-2004).

Synovial chondromatosis is a chronic, progressive disease of the synovial tissue in which free chondral loose bodies are formed after metaplasia (Bozkurt et al., 2007; Heybeli et al., 2009; Weiss et al., 1975).

A ganglion is a tumorlike, cystic lesion that arises from the joint, tendon sheath, or muscle (Miskovsky et al., 2004). It is a rare pathology at the PTFJ but can cause three different pathologies: asymptomatic mass, symptomatic fluctuant mass, and mass with peroneal nerve dysfunction (Forster et al., 2007).

PTFJ is affected similarly to other synovial joints in rheumatoid arthritis. Peroneal nerve dysfunction due to subluxation, dislocation of dextruded PTFJ is a pathologic entity of the joint with RA (Ishikawa & Hirohata, 1984). Moreover, it is possible to observe radiological deterioration of the PTFJ in patients with ankylosing spondylitis (Hong et al., 2009).

Pigmented villonodular synovitis (PVNS) is an uncommon proliferative disease of the synovium, which is usually monoarticular, presenting as chronic monoarthritis of the knee (Forster et al., 2007). PVNS is characterized by synovial hypertrophy with diffuse or focal hemosiderin deposition in the joint (Ryan et al., 2004). The disease can affect the PTFJ similarly to other synovial joints.

5. Importance of PTFJ in knees with arthroplasty

The PTFJ can be considered the fourth compartment of the knee joint because of its communication with the knee joint cavity (Bozkurt et al., 2003). PTFJ can be a source of lateral knee pain because of its pathologies. However, frequently it is overlooked because of its lack of emphasis in the literature (Forster et al., 2007). For this reason, the PTFJ should be carefully evaluated for osteoarthritis in patients being considered for a total knee arthroplasty operation. If it is overlooked as an etiology of a patient's lateral knee pain, pain from the diseased PTFJ may continue post-operatively. Furthermore, pathologies of the PTFJ may cause peroneal nerve dysfunction in patients with knee arthroplasty (Gibbon et al., 1999). PTFJ stability is important in cases with knee arthroplasty, especially in patients with rheumatoid arthritis. In those patients, the PTFJ may become unstable. Because of proximal movement of the fibular head, it may impinge to extruded bone cement under the tibial base

plate posterolaterally (Otani et al., 1998). This possibility should be considered during cementing of the tibial base plate; posterolateral cement excursion should be avoided in patients with inflammatory arthritis.

PTFJ pathologies can affect the arthroplasty results. Conversely, knee arthroplasty can produce PTFJ pathologies. It is possible to inadvertently destroy the PTFJ during an erroneous lower-level tibial cut; this may produce joint-related symptoms after a knee arthroplasty operation. Knees with aseptic loosening of the prosthesis produce inflammatory mediators and polyethylene particles can migrate to the PTFJ via communication between the knee and proximal tibiofibular joints. This can result in deterioration of the PTFJ and subsequent symptoms (Crawford et al., 1998).

6. Conclusion

The degree of osteoarthritis of the proximal tibiofibular joint strongly correlates with the degree of arthritis in tibiofemoral joints that have severe degenerative joint disease. Although primary degenerative disease of the PTFJ is commonly associated with primary degenerative disease of the knee joint, radiographic findings of the PTFJ in patients with severe degenerative knee osteoarthritis and varus misalignment do not correlate with clinical findings. It is possible to observe all disease at the PTFJ, similar to other synovial joints. Because of its close proximity to the knee joint, PTFJ may be the cause of lateral knee pain. This issue is important, especially in knees with arthroplasty. However, knee arthroplasty may exacerbate PTFJ. Consequently, the joint should be examined in detail before and after knee arthroplasty operations.

7. References

- Abdelwahab, IF., Poplaw, S., Abdul-Quader, M. & Naran, D. (2003). Tuberculous pseudotumor of the proximal end of the fibula. A case report. *Bull Hosp Jt Dis*, Vol.61, No.3-4, pp. 145-147, ISSN 0883-9344
- Barnett, CH. & Napier, JR. (1952) The axis of rotation of the ankle joint in man. Its influence upon the form of the talus and mobility of the fibula. *J Anat*, Vol.86, No.1, (January 1952), pp. 1-9, ISSN 1136-4890
- Boya, H., Ozcan, O. & Oztekin, HH. (2008). Radiological evaluation of the proximal tibiofibular joint in knees with severe primary osteoarthritis. *Knee Surg Sports Traumatol Arthrosc*, Vol.16, No.2, (February 2008), pp. 157-159, ISSN 0942-2056
- Bozkurt, M., Yilmaz, E., Atlihan, D., Tekdemir, I., Havitçioğlu, H. & Günel, I. (2003). The proximal tibiofibular joint: An anatomic study. *Clin Orthop Relate Res*, Vol.406, No.1 (January 2003), pp. 136-140, ISSN 0009-921X
- Bozkurt, M., Yilmaz, E., Akseki, D., Havitçioğlu, H. & Günel, I. (2004). The evaluation of the proximal tibiofibular joint for patients with lateral knee pain. *The Knee*, Vol.11, No.4, (August 2004), pp. 307-312, ISSN 0968-0160
- Bozkurt, M., Uğurlu, M., Doğan, M. & Tosun, N. (2007) Synovial chondromatosis of four compartments of the knee: medial and lateral tibiofemoral spaces, patellofemoral joint and proximal tibiofibular joint. *Knee Surg Sports Traumatol Arthrosc*, Vol.15, No.6, (June 2007), pp. 753-755, ISSN 0942-2056

- Crawford, R., Sabokbar, A., Wulke, A., Murray, DW. & Athanasou, NA. (1998). Expansion of an osteoarthritic cyst associated with wear debris: a case report. *J Bone Joint Surg Br*, Vol.80, No.6, (November 1998), pp. 990-993, ISSN 0301-620X
- De Franca, GG. (1992). Proximal tibiofibular joint dysfunction and chronic knee and low back pain. *J ManipPhysiol Ther*, Vol.15, No.6, (July-August 1992), pp. 382-387, ISSN 0161-4754
- Dirim, B., Wangwinyuvirat, M., Frank, A., Cink, V., Pretterklieber, ML., Pastore, D. & Resnick, D.(2008). Communication between the proximal tibiofibular joint and knee via the subpopliteal recess: MR arthrography with histologic correlation and stratigraphic dissection. *AJR Am J Roentgenol*, Vol.191, No.2, (August 2008), pp. W44-51, ISSN 1546-3141
- Eichenblat, M. & Nathan, H. (1983). The proximal tibiofibular joint. An anatomical study with clinical and pathological considerations. *Int Orthop*, Vol.7, No.1, pp. 31-39, ISSN 0341-2695
- Espregueira-Mendes, JD. & da Silva, MV. (2006). Anatomy of the proximal tibiofibular joint. *Knee Surg SportsTraumatol Arthrosc*, Vol.14, No.3, (March 2006), pp. 241-249, ISSN 0942-2056
- Forster, BB., Lee, JS., Kelly, S., O'Dowd, M., Munk, PL., Andrews, G. & Marchinkow, L. (2007) Proximal tibiofibular joint: an often-forgotten cause of lateral knee pain. *AJR Am J Roentgenol*, Vol.188, No.4, (April 2007) pp. W359-366, ISSN 1546-3141
- Gibbon, AJ., Wardell, SR. & Scott, RD. (1999). Synovial cyst of the proximal tibiofibular joint with peroneal nerve compression after total knee arthroplasty. *J Arthroplasty*, Vol.14, No.6, (September 1999), pp. 766-768, ISSN 0883-5403
- Gray, H. (1977). *The classic Collector's Edition. Gray's Anatomy*, Churchill Livingstone Inc, ISBN 978-051-7223-65-9, New York, USA
- Heybeli, N., Ozcan, M., Copuroğlu, C. & Yalniz, E. (2009). Isolated synovial chondromatosis of the proximal tibiofibular joint. *Acta Orthop Traumatol Turc*, Vol.43, No. 5, (November-December 2009), pp. 448-452, ISSN 1017-995X
- Hong, HP., Chung, HW., Choi, BK., Yoon, YC. & Choi, SH. (2009) Involvement of the proximal tibiofibular joint in ankylosing spondylitis. *Acta Radiol*, Vol.50, No.4, (May 2009), pp. 418-422, ISSN 0284-1851
- Ishikawa, H. & Hirohata K. (1984). Bilateral peroneal nerve palsy secondary to posterior dislocation of the proximaltibiofibular joint in rheumatoid arthritis. *Rheumatol Int*, Vol.5, No.1, pp. 45-47, ISSN 0172-8172
- Lambert KL. (1971). The weight-bearing function of the fibula. A strain gauge study. *J Bone Joint Surg Am*, Vol.53, No.3, (April 1971), pp. 507-513, ISSN 0021-9355
- Marshall, JG., Girgis, FG. & Zelko, RR. (1972). The Biceps Femoris tendon and its functional significance. *J Bone Joint Surg Am*, Vol.54, No.7, (October 1972), pp. 1444-1450, ISSN 0021-9355
- Miskovsky, S., Kaeding, C. & Weis, L. (2004) Proximal tibiofibular joint ganglion cysts: excision, recurrence, and joint arthrodesis. *Am J Sports Med*, Vol.32, No.4, (June 2004), pp.1022-1028, ISSN 0363-5465
- Mortazavi, SM., Farzan, M. & Asadollahi, S. (2006). Proximal tibiofibular joint synovial cyst-one pathology with three different presentations. *Knee Surg Sports Traumatol Arthrosc*, Vol.14, No.9, (September 2006), pp. 875-879, ISSN 0942-2056

- Ogden, JA. (1974). The anatomy and function of the proximal tibiofibular joint. *Clin Orthop*, Vol.101, No.6, (June 1974), pp. 186-191, ISSN 0009-921X
- Ogden, JA. (1984). Radiology of postnatal skeletal development. IX. Proximal tibia and fibula. *Skeletal Radiol*, Vol.11, No.3, pp. 169-177, ISSN 0364-2348
- Otani, T., Fujii, K., Ozawa, M., Kaechi, K., Funaki, K., Matsuba, T. & Ueno, H. (1998). Impingement after total knee arthroplasty caused by cement extrusion and proximal tibiofibular instability. *J Arthroplasty*, Vol.13, No.5, (August 1998), pp. 589-591, ISSN 0883-5403
- Özcan, O., Boya, H. & Oztekin, HH. (2009) Clinical evaluation of the proximal tibiofibular joint in knees with severe tibiofemoral primary osteoarthritis. *The Knee*, Vol.16, No.4, (August 2009), pp. 248-250, ISSN 0968- 0160
- Öztuna, V., Yıldız, A., Özer, C., Milcan, A., Kuyurtar, F. & Turgut, F. (2003). Involvement of the proximal tibiofibular joint in osteoarthritis of the knee. *The Knee*, Vol.10, No.4, (December 2003), pp. 347-349, ISSN 0968-0160
- Resnick, D., Newell, JD., Guerra, J Jr., Danzing, LA., Niwayama, G. & Goergen, TG. (1978) Proximal tibiofibular joint: Anatomic-pathologic-radiographic correlation. *Am J Roentgenol*, Vol.131, No.1, (July 1978), pp. 133-138, ISSN 0361-803
- Resnick, D. & Niwayama, G. (1995). Anatomy of individual joints, In: *Diagnosis of bone and joint disorders*, Resnick, D. & Niwayama, G. pp. 741-750, WB Saunders, ISBN 072165066X, 9780721650661, Philadelphia
- Ryan, RS., Louis, L., O'Connell, JX. & Munk, PL. (2004). Pigmented villonodular synovitis of the proximal tibiofibular joint. *Australas Radiol*, Vol.48, No.4, (December 2004), pp. 520-522, ISSN 0004-8461
- Ward, WG. & Eckardt, JJ. (1994) Ganglion cyst of the proximal tibiofibular joint causing anterior compartment syndrome. A case report and anatomical study. *J Bone Joint Surg Am*, Vol.76, No.10, (October 1994), pp. 1561- 1564, ISSN 0021-9355
- Weiss, C., Averbuch, PF., Steiner, GC. & Rusoff, JH. (1975). Synovial chondromatosis and instability of the proximal tibiofibular joint. *Clin Orthop Relat Res*, Vol.108, No.5, (May 1975), pp. 187-190, ISSN 0009-921X
- Veth, RP., Kingma, LM. & Nielsen, HK. (1984). The abnormal proximal tibiofibular joint. *Arch Orthop Trauma Surg*, Vol.102, No.3, pp. 167-171, ISSN 0936-8051

IntechOpen

Recent Advances in Hip and Knee Arthroplasty

Edited by Dr. Samo Fokter

ISBN 978-953-307-841-0

Hard cover, 452 pages

Publisher InTech

Published online 27, January, 2012

Published in print edition January, 2012

The purpose of this book is to offer an exhaustive overview of the recent insights into the state-of-the-art in most performed arthroplasties of large joints of lower extremities. The treatment options in degenerative joint disease have evolved very quickly. Many surgical procedures are quite different today than they were only five years ago. In an effort to be comprehensive, this book addresses hip arthroplasty with special emphasis on evolving minimally invasive surgical techniques. Some challenging topics in hip arthroplasty are covered in an additional section. Particular attention is given to different designs of knee endoprostheses and soft tissue balance. Special situations in knee arthroplasty are covered in a special section. Recent advances in computer technology created the possibility for the routine use of navigation in knee arthroplasty and this remarkable success is covered in depth as well. Each chapter includes current philosophies, techniques, and an extensive review of the literature.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Hakan Boya (2012). Proximal Tibiofibular Joint in Knees with Arthroplasty, Recent Advances in Hip and Knee Arthroplasty, Dr. Samo Fokter (Ed.), ISBN: 978-953-307-841-0, InTech, Available from:

<http://www.intechopen.com/books/recent-advances-in-hip-and-knee-arthroplasty/proximal-tibiofibular-joint-in-knees-with-arthroplasty>

INTech
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](https://creativecommons.org/licenses/by/3.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen