We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Soil Phosphorus Tests and Transformation Analysis to Quantify Plant Availability: A Review

Z.M. Zheng^{1,2} and T.Q. Zhang²

¹*School of Environmental Sciences,
University of Guelph, Guelph, Ontario,*

²*Greenhouse and Processing Crops Research Centre,
Agriculture and Agri-Food Canada, Harrow, Ontario,
Canada*

1. Introduction

Phosphorus (P) is regarded as the most important soil nutrient after nitrogen (N) for plant growth and development as it plays key roles in plant metabolism, structure and energy transformation. It has also been recognized as a potential pollutant in waters (Anderson, 1980). The P dynamics in soils and cycling in agro-ecosystems are of increased interest due to its contribution to the current environmental, agronomic and economic issues (Sharpley and Tunney, 2000).

Soil P tests involve extraction of P from soils with chemical or ion-sink extractants followed by a quantification of P in the extracting solution. Soil test is expected to determine the amount of P that can contribute to crop growth or water contamination. From the standpoint of availability to plants, soil P can be divided into functional pools of differing bioavailability (Tiessen et al., 1982). The information on soil P transformation between those pools is useful to predict P bioavailability as well as the risk of P transfer from soil to surface waters. However, soil P transformation has received less attention attributable to the difficulties associated with separation of inorganic P (P_i) and organic P (P_o) fractions and compositional identification of soil P_o pools. Such investigation is currently possible with an improved sequential fractionation procedure and adoption of advanced techniques such as nuclear magnetic resonance spectroscopy (NMR) and synchrotron-based techniques like X-ray absorption near-edge structure (XANES).

In this chapter, soil P tests with chemical and ion-sink methods and the influences of farming practices on P status and P transformation in soils were summarized. The emphasis was on description of the interrelationships between soil P pools of differing bioavailability. The analytical methods to assess P transformation in soils including advanced techniques such as path analysis, modeling and synchrotron-based techniques were described briefly. The limitations of methodologies of soil P tests and P transformation analyses were discussed and findings from our studies were integrated in this context.

2. Soil tests for available P

2.1 Conventional chemical extractions

Available P is the amount of P in soils that can be extracted or mined by plant roots and utilized by plant for its growth and development. It is a quantitative (extensive) parameter and is influenced by the prevailing soil conditions at a particular time and the plant's ability to extract P from soil solutions (Raven and Hossner, 1993; Holford, 1997). However, most people often use available P synonymously with P availability, an intensive parameter that does not reflect the amount or concentration of available P (White and Beckett, 1964).

Availability of P for plant utilization is not a function of its concentration in soil, but rather on the rate of its release from soil surface into soil solution (Abdu, 2006). Available P is composed of soil solution P and is replenished by P that enters the solution by desorption or dissolution of P_i associated with the soil solid phase or by mineralization of P_o (Hedley et al., 1982). The measurement of available P therefore needs to consider both the amount and rate of release of P from the solid phase. Very few appropriate methods have been developed. Isotopic dilution (^{32}P) techniques theoretically permit researchers to quantify the processes of P_o mineralization, dissolution of insoluble minerals and desorption of aggregate P, and could be likely used for this purpose. But errors involved in the measurement of change rates make it difficult to extrapolate the continuing release (or isotopic dilution) rates to a temporal scale corresponding to cropping seasons and growth cycles under field conditions (Tran et al., 1988; Sharpley et al., 1994). Thus, some limitations have to be overcome to give results that have practical application.

The most widely used soil P tests are chemical extractions that use chemical reagents to extract available P from soils. Water was probably the first extractant used to measure P in soils. The small amounts of soil P extracted by water and difficulties related to chemical analysis limit the use of water as an extractant. Bray and Kurtz (1945) used a combination of HCl and NH_4F to remove easily acid soluble P forms, largely Al- and Fe-phosphates. In 1953, Mehlich introduced a combination of HCl and H_2SO_4 acids (Mehlich 1) to extract P from soils in the north-central region of the U.S. In the early 1980s, Mehlich modified his initial soil test and developed a multi-element extractant (Mehlich 3) which is suitable for removing P and other elements in acid and neutral soils (Mehlich, 1984). Olsen et al. (1954) introduced 0.5 M sodium bicarbonate ($NaHCO_3$) solution at a pH of 8.5 to extract P from calcareous, alkaline, and neutral soils. The routine soil P tests may not give insight into the level of plant available P as the chemical reagents may solubilize non-labile P. For instance, the acidic Bray and Mehlich I extractants can dissolve Al- and Fe-phosphates, while Olsen extractant removes dissolved and adsorbed P on calcium carbonate and Fe-oxide surfaces (Mallarino, 1997). Moreover, these chemical extractants are not applicable over all soil types, which underscore the use of them for soil P extractions (Myers et al., 2005). Bray and Mehlich-3 extractants were designed to extract P from non-calcareous soils, whereas the Olsen method was meant for non-acidic soils. Furthermore, those conventional soil P tests derived from mineral soils may not necessarily be applicable for organic soils, although some routine soil P tests are being adopted to make agronomic recommendations in muck soils (Castillo and Wright, 2008; Wright, 2009). This, however, is an "alternative-than-never" choice, at this moment without specific test for organic soils available.

2.2 Ion-sink extractions

The ideal soil P test should be able to extract P in a similar manner as plant roots (Abdu, 2006). Ion sink tests have been developed to simulate how plant roots extract available P from soils. Methodologies were detailed in Sibbesen (1978), Schoenau and Huang (1991), Chardon et al. (1996) and Myers et al. (2005).

The ion-sink concept used in P extraction includes ionic exchange resin membranes, resin bags and FeO coated filter papers or strips. The exchange membrane resin is employed with either the batch or miscible displacement techniques. The batch technique involves the use of a wide soil to solution ratio, which varies the P concentration in the solution and the quantity of desorbed P as the reaction proceeds. If there is inadequate mixing of solution with the ion exchanger, a limited rate of reaction may occur (Sparks, 1985). This may also lead to a change in surface chemistry of the colloids and break down of soil particles (Barrow and Shaw, 1977). With miscible displacement technique, there can be error in dilution that leads to error in interpretation by altering the P concentration of soil (Sparks, 1999), which is more pronounced in colloids having low ion absorbing power (Carski and Sparks, 1985). Also, dispersion of soil colloids may not be fully achieved (Sparks, 1999). Even though anion exchange resin extracts more P than the FeO-coated papers, the additional P extracted may not be plant available (Robinson and Sharpley, 1994). Soil particles can contaminate the FeO-coated papers during shaking, which can lead to error in estimating desorbable P (Uusitalo and Yli-Halla 1999). This can, however, be minimized by the use of CaCl_2 solution as the background electrolyte to minimize soil dispersion (Myers et al., 2005), but reaction with CaCl_2 may reduce the amount of P extracted (Koopmans et al., 2001).

In spite of the aforementioned disadvantages, ion-sink methods, especially the anion exchange membranes, are still regarded as the most effective method of plant available P extraction. In addition to their capability to extract P from a variety of soil types, regardless of soil properties (Sharpley et al., 1994), the ion-sink methods simulate plant roots and extract P from soil without alteration of the soil chemical and physical characteristics (Raven and Hossner, 1993). Further more, the resin membranes can be re-used several times without losing its extracting power (Schoenau and Huang, 1991). This property makes it relatively cheaper than the FeO-coated papers. The compatibility of resins with soil solution chemistry and pH can be achieved by charging the resins with either HCO_3^- or Cl^- (Agbenin and Raij, 2001). The use of HCO_3^- -resin is more advocated than Cl^- -resin, as plant roots accumulate bicarbonate in the rhizosphere, leading to an increase in rhizosphere pH in acid to neutral soils and a decrease in rhizosphere pH in calcareous soils (Sibbesen, 1978), while when Cl^- -resin is used, the Cl^- accumulated in solution inhibits the exchange reaction (Myers et al., 2005).

2.3 Phosphorus determination

Analysis of extracted P is typically done by colorimetry, most notably the Murphy and Riley (1962) method. A specific ion reacts with the color developing reagents to form colorful complex (e.g. blue antimony phospho-molybdate), then light absorption by the formed complex is detected at a specific wavelength. Colorimetric procedures are sensitive, reproducible and enable to automated analysis. In addition, the methods can be accommodated to water samples, digested solutions and extracts (Pierzynski et al., 2008).

Inductively coupled plasma (ICP) spectrophotometry is also now commonly used for P determination, particularly in routine soil P tests offered by public and commercial laboratories. The use of ICP has increased as the use of multi-element soil extractants becomes more popular. Results from ICP are not always directly comparable to those from colorimetric analyses (Pierzynski et al., 2011), as ICP estimates the total P in a solution while the colorimetric procedures measure P that can react with the color developing reagents. Moreover, there are certain limitations that must be considered while evaluating data generated by ICP, such as the matrix effects, spectral interference, etc. (de Boer et al., 1998).

Nuclear magnetic resonance (NMR) is a physical phenomenon based upon the magnetic property of atomic nucleus. It is observed that magnetic nuclei, like ^1H , ^{13}C and ^{31}P , could absorb radio frequency when placed in a magnetic field with a specific strength, described as the resonance of the nucleus. Different atoms in a molecule resonate at different frequencies at a given field strength. This is a powerful method that allows researchers to determine the structure of chemical compounds. The use of solution ^{31}P NMR spectroscopy has allowed us to identify P forms in soils and residual materials, and confirm P forms estimated by commonly used chemical extractants, such as sequential fractionation schemes. This technique has enabled more accurate determination of organic forms of P in soils and residual materials (Zhang et al., 1999; Turner and Leytem, 2008). In addition, the use of synchrotron-based techniques (e.g. XANES) has provided insights into both P_i and P_o forms in soils and residual materials. Descriptions of these approaches were detailed by Beauchemin et al. (2003) and Shober et al. (2006). These analytical advances have been critical in gaining a more detailed understanding of soil P transformation and reaction products following land application of residual materials. This information has helped assess the fate, reactivity, behavior of specific forms of P and the environmental implications of land application of materials such as biosolids and animal manures (Pierzynski et al., 2008).

3. Soil P transformation

3.1 Soil P fractionation

From the standpoint of availability to plants, soil P is divided into three functional fractions each including both P_i and P_o forms: (1) readily labile P pools, (2) moderately labile P pools and (3) sparingly soluble P pools (Tiessen et al., 1982). Due to the technical difficulties involved in isolating specific P compounds from soils, the financial, practical and safety limitations of using P radioisotopes, and limited access to NMR spectroscopy and synchrotron-based techniques, most of field studies on P transformation in soils must rely on alternative methods of measuring changes in soil P fractions. One technique is to sequentially separate soil P into various P_i and P_o fractions based on their bioavailability and biological associations, by first removing labile P, then more stable forms. This sequential extraction method was originally presented by Chang and Jackson (1957) and was later modified by Petersen and Corey (1966) and Williams et al. (1967). Briefly, this soil P sequential procedure uses NH_4Cl to extract labile P_i , NH_4F to dissolve specifically Al-associated P_i followed by NaOH to extract Fe-bound P_i and by dithionite-citrate for reductant-soluble or 'occluded' P_i forms. The HCl dissolved Ca-bound P_i and the final residue is analyzed by Na_2CO_3 fusion for total P. However, the procedure presents many interpretational problems. Since P_i reprecipitates during the fluoride extraction, the separation of Al- and Fe-associated P_i is not reliable and the reductant-soluble or 'occluded' P_i is an ill-defined pool. Furthermore, the P_o fraction is ignored (Williams et al. 1967).

An improved P fractionation scheme was developed by Hedley et al. (1982). This sequential extraction aimed to quantify labile P_i , Ca-associated P_i , Fe- and Al- associated P_i , as well as labile and more stable forms of P_o . With this improved procedure, soil P is initially extracted with an anionic-exchange resin (resin-P), and then with NaHCO_3 ($\text{NaHCO}_3\text{-}P_i/P_o$). Resin-P and $\text{NaHCO}_3\text{-}P$ are assumed to be the readily plant-available fractions and generally regarded as P that is sorbed on carbonates, sesquioxides or crystalline minerals (Mattingly, 1975). Moderately labile P, sorbed on amorphous Fe and Al minerals and the 'protected P' that is occluded or contained within aggregates, is then extracted with NaOH ($\text{NaOH-}P_i/P_o$). The sparingly soluble apatite-type P minerals are extracted with HCl (HCl-P), and the residual P is dissolved by a H_2O_2 (or $\text{K}_2\text{S}_2\text{O}_8$)- H_2SO_4 digestion (residual $\text{H}_2\text{SO}_4\text{-}P$). The $\text{H}_2\text{SO}_4\text{-}P$ fraction, chemically stable and not readily available to plants, may consist of either P_o or P_i , or both. This approach is currently the only one to evaluate both available P_i and P_o in soils with moderate success (Tiessen and Moir, 1993), although modifications to the procedure are often study-specific, for example, the use of de-ionized water instead of anionic-exchange resins to extract readily labile P. It was proven useful for establishing the effects of long-term cropping practices on P_i and P_o fractions and transformation (O'Halloran et al., 1987a; Paniagua et al., 1995; Richards et al., 1995; Tran and N'dayegamiye, 1995; Zhang and Mackenzie, 1997a, 1997c; Zhang et al., 2004; Zheng et al. 2001, 2002, 2004a) and for assessing soil P status in field soils (Simard et al., 1995; Beauchemin and Simard, 2000; Zheng et al., 2004b). It was also valuable for quantifying changes in soil P fractions in short-term incubations (Hedley et al., 1982; Iyamuremye et al., 1996) and from greenhouse experiments (Ivarsson, 1990). In addition, it offers a useful index of the relative importance of P cycling by biological versus geochemical processes in soils at different stages of development (Cross and Schlesinger, 1995).

3.2 Conceptual model of soil P transformation

The P transformation in soils involves complex mineralogical, chemical and biological processes (Fig. 1). The P cycle in soil is a cohesive dynamic system under the influence of long-term chemical transformations and short-term changes due to plant uptake or cropping. The leaching of bases, the removal of carbonates and the increasing Fe and Al activity that accompany the development of soils cause a shift from primary to secondary P_i forms and also influence the stabilization of organic matter and its associated P_o (Walker and Syers, 1976). The abundance and activity of various P_i forms and the turnover of P_o in soils control the replenishment of labile solution P following plant uptake. Surface associated or amorphous P_i replenishes the labile P pool while more stable crystalline species act as a sink as well as long-term reservoir of P, depending on other soil properties such as pH (Murrman and Peech, 1969). In this way, hydroxide or acid extractable P_i ($\text{NaOH-}P_i$ or HCl-P) may act as the quantity factor that buffers the more labile P forms. The P_o may perform a similar function through controlled mineralization-immobilization processes (McGill and Cole, 1981). Soil organic carbon has a major role of promoting processes involved in P transformation, through its contribution as energy source for microbial activity (Stevenson, 1986). Microbes are heavily involved in P transformation in three ways: (1) by decomposition of P_o compounds, with release of available P_i ; (2) by immobilization available P into cellular material; and (3) by promoting the solubilization of fixed or insoluble mineral forms of P, such as through the production of chelating agents (Stevenson, 1986; Frossard et al., 2000).

inadequate to meet crop P requirements (Zhang and MacKenzie, 1997a, Zheng et al., 2001, 2004b). In many cases, the readily labile P increases more when the two P nutrient sources are applied together than when only one is applied (Paniagua et al., 1995). Consequently, the effects of P fertilization on different P pools in soils depend on nutrient sources and rates of P applied as chemical fertilizer or manure (O'Halloran, 1993; Zhang and MacKenzie, 1997a, 1997b; Zheng et al., 2001, 2002).

Long-term organic residues or manure application increases microbial activity and potential mineralization of soil organic matter (N'dayegamiye and Angers, 1990). Consequently, it may induce transformation of soil P_o to P_i fractions and increase the available P in surface and subsurface soil horizons (Sharpley et al., 1984; Tran and N'dayegamiye, 1995, Zheng et al., 2002, 2004a). This mineralization of P_o during the growing season is very important for P availability to plants. Nevertheless, the excessive application of manure or of P-rich organic amendments plays a significant role in decreasing the P sorption capacity in soils (Sharpley et al., 1993; Simard et al., 1995; Beauchemin et al., 1996; Zheng et al., 2001). Decreased P sorption capacities might increase the risk of contamination of the receiving water bodies.

3.3.2 Effects of tillage

The impacts of tillage on P forms and availability have been the subject of many investigations. There was an increase in total and labile P contents in surface soil layer due to no-till operations (Cruse et al., 1983; Weill et al., 1990; Selles et al., 1997). This was attributed to the enhanced microbial activity and mineralization of soil P_o (Follett and Peterson, 1988) or to a lack of soil mixing with fertilizer P thus reducing P fixation by soil colloids (Sweeney, 1993; Selles et al., 1997, 1999). As a result, extractable P in surface soils is increased in no-tillage compared to conventional tillage. Soil P_o is generally increased near the bottom of plow layer by tillage due to the incorporation of crop residues (Dick, 1983, Weill et al., 1990). Nevertheless, O'Halloran et al. (1987a) did not observe a significant difference in the size of soil P fractions except for NaOH- P_o between stubble mulch, bare sod and no-till practices in a grassland soil. On the other hand, Sharpley et al. (1993) found that no-till plots tended to have higher soil moisture content and lower temperature, which resulted in more dissolved P in the runoff than conventionally tilled plots. Blevins et al. (1990) observed that movement of P was reduced using a system of chisel plowing compared to direct seeding or conventional plowing. The chisel plowing was associated with a reduced runoff following the spreading of fertilizers on the soil surface. Therefore, as with fertilization, tillage can alter distribution of P fractions and affect the transformation and transfer of P in soils.

3.3.3 Effects of cropping systems

Cropping systems also have major effects on changes in P fractions in soils. As cropping systems change from less intensive systems (i.e. crop-summer fallow) to more intensive cropping sequences (i.e. continuous wheat and wheat-wheat-fallow), soil P availability and transformation become less predictable due to larger P_i and P_o components from greater residue and litter which are maintained on soil surface (O'Halloran, 1987a, b). Wagar et al. (1986) found a buildup of P_o occurred with and without added P fertilizer when cropping

systems changed from conventional winter wheat-fallow to more intensive rotations. Bowman and Halvorson (1997) observed that a change from wheat-fallow to continuous wheat cropping produced a significant increase of labile P fractions in 0-5 cm surface soils. Zheng et al. (2001) observed that forage-forage-barley rotation produced larger labile P fractions than barley monoculture in the 30-60 cm layer in a Labarre silty clay after 10 years of cultivation. Recently, Zhang et al. (2006) investigated the P status after 45 years of consistent cropping practices under three cropping systems of continuous corn, rotation corn and continuous bluegrass. Compared to the adjacent native soil in forest ecosystem, the study showed that continuous cropping without P fertilization decreased all P forms significantly except for water extractable P_o , with the largest decrease in labile P_i and moderately labile P_o ; continuous cropping with fertilization resulted in comparable total P concentrations in the continuous corn and rotation corn systems and increased total P in the continuous bluegrass sod. The study also indicated that long-term cropping significantly enhanced the rate of moderately labile P_o mineralization, regardless of P fertilization, especially for continuous corn cropping system. However, compared to monoculture, crop rotations showed no effect on nutrient contents in the 0-20 cm soil layer in an experiment with soybean, corn and wheat in an Oxisol and in a Rhodic Ferralsol (DeMaria et al., 1999).

3.4 Soil texture effects on soil P fractions and transformation

Soil texture can be related to changes in P fractions and transformation. O'Halloran et al. (1985) found that up to 90% of the spatial variability in total P content of a Mollisol was explained by texture. A similar study showed that significant proportions of variability of all P fractions, except for H_2SO_4 -P in a Brown Chernozemic loam, could be attributed to changes in sand content (O'Halloran et al., 1987b). Increasing silt plus clay content has been significantly correlated with larger soil resin-P, $NaHCO_3$ - P_i/P_o and $NaOH$ - P_i/P_o pools. HCl -P was positively correlated with sand content (O'Halloran et al., 1987b). There is evidence that P transformation in soils were closely linked to microbial activity and C dynamics, which affect P mineralization and immobilization (Hedley et al., 1982). For example, a loam soil supported higher microbial biomass than the sandy loam and sandy soils (Cooper and Warman, 1997). Huffman et al. (1996) indicated that soil texture had a greater effect on P transformation than did the combined effects of residue addition, residue placement and nutrient addition, because soil texture affected the labile P_i , labile P_o , and microbial P pools. Therefore, among those factors, the particle size of soil fractions tends to account for larger proportion of variability in soil P fractions. Those findings emphasize the importance of considering soil texture when planning studies of the effects of cropping practices on P fractions and transformation.

3.5 Profile-wise distribution of soil P fractions

Much effort has been devoted to investigate the vertical changes in forms and distribution of soil P as influenced by cropping practices. These changes in P forms were mainly limited to the top of 30 cm (O'Halloran, 1993; Sharpley et al., 1993; Bowman and Halvorson, 1997; Selles et al., 1997), or to depths of 65 cm (Reddy et al., 1980; Mozaffari and Sims, 1994; Richards et al., 1995), with a few studies extending to depths ≥ 90 cm (Chang et al., 1991;

Simard et al., 1995; Beauchemin et al., 1996; Zheng et al., 2001). These studies suggested that the impacts of cropping practices on soil P fractions extended deeper in soil profile than the depth distributed by primary tillage, and P profile-wise distribution was often complicated by P downward movement. The P_o movement in soils was greater than that of P_i (O'Halloran, 1993; Richards et al., 1995). This may be due to the fact that P_o was not retained as strongly as P_i by sorbing components of solid phase, or perhaps that P_o movement was driven by mineralization process. The P_o migrated to subsoil can be mineralized, since P_o was used by microorganisms as an energy source for their metabolism in reduced layers of poorly drained soils (Zheng et al., 2001).

The downward migration of P is associated with changes in soil P sorption capacity (Reddy et al., 1980). Simard et al. (1995) found that P sorption index (P_{si}) in agricultural soils was markedly decreased in all three horizons compared with forest soils from a watershed in Quebec. Accumulation of added P has lead to an increase in total labile P pool in the A horizon (0-30 cm), while resilient P pools were the major sinks for mobile P in the B (30-60 cm) and C (60-90 cm) horizons in agricultural soils (Simard et al., 1995; Beauchemin et al., 1996; Zheng et al., 2001). This finding was in line with previous studies suggesting that repeated additions of manure decreased P sorption capacity of soils and accelerated P downward movement (Reddy et al., 1980; Sharpley et al., 1993; Mozaffari and Sims, 1994).

4. Analysis of P transformation in soils

4.1 Path analysis of soil P transformation

Path analysis is a statistical technique used to examine interrelationships among variables that are often illustrated by a path diagram. It provides not only plausible explanations of observed correlations by constructing the cause-and-effect model, but also allows decomposition of observed correlations into direct and indirect effects (Johnson and Wichern, 1988). Path analysis was useful in linking changes in soil P fractions and clarifying concepts of P transformation in soils (Tiessen et al., 1984; Beck and Sanchez, 1994; Zhang and MacKenzie, 1997a; Zheng et al., 2002, 2004a). Using path analysis, Tiessen et al. (1984) found that in Mollisols, much of resin- P_i was derived from $NaHCO_3-P_i$ and $NaOH-P_i$ fractions, and in more weathered Ultisols, 80% of variability in labile P was accounted for by P_o forms. Beck and Sanchez (1994) showed that the $NaOH-P_i$ fraction acted as a major sink for fertilizer P in soils and P_o was a major primary source of plant-available P in unfertilized soils of Peru. Zhang and MacKenzie (1997a), using the same approach for soils receiving manure and fertilizer P, indicated that P_o accumulated as $NaOH-P_o$ through $NaHCO_3-P_i$. The $NaHCO_3-P_i$ and $NaOH-P_i$ were major sinks for added P. When mineral fertilizer was the only source of P, most $NaHCO_3-P_i$ was directly supplied from fertilizer P through $NaOH-P_i$. Zheng et al. (2002, 2004a) investigated the P transformation in a Labarre silty clay as affected by nutrient sources and cropping systems in a long term study and showed that the roles of P_o pools were more important than P_i pools for P transformation and $NaHCO_3-P_o$ was sensitive to P source and was likely acted as a transitory pool rather than as a sink or source of soil P (Fig. 2). Thus, path analysis can reflect the changes in P transformation depending on soil type, climatic conditions and cropping practices.

Fig. 2. Pathways of P transformation after 10 annual applications of mineral fertilizer (A) and liquid dairy manure (B) in the 0-15 cm soil of a Labarre silty clay under barley monoculture. The percentage value indicates the partial correlation between assed P_i and resin-P; numbers are path coefficients; NS, not significant at $P \leq 0.05$ (Source: Zheng et al., 2004a)

4.2 Modeling of soil P transformation

The P concentration in soil solution is in turn controlled by P transformation processes such as mineralization, precipitation and adsorption. The complexity of interactions among these processes has led to the use of both descriptive and mechanistic mathematical models to describe them. The interpretation of changes in P concentrations in soil solution has been modeled using a variety of approaches, such as adsorption model, sorption-transport model and multi-reaction model. Most of earlier simulations of P adsorption have been based on simple rate constants (Jones et al., 1984), such as Langmuir (Enfield et al., 1981), Freundlich (Shaviv and Shacher, 1989) and Elovich equations (Chien and Clayton, 1980). A few studies have used kinetic equations such as first- or second-order reaction rates to describe the P sorption in soils (Bowden et al., 1980; Beauchemin et al., 1996). Nevertheless, adsorption models did not fully describe the mechanisms of P transformation in soils, and hence were unlikely to be generally applicable. Van de Zee and Gjaltema (1992) proposed a sorption-transport model in which both sorption reversibility and precipitation irreversibility were taken into account, but reaction rates remained uncertain.

A mechanistic multi-reaction model was conceived to represent P concentrations under dynamic boundary conditions, and to explain temporal and spatial P distribution among water soluble and resin-, NaHCO_3 -, NaOH -, and HCl -P fractions (Grant and Heaney, 1997; Grant et al., 2004). The model describes adsorption-desorption, precipitation-dissolution and ion pairing. The model can explain the temporal and spatial distribution of soluble and solid fractions under specified changes in boundary conditions in different soil types. However, the model requires more detailed, explicit information about soil chemical composition, which may not be readily available in routine fashion for many soils. Thus, where necessary, some assumptions must be made. To date, despite several P transformation models have been developed to describe certain processes, the majority of existing models need to be verified in heterogeneous soils under field conditions. Furthermore, it is unlikely that one model can kinetically represent all processes involved in P transformation.

5. Summary

Phosphorus is an essential element for plant growth and development, as it plays key roles in plant metabolism, structure and energy transformation. It is also a potential pollutant of water. The P dynamics in plant-soil-water systems is of increased interest due to its importance for environmental, agronomic and economic issues.

Soil P test involves P extraction from soils followed by a quantification of the nutrient in the extracting solution. The most widely used soil P tests are chemical extractions, such as Bray 1 & 2, Olsen, Mehlich I & III methods. Those chemical extractants are not applicable over all soil types, which is a limitation for soil P extraction. The ion-sink extractions, including ionic exchange resin membranes, resin bags, FeO coated filter papers or strips, simulate plant roots to extract P from soils without alteration of soil characteristics and have the advantage of extracting P from variety of soil type regardless of soil properties. The mainstay of P determination is the use of colorimetric procedures, most notably Murphy and Riley (1962). Inductively coupled plasma (ICP) spectrophotometry is becoming more popular for multi-element determination. The isotopic dilution (^{32}P) techniques, the ^{31}P NMR solution and the synchrotron-based techniques can provide insights into P chemistry and forms in soil and

residual materials. Although these advanced analytical techniques are capable of gaining more detailed understanding of P dynamics in soils, unfortunately not all researchers have access to the required instrumentation.

Based on its availability to plants, soil P can be divided into functional groups of readily labile P pools, moderately labile P pools and sparingly soluble P pools, each includes both P_i and P_o forms. The P transformation in soils plays important roles in P bioavailability and mobility from soil to water, and was intensively affected by cropping practices, i.e. fertilization, tillage and cropping systems. It involves complex chemical, mineralogical and biological processes. The complexity of interactions among these processes has led to the use of both descriptive and mechanistic mathematical models to describe them. Path analysis is powerful in plausibly explaining interrelationships among P pools of differing bioavailability and clarifying changes in P fractions and transformation in soils. Mechanistic multi-reaction models that describe transformation processes of mineralization-mobilization, adsorption-desorption, precipitation-dissolution and ion pairing are available to interpret the temporal and spatial distribution of P fractions in soils. However, the models require more detailed, explicit information on soil chemical composition that are not routinely available for many soils. Furthermore, most mechanistic models need to be verified in heterogeneous soils under field conditions.

6. References

- Abdu, N. 2006. Soil-phosphorus extraction methodologies: A review. *Afr. J. Agric. Res.* 1: 159-161.
- Agbenin, J.O, B.V. Raij. 2001. Kinetics and energetics of phosphate release from tropical soils determined by mixed ion-exchange resin. *Soil Sci. Soc. Am. J.* 65:1108-1114.
- Anderson, G. 1980. Assessing organic phosphorus in soils. p. 411-431. *In* F.E. Khasawneh et al. (eds.) *The role of phosphorous in agriculture*. ASA, Madison, WI.
- Barrow, N.J., T.C. Shaw. 1977. Factors affecting the rate of phosphate extracted from soils by anion exchange resin. *Geoderma*. 18:309-323.
- Beauchemin, S., R.R. Simard, and D. Cluis. 1996. Phosphorus sorption-desorption kinetics of soil under contrasting land uses. *J. Environ. Qual.* 25:1317-1325.
- Beauchemin, S., and R.R. Simard. 2000. Phosphorus status of intensively cropped, neutral to calcareous soils of the St. Lawrence lowlands. *Soil Sci. Soc. Am. J.* 64:659-670.
- Beauchemin, S., D. Hesterbeg, J. Chou, M. Beauchemin, R.R. Simard, and D.E. Sayers. 2003. Speciation of phosphorus in phosphorus-enriched agricultural soils using x-ray absorption near-edge structure spectroscopy and chemical fractionation. *J. Environ. Qual.* 32:1809-1819.
- Beck, M.A., and P.A. Sanchez. 1994. Soil phosphorus fraction dynamics during 18 years of cultivation on a Typic Paleudult. *Soil Sci.* 34:1424-1431.
- Blevins, R.L., W.W. Frye, P.F. Baldwin, and S.D. Robertson. 1990. Tillage effects on sediment and soluble nutrient losses from a Maury silt loam. *J. Environ. Qual.* 19:683-686.
- Bowden, J.W., S. Nagarajah, N.J. Barrow, A.M. Posner, and J.P. Quirk. 1980. Describing the adsorption of phosphate, citrate and selenite on a variable charge mineral surface. *Austr. J. Soil Res.* 18:49-60.
- Bowman, R.A., and A.D. Halvorson. 1997. Crop rotation and tillage effects on phosphorus distribution in the Central Great Plains. *Soil Sci. Soc. Am. J.* 61:1418-1422.

- Bray, R.H., and L.T. Kurtz. 1945. Determination of total, organic, and available forms of phosphorus in soils. *Soil Sci.* 59:39-45.
- Carski, T.H., D.L. Sparks. 1985. A modified miscible displacement technique for investigating adsorption-desorption kinetics in soils. *Soil Sci. Soc. Am. J.* 49:1114-1116.
- Castillo, M.S., and A.L. Wright. 2008. Soil phosphorus pools for Histosols under sugarcane and pasture in the Everglades, USA. *Geoderma* 145:130-135.
- Chang, C., T.G. Sommerfeldt, and T. Entz. 1991. Soil chemistry after eleven applications of cattle feedlot manure. *J. Environ. Qual.* 20:475-480.
- Chang, S.C., and M.L. Jackson. 1957. Fractionation of soil phosphorus. *Soil Sci.* 84:133-144.
- Chardon, W.J., R.J. Menon, S.H. Chien. 1996. Iron oxide impregnated filter paper (Pi test): A review of its development and methodological research. *Nutr. Cycling Agroecosyst.* 46:41-51.
- Chien, S.H., and W.R. Clayton. 1980. Application of Elovich equation to the kinetics of phosphate release and sorption in soils. *Soil Sci. Soc. Am. J.* 44:265-268.
- Cooper, J.M., and P.R. Warman. 1997. Effects of three fertility amendments on soil dehydrogenase activity, organic C and pH. *Can. J. Soil Sci.* 77:28-283.
- Cross, A.F., and W.H. Schlesinger. 1995. A literature review and evaluation of the Hedley fractionation: Applications to the biogeochemical cycle of soil phosphorus in the natural ecosystems. *Geoderma* 64:197-214.
- Cruse, R.M., G.A. Yackle, T.C. Colvin, and D.R. Timmons. 1983. Tillage effects on corn and soybean production in farmer managed, university-monitored field plots. *J. Soil Water Conserv.* 38:512-514.
- De Boer, J.L.M., U. Kohlmeier, P.M. Breugem, and T. van der Velde-Koerts. 1998. Determination of total dissolved phosphorus in water samples by axial inductively coupled plasma emission spectrometry. *J. Anal. Chem.* 360:132-136.
- De Maria, I.C., P.C. Nnabude, and O.M. de Castro. 1999. Long-term tillage and crop rotation effects on soil chemical properties of a Rhodic Ferralsol in southern Brazil. *Soil Tillage Res.* 51:71-79.
- Dick, W.A. 1983. Organic carbon, nitrogen and phosphorus as influenced by tillage intensity. *Soil Sci. Soc. Am. J.* 47:102-107.
- Enfield, C.G., T. Phan, D.M. Walters, and R. Ellis. 1981. Kinetic models for phosphate transport and transformations in calcareous soils: 1. Kinetics of transformations. *Soil Sci. Soc. Am. J.* 45: 1059-1070.
- Follett, R.F., and G.A. Peterson. 1988. Surface soil nutrient distribution as affected by wheat-fallow tillage systems. *Soil. Sci. Soc. Am. J.* 52:141-147.
- Frossard, E., L.M. Condron, A. Oberson, S. Sinaj, and J.C. Fardeau. 2000. Processes governing phosphorus availability in temperate soils. *J. Environ. Qual.* 29:15-23.
- Grant, R.F., and D.J. Heaney. 1997. Inorganic phosphorus transformation and transport in soils: mathematical modeling in ecosystems. *Soil Sci. Soc. Am. J.* 61:752-764.
- Grant, R.F., M. Amrani, D.J. Heaney, R. Wright, and M. Zhang. 2004. Mathematical modelling of phosphorus losses from land application of hog and cattle manure. *J. Environ. Qual.* 33:210-233.
- Hedley, M.J., J.W.B. Stewart, and B.S. Chauhan. 1982. Changes in inorganic and organic phosphorus induced by cultivation practices and laboratory incubations. *Soil Sci. Soc. Am. J.* 46:970-976.

- Holford, J.C.R. 1997. Soil phosphorus: its measurement, and its uptake by plants. *Aust. J. Soil Res.* 35: 227-239.
- Huffman, S.A, C.V. Cole, and N.A. Scott. 1996. Soil texture and residue addition effects on soil phosphorus transformation. *Soil Sci. Soc. Am. J.* 60:1095-1101.
- Ivarsson, K. 1990. The long-term soil fertility experiments in southern Sweden. IV. Changes in inorganic and organic soil phosphorus after a pot trial. *Acta Agric. Scand.* 40:205-215.
- Jones, C.A., C.V. Cole, A.N. Sharpley, and J.R. Williams. 1984. Simplified soil and plant phosphorus model: I. Mobility of phosphorus. *Soil Sci. Soc. Am. Proc.* 38: 446-451.
- Johnson, R.A., and D.W. Wichern. 1988. Multivariate linear regression models. p. 273-333. *In* Applied multivariate statistical analysis. 2nd ed. Prentice Hall, Englewood Cliffs, NJ.
- Kamprath, E.J., and M.E. Watson. 1980. Conventional soil and tissue tests for assessing the phosphorus status of soils. p. 433-469. *In* F.H. Khasawneh et al. (eds.) The role of phosphorus in agriculture. ASA, CSSA, and SSSA, Madison, WI.
- Iyamuremye, F., R.P. Dick, and J. Baham. 1996. Organic amendments and phosphorus dynamics: II. Distribution of soil phosphorus fractions. *Soil Sci.* 161:436-443.
- Koopmans G.F., M.E. Van Der Zeeuw, P.F.M.A. Romkens, W.J. Chardon, O. Oenema 2001. Identification and characterization of phosphorus rich sandy soils. *Neth. J. Agric. Sci.* 49:369-384.
- Mallarino, A.P. 1997. Interpretation of soil phosphorus tests for corn in soils with varying pH and calcium carbonate content. *J. Prod. Agric.* 10:163-167.
- Mattingly, G.E.G. 1975. Labile phosphate in soils. *Soil Sci.* 119: 369-375.
- McGill, W.B., and C.V. Cole. 1981. Comparative aspects of C, N, S and P cycling through soil organic matter during pedogenesis. *Geoderma* 26:267-286.
- McKenzie, R.H., J.W.B. Stewart, J.F. Dormaar, and G.B. Schaalje. 1992a. Long-term crop rotation and fertilizer effects on phosphorus transformation: I. In a Chernozemic soil. *Can. J. Soil Sci.* 72:569-579.
- McKenzie, R.H., J.W.B. Stewart, J.F. Dormaar, and G.B. Schaalje. 1992b. Long-term crop rotation and fertilizer effects on phosphorus transformation: II. In a Luvisolic soil. *Can. J. Soil Sci.* 72:581-589.
- Mehlich, A. 1984. Mehlich no.3 extractant: A modification of Mehlich no.2 extractant. *Commun. Soil Sci. Plant Anal.* 15:1409-1416.
- Mozaffari, M., and J.T. Sims. 1994. Phosphorus availability and sorption in an atlantic coastal plain watershed dominated by animal-based agriculture. *Soil Sci.* 157:97-107.
- Murphy, J., and J.P. Riley. 1962. A modified single solution method for determination of phosphate in natural waters. *Anal. Chim. Acta.* 27:31-36.
- Murrman, R.P., and M. Peech. 1969. Relative significance of labile and crystalline phosphates in soil. *Soil Sci.* 107:149-155.
- Myers, R.G., A.N. Sharpley, S.J. Thien, G.M. Pierzynski. 2005. Ion-sink phosphorus extraction methods applied on 24 soils from the continental USA. *Soil Sci. Soc. Am. J.* 69: 511-521.
- N'dayegamiye, A., and D.A. Angers. 1990. Effets de l'apport prolongé de fumier de bovins sur quelques propriétés physiques et biologiques d'un loam limoneux Neubois sous culture de maïs. *Can. J. Soil Sci.* 70:259-262.

- O'Halloran, I.P., R.G. Kachanoski, and J.W.B. Stewart. 1985. Spatial variability of soil phosphorus as influenced by soil texture and management. *Can. J. Soil Sci.* 65:475-487.
- O'Halloran, I.P., J.W.B. Stewart, and E. De Jong. 1987a. Changes in P forms and availability as influenced by management practices. *Plant Soil* 100:113-126.
- O'Halloran, I.P., J.W.B. Stewart, and R.G. Kachanoski. 1987b. Influence of texture and management practices on the forms and distribution of soil phosphorus. *Can. J. Soil Sci.* 67:147-163.
- O'Halloran, I.P. 1993. Effect of tillage and fertilizer on the inorganic and organic phosphorus. *Can. J. Soil Sci.* 73:359-369.
- Olsen, S.R., C.V. Cole, F.S. Watanabe, and L.A. Dean. 1954. Estimation of available phosphorus by extraction with sodium bicarbonate. USDA Circ. 939, U.S. Govt. Print. Office, Washington, D.C.
- Paniagua, A., M.J. Mazzarino, D. Hass, L. Szott, and C. Fernandez. 1995. Soil phosphorus fractions under tropical agro-ecosystems on a Volcanic Soil. *Austr. J. Soil Res.* 33: 311-320.
- Petersen, G.W., and R.B. Corey. 1966. A modified Chang and Jackson procedure for Routine fractionation of inorganic soil phosphates. *Soil Sci. Soc. Am. Proc.* 30: 563-565.
- Pierzynski, G.M., A.N. Sharpley, and J.L. Kovar. 2008. Methods of phosphorus analyses for soils, sediments, residues and water: Introduction. p. 1-4. *In* J.L. Kovar and G.M. Pierzynski (eds.), *Methods for phosphorus analysis for soils, sediments, residuals and waters*. 2nd Edition. Southern Cooperative Series Bulletin No. 408.
- Pierzynski, G.M., H.L. Zhang, A. Wolf, P. Kleinman, A. Mallarino, and D. Sullivan. 2011. Phosphorus determination in waters and extracts of soils and by-products: inductively coupled plasma spectrometry versus colorimetric procedures. http://www.sera17.ext.vt.edu/SERA_17_Publications.htm. Accessed on July 13, 2011.
- Raven, K.P, and L.R. Hossner. 1993. Phosphate desorption quantity-intensity relationships in soil. *Soil Sci. Soc. Am. J.* 57:1505-1508.
- Reddy, K.R., M.R. Overcash, R. Khaleel, and P.W. Westerman. 1980. Phosphorus adsorption-desorption characteristics of two soils utilized for disposal of animal waste. *J. Environ. Qual.* 9: 86-92.
- Richards, J.E., T.E. Bates, and S.C. Sheppard. 1995. Change in the forms and distribution of soil phosphorus due to long-term corn production. *Can. J. Soil Sci.* 75:311-318.
- Robinson, J.S., and A.N. Sharpley. 1994. Effect of organic phosphorus on the sink characteristics of iron oxide-impregnated filter paper. *Soil Sci. Soc. Am. J.* 58:758-761.
- Schmidt, J.P., S.W. Buol, and E.J. Kamprath. 1996. Soil phosphorus dynamics during seventeen years of continuous cultivation: fractionation analysis. *Soil Sci. Am. J.* 60: 1168-1172.
- Schoenau, J.J., and W.N. Huang. 1991. Anion-exchange membrane, water and sodium bicarbonate extractions as soil tests for phosphorus. *Commun. Soil Sci. Plant Anal.* 22: 465-492.
- Selles, F., R.A. Kochann, J.E. Denardin, R.P. Zentner, and A. Faganello. 1997. Distribution of phosphorus fractions in a Brazilian Oxisol under different tillage systems. *Soil & Tillage Res.* 44:23-34.

- Selles, F., B.G. McConkey, and C.A. Campbell. 1999. Distribution and forms of P under cultivator and zero-tillage for continuous- and fallow-wheat cropping systems in the semi-arid Canadian prairies. *Soil Tillage Res.* 51:47-59.
- Sharpley, A.N., S.J. Smith, B.A. Stewart, and A.C. Mathers. 1984. Forms of phosphorus in soil receiving cattle feedlot waste. *J. Environ. Qual.* 13:211-215.
- Sharpley, A.N., S.J. Smith, and W.R. Bain. 1993. Nitrogen and phosphorus fate from long-term poultry litter applications to Oklahoma soils. *Soil Sci. Soc. Am. J.* 57:1131-1137.
- Sharpley, A.N., J.T. Sims, and G.M. Pierzynski. 1994. Innovative soil phosphorus availability indices: assessing inorganic phosphorus. p. 116-142. *In* Soil testing: prospects for improving nutrient recommendations, SSSA special publication 40. SSSA, Madison, WI.
- Sharpley, A.N., and H. Tunney. 2000. Phosphorus research strategies to meet agricultural and environmental challenges of 21st century. *J. Environ. Qual.* 29:176-181.
- Shaviv, A., and N. Shachar. 1989. A kinetic-mechanistic model of phosphorus sorption in calcareous soils. *Soil Sci.* 148:172-178.
- Shober, A.L., D.L. Hesterberg, J.T. Sims, and S. Gardner. 2006. Characterization of phosphorus species in biosolids and manures using XANES spectroscopy. *J. Environ. Qual.* 35:1983-1993.
- Sibbesen, E. 1978. An investigation of the anion-exchange resin method for soil phosphate extraction. *Plant Soil* 50:305-321.
- Simard, R.R., D. Cluis, G. Gangbazo, and S. Beauchemin. 1995. Phosphorus status of forest and agricultural soils from a watershed of high animal density. *J. Environ. Qual.* 24:1010-1017.
- Sparks, D.L. 1985. Kinetics of ionic reactions in clay minerals and soils. *Adv. Agron.* 38: 231-266.
- Sparks, D.L. 1999. Kinetics and mechanisms of chemical reactions at the soil mineral/water interface. p. 135-191. *In* D.L. Sparks (ed.) *Soil physical chemistry*. 2nd edition. CRC Press, Boca Raton, FL.
- Stevenson, F.J. 1986. The phosphorus cycle. p. 231-284. *In* *Cycles of soil: carbon, nitrogen, phosphorus, sulfur, micronutrients*. John Wiley & Sons, New York, NY.
- Sweeney, D.W. 1993. Fertilizer placement and tillage effects on grain sorghum growth and nutrient uptake. *Soil Sci. Soc. Am. J.* 57:532-537.
- Tiessen, H., J.W.B. Stewart, and J.R. Bettany. 1982. Cultivation effect on the amounts and concentration of carbon, nitrogen and phosphorus in grassland soils. *Agron. J.* 74:831-835.
- Tiessen, H., J.W. Stewart, and C.V.Cole. 1984. Pathways of phosphorus transformation in soils of differing pedogenesis. *Soil Sci. Soc. Am. J.* 48:853-858.
- Tiessen, H., and J.O. Moir. 1993. Characterization of available P by sequential extraction. p. 75-86. *In* M.R. Carter (ed.) *Soil sampling and methods of analysis*. Can. Soil Sci. Soc., Lewis Publ., Boca Raton, FL.
- Tran, T.S, J.C. Fardeau, and M. Giroux. 1988. Effects of soil properties on plant-available P determined by the isotopic dilution P-32 method. *Soil Sci. Soc. Am. J.* 52:1382-1390.
- Tran, T.S., and A. N'dayegamiye. 1995. Long-term effects of fertilizers and manure application on the forms and availability of soil phosphorus. *Can. J. Soil Sci.* 75:281-285.

- Turner, B.L., and A.B. Leytem. 2008. Phosphorus speciation in soils and manures by solution ^{31}P NMR spectroscopy. p. 95-101. In J.L. Kovar and G.M. Pierzynski (eds.) Methods for phosphorus analysis for soils, sediments, residuals and waters. 2nd Edition. Southern Cooperative Series Bulletin No. 408.
- Uusitalo, R, and M. Yli-Halla. 1999. Estimating errors associated with extracting phosphorus using iron oxide and resin methods. J. Environ. Qual. 28:1891-1897.
- Van der Zee, S.E.A.T.M., and A. Gjaltema. 1992. Simulation of phosphate transport in soil columns. I. Model development. Geoderma 52:87-109.
- Wagar, B.I., J.W.B. Stewart, and J.O. Moir. 1986. Changes with time in the form and availability of residual fertilizer phosphorus on Chernozemic soils. Can. J. Soil Sci. 66:105-119.
- Walker, T.W., and J.K. Syers. 1976. The fate of phosphorus during pedogenesis. Geoderma 15:1-19.
- Weill, A.N., G.R. Mehuys, and E. McKyes. 1990. Effect of tillage reduction and fertilizer type on soil properties during corn production. Soil & Tillage Res. 17:63-76.
- Williams, J.D.H., J.K. Syers, and T.W. Walker. 1967. Fractionation of soil inorganic phosphate by a modification of Chang and Jackson's procedure. Soil Sc. Soc. Am. Proc. 31:736-739.
- White, R.E., and P.H.T. Beckett. 1964. Studies on the phosphate potentials of soils. 1. The measurement of phosphate potential. Plant and Soil 20:1-16.
- Wright, A.L. 2009. Phosphorus sequestration in soil aggregates after long-term tillage and cropping. Soil & Tillage Res. 103:406-411.
- Zhang, T.Q., and A.F. MacKenzie. 1997a. Changes of soil phosphorus fractions under long-term corn monoculture. Soil Sci. Soc. Am. J. 61:485-493.
- Zhang, T.Q., and A.F. MacKenzie. 1997b. Phosphorus in zero-tension soil solution as influenced by long-term fertilization of corn. Can. J. Soil Sci. 77:685-691.
- Zhang, T.Q. and A.F. Mackenzie. 1997c. Changes of phosphorous fractions under continuous corn production in a temperate clay soil. Plant Soil. 192:133-139.
- Zhang, T.Q., A.F. Mackenzie, and F. Sauriol. 1999. Nature of soil organic phosphorus as affected by long-term fertilization under continuous corn (*Zea mays* L.): A ^{31}P NMR study. Soil Sci. 164:662-670.
- Zhang, T.Q., A.F. Mackenzie, B.C. Liang, and C.F. Drury. 2004. Soil test phosphorus and phosphorus fractions with long-term phosphorus addition and depletion. Soil Sci. Soc. Amer. J. 68:519-528.
- Zhang, T.Q., C.S. Tan, C.F. Drury, and W.D. Reynolds. 2006. Long-term (≥ 43 years) fate of soil phosphorus as related to cropping systems and fertilization. World Congress of Soil Science. Philadelphia, USA. July 9-16.
- Zheng, Z., R.R. Simard, J. Lafond, and L.E. Parent. 2001. Changes in phosphorus fractions of a Humic Gelysol as influenced by cropping systems and nutrient sources. Can. J. Soil Sci. 81:175-183.
- Zheng, Z., R.R. Simard, J. Lafond, and L.E. Parent. 2002. Pathways of soil phosphorus transformations after 8 years of cultivation under contrasting cropping practices. Soil Sci. Soc. Am. J. 66:999-1007.
- Zheng, Z., J.A. MacLoed, J.B. Sanderson, and J. Lafond. 2004a. Soil phosphorus dynamics during 10 annual applications of manure and mineral fertilizers: fractionation and path analyses. Soil Sci. 169:449-456.

Zheng, Z., J.A. MacLeod, and J. Lafond. 2004b. Phosphorus status of a Humic Cryaquept profile in frigid continental climate as influenced by cropping practices. *Biol. Fertil. Soils* 39:467-473.

IntechOpen

IntechOpen

Soil Fertility Improvement and Integrated Nutrient Management - A Global Perspective

Edited by Dr. Joann Whalen

ISBN 978-953-307-945-5

Hard cover, 306 pages

Publisher InTech

Published online 24, February, 2012

Published in print edition February, 2012

Soil Fertility Improvement and Integrated Nutrient Management: A Global Perspective presents 15 invited chapters written by leading soil fertility experts. The book is organized around three themes. The first theme is Soil Mapping and Soil Fertility Testing, describing spatial heterogeneity in soil nutrients within natural and managed ecosystems, as well as up-to-date soil testing methods and information on how soil fertility indicators respond to agricultural practices. The second theme, Organic and Inorganic Amendments for Soil Fertility Improvement, describes fertilizing materials that provide important amounts of essential nutrients for plants. The third theme, Integrated Nutrient Management Planning: Case Studies From Central Europe, South America, and Africa, highlights the principles of integrated nutrient management. Additionally, it gives case studies explaining how this approach has been implemented successfully across large geographic regions, and at local scales, to improve the productivity of staple crops and forages.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Z.M. Zheng and T.Q. Zhang (2012). Soil Phosphorus Tests and Transformation Analysis to Quantify Plant Availability: A Review, Soil Fertility Improvement and Integrated Nutrient Management - A Global Perspective, Dr. Joann Whalen (Ed.), ISBN: 978-953-307-945-5, InTech, Available from:
<http://www.intechopen.com/books/soil-fertility-improvement-and-integrated-nutrient-management-a-global-perspective/soil-phosphorus-tests-and-transformation-analysis-to-quantify-plant-availability-a-review>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](https://creativecommons.org/licenses/by/3.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen