

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

186,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Sex Steroids Modulate Fish Immune Response

Elena Chaves-Pozo¹, Isabel Cabas² and Alfonsa García-Ayala²

¹*Centro Oceanográfico de Murcia, Instituto Español de Oceanografía (IEO),
Carretera de la Azohía s/n. Puerto de Mazarrón, Murcia*

²*Department of Cell Biology and Histology, Faculty of Biology,
University of Murcia, Murcia
Spain*

1. Introduction

For some time behavioural and ecological studies have suggested that sex steroid hormones regulate several immune processes in fish. For example, the immunocompetence handicap hypothesis relates the heritability of parasite resistance with secondary sexual ornaments, which are determined and maintained by androgens. Such ornaments are probably a good indicator to potential mates of genetic resistance to infections (Dijkstra et al., 2007; Roberts et al., 2004). Among vertebrates, the prevalence and intensity of parasitic infections is higher in males than females (Klein, 2004). Some fish species show altered sex steroid hormones levels upon parasite infection. The main alterations recorded upon infection are decreases in androgen, estrogen and vitelogenin serum levels (Hecker & Karbe, 2005). For example, during an infective period of vibriosis, silver seabream showed gradually increasing testosterone serum levels, whereas serum estradiol levels significantly decreased at an early stage of infection and remained low until death. This process coincided with increasing macrophages phagocytic activity (Deane et al., 2001). Such field studies prompted immunologists to try to establish how sex steroid hormones are able to alter the functions of the circulating leukocytes. In fish, most existing information on reproductive-immune interactions deals with the modulation of immune responses by circulating hormones, including cortisol, growth hormone, prolactin and reproductive hormones and some proopiomelanocortin-derived peptides (Engelsma et al., 2002; Harris & Bird, 2000). Although the exact effect of these endocrine mediators depends on the species, in general, they are known to modulate immune responses by integrating the activities of all the systems. In this way they help to adapt the organism to its environment (Lutton & Callard, 2006).

From a reproductive biology point of view, the leukocytes located in mammalian gonads orchestrate important reproductive physiology processes, including gametogenesis and steroidogenesis. A long time has passed since leukocytes were first described in the gonad of teleosts. Since then, several types of leukocytes have been described in the testis of different teleost species using light and electron microscopy. Moreover, differences in the number and localization of leukocytes within the testis have also been observed during the different stages of the reproductive cycle (Besseau & Faliex, 1994; Billard, 1983; Bruslé-Sicard & Fourcault, 1997; Lo Nostro, 2004; Scott & Sumpter, 1989). Thus, in the gametogenic activity and spawning stages some macrophages have been described in the interstitial tissue of the rainbow trout testis (Loir et al., 1995), whereas in the post-spawning stage a

high population of phagocyte cells has been described in several teleost fish (Henderson, 1962; Loir et al., 1995; Scott & Sumpter, 1989; Shrestha & Khanna, 1976). Although macrophages, granulocytes and lymphocytes have been described in the testis of some sparid fish, only macrophages have been shown to be phagocytic cells (Besseau & Faliex, 1994; Bruslé-Sicard & Fourcalt, 1997; Micale et al., 1987).

The gilthead seabream (*Sparus aurata* L.) is a protandrous hermaphrodite seasonal breeding teleost with a bisexual gonad (Figure 1) that offers an interesting model for studying immune-reproductive interactions. This is because the remodelling events of the gonad, especially during the post-spawning and testicular involution stages, compromise the immune system. The specimens undergo sex change during the second or third year of life, depending on the natural environment of the populations studied (Lasserre, 1972). Our previous studies performed in the western Mediterranean area demonstrated that gilthead seabream are males during the first and second reproductive cycles although their gonads possess a non-developed ovarian area separated from the testicular area by connective tissue (Chaves-Pozo et al., 2005a; Liarte et al., 2007). The reproductive cycle of males is divided into four gonad stages: gametogenic activity, spawning, post-spawning and resting. Resting is replaced by a testicular involution stage when the fish are ready to undergo sex change (Chaves-Pozo et al., 2005a; Liarte et al., 2007).




Fig. 1. Section of the gonad of gilthead seabream (*Sparus aurata* L.) during the male phase. The bisexual gonad is formed of a testis and an ovary separated by a thin layer of connective tissue. The testis is constituted by seminiferous tubules which are in the initial stage of spermatogenesis and the ovary is occupied with pre-vitellogenic ovocytes. T, testis; Ov, ovary; CT, connective tissue. (Mallory trichromic) Magnification x 10.

During the first reproductive cycle, 11-ketotestosterone and testosterone, the main androgens in fish, play different and specific roles in the testicular physiology as they peak at different stages of the reproductive cycle. Moreover, the profiles of testosterone serum levels during the second reproductive cycle demonstrated that this androgen is not essential to the testicular regression process that occurs during this cycle. In contrast, changes in 17β -estradiol serum levels suggest that this hormone orchestrates the testicular regression process during both reproductive cycles. Moreover, the data suggest that there is a threshold level of 17β -estradiol that determines the initiation of ovarian development during the second reproductive cycle without promoting complete feminization (Chaves-Pozo et al., 2008a).




Fig. 2. Sections of the testis of gilthead seabream in the spermatogenesis (a), spawning (b), post-spawning (c) and resting (d) stages immunostained with a monoclonal antibody specific to gilthead seabream acidophilic granulocytes (Sepulcre et al., 2002). Acidophilic granulocytes (arrows) are seen in the blood vessels (a) during spermatogenesis, in the lumen of the tubules between the spermatozoa (b) and in the seminal epithelium in contact with germ cells (c) during spawning and post-spawning and in the interstitial tissue (d) during resting. Magnification x 400.

Few studies have dealt with the presence of leukocytes in the gonad of teleosts, their functions and the molecular pathways that regulate them. However, our studies in recent years have suggested that sex hormones might be key regulators of leukocyte functions in the gonad. For example, a massive infiltration of leukocytes, mainly acidophilic granulocytes (Figure 2), is orchestrated by gonadal factors including sex steroid hormones during post-spawning and testicular involution stages (Chaves-Pozo et al., 2003, 2005a, 2005b, 2007). The immune cells are produced in the head-kidney, the main haematopoietic organ in fish. However, when the acidophilic granulocytes infiltrate the testis, they show heavily impaired reactive oxygen intermediate production and phagocytic activity (hardly 1% of the testicular acidophilic granulocytes are able to phagocytise) (Figure 3) while the production of interleukin-1 β (IL-1 β) is sharply induced (Chaves-Pozo et al., 2003, 2005b, 2008a).




Fig. 3. An electron micrograph showing a testicular acidophilic granulocyte with the typical ultrastructure of acidophilic granulocytes and two phagosomes containing *Vibrio anguillarum* cells (Va). Magnification $\times 5000$.

Interestingly, it is the gonad itself which actively regulates the presence of these immune cells in the testis by stimulating their extravasation from the blood (Chaves-Pozo et al., 2005b). Moreover, 17 β -estradiol and testosterone seem to be related with the infiltration of acidophilic granulocytes and probably with the magnitude of the infiltration since both hormones peak when the infiltration of these cells into the gonad occurred (Chaves-Pozo et

al., 2008a). Moreover, the infiltration of acidophilic granulocytes was correlated with an increase in the expression of gonadal aromatase, the enzyme that transforms testosterone to 17β -estradiol. Such expression was seen to remain high during the period that acidophilic granulocytes are present in the gonad (Chaves-Pozo et al., 2005a, 2008b; Liarte et al., 2007). Moreover, experimentally induced increases of 17β -estradiol serum levels in spermatogenically active males triggered the migration of acidophilic granulocytes to the gonad in a way that resembles an inflammatory process (Chaves-Pozo et al., 2007). In the adult gilthead seabream gonad, macrophages and lymphocytes have also been observed in the interstitial tissue (Chaves-Pozo et al., 2008a; Liarte et al., 2007). However, the number of testicular macrophages remains steady throughout the reproductive cycle when the specimens are males, while no data related to lymphocytes are available (Chaves-Pozo et al., 2008a). Acidophilic granulocytes and B lymphocytes (Figure 4) also infiltrated the gonad and were located in the interstitial tissue and among the spermatozoa when fish were treated with an estrogenic endocrine disruptor, 17α -ethynylestradiol. This pharmaceutical compound, used for oral contraceptives and hormone replacement therapy, has a widespread presence in the aquatic environment (Ternes et al., 1999) and may reach concentrations of 0.5 to 62 ng/l in European sewage and surface waters (Hinteman et al., 2006; Johnson et al., 2005; Kuch and Ballschmiter, 2000).




Fig. 4. Sections of the gilthead seabream testis in the spermatogenesis stage of specimens control (a) and specimens treated with $5\text{ }\mu\text{g}$ of 17α -ethynylestradiol/g food (b) immunostained with a specific anti-gilthead seabream IgM serum. B lymphocytes can be seen in the interstitial tissue of the testis, the numbers slightly increasing after 17α -ethynylestradiol treatment. Magnification $\times 200$.

Testosterone administration *in vivo* modulates particular components of the physiological response of professional phagocytes such as respiratory burst, but does not alter their phagocytic activity. Testosterone is also able to regulate the gene expression profile of immune related molecules in head-kidney and other immune competent organs. This effect is characterized by a strong pro-inflammatory activation in the first week, after which it changes into an anti-inflammatory response (Águila et al., 2010).

These observations which, taken together, suggest that the presence of immune cells and cytokines in the gonad guarantees and modulates the reproductive functions (Figure 5),

2. Sex steroid hormones as regulators of the immune response

In mammals, androgens and estrogens exert their main long-term effects on cell growth, cell differentiation and cell functions through intracellular androgen receptors (AR) and estrogen receptors (ER), ER α and ER β , respectively, all of which belong to the nuclear receptor superfamily (Evans & Bergeron, 1988). These AR and ER are ligand-inducible transcription factors that cause the activation or repression of genes (Beato & Klug, 2000; Kumar & Tindall, 1998). In different mammalian models, the preponderance of ER α gene over the ER β gene is accepted as being one of the mechanism that control the effects of 17 β -estradiol on the immune system (Straub, 2007). The main effect of estrogens on the immune response involve enhancing the immune/inflammatory response by activating the nuclear factor κ B (NF κ B) signalling pathway (Cutolo et al., 2004) and stimulating the secretion of tumor necrosis factor (TNF) (Janele et al., 2006). Furthermore, using ER knock-out mice, researchers have shown that ER participates in the stimulation of interleukin (IL)-10 and immunoglobulin (Ig) M production. In accordance with these roles, a number of epidemiological studies have highlighted the relationship between plasma estrogen levels, IL production, and autoimmune disorders linked to some diseases (Cutolo et al., 2006). However, 17 β -estradiol also has an inhibitory effect on bone resorption and the suppression of inflammation in several animal models of chronic inflammatory diseases (Straub, 2007). Unlike estrogens, androgens are thought to be exclusively immunosuppressive in mammals. For example, androgens have a negative effect on the expression of inflammatory cytokines, increase apoptosis in human monocytes/macrophages, and inhibit lymphocyte proliferation (Cutolo & Straub, 2009; Cutolo et al., 2005; Lehmann et al., 1988).

In teleosts, the large number of different species and the genome duplications that have occurred during their phylogeny make it very difficult to assess the number of AR and ER existing in each specie. Depending on the species studied, three or four different ER genes have been described. Thus, in some species (gilthead seabream, atlantic croaker, zebrafish, goldfish) one ER α and two ER β have been cloned, while in others (rainbow trout and *Spinibarbus denticulatus*) two ER α and two ER β were found (Iwanowicz & Ottinger, 2009; Nagler et al., 2007). In order to determine whether immune tissues are potential targets for estrogens, several studies have looked at the expression of ER in immune tissues. In immature and mature male and female channel catfish, for example, ER α is expressed in spleen, blood and head-kidney, while ER β is only expressed in spleen (Xia et al., 2000). ER β is expressed in the spleen and head-kidney of male and female common solea (Caviola et al., 2007). In the gilthead seabream, *in vitro* long term treatment of head-kidney leukocytes with 17 β -estradiol revealed a suppressive effect on the production of reactive oxygen intermediates and the *Vibrio anguillarum* DNA (VaDNA)-stimulated production of IL-1 β (Chaves-Pozo et al., 2003). However, short term treatment with higher concentrations of 17 β -estradiol inhibited the phagocytic capability, while the percentage of phagocytic cells and the VaDNA-stimulated production of reactive oxygen intermediates and cell migration activity remained steady (Liarte et al., 2011b). In the case of AR, most vertebrates are believed to have one active form of nuclear AR with high specificity for the androgen 5 α -dihydrotestosterone, whereas there appear to be two subtypes of AR in some teleosts, AR α and AR β . These are differentially expressed in tissues and show high affinity for both testosterone and 11 β -hidroxytestosterone (review in Rempel & Schlenk, 2008). However, little is known about the expression of these AR in fish immune tissues, although in mammalian models AR are present in liver, spleen and thymus (Butts et al., 2011).

There is increasing evidence supporting the transcription-independent non-genomic actions of steroid hormones, including testosterone and 17 β -estradiol (Christ et al., 1997; Falkenstein et al., 2000). For example, mammalian mast cells, T and B cells and macrophages shows membrane AR and membrane ER (Benten et al., 1998, 2001, 2002; Zaitzu et al., 2007). A membrane ER (Pang et al., 2008) and a membrane AR (review in Thomas et al., 2006) have recently been cloned and characterized in atlantic croaker, although nothing is known about membrane AR in fish immune tissues. The complexity of the way in which sex steroid hormones act in fish through membrane and intracellular receptors, as well as the complexity of the systemic and gonadal immune responses and the several cell types involved, prompted researchers to characterize sex steroid hormone receptors and the effects of their ligands in purified immune cells and cell lines. However, since each cell type has its own response pattern these issues will be dealt with separately.

2.1 Macrophages

Macrophages are ubiquitous cells that play a central role in the innate immune response through the secretion of inflammatory cytokines, such as IL-1 β and TNF α , the production of cytotoxic reactive oxygen intermediates, and the secretion of leukostatic factors and other regulatory molecules. They are also important accessory cells for many other immune responses. In addition, during development, these cells are thought to have a trophic role through their remodelling capabilities and ability to produce cytokines. Interestingly, whereas a similar pattern of functioning has been demonstrated for macrophages in different tissues (Guillemin & Brew, 2004; Laskin et al., 2001; Stout & Suttles, 2004), testicular macrophages and their functions are largely determined by the local environment (Hedger, 1997, 2002), including not only cytokines and chemokines, but also steroid hormones. In mammals, it has been known for many years that ER are expressed in monocytes (Cunningham & Gilkeson). However, their response to estrogens and the predominance of ER α or ER β expression appear to be dependent on their stage of differentiation. For example, Mor et al. (2003) demonstrated that monocytes express more ER β and macrophages express more ER α . The behaviour of 17 β -estradiol functions in mammalian macrophages has been described as double-edge-sword (depending on 17 β -estradiol concentration). Thus, lower 17 β -estradiol concentrations stimulated IL-1 β production, whereas higher concentrations inhibited lipopolysaccharide (LPS)-induced TNF α production. This dichotomous effect of 17 β -estradiol on IL-1 β and TNF α at high and low concentrations is most probably due to inhibition of NF- κ B at high concentrations (review in Straub, 2007).

Gilthead seabream macrophages constitutively express only the ER α gene, although stimulation with VaDNA drastically up-regulates the expression of ER α , ER β 1 and ER β 2 genes, suggesting that the immune system is able to increase its sensitivity to 17 β -estradiol during development of the immune response (Liarte et al., 2011b). In long-term leukocyte cell lines of monocytes/macrophages from channel catfish, the expression of both ER α and ER β has been described (Iwanowicz & Ottinger, 2009). Although evidence conclusively demonstrates that fish leukocytes express ER genes, the literature in this respect does not deal with the possible differential roles of the two ER β genes (ER β 1 and ER β 2) present in fish. Our data in the gilthead seabream demonstrate for the first time that ER β 1 and ER β 2 are differentially regulated in macrophages. Thus, ER β 1 gene expression is only induced by VaDNA and its VaDNA-induced expression is slightly increased by 17 β -estradiol, in

contrast to ER β 2 gene whose expression is induced by both stimuli, which, moreover, have a synergic effect on ER β 2 gene expression (Liarte et al., 2011b).

The biological effect of 17 β -estradiol on fish head-kidney macrophages is mainly anti-inflammatory, although controversial data were observed depending on whether the studies were *in vivo* or *in vitro*. Intra-peritoneal injections of 17 β -estradiol in common carp inhibit phagocytosis and the production of reactive oxygen intermediates and reactive nitrogen intermediates by head-kidney macrophages in a dose-dependent manner (Watanuki et al., 2002). However, upon *in vitro* treatment, these head-kidney macrophages only showed impaired phagocytic capability (Yamaguchi et al., 2001) and, in goldfish macrophages, 17 β -estradiol inhibited the percentage of phagocytic cells (Wang & Belosevic, 1995). In the European flounder, microarray studies have revealed that 17 β -estradiol suppresses immune system-related transcripts in liver (Williams et al., 2007). In rainbow trout, 17 β -estradiol repressed the acute phase immune response genes (Tilton et al., 2006), as occurs in mammalian macrophages (Kramer & Wray, 2002). However, in gilthead seabream macrophages, 17 β -estradiol up-regulates some genes coding for key immune molecules, including inflammatory and anti-inflammatory molecules, innate immune receptors, molecules related to leukocyte infiltration, matrix metalloproteinases (MMP) and the anti-viral molecule *Myxovirus (influenza)* resistance protein (Mx). Moreover, the soluble factors produced by those 17 β -estradiol-stimulated macrophages modify the immune functions of head-kidney leukocytes (Liarte et al., 2011b), suggesting that the soluble factors produced by testicular macrophages in response to 17 β -estradiol contribute by blocking the phagocytic activity of testicular acidophilic granulocytes (Chaves-Pozo et al., 2005b). A suppression subtractive library was constructed to isolate and identify mRNA species up-regulated by a supra-physiological dose of 17 β -estradiol (50 ng/ml) to macrophages. Interestingly, this showed that 4% of up-regulated genes are related with the immune response, 6% with the stimulus response and 0.5% with physiological interactions between different organism categories, all of them probably involved in the interaction of immune cells with the immune stimulus. Although the number of identified genes within these categories was relatively low, other well-represented subcategories such as these related with biological regulation could contain genes whose functions may influence the behaviour of macrophages and thus affect their ability to respond to an immunological challenge upon exposure to estrogens (Liarte et al., 2011a; Xia & Yue, 2010).

Although less data are available for AR than for ER in mammalian species, several studies have demonstrated that testosterone alters macrophage functions in a complex manner, since it has both pro-inflammatory and anti-inflammatory effects. For example, wound healing is impaired in males, especially the elderly, which has been directly linked to a pro-inflammatory action of testosterone on tissue macrophages in the skin (Ashcroft & Mills, 2002). Moreover, castration increased macrophage-mediated damage at sites of injury in the skin, suggesting an anti-inflammatory role for testosterone (Ashcroft & Mills, 2002). Testosterone also inhibits inducible nitric oxide synthase and nitric oxide production in a mouse macrophage cell line (Friedl et al., 2000). The expression of AR in microglia, the brain macrophage, increases after injury and indicates that the innate immune cells of the brain may be modulated by androgens (García-Ovejero et al., 2002). Other data indicate that 5 α -dihydrotestosterone acts as an anti-inflammatory agent and depresses both nitric oxide and TNF α production in a dose-dependent fashion. However, testosterone treatment of microglia and peritoneal macrophages increased supernatant nitrite levels, suggesting a pro-inflammatory effect (Brown & Angel, 2005).

In fish, the data obtained show that androgens are also able to modulate the immune system responses. In common carp, intraperitoneal injections of 11-ketotestosterone inhibit phagocytosis and the production of reactive oxygen intermediates and reactive nitrogen intermediates by head-kidney macrophages in a dose-dependent manner (Watanuki et al., 2002). However, *in vitro* studies with head-kidney macrophages have demonstrated that this hormone inhibits phagocytosis and the production of reactive nitrogen intermediates and has no effect on the production of reactive oxygen intermediates (Yamaguchi et al., 2001). Interestingly, although gilthead seabream macrophages do not express the AR at a level detectable by real time polymerase chain reaction, both testosterone and 11-ketotestosterone up-regulated different immune genes, such as immune receptors and pro-inflammatory cytokines, and down-regulated the anti-inflammatory cytokine, transforming growth factor (TGF) β (Águila et al., 2011). Taking into account the complexity of sex steroid hormone signalling through intracellular and membrane receptors and sex steroid hormone conversion through transformation in other derivatives (such as reduced derivatives or even 17 β -estradiol) and bearing in mind that both testosterone and 11-ketotestosterone alter the macrophage gene expression and functions analyzed, it can not be discounted that macrophages convert testosterone into 11-ketotestosterone or another molecule capable of signalling through other receptors in this cell type. In this sense, mammalian macrophages lack AR but are able to respond to androgens through a membrane AR that triggers a Ca²⁺ influx (Benten et al., 2004). Moreover, mammalian testicular macrophages have a steroidogenic capability as they are able to produce and secrete 25-hydroxycholesterol, which affects Leydig cell steroidogenesis (Hales, 2002). In light of the above, further studies are needed to complete our understanding of the effect of androgens on fish innate immunity and macrophages.

2.2 Acidophilic granulocytes

The acidophilic granulocytes of gilthead seabream display some functions similar to human neutrophils despite their opposite staining pattern. In brief, they are the most abundant circulating granulocytes and are recruited from the head-kidney to the site of inflammation (Chaves-Pozo et al., 2004, 2005c), where they attach themselves to, internalize and kill bacteria through the production of reactive oxygen intermediates (Chaves-Pozo et al., 2004, 2005c; Meseguer et al., 1994; Sepulcre et al., 2002). However, they also show a monocyte/macrophage-like behaviour as they are able to specifically target a tissue and respond to physiological stimuli by displaying modified functions, as do the monocytes/macrophages of mammals (Chaves-Pozo et al., 2005c; Stout & Suttles, 2004). In fact, gilthead seabream acidophilic granulocytes infiltrate the testis in a way that resembles an inflammatory process triggered by physiological stimuli, and their main activities are strongly inhibited by the testicular microenvironment in order to preserve reproductive functions (Chaves-Pozo et al., 2005b). Previous data showed that 17 β -estradiol is related *in vivo* with the mobilization of acidophilic granulocytes from the head-kidney to the gonad and probably with the degree of this infiltration (Chaves-Pozo et al., 2007, 2008a). Interestingly, neither testicular nor head-kidney acidophilic granulocytes express any of the ER known in the gilthead seabream (Pinto et al., 2006; Liarte et al., 2011b). However, studies performed with conditioned medium from 17 β -estradiol-treated macrophages suggest that some, but not all, the acidophilic granulocyte functions modified by the testicular microenvironment might be regulated by the factors produced by 17 β -estradiol-treated

macrophages (Liarte et al., 2011b). In this sense, there is evidence that suggests a pro-inflammatory role for 17 β -estradiol in the gilthead seabream, since it is able to stimulate *in vivo* specific leukocyte migration and promote acidophilic granulocytes infiltration into the gonad (Chaves-Pozo et al., 2007). However, *in vitro*, 17 β -estradiol failed to promote chemotaxis in purified acidophilic granulocytes, although it is produced a positive migration of leukocytes when head-kidney suspensions were exposed to 17 β -estradiol (Liarte et al., 2011b).

In so far as androgens are concerned, acidophilic granulocytes constitutively express AR, the expression of which is modified by 11-ketotestosterone and testosterone, but only when the cells are co-stimulated with VaDNA (Águila et al., 2011). The effects of 11-ketotestosterone and testosterone on acidophilic granulocytes differ: while testosterone increased, 11-ketotestosterone decreased the expression of IL-1 β and toll-like receptors (TLRs), although both up-regulated the VaDNA-induced expression of IL-1 β (Águila et al., 2010).

2.3 Lymphocytes

T and B lymphocytes are the acknowledged cellular pillars of adaptative immunity. T cells are primarily responsible for cell-mediated immunity, while B lymphocytes are responsible for humoral immunity, but, in conjunction with other cell types, both mediate effective adaptive immunity (Pancer & Cooper, 2006). Recently, in long-term leukocyte cell lines of T-cells and B-cells from channel catfish, the differential expression of ER α and ER β was described. Thus, ER α is expressed in both cell types, while only T-cells express ER β 2 (Iwanowicz & Ottinger, 2009). In the gilthead seabream, lymphocytes only express the ER α gene (Liarte et al., 2011b). In mammals, however, B lymphocytes express both ER α and ER β genes, while there is debate as to whether or not T cells contain classical nuclear ER (Benten et al., 1998; Harkonen & Vaananen, 2006). *In vitro* functional assays demonstrated that 17 β -estradiol stimulates lymphocyte proliferation (Cook et al., 1994).

To determine whether fish lymphocytes respond to androgens, the classical chemical characterization of AR was performed in salmonid lymphocytes (Slater et al., 1995). In these species, 11-ketotestosterone inhibits lymphocyte proliferation, while testosterone reduces the number of antibody-producing cells and acts with cortisol to produce a greater inhibitory effect (Cook et al., 1994; Slater & Schreck, 1993).

2.4 Endothelial cells

Leukocyte recruitment is an early and pivotal event in any inflammatory response. Since gilthead seabream acidophilic granulocytes are recruited from the blood stream into the testis in a process that might be orchestrated by 17 β -estradiol (Chaves-Pozo et al., 2005b, 2007, 2008a), we investigated the role of the endothelium in this process. Leukocyte-endothelial interactions are a special case of cell sorting, in which the endothelium discriminates between circulating leukocytes in order to select cells for transmigration into surrounding tissue (Ebnet et al., 1996a). Endothelial cells play a singular role in this process, receiving information from the underlying tissue and transforming it into information that can be read rapidly by the passing leukocytes (Ebnet et al., 1996b). Accumulated evidence on mammalian models of cardiovascular disease points to the prominent role of estrogens in the ability of endothelial cells to trigger inflammation and participate in the leukocyte infiltration process (Nilsson, 2007; Straub, 2007). In mammals, endothelial cells constitutively express both ER α and ER β , although ER α plays a prominent role in the

vascular physiology (Ihionkhan et al., 2002; Straub, 2007). Gilthead seabream endothelial cells constitutively express ER α and ER β 1 but not ER β 2 (Liarte et al., 2011c). However, few studies have been carried out into the effect of 17 β -estradiol on endothelial cell physiology in fish. In the Japanese eel, 17 β -estradiol stimulated the production of vascular endothelial cell growth factor in endothelial cells (Huang et al., 2006). In the gilthead seabream endothelial cell cultures, 17 β -estradiol induced the expression of genes coding for chemokines, adhesion molecules and MMPs, which agrees with previous studies that demonstrated that 17 β -estradiol promotes acidophilic granulocyte infiltration into the testis (Chaves-Pozo et al., 2007). These effects contrast with that which occurs in mammals, where 17 β -estradiol inhibits *in vivo* the migration of leukocytes into inflamed areas and exerts tissue-protective activities through the down-regulation of adhesion molecules and the proforms of MMPs (Straub, 2007). On the other hand, 17 β -estradiol did not affect the expression in endothelial cells of the genes encoding major pro-inflammatory cytokines, such as IL-1 β , IL-6 and TNF α , which may prevent the detrimental effects of 17 β -estradiol-induced inflammation through leukocyte recruitment (Liarte et al., 2011c).

Little is known about androgens and their receptors in fish endothelial cells. There are very recent studies that suggest that androgens influence fish endothelial cell physiology, although further effort is needed to really understand how androgens affect endothelial cells and their molecular pathways. Trout testicular endothelial cells possess AR, as located by immunocytochemistry (Galas et al., 2009). In the gilthead seabream, recent studies determined that testosterone up-regulated TNF α , cyclooxygenase 2 (Cox2) and IL-1 β , and down-regulated TGF β and aromatase (the enzyme that transforms testosterone into 17 β -estradiol) gene expression (Águila et al., 2011).

3. Effects of endocrine disruptors on the immune response

In several species, the affinity for ER of several agonists, including natural estrogens like 17 β -estradiol, estrone or estriol, and estrogenic disruptor compounds, like 17 α -ethynilestradiol or diethylstilbestrol, has been tested. The different types of ER show differential binding preferences for ligands and their expression patterns are tissue-dependant (Iwanowicz & Ottinger, 2009). Taking into account that ER and AR are widely distributed in immune tissues, including the spleen, liver and anterior kidney (Lynn et al., 2008; Shved et al., 2009; Slater et al., 1995; Todo et al., 1999), the study of endocrine disruptor compounds as potential aquatic pollutants has taken on some importance for fish immunologists. Several anatomical and morphological changes were observed in lymphoid tissues following exposure to xenoestrogens and xenoandrogens. Spleno-somatic and hepato-somatic indices and thymus volume are affected by exposure to sex-steroids (androgens and estrogens) or to their related endocrine disruptor compounds (Grinwis et al., 2009; Kurtz et al., 2007; Tellez-Banuelos et al., 2009; van Ginneken et al., 2009). In the gilthead seabream the dietary intake of 17 α -ethynilestradiol promotes the up-regulation of several genes related with leukocyte recruitment (e.g. E-selectin (sele), the CC chemokine-like 4 (CCL4), TNF α and IL-8). Moreover, the heavy chain of IgM and IgT genes has also been seen to be up-regulated (Cabas et al., 2011). An increase in the spleno-somatic index was also recorded.

Surprisingly, *in vitro* 17 α -ethynilestradiol treatment of gilthead seabream endothelial cells dramatically reduces the expression of chemokines, adhesion molecules and MMPs in

VaDNA-activated endothelial cells unlike in 17β -estradiol-treated endothelial cells (see point 2.4). Although, the differential expression profile in stimulated 17α -ethynilestradiol-treated endothelial cells, compared with 17β -estradiol-treated endothelial cells, indicates that this compound would be able to impair the recruitment and activation of fish leukocytes, other molecular pathways might promote an inflammatory process in the gonad *in vivo*, as described by Cabas et al. (2011). These data show the complex effect of endocrine disruptor compounds on immune functions and the need to deepen our knowledge of their molecular action mechanism. As also occurs in mammals, 17β -estradiol, but not 17α -ethynilestradiol, significantly enhances nitric oxide production in gilthead seabream endothelial cells, indicating that some estrogens regulate nitric oxide production by endothelial cells from fish to mammals (Arnal et al., 1996; Liarte et al., 2011c; Nilsson, 2007). As far as we know, most studies on this topic have dealt with the effects of estrogenic and anti-androgenic disruptor compounds on reproductive functions. It is known that these disruptor compounds mainly affect several enzymes in the steroidogenic pathway, such as 20β -hydroxysteroid deshydrogenase, 17β -hydroxysteroid deshydrogenase and 11β -hydroxysteroid deshydrogenase, aromatase and 5α -reductase (Rempel & Schlenk, 2008). Further studies are needed into androgenic disruptor compounds as well as into estrogenic, anti-androgenic and androgenic disruptor compound mixtures to better understand how chemically and pharmaceutically polluted water might affect the reproductive and immune function of fish. Future studies and analyses along these times are being undertaken in our laboratory.

4. Conclusion

It is known that both estrogens and androgens modulate the fish immune response, although the molecular mechanisms by which they act are not completely understood. *In vivo* and *in vitro* analyses have demonstrated that gilthead seabream leukocyte (macrophages, acidophilic granulocytes and lymphocytes) express intracellular AR and/or ER, whose expression pattern upon stimulation depend on the cell type and the stimuli in question. Estrogens and androgens compromise the immune response, affecting cell types other than leukocytes. Thus, endothelial cells are involved in the leukocyte trafficking that occurs during the inflammatory process and their activities are also modulated by sex steroids. A wide variety of chemicals discharged from industrial and municipal sources has been reported to disrupt the endocrine system of animals via the food chain and contaminated water. Some of these contaminants have a widespread presence in the aquatic environment. Although, current knowledge concerning the sensitivity of marine fish to estrogenic and androgenic chemical in the environment is limited, we have seen that the most widespread (estrogenic) disruptor compound drastically affects leukocyte trafficking and recruitment into tissues. The short time of exposure (3 hours) used in our *in vitro* experiments suggests that, together with ER and AR activation, some transcription-independent non-genomic actions might be acting on sex steroid hormones-stimulated leukocytes. Taking all this into account, further effort will focus on the cloning and characterization of membrane AR and ER, their expression pattern in immune cells and the molecular characterization of the way of which estrogenic and androgenic compounds disrupt the molecular signalling pathways of intracellular and membrane androgen and estrogen receptors.

5. Acknowledgments

This work was supported by the Fundación Séneca, Coordination Center for Research, CARM (proyect 04538/GERM/06 to A.G.A and grant to I.C.), the Spanish Ministry of Science and Innovation (contract RYC-2009-05451 to E.C.P., project AGL2008-04575-C02-01 to A.G.A.). We thank the “Servicio de Apoyo a la Investigación” of the University of Murcia for their assistance with cell culture and gene expression analysis and the “Centro Oceanográfico de Murcia, Instituto Español de Oceanografía” for their assistance with fish care.

6. References

- Águila S.; Castillo-Briceño P.; Mulero V.; Meseguer J. & García-Ayala A. (2010). Testosterone modulates the immune response in vivo of the teleost fish gilthead seabream. First symposium of the European Organisation of Fish Immunology (EOFFI). Viterbo, Italy.
- Águila S.; Sánchez M.; Castillo Briceño P.; García Alcázar A.; Meseguer J.; Mulero V. & García Ayala A. (2011). Androgens modulate *in vitro* the activation of professional phagocytes from the bony fish gilthead seabream. GIA 2011. Heraklion, Crete, Greece.
- Arnal J. F.; Tack I.; Besombes J. P.; Pipy B. & Negre-Salvayre A. (1996). Nitric oxide and superoxide anion production during endothelial cell proliferation. American Journal of Physiology, Vol.271, No.5, (November 1996), pp. 1521-1526, ISSN 0002-9513.
- Ashcroft G. S. & Mills S. J. (2002). Androgen receptor-mediated inhibition of cutaneous wound healing. Journal of Clinical Investigation, Vol.110, No.5, (September 2002), pp. 615-624, ISSN 0021-9738.
- Beato M. & Klug J. (2000). Steroid hormone receptors: an update. Human Reproduction Update, Vol.6, No.3, (May-June 2000), pp. 225-236, ISSN 1355-4786.
- Benten W. P.; Guo Z.; Krucken J. & Wunderlich F. (2004). Rapid effects of androgens in macrophages. Steroids, Vol.69, No.8-9, (August 2004), pp. 585-590, ISSN 0039-128X.
- Benten W. P.; Lieberherr M.; Giese G. & Wunderlich F. (1998). Estradiol binding to cell surface raises cytosolic free calcium in T cells. FEBS Letters, Vol.422, No.3, (February 1998), pp. 349-53, ISSN 0014-5793.
- Benten W. P.; Stephan C.; Lieberherr M. & Wunderlich F. (2001). Estradiol signaling via sequestrable surface receptors. Endocrinology, Vol.142, No.4, (April 2001), pp. 1669-1677, ISSN 0013-7227.
- Benten W. P.; Stephan C. & Wunderlich F. (2002). B cells express intracellular but not surface receptors for testosterone and estradiol. Steroids, Vol.67, No.7, (June 2002), pp. 647-654, ISSN 0039-128X.
- Besseau L. & Faliex E. (1994). Resorption of unemitted gametes in *Lithognathus mormyrus* (Sparidae, Teleostei): a possible synergic action of somatic and immune cells. Cell and Tissue Research, Vol.276, No.1, (April 1994), pp. 123-132, ISSN 0302-766X.
- Billar R. (1983). A quantitative analysis of spermatogenesis in the trout, *Salmo trutta fario*. Cell and Tissue Research, Vol, 230, No.3, (April 1983), pp. 495-502, ISSN 0302-766X.

- Brown P. A. & Angel J. B. (2005). Granulocyte-macrophage colony-stimulating factor as an immune-based therapy in HIV infection. *Journal of Immune Based Therapies and Vaccines*, Vol.3, No.1, (May 2005), pp. 3-10, ISSN 1476-8518.
- Bruslé-Sicard S. & Fourcault B. (1997). Recognition of sex-inverting protandric *Sparus aurata*: ultrastructural aspects. *Journal of Fish Biology*, Vol.50, (December 1997), pp. 1094-1103, ISSN 0022-1112.
- Butts C. L.; Jones Y. L.; Lim J. K.; Salter C. E.; Belyavskaya E. & Sternberg E. M. (2011). Tissue expression of steroid hormone receptors is associated with differential immune responsiveness. *Brain, Behaviour and Immunity*, Vol.25, No.5, (July 2011), pp. 1000-1007, ISSN 1090-2139.
- Cabas I.; Liarte S.; Meseguer J.; Mulero V. & García-Ayala A. (2010). 17 α -ethynylestradiol *in vivo* treatment alters inflammatory gene expression profile in the gilthead seabream (*Sparus aurata* L.). 9th International Congress on the Biology of Fish. Barcelona, Spain.
- Cabas I.; Chaves-Pozo E.; García-Alcazar A.; Meseguer J.; Mulero V. & García-Ayala A. (2011). Dietary intake of 17 alpha-ethynylestradiol promotes leukocytes infiltration in the gonad of the hermaphrodite gilthead seabream. *Molecular Immunology*. ISSN 0161-5890. DOI:10.1016/j.molimm.2011.05.019.
- Caviola E.; Dalla Valle L.; Belvedere P. & Colombo L. (2007). Characterisation of three variants of estrogen receptor beta mRNA in the common sole, *Solea solea* L. (Teleostei). *General and Comparative Endocrinology*, Vol.153, No.1-3, (August-September 2007), pp. 31-39, ISSN 0016-6480.
- Cook G. A.; Elliott D.; Metwali A.; Blum A. M.; Sandor M.; Lynch R. & Weinstock J. V. (1994). Molecular evidence that granuloma T lymphocytes in murine schistosomiasis mansoni express an authentic substance P (NK-1) receptor. *Journal of Immunology*, Vol.152, No. 4, (February 1994), pp. 1830-1835, ISSN 0022-1767.
- Cunningham M. & Gilkeson G. (2011). Estrogen receptors in immunity and autoimmunity. *Clinical Reviews in Allergy and Immunology*, Vol.40, No.1, (February 2011), pp. 66-73, ISSN 1559-0267.
- Cutolo M.; Capellino S.; Montagna P.; Ghiorzo P.; Sulli A. & Villaggio B. (2005). Sex hormone modulation of cell growth and apoptosis of the human monocytic/macrophage cell line. *Arthritis Research and Therapy*, Vol.7, No.5, (July 2005), pp. 1124-1132, ISSN 1478-6362.
- Cutolo M.; Montecucco C. M.; Cavagna L.; Caporali R.; Capellino S.; Montagna P.; Fazzuoli L.; Villaggio B.; Serio B. & Sulli A. (2006). Serum cytokines and steroidal hormones in polymyalgia rheumatica and elderly-onset rheumatoid arthritis. *Annals of the Rheumatic Diseases*, Vol.65, No.11, (November 2006), pp. 1438-1443, ISSN 0003-4967.
- Cutolo M. & Straub R. H. (2009). Insights into endocrine-immunological disturbances in autoimmunity and their impact on treatment. *Arthritis Research and Therapy*, Vol.11, No.2, (April 2009), pp. 218-225, ISSN 1478-6362.
- Cutolo M.; Sulli A.; Capellino S.; Villaggio B.; Montagna P.; Serio B. & Straub R. H. (2004). Sex hormones influence on the immune system: basic and clinical aspects in autoimmunity. *Lupus*, Vol.13, No.9, (September, 2004) pp. 635-638, ISSN 0961-2033.
- Chaves-Pozo, E., Liarte, S., Fernández-Alacid, L., Abellán, E., Meseguer, J., Mulero, V. & García-Ayala, A. (2008a). Pattern of expression of immune-relevant genes in the

- gonad of a teleost, the gilthead seabream (*Sparus aurata* L.). *Molecular Immunology*, Vol.45, No.10, (May 2008), pp.2998-3011, ISSN 0161-5890.
- Chaves-Pozo E.; Arjona F. J.; García-López A.; García-Alcázar A.; Meseguer J. & García-Ayala A. (2008b). Sex steroids and metabolic parameter levels in a seasonal breeding fish (*Sparus aurata* L.). *General and Comparative Endocrinology*, Vol.156, No.3, (May 2008), pp. 531-536, ISSN 0016-6480.
- Chaves-Pozo E.; Castillo-Briceño P.; García-Alcázar A.; Meseguer J.; Mulero V. & García-Ayala A. (2008c). A role for matrix metalloproteinases in granulocyte infiltration and testicular remodeling in a seasonal breeding teleost. *Molecular Immunology*, Vol.45, No.10, (May 2008), pp. 2820-2830, ISSN 0161-5890.
- Chaves-Pozo E.; Liarte S.; Vargas-Chacoff L.; García-López A.; Mulero V.; Meseguer J.; Mancera J. M. & García-Ayala A. (2007). 17Beta-estradiol triggers postspawning in spermatogenically active gilthead seabream (*Sparus aurata* L.) males. *Biology of Reproduction*, Vol.76, No.1, (January 2007), pp. 142-148, ISSN 0006-3363.
- Chaves-Pozo E.; Mulero V.; Meseguer J. & García-Ayala A. (2005a). An overview of cell renewal in the testis throughout the reproductive cycle of a seasonal breeding teleost, the gilthead seabream (*Sparus aurata* L.). *Biology of Reproduction*, Vol.72, No.3, (March 2005), pp. 593-601, ISSN 0006-3363.
- Chaves-Pozo E.; Mulero V.; Meseguer J. & García-Ayala A. (2005b). Professional phagocytic granulocytes of the bony fish gilthead seabream display functional adaptation to testicular microenvironment. *Journal of Leukocyte Biology*, Vol.78, No.2, (August 2005), pp. 345-351, ISSN 0741-5400.
- Chaves-Pozo E.; Muñoz P.; López-Muñoz A.; Pelegrín P.; García-Ayala A.; Mulero V. & Meseguer J. (2005c). Early innate immune response and redistribution of inflammatory cells in the bony fish gilthead seabream experimentally infected with *Vibrio anguillarum*. *Cell and Tissue Research*, Vol.320, No.1, (April 2005), pp. 61-68, ISSN 0302-766X.
- Chaves-Pozo E.; Pelegrín P.; García-Castillo J.; García-Ayala A.; Mulero V. & Meseguer J. (2004). Acidophilic granulocytes of the marine fish gilthead seabream (*Sparus aurata* L.) produce interleukin-1beta following infection with *Vibrio anguillarum*. *Cell and Tissue Research*, Vol.316, No.2, (May 2004), pp. 189-195, ISSN 0302-766X.
- Chaves-Pozo E.; Pelegrín P.; Mulero V.; Meseguer J. & García-Ayala A. (2003). A role for acidophilic granulocytes in the testis of the gilthead seabream (*Sparus aurata* L., Teleostei). *Journal of Endocrinology*, Vol.179, No.2, (November 2003), pp. 165-174, ISSN 0022-0795.
- Christ M.; Haseroth K.; Falkenstein E. & Wehling M. (1997). Nongenomic steroid actions: fact or fantasy? *Vitamines and Hormones*, Vol.57, pp. 325-373, ISSN 0083-6729.
- Deane E. E.; Li J. & Woo N. Y. (2001). Hormonal status and phagocytic activity in sea bream infected with vibriosis. *Comparative Biochemistry and Physiology. Part B, Biochemistry and Molecular Biology*, Vol.129, No.2-3, (June 2001), pp. 687-693, ISSN 1096-4959.
- Dijkstra P. D.; Hekman R.; Schulz R. W. & Groothuis T. G. G. (2007). Social stimulation, nuptial colouration, androgens and immunocompetence in a sexual dimorphic cichlid fish. *Behavioural Ecology and Sociobiology*, Vol.61, No.2, (February 2007), pp. 599-609, ISSN 0340-5443.

- Ebnet K.; Kaldjian E. P.; Anderson A. O. & Shaw S. (1996a). Orchestrated information transfer underlying leukocyte endothelial interactions. *Annual Review of Immunology*, Vol.14, (April 1996), pp. 155-177, ISSN 0732-0582.
- Ebnet K.; Simon M. M. & Shaw S. (1996b). Regulation of chemokine gene expression in human endothelial cells by proinflammatory cytokines and *Borrelia burgdorferi*. *Annals of the New York Academy of Sciences*, Vol.797, (October 1996), pp. 107-117, ISSN 0077-8923.
- Engelsma M. Y.; Huising M. O.; van Muiswinkel W. B.; Flik G.; Kwang J.; Savelkoul H. F. & Verburg-van Kemenade B. M. (2002). Neuroendocrine-immune interactions in fish: a role for interleukin-1. *Veterinary Immunology*, Vol.87, No.3-4, (September 2002), pp. 467-479, ISSN 0165-2427.
- Evans W. H. & Bergeron J. J. (1988). Nuclear receptors: a re-evaluation. *Trends in Biochemical Sciences*, Vol.13, No.1, (January 1988), pp. 7-8, ISSN 0968-0004.
- Falkenstein E.; Tillmann H. C.; Christ M.; Feuring M. & Wehling M. (2000). Multiple actions of steroid hormones, a focus on rapid, nongenomic effects. *Pharmacological Reviews*, Vol.52, No.4, (December 2000), pp. 513-556, ISSN 0031-6997.
- Friedl R.; Brunner M.; Moeslinger T. & Spieckermann P. G. (2000). Testosterone inhibits expression of inducible nitric oxide synthase in murine macrophages. *Life Sciences*, Vol.68, No.4, (December 2000), pp. 417-429, ISSN 0024-3205.
- Galas J. F.; Hejmej A.; Glogowski J. & Bilinska B. (2009). Morphological and functional alterations in testes and efferent ducts of homogametic rainbow trout *Oncorhynchus mykiss* Walbaum. *Annals of the New York Academy of Sciences*, Vol.1163, (April 2009), pp. 398-401, ISSN 1749-6632.
- García-Ovejero D.; Veiga S.; García-Segura L. M. & DonCarlos L. L. (2002). Glial expression of estrogen and androgen receptors after rat brain injury. *The Journal of Comparative Neurology*, Vol.450, No.3, (August 2002), pp. 256-271, ISSN 0021-9967.
- Grinwis G. C.; Wester P. W. & Vethaak A. D. (2009). Histopathological effects of chronic aqueous exposure to bis(tri-n-butyltin)oxide (TBTO) to environmentally relevant concentrations reveal thymus atrophy in European flounder (*Platichthys flesus*). *Environmental Pollution*, Vol.157, No.10, (October 2009), pp. 2587-2593, ISSN 1873-6424.
- Guillemin G. J. & Brew B. J. (2004). Microglia, macrophages, perivascular macrophages, and pericytes: a review of function and identification. *Journal of Leukocyte Biology*, Vol.75, No.3, (March 2004), pp. 388-397, ISSN 0741-5400.
- Hales D. B. (2002). Testicular macrophage modulation of Leydig cell steroidogenesis. *Journal of Reproductive Immunology*, Vol.57, No.1-2, (October-November 2002), pp. 3-18, ISSN 0165-0378.
- Harkonen P. L. & Vaananen H. K. (2006). Monocyte-macrophage system as a target for estrogen and selective estrogen receptor modulators. *Annals of New York Academy of Sciences*, Vol.1089, (November 2006), pp. 218-227, ISSN 0077-8923.
- Harris J. & Bird D. J. (2000). Modulation of the fish immune system by hormones. *Veterinary Immunology and Immunopathology*, Vol.77, No.3-4, (December 2000), pp. 163-176, ISSN 0165-2427.

- Hecker M. & Karbe L. (2005). Parasitism in fish: an endocrine modulator of ecological relevance? *Aquatic Toxicology*, Vol.72, No.3, (April 2005), pp. 195-207, ISSN 0166-445X.
- Hedger M. P. (1997). Testicular leukocytes: what are they doing? *Review of Reproduction*, Vol.2, No.1, (January 1997), pp. 38-47, ISSN 1359-6004.
- Hedger M. P. (2002). Macrophages and the immune responsiveness of the testis. *Journal of Reproductive Immunology*, Vol.57, No.1-2, (October-November 2002), pp. 19-34, ISSN 0165-0378.
- Henderson N. E. (1962). The annual cycle in the testis of eastern brook trout *Salvelinus fontinalis* (Mitchell). *Canadian Journal of Zoology*, Vol.40, No. 4, pp. 631-641, ISSN 0008-4301.
- Hinteman T.; Schneider C.; Scholer H. F. & Schneider R. J. (2006). Field study using two immunoassays for the determination of estradiol and ethinylestradiol in the aquatic environment. *Water Research*, Vol.40, No.12, (July 2006), pp. 2287-2294, ISSN 0043-1354.
- Huang Y. S.; Huang W. L.; Lin W. F.; Chen M. C. & Jeng S. R. (2006). An endothelial-cell-enriched primary culture system to study vascular endothelial growth factor (VEGF A) expression in a teleost, the Japanese eel (*Anguilla japonica*). *Comparative Biochemistry and Physiology. Part A, Molecular and Integrative Physiology*, Vol.145, No.1, (September 2006), pp. 33-46, ISSN 1095-6433.
- Ihionkhan C. E.; Chambliss K. L.; Gibson L. L.; Hahner L. D.; Mendelsohn M. E. & Shaul P. W. (2002). Estrogen causes dynamic alterations in endothelial estrogen receptor expression. *Circulation Research*, Vol.91, No.9, (November 2002), pp. 814-820, ISSN 1524-4571.
- Iwanowicz L. R. & Ottinger C. A. (2009). Estrogens, estrogen receptors and their role as immunoregulators in fish, In: *Fish Defenses*, G. Zaccone; J. Meseguer; A. García-Ayala and B. G. Kapoor, (Eds), Vol. 1 Immunology, pp. 277-322, Science publisher, ISBN 978-1-57 808 -327 -5, Enfield.
- Janele D.; Lang T.; Capellino S.; Cutolo M.; Da Silva J. A. & Straub R. H. (2006). Effects of testosterone, 17beta-estradiol, and downstream estrogens on cytokine secretion from human leukocytes in the presence and absence of cortisol. *Annals of New York Academy of Sciences*, Vol.1069, (June 2006), pp. 168-182, ISSN 0077-8923.
- Johnson A. C.; Aerni H. R.; Gerritsen A.; Gibert M.; Giger W.; Hylland K.; Jurgens M.; Nakari T.; Pickering A.; Suter M. J.; Svenson A. & Wettstein F. E. (2005). Comparing steroid estrogen, and nonylphenol content across a range of European sewage plants with different treatment and management practices. *Water Research*, Vol.39, No.1, (January, 2005), pp. 47-58, ISSN 0043-1354.
- Klein S. L. (2004). Hormonal and immunological mechanisms mediating sex differences in parasite infection. *Parasite Immunology*, Vol.26, No.6-7, (June-July 2004), pp. 247-264, ISSN 0141-9838.
- Kramer P. R. & Wray S. (2002). 17-Beta-estradiol regulates expression of genes that function in macrophage activation and cholesterol homeostasis. *The Journal of Steroid Biochemistry and Molecular Biology*, Vol.81, No.3, (July 2002), pp. 203-216, ISSN 0960-0760.


- Kuch H. M. & Ballschmiter K. (2000). Determination of endogenous and exogenous estrogens in effluents from sewage treatment plants at the ng/L-level. *Fresenius Journal of Analytical Chemistry*, Vol.366, pp. 392-395, ISSN 1061-9348.
- Kumar M. V. & Tindall D. J. (1998). Transcriptional regulation of the steroid receptor genes. *Progress in Nucleic Acid Research and Molecular Biology*, Vol.59, pp. 289-306, ISSN 0079-6603.
- Kurtz J.; Kalbe M.; Langefors A.; Mayer I.; Milinski M. & Hasselquist D. (2007). An experimental test of the immunocompetence handicap hypothesis in a teleost fish: 11-ketotestosterone suppresses innate immunity in three-spined sticklebacks. *The American Naturalist*, Vol.170, No.4, (October 2007), pp. 509-519, ISSN 1537-5323.
- Lasserre G. (1972). Le coefficient de condition chez la daurade *Sparus auratus* L. 1758 de la région de Sète en 1971-1972. *Travaux du Laboratoire de Biologie Halieutique, Université de Rennes* 6, 141-150.
- Laskin D. L.; Weinberger B. & Laskin J. D. (2001). Functional heterogeneity in liver and lung macrophages. *Journal of Leukocyte Biology*, Vol.70, No.2, (August 2001), pp. 163-170, ISSN 0741-5400.
- Lehmann D.; Siebold K.; Emmons L. R. & Muller H. (1988). Androgens inhibit proliferation of human peripheral blood lymphocytes in vitro. *Clinical Immunology and Immunopathology*, Vol.46, No.1, (January 1988), pp. 122-128, ISSN 0090-1229.
- Liarte S.; Chaves-Pozo E.; Abellán E.; Meseguer J.; Mulero V.; Canario A. V. & García -Ayala A. (2011a). Estrogen-responsive genes in macrophages of the bony fish gilthead seabream: A transcriptomic approach. *Developmental and Comparative Immunology*, Vol.35, No.8 (March 2011), pp. 840-849, ISSN 1879-0089.
- Liarte S.; Chaves-Pozo E.; Abellán E.; Meseguer J.; Mulero V. & García-Ayala A. (2011b). 17beta-Estradiol regulates gilthead seabream professional phagocyte responses through macrophage activation. *Developmental and Comparative Immunology*, Vol.35, No.1(January 2011), pp. 19-27, ISSN 1879-0089.
- Liarte S.; Cabas I.; Chaves-Pozo E.; Arizcun M.; Meseguer J.; Mulero V. & García-Ayala A. (2011c). Natural and synthetic estrogens modulate the inflammatory response in the gilthead seabream (*Sparus aurata* L.) through the activation of endothelial cells. *Molecular Immunology*, ISSN 0161-5890. DOI:10.1016/j.molimm.2011.05.019.
- Liarte S.; Chaves-Pozo E.; García-Alcázar A.; Mulero V.; Meseguer J. & García-Ayala A. (2007). Testicular involution prior to sex change in gilthead seabream is characterized by a decrease in DMRT1 gene expression and by massive leukocyte infiltration. *Reproductive Biology and Endocrinology*, Vol.5, (June 2007), pp. 20-35, ISSN 1477-7827.
- Lo Nostro F. L.; Antoneli F. N.; Quagio-Grassiotto I. & Guerrero G. A. (2004). Testicular interstitial cells, and steroidogenic detection in the protogynous fish, *Synbranchus marmoratus* (Teleostei, Synbranchidae). *Tissue and Cell*, Vol.36, No.4, (August, 2004), pp. 221-231, ISSN 0040-8166.
- Loir M.; Sourdain P.; Mendis-Handagama S. M. & Jegou B. (1995). Cell-cell interactions in the testis of teleosts and elasmobranchs. *Microscopy Research and Technique*, Vol.32, No. 6, (December 1995), pp. 533-552, ISSN 1059-910X.
- Lutton B. & Callard I. (2006). Evolution of reproductive-immune interactions. *Integrative and Comparative Biology*, Vol.46, No.6, (December 2006), pp. 1060-1071, ISSN 1540-7063.

- Lynn S. G.; Birge W. J. & Shepherd B. S. (2008). Molecular characterization and sex-specific tissue expression of estrogen receptor alpha (esr1), estrogen receptor beta a (esr2a) and ovarian aromatase (cyp19a1a) in yellow perch (*Perca flavescens*). *Comparative Biochemistry and Physiology. Part B, Biochemistry and Molecular Biology*, Vol.149, No.1, (January 2008), pp. 126-147, ISSN 1096-4959.
- Meseguer J.; Esteban M. A.; Lopez-Ruiz A. & Bielek E. (1994). Ultrastructure of nonspecific cytotoxic cells in teleosts. I. Effector-target cell binding in a marine and a freshwater species (seabream: *Sparus aurata* L., and carp: *Cyprinus carpio* L.). *The Anatomical Record*, Vol.239, No.4, (August 1994), pp. 468-474, ISSN 0003-276X.
- Micale V.; Perdichizzi F. & Santangelo G. (1987). The gonadal cycle of captive white bream, *Diplodus sargus* (L.). *Journal of Fish Biology*, Vol.31, (May 1987), pp. 435-440, ISSN 002-1112.
- Mor G.; Sapi E.; Abrahams V. M.; Rutherford T.; Song J.; Hao X. Y.; Muzaffar S. & Kohen F. (2003). Interaction of the estrogen receptors with the Fas ligand promoter in human monocytes. *Journal of Immunology*, Vol.170, No.1, (January 2003), pp. 114-122, ISSN 0022-1767.
- Nagler J. J.; Cavileer T.; Sullivan J.; Cyr D. G. & Rexroad III C., 3rd. (2007). The complete nuclear estrogen receptor family in the rainbow trout: discovery of the novel ER α 2 and both ER β isoforms. *Gene*, Vol.392, No.1-2, (May 2007), pp. 164-173, ISSN 0378-1119.
- Nilsson B. O. (2007). Modulation of the inflammatory response by estrogens with focus on the endothelium and its interactions with leukocytes. *Inflammation Research*, Vol.56, No.7, (July 2007), pp. 269-273, ISSN 1023-3830.
- Pancer Z. & Cooper M. D. (2006). The evolution of adaptive immunity. *Annual Review of Immunology*, Vol.24, (April 2006), pp. 497-518, ISSN 0732-0582.
- Pang Y.; Dong J. & Thomas P. (2008). Estrogen signaling characteristics of Atlantic croaker G protein-coupled receptor 30 (GPR30) and evidence it is involved in maintenance of oocyte meiotic arrest. *Endocrinology*, Vol.149, No.7, (July 2008), pp. 3410-3426, ISSN 0013-7227.
- Pinto P. I.; Passos A. L.; Martins R. S.; Power D. M. & Canário A. V. (2006). Characterization of estrogen receptor betab in sea bream (*Sparus auratus*): phylogeny, ligand-binding, and comparative analysis of expression. *General and Comparative Endocrinology*, Vol.145, No.2, (January, 2006), pp. 197-207, ISSN 0016-6480.
- Rempel M. A. & Schlenk D. (2008). Effects of environmental estrogens and antiandrogens on endocrine function, gene regulation, and health in fish. *International Review of Cell and Molecular Biology*, Vol.267, (June 2008), pp. 207-252, ISSN 1937-6448.
- Roberts M. L.; Buchanan K. L. & Evans M. R. (2004). Testing the immunocompetence handicap hypothesis: a review of the evidence. *Animal Behaviour*, Vol.68, No. 2, (August 2004), pp. 227-239, ISSN 0003-3472.
- Scott A. P. & Sumpter J. P. (1989). Seasonal variations in testicular germ cell stages and in plasma concentrations of sex steroids in male rainbow trout (*Salmo gairdneri*) maturing at 2 years old. *General and Comparative Endocrinology*, Vol.73, No.1, (January 1989), pp. 46-58, ISSN 0016-6480.
- Sepulcre M. P.; Pelegrín P.; Mulero V. & Meseguer J. (2002). Characterisation of gilthead seabream acidophilic granulocytes by a monoclonal antibody unequivocally points

- to their involvement in fish phagocytic response. *Cell and Tissue Research*, Vol.308, No.1, (April 2002), pp. 97-102, ISSN 0302-766X.
- Shrestha T. K. & Khanna S. S. (1976). Histology and seasonal changes in the testes of a hill-stream fish *Schizothorax plagiostomus*. *Zeitschrift für Mikroskopisch-Anatomische Forschung*, Vol.90, No.4, pp. 749-761, ISSN 0044-3107.
- Shved N.; Berishvili G.; Hausermann E.; D'Cotta H.; Baroiller J. F. & Eppler E. (2009). Challenge with 17alpha-ethinylestradiol (EE2) during early development persistently impairs growth, differentiation, and local expression of IGF-I and IGF-II in immune organs of tilapia. *Fish and Shellfish Immunology*, Vol.26, No.3, (March 2009), pp. 524-530, ISSN 1095-9947.
- Slater C. H.; Fitzpatrick M. S. & Schreck C. B. (1995). Characterization of an androgen receptor in salmonid lymphocytes: possible link to androgen-induced immunosuppression. *General and Comparative Endocrinology*, Vol.100, No.2, (November 1995), pp. 218-225, ISSN 0016-6480.
- Slater C. H. & Schreck C. B. (1993). Testosterone alters the immune response of chinook salmon, *Oncorhynchus tshawytscha*. *General and Comparative Endocrinology*, Vol.89, No.2, (February 1993), pp. 291-298, ISSN 0016-6480.
- Stout R. D. & Suttles J. (2004). Functional plasticity of macrophages: reversible adaptation to changing microenvironments. *Journal of Leukocyte Biology*, Vol.76, No.3, (September 2004), pp. 509-513, ISSN 0741-5400.
- Straub R. H. (2007). The complex role of estrogens in inflammation. *Endocrine Reviews*, Vol.28, No.5, (August 2007), pp. 521-574, ISSN 0163-769X.
- Tellez-Bañuelos M. C.; Santerre A.; Casas-Solis J.; Bravo-Cuellar A. & Zaitseva G. (2009). Oxidative stress in macrophages from spleen of Nile tilapia (*Oreochromis niloticus*) exposed to sublethal concentration of endosulfan. *Fish and Shellfish Immunology*, Vol.27, No.2, (August 2009), pp. 105-111, ISSN 1095-9947.
- Ternes T. A.; Stumpf M.; Mueller J.; Haberer K.; Wilken R. D. & Servos M. (1999). Behavior and occurrence of estrogens in municipal sewage treatment plants--I. Investigations in Germany, Canada and Brazil. *The Science of the Total Environment*, Vol.225, No.1-2, (January, 1999), pp. 81-90, ISSN 0048-9697.
- Thomas P.; Dressing G.; Pang Y.; Berg H.; Tubbs C.; Benninghoff A. & Doughty K. (2006). Progestin, estrogen and androgen G-protein coupled receptors in fish gonads. *Steroids*, Vol.71, No.4, (April 2006), pp. 310-316, ISSN 0039-128X.
- Tilton S. C.; Givan S. A.; Pereira C. B.; Bailey G. S. & Williams D. E. (2006). Toxicogenomic profiling of the hepatic tumor promoters indole-3-carbinol, 17beta-estradiol and beta-naphthoflavone in rainbow trout. *Toxicological Sciences*, Vol.90, No.1, (March 2006), pp. 61-72, ISSN 1096-6080.
- Todo T.; Ikeuchi T.; Kobayashi T. & Nagahama Y. (1999). Fish androgen receptor: cDNA cloning, steroid activation of transcription in transfected mammalian cells, and tissue mRNA levels. *Biochemical and Biophysical Research Communications*, Vol.254, No.2, (January 1999), pp. 378-383, ISSN 0006-291X.
- van Ginneken V.; Palstra A.; Leonards P.; Nieveen M.; van den Berg H.; Flik G.; Spanings T.; Niemantsverdriet P.; van den Thillart G. & Murk A. (2009). PCBs and the energy cost of migration in the European eel (*Anguilla anguilla* L.). *Aquatic Toxicology*, Vol.92, No.4, (May 2009), pp. 213-220, ISSN 1879-1514.

- Wang R. & Belosevic M. (1995). The in vitro effects of estradiol and cortisol on the function of a long-term goldfish macrophage cell line. *Developmental and Comparative Immunology*, Vol.19, No.4, (July-August 1995), pp. 327-336, ISSN 0145-305X.
- Watanuki H.; Yamaguchi T. & Sakai M. (2002). Suppression in function of phagocytic cells in common carp *Cyprinus carpio* L. injected with estradiol, progesterone or 11-ketotestosterone. *Comparative Biochemistry and Physiology. Part C, Pharmacology, Toxicology and Endocrinology*, Vol.132, No.4, (August 2002), pp. 407-413, ISSN 1532-0456.
- Williams T. D.; Diab A. M.; George S. G.; Sabine V. & Chipman J. K. (2007). Gene expression responses of European flounder (*Platichthys flesus*) to 17-beta estradiol. *Toxicology Letters*, Vol.168, No.3, (February 2007), pp. 236-248, ISSN 0378-4274.
- Xia J. H. & Yue G. H. (2010). Identification and analysis of immune-related transcriptome in Asian seabass *Lates calcarifer*. *BMC Genomics*, Vol.11, (June 2010), pp. 356-368, ISSN 1471-2164.
- Xia Z.; Gale W. L.; Chang X.; Langenau D.; Patino R.; Maule A. G. & Densmore L. D. (2000). Phylogenetic sequence analysis, recombinant expression, and tissue distribution of a channel catfish estrogen receptor beta. *General and Comparative Endocrinology*, Vol.118, No.1, (April 2000), pp. 139-149, ISSN 0016-6480.
- Yamaguchi T.; Watanuki H. & Sakai M. (2001). Effects of estradiol, progesterone and testosterone on the function of carp, *Cyprinus carpio*, phagocytes in vitro. *Comparative Biochemistry and Physiology. Part C, Pharmacology, Toxicology & Endocrinology*, Vol.129, No.1, (May 2001), pp. 49-55, ISSN 1532-0456.
- Zaitzu M.; Narita S.; Lambert K. C.; Grady J. J.; Estes D. M.; Curran E. M.; Brooks E. G.; Watson C. S.; Goldblum R. M. & Midoro-Horiuti T. (2007). Estradiol activates mast cells via a non-genomic estrogen receptor-alpha and calcium influx. *Molecular Immunology*, Vol.44, No.8, (March 2007), pp. 1977-1985, ISSN 0161-5890.

IntechOpen



Sex Steroids

Edited by Dr. Scott M. Kahn

ISBN 978-953-307-857-1

Hard cover, 330 pages

Publisher InTech

Published online 27, January, 2012

Published in print edition January, 2012

This book, entitled "Sex Steroids", features a valuable collection of reviews and research articles written by experts in signal transduction, cellular biology, diseases and disorders. "Sex Steroids" is comprised of four sections, "The Biology of Sex Steroids", "Sex Steroids, Memory, and the Brain", "Sex Steroids and the Immune Response", and "Therapy"; individual chapters address a broad range of recognized and predicted functions and applications of sex steroids. "Sex Steroids" is intended to provide seasoned veterans as well as newcomers to this area of research with informative, resourceful, and provocative insights. Readers of "Sex Steroids" should emerge with an appreciation and understanding of the multitude and complexity of biologic processes attributed to these important hormones, and possible future directions of research in this fascinating and ever evolving field.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Elena Chaves-Pozo, Isabel Cabas and Alfonsa García-Ayala (2012). Sex Steroids Modulate Fish Immune Response, Sex Steroids, Dr. Scott M. Kahn (Ed.), ISBN: 978-953-307-857-1, InTech, Available from: <http://www.intechopen.com/books/sex-steroids/sex-steroids-modulate-fish-immune-response>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](https://creativecommons.org/licenses/by/3.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen