We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900
Open access books available

185,000
International authors and editors

200M
Downloads

Our authors are among the

154
Countries delivered to

TOP 1%
most cited scientists

12.2%
Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



7

Inflammatory Periprosthetic Bone Loss

Sang-Soo Lee[1], P. Edward Purdue[2] and Ju-Suk Nam[1]
[1]*Institute for Skeletal Aging & Orthopedic Surgery,*
Infectious Disease Medical Research Center,
Hallym University College of Medicine, Chuncheon,
[2]*Osteolysis Research Laboratory,*
Hospital for Special Surgery, New York,
[1]*Korea,*
[2]*US*

1. Introduction

Total hip arthroplasty [THA] is one of the most successful and effective procedures developed for the treatment of pain and lack of mobility associated with end-stage arthritis such as osteoarthritis and rheumatoid arthritis. Approximately 1.5 million joint arthroplastic operations are performed annually worldwide. THA, although considered an excellent surgical procedure, can be complicated by periprosthetic osteolysis. Periprosthetic osteolysis (also called 'Particle disease') is initiated by wear debris derived from the implant. In most long-term studies on hip arthroplasty, osteolysis related loosening, bone loss or periprosthetic fractures are the most frequent causes for revision surgeries (Talmo et al., 2006).

Osteolysis is a particle-induced biologic process at the metal–bone or cement–bone interface of prosthetic implants, manifesting radiographically as scalloped focal or linear endosteal radiolucencies due to bone loss and resulting in the loosening of implants. In the early days of hip arthroplasty, radiolucencies around implants were noticed and were thought to be related to curing of acrylic cement, infection or neoplastic process. These were first described by Charnley in association with Teflon cups, though later were also observed in patients with stable implants (Charnley, 1966). In 1977, Willert and Semlitsch demonstrated the presence of macrophages in response to wear debris and concluded that the particles accumulate macrophages in pericapsular lymph drainage, leading to a foreign body response and eventual loosening of the implant (Willert, 1977).

Goldring *et al.* described the synovial-like character of the interfacial membrane found and demonstrated the presence of prostaglandin E_2 [PGE_2] and collagenase secretion from the associated cells (Goldring et al., 1983). The early observations of osteolysis in cemented implants led to a general belief that osteolysis was related to the acrylic cement and the term 'Cement disease' was introduced. However, after the demonstration of lytic lesions in cementless implants, osteolysis is now considered to be a 'Particle disease', suggesting that wear-generated particulate debris is the main cause of periprosthetic osteolysis (Harris, 1995).

Biologic responses to implant debris, the basis of periprosthetic tissue destruction, are due to a wide variety of complex events. Aseptic failure occurs later as a secondary issue to the chronic granulomatous and inflammatory response, which is stimulated and maintained by

wear particles. This process is progressive and time dependant, ultimately leading to prosthetic loosening and failure (Keener et al., 2003). Wear debris can be generated from the articulating surfaces and bone cement. In general, higher wear rates are observed in patients with osteolysis (Dumbleton et al., 2002). However, the osteolytic process is a result of multiple factors, including physical and biologic components (Clohisy et al., 2004a).
Once macrophages are activated by particulate debris, they secrete various kinds of mediators to incite a complex cascade of events culminating in recruitment and maturation of osteoclasts, the bone resorbing cells directly responsible for the pathogenic bone loss in osteolysis (Glant et al., 1993). Other cell types also seem to be involved in production cytokines and inflammatory mediators during this process, such as osteoblasts and fibroblasts (Jacobs et al., 2001; Dorr et al., 1990). Matrix degradative enzymes and chemokines are also released from several types of cells (Jacobs et al., 2001; Takagi et al., 1998). The core of the biologic response that leads to osteolysis involves receptor activator of NF-κB ligand [RANKL]-RANK axis for osteoclast precursors, resulting in their differentiation and maturation (Abu-Amer, 2005; Khosla, 2001).

Category	Clinical manifestations
Soft tissue lesions	• Acute synovitis (Engler et al., 2001) • Particle-induced synovitis (Niki et al., 2007) • Heterotopic polyethylene granuloma (Walsh et al., 2011)
Osseous impairment	• Periprosthetic osteolysis (Lee et al., 2007) • Impaired osteogenesis (Wang et al., 2004) • Aseptic loosening (Harris, 1995) • Failure of implant (Clohisy et al., 2004)
Systemic reactions	• Metal hypersensitivity (Hallab et al., 2005)

Table 1. Clinical conditions related with wear debris-induced inflammation following total joint arthroplasty

Moreover, recent researches have uncovered the possibility that biological mechanism of osteolysis has to be extended to bone forming activity as well as resorption or dissolution of bone tissue. Recent datas suggest that bone-forming cells - osteoblasts, osteoprogenitors, and adult mesenchymal stem cells - may also contribute to osteolysis. As to date, there is no approved drug therapy to prevent or inhibit periprosthetic osteolysis, this concept will open up possibilities for the development of therapeutic agents that can enhance bone formation. This review presents novel insights into the current knowledge regarding how wear debris interact as an inflammatory process leading to periprosthetic osteolysis. The authors hope to outline potential perspectives for the future therapeutic strategies for this devastating complication.

2. Wear particle debris - the main cause of periprosthetic osteolysis

Wear-generated particulate debris is the main cause of periprosthetic osteolysis. Various kinds of cells have been implicated in the mechanisms leading to periprosthetic osteolysis in response to wear debris. They are indicative of a complex network of cellular pathogenesis (Drees et al., 2007). Several studies with retrieved implants, animal and *in-vitro* model suggest that wear-mediated periprosthetic osteolysis is unlikely to be caused solely by one particular cell type or particulate species, but is rather the cumulative consequence of a number of biological reactions (Wang et al., 2004).

Wear debris is formed at prosthetic joint articulations, modular interfaces, and nonarticulating interfaces (Goldring et al., 1993). The majority of particles are less than 5 um in diameter and exist in a range of shapes and sizes. Within a clinical context, polyethylene wear represents the dominant type of debris that leads to loss of prostheses. With regard to particle size, large particles are recognized as nondigestible foreign bodies. Particles within the broad size range of 0.2 – 10.0 μm are phagocytosed by macrophages leading to cellular activation. Although smaller particles are generally more pro-inflammatory, it is possible that extremely small submicron particles are less biologically active (Green et al., 1998). Particles beyond the size range of 0.2 – 10.0 μm can escape active phagocytosis, and fail to stimulate macrophages to produce high levels of proinflammatory and osteolytic cytokines. *In-vitro* studies of macrophage cultures clearly indicated that smaller [< 20 μm] polymethylmethacrylate [PMMA] and polyethylene particles [PE] elicited a significantly greater inflammatory cytokine response, as indicated by increased release of tumor necrosis factor [TNF-α], IL-1, IL-6, PGE_2, matrix metalloproteinases [MMPs], and other factors (Abbas et al., 2003; Gonzalez et al., 1996; Lee et al., 2003; O'Keefe et al., 1998; Shanbhag et al., 1994).

In addition to size of particles, the cellular response to wear debris depends on numerous other parameters of particles such as the composition (Haynes et al., 1998; Sethi et al., 2003), shape (Yang et al., 2002b), charge, number (Gonzalez et al., 1996; Sabokbar et al., 2003b), volume, and surface area (Shanbhag et al., 1994). Especially the amount of particle around implants exhibits a fair correlation with the severity of aseptic loosening, although certain cases shows an exaggerated biologic response to particulate debris (Abu-Amer et al., 2007). The relative numbers of particles and macrophages are also critical to the intensity of reaction. The extent of the reaction by macrophages was also affected by the particle: target cell ratio. Therefore, the association between particles and osteolysis represents a dose–response relationship (Wilkinson et al., 2005)

Interestingly, osteoblasts also can phagocyte small particles, causing potential adverse effects on viability, proliferation and function of osteoblast as well as on osteoclasts (Goodman et al., 2006; Lohmann et al., 2000). PE, PMMA or metallic particles reduce osteoblasts differentiation of bone marrow osteoprogenitor cells (Chiu et al., 2006), expression of collagens by osteoblasts (Vermes et al., 2001; Vermes et al., 2000), osteoblast viability by inducing apoptosis (Pioletti et al., 2002) characterized with decreased production of matrix, alkaline phosphatase and TGF-β by these cells (Dean et al., 1999). As for macrophages, such suppressive effects are also likely dependent on particle size, composition and dosage: different particle types can differentially affect osteoblast function (Lohmann et al., 2002).

The size and degree of clumping of particles are also important variables determining the biological response, especially in osteoblast. Smaller particles of nano-size have less detrimental effect on the functions of osteoblasts, compared to conventional particles (Granchi et al., 2005; Gutwein & Webster, 2004). The nano-sized particles were associated with increased cell viability, more normal cellular morphology and spreading compared to conventional particles, indicating nano-sized particles are less active (Gutwein & Webster, 2004). Therefore, roles of nano-sized wear debris in periprosthetic osteolysis deserve further testing.

3. Periprosthetic membrane in osteolysis around the implant

The tissue around osteolysis contains a synovial-like interface membrane between the prosthesis and the adjacent bone, called the periprosthetic membrane. Periprosthetic

membranes retrieved from patients contain macrophages, fibroblasts and multi nuclear giant cells such as osteoclasts. T lymphocytes and B lymphocytes are also seen.. The development of osteolysis is triggered by cellular and enzymatic processes within this membrane. The periprosthetic membrane is a histopathological hallmark of aseptic prosthesis loosening and shares some similarities with the hyperplastic synovium in patients with rheumatoid arthritis [RA] (Drees et al., 2007; Goldring et al., 1983; Harris, 1995). At a molecular level, RA synovial fibroblasts and prosthesis-loosening fibroblasts share several common features.




Fig. 1. Possible model of interplay between macrophages, fibroblasts, lymphocyte, osteoclasts and osteoblast in periprosthetic osteolysis. Osteoclasts develop from precursors under the influence of RANKL. The source of RANKL can be fibroblasts, osteoblasts, macrophages, or T cells. Particles may stimulate macrophage, fibroblasts and osteoblasts directly to induce RANKL and pro-inflammatory cytokines that can induce RANKL. It has been hypothesized that T cells stimulated by the pro-inflammatory microenvironment may also promote osteoclast formation, synergized with TNF-α, by secreting IL-17.Thus, RANKL, TNF-α, IL-1, IL-6, IL-17, and M-CSF may mediate the differentiation of myeloid precursor cells into multinucleated osteoclasts and development of impaired osteogenecity (Abu-Amer et al., 2007; Drees et al., 2007; Kotake et al., 1999; Tokuda et al., 2004)

Periprosthetic membranes, retrieved during revision surgery, produce a variety of factors including TNF-α, IL-1, IL-6, and PGs that are involved in mediating osteoclast biology (Chiba et al., 1994; Hirakawa et al., 1996; Jiranek et al., 1993; Margevicius et al., 1994; Shanbhag et al., 1995). These factors induce the final effector molecule, RANKL. It is generally accepted that

macrophage lineage cells do not express RANKL under normal conditions. In RA and periodontal disease, T cells have been known as the major source of RANKL. However, the relatively low numbers of T cells present near periprosthetic osteolysis make it unlikely that T cells are the major source of RANKL in periprosthetic osteolysis.
Studies of periprosthetic membranes of osteolysis patients revealed that fibroblast are the major source of RANKL (Haynes et al., 2004; Sakai et al., 2002), with possible involvement of macrophages and giant cells (Haynes et al., 2004; Sakai et al., 2002). Expression and secretion of MMPs are also elevated in macrophages exposed to wear debris in vitro. Elevated levels of degradation enzymes in periprosthetic osteolysis tissues were also observed (Kido et al., 2004). This array of chemokines, growth factors, pro-inflammatory and anti-inflammatory cytokines, and mediators demonstrate a potent ability of periprosthetic tissues to recruit and stimulate cells capable of inducing osteoclastic bone resorption and fibrous tissue formation (Talmo et al., 2006).

4. Inflammatory response in particle disease

The cellular response is dominated by macrophages (Archibeck et al., 2001; Lee et al., 2007; Neale & Athanasou, 1999; Quinn et al., 1992). Once macrophages are activated by particulate debris, they secrete various kinds of mediators to incite a complex cascade of events culminating in osteoclast maturation (Glant et al., 1993). This osteolytic response involves various cell types such as osteoclasts, fibroblasts, and osteoblasts/stromal cells, secreting a wide range of factors including cytokines, growth factors, and prostanoids (Dorr et al., 1990; Jacobs et al., 2001; Perry et al., 1995; Shanbhag et al., 1995). Matrix degradative enzymes and chemokines are also released from various cell types (Jacobs et al., 2001; Takagi et al., 1998).
Particle phagocytosis is the important component of the cellular response: the size of these particles is significant. Particles ranging from 0.2 to 10 μm in diameter undergo phagocytosis by macrophages (Gelb et al., 1994). The initial response of macrophage by particle is formation of fibrous tissue to encapsulate the implant. Often, synovial fluid and lining membranes are also formed, and granulomatous tissue is established. Such peripsrosthetic tissues have revealed an abundance of macrophages, fibroblasts and giant cells (Clohisy et al., 2004b; Ulrich-Vinther et al., 2002).
In addition, apart from massive recruitment of macrophages to the site of injury, some studies identified recruitment of lymphocytes (Abu-Amer, 2005; Arora et al., 2003; Gallo et al., 2002; Hallab et al., 2005; Lam et al., 2002; Purdue et al., 2007). Subsequently, pro-inflammatory response begins with secretion of factors, gelatinases, and proteases contributing to periprosthetic osteolysis, and thus causing failure of the implant (Abu-Amer et al., 2007). This inflammatory response is not restricted to the initial process, but rather it continues to appear in middle till late osteolytic stages of periprosthetic osteolysis (Abu-Amer et al., 2007).
Besides suppressing osteogenic activity, wear debris challenge can also affect the production of RANKL and OPG by osteoblasts. Osteoblast lineage cells can express RANKL, OPG, IL-1, TNF-α, IL-6, IL-11 and TGF-β (Hofbauer et al., 2000). Ultra high density molecular weight polyethylene [UHMWPE] increased the release of RANKL from human osteoblasts, while OPG was significantly inhibited. There was inductive also effects on the osteoclastogenesis with UHMWPE-human osteoblast-conditioned medium.
A study of the literature suggests that analysis of the involvement of osteoblasts in periprosthetic osteolysis has generally been limited to direct suppressive effect of particles on osteoblasts rather than through consideration of the possible effects of a pro-

inflammatory environment on osteoblast biology. Considering that TNF-α is also a potent inhibitor of osteoblast differentiation (Ghali et al., 2010; Yamazaki et al., 2009; Zhou et al., 2006; Karmakar et al. 2010), additional investigations into possible involvement of particle-activated macrophages in the impaired osteogencity mediated by proinflammatory cytokines including TNF-α would appear to be warranted. Although insufficient attention has been paid to the involvement of osteoblast, the cell type responsible for bone formation, more research should be conducted to delineate the potentially critical role of osteoblasts in periprosthetic osteolysis.

5. Roles of macrophages in particle disease

Since macrophages are the chief phagocytic cell in periprosthetic membranes, much attention has been focused on their role in cytokine production and osteoclast activation (Blaine et al., 1996; Nakashima et al., 1999b; Shanbhag et al., 1994). Macrophages are abundant in the periprosthetic tissues obtained from osteolysis patients, and are engaged in phagocytosis of wear particles as evidenced by the presence of such nondegradable particles within these cells. However, recent advances in osteoclast biology indicated that bone marrow-derived macrophages may play a dual role in periprosthetic osteolysis. First, as the major cell in host defense, they respond to particles through cytokine production. Second, macrophages have a role as precursors for the osteoclasts (Ingham & Fisher, 2005). Macrophages can phagocytose a variety of types of wear particles. Most notably, pro-inflammatory mediators such as PGE_2, TNF-α and IL-6 are generated in abundance by particle challenged macrophages.

Activation of macrophages by wear debris is a critical event in this process. It is believed that recognition of particles relies on phagocytosis of particles by macrophages and unidentified cell surface interactions. However, a little is known about the molecular mechanisms involved in particle recognition concerning the cell surface receptors that response to particles (Purdue, 2008). Although particle phagocytosis has been identified as a critical component of this biological response, recent studies in human macrophages indicate that direct interactions between particle and cell surface are sufficient to activate osteoclastogenic signaling pathways (Abu-Amer et al., 2007; Gallo et al., 2002; Gonzalez et al., 1996; Nakashima et al., 1999b). The latter interactions may include nonspecific physical induction of transmembrane proteins or recognition of cell surface molecules by particles. Recently this phenomenon was explained with the role of toll-like receptor (Takagi et al., 2007). However, the precise nature of stimulation of cells by particles remains unknown (Abu-Amer et al., 2007).

Recently, macrophages in periprosthetic space started to be defined as osteoclast precursors. *In-vitro* they have been shown to differentiate into osteoclasts in response to M-CSF and stromal cell-derived factors (Sabokbar et al., 1997): RANK ligand alone; or TNF-α and IL-1 in the absence of RANK ligand (Sabokbar et al., 2003a). Human arthroplasty–derived macrophages are capable of osteoclastic differentiation *in-vitro* in the presence of M-CSF and TNF-α (Ingham & Fisher, 2005; Sabokbar et al., 1997). Although recruitment of osteoclast precursor cells from the blood are more important as their source, the role of macrophages as osteoclast precusors in the periprosthetic space of osteolysis needs to be more clarified.

6. Involvement of lymphocyte in inflammatory osteolysis

The roles of lymphocytes in periprosthetic osteolysis remain to be delineated. Lymphocytes are generally absent or present in low numbers in the periprosthetic membranes. Mice

deficient in T cells, B cells, and natural killer cells develop osteolysis in response to wear particles as readily as wild type mice (Taki et al., 2005). However, the strongest evidence for the involvement of lymphocytes in aseptic loosening are a series of recent reports correlating a metal-specific lymphocyte response to poor implant performance and characterizing lymphocytic infiltration around metal-on-metal arthroplasties (Davies et al., 2005; Hallab et al., 2005). To promote osteoclastogenesis, activated T-cells positively regulate RANKL] and also negatively interferon-γ.

T-cell derived RANKL has been well known to play central role in inflammatory bone loss. In RA, the role of T cells has also been debated and unresolved as well as in periprosthetic osteolysis. An interesting new development has been the recognition of IL- 17 (Kolls & Linden, 2004). IL-17, produced predominantly by T-memory cells, acts synergistically with TNF-α to activate synovial fibroblast-like cells. T-helper cells producing IL-17 show a distinctive cytokine profile, which is consisted of IL-17, TNF-α and RANKL, but only low levels of IFN-γ and no IL-4 (Looney et al., 2006). Therfore, future work on the role of T cells in periprosthetic loosening should include evaluation of T cell signaling, related with the fact that inflammatory osteolysis do not produce much IFN-γ. It may be of special significance since IFN-γ has a potent inhibitory effect on osteoclast development and thus osteolysis (Looney et al., 2006; Takayanagi et al., 2000).

Cytokine	Effects on osteoblasts [OB]	Effects on osteoclasts [OC]
TNF	• Induces RANKL & M-CSF (Wei et al., 2005) • Inhibits OB differentiation & apoptosis (Gilbert et al., 2000; Jilka et al., 1998)	• Increases OC precursor numbers (Li et al., 2004; Yao et al., 2006) • Acts synergistically with RANKL (Lam et al., 2000)
IL-1	• IL-1α: Inhibits differentiation & matrix formation *in- vitro* (Tanabe et al., 2004) • IL-1β: Inhibits collagen synthesis *in-vitro* (Stashenko et al., 1987)	• Increases OC-genesis along with TNF-α (Wei et al., 2005) • Decreases apoptotic rate of OCs (Jimi et al., 1995)
IL-17	• Enhances TNF-α-stimulated IL-6 synthesis (Tokuda et al., 2004) • Increases RANKL/OPG in cells *in-vitro* (Kotake et al., 1999)	• Induces RANKL and RANK (Kotake et al., 1999; Lubberts et al., 2003) • Stimulate OC-genesis in RA (Kotake et al., 1999)

Table 2. Main effects of pro-inflammatory cytokines on osteoblast and osteoclast

7. Biological understandings of osteolytic response

The final cellular consequence of particle action is an excess of osteoclast activity, which results in progressive bone erosion. Osteoclasts are multinucleated cells derived from circulating osteoclast precursor cell of the monocyte/macrophage lineage, and represent the only cell type capable of bone resorption (Boyle et al., 2003). Osteoclast precursors are supplied from the periprosthetic space or recruited from the blood itself (Sabokbar et al., 1997). Wear debris probably increases osteoclast recruitment to periprosthetic tissues via the activation of chemokine [macrophage chemoattractant protein-1 ; IL-8] expression by macrophages and fibroblasts (Fritz et al., 2005; Nakashima et al., 1999a; Yaszay et al., 2001). In addition, macrophage lineage cells isolated from these tissues display a greatly increased propensity to differentiate into osteoclasts (Sabokbar et al., 1997; Sabokbar et al., 2003a). Osteoclasts can be differentiated by two critical cytokines, RANKL and M-CSF.

The molecular balance between RANK−RANKL and OPG has a key role in periprosthetic osteolysis. RANKL is the key cytokine regulator of osteoclast generation and activation. Interaction between RANK and RANKL constitutes a pivotal signaling pathway in the formation of osteoclasts. RANKL is expressed on the surface of activated T cells, marrow stoma cells, and osteoblasts as a 45-kDa transmembrane protein. It binds to RANK expressed on the surface of osteoclasts and also their precursors. This is necessary for the differentiation and maturation of osteoclasts in the presence of the survival factor M-CSF. By the binding of RANKL to RANK, the receptor recruits TNFR [TNF receptor]-associated cytoplasmic factor 6 [TRAF6]. This acts as a key adaptor for the assembling of signaling proteins, which directs osteoclast-specific gene expression and finally leads to their differentiation and activation.

OPG is a naturally occurring decoy receptor for RANKL secreted by stromal cells including osteoblasts as a soluble 110 kDa disulfide-linked homodimer. It down-regulates osteoclastogenesis by binding RANKL. Osteoclasts formation can be determined principally by the relative ratio of RANKL/OPG in the bone marrow microenvironment, and alterations in this ratio have been correlated with various bone disorders (Hofbauer & Schoppet, 2004).

Another important fact for regulation of osteoclastogenesis is that many pro-inflammatory and anti-inflammatory cytokines act directly to enhance or inhibit the RANKL/RANK axis (Abu-Amer et al., 2007). TNF-α also promotes osteoclastogenesis, particularly in the state of inflammatory osteolysis such as RA and periprosthetic osteolysis. Overexpression of TNF-α is sufficient to induce calvarial osteolysis even in the absence of added particles, emphasizing its pro-resorptive characteristics in mice (Schwarz et al., 2000). The molecular basis of increased RANKL in osteolysis is likely downstream of pro-inflammatory cytokines such as TNF-α and IL-1β, which are known to increase RANKL expression in several cell types (Purdue et al., 2007). RANKL and TNF-α seems to work in collaboration to induce osteoclast activation. Therefore, TNF-α and IL-1β, acting in concert with RANKL, can powerfully promote osteoclast recruitment, activation, and osteolysis (Romas et al., 2002).

During the past decade, the identification of several molecular pathways involved in bone loss raised hope for the development of therapeutic targets for periprosthetic osteolysis. TNF family members, especially RANKL, are prerequisites for osteoclast formation. The downstream signaling by wear particles, unsurprisingly, overlaps with that of TNF and RANKL. Notably, particle-induced pathways lead to the activation of kinases and transcription factors which are essential for osteoclastogenesis, such as activation of the tyrosine kinase c-src, mitogen-activated protein kinases [MAPK], and the NF-κB cascade (Abbas et al., 2003; Abu-Amer, 2005; Lam et al., 2002). Although activation of these pathways might be a secondary pathway, selective blockade of these downstream pathways reduces particle transmitted effects. The molecular targets described above need to be focused for selecting anti-resorptive therapeutic targets (Looney et al., 2006).

8. Impaired osteogenesis as an inflammatory reaction in periprosthetic osteolysis

The role of osteoblasts in periprosthetic osteolysis has received less attention than that of osteoclasts. Osteoblasts play important regulatory roles in bone remodeling. They produce and mineralize bone matrix, in addition to modulating differentiation and function of osteoclast by producing RANKL and OPG (Lorenzo et al., 2008). Osteoblasts are originated from MSCs and differentiated to matured cells. After maturation, osteoblasts diminish their

expression of RANKL and increase their expression of OPG, thereby creating a microenvironment that favors bone formation over bone loss (Atkins et al., 2003). Although osteoblasts have not been intensively investigated within the the field of periprosthetic osteolysis, more intensive research needs to be conducted to delineate the potentially critical role of osteoblasts based on their bone forming activity.

Most researches have limited their focus on *in-vitro* models for the study of direct interaction between osteoblast and particle (Dean et al., 1999; Gutwein & Webster, 2004; Lohmann et al., 2002; Pioletti et al., 2002; Yao et al., 1997). It has been postulated as a main mechanism of impaired osteogenesis that wear particles directly inhibit bone forming activity of osteoblast by altering typical osteogenic characters. For example, particles directly inhibit cell viability and proliferation, in addition to down-regulating the mRNA and protein level of bone formation markers. Particles less than 5 μm can also undergo phagocytosis by mature osteoblasts (Goodman et al., 2006), leading to potential adverse effects on cellular viability, proliferation and function. Along with particle size, composition and dosage can also effect these parameters (Lohmann et al., 2002). Moreover, it was reported that osteoblast challenged with particles can induce the expression of RANKL, OPG, IL-1, TNF-α, IL-6, IL-11, and TGF-β (Hofbauer et al., 2000).

MSCs and osteoprogenitors are also profoundly affected by wear particles (Drees et al., 2007; Goodman et al., 2006). Differentiation of osteoblasts from MSCs is also down-regulated by titanium particles (Wang et al., 2002). PMMA particles reduce osteoblast differentiation of bone marrow osteoprogenitor cells (Chiu et al., 2006). Titanium and zirconium oxide induce MSC apoptosis (Wang et al., 2003). Since MSCs and osteoprogenitors from the bone marrow are the precursors of osteoblasts, the reaction of these cells to wear particles is critical to both initial osseo-integration of implants and ongoing regeneration of the periprosthetic bed (Goodman et al., 2006). Future studies need to delineate the molecular mechanisms by which particles adversely affect bone cell lineage including MSCs and provide strategies to modulate these effects.

Recent research has uncovered the possibility that periprosthetic osteolysis likely involves multiple mechanisms including bone forming activity as well as bone resorption. It was reported that biologic effects on bone-forming cells - osteoblasts, osteoprogenitors, and adult MSCs - may also contribute to osteolysis (Chiu et al., 2009; Wang et al., 2002). These findings suggest that the following mechanisms of particle bioreactivity may contribute to osteolysis by means of exacerbated inflammation by reactive oxygen species [ROS] (Chiu et al., 2009) released from activated macrophages and osteoclasts, resulting to impaired periprosthetic bone formation with cytotoxic response and suppressed osteogenic differentiation of mesenchymal stem cells (Wang et al., 2004).

So far, most researches in terms of involvement of osteoblast in periprosthetic osteolyis have been limited to determine the direct suppressive effect of particle to osteoblast. However, the possibility that osteoblast can indirectly communicate with immune cells through many sectreted molecules such as TNF-α, IL-1, ROS requires further exploration (Ghali et al., 2010; Yamazaki et al., 2009; Zhou et al., 2006). Following phagocytosis of particles and the resultant pro-inflammatory reaction, the released cytokines from macrophages can be regarded as a potent inhibitor of osteoblast differentiation. Although insufficient attention has been paid to the involvement of osteoblasts, more extensive research should be conducted to delineate the potentially critical role of osteoblast in periprosthetic osteolysis. Modulation of bone forming activity in addition to existing anti-osteoclastic therapies, such as bisphosphonates and TNF-α blockade that inhibit bone destruction, represent a potential new therapeutic approach to this destructive disorder.

9. Molecular basis of inflammatory osteolysis

Inflammatory osteolysis is a major complication of conditions such as RA, periodontal disease, and orthopedic implant loosening. The persistence of these responses is often associated with skeletal pathology ranging from localized focal bone erosion and peri-articular osteolysis in the vicinity of inflamed area, to generalized osteopenia. This inflammatory osteolysis reflects increased osteoclast activity with enhanced osteoclast recruitment prompted by higher circulating levels of inflammatory mediators. Therefore, pathogenesis of inflammatory osteolysis is composed of distinct two primary components, inflammatory factors and regulation of osteoclasts. These are thought to operate through an ultimate common pathway of accelerated osteoclast recruitment and activation under the control of cytokines produced in the inflammatory environment.

As the only cell type capable of bone resorption, osteoclasts play a central to the pathogenesis of inflammatory osteolysis. Differentiation and activation of osteoclast are under the aegis of a variety of cytokines. Receptor activator of RANKL and M-CSF are the essential osteoclastogenic cytokines and are increased in inflammatory skeletal disease. The hyperplastic inflamed synovium also contains inflammatory cells such as lymphocytes, plasma cells, activated macrophages, and neutrophils. These cells can secrete a multitude of cytokines and growth factors including RANKL, TNF-α, IL-1, IL-6, PGE2, and IL-17 (Abu-Amer, 2009). This microenvironment is the evidence for recruitment and differentiation of osteoclasts that contribute to bone erosion.

The interaction of RANK and its ligand, RANKL is central to osteolytic responses on account of its critical role in osteoclast differentiation and survival. Interstingly, mouse models for the overexpression of OPG or administration of OPG-Fc are resistant to focal and systemic bone loss despite existence of the inflammatory response (Kong et al., 1999; Wong et al., 1999). These findings suggest that the osteoclast differentiation pathway, the RANKL/RANK signaling cascade, play a role as a target for other modulators for preventing bone resorption.

In addition, produced proinflammatory cytokines also play a vital role in the inflammatory osteolysis in RA, periprosthetic osteolysis, and periodontitis. Factors including TNF-α, IL-1, IL-17 and bacterial endotoxins also seem to impact osteoclastogenesis and bone resorption directly and indirectly (Abu-Amer, 2009). The dominant cytokine in the inflammatory osteolysis condition is TNF-α, primarily produced by activated T cells, macrophages and synoviocytes.

TNF-α is the most notable cytokine that can modulate both inflammatory and osteolytic process in the inflammatory osteolysis (Abu-Amer et al., 2008). Therefore, TNF-α can be regarded as the rate-limiting factor and it can be a target to eliminates both the inflammatory and osteoclastogenic components of these diseases (Wei & Siegal, 2008). However, in the most of researches, the role of TNF-α as the inflammatory mediator more than the osteolytic effector has been highlighted. This point is supported by studies in which inhibition of RANK signaling halted osteolysis whereas inflammation persisted. Nevertheless, TNF-α augments RANK/RANKL signaling tremendously leading to exacerbated osteoclastogenesis of RANKL-treated precursor cells. Therefore it appears that osteolytic activity of TNF-α requires RANKL/ RANK system in inflammatory disease (Abu-Amer, 2009). IL-1 also plays an essential role in the pathophysiology of inflammatory bone loss. Other prominent pro-inflammatory and pro-osteolytic factors include IL-17 and IL-6.

Regulation of pro-inflammatory cytokines appears to be a major function of IL-17. IL-17 directly upregulates IL-1 and TNF-α-induced inflammatory responses (Abu-Amer, 2009). IL-17, secreted by a distinct lymphocyte subset cells, plays an important role in

inflammation and bone erosion in a mouse model of CIA. Treatment with anti-IL-17, even after the onset of disease, markedly attenuates damage and inflammation of myocardium (Fan et al., 2011). In addition, IL-17 producing T cells are present in the synovium of RA patients (Page et al., 2004). Moreover IL-17 induces expression of RANKL by osteoblasts and synovial fibroblasts, leading to decreasing expression of OPG by stromal cells. Overall, a cascade from inflammatory cells lead to secretion of IL-17 which in turn up-regulates expression of RANKL, TNF-α and IL-1 and down-regulats expression of OPG, providing an intricate system supporting inflammation and subsequent osteolysis (Abu-Amer, 2009). Due to interdependence of TNF-α or IL-1, blockade of either TNF-α or IL-1 does not completely arrest the periarticular bone loss of inflammatory arthritis, however, inhibition of the two cytokines in combination is substantially more effective (Wei & Siegal, 2008).

The overall mechanism described above also can be applied to periprosthetic osteolysis from wear debris. Studies using animal model involving TNF-α blockade has been shown to significantly reduce wear debris-induced osteolysis (Childs et al., 2001a, b), but residual osteolysis still persists. In contrst, disruption of RANKL signaling via genetic ablation or high dose RANK-Fc treatment completely eliminates osteoclasts and bone resorption in this model (Childs et al., 2002). Similar effects were also achieved via OPG gene therapy (Goater et al., 2002; Ulrich-Vinther et al., 2002; Yang et al., 2002a).

It can be considered that the biological responsive pattern in periprosthetic osteolysis is similar to other modes of inflammatory osteolysis in that it is composed of two primary components, inflammatory factors and regulation of osteoclasts. This is thought to operate through common signaling pathways of cytokines such as TNF-α, IL-1 and RANKL to accelerate osteoclast recruitment and activation under the control of cytokines produced in the inflammatory environment against wear debris.

Understanding the mechanisms by which osteoclasts resorb bone, and the cytokines that regulate their differentiation and activity, provides mechanism-based candidate therapeutic targets to prevent inflammatory bone loss induced by wear debris from orthopedic implants. The success of anti-TNF-α and IL-1 therapy highlights the central role that these specific cytokines play in this disease except periprosthetic bone loss by wear debris. In addition, the interdependence of TNF-α, RANKL and IL-17 in the generation of osteoclasts also allows to explain the observation that combined blockade is more effective in preventing pathological bone loss in the inflammatory conditions including periprosthetic osteolysis (Buckland, 2011).

10. Conclusions

We hereby describe the biological mechanisms that are responsible for inflammatory bone loss in periprosthetic osteolysis, highlighting potential targets for further therapeutic approaches to prevent and minimize this devastating complication. As it is generally accepted that the inflammatory interaction between wear debris and activated macrophages is defined as a key event in periprosthetic osteolysis, much effort has been focused on this process and its role in osteoclast activation.

However, to date, despite extensive and complex research concerning periprosthetic osteolysis, there is no effective medical therapy to prevent or inhibit periprosthetic osteolysis. Therefore, an appreciation of the complex cellular and molecular signal network leading to cellular and inflammatory responses will form a foundation, on which several therapeutic interventions can be developed to overcome inflammatory periprosthetic bone loss. For the future direction, it seems to be reasonable that additional attention should be

equally paid to potentiate osteogenesis to overcome bone loss in the periprosthetic osteolysis.

11. Acknowledgment

This work was supported by Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education, Science and Technology (2011-000-6208 & 2011-001-4792).

12. References

Abbas, S.; Clohisy, J.C. & Abu-Amer, Y. (2003). Mitogen-activated protein (MAP) kinases mediate PMMA-induction of osteoclasts. *Journal of Orthopaedic Research*, Vol.21, No.6, (Nov), pp. 1041-1048

Abu-Amer, Y. (2005). Advances in osteoclast differentiation and function. *Curr Drug Targets Immune Endocr Metabol Disord*, Vol.5, No.3, (Sep), pp. 347-355, ISSN 1568-0088

Abu-Amer, Y. (2009). Inflammation, cancer, and bone loss. *Curr Opin Pharmacol*, Vol.9, No.4, (Aug), pp. 427-433, ISSN 1471-4973 (Electronic)

Abu-Amer, Y.; Darwech, I. & Clohisy, J.C. (2007). Aseptic loosening of total joint replacements: mechanisms underlying osteolysis and potential therapies. *Arthritis Res Ther*, Vol.9 Suppl 1, pp. S6, ISSN 1478-6362 (Electronic)

Abu-Amer, Y.; Darwech, I. & Otero, J. (2008). Role of the NF-kappaB axis in immune modulation of osteoclasts and bone loss. *Autoimmunity*, Vol.41, No.3, (Apr), pp. 204-211, ISSN 1607-842X (Electronic)

Archibeck, M.J.; Jacobs, J.J.; Roebuck, K.A. & Glant, T.T. (2001). The basic science of periprosthetic osteolysis. *Instructional Course Lectures*, Vol.50, pp. 185-195

Arora, A.; Song, Y.; Chun, L.; Huie, P.; Trindade, M.; Smith, R.L. & Goodman, S. (2003). The role of the TH1 and TH2 immune responses in loosening and osteolysis of cemented total hip replacements. *J Biomed Mater Res A*, Vol.64, No.4, (Mar 15), pp. 693-697

Atkins, G.J.; Kostakis, P.; Pan, B.; Farrugia, A.; Gronthos, S.; Evdokiou, A.; Harrison, K.; Findlay, D.M. & Zannettino, A.C. (2003). RANKL expression is related to the differentiation state of human osteoblasts. *Journal of Bone and Mineral Researc*h, Vol.18, No.6, (Jun), pp. 1088-1098, ISSN 0884-0431

Blaine, T.A.; Rosier, R.N.; Puzas, J.E.; Looney, R.J.; Reynolds, P.R.; Reynolds, S.D. & O'Keefe, R.J. (1996). Increased levels of tumor necrosis factor-alpha and interleukin-6 protein and messenger RNA in human peripheral blood monocytes due to titanium particles. *Journal of Bone and Joint Surgery*, Vol.78, No.8, (Aug), pp. 1181-1192

Boyle, W.J.; Simonet, W.S. & Lacey, D.L. (2003). Osteoclast differentiation and activation. *Nature*, Vol.423, No.6937, (May 15), pp. 337-342

Buckland, J. (2011). Experimental arthritis: Therapeutic promise of dual blockade of IL-17 and TNF in inflammatory arthritis. *Nat Rev Rheumatol*, Vol.7, No.6, (June), pp. 311, ISSN 1759-4804 (Electronic)

Charnley, J. (1966). Using Teflon in arthroplasty of the hip-joint. *Journal of Bone and Joint Surgery*, Vol.48, No.4, (Jun), pp. 819, ISSN 0021-9355

Chiba, J.; Rubash, H.E.; Kim, K.J. & Iwaki, Y. (1994). The characterization of cytokines in the interface tissue obtained from failed cementless total hip arthroplasty with and without femoral osteolysis. *Clin Orthop Relat Res*, No.300, (Mar), pp. 304-312

Childs, L.M.; Goater, J.J.; O'Keefe, R.J. & Schwarz, E.M. (2001a). Effect of anti-tumor necrosis factor-alpha gene therapy on wear debris-induced osteolysis. *J Bone Joint Surg Am*, Vol.83-A, No.12, (Dec), pp. 1789-1797, ISSN 0021-9355

Childs, L.M.; Goater, J.J.; O'Keefe, R.J. & Schwarz, E.M. (2001b). Efficacy of etanercept for wear debris-induced osteolysis. *Journal of Bone and Mineral Research*, Vol.16, No.2, (Feb), pp. 338-347

Childs, L.M.; Paschalis, E.P.; Xing, L.; Dougall, W.C.; Anderson, D.; Boskey, A.L.; Puzas, J.E.; Rosier, R.N.; O'Keefe, R.J.; Boyce, B.F. & Schwarz, E.M. (2002). In vivo RANK signaling blockade using the receptor activator of NF-kappaB:Fc effectively prevents and ameliorates wear debris-induced osteolysis via osteoclast depletion without inhibiting osteogenesis. *Journal of Bone and Mineral Research*, Vol.17, No.2, (Feb), pp. 192-199

Chiu, R.; Ma, T.; Smith, R.L. & Goodman, S.B. (2006). Polymethylmethacrylate particles inhibit osteoblastic differentiation of bone marrow osteoprogenitor cells. *J Biomed Mater Res A*, Vol.77, No.4, (Jun 15), pp. 850-856, ISSN 1549-3296

Chiu, R.; Ma, T.; Smith, R.L. & Goodman, S.B. (2009). Ultrahigh molecular weight polyethylene wear debris inhibits osteoprogenitor proliferation and differentiation in vitro. *J Biomed Mater Res A*, Vol.89, No.1, (Apr), pp. 242-247, ISSN 1552-4965 (Electronic)

Clohisy, J.C.; Calvert, G.; Tull, F.; McDonald, D. & Maloney, W.J. (2004a). Reasons for revision hip surgery: a retrospective review. *Clin Orthop Relat Res,* No.429, (Dec), pp. 188-192, ISSN 0009-921X

Clohisy, J.C.; Hirayama, T.; Frazier, E.; Han, S.K. & Abu-Amer, Y. (2004b). NF-kB signaling blockade abolishes implant particle-induced osteoclastogenesis. *Journal of Orthopaedic Research*, Vol.22, No.1, (Jan), pp. 13-20

Davies, A.P.; Willert, H.G.; Campbell, P.A.; Learmonth, I.D. & Case, C.P. (2005). An unusual lymphocytic perivascular infiltration in tissues around contemporary metal-on-metal joint replacements. *Journal of Bone and Joint Surgery*, Vol.87, No.1, (Jan), pp. 18-27

Dean, D.D.; Schwartz, Z.; Liu, Y.; Blanchard, C.R.; Agrawal, C.M.; Mabrey, J.D.; Sylvia, V.L.; Lohmann, C.H. & Boyan, B.D. (1999). The effect of ultra-high molecular weight polyethylene wear debris on MG63 osteosarcoma cells in vitro. *Journal of Bone and Joint Surgery*, Vol.81, No.4, (Apr), pp. 452-461

Dorr, L.D.; Bloebaum, R.; Emmanual, J. & Meldrum, R. (1990). Histologic, biochemical, and ion analysis of tissue and fluids retrieved during total hip arthroplasty. *Clin Orthop Relat Res*, No.261, (Dec), pp. 82-95, ISSN 0009-921X

Drees, P.; Eckardt, A.; Gay, R.E.; Gay, S. & Huber, L.C. (2007). Mechanisms of disease: Molecular insights into aseptic loosening of orthopedic implants. *Nat Clin Pract Rheumatol*, Vol.3, No.3, (Mar), pp. 165-171, ISSN 1745-8382

Dumbleton, J.H.; Manley, M.T. & Edidin, A.A. (2002). A literature review of the association between wear rate and osteolysis in total hip arthroplasty. *J Arthroplasty*, Vol.17, No.5, (Aug), pp. 649-661, ISSN 0883-5403

Engler, J.; Paul, H.; Gamardo, J. & Rodriguez, M.A. (2001). Acute synovitis, fever and rash possibly caused by metallic debris from a loosened knee prosthesis in a patient with rheumatoid arthritis. *J Clin Rheumatol,* Vol.7, No.4, (Aug), pp. 257-260, ISSN 1076-1608 (Print)

Fan, Y.; Weifeng, W.; Yuluan, Y.; Qing, K.; Yu, P. & Yanlan, H. (2011).Treatment with a neutralizing anti-murine interleukin-17 antibody after the onset of coxsackievirus

b3-induced viral myocarditis reduces myocardium inflammation. *Virol J*, Vol.8, pp. 17, ISSN 1743-422X (Electronic)

Fritz, E.A.; Jacobs, J.J.; Glant, T.T. & Roebuck, K.A. (2005). Chemokine IL-8 induction by particulate wear debris in osteoblasts is mediated by NF-kappaB. *Journal of Orthopaedic Research*, Vol.23, No.6, (Nov), pp. 1249-1257

Gallo, J.; Kaminek, P.; Ticha, V.; Rihakova, P. & Ditmar, R. (2002). Particle disease. A comprehensive theory of periprosthetic osteolysis: a review. *Biomed Pap Med Fac Univ Palacky Olomouc Czech Repub*, Vol.146, No.2, (Dec), pp. 21-28, ISSN 1213-8118

Gelb, H.; Schumacher, H.R.; Cuckler, J.; Ducheyne, P. & Baker, D.G. (1994). In vivo inflammatory response to polymethylmethacrylate particulate debris: effect of size, morphology, and surface area. *Journal of Orthopaedic Research*, Vol.12, No.1, (Jan), pp. 83-92

Ghali, O.; Chauveau, C.; Hardouin, P.; Broux, O. & Devedjian, J.C. (2010). TNF-alpha's effects on proliferation and apoptosis in human mesenchymal stem cells depend on RUNX2 expression. *Journal of Bone and Mineral Research*, Vol.25, No.7, (Jul), pp. 1616-1626, ISSN 1523-4681 (Electronic)

Gilbert, L.; He, X.; Farmer, P.; Boden, S.; Kozlowski, M.; Rubin, J. & Nanes, M.S. (2000). Inhibition of osteoblast differentiation by tumor necrosis factor-alpha. *Endocrinology*, Vol.141, No.11, (Nov), pp. 3956-3964, ISSN 0013-7227

Gilbert, L.; He, X.; Farmer, P.; Rubin, J.; Drissi, H.; van Wijnen, A.J.; Lian, J.B.; Stein, G.S. & Nanes, M.S. (2002). Expression of the osteoblast differentiation factor RUNX2 (Cbfa1/AML3/Pebp2alpha A) is inhibited by tumor necrosis factor-alpha. *Journal of Biological Chemistry*, Vol.277, No.4, (Jan 25), pp. 2695-2701, ISSN 0021-9258

Glant, T.T.; Jacobs, J.J.; Molnar, G.; Shanbhag, A.S.; Valyon, M. & Galante, J.O. (1993). Bone resorption activity of particulate-stimulated macrophages. *Journal of Bone and Mineral Research*, Vol.8, No.9, (Sep), pp. 1071-1079, ISSN 0884-0431

Goater, J.J.; O'Keefe, R.J.; Rosier, R.N.; Puzas, J.E. & Schwarz, E.M. (2002). Efficacy of ex vivo OPG gene therapy in preventing wear debris induced osteolysis. *Journal of Orthopaedic Research*, Vol.20, No.2, (Mar), pp. 169-173

Goldring, S.R.; Clark, C.R. & Wright, T.M. (1993). The problem in total joint arthroplasty: aseptic loosening. *Journal of Bone and Joint Surgery*, Vol.75, No.6, (Jun), pp. 799-801, ISSN 0021-9355

Goldring, S.R.; Schiller, A.L.; Roelke, M.; Rourke, C.M.; O'Neil, D.A. & Harris, W.H. (1983). The synovial-like membrane at the bone-cement interface in loose total hip replacements and its proposed role in bone lysis. *Journal of Bone and Joint Surgery*, Vol.65, No.5, (Jun), pp. 575-584, ISSN 0021-9355

Gonzalez, O.; Smith, R.L. & Goodman, S.B. (1996). Effect of size, concentration, surface area, and volume of polymethylmethacrylate particles on human macrophages in vitro. *Journal of Biomedical Materials Research*, Vol.30, No.4, (Apr), pp. 463-473, ISSN 0021-9304

Goodman, S.B.; Ma, T.; Chiu, R.; Ramachandran, R. & Smith, R.L. (2006). Effects of orthopaedic wear particles on osteoprogenitor cells. *Biomaterials*, Vol.27, No.36, (Dec), pp. 6096-6101, ISSN 0142-9612

Granchi, D.; Amato, I.; Battistelli, L.; Ciapetti, G.; Pagani, S.; Avnet, S.; Baldini, N. & Giunti, A. (2005). Molecular basis of osteoclastogenesis induced by osteoblasts exposed to wear particles. *Biomaterials*, Vol.26, No.15, (May), pp. 2371-2379

Green, T.R.; Fisher, J.; Stone, M.; Wroblewski, B.M. & Ingham, E. (1998). Polyethylene particles of a 'critical size' are necessary for the induction of cytokines by macrophages in vitro. *Biomaterials*, Vol.19, No.24, (Dec), pp. 2297-2302

Gutwein, L.G. & Webster, T.J. (2004). Increased viable osteoblast density in the presence of nanophase compared to conventional alumina and titania particles. *Biomaterials*, Vol.25, No.18, (Aug), pp. 4175-4183, ISSN 0142-9612

Hallab, N.J.; Anderson, S.; Stafford, T.; Glant, T. & Jacobs, J.J. (2005). Lymphocyte responses in patients with total hip arthroplasty. *Journal of Orthopaedic Research*, Vol.23, No.2, (Mar), pp. 384-391

Harris, W.H. (1995). The problem is osteolysis. *Clin Orthop Relat Res*, No.311, (Feb), pp. 46-53

Haynes, D.R.; Boyle, S.J.; Rogers, S.D.; Howie, D.W. & Vernon-Roberts, B. (1998). Variation in cytokines induced by particles from different prosthetic materials. *Clin Orthop Relat Res*, No.352, (Jul), pp. 223-230

Haynes, D.R.; Crotti, T.N. & Zreiqat, H. (2004). Regulation of osteoclast activity in peri-implant tissues. *Biomaterials*, Vol.25, No.20, (Sep), pp. 4877-4885

Hirakawa, K.; Bauer, T.W.; Stulberg, B.N. & Wilde, A.H. (1996). Comparison and quantitation of wear debris of failed total hip and total knee arthroplasty. *Journal of Biomedical Materials Research*, Vol.31, No.2, (Jun), pp. 257-263

Hofbauer, L.C.; Khosla, S.; Dunstan, C.R.; Lacey, D.L.; Boyle, W.J. & Riggs, B.L. (2000). The roles of osteoprotegerin and osteoprotegerin ligand in the paracrine regulation of bone resorption. *Journal of Bone and Mineral Research*, Vol.15, No.1, (Jan), pp. 2-12, ISSN 0884-0431

Hofbauer, L.C. & Schoppet, M. (2004). Clinical implications of the osteoprotegerin/ RANKL/RANK system for bone and vascular diseases. *JAMA*, Vol.292, No.4, (Jul 28), pp. 490-495

Ingham, E. & Fisher, J. (2005). The role of macrophages in osteolysis of total joint replacement. *Biomaterials*, Vol.26, No.11, (Apr), pp. 1271-1286

Jacobs, J.J.; Roebuck, K.A.; Archibeck, M.; Hallab, N.J. & Glant, T.T. (2001). Osteolysis: basic science. *Clin Orthop Relat Res*, No.393, (Dec), pp. 71-77

Jilka, R.L.; Weinstein, R.S.; Bellido, T.; Parfitt, A.M. & Manolagas, S.C. (1998). Osteoblast programmed cell death (apoptosis): modulation by growth factors and cytokines. *Journal of Bone and Mineral Research*, Vol.13, No.5, (May), pp. 793-802, ISSN 0884-0431

Jimi, E.; Shuto, T. & Koga, T. (1995). Macrophage colony-stimulating factor and interleukin-1 alpha maintain the survival of osteoclast-like cells. *Endocrinology*, Vol.136, No.2, (Feb), pp. 808-811, ISSN 0013-7227

Jiranek, W.A.; Machado, M.; Jasty, M.; Jevsevar, D.; Wolfe, H.J.; Goldring, S.R.; Goldberg, M.J. & Harris, W.H. (1993). Production of cytokines around loosened cemented acetabular components. Analysis with immunohistochemical techniques and in situ hybridization. *Journal of Bone and Joint Surgery*, Vol.75, No.6, (Jun), pp. 863-879

Karmakar, S.; Kay, J. & Gravallese, E.M. (2010). Bone damage in rheumatoid arthritis: mechanistic insights and approaches to prevention. *Rheumatic Diseases Clinics of North America*, Vol.36, No.2, (May), pp. 385-404, ISSN 1558-3163 (Electronic)

Keener, J.D.; Callaghan, J.J.; Goetz, D.D.; Pederson, D.R.; Sullivan, P.M. & Johnston, R.C. (2003). Twenty-five-year results after Charnley total hip arthroplasty in patients less than fifty years old: a concise follow-up of a previous report. *Journal of Bone and Joint Surgery*, Vol.85-A, No.6, (Jun), pp. 1066-1072

Khosla, S. (2001). Minireview: the OPG/RANKL/RANK system. *Endocrinology*, Vol.142, No.12, (Dec), pp. 5050-5055, ISSN 0013-7227

Kido, A.; Pap, G.; Nagler, D.K.; Ziomek, E.; Menard, R.; Neumann, H.W. & Roessner, A. (2004). Protease expression in interface tissues around loose arthroplasties. *Clin Orthop Relat Res*, No.425, (Aug), pp. 230-236

Kolls, J.K. & Linden, A. (2004). Interleukin-17 family members and inflammation. *Immunity*, Vol.21, No.4, (Oct), pp. 467-476, ISSN 1074-7613

Kong, Y.Y.; Yoshida, H.; Sarosi, I.; Tan, H.L.; Timms, E.; Capparelli, C.; Morony, S.; Oliveira-dos-Santos, A.J.; Van, G.; Itie, A.; Khoo, W.; Wakeham, A.; Dunstan, C.R.; Lacey, D.L.; Mak, T.W.; Boyle, W.J. & Penninger, J.M. (1999). OPGL is a key regulator of osteoclastogenesis, lymphocyte development and lymph-node organogenesis. *Nature*, Vol.397, No.6717, (Jan 28), pp. 315-323

Kotake, S.; Udagawa, N.; Takahashi, N.; Matsuzaki, K.; Itoh, K.; Ishiyama, S.; Saito, S.; Inoue, K.; Kamatani, N.; Gillespie, M.T.; Martin, T.J. & Suda, T. (1999). IL-17 in synovial fluids from patients with rheumatoid arthritis is a potent stimulator of osteoclastogenesis. *Journal of Clinical Investigation*, Vol.103, No.9, (May), pp. 1345-1352, ISSN 0021-9738

Lam, J.; Abu-Amer, Y.; Nelson, C.A.; Fremont, D.H.; Ross, F.P. & Teitelbaum, S.L. (2002). Tumour necrosis factor superfamily cytokines and the pathogenesis of inflammatory osteolysis. *Ann Rheum Dis*, Vol.61 Suppl 2, (Nov), pp. ii82-83, ISSN 0003-4967

Lam, J.; Takeshita, S.; Barker, J.E.; Kanagawa, O.; Ross, F.P. & Teitelbaum, S.L. (2000). TNF-alpha induces osteoclastogenesis by direct stimulation of macrophages exposed to permissive levels of RANK ligand. *Journal of Clinical Investigation*, Vol.106, No.12, (Dec), pp. 1481-1488, ISSN 0021-9738

Lee, S.S; Chang JD.: Purdue, E.P; Nestor, B.J.; Sculco, T.P. & Salvati, E.A. (2007). Roles of Cellular and Molecular Targets of Wear Debris In Periprosthetic Osteolysis *Proceedings of 12th Biolox®Symposiumon Bioceramic and alternative Bearing in joint arthroplasty*, pp. 19-30, ISBN 978-3-7985-1782-0, Seoul, Korea, Sep 7-8,2007.

Lee, S.S.; Woo, C.H.; Chang, J.D. & Kim, J.H. (2003). Roles of Rac and cytosolic phospholipase A2 in the intracellular signalling in response to titanium particles. *Cellular Signalling*, Vol.15, No.3, (Mar), pp. 339-345, ISSN 0898-6568

Li, P.; Schwarz, E.M.; O'Keefe, R.J.; Ma, L.; Looney, R.J.; Ritchlin, C.T.; Boyce, B.F. & Xing, L. (2004). Systemic tumor necrosis factor alpha mediates an increase in peripheral CD11bhigh osteoclast precursors in tumor necrosis factor alpha-transgenic mice. *Arthritis and Rheumatism*, Vol.50, No.1, (Jan), pp. 265-276, ISSN 0004-3591

Lohmann, C.H.; Dean, D.D.; Koster, G.; Casasola, D.; Buchhorn, G.H.; Fink, U.; Schwartz, Z. & Boyan, B.D. (2002). Ceramic and PMMA particles differentially affect osteoblast phenotype. *Biomaterials*, Vol.23, No.8, (Apr), pp. 1855-1863

Lohmann, C.H.; Schwartz, Z.; Koster, G.; Jahn, U.; Buchhorn, G.H.; MacDougall, M.J.; Casasola, D.; Liu, Y.; Sylvia, V.L.; Dean, D.D. & Boyan, B.D. (2000). Phagocytosis of wear debris by osteoblasts affects differentiation and local factor production in a manner dependent on particle composition. *Biomaterials*, Vol.21, No.6, (Mar), pp. 551-561

Looney, R.J.; Schwarz, E.M.; Boyd, A. & O'Keefe, R.J. (2006). Periprosthetic osteolysis: an immunologist's update. *Current Opinion in Rheumatology*, Vol.18, No.1, (Jan), pp. 80-87

Lorenzo, J.; Horowitz, M. & Choi, Y. (2008). Osteoimmunology: interactions of the bone and immune system. *Endocrine Reviews*, Vol.29, No.4, (Jun), pp. 403-440, ISSN 0163-769X

Lubberts, E.; van den Bersselaar, L.; Oppers-Walgreen, B.; Schwarzenberger, P.; Coenen-de Roo, C.J.; Kolls, J.K.; Joosten, L.A. & van den Berg, W.B. (2003). IL-17 promotes bone erosion in murine collagen-induced arthritis through loss of the receptor activator of NF-kappa B ligand/osteoprotegerin balance. *Journal of Immunology*, Vol.170, No.5, (Mar 1), pp. 2655-2662, ISSN 0022-1767

Margevicius, K.J.; Bauer, T.W.; McMahon, J.T.; Brown, S.A. & Merritt, K. (1994). Isolation and characterization of debris in membranes around total joint prostheses. *Journal of Bone and Joint Surgery*, Vol.76, No.11, (Nov), pp. 1664-1675

Nakashima, Y.; Sun, D.H.; Trindade, M.C.; Chun, L.E.; Song, Y.; Goodman, S.B.; Schurman, D.J.; Maloney, W.J. & Smith, R.L. (1999a). Induction of macrophage C-C chemokine expression by titanium alloy and bone cement particles. *Journal of Bone and Joint Surgery. British Volume*, Vol.81, No.1, (Jan), pp. 155-162

Nakashima, Y.; Sun, D.H.; Trindade, M.C.; Maloney, W.J.; Goodman, S.B.; Schurman, D.J. & Smith, R.L. (1999b). Signaling pathways for tumor necrosis factor-alpha and interleukin-6 expression in human macrophages exposed to titanium-alloy particulate debris in vitro. *Journal of Bone and Joint Surgery*, Vol.81, No.5, (May), pp. 603-615

Neale, S.D. & Athanasou, N.A. (1999). Cytokine receptor profile of arthroplasty macrophages, foreign body giant cells and mature osteoclasts. *Acta Orthopaedica Scandinavica*, Vol.70, No.5, (Oct), pp. 452-458, ISSN 0001-6470

Niki, Y.; Matsumoto, H.; Otani, T.; Tomatsu, T. & Toyama, Y. (2007). Five types of inflammatory arthritis following total knee arthroplasty. *J Biomed Mater Res A*, Vol.81, No.4, (Jun 15), pp. 1005-1010, ISSN 1549-3296 (Print)

O'Keefe, R.J.; Rosier, R.N.; Teot, L.A.; Stewart, J.M. & Hicks, D.G. (1998). Cytokine and matrix metalloproteinase expression in pigmented villonodular synovitis may mediate bone and cartilage destruction. *Iowa Orthopaedic Journal*, Vol.18, pp. 26-34, ISSN 1541-5457

Page, G.; Sattler, A.; Kersten, S.; Thiel, A.; Radbruch, A. & Miossec, P. (2004). Plasma cell-like morphology of Th1-cytokine-producing cells associated with the loss of CD3 expression. *American Journal of Pathology*, Vol.164, No.2, (Feb), pp. 409-417, ISSN 0002-9440

Perry, M.J.; Mortuza, F.Y.; Ponsford, F.M.; Elson, C.J. & Atkins, R.M. (1995). Analysis of cell types and mediator production from tissues around loosening joint implants. *Br J Rheumatol*, Vol.34, No.12, (Dec), pp. 1127-1134, ISSN 0263-7103

Pioletti, D.P.; Leoni, L.; Genini, D.; Takei, H.; Du, P. & Corbeil, J. (2002). Gene expression analysis of osteoblastic cells contacted by orthopedic implant particles. *Journal of Biomedical Materials Research*, Vol.61, No.3, (Sep 5), pp. 408-420

Purdue, P.E. (2008). Alternative macrophage activation in periprosthetic osteolysis. *Autoimmunity*, Vol.41, No.3, (Apr), pp. 212-217, ISSN 1607-842X (Electronic)

Purdue, P.E.; Koulouvaris, P.; Potter, H.G.; Nestor, B.J. & Sculco, T.P. (2007). The cellular and molecular biology of periprosthetic osteolysis. *Clin Orthop Relat Res*, Vol.454, (Jan), pp. 251-261

Quinn, J.; Joyner, C.; Triffitt, J.T. & Athanasou, N.A. (1992). Polymethylmethacrylate-induced inflammatory macrophages resorb bone. *J Bone Joint Surg Br*, Vol.74, No.5, (Sep), pp. 652-658, ISSN 0301-620X

Romas, E.; Gillespie, M.T. & Martin, T.J. (2002). Involvement of receptor activator of NFkappaB ligand and tumor necrosis factor-alpha in bone destruction in rheumatoid arthritis. *Bone*, Vol.30, No.2, (Feb), pp. 340-346, ISSN 8756-3282

Sabokbar, A.; Fujikawa, Y.; Neale, S.; Murray, D.W. & Athanasou, N.A. (1997). Human arthroplasty derived macrophages differentiate into osteoclastic bone resorbing cells. *Annals of the Rheumatic Diseases*, Vol.56, No.7, (Jul), pp. 414-420

Sabokbar, A.; Kudo, O. & Athanasou, N.A. (2003a). Two distinct cellular mechanisms of osteoclast formation and bone resorption in periprosthetic osteolysis. *Journal of Orthopaedic Research*, Vol.21, No.1, (Jan), pp. 73-80, ISSN 0736-0266

Sabokbar, A.; Pandey, R. & Athanasou, N.A. (2003b). The effect of particle size and electrical charge on macrophage-osteoclast differentiation and bone resorption. *J Mater Sci Mater Med*, Vol.14, No.9, (Sep), pp. 731-738, ISSN 0957-4530

Sakai, H.; Jingushi, S.; Shuto, T.; Urabe, K.; Ikenoue, T.; Okazaki, K.; Kukita, T.; Kukita, A. & Iwamoto, Y. (2002). Fibroblasts from the inner granulation tissue of the pseudocapsule in hips at revision arthroplasty induce osteoclast differentiation, as do stromal cells. *Annals of the Rheumatic Diseases*, Vol.61, No.2, (Feb), pp. 103-109, ISSN 0003-4967

Schwarz, E.M.; Lu, A.P.; Goater, J.J.; Benz, E.B.; Kollias, G.; Rosier, R.N.; Puzas, J.E. & O'Keefe, R.J. (2000). Tumor necrosis factor-alpha/nuclear transcription factor-kappaB signaling in periprosthetic osteolysis. *Journal of Orthopaedic Research*, Vol.18, No.3, (May), pp. 472-480

Sethi, R.K.; Neavyn, M.J.; Rubash, H.E. & Shanbhag, A.S. (2003). Macrophage response to cross-linked and conventional UHMWPE. *Biomaterials*, Vol.24, No.15, (Jul), pp. 2561-2573

Shanbhag, A.S.; Jacobs, J.J.; Black, J.; Galante, J.O. & Glant, T.T. (1994). Macrophage/particle interactions: effect of size, composition and surface area. *Journal of Biomedical Materials Research*, Vol.28, No.1, (Jan), pp. 81-90

Shanbhag, A.S.; Jacobs, J.J.; Black, J.; Galante, J.O. & Glant, T.T. (1995). Cellular mediators secreted by interfacial membranes obtained at revision total hip arthroplasty. *Journal of Arthroplasty*, Vol.10, No.4, (Aug), pp. 498-506

Stashenko, P.; Dewhirst, F.E.; Rooney, M.L.; Desjardins, L.A. & Heeley, J.D. (1987). Interleukin-1 beta is a potent inhibitor of bone formation in vitro. *Journal of Bone and Mineral Research*, Vol.2, No.6, (Dec), pp. 559-565, ISSN 0884-0431

Takagi, M.; Santavirta, S.; Ida, H.; Ishii, M.; Mandelin, J. & Konttinen, Y.T. (1998). Matrix metalloproteinases and tissue inhibitors of metalloproteinases in loose artificial hip joints. *Clin Orthop Relat Res*, No.352, (Jul), pp. 35-45, ISSN 0009-921X

Takagi, M.; Tamaki, Y.; Hasegawa, H.; Takakubo, Y.; Konttinen, L.; Tiainen, V.M.; Lappalainen, R.; Konttinen, Y.T. & Salo, J. (2007). Toll-like receptors in the interface membrane around loosening total hip replacement implants. *J Biomed Mater Res A*, Vol.81, No.4, (Jun 15), pp. 1017-1026, ISSN 1549-3296

Takayanagi, H.; Ogasawara, K.; Hida, S.; Chiba, T.; Murata, S.; Sato, K.; Takaoka, A.; Yokochi, T.; Oda, H.; Tanaka, K.; Nakamura, K. & Taniguchi, T. (2000). T-cell-mediated regulation of osteoclastogenesis by signalling cross-talk between RANKL and IFN-gamma. *Nature*, Vol.408, No.6812, (Nov 30), pp. 600-605

Taki, N.; Tatro, J.M.; Nalepka, J.L.; Togawa, D.; Goldberg, V.M.; Rimnac, C.M. & Greenfield, E.M. (2005). Polyethylene and titanium particles induce osteolysis by similar,

lymphocyte-independent, mechanisms. *Journal of Orthopaedic Research*, Vol.23, No.2, (Mar), pp. 376-383

Talmo, C.T.; Shanbhag, A.S. & Rubash, H.E. (2006). Nonsurgical management of osteolysis: challenges and opportunities. *Clin Orthop Relat Res*, Vol.453, (Dec), pp. 254-264, ISSN 0009-921X

Tanabe, N.; Ito-Kato, E.; Suzuki, N.; Nakayama, A.; Ogiso, B.; Maeno, M. & Ito, K. (2004). IL-1alpha affects mineralized nodule formation by rat osteoblasts. *Life Sciences*, Vol.75, No.19, (Sep 24), pp. 2317-2327, ISSN 0024-3205

Tokuda, H.; Kanno, Y.; Ishisaki, A.; Takenaka, M.; Harada, A. & Kozawa, O. (2004). Interleukin (IL)-17 enhances tumor necrosis factor-alpha-stimulated IL-6 synthesis via p38 mitogen-activated protein kinase in osteoblasts. *Journal of Cellular Biochemistry*, Vol.91, No.5, (Apr 1), pp. 1053-1061, ISSN 0730-2312

Ulrich-Vinther, M.; Carmody, E.E.; Goater, J.J.; K, S.b.; O'Keefe, R.J. & Schwarz, E.M. (2002). Recombinant adeno-associated virus-mediated osteoprotegerin gene therapy inhibits wear debris-induced osteolysis. *Journal of Bone and Joint Surgery*, Vol.84-A, No.8, (Aug), pp. 1405-1412

Vermes, C.; Chandrasekaran, R.; Jacobs, J.J.; Galante, J.O.; Roebuck, K.A. & Glant, T.T. (2001). The effects of particulate wear debris, cytokines, and growth factors on the functions of MG-63 osteoblasts. *Journal of Bone and Joint Surgery*, Vol.83-A, No.2, (Feb), pp. 201-211

Vermes, C.; Roebuck, K.A.; Chandrasekaran, R.; Dobai, J.G.; Jacobs, J.J. & Glant, T.T. (2000). Particulate wear debris activates protein tyrosine kinases and nuclear factor kappaB, which down-regulates type I collagen synthesis in human osteoblasts. *Journal of Bone and Mineral Research*, Vol.15, No.9, (Sep), pp. 1756-1765

Walsh, A.J.; Nikolaou, V.S. & Antoniou, J. (2011). Inflammatory Pseudotumor Complicating Metal-On-Highly Cross-Linked Polyethylene Total Hip Arthroplasty. *Journal of Arthroplasty*, (Apr 15), ISSN 1532-8406 (Electronic)

Wang, M.L.; Nesti, L.J.; Tuli, R.; Lazatin, J.; Danielson, K.G.; Sharkey, P.F. & Tuan, R.S. (2002). Titanium particles suppress expression of osteoblastic phenotype in human mesenchymal stem cells. *Journal of Orthopaedic Research*, Vol.20, No.6, (Nov), pp. 1175-1184

Wang, M.L.; Sharkey, P.F. & Tuan, R.S. (2004). Particle bioreactivity and wear-mediated osteolysis. *Journal of Arthroplasty*, Vol.19, No.8, (Dec), pp. 1028-1038, ISSN 0883-5403

Wang, M.L.; Tuli, R.; Manner, P.A.; Sharkey, P.F.; Hall, D.J. & Tuan, R.S. (2003). Direct and indirect induction of apoptosis in human mesenchymal stem cells in response to titanium particles. *Journal of Orthopaedic Research*, Vol.21, No.4, (Jul), pp. 697-707

Wei, S.; Kitaura, H.; Zhou, P.; Ross, F.P. & Teitelbaum, S.L. (2005). IL-1 mediates TNF-induced osteoclastogenesis. *Journal of Clinical Investigation*, Vol.115, No.2, (Feb), pp. 282-290, ISSN 0021-9738

Wei, S. & Siegal, G.P. (2008). Mechanisms modulating inflammatory osteolysis: a review with insights into therapeutic targets. *Pathology, Research and Practice*, Vol.204, No.10, pp. 695-706, ISSN 0344-0338

Wilkinson, J.M.; Hamer, A.J.; Stockley, I. & Eastell, R. (2005). Polyethylene wear rate and osteolysis: critical threshold versus continuous dose-response relationship. *Journal of Orthopaedic Research*, Vol.23, No.3, (May), pp. 520-525

Willert, H.G. (1977). Reactions of the articular capsule to wear products of artificial joint prostheses. *Journal of Biomedical Materials Research*, Vol.11, No.2, (Mar), pp. 157-164

Wong, B.R.; Besser, D.; Kim, N.; Arron, J.R.; Vologodskaia, M.; Hanafusa, H. & Choi, Y. (1999). TRANCE, a TNF family member, activates Akt/PKB through a signaling complex involving TRAF6 and c-Src. *Molecular Cell*, Vol.4, No.6, (Dec), pp. 1041-1049, ISSN 1097-2765

Yamazaki, M.; Fukushima, H.; Shin, M.; Katagiri, T.; Doi, T.; Takahashi, T. & Jimi, E. (2009). Tumor necrosis factor alpha represses bone morphogenetic protein (BMP) signaling by interfering with the DNA binding of Smads through the activation of NF-kappaB. *Journal of Biological Chemistry*, Vol.284, No.51, (Dec 18), pp. 35987-35995, ISSN 1083-351X (Electronic)

Yang, S.; Wu, B.; Mayton, L.; Evans, C.H.; Robbins, P.D. & Wooley, P.H. (2002a). IL-1Ra and vIL-10 gene transfer using retroviral vectors ameliorates particle-associated inflammation in the murine air pouch model. *Inflammation Research*, Vol.51, No.7, (Jul), pp. 342-350

Yang, S.Y.; Ren, W.; Park, Y.; Sieving, A.; Hsu, S.; Nasser, S. & Wooley, P.H. (2002b). Diverse cellular and apoptotic responses to variant shapes of UHMWPE particles in a murine model of inflammation. *Biomaterials*, Vol.23, No.17, (Sep), pp. 3535-3543


Yao, J.; Cs-Szabo, G.; Jacobs, J.J.; Kuettner, K.E. & Glant, T.T. (1997). Suppression of osteoblast function by titanium particles. *Journal of Bone and Joint Surgery*, Vol.79, No.1, (Jan), pp. 107-112

Yao, Z.; Li, P.; Zhang, Q.; Schwarz, E.M.; Keng, P.; Arbini, A.; Boyce, B.F. & Xing, L. (2006). Tumor necrosis factor-alpha increases circulating osteoclast precursor numbers by promoting their proliferation and differentiation in the bone marrow through up-regulation of c-Fms expression. *Journal of Biological Chemistry*, Vol.281, No.17, (Apr 28), pp. 11846-11855, ISSN 0021-9258

Yaszay, B.; Trindade, M.C.; Lind, M.; Goodman, S.B. & Smith, R.L. (2001). Fibroblast expression of C-C chemokines in response to orthopaedic biomaterial particle challenge in vitro. *Journal of Orthopaedic Research*, Vol.19, No.5, (Sep), pp. 970-976

Zhou, F.H.; Foster, B.K.; Zhou, X.F.; Cowin, A.J. & Xian, C.J. (2006). TNF-alpha mediates p38 MAP kinase activation and negatively regulates bone formation at the injured growth plate in rats. *Journal of Bone and Mineral Research*, Vol.21, No.7, (Jul), pp. 1075-1088, ISSN 0884-0431

IntechOpen



Inflammatory Diseases - Immunopathology, Clinical and Pharmacological Bases

Edited by Dr Mahin Khatami

ISBN 978-953-307-911-0
Hard cover, 396 pages
Publisher InTech
Published online 10, February, 2012
Published in print edition February, 2012

This book is a collection of comprehensive reviews contributed by experts in the diverse fields of acute and chronic inflammatory diseases, with emphasis on current pharmacological and diagnostic options. Interested professionals are also encouraged to review the contributions made by experts in a second related book entitled "Inflammation, Chronic Diseases and Cancer"; it deals with immunobiology, clinical reviews, and perspectives of the mechanisms of immune inflammatory responses that are involved in alterations of immune dynamics during the genesis, progression and manifestation of a number of inflammatory diseases and cancers, as well as perspectives for diagnosis, and treatment or prevention of these disabling and potentially preventable diseases, particularly for the growing population of older adults around the globe.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Sang-Soo Lee, P. Edward Purdue and Ju-Suk Nam (2012). Inflammatory Periprosthetic Bone Loss, Inflammatory Diseases - Immunopathology, Clinical and Pharmacological Bases, Dr Mahin Khatami (Ed.), ISBN: 978-953-307-911-0, InTech, Available from: http://www.intechopen.com/books/inflammatory-diseases-immunopathology-clinical-and-pharmacological-bases/biology-of-inflammatory-periprosthetic-bone-loss

INTECH
open science | open minds

InTech Europe
University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China
Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the Creative Commons Attribution 3.0 License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen