

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

186,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Assessing the Influence of Neuroinflammation on Neurogenesis: *In Vitro* Models Using Neural Stem Cells and Microglia as Valuable Research Tools

Bruno P. Carreira¹, Maria Inês Morte¹,
Caetana M. Carvalho¹ and Inês M. Araújo^{1,2}

¹*Center for Neuroscience and Cell Biology, Neuroendocrinology and Neurogenesis Group,
University of Coimbra, Coimbra,*

²*Regenerative Medicine Program, Department of Biomedical Sciences and Medicine
University of Algarve, Faro,
Portugal*

1. Introduction

1.1 Neural stem cells

Neural stem cells are localized in two limited regions of the adult mammalian brain: the subgranular zone of the dentate gyrus (DG) of the hippocampus, a cell layer located between the granule cell layer and the hilus (Eriksson *et al.*, 1998; Limke and Rao, 2002), and the subventricular zone (SVZ), located next to the ependyma of the lateral walls of the lateral ventricles (Doetsch and Scharff, 2001; Curtis *et al.*, 2007). These regions are thought to provide a specific microenvironment, the stem cell niche, characterized by the presence of several agents involved in the maintenance of self-renewal and/or multipotency of neural stem cells (Alvarez-Buylla and Lim, 2004).

Although neurogenesis has been intensively studied over the past decades, only recently it has been established that newly formed neurons in the adult mammalian brain are functional and integrate into the existing neuronal network (Carlen *et al.*, 2002). The several stages of adult neurogenesis include proliferation of adult neural stem cells, fate determination, migration, integration and maturation of the newborn neurons. Using specific cell markers it is possible to independently investigate the different phases of development. Hippocampal neurogenesis plays an important role in normal hippocampal function, learning and memory (Gould *et al.*, 1999a; Shors *et al.*, 2001; Drapeau *et al.*, 2007). Newborn cells emerging from the SVZ migrate through the rostral migratory stream and integrate into the neuronal network of the olfactory bulb, establish functional synaptic connections and develop electrophysiological properties of mature neurons (Carlen *et al.*, 2002; Petreanu and Alvarez-Buylla, 2002; Belluzzi *et al.*, 2003). Furthermore, neurogenesis in the olfactory bulb is involved in important functions such as odor memory and discrimination (Gheusi *et al.*, 2000; Rochefort *et al.*, 2002; Shingo *et al.*, 2003). Under

physiological conditions, neural stem cells are tightly controlled contributing for the maintenance of brain homeostasis (Morshead *et al.*, 1994; Morshead *et al.*, 1998), however they seem to be also involved in neuronal replacement in response to pathophysiological conditions, particularly in conditions associated with neuroinflammation. Although little is known about the molecular mechanisms involved in the regulation of neural stem cells, several factors, both intrinsic and extrinsic, have been described to modulate the neurogenic process, such as hormones, trophic factors, neurotransmitters, neuromodulators and glial cells (for review see Ming and Song, 2005).

The existence of neurogenesis in areas beyond the SVZ and the DG of the adult mammalian brain have also been reported, namely in the neocortex (Gould *et al.*, 1999b; Dayer *et al.*, 2005), striatum, amygdala (Bernier *et al.*, 2002), hypothalamus (Gould *et al.*, 2001; Xu *et al.*, 2005), mesencephalon (Zhao *et al.*, 2003) and spinal cord (Yamamoto *et al.*, 2001). However, these findings need further experimental support, thus more studies need to be conducted.

1.2 Neuroinflammation

The central nervous system (CNS) was considered an immunologically privileged site, not susceptible to immune activation, due to its protection by the blood-brain barrier, which selectively allows certain inflammatory agents to enter and/or exit (Lucas *et al.*, 2006). Nowadays it is well established that immune surveillance takes place in the CNS due to the selective permeability of the blood-brain barrier to immune cells such as T cells, macrophages and dendritic cells (Hickey, 1999). Following injury or exposure to pathogens, an inflammatory response is driven by the activation of two types of immune cells: CNS resident cells, such as microglial cells and astrocytes, and CNS infiltrating cells, such as lymphocytes, monocytes and macrophages from the hematopoietic system (Stoll and Jander, 1999; Streit *et al.*, 1999). The activation of immune cells leads to the production and release of a plethora of regulatory substances, like cytokines, chemokines, neurotransmitters, reactive oxygen species and reactive nitrogen species (reviewed by Whitney *et al.*, 2009). These inflammatory mediators are essential for the recruitment of immune cells, particularly microglial cells, but also for changing the permeability of the blood-brain barrier and recruitment of monocytes and lymphocytes from the hematopoietic system to the compromised area (Hickey, 1999; Lossinsky and Shivers, 2004; Taupin, 2008), which creates a positive feedback loop to the inflammatory response.

Microglia, frequently referred to as the resident macrophages of the brain parenchyma, play a central role in the inflammatory response. Unlike astrocytes, oligodendrocytes and ependymal cells, microglial cells derive from the mesodermal germ layer. During adult life, the microglial cell pool is renewed by division of CNS resident cells. Moreover, microglia are distributed throughout the CNS with distinct densities (Lawson *et al.*, 1990). In the healthy brain, microglia are present in a resting state assuming a typical and dynamic morphology, whose function has been clarified by different studies (Davalos *et al.*, 2005; Nimmerjahn *et al.*, 2005; Davalos *et al.*, 2008). This resting state consists of a constant surveillance activity of the brain parenchyma, which enables microglial cells to screen different brain regions without disturbing the neuronal network (Hanisch and Kettenmann, 2007). Therefore, microglial cells can rapidly react to subtle homeostatic variations by changing morphology and acquiring an array of functions that allow the targeted migration

into a site of injury and release of inflammatory mediators (Gehrmann, 1996; Kreutzberg, 1996; Haynes *et al.*, 2006). Reactive microglia have the ability to rapidly upregulate a large number of receptor types, like cytokine receptors, toll-like receptors or cell adhesion molecules, but also to release a plethora of inflammatory agents (for review see Block and Hong, 2005). In fact, chemokines released by reactive microglial cells attract more microglia that, following activation, contribute to further propagate the neuroinflammatory event (Whitney *et al.*, 2009).

Astrocytes constitute the majority of glial cells in the CNS, and play an important structural function, providing support for neurons, playing also regulatory functions, including maintenance of extracellular ion balance, signaling to neurons, repair and scarring process of the CNS (Svendsen, 2002). During inflammation, astrocytes also become activated and release inflammatory factors, growth factors and excitatory amino acids, such as glutamate, which are involved in the regulation of the inflammatory response (Song *et al.*, 2002).

1.3 Neuroinflammation and neurogenesis

Neuroinflammation is a complex event with different outcomes in the neurogenic process, which can therefore enhance or suppress neurogenesis. The secreted products during inflammation have been shown to act as pro- or anti-neurogenic agents, contributing to beneficial or detrimental outcomes of neuroinflammation on the different steps of neurogenesis. Moreover, these effects seem to be particularly dependent on how and for how long microglial cells are activated. Inflammation and microglia activation were initially thought to inhibit adult neurogenesis (Ekdahl *et al.*, 2003; Monje *et al.*, 2003), while recent evidence indicates that microglia under certain circumstances can support neurogenic events (reviewed by Hanisch and Kettenmann, 2007). It has been suggested that mediators released by reactive microglia, such as cytokines and nitric oxide (NO), can inhibit adult neurogenesis in inflammatory conditions (Vallieres *et al.*, 2002; Monje *et al.*, 2003; Liu *et al.*, 2006). On the other hand, neurogenesis seems to be induced by microglial cells activated by IL-4 or low level of IFN-gamma, which has been associated with increased neuroprotection (Wong *et al.*, 2004; Song *et al.*, 2005; Baron *et al.*, 2008). Moreover, some inflammatory mediators like NO seem to have opposite roles in regulating neurogenesis in inflammatory conditions (Carreira *et al.*, 2010). Apparently, microglial cells and the factors they release play a dual role in neurogenesis acting as antiproliferative or proliferative agents. Indeed, self-renewal, proliferation, migration, differentiation, integration and, more importantly, survival of newborn neurons is modulated by the local microenvironment characterizing the neuroinflammatory response. Neural stem cells become “activated” following brain injury and migrate into the lesioned areas, which suggests that mediators present in the inflammatory microenvironment can guide the migration of newborn cells (Arvidsson *et al.*, 2002; Nakatomi *et al.*, 2002).

The role of neuroinflammation in regulating neurogenesis and neuroprotection is not clear yet, and is the subject of numerous studies (for comprehensive review see Whitney *et al.*, 2009; and Gonzalez-Perez *et al.*, 2010). There is, however, evidence for some of the most important mediators of the inflammatory response in their role in the regulation of neurogenesis and neuroprotection (Table 1).

Inflammatory factor	Neurogenesis	Neuroprotection	References
IFN-gamma	Pro-neurogenic	Decreased	(Ben-Hur <i>et al.</i> , 2003; Wong <i>et al.</i> , 2004; Butovsky <i>et al.</i> , 2006; Johansson <i>et al.</i> , 2008)
Interleukin-6	Anti-neurogenic	Decreased	(Ekdahl <i>et al.</i> , 2003; Liu <i>et al.</i> , 2005; Nakanishi <i>et al.</i> , 2007; Koo and Duman, 2008; Bauer, 2009; Islam <i>et al.</i> , 2009)
Interleukin-18			
Nitric oxide	Anti-neurogenic (nNOS) Pro-astrogliogenic (iNOS)	Decreased	(Contestabile <i>et al.</i> , 2003; Moreno-Lopez <i>et al.</i> , 2004; Matarredona <i>et al.</i> , 2005; Ciani <i>et al.</i> , 2006; Covacu <i>et al.</i> , 2006; Fritzen <i>et al.</i> , 2007; Luo <i>et al.</i> , 2007; Carreira <i>et al.</i> , 2010)
TNF-alpha	Anti-neurogenic (TNF-R1) Pro-neurogenic (TNF-R2)	Decreased or Increased	(Ben-Hur <i>et al.</i> , 2003; Wong <i>et al.</i> , 2004; Cacci <i>et al.</i> , 2005; Heldmann <i>et al.</i> , 2005; Liu <i>et al.</i> , 2005; Iosif <i>et al.</i> , 2006; Bernardino <i>et al.</i> , 2008)

Table 1. Effect of some inflammatory factors on neurogenesis and their neuroprotective role.

We are only beginning to understand how inflammatory factors and microglial cells influence neurogenesis in an inflammatory scenario, and the mechanisms, function and modulation of neurogenesis during inflammation require further investigation. This field of work is of particular interest for a better understanding of the mechanisms underlying the effects of neuroinflammation on neurogenesis, and further studies need to be conducted to increase the potential therapeutic value of regulating neuroinflammation in cellular regeneration in the diseased brain.

1.4 Brain repair and stem cell based therapies

Repair of damaged tissues is essential for the survival of living organisms. Each tissue or organ has an intrinsic, albeit limited ability for the replacement of dead cells, and correct integration of the newborn cells that, ideally, should restore the original structure. Cell replacement and correct integration of the newborn cells in the CNS is not so efficient as in other tissues such as skin or bone, which present a higher cell turnover. The CNS, on the other hand, has weak capabilities for both endogenous cell replacement and pattern repair. Some approaches have been used to attempt to develop therapeutic strategies for brain repair, namely transplantation of neural stem cells, stimulation of endogenous neurogenesis, neuroprotective strategies and anti-inflammatory approaches.

Transplantation of neural stem cells is one of the promising methods in study to be used in the reconstruction of neuronal circuits. However, the cells to be transplanted should be phenotypically plastic and able to proliferate *ex vivo* in response to external stimulus (Wang *et al.*, 1998; Sheen *et al.*, 1999). Intracerebral transplantation of SVZ-derived neural stem cells

has been successfully used in experimental models of Parkinson's disease (Zigova *et al.*, 1998; Richardson *et al.*, 2005), Huntington's disease (Vazey *et al.*, 2006), and in Multiple Sclerosis (Cayre *et al.*, 2006). Cell replacement could also be achieved by inducing endogenous neural stem cells to differentiate into neurons in the adult CNS, which consists in a less invasive strategy when compared to cell transplantation.

Indeed, *in situ* stimulation of endogenous adult neural stem cells and modulation of injury-induced neurogenesis is a therapeutic strategy, developed to upregulate endogenous neurogenesis, for instance through the control of the inflammatory response in a safe and efficient way. This approach seems to be a more advantageous strategy for multifocal diseases such as Alzheimer's disease, when compared to grafting strategies. Therefore, increased neurogenesis has been achieved by different strategies, such as administration of mitotic agents or trophic factors (Craig *et al.*, 1996; Kuhn *et al.*, 1997; Zigova *et al.*, 1998), treatment with neuroleptics like olanzepine (Green *et al.*, 2006), administration of NO donors or 5-phosphodiesterase inhibitors (Zhang *et al.*, 2003; Imitola *et al.*, 2004; Sun *et al.*, 2004; Sun *et al.*, 2006).

Other strategies designed to improve brain repair are being investigated, such as neuroprotective approaches consisting in the administration of radical scavengers, apoptosis inhibitors, neurotrophic agents, metal ions chelators and gene therapy, which seem to be useful to limit injury-induced lesion, but also for the enhancement of the survival of newborn cells (Polazzi and Monti, 2010). The use of anti-inflammatory drugs as a strategy to promote neurogenesis has also been explored and, although the chronic use of nonsteroidal anti-inflammatory drugs is detrimental for the gastrointestinal tract, it has also been associated with a decreased risk for neurodegenerative diseases (McGeer and McGeer, 1995; Lim *et al.*, 2000; Chen *et al.*, 2003). In fact, control of the inflammatory response seems to be an important strategy to increase proliferation of neural stem cells and/or differentiation of newborn neurons.

Strategies to promote regeneration of lesioned areas or cell replacement therapies will have to take into account the effects of inflammation on the formation and survival of newly generated neurons, either from the brain's own pool of neural stem cells, or from transplanted neural stem cells. Thus, the understanding of the mechanisms underlying the effect of neuroinflammation in proliferation, fate determination, migration and differentiation of neural stem cells is the first step in the development of specific strategies that could target the deleterious effect of inflammation in neurogenesis. Since the neuroinflammatory event is mostly characterized by the activation of resident microglial cells, the use of *in vitro* models that allow the study of the effects of microglia activation in the modulation of neural stem cells proliferation, fate determination, migration and differentiation into neurons is of high importance for the development of therapeutic strategies.

2. *In vitro* models to assess the crosstalk of neurogenesis and neuroinflammation

In vitro culture systems are critical tools for the study of various aspects related to the mechanisms that regulate biological functions. The removal of cells from their native microenvironment allows the study in a more focused way without the restrictions or

control of other cell types. When using *in vitro* systems it is essential to recognize that some of the isolated cells must be studied within a short period of time following isolation, or instead, the experimental model must reproduce the microenvironment of the CNS from where cells were isolated. These limitations can, however, be useful to investigate the factors that regulate the phenotype of isolated cells. Different *in vitro* models using neural stem cells and microglial cells may be used, to better understand how inflammation affects the formation of new neurons from neural stem cells.

2.1 Neural stem cell cultures

Reynolds and collaborators performed the first adult neural stem cell culture in the 90's (Reynolds *et al.*, 1992; Reynolds and Weiss, 1992), as free floating cell clusters, commonly referred to as neurospheres. These adult neural stem cells found *in vivo* were dissociated *in vitro* and kept their main properties: self-renewal capacity and multipotency, when in presence of mitogens such as basic fibroblast growth factor (bFGF) and epidermal growth factor (EGF). This cell culture system is extensively used by researchers in neural stem cell biology, and models based on adherent adult neural stem cells cultured in a monolayer on matrix are also widely used (Pollard *et al.*, 2006).

The neural stem cell cultures can be obtained from different regions from the neuroaxis of the adult mammalian CNS, from the olfactory bulb to the spinal cord, and kept in uncoated dishes under serum-free conditions plus mitogens and other essential supplements (Golmohammadi *et al.*, 2008). These adult neural stem cells can be identified based on the expression of specific protein markers such as the transcription factor Sox2, nestin, musashi-1 and the EGF receptor, among others (Kaneko *et al.*, 2000; Ming and Song, 2005). After removal of mitogens these cells can give rise to three different cell types, namely neurons, astrocytes and oligodendrocytes (Levison and Goldman, 1997; Luskin *et al.*, 1997; Palmer *et al.*, 2001; Sanai *et al.*, 2004). Thus, in cultures we can find cells expressing the referred markers but also cells expressing other specific markers, such as glial fibrillary acidic protein (GFAP), polysialylated-neural cell adhesion molecule (PSA-NCAM) and beta-III tubulin (Suslov *et al.*, 2002; Ming and Song, 2011).

It is believed that the neurosphere culture may closer resemble the *in vivo* architecture than adherent cultures since it is believed that the stem cell niche is created by clustered cells. On the other hand, the sphere size can be a limitation of this culture in comparison to adherent neural stem cell cultures since the cells that are in the sphere core can have lower access to the nutrients and oxygen, thus undergoing cell death (Ostenfeld *et al.*, 2002; Bez *et al.*, 2003).

Adult neural stem cell culturing systems have been a relevant tool in the study of biological processes within the mammalian nervous system such as neurogenesis and their distinct phases. Cultures are good platforms for expansion of adult neural stem cells, being easily manipulated without loss of function. Additionally, they can be used as experimental models for the study of differentiation and intrinsic specification, and also for screening of drugs with the potential to enhance neurogenesis. However, further investigation should be performed for characterization of stem cells in these models, since a specific marker for neural stem cells is still lacking.

On the other hand, adult stem cell cultures have some limitations, as described next. Cells are sensitive to the culturing protocols, namely the overall number of passages, mitogen

concentration and also to the methodology adopted to dissociate spheres – mechanically or by enzymatic digestion (Caldwell, 2001; Caldwell *et al.*, 2001; Morshead *et al.*, 2002; Irvin *et al.*, 2003). The overall size of spheres has been linked to the heterogeneity of sphere composition, since it increases with sphere size, the artificiality of the cell cultures, since cells propagate without instructions of their niche, and the fact that all dividing cells propagate resulting in a mixture of different cell types, are all limitations of the neurosphere culture (Reynolds and Weiss, 1996; Suslov *et al.*, 2002; Parmar *et al.*, 2003). Moreover, the non-limited expansion of cultures could be a disadvantage once the proliferative capacity could be lost by fast dividing cells over multipotent cells or by loss of stem cell capacity over the number of passages. This situation may occur at the expense of differentiation. Moreover, long-term culturing emphasizes the tendency for neural stem cells to adopt an astrocytic phenotype, with reduced capacity to generate oligodendrocytes and neurons (Chang *et al.*, 2004; Vukicevic *et al.*, 2010). Despite these limitations, free floating neural stem cell culturing systems have several advantages and are by far the most used tool concerning the study of neural stem cell biology. The use of neural stem cell cultures allows the easy access to different stages of adult neurogenesis, including proliferation of neural stem cells or progenitors, differentiation and fate determination of progenitor cells, migration of newborn cells and cell survival. By choosing the right tools and correct techniques, these different stages can be independently studied *in vitro*.

Adult neurogenesis was initially reported *in vivo* using autoradiography to track tritiated ($[^3\text{H}]$)-thymidine. $[^3\text{H}]$ -thymidine is incorporated in the DNA of dividing cells, thus proving evidence for the existence of newborn cells in the hippocampus (Altman and Das, 1965) and later, in the olfactory bulb (Altman, 1969). Proliferation of neural stem cells, the first stage of neurogenesis, can be also detected *in vitro*. Different methods have been developed since, such as the evaluation of 5-bromo-2'-deoxyuridine (BrdU) incorporation, a thymidine analogue that can be incorporated by S-phase cells during DNA synthesis, to detect cell proliferation instead of $[^3\text{H}]$ -thymidine (Gratzner, 1982; Nowakowski *et al.*, 1989). BrdU has been the golden standard in the detection of cell proliferation for the last 20 years both *in vivo* and *in vitro*. Detection of BrdU can be easily performed with antibodies, either by immunocytochemistry, microplate assay or by flow cytometry. However, BrdU detection requires aggressive treatment for DNA denaturation, in order to allow exposure of the incorporated BrdU to antibodies. Such harsh treatment can be a major drawback in the technique, as heat or acid treatment can destroy several epitopes, thus precluding multiplex labeling with other antibodies, and DNA denaturation causes the loss of binding sites for cell cycle dyes.

The use of 5-ethynyl-2'-deoxyuridine (EdU) has recently been proposed as an alternative to BrdU, since EdU detection does not require DNA denaturation, thus improving DNA structural preservation (Salic and Mitchison, 2008). EdU is also a thymidine analog that is incorporated into DNA by dividing cells during active DNA synthesis, and can be used *in vitro* as well as *in vivo* (Rostovtsev *et al.*, 2002). EdU detection is based on click chemistry, via the copper-mediated covalent coupling of the ethynyl group of EdU to a fluorescent dye-conjugated azide (Rostovtsev *et al.*, 2002). Detection can be performed by microscopy, high-throughput analysis equipment or flow cytometry. Particularly, flow cytometry is extremely useful for fast cell cycle analysis together with detection of EdU incorporation, while at same time it is possible to co-label the proliferative cells with other cell-type specific markers. The use of cell cycle markers (described next) complement detection of proliferation by ^3H -thymidine, BrdU or EdU, allowing for a more accurate timing of the birth of newborn cells

(Eisch and Mandyam, 2007). Other thymidine analogues that can be detected with antibodies are also available, such as iododeoxyuridine (IdU) and chlorodeoxyuridine (CldU).

Proteins related to the cell cycle have different expression patterns in the neurogenic regions accordingly to the phases of the cell cycle: retinoblastoma protein (Rb), a nuclear protein involved in the control of cell cycle progression, has a functional domain that binds to transcription factors and is expressed mostly in late G1 phase (Yoshikawa, 2000). Proliferating cell nuclear antigen (PCNA), a catalytic nuclear protein associated with DNA polymerase δ , is detected throughout all four phases of the cell cycle, however it is most abundant at late G1 and early S and scarce during G2 and M (Kawabe *et al.*, 2002). Ki-67, a nonhistone nuclear protein, is present during G1, S, G2 and M phase (Gerdes *et al.*, 1984). Cyclin-dependent kinase 1 (CDK1) or Cdc2 (the p34cdc2) is one of the mitosis-promoting factors and has an important role in the initiation of mitosis (Draetta *et al.*, 1988; Okano *et al.*, 1993).

Multi-labeling cells with specific cell markers and proliferation makers could easily identify newly generated neurons and glial cells, such as astrocytes and oligodendrocytes, which allows the distinction between these cell types. Proteins such as RNA-binding protein Hu and musashi-1 are exclusively expressed in mitotic active neural precursor cells, and they are absent in fully differentiated neuronal cells (Sakakibara *et al.*, 1996; Akamatsu *et al.*, 1999). The expression pattern of these markers can be detected by immunolabeling or quantitative real-time PCR (qRT-PCR). Mature neurons can be identified by assessing the presence of markers such as beta-III-tubulin, which contributes to microtubule stability in neuronal cell bodies and axons (Lee *et al.*, 1990; Memberg and Hall, 1995), or by evaluating the presence of neuronal nuclear antigen (NeuN) (Mullen *et al.*, 1992). Also the transcription factor NeuroD can be used since it is expressed throughout maturation until new neurons develop dendrites (Seki, 2002). Other markers that are commonly used can also be found in non-neuronal cells, namely PSA-NCAM (Seki and Arai, 1993; Kiss and Rougon, 1997); nestin, which is expressed in newly generated cells that still have the capacity to divide and differentiate into neurons or astrocytes (Reynolds and Weiss, 1992; Daniel *et al.*, 2008); Sox2, a transcription factor essential to maintain self-renewal of stem cells (Pevny and Placzek, 2005); and doublecortin (DCX) which has a transient expression in proliferating progenitor cells and newly generated neuroblasts or glial cells (Brown *et al.*, 2003; Kempermann *et al.*, 2003; Rao and Shetty, 2004). Oligodendrocytes are easily identified by immunolabeling against 2', 3'-cyclic nucleotide 3'-phosphodiesterase (CNPase), APC or O4 (Vernadakis *et al.*, 1984; Wu *et al.*, 2008; Girolamo *et al.*, 2010), while astrocytes can be identified by immunolabeling against GFAP, a specific protein for astrocytes (Bock *et al.*, 1977).

Concerning the migration of newly formed cells, it has been extensively studied *in vivo* (Kempermann *et al.*, 2003; Rao and Shetty, 2004), but also *in vitro*, by measuring DCX immunoreactivity (Francis *et al.*, 1999; Cohen *et al.*, 2008). DCX is a microtubule-associated protein having an important role in neuronal migration, by stabilizing microtubules and causing bundling (Sapir *et al.*, 2000). While immunolabeling is currently used, other assays have been developed in order to evaluate migration and simultaneously the mechanisms controlling cell migration, cell protrusion and cell polarization, such as the scratch-wound migration assay (Etienne-Manneville, 2006). Additionally, Durbec and collaborators compared three different assays to evaluate migration of neural stem cells *in vitro*: matrigel, a three-dimensional substrate mimicking the *in vivo* extracellular matrix, detection of soluble factors influencing radial migration and the chemotaxis chamber assay, where the researcher can evaluate whether the cells prefer or not a chemical factor (Durbec *et al.*, 2008).

When mature, not all neurons in culture are functional or survive. It is important to check their viability, namely identify functional synapses by morphological, electrophysiological and immunological characterization (Hartley *et al.*, 1999). Several methods have been used, including immunocytochemical assays, Western blotting and qRT-PCR which allow identification and quantification of proteins, neurotransmitters, neurotrophic factors, among others, involved in neuronal or glial neurotransmitter systems (Hartley *et al.*, 1999; Elmariah *et al.*, 2005; Goodfellow *et al.*, 2011). Using patch-clamp techniques *in vitro* the electrophysiological characterization of neural stem cell cultures can be performed by evaluating the formation of action potentials and activity patterns (Li *et al.*, 2008; Cheyne *et al.*, 2011). Also single-cell calcium currents may be evaluated to discriminate neuronal profile and viability in response to different stimuli, as reported by Bernardino and collaborators (Bernardino *et al.*, 2008).

2.2 Microglial cell cultures

Microglial cells may be obtained for culturing by several methods. One of the most used models for the study of microglial cell function consists in the isolation and expansion of microglia from the neonatal brain. However, there are several limitations and criticisms to this approach since it consists in the isolation of microglial cells from the neonatal brain, not the adult brain. One of the main problems associated with the use of microglial cells *in vitro* is related to the characterization of microglia phenotype. Since there are no truly, unique and specific microglial cell markers, microglia phenotype is defined through a combined analysis of morphology and presence or absence of certain antigens. Several works lack a proper evaluation of microglia phenotype that would allow to distinguish microglia from macrophages. In most studies, the presence of microglial cell markers is excluded from cells that are positive for astrocytic or neuronal markers, but do not distinguish between microglia or macrophages. One of the most used immunocytochemical marker of microglial cells that is the ionized calcium binding adapter molecule 1 (Iba1) (Ito *et al.*, 1998). Other markers that have been identified include the beta-integrin marker CD11b (Ling and Wong, 1993; Gonzalez-Scarano and Baltuch, 1999), the glucose transporter 5 (GLUT5) (Sasaki *et al.*, 2004), CD163 (Roberts *et al.*, 2004; Borda *et al.*, 2008), CCR2 (Albright *et al.*, 1999; Zhang *et al.*, 2007), CD34 (Asheuer *et al.*, 2004; Ladeby *et al.*, 2005) and C-type lectin CD209b (Park *et al.*, 2009). Toll-like receptor 2 (TLR2) and Toll-like receptor 4 (TLR4) have been also used as markers of microglial cells as they appear to be involved in determining the phenotype and function of microglia (Li *et al.*, 2009). A combination of several of these markers would allow for a better characterization of microglia phenotype, rather than the use of a single marker, which is the current standard. The use of multiplex detection systems would be the best approach for a full molecular characterization of microglia (Albright and Gonzalez-Scarano, 2004; Duke *et al.*, 2004; Gebicke-Haerter, 2005; Glanzer *et al.*, 2007; Moran *et al.*, 2007).

The most popular protocol to isolate microglial cells is the shaking method described by Guilian and Baker (Giulian and Baker, 1986) and Frei and colleagues (Frei *et al.*, 1986). In this method, microglial cells are separated from confluent primary mixed glial cultures, isolated from the rodent neonatal cortex, by agitation in an orbital shaker. Although this method allows the preparation of highly pure microglial cultures, the yield of this protocol is low. Saura and colleagues described a method to isolate microglial cells from primary mixed glial cultures of rodent brain by a mild trypsinization protocol, which allows the preparation of

high purity microglial cultures, with a higher yield when compared to the shaking method (Saura *et al.*, 2003). Similarly to the shaking method, several works describe the isolation of microglia from adult rodents, and the large majority of these studies take advantage from the astrocyte-microglia interaction for the success of cell cultures (Rosenstiel *et al.*, 2001; Ponomarev *et al.*, 2005). These studies showed that microglial cells, when grown on a monolayer of astrocytes, develop a highly branched morphology which seems to be associated with the downregulation of the nuclear factor kappa B (NF-kappaB) (Rosenstiel *et al.*, 2001). It has been shown that microglial cells isolated from the neonatal or adult brain are sensitive to the treatment with granulocyte macrophage colony-stimulating factor (GM-CSF), which induced a differentiation into a phenotype more similar to those of dendritic cells (Suzumura *et al.*, 1990; Aloisi, 2001). On the other hand, the isolation of adult microglial cells and subsequent culture with low concentrations of macrophage colony-stimulating factor (M-CSF) leads to increased proliferation and survival of cells that persists for several weeks (Suzumura *et al.*, 1990; Ponomarev *et al.*, 2005). M-CSF seems to be a key factor for the maintenance and survival of microglial cells *in vitro*, and has been used in several works (Wegiel *et al.*, 1998; Ponomarev *et al.*, 2005; Carreira *et al.*, 2010). Other methods are also described for the isolation of microglial cells, which include isolation from CNS tissue by Percoll gradient (Dick *et al.*, 1995; Ford *et al.*, 1995), isolation from primary cultures by nutritional deprivation (Hao *et al.*, 1991) or by collecting floating cells in mixed glial cultures (Ganter *et al.*, 1992), but the yield is generally very low.

The use of *in vitro* models allows for the understanding of many aspects of the dynamics associated with the biological functions of microglial cells in a quick and simple manner. However, one cannot overlook that the relevance of the observations obtained can only be extrapolated following *in vivo* studies. Several groups work with microglial cell lines, such as BV-2, HAPI or N9, however the use of microglial cell lines should be carefully considered since immortalization could significantly affect cell biology when compared to the use of primary microglial cultures (Corradin *et al.*, 1993; Lockhart *et al.*, 1998; Horvath *et al.*, 2008).

Concerning primary cultures of microglial cells it is always important to assess the purity of the cultures, this parameter being intrinsically linked to the method of isolation adopted. The isolation method described by Saura and collaborators is, therefore, one of the methods that seems to offer the best value yield/purity (Saura *et al.*, 2003). We favor the isolation of microglial cells by shaking from mixed glial cultures treated with low levels of M-CSF as an alternative to the method of Saura (Saura *et al.*, 2003), with a high purity of the microglia obtained (>90%) and, unlike previous methods, with a high yield (Carreira *et al.*, 2010).

When microglial cells become activated in response to immunologic stimuli or brain injury, activation is characterized by changes in microglia morphology (Streit *et al.*, 1988; Kreutzberg, 1996; Streit *et al.*, 1999; Liu and Hong, 2003), from resting ramified into activated amoeboid microglia (Kreutzberg, 1996). There is also a complex cellular response after activation of microglial cells, which is characterized by upregulation of surface molecules, such as complement receptors and major histocompatibility complex molecules (Oehmichen and Gencic, 1975; Graeber *et al.*, 1988). In addition, activated microglia release a large variety of soluble factors, with a pro- or anti-inflammatory nature and potentially cytotoxic (for review see Block and Hong, 2005). It is therefore important, when establishing primary

cultures of microglia, to assess whether microglial cells *in vitro* are also responsive to inflammatory stimuli similarly to what occurs *in vivo*. Microglial cells can be challenged with different stimuli *in vitro*, and by far the most widely used stimulus in primary cultures of microglia isolated from rodents is the bacterial endotoxin lipopolysaccharide (LPS) (Qin *et al.*, 2005a; Qin *et al.*, 2005b; Pei *et al.*, 2007). LPS mimics the infection by Gram-negative bacteria, which induces an increase in the synthesis of inflammatory mediators, namely cytokines, such as IL-1, IL-6 and tumor necrosis factor-alpha (TNF-alpha), chemokines, such as stromal derived factor-1 alpha (SDF-1alpha), free radicals and nitric oxide (Block and Hong, 2005). Other stimuli may consist in the use of ATP, interleukins, IFN-gamma or LPS plus IFN-gamma (Wollmer *et al.*, 2001; Saura *et al.*, 2003).

To characterize the activation of microglial cells after an inflammatory stimulus, we suggest to define at least three parameters to evaluate the activation of microglial cells following exposure to an inflammatory stimulus, including: change to an amoeboid morphology (Suzumura *et al.*, 1991; Wollmer *et al.*, 2001), the expression of NF-kappaB (Heyen *et al.*, 2000; Wollmer *et al.*, 2001), expression of the inducible nitric oxide synthase (iNOS) and subsequent evaluation of the production of NO (Boje and Arora, 1992; Chao *et al.*, 1992b), or the release of TNF-alpha (Sawada *et al.*, 1989; Chao *et al.*, 1992a). The various mechanisms by which microglial cells are activated and the identity of the inflammatory factors released by microglia have been studied and characterized, but there still is a great controversy whether these factors are neuroprotective or neurotoxic when released. The hypothesis that seems to be more acceptable is that, depending on the aggressiveness of the inflammatory response, the activation of microglial cells may shift from a beneficial to a harmful outcome for neurogenesis.

2.3 Combination of neural stem cells and microglial cell cultures

The study of the link between brain inflammation and neurogenesis, in particular the role of microglia in the modulation of the various steps of the neurogenic process, is of particular relevance. In order to operate at a therapeutic level there is an urgent need to understand the crosstalk between microglia and neural stem cells and the implications of the inflammatory response for the neurogenic outcome. Several studies *in vivo* have been developed in recent years, but the potential of *in vitro* studies becomes indisputable when the aim is to study the effect of a particular inflammatory factor or a very specific parameter related to the inflammatory response and its effect on neurogenesis. Whether the function of microglial cells is pro- or anti-neurogenic and whether it is possible to control microglial activation in order to reach a beneficial effect are important questions that need to be answered. Thus, the development of basic models for the *in vitro* study of these issues is an asset to the studies in this area. The use of combined primary neuronal and microglial cell cultures has been a very useful tool in studying the effect of the inflammatory response on neurons from different brain regions. In fact, there are numerous published studies where different approaches have been adopted for the study of the crosstalk between microglial cells and neurons *in vitro* (Boje and Arora, 1992; Lambertsen *et al.*, 2009). Here we describe the use of three different *in vitro* models, which address different aspects of the effects of inflammatory factors released by microglial cells in the neurogenic process.

2.3.1 Co-cultures of neural stem cells with microglia

The inflammatory response has been identified as responsible for the down-regulation of neurogenesis. This hypothesis has been supported by several studies *in vivo* (Ekdahl *et al.*, 2003; Monje *et al.*, 2003), but also by *in vitro* studies where the survival of new neurons is compromised when these are co-cultured with microglial cells activated by LPS (Monje *et al.*, 2003; Cacci *et al.*, 2005; Liu *et al.*, 2005; Cacci *et al.*, 2008). Co-cultures of neural stem cells with microglia, without physical contact between the two cell types, is an experimental model that allows the researcher to assess the role of soluble neuroinflammatory factors using co-cultures of microglial cells seeded in membrane inserts placed on top of multiwell plates containing neural stem cells. The use of techniques of immunodepletion, but also the use of genetically modified animals, allowed to correlate this anti-neurogenic inflammatory response to different interleukins produced during the activation of microglial cells, including IL-6 and IL-1beta (Vallieres *et al.*, 2002; Monje *et al.*, 2003; Nakanishi *et al.*, 2007; Goshen *et al.*, 2008; Koo and Duman, 2008; Spulber *et al.*, 2008). Other factors involved in the inflammatory response appear to contribute to the inhibition of neurogenesis. For example, the increased production of TNF-alpha by microglial cells appears to reduce the survival and differentiation of neural stem cells (Vezzani *et al.*, 2002; Monje *et al.*, 2003; Liu *et al.*, 2005; Iosif *et al.*, 2006).

Although some studies have described IFN-gamma as having a deleterious effect on neurogenesis, it has been demonstrated that microglia stimulated with low levels of IFN-gamma can support the neurogenic process, promoting neuronal differentiation *in vitro* (Butovsky *et al.*, 2006). In other studies it was observed that IFN-gamma is involved in the modulation of proliferation and differentiation of neural stem cells into neurons (Wong *et al.*, 2004; Song *et al.*, 2005; Baron *et al.*, 2008). Recent *in vitro* studies based on the establishment of co-cultures of microglia and neural stem cells, without physical contact between cells, reported that microglia might have a more complex role in neurogenesis contrarily to initial thoughts. Microglia seems to play a dual role in adult neurogenesis, being detrimental or beneficial and support the different steps in neurogenesis, such as stem cell proliferation, differentiation, migration and survival (reviewed in Ekdahl *et al.*, 2009). This dual effect becomes associated to different soluble factors produced by activated microglial cells, such as TNF-alpha or nitric oxide.

The establishment of experimental models such as co-cultures of microglia and neural stem cells allows to mimic the chemical microenvironment that surrounds the SVZ and/or the DG during inflammatory conditions when microglial cells are recruited and activated. On the other hand, the fact that both cell types share the same culture environment is important to determine the effect of factors produced by microglial cells on neural stem cells. The fact that this is a system without physical contact between the two cell types also allows determining more quickly, and using more economic approaches, the modulation of the multistep neurogenic process mediated by the inflammatory response. Thus, experimental approaches to determine cell proliferation and cell cycle, such as flow cytometry, cell migration, could be performed without the need for prior characterization to distinguish neural stem cells from microglial cells as in mixed cultures. Moreover, signaling pathways present in both cell types can be studied this way, as is the case of TLR4 that directly modulates self-renewal and the decision-cell-fate in neural stem cells (Rolls *et al.*, 2007) and in microglial cells is involved in its activation, particularly in the regulation of gene expression of iNOS (Graeber and Streit, 2010).

However, there are also some disadvantages associated with the use of this experimental methodology. Firstly, the fact that it does not allow an easy processing of microglia cells, which are placed in membrane inserts, after experimental treatment. In fact, simple experimental procedures such as protein, RNA or DNA extraction from microglial cells becomes difficult to perform. On the other hand, it is not possible to perform immunostaining techniques for subsequent microscopic analysis of microglial cells plated in inserts. In addition, this model does not answer a question that seems to be increasingly important which is the influence of cell-to-cell contact in the modulation of neurogenesis by the inflammatory response (Song *et al.*, 2002; Aarum *et al.*, 2003; Alvarez-Buylla and Lim, 2004). Despite these disadvantages, the use of co-cultures of neural stem cells with microglia, without physical contact between the two cell types, is a good approach for some studies.

2.3.2 Neural stem cell cultures exposed to microglia-conditioned medium

The production of cytokines and other molecules by activated microglial cells with implications in cellular processes has been demonstrated in many studies based on *in vitro* models (Banati *et al.*, 1993; Minghetti and Levi, 1998; Gebicke-Haerter *et al.*, 2001; Hanisch, 2002; Hausler *et al.*, 2002). However, there is still much to be learned about how cellular pathways in neural stem cells are regulated by these soluble factors from microglial origin. It is therefore important to assess how these diffusible factors influence phenomena as diverse as proliferation, differentiation, migration or cell survival.

Culturing neural stem cells with microglia conditioned medium, obtained from a separate microglia culture, allows the isolation of the unidirectional communication between activated microglia and neural stem cells, with further investigation of soluble inflammatory factors. According to studies using this experimental model, the conditioned medium of microglial cells acutely challenged with LPS reduced the survival of neural stem cells, preventing their differentiation into neurons (Monje *et al.*, 2003; Cacci *et al.*, 2008). One of the inflammatory agents reported to be responsible for this antineurogenic effect is the cytokine IL-6, as evidenced by the works of Monje and collaborators or Nakanishi and colleagues that by using a specific antibody against IL-6 rescued neurogenesis (Monje *et al.*, 2003; Nakanishi *et al.*, 2007). On the other hand, several *in vitro* studies described a pro-neurogenic effect of microglial cells and their conditioned medium, in which neural stem cells grow (Aarum *et al.*, 2003; Morgan *et al.*, 2004; Walton *et al.*, 2006; Nakanishi *et al.*, 2007).

Despite the advantages of this experimental model, namely the fact that it allows a study of the unidirectional effect of microglia on neural stem cells, there are also some disadvantages. This model does not allow inferring any conclusion about the influence of cell-to-cell contact between microglia and neural stem cells, an event that has been described to occur between glial cells and neural stem cells (Song *et al.*, 2002; Aarum *et al.*, 2003; Alvarez-Buylla and Lim, 2004). On the other hand, this model completely neglects the fact that some of the factors released by microglial cells have physical characteristics that do not allow their study in a conditioned medium transferred from a cell culture to another. Particularly nitric oxide, a gaseous molecule with a short half-life, cannot be studied because it is highly reactive in aqueous solution at 37 °C and physiological pH

(pH = 7.4). Thus, although stable end products of NO can be detected in conditioned medium from activated microglial cell cultures, the effect of NO in the neural stem cells cannot be analyzed. These are negative aspects that must be taken into account when a researcher decides to select this experimental model. Despite these aspects, the use of conditioned medium of microglia in cultures of neural stem cells is a good model to further study the influence of inflammation on neurogenesis. This model is useful to complement other *in vitro* approaches, including co-cultures of microglia and neural stem cells, with or without physical contact.

2.3.3 Mixed cultures of neural stem cells with microglia

The progression of the neurogenic process until the differentiation of neural stem cells into neurons appears to be regulated by the inflammatory microenvironment but also by cell-to-cell interactions involved (Arvidsson *et al.*, 2002; Nakatomi *et al.*, 2002; Ben-Hur *et al.*, 2003; Thored *et al.*, 2006; Thored *et al.*, 2009). Therefore, the optimization of an *in vitro* system that allows the study of physical interactions between microglia and neural stem cells is of great interest to understand how both cell types crosstalk in inflammatory conditions.

Mixed cultures are co-cultures of neural stem cells with microglia with physical contact between the two cell types. In this culture model, the role of physical contacts between microglia and neural stem cells can be studied. The mixed culture system is, probably, the *in vitro* approach that more closely mimics what happens *in vivo*, where microglial cells physically contact with the neural stem cells from neurogenic areas. Adopting this experimental model, the researcher can study the influence of the inflammatory response on the several steps of the neurogenic process, but also cell-cell interactions, which is an advantage compared to the *in vitro* models already described. An example of a mixed culture of neural stem cells cultured together with forebrain microglia is shown in Fig. 1. Enhanced green fluorescent protein (EGFP)-positive SVZ cells were isolated from the SVZ of postnatal day 1-3 actin-EGFP C57Bl6 mice, thus being readily distinguishable from microglia isolated from wild-type mice (Fig. 1A).

The mixed culture model allows simultaneous evaluation of microglia and neural stem cells. Thus, following stimulation of microglial cells, the researcher can evaluate the activation of these cells as well as several biological processes of neural stem cells, such as proliferation, differentiation and/or survival. Moreover, multi-labeling experiments of proliferation markers, such as BrdU or EdU (Fig. 1B), with microglia-specific (Iba-1 or CD11b), neuron-specific (NeuN or Tuj-1) or glia-specific (GFAP) proteins by confocal microscopy or flow cytometry are a good way to determine the phenotype of proliferating cells (Nixon and Crews, 2004). In addition, it is also possible to evaluate the effect of diffusible factors that are produced following activation of microglial cells. Separation of the two cell populations for posterior analysis (e.g. of protein or nucleic acids) is possible using a cell sorter. The researcher can confirm whether the effects observed in mixed cultures are caused by physical interactions or by diffusible factors released by microglial cells by combining such experiments with a comparative study using co-cultured cells without physical contact.




Fig. 1. Mixed cultures of primary microglial cells and subventricular zone (SVZ)-derived neural stem cells. SVZ cells (isolated from transgenic mice expressing green fluorescence protein (GFP) under the actin promoter (shown in white) are readily distinguishable from CD11b-positive microglia (red) (A). Microglia (red) cultured with GFP-positive SVZ cells (white) show immunoreactivity for inducible nitric oxide synthase (iNOS, green), following treatment with lipopolysaccharide (LPS; 100 ng/ml) plus interferon-gamma (IFN-gamma; 0.5 ng/ml), for 24 h. Nuclei are labeled with Hoechst 33342 (blue). Scale bar: 20 μm. B) Stimulation with LPS plus IFN-gamma decreases the proliferation of GFP-positive SVZ-derived neural stem cells (green), in mixed cultures of SVZ and microglia obtained from wild type mice (iNOS^{+/+}), which are CD11b-positive (red). Cell proliferation was assessed by 5-ethynyl-2'-deoxyuridine (EdU) incorporation (white). The antiproliferative effect of LPS plus IFN-gamma on EdU incorporation is abolished in mixed cultures in which the microglia was obtained from iNOS-knockout mice (iNOS^{-/-}). Scale bar: 20 μm.

3. Summary and future directions

Microglial cells may cause different effects on the neurogenic process, promoting or inhibiting it. Experimental evidence has been presented indicating that microglia, depending on their activation status and phenotype, could favor or hinder adult neurogenesis, in physiological or pathophysiological conditions. In fact, microglia can have a dual role in different steps of the neurogenic process, namely in the formation, maturation and integration of newly formed neurons. Therefore the need to explore in more detail how microglia regulate adult neurogenesis in physiological and pathophysiological conditions is of particular importance (Graeber and Streit, 2010).

Genetic mouse models in which the researcher can selectively ablate genes have already been described as useful strategies to study the involvement of particular effectors of the neuroinflammatory response on neural stem cells. Experimental models may have as an objective the determination of how modulation of microglial cell activation can be used as a therapeutic target to regulate neurogenesis in the adult brain (Ekdahl *et al.*, 2009; Whitney *et al.*, 2009; Polazzi and Monti, 2010). These models are suitable to evaluate the neurogenic potential of anti-inflammatory drugs or identify pro-neurogenic targets. Thus, these experimental approaches will allow the design of therapeutic strategies to enhance the formation, proper migration, differentiation, integration and survival of new neuronal cells in the injured nervous system. Moreover, all culture models are suitable for pharmacological or genetic manipulation, including obtaining the cells used in the cultures from wild-type or genetically modified animals, and can be adapted for high-throughput analysis and drug screening. The use of anti-inflammatory drugs with a selective mechanism of action at the level of microglial cells, or the use of anti-inflammatory drugs which may release molecules that may enhance the neurogenesis are strategies under investigation (Keeble and Moore, 2002; Napoli and Ignarro, 2003; Ajmone-Cat *et al.*, 2008; Koc and Kucukguzel, 2009). In order to develop more specific therapeutic interventions in the future, it is necessary to identify the mechanisms and factors that regulate the switch between the enhancing or detrimental effect of the inflammatory response on neurogenic events. The *in vitro* strategies discussed here are important as a first step in identifying and characterizing these events (Table 2).

Experimental model	Parameters evaluated			
	Diffusible/soluble factors	Cell-to-cell interaction	Cellular characterization	Protein, RNA and DNA content
Co-culture	Very Good	-	Very Good	Very Good
Conditioned medium	Good	-	Very Good	Very Good
Mixed culture	Very Good	Very Good	Good (requires multiplex analysis)	Good (requires cell sorting)

Table 2. Evaluation of experimental *in vitro* models using neural stem cells and microglial cells as research tools to evaluate the effect of neuroinflammation in the neurogenesis.

4. References

- Aarum J, Sandberg K, Haeberlein SL and Persson MA (2003). "Migration and differentiation of neural precursor cells can be directed by microglia." *Proc Natl Acad Sci U S A* 100(26): 15983-8.
- Ajmone-Cat MA, Cacci E and Minghetti L (2008). "Non steroidal anti-inflammatory drugs and neurogenesis in the adult mammalian brain." *Curr Pharm Des* 14(14): 1435-42.
- Akamatsu W, Okano HJ, Osumi N, Inoue T, Nakamura S, Sakakibara S, Miura M, Matsuo N, Darnell RB and Okano H (1999). "Mammalian ELAV-like neuronal RNA-binding proteins HuB and HuC promote neuronal development in both the central and the peripheral nervous systems." *Proc Natl Acad Sci U S A* 96(17): 9885-90.
- Albright AV and Gonzalez-Scarano F (2004). "Microarray analysis of activated mixed glial (microglia) and monocyte-derived macrophage gene expression." *J Neuroimmunol* 157(1-2): 27-38.
- Albright AV, Shieh JT, Itoh T, Lee B, Pleasure D, O'Connor MJ, Doms RW and Gonzalez-Scarano F (1999). "Microglia express CCR5, CXCR4, and CCR3, but of these, CCR5 is the principal coreceptor for human immunodeficiency virus type 1 dementia isolates." *J Virol* 73(1): 205-13.
- Aloisi F (2001). "Immune function of microglia." *Glia* 36(2): 165-79.
- Altman J (1969). "Autoradiographic and histological studies of postnatal neurogenesis. IV. Cell proliferation and migration in the anterior forebrain, with special reference to persisting neurogenesis in the olfactory bulb." *J Comp Neurol* 137(4): 433-57.
- Altman J and Das GD (1965). "Autoradiographic and histological evidence of postnatal hippocampal neurogenesis in rats." *J Comp Neurol* 124(3): 319-35.
- Alvarez-Buylla A and Lim DA (2004). "For the long run: maintaining germinal niches in the adult brain." *Neuron* 41(5): 683-6.
- Arvidsson A, Collin T, Kirik D, Kokaia Z and Lindvall O (2002). "Neuronal replacement from endogenous precursors in the adult brain after stroke." *Nat Med* 8(9): 963-70.
- Asheuer M, Pflumio F, Benhamida S, Dubart-Kupperschmitt A, Fouquet F, Imai Y, Aubourg P and Cartier N (2004). "Human CD34+ cells differentiate into microglia and express recombinant therapeutic protein." *Proc Natl Acad Sci U S A* 101(10): 3557-62.
- Banati RB, Gehrmann J, Schubert P and Kreutzberg GW (1993). "Cytotoxicity of microglia." *Glia* 7(1): 111-8.
- Baron R, Nemirovsky A, Harpaz I, Cohen H, Owens T and Monsonego A (2008). "IFN-gamma enhances neurogenesis in wild-type mice and in a mouse model of Alzheimer's disease." *Faseb J* 22(8): 2843-52.
- Bauer S (2009). "Cytokine control of adult neural stem cells." *Ann N Y Acad Sci* 1153: 48-56.
- Belluzzi O, Benedusi M, Ackman J and LoTurco JJ (2003). "Electrophysiological differentiation of new neurons in the olfactory bulb." *J Neurosci* 23(32): 10411-8.
- Ben-Hur T, Ben-Menachem O, Furer V, Einstein O, Mizrachi-Kol R and Grigoriadis N (2003). "Effects of proinflammatory cytokines on the growth, fate, and motility of multipotential neural precursor cells." *Mol Cell Neurosci* 24(3): 623-31.
- Bernardino L, Agasse F, Silva B, Ferreira R, Grade S and Malva JO (2008). "Tumor necrosis factor-alpha modulates survival, proliferation, and neuronal differentiation in neonatal subventricular zone cell cultures." *Stem Cells* 26(9): 2361-71.

- Bernier PJ, Bedard A, Vinet J, Levesque M and Parent A (2002). "Newly generated neurons in the amygdala and adjoining cortex of adult primates." *Proc Natl Acad Sci U S A* 99(17): 11464-9.
- Bez A, Corsini E, Curti D, Biggiogera M, Colombo A, Nicosia RF, Pagano SF and Parati EA (2003). "Neurosphere and neurosphere-forming cells: morphological and ultrastructural characterization." *Brain Res* 993(1-2): 18-29.
- Block ML and Hong JS (2005). "Microglia and inflammation-mediated neurodegeneration: multiple triggers with a common mechanism." *Prog Neurobiol* 76(2): 77-98.
- Bock E, Moller M, Nissen C and Sensenbrenner M (1977). "Glial fibrillary acidic protein in primary astroglial cell cultures derived from newborn rat brain." *FEBS Lett* 83(2): 207-11.
- Boje KM and Arora PK (1992). "Microglial-produced nitric oxide and reactive nitrogen oxides mediate neuronal cell death." *Brain Res* 587(2): 250-6.
- Borda JT, Alvarez X, Mohan M, Hasegawa A, Bernardino A, Jean S, Aye P and Lackner AA (2008). "CD163, a marker of perivascular macrophages, is up-regulated by microglia in simian immunodeficiency virus encephalitis after haptoglobin-hemoglobin complex stimulation and is suggestive of breakdown of the blood-brain barrier." *Am J Pathol* 172(3): 725-37.
- Brown JP, Couillard-Despres S, Cooper-Kuhn CM, Winkler J, Aigner L and Kuhn HG (2003). "Transient expression of doublecortin during adult neurogenesis." *J Comp Neurol* 467(1): 1-10.
- Butovsky O, Ziv Y, Schwartz A, Landa G, Talpalar AE, Pluchino S, Martino G and Schwartz M (2006). "Microglia activated by IL-4 or IFN-gamma differentially induce neurogenesis and oligodendrogenesis from adult stem/progenitor cells." *Mol Cell Neurosci* 31(1): 149-60.
- Cacci E, Ajmone-Cat MA, Anelli T, Biagioni S and Minghetti L (2008). "In vitro neuronal and glial differentiation from embryonic or adult neural precursor cells are differently affected by chronic or acute activation of microglia." *Glia* 56(4): 412-25.
- Cacci E, Claasen JH and Kokaia Z (2005). "Microglia-derived tumor necrosis factor- α exaggerates death of newborn hippocampal progenitor cells in vitro." *J Neurosci Res* 80(6): 789-97.
- Caldwell MA (2001). "Recent advances in neuralstem cell technologies." *Trends Neurosci* 24(2): 72-4.
- Caldwell MA, He X, Wilkie N, Pollack S, Marshall G, Wafford KA and Svendsen CN (2001). "Growth factors regulate the survival and fate of cells derived from human neurospheres." *Nat Biotechnol* 19(5): 475-9.
- Carlen M, Cassidy RM, Brismar H, Smith GA, Enquist LW and Frisen J (2002). "Functional integration of adult-born neurons." *Curr Biol* 12(7): 606-8.
- Carreira BP, Morte MI, Inacio A, Costa G, Rosmaninho-Salgado J, Agasse F, Carmo A, Couceiro P, Brundin P, Ambrosio AF, Carvalho CM and Araujo IM (2010). "Nitric oxide stimulates the proliferation of neural stem cells bypassing the epidermal growth factor receptor." *Stem Cells* 28(7): 1219-30.
- Cayre M, Bancila M, Virard I, Borges A and Durbec P (2006). "Migrating and myelinating potential of subventricular zone neural progenitor cells in white matter tracts of the adult rodent brain." *Mol Cell Neurosci* 31(4): 748-58.

- Chang MY, Park CH, Lee SY and Lee SH (2004). "Properties of cortical precursor cells cultured long term are similar to those of precursors at later developmental stages." *Brain Res Dev Brain Res* 153(1): 89-96.
- Chao CC, Hu S, Close K, Choi CS, Molitor TW, Novick WJ and Peterson PK (1992a). "Cytokine release from microglia: differential inhibition by pentoxifylline and dexamethasone." *J Infect Dis* 166(4): 847-53.
- Chao CC, Hu S, Molitor TW, Shaskan EG and Peterson PK (1992b). "Activated microglia mediate neuronal cell injury via a nitric oxide mechanism." *J Immunol* 149(8): 2736-41.
- Chen H, Zhang SM, Hernan MA, Schwarzschild MA, Willett WC, Colditz GA, Speizer FE and Ascherio A (2003). "Nonsteroidal anti-inflammatory drugs and the risk of Parkinson disease." *Arch Neurol* 60(8): 1059-64.
- Cheyne JE, Grant L, Butler-Munro C, Foote JW, Connor B and Montgomery JM (2011). "Synaptic integration of newly generated neurons in rat dissociated hippocampal cultures." *Mol Cell Neurosci* 47(3): 203-14.
- Ciani E, Calvanese V, Crochemore C, Bartesaghi R and Contestabile A (2006). "Proliferation of cerebellar precursor cells is negatively regulated by nitric oxide in newborn rat." *J Cell Sci* 119(Pt 15): 3161-70.
- Cohen D, Segal M and Reiner O (2008). "Doublecortin supports the development of dendritic arbors in primary hippocampal neurons." *Dev Neurosci* 30(1-3): 187-99.
- Contestabile A, Monti B and Ciani E (2003). "Brain nitric oxide and its dual role in neurodegeneration/neuroprotection: understanding molecular mechanisms to devise drug approaches." *Curr Med Chem* 10(20): 2147-74.
- Corradin SB, Mauer J, Donini SD, Quattrocchi E and Ricciardi-Castagnoli P (1993). "Inducible nitric oxide synthase activity of cloned murine microglial cells." *Glia* 7(3): 255-62.
- Covacu R, Danilov AI, Rasmussen BS, Hallen K, Moe MC, Lobell A, Johansson CB, Svensson MA, Olsson T and Brundin L (2006). "Nitric oxide exposure diverts neural stem cell fate from neurogenesis towards astrogliogenesis." *Stem Cells* 24(12): 2792-800.
- Craig CG, Tropepe V, Morshead CM, Reynolds BA, Weiss S and van der Kooy D (1996). "In vivo growth factor expansion of endogenous subependymal neural precursor cell populations in the adult mouse brain." *J Neurosci* 16(8): 2649-58.
- Curtis MA, Eriksson PS and Faull RL (2007). "Progenitor cells and adult neurogenesis in neurodegenerative diseases and injuries of the basal ganglia." *Clin Exp Pharmacol Physiol* 34(5-6): 528-32.
- Daniel C, Albrecht H, Ludke A and Hugo C (2008). "Nestin expression in repopulating mesangial cells promotes their proliferation." *Lab Invest* 88(4): 387-97.
- Davalos D, Grutzendler J, Yang G, Kim JV, Zuo Y, Jung S, Littman DR, Dustin ML and Gan WB (2005). "ATP mediates rapid microglial response to local brain injury in vivo." *Nat Neurosci* 8(6): 752-8.
- Davalos D, Lee JK, Smith WB, Brinkman B, Ellisman MH, Zheng B and Akassoglou K (2008). "Stable in vivo imaging of densely populated glia, axons and blood vessels in the mouse spinal cord using two-photon microscopy." *J Neurosci Methods* 169(1): 1-7.
- Dayer AG, Cleaver KM, Abouantoun T and Cameron HA (2005). "New GABAergic interneurons in the adult neocortex and striatum are generated from different precursors." *J Cell Biol* 168(3): 415-27.

- Dick AD, Ford AL, Forrester JV and Sedgwick JD (1995). "Flow cytometric identification of a minority population of MHC class II positive cells in the normal rat retina distinct from CD45^{low}CD11b/c+CD4^{low} parenchymal microglia." *Br J Ophthalmol* 79(9): 834-40.
- Doetsch F and Scharff C (2001). "Challenges for brain repair: insights from adult neurogenesis in birds and mammals." *Brain Behav Evol* 58(5): 306-22.
- Draetta G, Brizuela L, Moran B and Beach D (1988). "Regulation of the vertebrate cell cycle by the cdc2 protein kinase." *Cold Spring Harb Symp Quant Biol* 53 Pt 1: 195-201.
- Drapeau E, Montaron MF, Aguerre S and Abrous DN (2007). "Learning-induced survival of new neurons depends on the cognitive status of aged rats." *J Neurosci* 27(22): 6037-44.
- Duke DC, Moran LB, Turkheimer FE, Banati R and Graeber MB (2004). "Microglia in culture: what genes do they express?" *Dev Neurosci* 26(1): 30-7.
- Durbec P, Franceschini I, Lazarini F and Dubois-Dalcq M (2008). "In vitro migration assays of neural stem cells." *Methods Mol Biol* 438: 213-25.
- Eisch AJ and Mandyam CD (2007). "Adult neurogenesis: can analysis of cell cycle proteins move us "Beyond BrdU"?" *Curr Pharm Biotechnol* 8(3): 147-65.
- Ekdahl CT, Claassen JH, Bonde S, Kokaia Z and Lindvall O (2003). "Inflammation is detrimental for neurogenesis in adult brain." *Proc Natl Acad Sci U S A* 100(23): 13632-7.
- Ekdahl CT, Kokaia Z and Lindvall O (2009). "Brain inflammation and adult neurogenesis: the dual role of microglia." *Neuroscience* 158(3): 1021-9.
- Elmariah SB, Oh EJ, Hughes EG and Balice-Gordon RJ (2005). "Astrocytes regulate inhibitory synapse formation via Trk-mediated modulation of postsynaptic GABAA receptors." *J Neurosci* 25(14): 3638-50.
- Eriksson PS, Perfilieva E, Bjork-Eriksson T, Alborn AM, Nordborg C, Peterson DA and Gage FH (1998). "Neurogenesis in the adult human hippocampus." *Nat Med* 4(11): 1313-7.
- Etienne-Manneville S (2006). "In vitro assay of primary astrocyte migration as a tool to study Rho GTPase function in cell polarization." *Methods Enzymol* 406: 565-78.
- Ford AL, Goodsall AL, Hickey WF and Sedgwick JD (1995). "Normal adult ramified microglia separated from other central nervous system macrophages by flow cytometric sorting. Phenotypic differences defined and direct ex vivo antigen presentation to myelin basic protein-reactive CD4⁺ T cells compared." *J Immunol* 154(9): 4309-21.
- Francis F, Koulakoff A, Boucher D, Chafey P, Schaar B, Vinet MC, Friocourt G, McDonnell N, Reiner O, Kahn A, McConnell SK, Berwald-Netter Y, Denoulet P and Chelly J (1999). "Doublecortin is a developmentally regulated, microtubule-associated protein expressed in migrating and differentiating neurons." *Neuron* 23(2): 247-56.
- Frei K, Bodmer S, Schwerdel C and Fontana A (1986). "Astrocyte-derived interleukin 3 as a growth factor for microglia cells and peritoneal macrophages." *J Immunol* 137(11): 3521-7.
- Fritzen S, Schmitt A, Koth K, Sommer C, Lesch KP and Reif A (2007). "Neuronal nitric oxide synthase (NOS-I) knockout increases the survival rate of neural cells in the hippocampus independently of BDNF." *Mol Cell Neurosci* 35(2): 261-71.
- Ganter S, Northoff H, Mannel D and Gebicke-Harter PJ (1992). "Growth control of cultured microglia." *J Neurosci Res* 33(2): 218-30.

- Gebicke-Haerter PJ (2005). "Microarrays and expression profiling in microglia research and in inflammatory brain disorders." *J Neurosci Res* 81(3): 327-41.
- Gebicke-Haerter PJ, Spleiss O, Ren LQ, Li H, Dichmann S, Norgauer J and Boddeke HW (2001). "Microglial chemokines and chemokine receptors." *Prog Brain Res* 132: 525-32.
- Gehrmann J (1996). "Microglia: a sensor to threats in the nervous system?" *Res Virol* 147(2-3): 79-88.
- Gerdes J, Lemke H, Baisch H, Wacker HH, Schwab U and Stein H (1984). "Cell cycle analysis of a cell proliferation-associated human nuclear antigen defined by the monoclonal antibody Ki-67." *J Immunol* 133(4): 1710-5.
- Gheusi G, Cremer H, McLean H, Chazal G, Vincent JD and Lledo PM (2000). "Importance of newly generated neurons in the adult olfactory bulb for odor discrimination." *Proc Natl Acad Sci U S A* 97(4): 1823-8.
- Girolamo F, Strippoli M, Errede M, Benagiano V, Roncali L, Ambrosi G and Virgintino D (2010). "Characterization of oligodendrocyte lineage precursor cells in the mouse cerebral cortex: a confocal microscopy approach to demyelinating diseases." *Ital J Anat Embryol* 115(1-2): 95-102.
- Giulian D and Baker TJ (1986). "Characterization of ameboid microglia isolated from developing mammalian brain." *J Neurosci* 6(8): 2163-78.
- Glanzer JG, Enose Y, Wang T, Kadiu I, Gong N, Rozek W, Liu J, Schlautman JD, Ciborowski PS, Thomas MP and Gendelman HE (2007). "Genomic and proteomic microglial profiling: pathways for neuroprotective inflammatory responses following nerve fragment clearance and activation." *J Neurochem* 102(3): 627-45.
- Golmohammadi MG, Blackmore DG, Large B, Azari H, Esfandiary E, Paxinos G, Franklin KB, Reynolds BA and Rietze RL (2008). "Comparative analysis of the frequency and distribution of stem and progenitor cells in the adult mouse brain." *Stem Cells* 26(4): 979-87.
- Gonzalez-Perez O, Jauregui-Huerta F and Galvez-Contreras AY (2010). "Immune system modulates the function of adult neural stem cells." *Curr Immunol Rev* 6(3): 167-173.
- Gonzalez-Scarano F and Baltuch G (1999). "Microglia as mediators of inflammatory and degenerative diseases." *Annu Rev Neurosci* 22: 219-40.
- Goodfellow CE, Graham SE, Dragunow M and Glass M (2011). "Characterization of NTera2/D1 cells as a model system for the investigation of cannabinoid function in human neurons and astrocytes." *J Neurosci Res*.
- Goshen I, Kreisel T, Ben-Menachem-Zidon O, Licht T, Weidenfeld J, Ben-Hur T and Yirmiya R (2008). "Brain interleukin-1 mediates chronic stress-induced depression in mice via adrenocortical activation and hippocampal neurogenesis suppression." *Mol Psychiatry* 13(7): 717-28.
- Gould E, Beylin A, Tanapat P, Reeves A and Shors TJ (1999a). "Learning enhances adult neurogenesis in the hippocampal formation." *Nat Neurosci* 2(3): 260-5.
- Gould E, Reeves AJ, Graziano MS and Gross CG (1999b). "Neurogenesis in the neocortex of adult primates." *Science* 286(5439): 548-52.
- Gould E, Vail N, Wagers M and Gross CG (2001). "Adult-generated hippocampal and neocortical neurons in macaques have a transient existence." *Proc Natl Acad Sci U S A* 98(19): 10910-7.

- Graeber MB and Streit WJ (2010). "Microglia: biology and pathology." *Acta Neuropathol* 119(1): 89-105.
- Graeber MB, Streit WJ and Kreutzberg GW (1988). "The microglial cytoskeleton: vimentin is localized within activated cells in situ." *J Neurocytol* 17(4): 573-80.
- Gratzner HG (1982). "Monoclonal antibody to 5-bromo- and 5-iododeoxyuridine: A new reagent for detection of DNA replication." *Science* 218(4571): 474-5.
- Green W, Patil P, Marsden CA, Bennett GW and Wigmore PM (2006). "Treatment with olanzapine increases cell proliferation in the subventricular zone and prefrontal cortex." *Brain Res* 1070(1): 242-5.
- Hanisch UK (2002). "Microglia as a source and target of cytokines." *Glia* 40(2): 140-55.
- Hanisch UK and Kettenmann H (2007). "Microglia: active sensor and versatile effector cells in the normal and pathologic brain." *Nat Neurosci* 10(11): 1387-94.
- Hao C, Richardson A and Fedoroff S (1991). "Macrophage-like cells originate from neuroepithelium in culture: characterization and properties of the macrophage-like cells." *Int J Dev Neurosci* 9(1): 1-14.
- Hartley RS, Margulis M, Fishman PS, Lee VM and Tang CM (1999). "Functional synapses are formed between human NTera2 (NT2N, hNT) neurons grown on astrocytes." *J Comp Neurol* 407(1): 1-10.
- Hausler KG, Prinz M, Nolte C, Weber JR, Schumann RR, Kettenmann H and Hanisch UK (2002). "Interferon-gamma differentially modulates the release of cytokines and chemokines in lipopolysaccharide- and pneumococcal cell wall-stimulated mouse microglia and macrophages." *Eur J Neurosci* 16(11): 2113-22.
- Haynes SE, Hollopeter G, Yang G, Kurpius D, Dailey ME, Gan WB and Julius D (2006). "The P2Y₁₂ receptor regulates microglial activation by extracellular nucleotides." *Nat Neurosci* 9(12): 1512-9.
- Heldmann U, Thored P, Claassen JH, Arvidsson A, Kokaia Z and Lindvall O (2005). "TNF- α antibody infusion impairs survival of stroke-generated neuroblasts in adult rat brain." *Exp Neurol* 196(1): 204-8.
- Heyen JR, Ye S, Finck BN and Johnson RW (2000). "Interleukin (IL)-10 inhibits IL-6 production in microglia by preventing activation of NF- κ B." *Brain Res Mol Brain Res* 77(1): 138-47.
- Hickey WF (1999). "Leukocyte traffic in the central nervous system: the participants and their roles." *Semin Immunol* 11(2): 125-37.
- Horvath RJ, Natile-McMenemy N, Alkaitis MS and Deleo JA (2008). "Differential migration, LPS-induced cytokine, chemokine, and NO expression in immortalized BV-2 and HAPI cell lines and primary microglial cultures." *J Neurochem* 107(2): 557-69.
- Imitola J, Raddassi K, Park KI, Mueller FJ, Nieto M, Teng YD, Frenkel D, Li J, Sidman RL, Walsh CA, Snyder EY and Khoury SJ (2004). "Directed migration of neural stem cells to sites of CNS injury by the stromal cell-derived factor 1 α /CXC chemokine receptor 4 pathway." *Proc Natl Acad Sci U S A* 101(52): 18117-22.
- Iosif RE, Ekdahl CT, Ahlenius H, Pronk CJ, Bonde S, Kokaia Z, Jacobsen SE and Lindvall O (2006). "Tumor necrosis factor receptor 1 is a negative regulator of progenitor proliferation in adult hippocampal neurogenesis." *J Neurosci* 26(38): 9703-12.
- Irvin DK, Dhaka A, Hicks C, Weinmaster G and Kornblum HI (2003). "Extrinsic and intrinsic factors governing cell fate in cortical progenitor cultures." *Dev Neurosci* 25(2-4): 162-72.

- Islam O, Gong X, Rose-John S and Heese K (2009). "Interleukin-6 and neural stem cells: more than gliogenesis." *Mol Biol Cell* 20(1): 188-99.
- Ito D, Imai Y, Ohsawa K, Nakajima K, Fukuuchi Y and Kohsaka S (1998). "Microglia-specific localisation of a novel calcium binding protein, Iba1." *Brain Res Mol Brain Res* 57(1): 1-9.
- Johansson S, Price J and Modo M (2008). "Effect of inflammatory cytokines on major histocompatibility complex expression and differentiation of human neural stem/progenitor cells." *Stem Cells* 26(9): 2444-54.
- Kaneko Y, Sakakibara S, Imai T, Suzuki A, Nakamura Y, Sawamoto K, Ogawa Y, Toyama Y, Miyata T and Okano H (2000). "Musashi1: an evolutionally conserved marker for CNS progenitor cells including neural stem cells." *Dev Neurosci* 22(1-2): 139-53.
- Kawabe T, Suganuma M, Ando T, Kimura M, Hori H and Okamoto T (2002). "Cdc25C interacts with PCNA at G2/M transition." *Oncogene* 21(11): 1717-26.
- Keeble JE and Moore PK (2002). "Pharmacology and potential therapeutic applications of nitric oxide-releasing non-steroidal anti-inflammatory and related nitric oxide-donating drugs." *Br J Pharmacol* 137(3): 295-310.
- Kempermann G, Gast D, Kronenberg G, Yamaguchi M and Gage FH (2003). "Early determination and long-term persistence of adult-generated new neurons in the hippocampus of mice." *Development* 130(2): 391-9.
- Kiss JZ and Rougon G (1997). "Cell biology of polysialic acid." *Curr Opin Neurobiol* 7(5): 640-6.
- Koc and Kucukguzel SG (2009). "Medicinal chemistry and anti-inflammatory activity of nitric oxide-releasing NSAID drugs." *Mini Rev Med Chem* 9(5): 611-9.
- Koo JW and Duman RS (2008). "IL-1 β is an essential mediator of the antineurogenic and anhedonic effects of stress." *Proc Natl Acad Sci U S A* 105(2): 751-6.
- Kreutzberg GW (1996). "Microglia: a sensor for pathological events in the CNS." *Trends Neurosci* 19(8): 312-8.
- Kuhn HG, Winkler J, Kempermann G, Thal LJ and Gage FH (1997). "Epidermal growth factor and fibroblast growth factor-2 have different effects on neural progenitors in the adult rat brain." *J Neurosci* 17(15): 5820-9.
- Ladeby R, Wirenfeldt M, Dalmau I, Gregersen R, Garcia-Ovejero D, Babcock A, Owens T and Finsen B (2005). "Proliferating resident microglia express the stem cell antigen CD34 in response to acute neural injury." *Glia* 50(2): 121-31.
- Lambertsen KL, Clausen BH, Babcock AA, Gregersen R, Fenger C, Nielsen HH, Haugaard LS, Wirenfeldt M, Nielsen M, Dagnaes-Hansen F, Bluethmann H, Faergeman NJ, Meldgaard M, Deierborg T and Finsen B (2009). "Microglia protect neurons against ischemia by synthesis of tumor necrosis factor." *J Neurosci* 29(5): 1319-30.
- Lawson LJ, Perry VH, Dri P and Gordon S (1990). "Heterogeneity in the distribution and morphology of microglia in the normal adult mouse brain." *Neuroscience* 39(1): 151-70.
- Lee MK, Rebhun LI and Frankfurter A (1990). "Posttranslational modification of class III beta-tubulin." *Proc Natl Acad Sci U S A* 87(18): 7195-9.
- Levison SW and Goldman JE (1997). "Multipotential and lineage restricted precursors coexist in the mammalian perinatal subventricular zone." *J Neurosci Res* 48(2): 83-94.
- Li T, Jiang L, Chen H and Zhang X (2008). "Characterization of excitability and voltage-gated ion channels of neural progenitor cells in rat hippocampus." *J Mol Neurosci* 35(3): 289-95.

- Li W, Gao G, Guo Q, Jia D, Wang J, Wang X, He S and Liang Q (2009). "Function and phenotype of microglia are determined by toll-like receptor 2/toll-like receptor 4 activation sequence." *DNA Cell Biol* 28(10): 493-9.
- Lim GP, Yang F, Chu T, Chen P, Beech W, Teter B, Tran T, Ubeda O, Ashe KH, Frautschy SA and Cole GM (2000). "Ibuprofen suppresses plaque pathology and inflammation in a mouse model for Alzheimer's disease." *J Neurosci* 20(15): 5709-14.
- Limke TL and Rao MS (2002). "Neural stem cells in aging and disease." *J Cell Mol Med* 6(4): 475-96.
- Ling EA and Wong WC (1993). "The origin and nature of ramified and amoeboid microglia: a historical review and current concepts." *Glia* 7(1): 9-18.
- Liu B and Hong JS (2003). "Role of microglia in inflammation-mediated neurodegenerative diseases: mechanisms and strategies for therapeutic intervention." *J Pharmacol Exp Ther* 304(1): 1-7.
- Liu BF, Gao EJ, Zeng XZ, Ji M, Cai Q, Lu Q, Yang H and Xu QY (2006). "Proliferation of neural precursors in the subventricular zone after chemical lesions of the nigrostriatal pathway in rat brain." *Brain Res* 1106(1): 30-9.
- Liu YP, Lin HI and Tzeng SF (2005). "Tumor necrosis factor- α and interleukin-18 modulate neuronal cell fate in embryonic neural progenitor culture." *Brain Res* 1054(2): 152-8.
- Lockhart BP, Cressey KC and Lepagnol JM (1998). "Suppression of nitric oxide formation by tyrosine kinase inhibitors in murine N9 microglia." *Br J Pharmacol* 123(5): 879-89.
- Lossinsky AS and Shivers RR (2004). "Structural pathways for macromolecular and cellular transport across the blood-brain barrier during inflammatory conditions. Review." *Histol Histopathol* 19(2): 535-64.
- Lucas SM, Rothwell NJ and Gibson RM (2006). "The role of inflammation in CNS injury and disease." *Br J Pharmacol* 147 Suppl 1: S232-40.
- Luo CX, Zhu XJ, Zhou QG, Wang B, Wang W, Cai HH, Sun YJ, Hu M, Jiang J, Hua Y, Han X and Zhu DY (2007). "Reduced neuronal nitric oxide synthase is involved in ischemia-induced hippocampal neurogenesis by up-regulating inducible nitric oxide synthase expression." *J Neurochem* 103(5): 1872-82.
- Luskin MB, Zigova T, Soteres BJ and Stewart RR (1997). "Neuronal progenitor cells derived from the anterior subventricular zone of the neonatal rat forebrain continue to proliferate in vitro and express a neuronal phenotype." *Mol Cell Neurosci* 8(5): 351-66.
- Matarredona ER, Murillo-Carretero M, Moreno-Lopez B and Estrada C (2005). "Role of nitric oxide in subventricular zone neurogenesis." *Brain Res Brain Res Rev* 49(2): 355-66.
- McGeer PL and McGeer EG (1995). "The inflammatory response system of brain: implications for therapy of Alzheimer and other neurodegenerative diseases." *Brain Res Brain Res Rev* 21(2): 195-218.
- Memberg SP and Hall AK (1995). "Dividing neuron precursors express neuron-specific tubulin." *J Neurobiol* 27(1): 26-43.
- Ming GL and Song H (2005). "Adult neurogenesis in the mammalian central nervous system." *Annu Rev Neurosci* 28: 223-50.
- Ming GL and Song H (2011). "Adult neurogenesis in the mammalian brain: significant answers and significant questions." *Neuron* 70(4): 687-702.
- Minghetti L and Levi G (1998). "Microglia as effector cells in brain damage and repair: focus on prostanoids and nitric oxide." *Prog Neurobiol* 54(1): 99-125.

- Monje ML, Toda H and Palmer TD (2003). "Inflammatory blockade restores adult hippocampal neurogenesis." *Science* 302(5651): 1760-5.
- Moran LB, Duke DC and Graeber MB (2007). "The microglial gene regulatory network activated by interferon-gamma." *J Neuroimmunol* 183(1-2): 1-6.
- Moreno-Lopez B, Romero-Grimaldi C, Noval JA, Murillo-Carretero M, Matarredona ER and Estrada C (2004). "Nitric oxide is a physiological inhibitor of neurogenesis in the adult mouse subventricular zone and olfactory bulb." *J Neurosci* 24(1): 85-95.
- Morgan SC, Taylor DL and Pocock JM (2004). "Microglia release activators of neuronal proliferation mediated by activation of mitogen-activated protein kinase, phosphatidylinositol-3-kinase/Akt and delta-Notch signalling cascades." *J Neurochem* 90(1): 89-101.
- Morshead CM, Benveniste P, Iscove NN and van der Kooy D (2002). "Hematopoietic competence is a rare property of neural stem cells that may depend on genetic and epigenetic alterations." *Nat Med* 8(3): 268-73.
- Morshead CM, Craig CG and van der Kooy D (1998). "In vivo clonal analyses reveal the properties of endogenous neural stem cell proliferation in the adult mammalian forebrain." *Development* 125(12): 2251-61.
- Morshead CM, Reynolds BA, Craig CG, McBurney MW, Staines WA, Morassutti D, Weiss S and van der Kooy D (1994). "Neural stem cells in the adult mammalian forebrain: a relatively quiescent subpopulation of subependymal cells." *Neuron* 13(5): 1071-82.
- Mullen RJ, Buck CR and Smith AM (1992). "NeuN, a neuronal specific nuclear protein in vertebrates." *Development* 116(1): 201-11.
- Nakanishi M, Niidome T, Matsuda S, Akaike A, Kihara T and Sugimoto H (2007). "Microglia-derived interleukin-6 and leukaemia inhibitory factor promote astrocytic differentiation of neural stem/progenitor cells." *Eur J Neurosci* 25(3): 649-58.
- Nakatomi H, Kuriu T, Okabe S, Yamamoto S, Hatano O, Kawahara N, Tamura A, Kirino T and Nakafuku M (2002). "Regeneration of hippocampal pyramidal neurons after ischemic brain injury by recruitment of endogenous neural progenitors." *Cell* 110(4): 429-41.
- Napoli C and Ignarro LJ (2003). "Nitric oxide-releasing drugs." *Annu Rev Pharmacol Toxicol* 43: 97-123.
- Nimmerjahn A, Kirchhoff F and Helmchen F (2005). "Resting microglial cells are highly dynamic surveillants of brain parenchyma in vivo." *Science* 308(5726): 1314-8.
- Nixon K and Crews FT (2004). "Temporally specific burst in cell proliferation increases hippocampal neurogenesis in protracted abstinence from alcohol." *J Neurosci* 24(43): 9714-22.
- Nowakowski RS, Lewin SB and Miller MW (1989). "Bromodeoxyuridine immunohistochemical determination of the lengths of the cell cycle and the DNA-synthetic phase for an anatomically defined population." *J Neurocytol* 18(3): 311-8.
- Oehmichen W and Gencic M (1975). "Experimental studies on kinetics and functions of mononuclear phagocytes of the central nervous system." *Acta Neuropathol Suppl* 6: 285-90.
- Okano HJ, Pfaff DW and Gibbs RB (1993). "RB and Cdc2 expression in brain: correlations with 3H-thymidine incorporation and neurogenesis." *J Neurosci* 13(7): 2930-8.


- Ostenfeld T, Joly E, Tai YT, Peters A, Caldwell M, Jauniaux E and Svendsen CN (2002). "Regional specification of rodent and human neurospheres." *Brain Res Dev Brain Res* 134(1-2): 43-55.
- Palmer TD, Schwartz PH, Taupin P, Kaspar B, Stein SA and Gage FH (2001). "Cell culture. Progenitor cells from human brain after death." *Nature* 411(6833): 42-3.
- Park JY, Choi HJ, Prabagar MG, Choi WS, Kim SJ, Cheong C, Park CG, Chin CY and Kang YS (2009). "The C-type lectin CD209b is expressed on microglia and it mediates the uptake of capsular polysaccharides of *Streptococcus pneumoniae*." *Neurosci Lett* 450(3): 246-51.
- Parmar M, Sjoberg A, Bjorklund A and Kokaia Z (2003). "Phenotypic and molecular identity of cells in the adult subventricular zone. in vivo and after expansion in vitro." *Mol Cell Neurosci* 24(3): 741-52.
- Pei Z, Pang H, Qian L, Yang S, Wang T, Zhang W, Wu X, Dallas S, Wilson B, Reece JM, Miller DS, Hong JS and Block ML (2007). "MAC1 mediates LPS-induced production of superoxide by microglia: the role of pattern recognition receptors in dopaminergic neurotoxicity." *Glia* 55(13): 1362-73.
- Petreanu L and Alvarez-Buylla A (2002). "Maturation and death of adult-born olfactory bulb granule neurons: role of olfaction." *J Neurosci* 22(14): 6106-13.
- Pevny L and Placzek M (2005). "SOX genes and neural progenitor identity." *Curr Opin Neurobiol* 15(1): 7-13.
- Polazzi E and Monti B (2010). "Microglia and neuroprotection: from in vitro studies to therapeutic applications." *Prog Neurobiol* 92(3): 293-315.
- Pollard SM, Conti L, Sun Y, Goffredo D and Smith A (2006). "Adherent neural stem (NS) cells from fetal and adult forebrain." *Cereb Cortex* 16 Suppl 1: i112-20.
- Ponomarev ED, Novikova M, Maresz K, Shriver LP and Dittel BN (2005). "Development of a culture system that supports adult microglial cell proliferation and maintenance in the resting state." *J Immunol Methods* 300(1-2): 32-46.
- Qin H, Wilson CA, Lee SJ, Zhao X and Benveniste EN (2005a). "LPS induces CD40 gene expression through the activation of NF-kappaB and STAT-1alpha in macrophages and microglia." *Blood* 106(9): 3114-22.
- Qin L, Li G, Qian X, Liu Y, Wu X, Liu B, Hong JS and Block ML (2005b). "Interactive role of the toll-like receptor 4 and reactive oxygen species in LPS-induced microglia activation." *Glia* 52(1): 78-84.
- Rao MS and Shetty AK (2004). "Efficacy of doublecortin as a marker to analyse the absolute number and dendritic growth of newly generated neurons in the adult dentate gyrus." *Eur J Neurosci* 19(2): 234-46.
- Reynolds BA, Tetzlaff W and Weiss S (1992). "A multipotent EGF-responsive striatal embryonic progenitor cell produces neurons and astrocytes." *J Neurosci* 12(11): 4565-74.
- Reynolds BA and Weiss S (1992). "Generation of neurons and astrocytes from isolated cells of the adult mammalian central nervous system." *Science* 255(5052): 1707-10.
- Reynolds BA and Weiss S (1996). "Clonal and population analyses demonstrate that an EGF-responsive mammalian embryonic CNS precursor is a stem cell." *Dev Biol* 175(1): 1-13.
- Richardson RM, Broadbush WC, Holloway KL and Fillmore HL (2005). "Grafts of adult subependymal zone neuronal progenitor cells rescue hemiparkinsonian behavioral decline." *Brain Res* 1032(1-2): 11-22.

- Roberts ES, Masliah E and Fox HS (2004). "CD163 identifies a unique population of ramified microglia in HIV encephalitis (HIVE)." *J Neuropathol Exp Neurol* 63(12): 1255-64.
- Rocheffort C, Gheusi G, Vincent JD and Lledo PM (2002). "Enriched odor exposure increases the number of newborn neurons in the adult olfactory bulb and improves odor memory." *J Neurosci* 22(7): 2679-89.
- Rolls A, Shechter R, London A, Ziv Y, Ronen A, Levy R and Schwartz M (2007). "Toll-like receptors modulate adult hippocampal neurogenesis." *Nat Cell Biol* 9(9): 1081-8.
- Rosenstiel P, Lucius R, Deuschl G, Sievers J and Wilms H (2001). "From theory to therapy: implications from an in vitro model of ramified microglia." *Microsc Res Tech* 54(1): 18-25.
- Rostovtsev VV, Green LG, Fokin VV and Sharpless KB (2002). "A stepwise Huisgen cycloaddition process: copper(I)-catalyzed regioselective "ligation" of azides and terminal alkynes." *Angew Chem Int Ed Engl* 41(14): 2596-9.
- Sakakibara S, Imai T, Hamaguchi K, Okabe M, Aruga J, Nakajima K, Yasutomi D, Nagata T, Kurihara Y, Uesugi S, Miyata T, Ogawa M, Mikoshiba K and Okano H (1996). "Mouse-Musashi-1, a neural RNA-binding protein highly enriched in the mammalian CNS stem cell." *Dev Biol* 176(2): 230-42.
- Salic A and Mitchison TJ (2008). "A chemical method for fast and sensitive detection of DNA synthesis in vivo." *Proc Natl Acad Sci U S A* 105(7): 2415-20.
- Sanai N, Tramontin AD, Quinones-Hinojosa A, Barbaro NM, Gupta N, Kunwar S, Lawton MT, McDermott MW, Parsa AT, Manuel-Garcia Verdugo J, Berger MS and Alvarez-Buylla A (2004). "Unique astrocyte ribbon in adult human brain contains neural stem cells but lacks chain migration." *Nature* 427(6976): 740-4.
- Sapir T, Horesh D, Caspi M, Atlas R, Burgess HA, Wolf SG, Francis F, Chelly J, Elbaum M, Pietrokovski S and Reiner O (2000). "Doublecortin mutations cluster in evolutionarily conserved functional domains." *Hum Mol Genet* 9(5): 703-12.
- Sasaki A, Yamaguchi H, Horikoshi Y, Tanaka G and Nakazato Y (2004). "Expression of glucose transporter 5 by microglia in human gliomas." *Neuropathol Appl Neurobiol* 30(5): 447-55.
- Saura J, Tusell JM and Serratosa J (2003). "High-yield isolation of murine microglia by mild trypsinization." *Glia* 44(3): 183-9.
- Sawada M, Kondo N, Suzumura A and Marunouchi T (1989). "Production of tumor necrosis factor- α by microglia and astrocytes in culture." *Brain Res* 491(2): 394-7.
- Seki T (2002). "Hippocampal adult neurogenesis occurs in a microenvironment provided by PSA-NCAM-expressing immature neurons." *J Neurosci Res* 69(6): 772-83.
- Seki T and Arai Y (1993). "Highly polysialylated neural cell adhesion molecule (NCAM-H) is expressed by newly generated granule cells in the dentate gyrus of the adult rat." *J Neurosci* 13(6): 2351-8.
- Sheen VL, Arnold MW, Wang Y and Macklis JD (1999). "Neural precursor differentiation following transplantation into neocortex is dependent on intrinsic developmental state and receptor competence." *Exp Neurol* 158(1): 47-62.
- Shingo T, Gregg C, Enwere E, Fujikawa H, Hassam R, Geary C, Cross JC and Weiss S (2003). "Pregnancy-stimulated neurogenesis in the adult female forebrain mediated by prolactin." *Science* 299(5603): 117-20.
- Shors TJ, Miesegaes G, Beylin A, Zhao M, Rydel T and Gould E (2001). "Neurogenesis in the adult is involved in the formation of trace memories." *Nature* 410(6826): 372-6.

- Song H, Stevens CF and Gage FH (2002). "Astroglia induce neurogenesis from adult neural stem cells." *Nature* 417(6884): 39-44.
- Song JH, Wang CX, Song DK, Wang P, Shuaib A and Hao C (2005). "Interferon gamma induces neurite outgrowth by up-regulation of p35 neuron-specific cyclin-dependent kinase 5 activator via activation of ERK1/2 pathway." *J Biol Chem* 280(13): 12896-901.
- Spulber S, Oprica M, Bartfai T, Winblad B and Schultzberg M (2008). "Blunted neurogenesis and gliosis due to transgenic overexpression of human soluble IL-1ra in the mouse." *Eur J Neurosci* 27(3): 549-58.
- Stoll G and Jander S (1999). "The role of microglia and macrophages in the pathophysiology of the CNS." *Prog Neurobiol* 58(3): 233-47.
- Streit WJ, Graeber MB and Kreutzberg GW (1988). "Functional plasticity of microglia: a review." *Glia* 1(5): 301-7.
- Streit WJ, Walter SA and Pennell NA (1999). "Reactive microgliosis." *Prog Neurobiol* 57(6): 563-81.
- Sun L, Lee J and Fine HA (2004). "Neuronally expressed stem cell factor induces neural stem cell migration to areas of brain injury." *J Clin Invest* 113(9): 1364-74.
- Sun Y, Jin K, Childs JT, Xie L, Mao XO and Greenberg DA (2006). "Vascular endothelial growth factor-B (VEGFB) stimulates neurogenesis: evidence from knockout mice and growth factor administration." *Dev Biol* 289(2): 329-35.
- Suslov ON, Kukekov VG, Ignatova TN and Steindler DA (2002). "Neural stem cell heterogeneity demonstrated by molecular phenotyping of clonal neurospheres." *Proc Natl Acad Sci U S A* 99(22): 14506-11.
- Suzumura A, Marunouchi T and Yamamoto H (1991). "Morphological transformation of microglia in vitro." *Brain Res* 545(1-2): 301-6.
- Suzumura A, Sawada M, Yamamoto H and Marunouchi T (1990). "Effects of colony stimulating factors on isolated microglia in vitro." *J Neuroimmunol* 30(2-3): 111-20.
- Svendsen CN (2002). "The amazing astrocyte." *Nature* 417(6884): 29-32.
- Taupin P (2008). "Adult neurogenesis, neuroinflammation and therapeutic potential of adult neural stem cells." *Int J Med Sci* 5(3): 127-32.
- Thored P, Arvidsson A, Cacci E, Ahlenius H, Kallur T, Darsalia V, Ekdahl CT, Kokaia Z and Lindvall O (2006). "Persistent production of neurons from adult brain stem cells during recovery after stroke." *Stem Cells* 24(3): 739-47.
- Thored P, Heldmann U, Gomes-Leal W, Gisler R, Darsalia V, Taneera J, Nygren JM, Jacobsen SE, Ekdahl CT, Kokaia Z and Lindvall O (2009). "Long-term accumulation of microglia with proneurogenic phenotype concomitant with persistent neurogenesis in adult subventricular zone after stroke." *Glia* 57(8): 835-49.
- Vallieres L, Campbell IL, Gage FH and Sawchenko PE (2002). "Reduced hippocampal neurogenesis in adult transgenic mice with chronic astrocytic production of interleukin-6." *J Neurosci* 22(2): 486-92.
- Vazey EM, Chen K, Hughes SM and Connor B (2006). "Transplanted adult neural progenitor cells survive, differentiate and reduce motor function impairment in a rodent model of Huntington's disease." *Exp Neurol* 199(2): 384-96.
- Vernadakis A, Mangoura D, Sakellaridis N and Linderholm S (1984). "Glial cells dissociated from newborn and aged mouse brain." *J Neurosci Res* 11(3): 253-62.

- Vezzani A, Moneta D, Richichi C, Aliprandi M, Burrows SJ, Ravizza T, Perego C and De Simoni MG (2002). "Functional role of inflammatory cytokines and antiinflammatory molecules in seizures and epileptogenesis." *Epilepsia* 43 Suppl 5: 30-5.
- Vukicevic V, Jauch A, Dinger TC, Gebauer L, Hornich V, Bornstein SR, Ehrhart-Bornstein M and Muller AM (2010). "Genetic instability and diminished differentiation capacity in long-term cultured mouse neurosphere cells." *Mech Ageing Dev* 131(2): 124-32.
- Walton NM, Sutter BM, Laywell ED, Levkoff LH, Kearns SM, Marshall GP, 2nd, Scheffler B and Steindler DA (2006). "Microglia instruct subventricular zone neurogenesis." *Glia* 54(8): 815-25.
- Wang Y, Sheen VL and Macklis JD (1998). "Cortical interneurons upregulate neurotrophins in vivo in response to targeted apoptotic degeneration of neighboring pyramidal neurons." *Exp Neurol* 154(2): 389-402.
- Wegiel J, Wisniewski HM, Dziewiatkowski J, Tarnawski M, Kozielski R, Trenkner E and Wiktor-Jedrzejczak W (1998). "Reduced number and altered morphology of microglial cells in colony stimulating factor-1-deficient osteopetrotic op/op mice." *Brain Res* 804(1): 135-9.
- Whitney NP, Eidem TM, Peng H, Huang Y and Zheng JC (2009). "Inflammation mediates varying effects in neurogenesis: relevance to the pathogenesis of brain injury and neurodegenerative disorders." *J Neurochem* 108(6): 1343-59.
- Wollmer MA, Lucius R, Wilms H, Held-Feindt J, Sievers J and Mentlein R (2001). "ATP and adenosine induce ramification of microglia in vitro." *J Neuroimmunol* 115(1-2): 19-27.
- Wong G, Goldshmit Y and Turnley AM (2004). "Interferon-gamma but not TNF alpha promotes neuronal differentiation and neurite outgrowth of murine adult neural stem cells." *Exp Neurol* 187(1): 171-7.
- Wu J, Ohlsson M, Warner EA, Loo KK, Hoang TX, Voskuhl RR and Havton LA (2008). "Glial reactions and degeneration of myelinated processes in spinal cord gray matter in chronic experimental autoimmune encephalomyelitis." *Neuroscience* 156(3): 586-96.
- Xu Y, Tamamaki N, Noda T, Kimura K, Itokazu Y, Matsumoto N, Dezawa M and Ide C (2005). "Neurogenesis in the ependymal layer of the adult rat 3rd ventricle." *Exp Neurol* 192(2): 251-64.
- Yamamoto S, Yamamoto N, Kitamura T, Nakamura K and Nakafuku M (2001). "Proliferation of parenchymal neural progenitors in response to injury in the adult rat spinal cord." *Exp Neurol* 172(1): 115-27.
- Yoshikawa K (2000). "Cell cycle regulators in neural stem cells and postmitotic neurons." *Neurosci Res* 37(1): 1-14.
- Zhang J, Shi XQ, Echeverry S, Mogil JS, De Koninck Y and Rivest S (2007). "Expression of CCR2 in both resident and bone marrow-derived microglia plays a critical role in neuropathic pain." *J Neurosci* 27(45): 12396-406.
- Zhang RL, Zhang L, Zhang ZG, Morris D, Jiang Q, Wang L, Zhang LJ and Chopp M (2003). "Migration and differentiation of adult rat subventricular zone progenitor cells transplanted into the adult rat striatum." *Neuroscience* 116(2): 373-82.

- Zhao M, Momma S, Delfani K, Carlen M, Cassidy RM, Johansson CB, Brismar H, Shupliakov O, Frisen J and Janson AM (2003). "Evidence for neurogenesis in the adult mammalian substantia nigra." *Proc Natl Acad Sci U S A* 100(13): 7925-30.
- Zigova T, Pencea V, Betarbet R, Wiegand SJ, Alexander C, Bakay RA and Luskin MB (1998). "Neuronal progenitor cells of the neonatal subventricular zone differentiate and disperse following transplantation into the adult rat striatum." *Cell Transplant* 7(2): 137-56.



Neural Stem Cells and Therapy

Edited by Dr. Tao Sun

ISBN 978-953-307-958-5

Hard cover, 440 pages

Publisher InTech

Published online 15, February, 2012

Published in print edition February, 2012

This book is a collective work of international experts in the neural stem cell field. The book incorporates the characterization of embryonic and adult neural stem cells in both invertebrates and vertebrates. It highlights the history and the most advanced discoveries in neural stem cells, and summarizes the mechanisms of neural stem cell development. In particular, this book provides strategies and discusses the challenges of utilizing neural stem cells for therapy of neurological disorders and brain and spinal cord injuries. It is suitable for general readers, students, doctors and researchers who are interested in understanding the principles of and new discoveries in neural stem cells and therapy.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Bruno P. Carreira, Maria Inês Morte, Caetana M. Carvalho and Inês M. Araújo (2012). Assessing the Influence of Neuroinflammation on Neurogenesis: In Vitro Models Using Neural Stem Cells and Microglia as Valuable Research Tools, *Neural Stem Cells and Therapy*, Dr. Tao Sun (Ed.), ISBN: 978-953-307-958-5, InTech, Available from: <http://www.intechopen.com/books/neural-stem-cells-and-therapy/assessing-the-influence-of-neuroinflammation-on-neurogenesis-in-vitro-models-using-neural-stem-cells>

INTech
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](https://creativecommons.org/licenses/by/3.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen