We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Inner Blood-Retinal Barrier Transporters: Relevance to Diabetic Retinopathy

Yoshiyuki Kubo and Ken-ichi Hosoya
*University of Toyama
Japan*

1. Introduction

Diabetic retinopathy is a frequent complication of diabetes, and leads to acquired blindness. A variety of gene and molecules have been studied to clarify its pathophysiological mechanism. The retina is involved in the vision process and contains neuronal cells that are the most sensitive to changes in the retinal environment, and the physiological barrier structure is important for maintaining optimal retinal homeostasis. The blood-retinal barrier (BRB) is composed of two cellular barriers, the tight junction of the retinal capillary endothelial cells (inner BRB) and the retinal pigment epithelial cells (outer BRB), restricting nonspecific material transport between the circulating blood and the retina. However, the specific transport of low molecular weight compounds, that is the supply of nutrients and elimination of undesired toxic compounds, is carried out by membrane transporter molecules at the BRB, suggesting that they are closely related to diabetic retinopathy. Since it is also known that two thirds of the human retina is nourished by the inner BRB (Cunha-Vaz, 2004; Hosoya & Tomi, 2005), in this chapter, we will describe the relationship between diabetic retinopathy and the membrane transporter molecules expressed at the inner BRB.

2. Structure and function of the inner BRB

In 1913, Schnaudigel was the first to propose the concept of the BRB. In his experiment, the retina showed similarity to the blood-brain barrier (BBB), that is, the retina was not stained with dye injected intravenously although peripheral tissues were stained (Schnaudigel, 1913). The tight junctions of retinal pigment epithelial (RPE) cells form the outer BRB, and the choriocapillaries are fenestrated while the inner BRB consists of multiple cells, retinal endothelial cells, pericytes and glial cells, and Müller cells are representative retinal glial cells. The inner BRB is formed by tight junctions of the retinal endothelial cells that are covered by pericytes and glial cells (Figure 1). Since the endothelial barrier is formed by a network complex including neurons, Müller cells/astrocytes and endothelial cells which control the function of retinal capillaries, the inner BRB can be thought of as a 'glio-vascular unit' (Kim et al., 2006).

2.1 Cellular interaction

The functional properties of the inner BRB are inducible by paracrine interactions with pericytes and glial cells. For example, in the endothelial cells, several barrier properties and

host-derived angiogenesis are induced by injection of type I astrocytes into the anterior eye chamber of rats (Janzer & Raff, 1987; Janzer, 1997), and the barrier properties of bovine-derived retinal endothelial cells are increased by co-culture with rat brain-derived astrocytes (Gardner et al., 1997). These pieces of evidence suggest that retinal glial cells (Müller cells and astrocytes) acts in a similar manner to astrocytes in the brain, and imply that the barrier function of the inner BRB is modified by several factors secreted from astrocytes. It is known that Müller cells produce several factors to enhance the barrier function of blood vessels in the retina (Tout et al., 1993). In experiments with TR-MUL cells, conditionally immortalized rat Müller cells, TR-MUL cells produce transforming growth factor-beta (TGF- β) to increase the activity of barrier marker proteins expressed in the TR-iBRB cells, conditionally immortalized rat retinal capillary endothelial cells, suggesting that Müller cells contribute to the regulation of barrier function (Abukawa et al., 2009). In addition, it has been reported that there is involvement of glia cell-derived neurotrophic growth factors in the TGF- β family, interleukin-6, and basic fibroblast growth factor (bFGF), in barrier regulation (Abbott, 2002). Pericytes produce angiopoietin-1 to modify the barrier function of endothelial cells (Hori et al., 2004). The gap junction-mediated interaction between pericytes, endothelial cells and contractile cells is involved in the regulation of blood flow (Bandopadhyay et al., 2001). It is also known that pericytes exhibit contraction in the presence of endothelin-1, angiotensin II, ATP and hypoxia, and relaxation in the presence of CO₂, NO, and adenosine (Matsugi et al., 1997a; Matsugi et al., 1997b; Chen & Anderson, 1997).

Fig. 1. Structure of inner Blood-Retinal Barrier (BRB)

2.2 Molecular aspects of the barrier structure

At the inner BRB, the retinal endothelial cells form a tightly sealed monolayer, separating the abluminal (retina side) and luminal (blood side) domains of the retinal endothelium, and prevent paracellular transport of materials across endothelial cells between the retina and

circulating blood (Wolburg et al., 2009). In particular, D-mannitol, a representative non-permeable paracellular marker, exhibits very low blood-to-retina influx permeability while D-glucose and amino acids, the substrates of membrane transporter molecules, exhibit over a 300-fold higher permeability than that of D-mannitol (Puchowicz et al., 2004; Hosoya & Tachikawa, 2009). These pieces of evidence strongly suggest that the inner BRB is a selective barrier for the retina. In order to maintain the tightly sealed monolayer, it is important for the retinal endothelial cells to be connected via a junctional complex including adherens junctions and tight junctions. Tight junctions are formed by signaling, scaffolding and transmembrane proteins, and it is known that the junctional adhesion molecules (JAM), occludin and claudin play a role in the tight junctions (Hirase et al., 1997; Furuse et al., 1998; Bazzoni et al., 2005). The quantitative transcript analysis of rodent retinal endothelial cells shows markedly higher expressions of claudin-5, occludin, and JAM-1 than non-retinal endothelial cells (Tomi & Hosoya, 2004; Tachikawa et al., 2008). ZO-1, ZO-2 and ZO-3, accessory proteins, belong to the zonula occludens family, and are involved in linking the actin cytoskeleton and the cytoplasmic tails of the occludin and claudin complex (Anderson et al., 1995; Haskins et al., 1998). In addition, the seal between the retinal endothelial cells is enhanced by catenins and vascular endothelial cadherin (VE-cadherin) (Bazzoni & Dejana, 2004). An anti-sense nucleotide study has suggested that occludin plays an important role in the functional regulation of the inner BRB since the barrier permeability is increased by a reduction in occludin expression (Kevil et al., 1998). In addition, the expression of occludin is reduced in experimentally conditioned-diabetes, suggesting a change in retinal barrier function in patients with diabetic retinopathy (Antonetti et al., 1998). Other reports have suggested that vascular endothelial growth factor (VEGF) and nitric oxide (NO) have an effect on the increase in retinal barrier permeability. In the presence of vascular endothelial growth factor (VEGF), the cultured endothelial cells exhibit reduced occludin expression and increased barrier permeability across the endothelial cell monolayers (Yaccino et al., 1997), and the NO synthesis or release has been reported to increase the vascular permeability (Mark et al., 2004). Thus, it is thought that VEGF and NO are closely involved in retinal barrier function in diabetic retinopathy since the retina exhibits enhanced production of these factors in hypoxia (Kaur et al., 2006). In addition, hypoxia-ischemia leads to the production of reactive oxygen species (ROS) that cause oxidative stress and affect neovascularization in the diabetic eyes and retinopathy of prematurity (ROP) (Augustin et al., 1993; Saugstad & Rognum, 1988). Therefore, the physiological and pathological roles of VEGF, NO and ROS are important in retinal barrier function, and the suppression of their production or function will help in the clinical treatment of diabetic retinopathy, retinal hypoxia, ischemic central retinal vein occlusion, and other conditions (Kaur et al., 2008).

2.3 Transport system across the barrier

The paracellular impermeability of hydrophilic molecules is governed by the tight junctions in the retinal capillary endothelium. However, it is essential that retinal neural cells, such as photoreceptor cells, are able to take up sources of energy and eliminate undesired materials. Thus, transcellular transport by retinal capillary endothelial cells is needed for a variety of low molecular weight compounds, such as D-glucose, amino acids and their metabolites (Niemeyer, 1997; Tachikawa et al., 2007). Regarding the mechanisms of transcellular

transport, there are three transport systems at the inner BRB, namely, passive diffusion, receptor-mediated transport and carrier-mediated transport (Figure 2). In particular, carrier-mediated transport is the most important for the uptake of essential nutrients and elimination of discarded metabolites, and this can be subdivided into facilitated transport, primary active efflux transport and secondary active influx and efflux transport (Hosoya & Tachikawa, 2009). In general, the membrane transporter is the 12 membrane-spanning membrane protein widely found in a variety of species ranging from bacteria to humans (Kubo et al., 2000; Kubo et al., 2005), and it is protein responsible for carrier-mediated transport. ATP-binding cassette (ABC) transporter and Solute carrier (SLC) transporter are involved in primary active efflux transport and secondary influx and efflux transport, respectively (Figure 3). It has been shown that a variety of influx membrane transporters, such as GLUT1 for D-glucose, are expressed in retinal capillary endothelial cells (Table 1), and they contribute to the retinal uptake of essential nutrients. It is also important to eliminate unwanted metabolites and toxic compounds from the retina. While facilitative and secondary active influx transport systems mediated by influx membrane transporters contribute to the influx of nutrients at the inner BRB, the elimination involves primary active and secondary active efflux transport systems. The efflux transport systems are mediated by ABC transporters, such as MDR1 (P-gp), and SLC transporters, such as OAT3 (Table 1). Research of membrane transporters uses a variety of analytical methods (Kubo et al., 2007). In particular, in the study of the inner BRB, integration plot and retinal uptake index analyses are available to study the *in vivo* blood-to-retina transport (Hosoya & Tomi, 2008), and TR-iBRB cells, the model cell line of retinal capillary endothelial cells, are useful for studying *in vitro* transport mechanisms (Hosoya & Tomi, 2005; Hosoya et al., 2001a).

Fig. 2. Transport Systems in the inner BRB

Fig. 3. Structure of Membrane Transporter

3. Hyperglycemia and glucose transporter (GLUT1)

Hyperglycemia (an increased blood D-glucose concentration) is the most important symptom exhibited by diabetic patients, and this has severe effects on the development of diabetic retinopathy (Cai & Boulton, 2002). At the inner BRB, the retinal endothelial cells express facilitative glucose transporter, GLUT1 that recognizes hexoses and dehydroascorbic acid (DHA) as substrates (Vera et al., 1993). GLUT1 mainly mediates the influx transport of D-glucose across the inner BRB. GLUT1 exhibits an asymmetrical localization at the inner BRB, and the abluminal expression of GLUT1 protein is 2- and 3-times higher than that on the luminal membrane (Takata K et al., 1992; Fernandes et al., 2003), suggesting that GLUT1 suppresses glucose accumulation in the retinal interstitial fluid. Regarding the influx permeability rate, the blood-to-retina transport is 544 and 2440 $\mu\text{L}/(\text{min} \cdot \text{g retina})$ for D-glucose and DHA, respectively (Puchowicz et al., 2004; Hosoya et al., 2004). DHA is the oxidized form of ascorbic acid (vitamin C), one of the representative antioxidants, and rapidly undergoes cellular reduction to ascorbic acid, resulting in the higher permeability rate of DHA (Hosoya et al., 2008b). According to the K_m values of GLUT1 for D-glucose (5 mM) and DHA (93 μM) and the physiological plasma concentration of D-glucose (~ 5 mM) and DHA (~ 10 μM) (Hosoya et al., 2004; Ennis et al., 1982), GLUT1-mediated DHA influx transport across the inner BRB is not inhibited completely under normal and healthy conditions. However, under diabetic conditions, the elevated plasma concentration of D-glucose (>20 mM, hyperglycemia) causes a reduction in GLUT1-mediated DHA transport from the blood to the retina (Minamizono et al., 2006), showing that the retina of diabetic patients is subject to increased oxidative stress.

4. Hyperosmolality mediated by GLUT1

For all cells in the body, it is important to maintain a physiologically optimal osmolality. Sorbitol, a popular sweetener, works as a common organic osmolyte in cells. In the retinal cells, it is known that the cellular polyol pathway is responsible for sorbitol production from D-glucose (Vinores et al., 1993). The rate-limiting enzyme involved in sorbitol production is aldose reductase encoded by the *AR2* gene located on 7q35 which is thought to be a possible susceptible region for diabetic retinopathy (Patel et al., 1996). Under diabetic conditions, hyperglycemia enhances the intracellular accumulation of sorbitol because of the increased GLUT1-mediated facilitative D-glucose transport to the retina and stimulated cellular aldose reductase activity (Iannello et al., 1999). The elevated concentration of sorbitol causes hyperosmolality which stimulates lactate production and intracellular water and reduces the uptake of O₂ (Stevens et al., 1993; Lim et al., 2001). Therefore, GLUT1 is closely involved in the dysfunction and loss of retinal cells, including the capillary endothelial cells in diabetes. Although the change in GLUT1 expression also needs to be considered for a better understanding of the pathological and therapeutic aspects of diabetic retinopathy, both up- and down-regulation of GLUT1 have been reported in the retina with diabetes (Kumagai et al., 1996; Badr et al., 2000), and this remains controversial.

5. Advanced glycation end products (AGEs)

Advanced glycation end products (AGEs) are the result of a chemical chain reaction (non-enzymatic reaction). During normal aging and metabolism, glucose binds to the amino groups of proteins, through the Maillard reaction, Schiff base, and Amadori rearrangement, to produce Amadori products such as glycolhemoglobin (HbA1c) and glycolalbumin that are used to diagnose diabetes. The Amadori products undergo dehydration, hydrolysis and cleavage to form alpha-dicarbonyl compounds, such as glyoxal, methylglyoxal and 3-deoxyglucosone, that have a much greater ability than glucose to accelerate protein glycation. After further reactions, such as oxidation and degradation, irreversible AGEs are produced finally (Brownlee et al., 1988; Takeuchi & Makita Z, 2001).

5.1 AGE effects on the inner BRB

AGEs is the generic term that includes a number of compounds such as pentosidine, pyrraline, crossline, and N (epsilon)-(carboxymethyl) lysine. Interestingly, there are reports of the expression of receptors for AGEs, such as RAGE, on the cellular surface (Schmidt et al., 1992; Neeper et al., 1992). Under diabetic conditions, hyperglycemia promotes the production and accumulation of AGEs, and it is suggested that AGEs are closely related to the pericytes loss in diabetic retinopathy (Brownlee et al., 1988). As described previously, the retinal capillary is composed of endothelial cells, pericytes and glial cells, and it has been reported that the pericytes interacts with the endothelial cells to suppress the undesirable proliferation and prostacyclin production of endothelial cells and to protect these endothelial cells from harmful events (Yamagishi et al., 1993a; Yamagishi et al., 1993b). Therefore, the loss of pericytes, observed during the early stage of diabetic retinopathy, can be an exacerbating factor leading to the induction of neoangiogenesis via VEGF production, thrombus and hypoxia via prostacyclin suppression in the retinal capillary endothelial cells. According to a recent report, the loss of pericytes is caused by AGEs and their receptors that inhibit the proliferation of pericytes and induce their apoptosis (Yamagishi et al., 1995; Yamagishi et al., 2002).

5.2 Taurine homeostasis and TAUT

It has been reported that the reactivity of AGEs can be blocked by the administration of taurine (Nandhini et al., 2004). Taurine is a non-essential amino acid which is thought to have a neuroprotective role as an osmolyte and antioxidant in the retina. In the body, taurine is synthesized from L-cysteine, and cysteine sulfinic acid decarboxylase is the rate-limiting enzyme involved in taurine biosynthesis. Interestingly, it is known that the retina is rich in taurine although the activity of cysteine sulfinic acid decarboxylase is low (Lin et al., 1985). This suggests the physiological importance of blood-to-retina taurine transport across the inner BRB for the maintenance of retina homeostasis. The taurine transport system is mediated by TAUT, which accepts taurine ($K_m = 22.2 \text{ } \mu\text{M}$) for Na^+ - and Cl^- -dependent transport (Tomi et al., 2007). The expression of TAUT has been demonstrated in human primary retinal endothelial cells and TR-iBRB cells. Regarding the influx permeability rate, the blood-to-retinal transport is $259 \text{ } \mu\text{L}/(\text{min} \cdot \text{g retina})$ for taurine (Tomi et al., 2007), and it has been confirmed that the substrates of TAUT have inhibitory effects on retinal taurine uptake (Törnquist et al., 1986). In a study with knockout mice, *taut*^{-/-} mice exhibited an 80 to 90% reduction in taurine levels in the retina when compared with wild-type mice, showing that TAUT is responsible for the retinal homeostasis of taurine (Warskulat et al., 2007). Diabetic patients exhibit taurine deficiency, and a recent report shows that a reduced level of taurine in the retina causes the loss of cone photoreceptor and retinal ganglion cells, suggesting that retinal taurine deficiency is one of the exacerbating factors for diabetic retinopathy (Franconi et al., 1995; Jammoul et al., 2010). Reports have been published describing that taurine administration reduces the severity of the symptoms of diabetes (Barber, 2003; Moloney et al., 2010; Nakamura et al., 1999).

6. Oxidative stress

Oxidative stress is one of the exacerbating factors of diabetic retinopathy. Under normal conditions, it is important to protect the retina from light-induced oxidative stress, and the cellular uptake and synthesis of antioxidants can contribute to prevent the development of diabetic retinopathy. Catalase, superoxide dismutase (SOD) and glutathione peroxidase are representative cellular enzymatic systems that combat oxidative stress (Roginsky et al., 2001; Sozmen et al., 2001; Mates et al., 1999). Under diabetic conditions, down-regulation of SOD and glutathione peroxidase have been reported (Agardh et al., 1998; Agardh et al., 2000; Kern et al., 1994; Kowluru et al., 1997), and ROS are supposed to be generated by the production of AGE signaling via receptors for AGEs, the polyol pathway and enhanced metabolism of eicosanoid (Nourooz-Zadeh & Pereira 2000). Recently, TR-iBRB cells have been reported to show ROS-induced down-regulation of GLUT1 protein expression at the cellular plasma membrane, and proteasome and protein kinase B have been shown to be involved in this mechanism, suggesting that ROS disrupt glucose homeostasis in the retina (Fernandes et al., 2011). Regarding the enzymatic availability of NADPH, glutathione reductase, reducing the oxidized glutathione (GSSG) to glutathione (GSH), competes with aldose reductase in the polyol pathway, suggesting inhibitory effects on the retinal enzymes (Sato et al., 1999; Bravi et al., 1997). In glutathione synthesis, xCT, the membrane transporter expressed in retinal capillary endothelial cells, plays an important role in transporting L-cystine across the inner BRB from the circulating blood. xCT is the representative molecule for the system Xc⁻ and forms a heterodimer with 4F2hc to transport L-cystine and L-glutamate. TR-iBRB cells exhibit Na^+ -independent L-cystine uptake ($K_m=9.2 \text{ } \mu\text{M}$), which is inhibited

by substrates of xCT. xCT is one of the important molecules involved in the biosynthesis of GSH which is a potent endogenous antioxidant. The expression and activity of xCT has been reported to be up-regulated in response to oxidative conditions (Tomi et al., 2002), and it is expected that the expressional and functional alterations of xCT will have an effect on the development of the diabetic retinopathy, regulating the retinal GSH level.

Transporter	Alias	Substrates	Transport Direction	References
SLC2A1	GLUT1	D-Glucose	Influx	Puchowicz et al., 2004
		DHA	Influx	Hosoya et al., 2004
SLC5A6	SMVT	Biotin	Influx	Ohkura et al., 2010
SLC6A6	TAUT	Taurine	Influx	Törnquist et al, 1986
		GABA	Influx	Tomi et al., 2007
SLC6A8	CRT	Creatine	Influx	Nakashima et al., 2004
SLC6A9	GlyT	Glycine	Influx	Okamoto et al., 2009
SLC7A1	CAT1	L-Arginine	Influx	Tomi et al., 2009
SLC7A5	LAT1	L-Leucine	Influx	Törnquist et al, 1986
				Tomi et al., 2005
SLC7A11	xCT	L-Cystine	Influx	Tomi et al., 2002
		L-Glutamate	Influx	Hosoya et al., 2001b
SLC16A1	MCT1	L-Lactate	Influx	Gerhart et al., 1999
				Alm et al., 1985
				Hosoya et al., 2001c
SLC19A1	RFC1	MTF	Influx	Hosoya K et al., 2008a
SLC22A5	OCTN2	L-Carnitine	Influx	Tachikawa et al., 2010
SLC22A8	OAT3	Organic anions	Efflux	Kikuchi et al., 2003
				Somervaille et al., 2003
				Hosoya et al., 2009
SLC29A2	ENT2	Nucleosides	Influx	Nagase et al., 2006
SLC38A2	ATA2	L-Proline	Efflux	Baldwin et al., 2004,
	SNAT2	L-Alanine	Efflux	Yoneyama et al., 2010
				LaNoue et al., 2001
				Levkovitch-Verbin et al., 2002
SLCO1A4	OATP1A4	Organic anions	Efflux	Nakakariya et al., 2008
	oatp2			Katayama et al., 2006
				Noé et al., 1994
				Gao et al., 2002
ABCB1	MDR1	Lipophilic drugs	Efflux	Sugiyama et al., 2001
	P-gp	Organic cations		Hosoya et al., 2001a
				Maines et al., 2005
				Shen et al., 2003
				BenEzra et al., 1990a
				BenEzra et al., 1990b
ABCC4	MRP4	Organic anions	Efflux	Tagami et al., 2009
				Smeets et al., 2004
				Uchida et al., 2007
ABCG2	BCRP	Organic anions	Efflux	Asashima et al., 2006
	MTX			Boulton et al., 2001

DHA dehydroascorbic acid; GABA gamma-aminobutyric acid; MTF methyltetrahydrofolate

Table 1. Membrane Transporters Identified at the inner BRB

7. Conclusion

In this chapter, we have described membrane transporters, such as GLUT1 for D-glucose and DHA, TAUT for taurine, and xCT for L-cystine and L-glutamate, which are mainly involved in the uptake of nutrients across the inner BRB under normal physiological conditions. However, under diabetic conditions, these membrane transporters have accelerating or decelerating roles in retinal capillary endothelial cells, and precise quantification of their expressional alteration in diabetes will provide information about the detailed pathological features of diabetic retinopathy. To date, although it has been shown that a variety of membrane transporters are expressed in retinal capillary endothelial cells (Table 1), there is still insufficient information about them to allow us to have a complete picture of retinal homeostasis, and further studies are needed. Therefore, there is still the possibility that several known membrane transporters play roles in diabetic retinopathy. Regarding the drug treatment of diabetic retinopathy, the membrane transporters are expected to be used in pharmacokinetic predictions and retina-specific drug delivery systems. At the inner BRB, OCTN2 and MCT1 are thought to accept drugs as their substrates (Ohashi et al., 1999; Tamai et al., 1999), and novel drug transport systems have also been suggested (Hosoya et al., 2010). Therefore, retina-specific delivery is a potential for aldose reductase inhibitors, such as sorbinil, ranirestat and epalrestat, that can suppress the cell death of the retinal capillary endothelium (Goldfarb et al., 1991; Narayanan et al., 1993). Furthermore, over 400 identified gene/protein molecules belong to the membrane transporter family, and over 100 molecules are 'orphan transporters' and their expression, localization, function, substrates and roles need to be fully identified. In addition, it is thought that there are also a number of unidentified membrane transporter genes, and new research reports on novel membrane transporters can be seen even now (Kawahara et al., 2009). Therefore, new discoveries and findings will be made as a result of the study of the membrane transporters expressed at the inner BRB, and advances in this field will contribute to our understanding of the pathological and therapeutic aspects of diabetic retinopathy.

8. References

- Abbott, N.J. (2002). Astrocyte-endothelial interactions and blood-brain barrier permeability. *Journal of Anatomy*, Vol.200, pp. 629-638
- Abukawa, H.; Tomi, M.; Kiyokawa, J.; Hori, S.; Kondo, T.; Terasaki, T. & Hosoya, K. (2009). Modulation of retinal capillary endothelial cells by Müller glial cell-derived factors. *Molecular Vision*, Vol.15, pp. 451-457
- Agardh, C.D.; Agardh, E.; Hultberg, B.; Qian, Y. & Ostenson, C.G. (1998). The glutathione levels are reduced in Goto-Kakizaki rat retina, but are not influenced by aminoguanidine treatment. *Current Eye Research*, Vol.17, pp. 251-256
- Agardh, E.; Hultberg, B. & Agardh, C. (2000). Effects of inhibition of glycation and oxidative stress on the development of cataract and retinal vessel abnormalities in diabetic rats. *Current Eye Research*, Vol.21, pp. 543-549
- Alm, A. & Törnquist, P. (1985). Lactate transport through the blood-retinal and the blood-brain barrier in rats. *Ophthalmic Research*, Vol.17, pp. 181-184

- Anderson, J.M.; Fanning, A.S.; Lapierre, L. & Van Itallie C.M. (1995). Zonula occludens (ZO)-1 and ZO-2: membrane-associated guanylate kinase homologues (MAGuKs) of the tight junction. *Biochemical Society Transactions*, Vol.23, pp. 470-475
- Antonetti, D.A.; Barber, A.J.; Khin, S.; Lieth, E.; Tarbell, J.M. & Gardner, T.W. (1998). Vascular permeability in experimental diabetes is associated with reduced endothelial occludin content: vascular endothelial growth factor decreases occludin in retinal endothelial cells. *Diabetes*, Vol.47, pp. 1953-1959
- Asashima, T.; Hori, S.; Ohtsuki, S.; Tachikawa, M.; Watanabe, M.; Mukai, C.; Kitagaki, S.; Miyakoshi, N. & Terasaki, T. (2006). ATP-binding cassette transporter G2 mediates the efflux of phototoxins on the luminal membrane of retinal capillary endothelial cells. *Pharmaceutical Research*, Vol.23, pp. 1235-1242
- Augustin, A.J.; Breipohl, W.; Böker, T.; Lutz, J. & Spitznas, M. (1993). Increased lipid peroxide levels and myeloperoxidase activity in the vitreous of patients suffering from proliferative diabetic retinopathy. *Graefe's Archive for Clinical and Experimental Ophthalmology*, Vol.231, pp. 647-650
- Badr, G.A.; Tang, J.; Ismail-Beigi, F. & Kern, T.S. (2000). Diabetes downregulates GLUT1 expression in the retina and its microvessels but not in the cerebral cortex or its microvessels. *Diabetes*, Vol.49, pp. 1016-1021
- Baldwin, S.A.; Beal, P.R.; Yao, S.Y.; King, A.E.; Cass, C.E. & Young, J.D. (2004). The equilibrative nucleoside transporter family, SLC29. *Pflügers Archiv*, Vol.447, pp. 735-743
- Bandopadhyay, R. ; Orte, C. ; Lawrenson, J.G. ; Reid, A.R. ; De Silva, S. & Allt, G. (2001). Contractile proteins in pericytes at the blood-brain and blood-retinal barriers. *Journal of Neurocytology*, Vol.30, pp. 35-44
- Barber, A. J. (2003). A new view of diabetic retinopathy: a neurodegenerative disease of the eye. *Progress in Neuro-Psychopharmacology and Biological Psychiatry*, Vol.27, pp. 283-290
- Bazzoni, G. & Dejana, E. (2004). Endothelial cell-to-cell junctions: molecular organization and role in vascular homeostasis. *Physiological Reviews*, Vol.84, pp. 869-901
- Bazzoni, G.; Tonetti, P.; Manzi, L.; Cera, M.R.; Balconi, G. & Dejana, E. (2005). Expression of junctional adhesion molecule-A prevents spontaneous and random motility. *Journal of Cell Science*, Vol.118, pp. 623-632
- BenEzra, D. & Maftzir, G. (1990a). Ocular penetration of cyclosporin A. The rabbit eye. *Investigative Ophthalmology and Vision Science*, Vol.31, pp. 1362-1366
- BenEzra, D. & Maftzir, G. (1990b). Ocular penetration of cyclosporine A in the rat eye. *Archives of Ophthalmology*, Vol.108, pp. 584-587
- Boulton, M; Rozanowska, M. & Rozanowski, B. (2001). Retinal photodamage. *Journal of Photochemistry and Photobiology B*, Vol.64, pp. 144-161
- Bravi, M.C.; Pietrangeli, P.; Laurenti, O.; Basili, S.; Cassone-Faldetta, M.; Ferri, C. & De Mattia, G. (1997). Polyol pathway activation and glutathione redox status in non-insulin- dependent diabetic patients. *Metabolism*, Vol.46, pp. 1194-1198
- Brownlee, M.; Cerami, A. & Vlassara, H. (1988). Advanced glycosylation end products in tissue and the biochemical basis of diabetic complications. *New England Journal of Medicine*, Vol.318, pp. 1315-1321
- Cai, J. & Boulton, M. (2002). The pathogenesis of diabetic retinopathy: old concepts and new questions. *Eye*, Vol.16, pp. 242-260

- Chen, Q. & Anderson, D.R. (1997). Effect of CO₂ on intracellular pH and contraction of retinal capillary pericytes. *Investigative Ophthalmology and Vision Science*, Vol.38, pp. 643-651
- Cunha-Vaz, J.G. (2004). The blood-retinal barriers system. Basic concepts and clinical evaluation. *Experimental Eye Research*, Vol.78, pp. 715-721
- Ennis, S.R.; Johnson, J.E. & Pautler, E.L. (1982). In situ kinetics of glucose transport across the blood-retinal barrier in normal rats and rats with streptozocin-induced diabetes. *Investigative Ophthalmology and Vision Science*, Vol.23, pp. 447-456
- Fernandes, R.; Suzuki, K. & Kumagai, A.K. (2003). Inner blood-retinal barrier GLUT1 in long-term diabetic rats: an immunogold electron microscopic study. *Investigative Ophthalmology and Vision Science*, Vol.44, pp. 3150-3154
- Fernandes, R.; Hosoya, K. & Pereira, P. (2011). Reactive oxygen species downregulate glucose transport system in retinal endothelial cells. *American Journal of Physiology-Cell Physiology*, Vol.300, pp. 927-936
- Franconi, F.; Bernardini, F.; Mattana, A.; Miceli, M.; Ciuti, M.; Mian, M.; Gironi, A.; Anichini, R. & Seghieri, G. (1995). Plasma and platelet taurine are reduced in subjects with insulindependent diabetes mellitus: effects of taurine supplementation. *American Journal of Clinical Nutrition*, Vol.61, pp. 1115-1119
- Furuse, M.; Fujita, K.; Hiiragi, T.; Fujimoto, K. & Tsukita, S. (1998). Claudin-1 and -2: novel integral membrane proteins localizing at tight junctions with no sequence similarity to occludin. *Journal of Cell Biology*, Vol.141, pp. 1539-1550
- Gao, B.; Wenzel, A.; Grimm, C.; Vavricka, S.R.; Benke, D.; Meier, P.J. & Remè, C.E. (2002). Localization of organic anion transport protein 2 in the apical region of rat retinal pigment epithelium. *Investigative Ophthalmology and Vision Science*, Vol.43, pp. 510-514
- Gerhart, D.Z.; Leino, R.L. & Drewes, L.R. (1999). Distribution of monocarboxylate transporters MCT1 and MCT2 in rat retina. *Neuroscience*, Vol.92, pp. 367-375
- Gardner, T.W.; Lieth, E.; Khin, S.A.; Barber, A.J.; Bonsall, D.J.; Leshner, T.; Rice, K. & Brennan, W.A. Jr. (1997). Astrocytes increase barrier properties and ZO-1 expression in retinal vascular endothelial cells. *Investigative Ophthalmology and Vision Science*, Vol.38, pp. 2423-2427
- Goldfarb, S.; Ziyadeh, F.N.; Kern, E.F. & Simmons, D.A. (1991). Effects of polyol-pathway inhibition and dietary myo-inositol on glomerular hemodynamic function in experimental diabetes mellitus in rats. *Diabetes*, Vol.40, pp. 465-471
- Haskins, J.; Gu, L.; Wittchen, E.S.; Hibbard, J. & Stevenson, B.R. (1998). ZO-3, a novel member of the MAGUK protein family found at the tight junction, interacts with ZO-1 and occludin. *Journal of Cell Biology*, Vol.141, pp. 199-208
- Hirase, T.; Staddon, J.M.; Saitou, M.; Ando-Akatsuka, Y.; Itoh, M.; Furuse, M.; Fujimoto, K.; Tsukita, S. & Rubin, L.L. (1997). Occludin as a possible determinant of tight junction permeability in endothelial cells. *Journal of Cell Science*, Vol.110, pp. 1603-1613
- Hori, S.; Ohtsuki, S.; Hosoya, K.; Nakashima, E. & Terasaki, T. (2004). A pericyte-derived angiopoietin-1 multimeric complex induces occludin gene expression in brain capillary endothelial cells through Tie-2 activation in vitro. *Journal of Neurochemistry*, Vol.89, pp. 503-513

- Hosoya, K.; Tomi, M.; Ohtsuki, S.; Takanaga, H.; Ueda, M.; Yanai, N.; Obinata, M. & Terasaki, T. (2001a). Conditionally immortalized retinal capillary endothelial cell lines (TR-iBRB) expressing differentiated endothelial cell functions derived from a transgenic rat. *Experimental Eye Research*, Vol.72, pp. 163-172
- Hosoya, K.; Saeki, S. & Terasaki, T. (2001b). Activation of carrier-mediated transport of L-cystine at the blood-brain and blood-retinal barriers in vivo. *Microvascular Research*, Vol.62, pp. 136-142
- Hosoya, K.; Kondo, T.; Tomi, M.; Takanaga, H.; Ohtsuki, S. & Terasaki, T. (2001c). MCT1-mediated transport of L-lactic acid at the inner blood-retinal barrier: a possible route for delivery of monocarboxylic acid drugs to the retina. *Pharmaceutical Research*, Vol.18, pp. 1669-1676
- Hosoya, K.; Minamizono, A.; Katayama, K.; Terasaki, T. & Tomi, M. (2004). Vitamin C transport in oxidized form across the rat blood-retinal barrier. *Investigative Ophthalmology and Vision Science*, Vol.45, pp. 1232-1239
- Hosoya, K. & Tomi, M. (2005). Advances in the cell biology of transport via the inner blood-retinal barrier: establishment of cell lines and transport functions. *Biological Pharmaceutical Bulletin*, Vol.28, pp. 1-8
- Hosoya, K. & Tomi, M. (2008). Inner blood-retinal barrier: transport biology and methodology. In: *Drug Absorption Studies-In Situ, In Vitro and In Silico Models*, Ehrhardt, C. & Kim, K.J. 321-338, AAPS Press-Springer, ISBN978-0-387-74900-6, New York
- Hosoya, K.; Fujita, K. & Tachikawa, M. (2008a). Involvement of reduced folate carrier 1 in the inner blood-retinal barrier transport of methyltetrahydrofolate. *Drug Metabolism and Pharmacokinetics*, Vol.23, pp. 285-292
- Hosoya, K.; Nakamura, G.; Akanuma, S.; Tomi, M. & Tachikawa, M. (2008b). Dehydroascorbic acid uptake and intracellular ascorbic acid accumulation in cultured Müller glial cells (TR-MUL). *Neurochemistry International*, Vol.52, pp. 1351-1357
- Hosoya, K.; Makihara, A.; Tsujikawa, Y.; Yoneyama, D.; Mori, S.; Terasaki, T.; Akanuma, S.; Tomi, M. & Tachikawa, M. (2009). Roles of inner blood-retinal barrier organic anion transporter 3 in the vitreous/retina-to-blood efflux transport of p-aminohippuric acid, benzylpenicillin, and 6-mercaptopurine. *Journal of Pharmacology and Experimental Therapeutics*, Vol.329, pp. 87-93
- Hosoya, K. & Tachikawa, M. (2009). Inner blood-retinal barrier transporters: role of retinal drug delivery. *Pharmaceutical Research*, Vol.26, pp. 2055-2065
- Hosoya, K.; Yamamoto, A.; Akanuma, S. & Tachikawa, M. (2010). Lipophilicity and transporter Influence on blood-retinal barrier permeability: a comparison with blood-brain barrier permeability. *Pharmaceutical Research*, Vol.27, pp. 2715-2724
- Iannello, S.; Cavaleri, A.; Camuto, M. & Belfiore, F. (1999). In vitro inhibition of glucose phosphorylation by an aldosereductase inhibitor (Tolrestat) in some non-insulin sensitive rabbit tissues. *Journal of Diabetes Complications*, Vol.13, pp. 68-73
- Jammoul, F.; Dégardin, J.; Pain, D.; Gondouin, P.; Simonutti, M.; Dubus, E.; Caplette, R.; Fouquet, S.; Craft, C.M.; Sahel, J.A. & Picaud, S. (2010). Taurine deficiency damages photoreceptors and retinal ganglion cells in vigabatrin-treated neonatal rats. *Molecular and Cellular Neuroscience*, Vol.43, pp. 414-421

- Janzer, R.C. & Raff, M.C. (1987). Astrocytes induce blood-brain barrier properties in endothelial cells. *Nature*, Vol.325, pp. 253-257
- Janzer, R.C. (1997). The blood-brain barrier: cellular basis. *Journal of Inherited Metabolic Disease*, Vol.16, pp. 639-647
- Katayama, K.; Ohshima, Y.; Tomi, M. & Hosoya, K. (2006). Application of microdialysis to evaluate the efflux transport of estradiol 17-beta glucuronide across the rat blood-retinal barrier. *Journal of Neuroscience Methods*, Vol.156, pp. 249-256
- Kaur, C.; Sivakumar, V. & Foulds, W.S. (2006). Early response of neurons and glial cells to hypoxia in the retina. *Investigative Ophthalmology and Vision Science*, Vol.47, pp. 1126-1141
- Kaur, C.; Foulds, W.S. & Ling, E.A. (2008). Blood-retinal barrier in hypoxic ischaemic conditions: basic concepts, clinical features and management. *Progress in Retinal and Eye Research*, Vol.27, pp. 622-647
- Kawahara, A.; Nishi, T.; Hisano, Y.; Fukui, H.; Yamaguchi, A. & Mochizuki, N. (2009). The sphingolipid transporter spns2 functions in migration of zebrafish myocardial precursors. *Science*, Vol.323, pp. 524-527
- Kern, T.S.; Kowluru, R.A & Engerman, R.L. (1994). Abnormalities of retinal metabolism in diabetes or galactosemia: ATPases and glutathione. *Investigative Ophthalmology and Visual Science*, Vol.35, pp. 2962-2967
- Kevil, C.G.; Okayama, N.; Trocha, S.D.; Kalogeris, T.J.; Coe, L.L.; Specian, R.D.; Davis, C.P. & Alexander J.S. (1998). Expression of zonula occludens and adherens junctional proteins in human venous and arterial endothelial cells: role of occludin in endothelial solute barriers. *Microcirculation*, Vol.5, pp. 197-210
- Kikuchi, R.; Kusuhara, H.; Sugiyama, D. & Sugiyama, Y. (2003). Contribution of organic anion transporter 3 (Slc22a8) to the elimination of *p*-aminohippuric acid and benzylpenicillin across the blood-brain barrier. *Journal of Pharmacology and Experimental Therapeutics*, Vol.306, pp. 51-58
- Kim, J.H.; Kim, J.H.; Park, J.A.; Lee, S.W.; Kim, W.J.; Yu, Y.S. & Kim, K.W. (2006). Blood-neural barrier: intercellular communication at glio-vascular interface. *Journal of Biochemistry and Molecular Biology*, Vol.39, pp. 339-345
- Kowluru, R.N.; Keern, T.S. & Engerman, R.L. (1997). Abnormalities of retinal metabolism in diabetes or experimental galactosemia. IV. Antioxidant defense system. *Free Radical Biology and Medicine*, Vol.22, pp. 587-592
- Kubo, Y.; Konishi, S.; Kawabe, T; Nada, S. & Yamaguchi, A. (2000). Proximity of periplasmic loops in the metal-Tetracycline/H(+) antiporter of *Escherichia coli* observed on site-directed chemical cross-linking. *Journal of Biological Chemistry*, Vol.275, pp. 5270-5274
- Kubo, Y.; Sekiya, S.; Ohigashi, M.; Takenaka, C.; Tamura, K.; Nada, S.; Nishi, T.; Yamamoto, A. & Yamaguchi, A. (2005). ABCA5 resides in lysosomes, and ABCA5 knockout mice develop lysosomal disease-like symptoms. *Molecular Cellular Biology*, Vol.25, pp. 4138-4149
- Kubo, Y.; Kato, Y. & Tsuji, A. (2007). Experimental Approaches to Study Drug Transporters. In: *Drug Transporters: Molecular Characterization and Role in Drug Disposition* (Wiley Series in Drug Discovery and Development, You, G. & Morris, M.E. 533-556, Wiley, ISBN 978-0-471-78491-3, New Jersey

- Kumagai, A.K.; Vinores, S.A. & Pardridge, W.M. (1996). Pathological upregulation of inner blood-retinal barrier Glut1 glucose transporter expression in diabetes mellitus. *Brain Research*, Vol.706, pp. 313-317
- LaNoue, K.F.; Berkich, D.A.; Conway, M., Barber, A.J.; Hu, L.Y.; Taylor, C. & Hutson, S. (2001). Role of specific aminotransferases in de novo glutamate synthesis and redox shuttling in the retina. *Journal of Neuroscience Research*, Vol.66, pp. 914-922
- Levkovitch-Verbin, H.; Martin, K.R.; Quigley, H.A.; Baumrind, L.A.; Pease, M.E. & Valenta, D. (2002). Measurement of amino acid levels in the vitreous humor of rats after chronic intraocular pressure elevation or optic nerve transection. *Journal of Glaucoma*, Vol.11, pp. 396-405
- Lim, S.S.; Jung, S.H.; Ji, J.; Shin, K.H. & Keum, S.R. (2001). Synthesis of flavenoids and their effects on aldose reductase and sorbitol accumulation in streptozotocin-induced diabetic rat tissues. *Journal of Pharmacy and Pharmacology*, Vol.53, pp. 653-668
- Lin, C.T.; Song, G.X. & Wu, J.Y. (1985). Ultrastructural demonstration of L-glutamate decarboxylase and cysteinesulfinic acid decarboxylase in rat retina by immunocytochemistry. *Brain Research*, Vol.331, pp. 71-80
- Maines, L.W.; Antonetti, D.A.; Wolpert, E.B. & Smith, C.D. (2005). Evaluation of the role of P-glycoprotein in the uptake of paroxetine, clozapine, phenytoin and carbamazepine by bovine retinal endothelial cells. *Neuropharmacology*, Vol.49, pp. 610-617
- Mark, K.S.; Burroughs, A.R.; Brown, R.C.; Huber, J.D. & Davis, T.P. (2004). Nitric oxide mediates hypoxia-induced changes in paracellular permeability of cerebral microvasculature. *American Journal of Physiology Heart and Circulatory Physiology*, Vol.286, pp. 174-180
- Mates, J.M.; Perez-Gomez, C. & Nunez de Castro, I. (1999). Antioxidant enzymes and human diseases. *Clinical Biochemistry*, Vol.32, pp. 595-603
- Matsugi, T.; Chen, Q. & Anderson, D.R. (1997a). Contractile responses of cultured bovine retinal pericytes to angiotensin II. *Archives of Ophthalmology*, Vol.115, pp. 1281-1285
- Matsugi, T.; Chen, Q. & Anderson, D.R. (1997b). Adenosine-induced relaxation of cultured bovine retinal pericytes. *Investigative Ophthalmology and Vision Science*, Vol.38, pp. 2695-2701
- Minamizono, A.; Tomi, M. & Hosoya, K. (2006). Inhibition of dehydroascorbic acid transport across the rat blood-retinal and -brain barriers in experimental diabetes. *Biological Pharmaceutical Bulletin*, Vol.29, pp. 2148-2150
- Moloney, M.A.; Casey, R.G.; O'Donnell, D.H.; Fitzgerald, P.; Thompson, C. & Bouchier-Hayes, D.J. (2010). Two weeks taurine supplementation reverses endothelial dysfunction in young male type 1 diabetics. *Diabetes and Vascular Disease Research*, Vol.7, pp. 300-301
- Nagase, K.; Tomi, M.; Tachikawa, M. & Hosoya, K. (2006). Functional and molecular characterization of adenosine transport at the rat inner blood-retinal barrier. *Biochimica et Biophysica Acta*, Vol.1758, pp. 13-19
- Nakakariya, M.; Shimada, T.; Irokawa, M.; Koibuchi, H.; Iwanaga, T.; Yabuuchi, H.; Maeda, T. & Tamai, I. (2008). Predominant contribution of rat organic anion transporting polypeptide-2 (Oatp2) to hepatic uptake of beta-lactam antibiotics. *Pharmaceutical Research*, Vol.25, pp. 578-585

- Nakamura, T.; Ushiyama, C.; Suzuki, S.; Shimada, N.; Ohmuro, H.; Ebihara, I. & Koide, H. (1999). Effects of taurine and vitamin E on microalbuminuria, plasma metalloproteinase-9, and serum type IV collagen concentrations in patients with diabetic nephropathy. *Nephron*, Vol.83, pp. 361-362
- Nakashima, T.; Tomi, M.; Katayama, K.; Tachikawa, M.; Watanabe, M.; Terasaki, T. & Hosoya, K. (2004). Blood-to-retina transport of creatine via creatine transporter (CRT) at the rat inner blood-retinal barrier. *Journal of Neurochemistry*, Vol.89, pp. 1454-1461
- Nandhini, A.T.; Thirunavukkarasu, V. & Anuradha, C.V. (2004). Stimulation of glucose utilization and inhibition of protein glycation and AGE products by taurine. *Acta Physiologica Scandinavica*, Vol.181, pp. 297-303.
- Narayanan, S. (1993). Aldose reductase and its inhibition in the control of diabetic complications. *Annals of Clinical and Laboratory Science*, Vol.23, pp. 148-158
- Neeper, M.; Schmidt, A.M.; Brett, J.; Yan, S.D.; Wang, F.; Pan, Y.C.; Elliston, K.; Stern, D. & Shaw, A. (1992). Cloning and expression of a cell surface receptor for advanced glycosylation end products of proteins. *Journal of Biological Chemistry*, Vol.267, pp. 14998-15004
- Niemeyer, G. (1997). Glucose concentration and retinal function. *Clinical Neuroscience*, Vol.4, pp. 327-335
- Noé, B.; Hagenbuch, B.; Stieger, B. & Meier, P.J. (1994). Isolation of a multispecific organic anion and cardiac glycoside transporter from rat brain. *PNAS*, Vol.94, pp. 10346-10350.
- Nourooz-Zadeh, J. & Pereira, P. F. (2000). isoprostanes, potential specific markers of oxidative damage in human retina. *Ophthalmic Research*, Vol.32, pp. 133-137
- Ohashi, R.; Tamai, I.; Yabuuchi, H.; Nezu, J.I.; Oku, A.; Sai, Y.; Shimane, M. & Tsuji, A. (1999). Na(+)-dependent carnitine transport by organic cation transporter (OCTN2): its pharmacological and toxicological relevance. *Journal of Pharmacology and Experimental Therapeutics*, Vol.291, pp. 778-784
- Ohkura, Y.; Akanuma, S.I.; Tachikawa, M. & Hosoya, K. (2010). Blood-to-retina transport of biotin via Na(+)-dependent multivitamin transporter (SMVT) at the inner blood-retinal barrier. *Experimental Eye Research*, Vol.91, pp. 387-392
- Okamoto, M.; Akanuma, S.; Tachikawa, M. & Hosoya, K. (2009). Characteristics of glycine transport across the inner blood-retinal barrier. *Neurochemistry International*, Vol.55, pp. 789-795
- Patel, A.; Hibberd, M.L.; Millward, B.A. & Demaine, A.G. (1996). Chromosome 7q35 and susceptibility to diabetic micro-vascular complications. *Journal of Diabetes Complications*, Vol.10, pp. 62-67
- Puchowicz, M.A.; Xu, K.; Magness, D.; Miller, C.; Lust, W.D.; Kern, T.S. & LaManna, J.C. (2004). Comparison of glucose influx and blood flow in retina and brain of diabetic rats. *Journal of Cerebral Blood Flow and Metabolism*, Vol.24, pp. 449-457
- Roginsky, V. & Barsukova, T. (2001). Superoxide dismutase inhibits lipid peroxidation in micelles. *Chemistry and Physics of Lipids*, Vol.111, pp. 87-91
- Sato, S.; Secchi, E.F.; Lizak, M.J.; Fukase, S.; Ohta, N.; Murata, M.; Tsai, J.Y. & Kador, P.F. (1999). Polyol formation and NADPH-dependent reductases in dog retinal capillary pericytes and endothelial cells. *Investigative Ophthalmology and Visual Science*, Vol.40, pp. 697-704

- Saugstad, O.D. & Rognum, T.O. (1988). High postmortem levels of hypoxanthine in the vitreous humor of premature babies with respiratory distress syndrome. *Pediatrics*, Vol.81, pp. 395-398.
- Schmidt, A.M.; Vianna, M.; Gerlach, M.; Brett, J.; Ryan, J.; Kao, J.; Esposito, C.; Hegarty, H.; Hurley, W.; Clauss, M. et al. (1992). Isolation and characterization of two binding proteins for advanced glycosylation end products from bovine lung which are present on the endothelial cell surface. *Journal of Biological Chemistry*, Vol.267, pp. 14987-14997
- Schnaudigel O. (1913). Die vitale farbung mit trypanblau an auge. *Graefe's Archive for Clinical and Experimental Ophthalmology*, Vol.86, pp. 93-105
- Shen, J.; Cross, S.T.; Tang-Liu, D.D. & Welty, D.F. (2003). Evaluation of an immortalized retinal endothelial cell line as an in vitro model for drug transport studies across the blood-retinal barrier. *Pharmaceutical Research*, Vol.20, pp. 1357-1363
- Smeets, P.H.; van Aubel, R.A.; Wouterse, A.C.; van den Heuvel, J.J. & Russel, F.G. (2004). Contribution of multidrug resistance protein 2 (MRP2/ABCC2) to the renal excretion of p-aminohippurate (PAH) and identification of MRP4 (ABCC4) as a novel PAH transporter. *Journal of the American Society of Nephrology*, Vol.15, pp. 2828-2835
- Somervaille, T.C.; Hann, I.M.; Harrison, G.; Eden, T.O.; Gibson, B.E.; Hill, F.G.; Mitchell, C.; Kinsey, S.E.; Vora, A.J. & Lilleyman, J.S. (MRC Childhood Leukaemia Working Party). (2003). Intraocular relapse of childhood acute lymphoblastic leukaemia. *British Journal of Haematology*, Vol.121, pp. 280-288
- Sozmen, E.Y.; Sozmen, B.; Delen, Y. & Onat, T. (2001). Catalase/superoxide dismutase (sod) and catalase/paraonase (pon) ratios may implicate poor glycemic control. *Archives of Medical Research*, Vol.32, pp. 283-287
- Stevens, M.J.; Henry, D.N.; Thomas, T.P.; Killen, P.D. & Greene, D.A. (1993). Aldose reductase gene expression and osmotic dysregulation in cultured human retinal pigment epithelial cells. *American Journal of Physiology*, Vol.265, pp. E428-E438
- Sugiyama, D.; Kusuhara, H.; Shitara, Y.; Abe, T.; Meier, P.J.; Sekine, T.; Endou, H.; Suzuki, H. & Sugiyama Y. (2001). Characterization of the efflux transport of 17beta-estradiol-D-17beta-glucuronide from the brain across the blood-brain barrier. *Journal of Pharmacology and Experimental Therapeutics*, Vol.298, pp. 316-322
- Tachikawa, M.; Hosoya, K.; Ohtsuki, S. & Terasaki, T. (2007). A novel relationship between creatine transport at the blood-brain and blood-retinal barriers, creatine biosynthesis, and its use for brain and retinal energy homeostasis. *Subcellular Biochemistry*, Vol.46, pp. 83-98
- Tachikawa, M.; Toki, H.; Tomi, M. & Hosoya, K. (2008). Gene expression profiles of ATP-binding cassette transporter A and C subfamilies in mouse retinal vascular endothelial cells. *Microvascular Research*, Vol.75, pp. 68-72
- Tachikawa, M.; Takeda, Y.; Tomi, M. & Hosoya, K. (2010). Involvement of OCTN2 in the transport of acetyl-L-carnitine across the inner blood-retinal barrier. *Investigative Ophthalmology and Vision Science*, Vol.51, pp. 430-436
- Tagami, M.; Kusuhara, S.; Honda, S.; Tsukahara, Y. & Negi, A. (2009). Expression of ATP-binding cassette transporters at the inner blood-retinal barrier in a neonatal mouse model of oxygen-induced retinopathy. *Brain Research*, Vol.1283, pp. 186-193

- Takata, K.; Kasahara, T.; Kasahara, M.; Ezaki, O. & Hirano, H. (1992). Ultracytochemical localization of the erythrocyte/HepG2-type glucose transporter (GLUT1) in cells of the blood-retinal barrier in the rat. *Investigative Ophthalmology and Vision Science*, Vol.33, pp. 377-383
- Takeuchi, M. & Makita, Z. (2001). Alternative routes for the formation of immunochemically distinct advanced glycation end-products in vivo. *Current Molecular Medicine*, Vol.1, pp. 305-315
- Tamai, I.; Sai, Y.; Ono, A.; Kido, Y.; Yabuuchi, H.; Takanaga, H.; Satoh, E.; Ogihara, T.; Amano, O.; Izeki, S. & Tsuji, A. (1999). Immunohistochemical and functional characterization of pH-dependent intestinal absorption of weak organic acids by the monocarboxylic acid transporter MCT1. *Journal of Pharmacy and Pharmacology*, Vol.51, pp. 1113-1121
- Tomi, M.; Hosoya, K.; Takanaga, H.; Ohtsuki, S. & Terasaki, T. (2002). Induction of xCT gene expression and L-cystine transport activity by diethyl maleate at the inner blood-retinal barrier. *Investigative Ophthalmology and Vision Science*, Vol.43, pp. 774-779
- Tomi, M. & Hosoya, K. (2004). Application of magnetically isolated rat retinal vascular endothelial cells for the determination of transporter gene expression levels at the inner blood-retinal barrier. *Journal of Neurochemistry*, Vol. 91, pp. 1244-1248
- Tomi, M.; Mori, M.; Tachikawa, M.; Katayama, K.; Terasaki, T. & Hosoya, K. (2005). L-type amino acid transporter 1-mediated L-leucine transport at the inner blood-retinal barrier. *Investigative Ophthalmology and Vision Science*, Vol.46, pp. 2522-2530
- Tomi, M.; Terayama, T.; Isobe, T.; Egami, F.; Morito, A.; Kurachi, M.; Ohtsuki, S.; Kang, Y.S.; Terasaki, T. & Hosoya, K. (2007). Function and regulation of taurine transport at the inner blood-retinal barrier. *Microvascular Research*, Vol.73, pp. 100-106
- Tomi, M.; Kitade, N.; Hirose, S.; Yokota, N.; Akanuma, S.; Tachikawa, M. & Hosoya, K. (2009). Cationic amino acid transporter 1-mediated L-arginine transport at the inner blood-retinal barrier. *Journal of Neurochemistry*, Vol.111, pp. 716-725
- Törnquist, P. & Alm, A. (1986). Carrier-mediated transport of amino acids through the blood-retinal and the blood-brain barriers. *Graefe's Archive for Clinical and Experimental Ophthalmology*, Vol.224, pp. 21-25
- Tout, S.; Chan-Ling, T.; Holländer, H. & Stone, J. (1993). The role of Müller cells in the formation of the blood-retinal barrier. *Neuroscience*, Vol.55, pp. 291-301.
- Uchida, Y.; Kamiie, J.; Ohtsuki, S. & Terasaki, T. (2007). Multichannel liquid chromatography-tandem mass spectrometry cocktail method for comprehensive substrate characterization of multidrug resistance-associated protein 4 transporter. *Pharmaceutical Research*, Vol.24, pp. 2281-2296
- Vera, J.C.; Rivas, C.I.; Fischbarg, J. & Golde, D.W. (1993). Mammalian facilitative hexose transporters mediate the transport of dehydroascorbic acid. *Nature*, Vol.364, pp. 79-82
- Vinores, S.A.; Van Niel, E.; Swerdloff, J.L. & Campochiaro, P.A. (1993). Electron microscopic immunocytochemical demonstration of blood-retinal barrier breakdown in human diabetics and its association with aldose reductase in retinal vascular endothelium and retinal pigment epithelium. *Histochemical Journal*, Vol.25, pp. 648-663
- Warskulat, U.; Heller-Stilb, B.; Oermann, E.; Zilles, K.; Haas, H.; Lang, F. & Häussinger, D. (2007). Phenotype of the taurine transporter knockout mouse. *Methods in Enzymology*, Vol.428, pp. 439-458

- Wolburg, H.; Noell, S.; Mack, A.; Wolburg-Buchholz, K. & Fallier-Becker, P. (2009). Brain endothelial cells and the glio-vascular complex. *Cell and Tissue Research*, Vol.335, pp. 75-96
- Yaccino, J.A.; Chang, Y.S.; Hollis, T.M.; Gardner, T.W. & Tarbell J.M. (1997). Physiological transport properties of cultured retinal microvascular endothelial cell monolayers. *Current Eye Research*, Vol.16, pp. 761-768
- Yamagishi, S.; Kobayashi, K. & Yamamoto, H. (1993a). Vascular pericytes not only regulate growth, but also preserve prostacyclin-producing ability and protect against lipid peroxide-induced injury of co-cultured endothelial cells. *Biochemical and Biophysical Research Communications*, Vol.190, pp. 418-425
- Yamagishi, S.; Hsu, C.C.; Kobayashi, K. & Yamamoto, H. (1993b). Endothelin 1 mediates endothelial cell-dependent proliferation of vascular pericytes. *Biochemical and Biophysical Research Communications*, Vol.191, pp. 840-846
- Yamagishi, S.; Hsu, C.C.; Taniguchi, M.; Harada, S.; Yamamoto, Y.; Ohsawa, K.; Kobayashi, K. & Yamamoto, H. (1995). Receptor-mediated toxicity to pericytes of advanced glycosylation end products: a possible mechanism of pericyte loss in diabetic microangiopathy. *Biochemical and Biophysical Research Communications*, Vol.213, pp. 681-687
- Yamagishi, S.; Amano, S.; Inagaki, Y.; Okamoto, T.; Koga, K.; Sasaki, N.; Yamamoto, H.; Takeuchi, M. & Makita, Z. (2002). Advanced glycation end products-induced apoptosis and overexpression of vascular endothelial growth factor in bovine retinal pericytes. *Biochemical and Biophysical Research Communications*, Vol.290, pp. 973-978
- Yoneyama, D.; Shinozaki, Y.; Lu, W.L.; Tomi, M.; Tachikawa, M. & Hosoya, K. (2010). Involvement of system A in the retina-to-blood transport of L-proline across the inner blood-retinal barrier. *Experimental Eye Research*, Vol.90, pp. 507-513

IntechOpen

Diabetic Retinopathy

Edited by Dr. Mohammad Shamsul Ola

ISBN 978-953-51-0044-7

Hard cover, 356 pages

Publisher InTech

Published online 24, February, 2012

Published in print edition February, 2012

The aim of this book is to provide a comprehensive overview of current concepts in pathogenesis, diagnosis and treatments of diabetic retinopathy. It provides a collection of topics written by excellent authors, covering discussions on advances in understanding of pathophysiology, immunological factors and emerging concepts, relating to clinical aspects and treatment strategies. The contents of the book will not only provide a resource for our knowledge but also improve diagnosis and treatment options for those patients who suffer vision loss due to diabetic retinopathy.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Yoshiyuki Kubo and Ken-ichi Hosoya (2012). Inner Blood-Retinal Barrier Transporters: Relevance to Diabetic Retinopathy, *Diabetic Retinopathy*, Dr. Mohammad Shamsul Ola (Ed.), ISBN: 978-953-51-0044-7, InTech, Available from: <http://www.intechopen.com/books/diabetic-retinopathy/inner-blood-retinal-barrier-transporters-relevance-to-diabetic-retinopathy>

INTech
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](https://creativecommons.org/licenses/by/3.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen