

# We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

186,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index  
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?  
Contact [book.department@intechopen.com](mailto:book.department@intechopen.com)

Numbers displayed above are based on latest data collected.  
For more information visit [www.intechopen.com](http://www.intechopen.com)



# Molecular Biodiversity Inventory of the Ichthyofauna of the Czech Republic

Jan Mendel et al.\*

*Institute of Vertebrate Biology  
Academy of Sciences of the Czech Republic, Brno  
Czech Republic*

## 1. Introduction

The term biological diversity (biodiversity) covers the variability of life on Earth. In 1989, the World Wildlife Fund defined biodiversity as “the richness of life on Earth – millions of plants, animals and microorganisms, including the genes which they carry, and complex ecosystems that create the environment” (Primack et al., 2001).

The degree of biodiversity was dynamically changing during the centuries and millennia, according to available resources it was gradually growing. In the course of evolution the periods of intensive speciation and relative “speciation rest” were alternating, with five major episodes of mass extinction (Wilson, 1989; Raup, 1992). The worst extinction event took place at the end of the Permian period, 250 million years ago (Primack et al., 2001), the most recent and also the best known mass extinction event occurred in the late Cretaceous period, i. e. 65 million years ago (Freeland, 2005). From then on, the rate of speciation was in equilibrium with the rate of extinction, or was even higher. Currently, the rate of extinction is 100-1000 times higher, namely almost exclusively in consequence of human activity. Therefore a lot of experts call the present situation the sixth mass extinction (Primack et al., 2001). At present, only approximately 1.5 million species are described, of which the majority is represented by insects and plants (Wilson, 1992). According to various estimates the number of non-described species amounts to 10 million or even 30-150 million (Hammond, 1992).

In recent years, when the effort to map and at the same time to protect global biodiversity has been intensifying, it is becoming clear that the existing morphological approach to species classification is not sufficient. Therefore, interest has focused on the methods of molecular genetics.

In 2003, a global project aimed at the mapping and protection of global biodiversity using a new taxonomic method called “DNA barcoding” was started. The key personality in this

---

\* Eva Marešová<sup>1</sup>, Ivo Papoušek<sup>1</sup>, Karel Halačka<sup>1</sup>, Lukáš Vetešník<sup>1</sup>, Radek Šanda<sup>2</sup>, Milena Koníčková<sup>1</sup> and Soňa Urbánková<sup>1</sup>

<sup>1</sup>*Institute of Vertebrate Biology, Academy of Sciences of the Czech Republic, v.v.i. Brno*

<sup>2</sup>*Department of Zoology, National Museum, Prague  
Czech Republic*

initiative was Canadian professor Paul D. N. Hebert from the Biodiversity Institute of Ontario (BIO), who proposed the creation of a global public library of DNA barcodes for all living organisms, also known as BoLD (Barcode of Life Database). The Barcode of Life Data Systems is an informatics workbench for the acquisition, storage, intercontinental comparison and publication of DNA barcode records (Fig. 1).




Fig. 1. Multifunctionality of the BoLD database (adopted from <http://ibol.org/resources/barcode-library/>)

At the same time it bridges a long-standing bioinformatics chasm between molecular, morphological and distributional data. BoLD is freely available to anyone interested in these problems, the identification and preservation of all life on Earth.

## 2. DNA barcoding

The basic assumption behind this method is that every biological species has a short sequence in its genome which is unique to that species, like a fingerprint to every human being. Its mutation rate should be sufficiently fast to enable the divergence of the sequences of closely related species and at the same time slow enough to enable the minimization of the differences between members of the same species. The implementation of a comparative sequence analysis of this carefully selected sequence should enable mutual identification of the individual species.

The term DNA barcoding is based on the analogy with EAN barcodes on goods which enable safe identification of the individual products (Hebert et al., 2003a). EAN barcode (Fig. 2) is composed of 10 digits at 11 positions, which gives  $10^{11}$  possible combinations. The genetic code has only 4 letters (DNA bases) for one position, but the length of that code is incomparably longer. Only with the sequence length of 15 nucleotides, theoretically  $4^{15}$  various combinations are available, which is a number of unique combinations far exceeding even the boldest estimates of the number of biological species living on Earth (Hebert et al., 2003a).

The technique of DNA barcoding provides a standardized method based on the mapping of a single gene in all the species on Earth (Hebert et al., 2003a). At present, the fragment of the gene for the first cytochrome c oxidase (COI) subunit appears to be the major marker for species of the animal kingdom. This segment is approximately 650 bp long and the protein it codes is part of the respiration chain. The results suggest that this global standard can also be easily obtained from phylogenetically very distant taxa by using a relatively small set of primers and that it is effective in the differentiation of closely related animal species belonging to a great number of groups of invertebrates and vertebrates (Hajibabaei et al., 2007; Hebert et al., 2003a, 2003 ; Hebert & Barrett, 2005; Ivanova et al., 2007).



Fig. 2. EAN barcode (adopted from Brian Krueger).

Hebert et al. (2003b) proposed a limit of a 2% deviation in the barcode sequence for the identification of intraspecific variability. The intraspecific deviation only rarely exceeds 2% and in most cases does not reach 1% (Avice, 2000). They also proposed an experiential rule that the divergence between the species sequences should be 10 times higher than that within one species (Hebert et al., 2004).

For the identification of samples with degraded DNA, Hajibabaei et al. (2006) and Meusnier et al. (2008) proposed using shorter COI segments, the so-called “universal mini-barcodes” of a length smaller than 150 bp. Thus they provided a reliable solution for the identification of older tissues, badly preserved museum specimens, partly digested pieces of food in animal stomachs, etc. The results indicate that although the selection of a concrete position of the mini-barcode plays an important role concerning the success of identification, generally it can be stated that the mini-barcodes analysis provides virtually the same results as the specific full-barcode analysis.

## 2.1 Global progression of DNA barcoding

In April 2004, the Consortium for Barcode of Life (CBOL) was established (Schindel & Miller, 2005). It is an international collaborative effort of more than 160 member organizations from more than 50 countries on six continents (<http://barcoding.si.edu>, Hebert et al., 2004). CBOL’s mission is to develop the potential of DNA barcoding as a practical and financially affordable tool for taxonomic research, the study and preservation of biodiversity and the development of applications which will utilize the taxonomic information for the benefit of science and society.

Since 2004, a number of campaigns and projects striving to map global biodiversity of all life on Earth have been started (<http://ibol.org/about-us/campaigns/>):

- **Formicidae Barcode of Life** - a campaign aimed at barcoding all of the world’s more than 12,000 ant species.
- **Bee Barcode of Life Initiative (Bee-BOL)** - a global effort to assembly the barcodes of all 20,000 bee species.

- **All Birds Barcoding Initiative (ABBI)** – the aim is to collect DNA barcodes of all of the approximately 10,000 known bird species. The genetic analyses made within the project show that there are hundreds of bird species which have not been described yet.
- **Trichoptera Barcode of Life** – the aim of the project is to barcode the world's approximately 13,000 species of caddisflies.
- **Coral Reef Barcode of Life** – a detailed study of fishes living at one site of the Great Barrier Reef.
- **Fish Barcode of Life Initiative (FISH-BOL)** – a project aimed at the mapping of global ichthyofauna involving 31,200 known fish species (Ward et al., 2009). This initiative also includes the Czech project „Molecular biodiversity inventory of the ichthyofauna of the Czech Republic“ which has been introduced here.
- **All Fungi Barcoding** – a project associating initiatives mapping the global diversity of fungi.
- **HealthBOL** – an initiative coordinating the barcoding of vectors, pathogens, and parasites for the improvement of human health around the world.
- **Lepidoptera Barcode of Life** – a campaign mapping all butterflies, with an associated sub-campaign for selected families of Australia and North America.
- **Mammal Barcode of Life** – a project studying global mammal fauna is part of the initiative encompassing all vertebrates.
- **Mosquito Barcoding Initiative** – an international effort aimed at the identification of approximately 3,200 known mosquito species, the disease-bearing species being the priority.
- **Marine Barcode of Life campaign (MarBOL)** – a joint project of the Consortium for the Barcode of Life (CBOL) and the Census of Marine Life (CoML) is aimed at the comprehensive mapping of the diversity of the world's oceans.
- **Polar Barcode of Life campaign** – a campaign studying the biodiversity of the Arctic and Antarctic. It includes marine, freshwater and terrestrial ecosystems.
- **Shark Barcode of Life** – a project aimed at the barcoding of sharks. It plans for 1,000 marine and 100 freshwater shark species.
- **Sponge Barcoding Project** – it is the first global project studying diploblastic species using DNA barcoding. It covers the complete taxonomic range of Porifera.

The existing projects proceed successfully and have already brought the first results regarding various groups of plants and animals, e. g., birds (Hebert et al., 2004), gorillas (Thalmann et al., 2004), tropical beetles (Monaghan et al., 2005), spiders (Barrett & Hebert., 2005), ants (Fisher, 2006), flowering plants (Kress et al., 2005), amphibians (Vences et al., 2005), fishes (Ward et al., 2005), etc.

Currently, as many as 1,371,809 DNA barcodes are identified and stored, which corresponds to approximately 113,435 denominated species (as of September 30, 2011). The collaborating scientific teams have set a goal by 2015 to collect 5 million barcodes/specimens for 500,000 species.

### 2.1.1 Fish DNA barcoding

In 2005, the FISH-BOL initiative was established. It is an expression of the global effort to develop and coordinate a standardised library of sequences for all fish species. For this



purpose a guideline containing DNA barcodes, photographs of the studied specimens, geospatial coordinates of the locations of the finds, etc. was issued. It also contains references to type specimens, information on species distribution, nomenclature, author taxonomic information, literature citations, etc. Thus, it complements and enhances the information resources available so far – the genome databases (GenBank, EMBL, DDBJ), including the internet encyclopaedia FishBase.

Barcoding in fish has multiple usages, of which we select only some:

- Identifying endangered and protected species (Ward et al., 2008)
- Identifying historical and museum material (Meusnier et al., 2008)
- Identifying new and cryptic species and possible fusions of existing taxa, and insight into phylogenetic relationships (Pyle et al., 2008)
- The development of a reference library for known species (Ratnasingham & Hebert, 2007)

Fishes are the most diversified group of vertebrates; at present, there are about 30,000 known species, including 15,758 (53.3%) marine species, 13,779 (46.4%) freshwater species and 86 (0.3%) brackish species (Ward et al., 2005). By September 2011, DNA barcodes were obtained from 8,293 species (27%) in the world, and from 503 species (25%) in Europe (<http://www.fishbol.org/progress.php>).

Some of the best-known projects utilizing the DNA barcoding in fishes are:

1. “Barcoding Marine Fishes: A Three-Ocean Perspective Project”. The goal of this project is to obtain DNA barcodes of all marine species in the Pacific, Indian and Atlantic oceans. Participating countries include Canada, Australia, South Africa and Portugal.
2. “Planetary Biodiversity Inventories – Catfishes”. It incorporates 300 participants from 40 countries whose work is taking an inventory of catfishes (Siluriformes).
3. “The Cypriniformes Tree of Life Initiative” (CToL), an American initiative aimed at the collection and analysis of DNA samples and reference specimens of nearly all North American freshwater species (~ 1100 species).

### **2.1.2 The state of the molecular biodiversity inventory of the ichthyofauna of the Czech Republic and introduction of the project**

The study and recognition of genetic diversity in the Czech Republic has an approximately twenty-year history. Although the methods of biochemical genetics which were used in the past brought significant knowledge, only the application of molecular-genetic methods allowed significant detailed recognition and disclosure of the genetic diversity of species at the level of populations (Lusk & Hanel, 2008). Existing information about the genetic diversity of the individual species of the native ichthyofauna of the Czech Republic was and still is insufficient. The result of the research project VaV – SM/6/3/05 “Genetic diversity of endangered fish species - the essential basis for effective biodiversity protection” brought a breakthrough in the recognition of genetic diversity, especially of the protected and endangered species. The project was conducted in 2005-2007 in cooperation between our department and the research workers from the Institute of Animal Physiology and Genetics, v.v.i., Liběchov. This allowed the presentation of the first taxonomic approaches to some species of the Czech ichthyofauna (Bartoňová et al., 2008; Mendel et al., 2005; Mendel et al.,

2008a; Papoušek et al., 2008a, 2008b; Vetešník et al., 2007). Other taxonomic studies of our team are included in the proceedings "Biodiversity of fishes in the Czech Republic VII", which paid attention to the problems of the genetic diversity of some protected and rare fish species of the Czech Republic. Apart from the study of actual intra-species diversity, attention in this book was also paid to the distribution or specification of the occurrence of the studied species (Bartoňová et al., 2008; Lusk et al., 2008a, 2008b; Lusková et al., 2008a, 2008b; Mendel et al., 2008b, 2008c; Papoušek et al., 2008b).

The main goal of the presented project is the mapping and inventory making of all Czech ichthyofauna using the DNA barcoding technique. In addition to that, in the case of archived specimens forming new museum type series a comprehensive approach was used (morphology and nuclear genome analysis).

A study of this type has been lacking in the Czech Republic. It is unique due to its comprehensiveness using both the classical and the newest taxonomic tools and due to its future potential – the intercontinental mapping of the diversity of life on Earth. Thanks to our cooperation with the Biodiversity Institute of Ontario the Czech Republic became actively involved in the international iBOL project (International Barcode of Life Project), namely in its part which was already mentioned above – FISH-BOL (Fish Barcode of Life Initiative). Thus, the Czech study took the side of already finished or currently running studies which take place within whole countries and continents, e. g.:

- Fishes of Australia (Ward et al., 2005)
- Barcoding of Canadian freshwater fishes (Hubert et al., 2008)
- DNA barcoding of fish of the Antarctic Scotia Sea (Rock et al., 2008)
- Freshwater fishes from Mexico and Guatemala (Valdez-Moreno et al., 2009)
- Aquarium Imports (Steinke et al., 2009)
- Fishes of Alaska and the Pacific Arctic (Mecklenburg et al., 2011)
- Amazon fishes (Ardura et al., 2010)
- DNA Barcoding of Indian Marine Fish (Lakra et al., 2011)
- Fishes of Japan (Zhang & Hanner 2011)
- Freshwater Fishes of North America (April et al., 2011)
- Barcoding Fishes of Eastern Nigeria (Nwani et al., 2011)
- Fishes of Argentina (running project)
- Fishes from South China Sea (running project)
- Etc.

More information can be found at: <http://www.boldsystems.org/views/projectlist.php>

### 3. Material and methods

The inventory and subsequent cataloguing were focused on recent indigenous and non-indigenous fishes and lampreys living in the natural waters of the Czech Republic. It concerned 11 orders, 17 families and 72 species. The selection of collection localities covered the main distribution areas of the studied taxa according to comprehensive surveys and the most recent taxonomic and phylogenetic studies (Baruš & Oliva, 1995; Hanel & Lusk, 2005; Kottelat & Freyhof, 2007; Janko et al., 2007; Mendel et al., 2008d, etc). The collection of samples was divided into three levels:

- classification of the first level is based on the hydrologic position of the Czech Republic (Fig. 3)

Concerning hydrologic division, the Czech Republic belongs to three sea drainage areas (the North Sea, Baltic Sea and Black Sea drainage areas).



Fig. 3. Collection localities – 1<sup>st</sup> level of classification.

- classification of the second level (Fig. 4)

The hydrologic network of the Czech Republic was further divided into six areas, which for the most part correspond with the division of state administration: I. the Ohře River basin (the North Sea drainage area, the actual Ohře River basin and parts of other tributaries of the Elbe), II. the Vltava River basin (the North Sea drainage area, without the Ohře and the Elbe River basins), III. the Elbe River basin (the North Sea drainage area, without the Ohře and the Vltava River basins), IV. the Dyje River basin and the lower course of the Morava River, V. the Morava River basin (the Black Sea drainage area, the upper and middle courses of the Morava River), VI. the Odra River basin (the Baltic Sea drainage area).



Fig. 4. Collection localities – 2<sup>nd</sup> level of classification.

- classification of the third level

The selection of the sampling localities for the given species was further specified according to up-to-date information concerning its distribution, according to the specific features of the given species (for example, the Natura 2000 localities, etc.) and with regard to the



elimination of “family sampling”. In the case of threatened and critically endangered fish species (*Pelecus cultratus*, *Zingel zingel*, *Zingel streber*, etc.) the database of the tissue samples collected by our workplace in the past was used.

The capturing of adult specimens was done using electro-fishing gear (direct pulsed current) in collaboration with the Czech and Moravian Fishing Unions and the representatives of the Agency for Nature Conservation and Landscape Protection of the Czech Republic. Approximately 1,400 specimens were taken for subsequent molecular-genetic analyses. From the captured specimens (about five individuals per location) a part of the pectoral fin (fin border) was taken and then they were released back into the stream. Only an indispensable number of complete type specimens were taken for morphological description and museum collections.

The identification of the individual species was made on the background of the study of morphological characteristics of the species using a multilocus and two-genome comparative approach (analysis of both mt and nDNA markers). Thus some limitations of the DNA barcoding method were eliminated, e. g. a) occurrence of hybrids; b) maternal contribution to COI marker as only one part of the information necessary for a valid description of a species. For reasons of saving space we do not present here the morphological characteristics of the individual species and the detailed results of the analyses of the nuclear marker – 1<sup>st</sup> intron of the S7 r-protein (RP1).

The preparation of the DNA barcodes conformed to the standardized protocols developed by the CCDB Centre, the FISH-BOL initiative and the BoLD database (<http://www.ccdb.ca/pa/ge/research/protocols>; Ratnasingham & Hebert, 2007; Hajibabaei et al., 2005; deWaard et al., 2008).

DNA isolation was performed by using a commercial kit - the Genomic DNA Mini kit (KRD) - according to the instructions for use. Cocktails of primers from the genetic study (Ivanova et al., 2007) were used for PCR amplification of the COI gene fragment (Fig. 5). For sequencing PCR, mainly M13F and M13R primers (Messing, 1983) were used, in a small number of cases the primers described by Ivanova et al. (2007) were used.




Fig. 5. Mitochondrial DNA – the area of the COI marker – adopted from <http://www.barcodeoflife.org>.

At the Canadian workplace, the subsequent steps were performed according to the above mentioned protocol. At the Czech workplace, the PCR products were re-purified by precipitation with PEG/Mg/NaAc solution (26% Polyethylene glycol, 6.5 mM  $\text{MgCl}_2 \cdot 6\text{H}_2\text{O}$ , 0.6 M  $\text{NaAc} \cdot 3\text{H}_2\text{O}$ ) and subsequently used for the sequencing reaction using BigDye™ Terminator Cycle Sequencing kit v 1.1. (Applied Biosystems), performed according to the manufacturer's instructions. The products of the sequencing reaction were re-purified with ethanol precipitation. For the COI sequencing, the instrumentation of the Canadian BIO workplace (ABI 3730XL; Applied Biosystems) was used and for test and complementary COI sequencing and nuclear marker analysis also the instrument ABI PRISM 310 (Applied Biosystems) at the Czech workplace was used. Also, commercial sequencing (Macrogen, Korea) was used in optimized samples. The analyzed fragment of mitochondrial DNA from each sample was sequenced from both directions. Chromatograms were adjusted and developed using the computer module SeqManII (part of the program package Lasergene v. 6.0; DNASTAR Inc.) and MEGA software.

The developed sequences were visually checked for errors and subsequently compared using the ClustalW algorithm. The correct taxonomic classification of the sequences was confirmed using comparison with the GenBank and BoLD databases. An electropherogram depicting the order of nucleotides in the sequence was obtained through sequencing. With its subsequent processing the DNA barcode was produced (Fig. 6).



Fig. 6. DNA barcode.

The individual steps of the barcoding process are shown in Figure 7.

The morphologic characteristics were identified in selected adult specimens to enable the establishment of a reference type series. A holotype was selected as a nomenclatoric type and other specimens from the described group were assigned to it as the documentary material (paratypes).

For the purpose of the cataloguing of the taxa of the ichthyofauna of the Czech Republic using the DNA barcoding method, the basic data concerning each specimen were collected and completed, including the body length (SL), weight and sex, photo documentation, locality description (including the GPS coordinates), date of collection and the person who caught or identified the specimen, the catalogue number and place of storage. The COI sequences of a length of at least 500 bp and the sequences of used PCR primers were recorded as well. For further details see the instructions of the National Museum and the standardized protocol of the BoLD database (<http://www.boldsystems.org/docs/boldmas.html>; Ratnasingham & Hebert, 2007).

In all archived specimens of each species a sequencing of RP1 marker from nuclear genome was also performed in order to exclude the presence of hybrid specimens. The correct nuclear identification is ensured by sequence comparison with the results of various studies from both domestic and foreign workplaces (e. g., He et al., 2008; Mayden et al., 2008; Mendel et al., 2008d; etc.) and international initiatives (Cypriniformes Tree of Life; Assembling the Tree of Life; etc.)




Fig. 7. The barcoding pipeline (<http://www.barcodeoflife.org/content/about/what-dna-barcoding>).

The platform of the BoLD system (<http://www.barcodinglife.org>) with three identification-statistical modules - MAS, IDS and ECS - was used for the organization of the results. The obtained sequences are saved by the CBOL consortium primarily in the BoLD datasystem and then automatically sent, within INSDC (International Nucleotide Sequence Database Collaboration), to the interconnected databases: GenBank, EMBL and DDBJ. For more information see the standardized protocol of the CBOL consortium: <http://www.dnabarcoding.ca/page/research/protocols/>.

#### 4. Results

An analysis of the COI gene sequences of 500-652 bp in length was performed on 820 individuals from 67 species, which represent 93.1% of the Czech ichthyofauna (Table 1). The map of sampling distribution (Fig. 8) clearly shows that we have evenly sampled the whole territory of the Czech Republic. Altogether, 109 localities were sampled. The number of individuals per species ranged from 1 to 31, with an average number of 12 individuals per species. Out of a total number of 72 species, the following species, which occur very rarely, were not analysed: *Micropterus salmoides*, *Romanogobio banaticus*, *Romanogobio belingi*, *Ameirus melas* and *Acipenser baeri*. Only one or two individuals were analysed in some critically endangered (*Zingel zingel*, *Zingel streber*, *Sabanejewia balcanica*, *Eudontomyzon mariae*, *Pelecus cultratus*, *Ballerus sapa*, *Gymnocephalus schraetser*), and endangered (*Misgurnus fossilis*) species. Neither insertions/deletions nor stop codons were found, thus encouraging the view that all amplified sequences represent functional COI gene sequences.

Family	Species	N	No of haplotypes	Intraspecies difference range (%)
Acipenseridae	<i>Acipenser gueldenstaedtii</i>	3 (+1)	1 (+1)*	0 (4.484)
	<i>Acipenser ruthenus</i>	4	1	0
	<i>Acipenser stellatus</i>	5	1	0
Anguillidae	<i>Anguilla anguilla</i>	5	5	0.154 – 3.494
Balitoridae	<i>Barbatula barbatula</i>	27	13	0 – 5.967
Centrarchidae	<i>Lepomis gibbosus</i>	10	1	0
Cobitidae	<i>Cobitis</i> sp.	3	2	0 – 0.677
	<i>Misgurnus fossilis</i>	1	1	-
	<i>Sabanejewia balcanica</i>	1	1	-
Cottidae	<i>Cottus gobio</i>	21	4	0 – 0.773
	<i>Cottus poecilopus</i>	15	6	0 – 2.681
Cyprinidae	<i>Abramis brama</i>	29	4	0 – 0.617
	<i>Alburnoides bipunctatus</i>	14	5	0 – 0.617
	<i>Alburnus alburnus</i>	26	4	0 – 0.308
	<i>Aspius aspius</i>	14	2	0 – 0.308
	<i>Ballerus ballerus</i>	3	1	0
	<i>Ballerus sapa</i>	2	1	0
	<i>Barbus barbus</i>	25	2	0 – 0.163
	<i>Blicca bjoerkna</i>	14 (+1)	4 (+1)**	0 – 0.308 (4.316)
	<i>Carassius carassius</i>	16	5	0 – 1.266
	<i>Carassius gibelio</i>	27	6	0 – 0.621
	<i>Carassius langsdorfii</i>	3	1	0
	<i>Ctenopharyngodon idella</i>	3	1	0
	<i>Cyprinus carpio</i>	10	2	0 – 0.31
	<i>Gobio gobio</i>	27 (+1)	3 (+1)***	0 – 0.308 (11.420)
	<i>Gobio obtusirostris</i>	5	3	0 – 0.308
	<i>Hypophthalmichthys molitrix</i>	3	1	0
	<i>Chondrostoma nasus</i>	25	5	0 – 0.308
	<i>Leucaspis delineatus</i>	5	2	0 – 3.808
	<i>Leuciscus idus</i>	13	4	0 – 0.621
	<i>Leuciscus leuciscus</i>	21 (+5)	2 (+1)†	0 – 0.161 (0.933)
	<i>Pelecus cultratus</i>	2	1	0
	<i>Phoxinus phoxinus</i>	25	10	0 – 3.973
	<i>Pseudorasbora parva</i>	22	7	0 – 1.084
	<i>Rhodeus amarus</i>	18	7	0 – 1.244
	<i>Romanogobio vladykovi</i>	7	3	0 – 0.618
	<i>Rutilus rutilus</i>	31	7	0 – 0.795
	<i>Scardinius erythrophthalmus</i>	16	1	0
	<i>Squalius cephalus</i>	28	4	0 – 5.252
	<i>Tinca tinca</i>	24	3	0 – 1.521
	<i>Vimba vimba</i>	4	1	0
Esocidae	<i>Esox lucius</i>	22	4	0 – 0.772
Gasterosteidae	<i>Gasterosteus aculeatus</i>	2	1	0
Gobiidae	<i>Neogobius melanostomus</i>	5	1	0
	<i>Proterorhinus semilunaris</i>	6	2	0 – 1.087
Ictaluridae	<i>Ameiurus nebulosus</i>	2	1	0
Lotidae	<i>Lota lota</i>	16	3	0 – 0.308



Percidae	<i>Gymnocephalus baloni</i>	1	1	-
	<i>Gymnocephalus cernua</i>	16	5	0 - 0.798
	<i>Gymnocephalus schraetser</i>	2	1	0
	<i>Perca fluviatilis</i>	29	5	0 - 0.772
	<i>Sander lucioperca</i>	16	1	0
	<i>Sander volgensis</i>	2	1	0
	<i>Zingel streber</i>	1	1	-
	<i>Zingel zingel</i>	1	1	-
Petromyzontidae	<i>Eudontomyzon mariae</i>	2	2	0.197
	<i>Lampetra planeri</i>	14	3	0 - 0.308
Salmonidae	<i>Coregonus maraena</i>	4 (+1)	1 (+1)††	0 (1.558)
	<i>Coregonus peled</i>	5	4	0 - 0.618
	<i>Hucho hucho</i>	2	1	0
	<i>Oncorhynchus mykiss</i>	24	5	0 - 0.308
	<i>Salmo salar</i>	3	2	0 - 0.154
	<i>Salmo trutta</i>	31	4	0 - 1.087
	<i>Salvelinus fontinalis</i>	16	4	0 - 0.617
	<i>Thymallus thymallus</i>	16	2	0 - 0.155
Siluridae	<i>Silurus glanis</i>	11	1	0
Umbridae	<i>Umbra krameri</i>	5	1	0

Table 1. List of species analysed in the current study. N - number of studied individuals (further individuals with haplotypes shared with another species in parentheses). No of haplotypes - number of COI haplotypes found (further haplotypes shared with another species in parentheses). \* - shared haplotype with *Acipenser ruthenus*. \*\* - shared haplotype with *Abramis brama*. \*\*\* - shared haplotype with *Romanogobio vladykovi*. † - shared haplotype with *Leuciscus idus*. †† - shared haplotype with *Coregonus peled*. Intraspecies difference range - range of intraspecific genetic distances excluding haplotypes shared with another species (maximum genetic distance including shared haplotypes in parentheses).




Fig. 8. Overview of sampling localities in the Czech Republic. The borders of the Czech Republic are schematically drawn in red.



As we expected, we found a growing genetic divergence with raising taxonomical level. On the intra-species level, the distances ranged 0-5.97% (Fig. 9), not including inter-specific shared haplotypes (hybrids, misidentifications etc.). Intra-specific diversities exceeding 5% were detected in *Barbatula barbatula* (the highest value of 5.97% was found here), and in *Squalius cephalus* (0-5.25%), values exceeding 2% were found in *Phoxinus phoxinus* (0-3.97%), *Anguilla anguilla* (0.15-3.49%), *Leucaspis delineatus* (0-3.81%) and *Cottus poecilopus* (0-2.68%). In these six mentioned species, there was an apparent deep divergence of the lineages that had been assigned to a single species. Two or three (*B. barbatula*, *S. cephalus*) lineages were distinguished in each species, both from the phylogenetic tree (Fig. 10) and the list of variable sites (Fig. 11). Two lineages can also be distinguished in the following species, although they possess lower genetic distances: *Tinca tinca* (0-1.52%) and *Carassius carassius* (0-1.27%). In *Rhodeus amarus*, a single individual diverges from all other, which raises the mean intra-species diversity from 0.46 to 1.09%.




Fig. 9. Plot of an average intraspecific divergence. Species that exceeded 2% are listed.

An analysis of distribution of the nearest-neighbour distance (NND, the minimum distance between a species and its closest – usually congeneric – relative, Table 2) has shown that only a single species pair exhibited NND lower than 1%, that being namely: *Leuciscus leuciscus* and *L. idus*. The NND averaged 9.67%, which was 40 times higher than the mean within species distance (ca. 0.25%) and 14 times higher than the mean maximum intraspecific distance (0.70%).

We have detected altogether nine cases of shared barcode haplotypes: one *Acipenser gueldenstaedtii* individual possessed an *A. ruthenus* haplotype, one *Blicca bjoerkna* individual possessed an *Abramis brama* haplotype, one *Gobio gobio* individual possessed a *R. vladykovi* haplotype, one *Coregonus maraena* possessed a *C. peled* haplotype, and five *Leuciscus leuciscus* carried a *L. idus* haplotype.

A phylogenetic tree based on all sequences longer than 500 bp, employing the Kimura-2-parameter (K2P) model is available on <http://www.ivb.cz/projects-molecular->

biodiversity-inventory-of-the-ichthyofauna-of-the-czech-republic.html. Subsequently, we selected the representatives of all species or intra-species lineages with intra-species distance at least three times higher (1.176%) than the mean distance within the species (0.392%; Hubert et al., 2008). Based on these sequences, a reduced phylogenetic tree was constructed and put through a bootstrap analysis of 1000 replications. The reduced phylogenetic tree (Fig. 10) documents strong bootstrap support both on the level of genera and on family level. Only the most numerous family Cyprinidae has exhibited moderate support. The *Neogobius* and *Proterorhinus* genera were not supported in forming a single clade of the Gobiidae family.

The phylogenetic tree supports the most recently proposed changes in the scientific taxonomical nomenclature. According to recent taxonomical opinions and studies (Froese & Pauly, 2010; Kottelat & Freyhof, 2007; Perea et al., 2010), *A. sapa* and *A. ballerus* have been currently sorted into the genus *Ballerus*; *Abramis bjoerkna* now belongs again to the *Blicca* genus, *L. cephalus* into *Squalius* and it is suggested that *Aspius aspius* be included into *Leuciscus*. All these proposales are reflected in the tree, as the above mentioned species always cluster together.

Order	Family	N	< 0.1	0.1 - 1.0	1.0 - 2.7	> 2.7
Acipenseriformes	Acipenseridae	3				3
Cypriniformes	Cobitidae	3				3
	Cyprinidae	30		2*	2**	26
	Balitoridae	1				1
Scorpaeniformes	Cottidae	2				2
Salmoniformes	Salmonidae	8			2***	6
Petromyzontiformes	Petromyzontidae	2				2
Esociformes	Umbridae	1				1
	Esocidae	1				1
Perciformes	Gobiidae	2				2
	Percidae	8				8
	Centrarchidae	1				1
Gadiformes	Lotidae	1				1
Anguilliformes	Anguillidae	1				1
Siluriformes	Siluridae	1				1
	Ictaluridae	1				1
Gasterosteiformes	Gasterosteidae	1				1
Totals:		67	0	2	4	61

Table 2. Summary of Czech ichthyofauna diversity and distribution of genetic distance of each species analysed to nearest neighbour (in per cent). N - number of species in family.  
\* - *Leuciscus idus* and *Leuciscus leuciscus* (0.62%). \*\* - *Blicca bjoerkna* and *Vimba vimba* (2.51%).  
\*\*\* - *Coregonus peled* and *Coregonus maraena* (1.4%).




Fig. 10. K2P/NJ tree of representatives of each species, or lineage within species. Bootstrap values are listed near the nodes, only values  $\geq 75\%$  are shown. Families are listed in colour. Species showing an internal divergence more than three times higher than the mean distance within the species are listed in bold italics.

[	12222222	2223333333	4444445555	5555555666	666]	[	1222333	3444455556	666]
[	2389223345	6890112379	0113680015	6667899011	22]	[	2247128114	7013802380	012]
[	5150035870	5081362408	9250645810	2581038469	25]	[	5605796030	0059426001	438]
#IFCZE0815 Barbatula_barbatula	ACCCACCGCG	CGCTACGGAAC	CGTAAAGGATC	TGCACCCCGG	GA	#IFCZE0248 Anguilla_anguilla	AGCTCAACAA	GCTCGTCTGG	CTA
#IFCZE0556 Barbatula_barbatula	.....G.....	.....	.....	.....	..	#IFCZE0603 Anguilla_anguilla	GA.....	.....	A..
#IFCZE0802 Barbatula_barbatula	.....G.....	.....	.....	.....	..	#IFCZE0771 Anguilla_anguilla	A.....	.....	A..
#IFCZE0809 Barbatula_barbatula	.....G.....	.....	.....	.....	..	#IFCZE0604 Anguilla_anguilla	..TCT..TGG	ATCTACTCAA	..CG
#IFCZE11111 Barbatula_barbatula	.....C.....	.....	.....	.....	..	#IFCZE0772 Anguilla_anguilla	..TCTGGTG.	ATCTACTCAA	..CG
#IFCZE0555 Barbatula_barbatula	G.....T.....	GA..GT	..G..A..	.....	A..	[	2223333	3444455556]	
#IFCZE0835 Barbatula_barbatula	.....T.....	GA..GT	..G.....	.....	A..	[	1560490147	702394561]	
#IFCZE0849 Barbatula_barbatula	.....T.....	GA..GT	..G..A..	.....	A..	[	0548181300	397067329]	
#IFCZE11031 Barbatula_barbatula	.....T.....	GA..GT	..G..A..	.....	G....A..	#IFCZE0187 Cottus_poecilopus	AAGAAAAATA	CATATTGCC	
#IFCZE11071 Barbatula_barbatula	.....T.....	GA..GT	..G..A..	.....	A..	#IFCZE1132 Cottus_poecilopus	G.....	.....	
#IFCZE0726 Barbatula_barbatula	..TGTCAT..A	ATA..AAG..	TAC..A.GCT	CTTCT.TAGA	A..	#IFCZE1133 Cottus_poecilopus	.....T.....	.....	
#IFCZE0727 Barbatula_barbatula	..TGTCAT..A	ATA..AAG..	TAC..A.GCT	CTTCT.TAGA	AG	#IFCZE1137 Cottus_poecilopus	.....G.....	.....	
#IFCZE0730 Barbatula_barbatula	..TGTCAT..A	ATA..AAG..	T.C..A.GCT	CTTCT.TAGA	A..	#IFCZE0746 Cottus_poecilopus	..CAG.GGGAG	..GG.CCAAT	
[	12222223	333333444	4444445555	556666	6]	#IFCZE0750 Cottus_poecilopus	..CAG.GGGAG	..GGGCCAAT	
[	4600144700	2223457000	0368991235	71235]		[	111223345]		
[	7028747447	2581020036	9334797023	10242]		[	089041210]		
IFCZE0830 Squalius_cephalus	CAGAAATTATG	GGAGACACTA	GGAAACCAAT	ACGAA		[	076249259]		
IFCZE0680 Squalius_cephalus	.....A..G..	A.....G..	G....			#IFCZE0264 Tinca_tinca	TACGTAGGC		
IFCZE0785 Squalius_cephalus	.....G.....					#IFCZE0853 Tinca_tinca	.....T		
IFCZE0534 Squalius_cephalus	TCAGGCCGAA	AA.AGT.TCG	..AGGTAGGCC	..TAGG		#IFCZE0394 Tinca_tinca	CGTACGAT.		
[	111	2222333333	4444445556	6]		[	2222334	56]	
[	3448899099	3679144456	0389903561	4]		[	2550277256	54]	
[	7032512836	2515903627	3643655953	3]		[	0563917550	09]	
#IFCZE0682 Phoxinus_phoxinus	GGCGGCTGT	CAATATCCCA	CCGATCTTTT	C		#IFCZE0048 Rhodeus_amarus	CTCCACCCC	GC	
#IFCZE0684 Phoxinus_phoxinus	A...A....	T...CA....	A.....			#IFCZE0050 Rhodeus_amarus	G.....		
#IFCZE0686 Phoxinus_phoxinus	A...A....	T.G..CA....	A.....			#IFCZE0494 Rhodeus_amarus	GCT.....		
#IFCZE0767 Phoxinus_phoxinus	.....A....	T...GCA..G	A.....			#IFCZE0633 Rhodeus_amarus	G.....A.		
#IFCZE0805 Phoxinus_phoxinus	.....AT....	T...CA....	A.....			#IFCZE0822 Rhodeus_amarus	G...T.....		
#IFCZE0816 Phoxinus_phoxinus	AA..A....	T...CA....	A.....			#IFCZE10047 Rhodeus_amarus	G.....G..		
#IFCZE0127 Phoxinus_phoxinus	..TTATT.A.	TG.CGCGTT.	TT.CGCGCA	T		#IFCZE0076 Rhodeus_amarus	G..T.GTT.T	T	
#IFCZE0255 Phoxinus_phoxinus	..TCATTCA.	TG.CGCGTT.	TT.CGCGCAC	T		[	1223344]		
#IFCZE0551 Phoxinus_phoxinus	..TT.TT.A.	TG.CGCGTT.	TT.CGCGCAC	T		[	68272478]		
#IFCZE0834 Phoxinus_phoxinus	..TTATT.A.	TG.CGCGTT.	TT.CGCGCAC	T		[	20012077]		
#IFCZE0914 Phoxinus_phoxinus	..TTATT.AG	TG.CGCGTT.	TT.CGCGCAC	T		[			
[	1111222	2333334455	5556]			#IFCZE0571 Carassius_carassius	CATATTGC		
[	4780037013	7122895612	4572]			#IFCZE10150 Carassius_carassius	G.....		
[	9926938378	7958877640	1015]			#IFCZE0392 Carassius_carassius	..CGCCAT		
#IFCZE0936 Leucaspisus_delineatus	GGTGTCOCCT	CGGGGATGAA	CTAA			#IFCZE0821 Carassius_carassius	..C.CCAT		
#IFCZE0938 Leucaspisus_delineatus	AACACTATAG	TAAAGCGCTG	TAGG			#IFCZE10051 Carassius_carassius	CGCCAT		

Fig. 11. Summary of variable sites in the COI gene in haplotypes of nine species showing an internal divergence more than three times higher than the mean distance within the species. Numbers show variable position of nucleotide in whole 652 bp sequence. Dots represent nucleotide identical with first sequence.

## 5. Discussion

This study presents the first molecular screening of the whole Czech ichthyofauna. Indeed, the position of our project, „The Ichthyofauna of the Czech Republic“ (abbreviated IFCZE in the BoLD database), is in a certain sense unique. Table 3, which compares genetic divergences on various taxonomic levels in projects spanning countries across continents from North America to Asia, Australia and Africa, contains only a single project representing the European ichthyofauna – Czech project. It is the first and currently the only public project in BoLD database dealing with the barcoding of European ichthyofauna on this scale.

Among the analyzed species, there was no species endemic to the Czech Republic. Aside from indigenous species, further 18 non-indigenous species introduced in the past to the Czech ichthyofauna were also tested. One species is considered to be regionally extinct in the wild, 12 species are considered to be critically endangered, five to be endangered and six to be vulnerable according to the IUCN classification (Lusk et al., 2011). Also included in the project was the *Umbra krameri*, while its indigenuity to the Czech Republic is currently disputed.



The amplification of the 5′ region of the COI gene (up to 652 bp) was successful for all 820 tested individuals, while utilizing a single primer set for amplification of all tested species. No nuclear copies of the COI gene were detected, neither frame shifts nor mutations causing the emergence of stop codons. By means of DNA barcoding, all tested species could be successfully distinguished. The DNA barcodes were acquired for 67 species. An example of such a barcode record can be seen in Figure 12.

	CAN	MEX	NorthAm.	IND	PHL	NGA	AUS	CZE	
	min%								
within species	0	0	0	0	0	0	0	0	
within genus	0	0	0	0.10	10.0	1.80	0	0	
within family	2.67	1.804	0.185	0.20	5.50	0.46	1.39	0	
within order	14.25	17.24	14.53	8.00	16.20	15.52	9.55	18.07	
within class	17.49	17.88	16.42	-	-	18.46	14.33	18.36	
	mean dist%								
within species	0.27	0.45	2.52	0.30	0.60	0.30	0.39	0.39	
within genus	8.37	5.10	14.10	6.60	11.70	1.90	9.93	5.00	
within family	15.38	13.57	16.00	9.91	17.67	17.19	15.46	14.85	
within order	20.60	23.38	21.50	16.00	24.80	21.62	22.18	24.62	
within class	24.57	25.34	24.80	-	-	25.40	23.27	24.87	
	max dist%								
	S.E.								
within species	7.42	2.11	15.45	0.30	9.30	22.38	14.80	11.61	0.013
within genus	19.33	16.67	23.68	11.90	15.30	36.61	20.63	11.20	0.067
within family	23.22	28.48	29.92	23.10	22.60	24.82	35.7	24.46	0.011
within order	29.44	28.77	30.95	23.40	31.50	28.12	37.52	28.84	0.013
within class	31.20	31.16	37.50	-	-	31.90	37.39	30.77	0.004
species	190	76	605	98	23	59	207	61	
genus	85	56	134	78	21	33	122	51	
family	28	32	36	53	17	19	?	17	
order	20	11	18	24	9	8	?	11	
class	2	2	2	4	-	2	?	2	

Table 3. Comparison of genetic divergences (K2P model used) within various taxonomic levels among selected freshwater and marine fishes BoLD projects. (CAN-Barcoding of Canadian freshwater fishes, MEX-Freshwater Fishes of Mexico, NorthAm-Freshwater Fishes of North America, IND-Marine Fishes of India, PHL-Freshwater Fish of Taal Lake, Philippines, NGA- Barcoding Freshwater Fishes of Eastern Nigeria, AUS-Marine Fishes of Australia, CZE-Ichthyofauna of the Czech Republic).

The barcoding of the Czech ichthyofauna has not ended yet: several further species (*M. salmoides*, *R. banaticus*, *R. belingi*, *A. baerii*, and *A. melas*) are currently being tested as well. Therefore, the analyzed sample set currently covers 67 of 72 species, or 93.1% of the Czech ichthyofauna. Certain species are present in all six investigated regions and were therefore analysed in greater numbers and in some cases exhibited greater haplotype variability (e.g. *B. barbatula*, 27 individuals/13 haplotypes or *P. phoxinus*, 25 individuals/10 haplotypes. For more details see Table 1.



Overall, the mean congeneric divergence was ca. 13 times higher than the mean conspecific divergence (5.00 vs 0.39%, Table 3). These values are in good concordance with the results of Valdez-Moreno et al. (2009), which studied the diversity of ichthyofauna in Mexico and Guatemala (5.10 vs 0.45%, Table 3). The divergence ratio was higher in studies dealing with the ichthyofauna of Australia (25x, Ward et al., 2005), Canada (31x, Hubert et al., 2008) and the Philippines (19x, Aquilino et al., 2011) (Table 3). Such a low divergence in our project is probably caused by overall more recent speciation of the freshwater species in comparison to their marine counterparts (which are included in the mentioned projects). This is also supported by the results of the marine project from India (27x, Lakra et al., 2011), China (50x, a running project) and Japan (59x, Zhang & Hanner, 2011).

The overlapping of conspecific and congeneric levels of divergence was minimal: only two genera have shown inter-specific distance between the nearest neighbours lower than 2.7%: *Leuciscus* – *L. leuciscus* and *L. idus* (0.62%) and *Coregonus* – *C. peled* and *C. maraena* (1.4%). Interestingly, in one case an inter-specific distance lower than 2.7% occurred between two non-congeneric species: *B. bjoerkna* and *Vimba vimba* (2.51%, Table 2). Zardoya & Doadrio (1999) and Cunha et al. (2002) observed the same effect, based on the sequencing of a different marker. All congeneric species always clustered together in the phylogenetic analysis. Furthermore, several species pairs (not always congeneric) have been shown to possess shared haplotypes. In most cases, a single individual of one species bore a haplotype typical of a close (sister) species. Such was the case of a single *B. bjoerkna* among *A. brama*, *G. gobio* among *R. vladykovi*, *C. maraena* among *C. peled*, and *A. gueldenstaedtii* among *A. ruthenus*. All these cases can be easily explained as simple hybridization, in the case of *Coregonus* even human-driven intentional hybridization, (with the second species being the maternal one due to the maternal inheritance of mtDNA), which was confirmed by the sequencing of the nuclear marker (first intron of the S7 ribosomal protein gene).

The last species pair, *L. leuciscus* and *L. idus*, represented an exception. We found five individuals classified as *L. leuciscus* bearing the haplotype of *L. idus*. These two species also presented the lowest mutual genetic distance from all analyzed species pairs (0.62%). We therefore conclude that in these two species, not only hybridization (as confirmed by the sequencing of S7) but also probably recent speciation might have taken place in the genetic similarity of both species.

Analyses have revealed a significant deep intraspecies divergence in several taxa. In most cases, those are species with large areas (Kottelat & Freyhof, 2007), such as *B. barbatula*, *L. delineatus*, *P. phoxinus* or *S. cephalus*.

The most pronounced example is the stone loach (*B. barbatula*). The DNA barcoding technique has revealed high intra-specific divergence (up to 5.97%, Table 1). We have tested 27 individuals from six regions and we found 13 haplotypes diverging into three lineages (Table 1, Fig. 10, Fig. 11). Šedivá et al. (2008) has described the presence of the lineage V in the Labe River basin and the sublineage V<sub>a</sub> in the Morava River basin, both of the lineages belonging to the Danubian clade. Our results are in good concordance with these findings. Furthermore, we have distinguished a third, the most divergent lineage, occurring in the Odra River basin. This lineage possibly belongs to the Eastern clade similarly to previously detected individuals from the same basin in Poland (Šedivá et al., 2008).




Fig. 12. An example of barcode record of the species.

The second deepest divergence has been detected in chub (*S. cephalus*). We have tested 28 individuals and found four haplotypes, diverging into two lineages (Table 1, Fig. 10, Fig. 11), separated by 5.25%. The less abundant lineage occurred only in the North Sea drainage area (Labe and Vltava River basins), the second, more numerous lineage was found across all river basins. Seifertová et al. (2008) in their large study of phylogenetic diversity of chub across its European range distinguished four different haplogroups, corresponding to four lineages, i.e. the Western, Eastern, Aegean and Adriatic (*sensu*

Durand et al., 1999) ones, with the populations collected in the Czech Republic belonging to the Western Lineage. It is probable that our more prevalent lineage corresponds to the Western lineage, sharing its distribution pattern. It is unclear, but possible that the second of our lineages corresponds to the Eastern lineage. However, the Eastern lineage was not found by Seifertová et al. (2008) in the North Sea basin, but in populations from the Vistula River basin in Poland and in Finland. Therefore we cannot exclude the possibility that we have found a new, fifth lineage. A definite conclusion about the identity of lineages can not be drawn due to different markers utilized in both studies. The existence of other deeply divergent lineages of chub is also evident e.g. in Perea et al. (2010), who found two lineages of chub in their study of phylogenetic relationships in the Circum-Mediterranean subfamily Leuciscinae. It is clear that the taxonomy of *S. cephalus* needs to be further addressed; both on the level of additional samples, which we have already collected, and by the unification of utilized markers.

The DNA barcoding further discriminated two deep lineages in two other cyprinid species, *P. phoxinus* (lineages separated by 3.97%) and *L. delineatus* (lineages separated by 3.81%). In the most recent Red List of the Czech Republic, the minnow (*P. phoxinus*) is classified as vulnerable (Lusk et al., 2011). The priority task in the protection of minnow lies in the long-term sustenance of water quality, the segmentation of river bed and banks together with thoughtful fishing and angling management (Hanel & Lusk 2005). We have tested 25 individuals from six regions and we distinguished 10 haplotypes diverging into two lineages (Table 1, Fig. 10, Fig. 11). Lineage I is located in water courses flowing into the North and Black Sea drainage areas and the lineage II is located in the Baltic Sea drainage area (including the Bečva River). While the Bečva River currently belongs to the Black Sea drainage area, here we can consider an ancient proximity or identical origin of the two populations, as during the Pleistocene (e.g. the Elster glaciation) the waters of the Odra flowed towards the south into the drainage area of the river Bečva (Czudek, 1997). The systematics of the genus *Phoxinus* has not been solved yet.

The belica (*L. delineatus*) is currently classified among the critically endangered species in the Red List of the Czech Republic (Lusk et al., 2011). Apart from food competition and the continuous forcing out of original localities by invasive species, its vanishing is also probably influenced by the spread of a disease caused by the pathogen *Sphaerothecum destruens*, which came to Europe through an invasion of the invasive fish species *Pseudorasbora parva* with the import of herbivorous fish fry (Gozlan et al., 2005; Hanel & Lusk 2005). Interestingly, both our lineages of the belica coexist in the same locality. Those five tested individuals came from one fish hatchery in the Ohře River basin. Currently, there is no detailed phylogenetic study of this species. Only Perea et al. (2010) dealt with the taxonomy and systematics of this genus, but based on two individuals only and in the context of general phylogenetic relationships and biogeographical patterns in the Circum-Mediterranean subfamily Leuciscinae. Our studies of this species continue in order to collect individuals from further localities. We also aim to evaluate hybrid events in the scope of uncovered lineages based on a nuclear marker.

Two separate lineages were also detected in eel (*A. anguilla*) (3.49%) and Siberian bullhead (*C. poecilopus*) (2.68%). The intra-specific diversity of eel on a molecular level has not been studied yet. Both of our two lineages were found in North Sea and Baltic Sea drainage areas. However, when performing a comparison of our sequences with other sequences stored in



BoLD, one of these lineages is identical to samples of the American eel (*A. rostrata*). Kottelat & Freyhof (2007) mention that the American eel has been stocked in Eastern Europe; our findings confirm their suspicion that the presence of isolated individuals of American eel cannot be ruled out in our free waters. In order to confirm these findings, further samples will be evaluated.

In the Siberian bullhead, Lusk et al. (2008a) distinguished two complexes based on the sequencing of the *cyt b* gene and mitochondrial control region: the “baltica” complex, which consists of the populations from the Baltic Sea drainage area, and the “danubialis” complex, which consists of the populations from the Black Sea drainage area. An existence of the lineages uncovered in this study further supports these findings: our tested populations from the Odra River basin fall into the “baltica” complex, the individuals from the Morava and Dyje River basins fall into the “danubialis” complex. Lusk et al. (2008a) also further discuss the necessity of a more complex genetic analysis (utilizing a nuclear marker in combination with morphological characteristics) in order to properly evaluate the possible taxonomic status of these complexes.

All these findings (visualized in a bootstrapped NJ tree, Fig. 10) are based on the sequencing of a single marker. Thus they are not *per se* a sufficient tool and criterion for a definitive taxonomical and phylogenetic consequences. For this reason, our project has employed a more complex approach: based on these findings, a sequencing analysis of a nuclear marker must be performed, and comparisons of the morphological characteristics of the lineages must be evaluated. Such complex evaluations will subsequently serve for the definitive confirmation of taxonomic conclusions, and also for the reference collection of the fish which will be stored in the National Museum of Natural History, Prague.

## 6. Conclusion

The presented study provides a clear example of the usefulness and suitability of DNA barcoding for the cataloguing of the biodiversity of the Czech freshwater fishes. The lineages newly uncovered in our study are being subjected to further detailed research in order to evaluate their correct taxonomical status and systematic position. Definitive results of the project, once obtained, can probably serve as a good starting point for subsequent comparative phylogenetic studies in an Euroasian context. The presented project also enabled us to focus our attention also on taxa which have never been in the scope of molecular-genetic investigation (e.g. *L. delineatus*, *P. phoxinus*, *Gasterosteus aculeatus*, *A. anguilla*, etc.).

New taxonomically and systematically important knowledge may (and hopefully will) contribute to the updating of the background information for the monitoring of the NATURA 2000 system. It also brings crucial information and recommendations to the Agency for Nature Conservation and Landscape Protection of the Czech Republic. A new design of the species collection of type individuals, constructed on the basis of most recent molecular methods and including morphological data, as well as organizing a unified voucher, is also a great benefit for the National Museum of Natural Science in Prague.

By employing a complex approach (analysis of both the nuclear and mitochondrial genomes plus morphological analyses) we were able to eliminate the danger of including a hybrid individual as a representative of a particular species into the type series. Combined

preservation approaches (whole individuals stored in formaldehyde, fin clippings in ethanol, DNA isolates) targeted at individuals from the new type collections will also enable possible comparative DNA analyses in future, whenever needed. All the results are clearly and concisely assembled in the BoLD database, which allows very simple and quick on-line access to informations for the broadest scientific and for the general public.

The UN has declared the year, 2010, to be the International Year of Biodiversity. Let us all wish much success to all future projects and initiatives dealing with biodiversity; for every successful project will bring us closer to the ultimate goal of the knowledge and understanding of the biodiversity.

## 7. Acknowledgement

This study was carried out within the framework of research projects no. M200930901 supported by the Program of Internal Support for International Collaborative Projects of the Academy of Sciences of the Czech Republic and no. 206/09/P608 supported by the Grant Agency of the Czech Republic. We would like to thank our colleagues from the Biodiversity Institute of Ontario, University of Guelph for their financial, laboratory and informatics support, the Czech and Moravian Fishing Unions, the representatives of the water-resources authorities (pond management companies, river basin management agencies), the management of the Protected Landscape Areas, our colleagues from universities and other people for their help with sample collection.

### Data accessibility

Cytochrome c oxidase subunit I gene sequences, trace files, digital images of fish and other metadata were submitted to Barcoding of Life Data systems (BoLD - <http://www.barcodinglife.com>) and were given BoLD Process ID's IFCZE001-10 to IFCZE701-10 and IFCZE702-11 to IFCZE852-11, all being part of the project „The ichthyofauna of the Czech Republic (IFCZE). The complete NJ/K2P tree can be found on the website of IVB on <http://www.ivb.cz/projects-molecular-biodiversity-inventory-of-the-ichthyofauna-of-the-czech-republic.html>. Acquired DNA sequences are also being stored in GenBank with accession numbers HQ960417-HQ961093.

## 8. References

- April, J., Mayden, R. L., Hanner, R. H. & Bernatchez, L. (2011). Genetic calibration of species diversity among North America's freshwater fishes. *Proceedings of the National Academy of Sciences of the United States of America, In Press*
- Aquilino, S. L., Tango, J. M., Fontanilla, I. C., Pagulayan, R. C., Basiao, Z. U., Ong, P. S. & Quilang, J. P. (2011). DNABarcoding of the ichthyofauna of Taal Lake, Philippines. *Molecular Ecology Resources*, Vol. 11, No. 4, pp. 612-619, ISSN 1755-0998
- Ardura, A., Linde, A. R., Moreira, J. C. & Garcia-Vazquez, E. (2010). DNA barcoding for conservation and management of Amazonian commercial fish. *Biological Conservation*, Vol. 143, No. 6, pp. 1438-1443, ISSN 0006-3207
- Avise, J. C. (2000). *Phylogeography. The history and formation of species*. Harvard University Press, ISBN 978-0674666382, Cambridge



- Barrett, R. D. H. & Hebert, P. D. N. (2005). Identifying spiders through DNA barcodes. *Canadian Journal of Zoology*, Vol. 83, No. 3, pp. 481-491, ISSN 1480-3283
- Bartoňová, E., Papoušek, I., Lusková, V., Koščo, J., Lusk, S., Halačka, K., Švátora, M. & Vetešník, L. (2008). Genetic diversity and taxonomy of *Sabanejewia balcanica* (Osteichthyes: Cobitidae) in the waters of the Czech Republic and Slovakia. *Folia Zoologica*, Vol. 57, No. 1-2, pp. 60-70, ISSN 0139-7893
- Baruš, V. & Oliva, O. (1995). *Mihulovci a Ryby*. Academia, ISBN 80-200-0501-3, Praha
- Cunha, C., Mesquita, N., Dowling, T. E., Gilles, A. & Coelho, M. M. (2002). Phylogenetic relationships of Eurasian and American cyprinids using cytochrome *b* sequences. *Journal of Fish Biology* Vol. 61, No. 4, pp. 929-944. ISSN 1095-8649
- Czudek, T. (1997). *Reliéf Moravy a Slezska v kvartéru*. Sursum, ISBN 80-85799-27-8, Tišnov, (in Czech)
- deWaard, J. R., Ivanova, N. V., Hajibabaei, M. & Hebert, P. D. N. (2008). Assembling DNA barcodes. Analytical protocols. *Methods in Molecular Biology*, Vol. 410, pp. 275-293, ISSN 1064-3745
- Durand, J. D., Persat, H. & Bouvet, Y. (1999). Phylogeography and postglacial dispersion of the chub (*Leuciscus cephalus*) in Europe. *Molecular Ecology*, Vol. 8, No. 6, pp. 989-997, ISSN 0962-1083
- Fisher, B. L. (2006). DNA Barcoding: Tomorrow Is Too Late. *California Wild*, Vol. 59, No. 1, pp. 1-3
- Freeland, J. R. (2005). *Molecular ecology*. John Wiley & Sons Ltd, ISBN 978-0-470-09062-6, Chichester
- Froese, R. & Pauly, D. (2010). FishBase. World Wide Web electronic publication. Available from: [www.fishbase.org](http://www.fishbase.org)
- Gozlan, R. E., St-Hilaire, S., Feist, S. W., Martin, P. & Kent, M. L. (2005). Biodiversity: Disease threat to European fish. *Nature*, Vol. 435, No. 7045, p. 1046, ISSN 0028-0836
- Hajibabaei, M., deWaard, J. R., Ivanova, N. V., Ratnasingham, S., Dooh, R. T., Kirk, S. L., Mackie, P. M. & Hebert, P. D. N. (2005). Critical factors for assembling a high volume of DNA barcodes. *Philosophical Transactions of the Royal Society B: Biological Science*, Vol. 360, No. 1462, pp. 1959-1967, ISSN 1471-2970
- Hajibabaei, M., Smith, M. A., Janzen, D. H., Rodriguez, J. J., Whitfield, J. B. & Hebert, P. D. N. (2006). A minimalist barcode can identify a specimen whose DNA is degraded. *Molecular Ecology Notes*, Vol. 6, No. 4, pp. 959-964, ISSN 1471-8286
- Hajibabaei, M., Singer, G. A. C., Hebert, P. D. N. & Hickey, D. A. (2007). DNA barcoding: how it complements taxonomy, molecular phylogenetics and population genetics. *Trends in Genetics*, Vol. 23, No. 4, pp. 167-172, ISSN 0168-9525
- Hammond, P. M. (1992). Species inventory, In: *Global Biodiversity: status of the Earth's living resources*, B. Groombridge (ed.), pp. 17-39, Chapman and Hall, ISBN 0-41247240-6, London
- Hanel, L. & Lusk, S. (2005). *Ryby a mihule České republiky, rozšíření a ochrana*. ČSOP Vlašim, ISBN 80-86327-49-3, Vlašim
- He, S., Mayden, R. L., Wang, X., Wang, W., Tang, K. L., Chen, W. J. & Chen, Y. (2008). Molecular phylogenetics of the family Cyprinidae (Actinopterygii: Cypriniformes) as evidenced by sequence variation in the first intron of S7 ribosomal protein-coding gene: Further evidence from a nuclear gene of the systematic chaos in the

- family. *Molecular Phylogenetics and Evolution*, Vol. 46, No. 3, pp. 818-829, ISSN 1055-7903
- Hebert, P. D. N., Cywinska, A., Ball, S. L. & deWaard, J. R. (2003a). Biological identifications through DNA barcodes. *Proceedings of the Royal Society B: Biological Science*, Vol. 270, pp. 313-321
- Hebert, P. D. N., Ratsingham, S. & Dewaard, J. R. (2003b). Barcoding animal life: cytochrome c oxidase subunit 1 divergences among closely related species. *Proceedings of the Royal Society B: Biological Science*, Vol. 270 (Suppl 1), pp. 96-99, ISSN 0962-8452
- Hebert, P. D. N., Stoeckle, M. Y., Zemlak, T. S. & Francis, Ch. M. (2004). Identification of Birds through DNA Barcodes. *PLoS Biology*, Vol. 2, No. 10, pp. e312, ISSN 1545-7885
- Hebert, P. D. N. & Barrett, R. D. H. (2005). Reply to the comment by L. Prendini on "Identifying spiders through DNA barcodes". *Canadian Journal of Zoology*, Vol. 83, No.3, pp. 505-506, ISSN 1480-3283
- Hubert, N., Hanner, R., Holm, E., Mandrak, N. E., Taylor, E., Burridge, M., Watkinson, D., Dumont, P., Curry, A., Bentzen, P., Zhang, J., April, J. & Bernatchez, L. (2008). Identifying Canadian Freshwater Fishes through DNA Barcodes. *PloS One*, Vol. 3, No. 6, pp. e2490, ISSN 1932-6203
- Ivanova, N. V., Zemlak, T. S., Hanner, R. H. & Hebert, P. D. N. (2007). Universal primer cocktails for fish DNA barcoding. *Molecular Ecology Notes*, Vol. 7, No. 4, pp. 544-548, ISSN 1471-8286
- Janko, K., Flajšhans, M., Choleva, L., Bohlen, J., Šlechtová, V., Rábová, M., Lajbner, Z., Šlechta, V., Ivanova, P., Dobrovolov, I., Culling, M., Persat, H., Kotusz, J. & Ráb, P. (2007). Diversity of European spined loaches (genus *Cobitis* L.): an update of the geographic distribution of the *Cobitis taenia* hybrid complex with a description of new molecular tools for species and hybrid determination. *Journal of Fish Biology*, Vol. 71, No. 2, pp. 387- 408, ISSN 1095-8649
- Kottelat, M. & Freyhof, J. (2007). *Handbook of European freshwater fishes*. Kottelat, Cornol, Switzerland and Freyhof, ISBN 978-2-8399-0298-4, Berlin
- Kress, W. J., Wurdack, K. J., Zimmer, E. A., Weigt, L. A. & Janzen, D. H. (2005). Use of DNA barcodes to identify flowering plants. *PNAS*, Vol. 102, No. 23, pp. 8369-8374, ISSN 0027-8424
- Lakra, W., Verma, M. & Goswami, M. (2011). DNA barcoding Indian marine fishes. *Molecular Ecology Resources*, Vol. 11, No. 1, pp. 60-71, ISSN 1755-0998
- Lusk, S., Bartoňová E., Lusková, V., Lojkásek, B. & Koščo, J. (2008a). Vranka pruhoploutvá *Cottus poecilopus* – rozšíření a genetická diverzita v povodí řek Morava, Odra (Česká republika) a Hornád (Slovensko) [Siberian sculpin *Cottus poecilopus*: distribution and genetic diversity in the Morava, Odra (Czech Republic), and Hornád (Slovakia) drainage areas], In: *Biodiverzita ichtyofauny ČR (VII)*, S. Lusk, V. Lusková (Ed.), pp. 67-80, Ústav biologie obratlovců AV ČR, v.v.i, ISBN 978-80-87189-01-6, Brno, (in Czech with English summary)
- Lusk, S. & Hanel, L. (2008). Foreword. In: *Biodiverzita ichtyofauny ČR (VII)*, S. Lusk, V. Lusková (Ed.), pp. 3-5, Ústav biologie obratlovců AV ČR, v.v.i, ISBN 978-80-87189-01-6, Brno
- Lusk, S., Papoušek, I., Lusková, V., Halačka, K. & Koščo, J. (2008b). Výskyt a genetická diverzita drska menšího a drska většího v povodí Moravy (Česká republika)

- [Occurrence and genetic diversity of Streber *Zingel streber* and Zingel *Zingel zingel* in the Morava river drainage area (Czech Republic)], In: *Biodiverzita ichtyofauny ČR (VII)*, S. Lusk, V. Lusková (Ed.), pp. 81-87, Ústav biologie obratlovců AV ČR, v.v.i, ISBN 978-80-87189-01-6, Brno. (in Czech with English summary)
- Lusk, S., Lusková, V., Hanel, L., Lojkásek, B. & Hartvich, P. (2011). Červený seznam mihulí a ryb České republiky – verze 2010, [Red List of lampreys and fishes of the Czech Republic – Version 2010], In: *Biodiverzita ichtyofauny ČR (VII)*, S. Lusk, V. Lusková (Ed.), pp. 68-78, Ústav biologie obratlovců AV ČR, v.v.i, ISBN 978-80-87189-08-5, Brno. (in Czech with English summary)
- Lusková, V., Bartoňová, E. & Lusk, S. (2008a). Ostrucha křivočará *Pelecus cultratus* – vzácný objekt ochrany v České republice [The Ziege *Pelecus cultratus*, a rare object of protection in the Czech Republic], In: *Biodiverzita ichtyofauny ČR (VII)*, S. Lusk, V. Lusková (Ed.), pp. 38-45, Ústav biologie obratlovců AV ČR, v.v.i, ISBN 978-80-87189-01-6, Brno. (in Czech with English summary)
- Lusková, V., Bartoňová, E. & Lusk, S. (2008b). Karas obecný *Carassius carassius* Linnaeus, 1758 v minulosti obecně rozšířený a v současnosti ohrožený druh v České republice [Crucian carp, *Carassius carassius* Linnaeus, 1758, a common species in the Czech Republic in the past yet endangered at present], In: *Biodiverzita ichtyofauny ČR (VII)*, S. Lusk, V. Lusková (Ed.), pp. 46-53, Ústav biologie obratlovců AV ČR, v.v.i, ISBN 978-80-87189-01-6, Brno. (in Czech with English summary)
- Mayden, R. L., Tang, K. L., Wood, R. M., Chen, W.-J., Agnew, M. K., Conway, K. W., Yang, L., Simons, A. M., Bart, H. L., Harris, P. M., Li, J., Wang, X., Saitoh, K., He, S., Liu, H., Chen, Y., Nishida, M. & Miya, M. (2008). Inferring the Tree of Life of the order Cypriniformes, the earth's most diverse clade of freshwater fishes: Implications of varied taxon and character sampling. *Journal of Systematics and Evolution*, Vol. 46, No. 3, pp. 424-438, ISSN 1759-6831
- Mecklenburg, C. W., Moller, P. R. & Steinke, D. (2011). Biodiversity of arctic marine fishes: taxonomy and zoogeography. *Marine Biodiversity*, Vol. 41, No. 1, pp. 109-140, ISSN 1867-1624
- Mendel, J., Lusková, V., Halačka, K., Lusk, S. & Vetešník, L. (2005). Genetic diversity of *Gobio gobio* populations in the Czech Republic and Slovakia, based on RAPD markers. *Folia Zoologica*, Vol. 54, No. 1, pp. 13-24, ISSN 0139-7893
- Mendel, J., Lusk, S., Halačka, K., Lusková, V., Vetešník, L., Čáleta M. & Ruchin, A. (2008b). Genetická diverzita a poznámky k výskytu ouklejky pruhované *Alburnoides bipunctatus* [Genetic diversity of Spirlin, *Alburnoides bipunctatus*, with notes on its occurrence]. In: *Biodiverzita ichtyofauny ČR (VII)*, S. Lusk, V. Lusková (Ed.), pp. 25-37, Ústav biologie obratlovců AV ČR, v.v.i, ISBN 978-80-87189-01-6, Brno. (in Czech with English summary)
- Mendel, J., Lusk, S., Koščo, J., Vetešník, L., Halačka, K. & Papoušek, I. (2008a). Genetic diversity of *Misgurnus fossilis* populations from the Czech Republic and Slovakia. *Folia zoologica*, Vol. 57, No. 1-2, pp. 90-99, ISSN 0139-7893
- Mendel, J., Lusk, S., Lusková, V., Koščo, J., Vetešník, L., Halačka, K. (2008c). Druhová pestrost hrouzků rodů *Gobio* a *Romanogobio* na území České republiky a Slovenska [The species diversity of gudgeon of the genera *Gobio* and *Romanogobio* in the territory of the Czech Republic and Slovakia], In: *Biodiverzita ichtyofauny ČR (VII)*, S.

- Lusk, V. Lusková (Ed.), pp. 17-24, Ústav biologie obratlovců AV ČR, v.v.i, ISBN 978-80-87189-01-6, Brno, (in Czech with English summary)
- Mendel, J., Lusk, S., Vasil'eva, E. D., Vasil'ev, V. P., Lusková, V., Ekmekci, F. G., Erk'akan, F., Ruchin, A., Koščo, J., Vetešník, L., Halačka, K., Šanda, R., Pashkov, A. N. & Reshetnikov, S. I. (2008d). Molecular phylogeny of the genus *Gobio* Cuvier, 1816 (Teleostei: Cyprinidae) and its contribution to taxonomy. *Molecular Phylogenetics and Evolution*, Vol. 47, No. 3, pp. 1061-1075, ISSN 1055-7903
- Messing, J. (1983). New M13 vectors for cloning. *Methods in Enzymology*, Vol. 101, pp. 20-78
- Meusnier, I., Singer, G. A. C., Landry, J. F., Hickey, D. A., Hebert, P. D. N. & Hajibabaei, M. (2008). A universal DNA mini-barcode for biodiversity analysis. *BMC Genomics*, Vol. 9, pp. 214, ISSN 1471-2164
- Monaghan, M. T., Balke, M., Gregory, T. R. & Vogler, A. P. (2005). DNA-based species delineation in tropical beetles using mitochondrial and nuclear markers In: *Philosophical Transactions of the Royal Society B: Biological Science*, Available from: doi:10.1098/rstb.2005.172.4
- Nwani, D. W., Becker, S., Braid, H. E., Ude, E. F., Okogwu, O. I. & Hanner R. (2011). DNA barcoding discriminates freshwater fishes from southeastern Nigeria and provides river system-level phylogeographic resolution within some species. *Mitochondrial DNA*, In Press
- Papoušek, I., Lusková, V., Koščo, J., Lusk, S., Halačka, K., Povž, M. & Šumer, Z. (2008a). Genetic diversity of *Cobitis* spp. (Cypriniformes: Cobitidae) from different drainage areas. *Folia Zoologica*, Vol. 57, No. 1-2, pp. 83-89, ISSN 0139-7893
- Papoušek, I., Lusková, V., Lusk, S. & Koščo, J. (2008b). Výskyt a genetická diverzita ježdíka žlutého a ježdíka dunajského v České republice. [Occurrence and genetic diversity of Stripped ruffe *Gymnocephalus schraetser* and Danube ruffe *Gymnocephalus baloni* in the Czech Republic], In: *Biodiverzita ichtyofauny ČR (VII)*, S. Lusk, V. Lusková (Ed.), pp. 88-95, Ústav biologie obratlovců AV ČR, v.v.i, ISBN 978-80-87189-01-6, Brno, (in Czech with English summary)
- Perea, S., Böhme, M., Zupančič, P., Freyhof, J., Šanda, R., Özuluğ, M., Abdoli, A. & Doadrio, I. (2010). Phylogenetic relationships and biogeographical patterns in Circum-Mediterranean subfamily Leuciscinae (Teleostei, Cyprinidae) inferred from both mitochondrial and nuclear data. *BMC Evolutionary Biology*. Vol. 10, pp. 265, ISSN 1471-2148
- Primack, R. B., Kindlmann, P. & Jersáková, J. (2001). Biological principles of nature conservation, Portál, ISBN 80-7178-552-0, Praha
- Pyle, R. L., Earle, J. L. & Greene, B. D. (2008). Five new species of the damselfish genus *Chromis* (Perciformes: Labroidae: Pomacentridae) from deep coral reefs in the tropical western Pacific. *Zootaxa*, Vol. 1671, pp. 3-31, ISSN 1175-5334
- Ratnasingham, S. & Hebert, P. D. N. (2007). BOLD: the Barcode of Life Data System (www.barcodinglife.org). *Molecular Ecology Notes*, Vol. 7, No. 3, pp. 355-364, ISSN 1471-8286
- Raup, D. M. (1992). *Extinction: bad genes or bad luck?* W. W. Norton & Company, ISBN: 978-0393309270, New York
- Rock, J., Costa, F. O., Walker, D. I., North, A. W., Hutchinson, W. F. & Carvalho, G. R. (2008). DNA barcodes of fish of the Scotia Sea, Antarctica indicate priority groups for




- taxonomic and systematic focus. *Antarctic Science*, Vol. 20, No. 3, pp. 253–262, ISSN 1365-2079
- Schindel, D. E. & Miller, S. E. (2005). DNA barcoding a useful tool for taxonomists. *Nature*, Vol. 435, pp. 17, ISSN 1476-4687
- Seifertová, M., Vyskočilová, M., Morand, S. & Šimková, A. (2008). Metazoan parasites of freshwater cyprinid fish (*Leuciscus cephalus*): testing biogeographical hypotheses of species diversity. *Parasitology* Vol. 135, No. 12, pp. 1417–35, ISSN 0031-1820.
- Steinke, D., Zemlak, T. S. & Hebert, P. D. N. (2009). Barcoding Nemo: DNA-Based identifications for the ornamental fish Trade. *PLoS One*, Vol. 4, No. 7, pp. e6300, ISSN 1932-6203
- Šedivá, A., Janko, K., Šlechtová, V., Kotlík, P., Simonović, P. Delic, A. & Vassilev, M. (2008). Around or across the Carpathians: colonization model of the Danube basin inferred from genetic diversification of stone loach (*Barbatula barbatula*) populations. *Molecular Ecology*, Vol. 17, No. 5, pp. 1277–1292, ISSN 0962-1083
- Thalmann, O., Hebler, J., Poinar, H. N., Pääbo, S. & Vigilant, L. (2004). Unreliable mtDNA data due to nuclear insertions: a cautionary tale from analysis of humans and other great apes. *Molecular Ecology*, Vol. 13, No. 2, pp. 321–335, ISSN 0962-1083
- Valdez-Moreno, M., Ivanova, N. V., Elías-Gutiérrez, M., Contreras-Balderas, S. & Hebert, P. D. N. (2009). Probing diversity in freshwater fishes from Mexico and Guatemala with DNA barcodes. *Journal of Fish Biology*, Vol. 74, No. 2, pp. 377–402, ISSN 1095-8649
- Vences, M., Thomas, M., van der Meijden, A., Chiari, Y. & Vieites, D. R. (2005). Comparative performance of the 16S rRNA gene in DNA barcoding of amphibians. *Frontiers in Zoology*, Vol. 2, pp. 5, ISSN 1742-9994
- Vetešník, L., Papoušek, I., Halačka, K., Lusková, V. & Mendel, J. (2007). Morphometric and genetic analysis of the *Carassius auratus* complex from an artificial wetland in floodplain of the Morava River (Czech Republic). *Fisheries Science*, Vol. 73, No. 4, pp. 817–822, ISSN 1444-2906
- Ward, R. D., Zemlak, T. S., Innes, B. H., Last, P. R. & Hebert, P. D. N. (2005). DNA barcoding Australia's fish species. *Philosophical Transactions of the Royal Society B: Biological Science*, Vol. 360, pp. 1847–1857, ISSN 1471-2970
- Ward, R. D., Holmes, B. H. & Yearsley, G. K. (2008). DNA barcoding reveals a likely second species of Asian seabass (barramundi) (*Lates calcarifer*). *Journal of Fish Biology*, Vol. 72, No. 2, pp. 458–463, ISSN 1095-8649
- Ward, R. D., Hanner, R. & Hebert, P. D. N. (2009). The campaign to DNA barcode all fishes, FISH-BOL. *Journal of Fish Biology*, Vol. 74, No. 2, pp. 329–356, ISSN 1095-8649
- Wilson, E. O. (1989). Threats to biodiversity. *Scientific American*, Vol. 261, pp. 108–116, ISSN 0036-8733
- Wilson, E. O. (1992). *The Diversity of Life*. The Belknap Press of Harvard University Press, ISBN 978-0393319408, Cambridge
- Zardoya, R. & Doadrio, I. (1999). Molecular Evidence on the Evolutionary and Biogeographical Patterns of European Cyprinids. *Journal of Molecular Biology*, Vol. 49, pp. 227–237, ISSN 0022-2836



Zhang, J. & Hanner, R. (2011). DNA barcoding is a useful tool for the identification of marine fishes from Japan. *Biochemical Systematics and Ecology*, Vol. 39, No. 1, pp. 31–42, ISSN 0305-1978

IntechOpen

IntechOpen



## **Analysis of Genetic Variation in Animals**

Edited by Prof. Mahmut Caliskan

ISBN 978-953-51-0093-5

Hard cover, 360 pages

**Publisher** InTech

**Published online** 29, February, 2012

**Published in print edition** February, 2012

Analysis of Genetic Variation in Animals includes chapters revealing the magnitude of genetic variation existing in animal populations. The genetic diversity between and within populations displayed by molecular markers receive extensive interest due to the usefulness of this information in breeding and conservation programs. In this concept molecular markers give valuable information. The increasing availability of PCR-based molecular markers allows the detailed analyses and evaluation of genetic diversity in animals and also, the detection of genes influencing economically important traits. The purpose of the book is to provide a glimpse into the dynamic process of genetic variation in animals by presenting the thoughts of scientists who are engaged in the generation of new idea and techniques employed for the assessment of genetic diversity, often from very different perspectives. The book should prove useful to students, researchers, and experts in the area of conservation biology, genetic diversity, and molecular biology.

### **How to reference**

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Jan Mendel, Eva Marešová, Ivo Papoušek, Karel Halačka, Lukáš Vetešník, Radek Šanda, Milena Koničková and Soňa Urbánková (2012). Molecular Biodiversity Inventory of the Ichthyofauna of the Czech Republic, Analysis of Genetic Variation in Animals, Prof. Mahmut Caliskan (Ed.), ISBN: 978-953-51-0093-5, InTech, Available from: <http://www.intechopen.com/books/analysis-of-genetic-variation-in-animals/molecular-biodiversity-inventory-of-the-ichthyofauna-of-the-czech-republic>

**INTECH**  
open science | open minds

### **InTech Europe**

University Campus STeP Ri  
Slavka Krautzeka 83/A  
51000 Rijeka, Croatia  
Phone: +385 (51) 770 447  
Fax: +385 (51) 686 166  
[www.intechopen.com](http://www.intechopen.com)

### **InTech China**

Unit 405, Office Block, Hotel Equatorial Shanghai  
No.65, Yan An Road (West), Shanghai, 200040, China  
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元  
Phone: +86-21-62489820  
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](https://creativecommons.org/licenses/by/3.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen