We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Multidrug Resistance and Breast Cancer

Gengyin Zhou and Xiaofang Zhang
Shandong University of Medicine,
China

1. Introduction

Millions of new cancer patients are diagnosed each year and over half of these patients die from this disease. As the second leading cause of cancer deaths, breast cancer is estimated to be diagnosed in over one million people worldwide and to cause more than 400,000 deaths each year [1]. Chemotherapy is part of a successful treatment to many cases; however, the development of multidrug resistance (MDR) to it becomes a major obstacle so as to as few as half of the breast cancer patients treated benefit from chemotherapy.

MDR is a term used to describe the phenomenon characterized by the ability of drug resistant tumors to exhibit simultaneous resistance to a number of structurally and functionally unrelated chemotherapeutic agents [2]. The cytotoxic drugs that are most frequently associated with MDR are hydrophobic, amphipathic natural products, such as the taxanes (paclitaxel and docetaxel), vinca alkaloids (vinorelbine, vincristine, and vinblastine), anthracyclines (doxorubicin, daunorubicin, and epirubicin), epipodophyllotoxins (etoposide and teniposide), antimetabolites (methorexate, fluorouracil, cytosar, 5-azacytosine, 6-mercaptopurine, and gemcitabine) topotecan, dactinomycin, mitomycin C and so on[3].

At present, many mechanisms have been found to be responsible for it, including overexpression of the members of the adenosine triphosphate (ATP)-binding cassette (ABC) membrane transporter family, changes of apoptosis-related genes, the alteration of DNA-repair gene, cancer stem cells and so on. And up to date, many methods were adopted to overcome MDR, for example natural drugs, chemical drugs and genetic therapy.

Herein, we will introduce the mechanisms and therapy of MDR of breast cancer briefly.

2. Mechanisms of MDR

2.1 The adenosine triphosphate (ATP)-binding cassette (ABC) membrane transporter family

Elevated expression of ATP-binding cassette (ABC) transporters is considered to be the main cause of MDR in breast cancer. ATP-binding cassette (ABC) transporters are a family of transporter proteins that contribute to drug resistance via adenosinetriphosphate (ATP)-dependent drug efflux pumps, which can result in an increased efflux of the cytotoxic drugs from the cancer cells, thus lowering their intracellular concentrations [4]. Up to date, more than 100 ABC transporters from prokaryotes to humans and 48 human ABC genes have been identified that share sequence and structural homology [3]. The proteins which are related to the MDR in breast cancer are mainly including p-glycoprotein (p-gp), multidrug resistance-related protein (MRP) and breast cancer resistance protein (BCRP).

In mammals, the functionally active typical ABC proteins consist of at least four core domains, two transmembrane domains (TMDs) and two nucleotide-binding domains (NBDs). The two TMDs of each ABC transporter consist of multiple membrane-spanning α -helices (typically, but not always, six α -helices per domain) and form the pathway through which substrate crosses the membrane. The two NBDs play a role in cleaving ATP (hydrolysis) to derive energy necessary for transporting cell nutrients such as sugars, amino acids, ions and small peptides [5]. Normally, they have important physiological function, such as the excretion of toxins from the liver, kidneys, and gastrointestinal tract [6]. Overexpression of these transporters has been observed in many types of human malignancies and correlated with poor responses to chemotherapeutic agents.

2.1.1 P-glycoprotein

In 1970s, a carbohydrate-containing protein, 170 kDa in molecular weight, was found in multidrug-resistant Chinese hamster ovary cells. The glycoprotein was named P-glycoprotein (P-gp) because the protein can modulate membrane permeability with respect to a number of apparently unrelated drugs including actinomycin D, methotrexate, daunomycin, and colchicine [7]. The MDR mediated by P-gp is also called "classical MDR". Gene sequence membrane analysis for mammalian P-gp has revealed the presence of two similar halves, each containing 6 putative transmembrane segments, and an ATP-binding consensus motif. The human protein is comprised of 1280 amino acids with 12 transmembrane domains and 43% sequence homology between the two halves. Three glycosylation sites on the first extracytoplasmic domain are present [8]. The gene encoding p-glycoprotein was termed *mdr1*. The gene with 28 exon and 1.2 kb and is located on chromosome 7q21.12.

There are three known isoforms of P-gp, namely, class I, II and III. Rodent cells have all three P-gp genes, whereas human cells only have class I and III P-gp [9]. Classes I and II P-gp genes confer MDR when transfected into sensitive wild type (WT) cells, whereas the class III P-gp gene is not shown to be associated with drug resistance. All three types of P-gp expressed in several normal tissues. In mammalian tissues, class I P-gp is found in epithelium, intestinal, endothelial cells, bone marrow progenitor peripheral blood lymphocytes, natural killer cells and so on. The class III P-GP is localized in hepatocytes, cardiac and striated muscle [10]. The distribution displays that p-gp plays an important role in normal physiological function. Evident confirmed that P-gp take part in the transepithelial secretion of substrates into bile, urine, or gastrointestinal tract lumen. P-gp may also confer a protective role to mediate xenobiotic efflux in tissues such as the brain, testis, and placenta.

P-gp substrates are widespread. Although their structures are very different, they share many physical properties including high hydrophobicity, an amphiphilic nature and a net positive charge [11]. It is important for us to understanding the modulation of P-gp. Evident shows that p-gp is phosphorylated by protein kinase C (PKC) and PKC blockers can reduce P-gp phosphorylation and increase drug accumulation. However, there is evidence that PKC inhibitors directly interact with P-gp and inhibit drug transport by a mechanism independent of P-gp phosphorylation [12]. Experiments using transient transfection of the MDR1 promoter (linked to a reporter gene) into the cells as well as stable transfection of some other genes showed that genes p53, ras and raf can influence the activity of introduced MDR1 promoter or the expression of the endogenous cellular MDR1. Genes c-fos and c-jun also were shown to confer the regulation of MDR1 activity [13].

For a long time, P-gp was believed to be the only protein capable of conferring MDR in mammalian tumor cells. Over 50% breast cancer expressed P-gp [14]. Moreover, prior exposure to chemotherapy or hormonal therapy has been shown to increase the proportion of breast cancers expressing P-gp by 1.8-fold [15]. However, pre-chemotherapy P-gp expression showed no association with shorter progression-free survival (PFS) so the clinical relevance of this observation in terms of screening patients and treatment selection remains unclear[16].

2.1.2 Multidrug resistance- related protein 1(MRP)

In 1992, Susan Cole and Roger Deeley observed amplification and increased expression of a novel gene in non-P-gp expressing small cell lung cancer DOX resistant cell lines and this is the MRP1 (ABCC1) (MDR related protein) gene. The following study shows that the protein encoded by this gene is also a member of ABC transporters [17].

The multidrug-resistance-associated protein (MRP or MRP1) is a 190 kDa protein and is constituted by 1531 amino acids. Like other members of ABC transporters, MRP1 has 3 membrane spanning domains, 2 NBDs and extracellular N-terminal. Up to now, several isoforms of MRP1 have been identified. Included among these are five human MRP1-related proteins, designated MRP2, MRP3, MRP4, MRP5 and MRP6. MRP7, MRP8 and MRP9 are recent additions to the family which have not yet been characterised[18].

Physiologically, MRP1 also plays a normal role in the ATP-dependent unidirectional membrane transport of glutathione conjugates, such as leukotriene C₄, S-(2,4-dinitrophenyl)glutathione and leukotriene receptor antagonists could inhibit this function[19]. Besides multidrug-resistance cancer cells, MRP is also expressed in normal human tissues, such as muscle, lung, spleen, bladder, adrenal gland and gall bladder [2]. MRP2 (or canalicular multispecific organic anion transporter or cMOAT) was first shown to be expressed in the liver which functions in the excretion of glutathione and glucuronate conjugates across the canalicular membrane into bile. In addition, MRP2/ cMOAT is also expressed in the human kidney proximal tubule epithelia on the apical side. Therefore, it is implicated that MRP2 may play a role in the renal excretion of endogenous substances and xenobiotics, in normal conditions. MRP3 is expressed in liver and involved in the efflux of organic anions from the liver into the blood in case of biliary obstruction. MRP4 and MRP5 transport nucleosides and confer resistance to antiretroviral nucleoside analogs. MRP6 is a lipophilic anion pump with a wide spectrum of drug resistance. Among the members of MRP family, only MRP1 has been widely accepted to cause clinical drug resistance[3].

Like other members of ABC transporters, MRP1 can pump anti-tumor drugs out of the tumor cells, cause reduced intracellular accumulation of drugs and lead to resistance. Whereas P-gp transports neutral and positively charged molecules in their unmodified form, MRP1 overexpression is associated with an increased ATP-dependent glutathione-S conjugate transport activity. Reduced glutathione (GSH) has been suggested as an important component of MRP mediated MDR and drug transport. MRP1 is able to transport a range of substrates as such or conjugated to GSH, glucuronide, and sulfate [20]. The anticancer drugs that are substrates of MRP1 mainly include anthracyclines such as doxorubicin and daunorubicin, vinca alkaloids and etoposide. Several findings indicate that MRP1 reduces drug accumulation by effluxing drugs by a GSH co-transport mechanism or after their conjugation to GSH [21]. But the mechanism by which GSH facilitates transport of some compounds by MRP1 is still a matter of debate.

2.1.3 Breast cancer resistance protein (BCRP)

Breast cancer resistance protein (BCRP) is the latest ABC transporter involved in MDR and it was cloned by Ross and Doyle in 1998 from a mitoxantrone-resistant subline of the breast cancer cell line MCF-7/Adr/Vp which does not express other known multidrug efflux transporters such as P-glycoprotein (P-gp) or the multidrug resistance protein 1 (MRP1) [22]. Two almost identical proteins as BCRP with only a few amino acid differences were later discovered independently by other laboratories from mitoxantrone-resistant human cancer cell lines (so named as MXR) and humanplacenta (so named as ABCP)[23].

The BCRP gene is located on chromosome 4q22. The full length of BCRP gene is 66kb and the length of mRNA is about 2.4kb [23]. The product of the gene is a 72KD protein with 655 amino acid that contains an ATP-binding domain and six transmembrane domains, and it is a half transporter member of the ABCG subfamily [24]. As a half transporter, BCRP functions as a homodimeric/oligomeric efflux pump [25], and in a manner that is similar to other ABC transporters. Besides that, BCRP can also transport hydrophilic conjugated organic anions, particularly the sulfated conjugates with high affinity, for example BCRP can detoxify irinotecan and SN-38 by glucuronidation via the activity of UDP-glucuronyltransferase [26]. BCRP substrates include not only chemotherapeutic agents such as mitoxantrone, methotrexate, topotecan, irinotecan and its active analog SN-38, and tyrosine kinase inhibitors imatinib and gefitinib, but non-chemotherapy drugs such as prazosin, glyburide, nitrofurantoin, dipyridamole, statins, and cimetidine as well as nontherapeutic compounds such as the dietary flavonoids, porphyrins, estrone 3-sulfate (E1S), and the carcinogen PhIP [27].

Similar to P-gp and MRP1, BCRP is widely expressed in normal cells and tissues including the capillary endothelial cells, the hematopoietic stem cells [28], the maternal - fetal barrier of the placenta and the blood-brain barrier [29]. In these tissues, BCRP play a protective role against xenobiotics and their metabolites. Whereas, the apical localization of BCRP in the intestinal epithelium and in the bile canalicular membrane also suggests the intestinal absorption and hepatobiliary excretion of BCRP substrates [30].

Unexpectedly, many mutant forms of BCRP proteins were found in drug-selected cells such as those of the S1-M1-80 and MCF7/AdVp3000 cell lines and up to now, more than 50 mutations have been identified including natural variants and non-natural mutations [27]. The most important natural variant is Q141K, which occurs in Japanese and Chinese populations at high allele frequencies (30 -60%) and in Caucasians and African-American populations at relatively low allele frequencies (5 - 10%) [27]. Several studies consistently revealed that Q141K had a lower protein expression level than wild-type BCRP in both transfected cells and human tissues. A recent study has revealed that Q141K undergoes increased lysosomal and proteasomal degradations than wild-type BCRP, possibly explaining the lower level of protein expression of the variant [31]. The R482T and R482G variants of BCRP detected from MCF7/ AdVp3000 and S1-M1-80 cells belong to non-natural mutants. The non-natural mutants have different effects on BCRP expression, distribution and functions. Some mutations do not affect plasma membrane expression, but alter substrate specificity and/or overall transport activity. For example, the R482T and R482G lose their methotrexate-transporting activity but at the same time confer increased mitoxantrone resistance, so they are highly resistant to both mitoxantrone and doxorubicin[32]. Wild-type BCRP does not transport Rhodamine 123 and Lyso-Tracker Green; however, the mutants R482T and R482G do [33]. These findings confirmed that the

transmembrane region of BCRP plays important roles in its activity. Some mutations affect biogenesis with decreased stability, lower expression and/or altered subcellular distribution of BCRP. A typical example is mutations of Arg383 which results in a significant decrease in the protein level, partial retention in the endoplasmic reticulum, and altered glycosylation and the treatment with mitoxantrone assisted in protein maturation [34]. Some mutations influence the chemical modifications of BCRP such as N-linked glycosylation or disulfide bond formation in BCRP such as the mutation Asn596. Otherwise, there are also many mutations which do not have major effects on both plasma membrane expression and function of BCRP, including K473A and H630X. The research on the mutations of BCRP could help us to further understand the structures and functions of ABC transporters.

Up to date, BCRP was detected in many resistance tumor cells such as human colon cancer cell line S1-M1-80, prostate cancer cell lines and breast cancer cell line MCF7/AdVp3000 [35]. Many clinical sample were also found BCRP expression, including acute myelogenous leukemia (AML), acute lymphocytic leukemia (ALL), non-small cell lung cancer and so on [36-38]. And it has been suggested that the expression of BCRP is associated with a poor response to cancer chemotherapy and may be responsible for clinical drug resistance. However, the studies on the expression characters of BCRP in breast cancer clinical samples are still very few.

2.2 Apoptosis and MDR

2.2.1 P53

As a tumor suppressor, p53 plays a pivotal role in inducing apoptosis in response to cellular damage, including DNA damage. However, mutant P53 plays an opposite role in the regulation of apoptosis, that is mutant p53 is an anti-apoptosis factor. In a study from the National Cancer Institute (NCI), the majority of breast cancer cell lines were mutant for p53 [39]. About 50% of all tumours have an approximately 25% occurrence of deletions and point mutations in sporadic breast cancers [40]. Many anti-tumor drugs can lead cellular death by inducing cellular apoptosis. When p53 mutations or deletions occur, the cellular apoptosis can be inhibited and the cells exhibit MDR phenotype. Mutations in p53 have been verified to be related with resistance to doxorubicin in breast cancer patients [41].

Many data show the correlation of P53 and ABC transporters. The first experiments implied that a mutant p53 (mtP53) specifically stimulated the MDR1 promoter and wild-type p53 (wtP53) exerted specific repression [42]. In the follow-up study, a p53 consensus binding sequence was also found in the promoter of the rat ABCB1 gene. Both promoter function and endogenous *mdr1b* expression were shown to be up-regulated by wtp53 [43]. More studies displayed that the mutations of p53 can dramatically activate the ABCB1 promoter in multiple cell lines including Saos-2, Caco-2, MCF-7 and so on [44]. Linn et al. assessed the status of p53 and ABCB1 in both primary operable and advanced-staged tumors and their results revealed that nuclear p53 accumulation and coexpression of ABCB1 were more prevalent in locally advanced breast cancers and that these markers provided a strong prognostic indication of shorter survival [45,46]. Similar results were also found in other studies.

Recently, Wang et al investigated the effects of wild-type and mutant p53, and nuclear factor kappa-B (NF-kappaB) (p50) on BCRP promoter activity in MCF-7 cells, and the results show that wild-type p53 induced transcriptional suppression of breast cancer resistance protein (BCRP) through the NF-kappaB pathway in MCF-7 cells [47].

2.2.2 Other apoptosis related genes and MDR

Mitochondrial (intrinsic pathway) and cell surface receptor (Fas) mediated (extrinsic pathway) apoptosis are the two main routes leading to programmed cell death. MCF-7 cells can undergo apoptosis by the sequential activation of caspases-9 (associated with mitochondrial mediated apoptosis), -7, and -6. Recently, a splice variant form of caspase-3 has been shown to be overexpressed in chemoresistant, locally advanced breast cancers, and is particularly associated with response to cyclophosphamide[48].

Bcl-2 is a member of a large family of genes coding both anti-apoptotic proteins (for example, Bcl-2, Bcl-XL) and pro-apoptotic proteins (Bax, Bad, Bcl, etc.). Bcl-2 protein is able to inhibit the apoptosis induced by p53 in response to genotoxic stress. There are data showing that Bcl-2 overexpression results in the resistance of cells to different drugs, including DOX, taxol, etoposide, camptothecin, mitoxantrone and cisplatin[49]. When Bcl-2 is over expressed and contributes as a resistance mechanism, it has been shown that the anticancer drugs promote cell cycle arrest; however, their effects are cytostatic rather than cytotoxic[2]. The phosphorylation state of the Bcl-2 oncoproteins has been shown to modulate response to taxanes[50].

Survivin is another apoptosis-related gene which has been confirmed to confer MDR in tumors. It is a structurally unique inhibitor of apoptosis (IAP), substances which block apoptosis induced by a variety of nonrelated apoptosis triggers. Survivin is known to directly or indirectly bind and inhibit the terminal effector cell death protease cascades, caspase 3 and 7, as well as inhibit the activation of caspase 9[51]. Furthermore, it has been reported that the expression of survivin was significantly higher after treatment with anti-cancer drugs in many cancer cells and may be involved in radio- and chemo-resistance[52]. Liu et al. documented that survivin might modulate the turnover of P-gp or transport by P-gp in the cell, which then resulted in anti-apoptosis and drug resistance in breast cancer cells[51]. However, the role of survivin in MDR breast cancer in the presence of P-gp is still not clear.

In addition, some other apoptosis-related genes were found to take part in the regulation of MDR, such as CD95, TRAIL and so on.

2.3 MDR-related enzyme

2.3.1 Glutathione S-transferase (GST)

GST is a member of phase II detoxification enzymes that catalyses the conjugation of glutathione (GSH) to a wide variety of endogenous and exogenous electrophilic compounds. Because of their capacity to react with electrophiles, radicals and reactive oxygen species, GSTs, together with GSH, have a major role in the protection against oxidative stress [53].

GSTs are divided into two super-family members: the membrane microsomal and cytosolic GSTs(c-GSTs). Microsomal GSTs (m-GSTs) are structurally distinct from the cytosolic in that they homo- and heterotrimerize rather than dimerize to form a single active site and the microsomal GSTs are mainly involved in the metabolism of endogenous compounds, like leukotrienes and prostaglandins. The cytosolic GSTs also conjugate exogenous compounds and the cytosolic GSTs are subject to significant genetic polymorphisms in human populations. Up to date, the cytosolic GSTs are divided into seven classes, Alpha (A), Mu (M), Omega (O), Pi (P), Sigma (S), Teta (T) and Zeta (Z) which have a promiscuous substrate specificity and are localized in different tissues with organ specific expression patterns [54]. The GST-Pi have been confirmed to be closely related with MDR.

Many data show that GST confers the development and expression of MDR. Increased expression of GSTpi—detected as strong immunoreactivity—has been documented to contribute to drug resistance of ovarian carcinomas, head and neck cancer, lung squamous-cell carcinoma, breast cancers and so on[55]. Cells with GST isozyme transfections yield mild increases in resistance (mostly in the 2-5 fold range) to a number of different anticancer drugs [54]. While inhibition of GST expression by antisense cDNA increased the sensitivity to several anticancer drugs [56]. Besides, exposure of cells to a specific inhibitor of c-GCS, buthionine sulfoximine (BSO), decreases multidrug resistance to doxorubicin and vincristine[57]. The substrates of GST reported include chlorambucil, melphalan, nitrogen mustard, phosphoramidate mustard, acrolein, carmustine, hydroxyalkenals, ethacrynic acid and steroids. And the MDR mediated by GST is related to mitomycin C, adriamycin, cisplatin and carboplatin.

How GSTs affect MDR in tumor cells? There are mainly two mechanisms found now. First, GST-Pi plays a key role in regulating the MAP kinase pathway via protein: protein interactions. GST-Pi was shown to be an endogenous inhibitor of c-Jun N-terminal kinase 1 (JNK1), a key member of MAP pathway which involved in stress response, apoptosis, and cellular proliferation [58]. In nonstressed cells, low JNK activity is observed due to the sequestration of the protein in a GST- Pi : JNK complex. Direct protein : protein interactions between the C-terminal of JNK and GST-pi were reported with a binding constant of approximately 200 nm. The second, there is a coordinate action of phase II enzymes and MRP in MDR[59]. The already mentioned connection between a MRP drug resistance profile and an increased GST-pi expression, shown in many cell lines, is indeed indicative for a shared regulatory mechanism of MRP and GSTs expression [60]. Studies demonstrated that Nrf2 may play a key role between MRP and GSTs. A study on Nrf2 knockout mice displayed that: disruption of the Nrf2 gene decreased both the constitutive as well as the inducible expression of class Alpha, class Mu and class Pi glutathione transferases[61]. Meanwhile, Nrf2 was also shown to be necessary for the constitutive and inducible expression of MRP1 in mouse embryo fibroblast[62].

GSTpi immunoreactivity was reported not to correlate with response to chemotherapy in cervical carcinoma, but many data show that in primary breast cancers, expression of GST-Pi was associated with poor prognosis. Fengxi Su et al analyzed the relationship between GST-Pi and the FAM (5-fluorouracil, adriamycin, mitomycin) protocol and the result showed that the presence of GSTpi in breast cancer tissue was a bad prognostic indicator, and these tumors were largely resistant to chemotherapy[55]. In cultured breast cancer cells, GST-pi is exclusively expressed in estrogen receptor-negative (ER-) cells but not in receptor-positive (ER+) cells [63]. In 1997, Mona S. Jhaveri verified that that methylation status of the promoter contributes significantly to the levels of GSTP1 expressed in ER- and ER+ breast cancer cell lines [64].

2.3.2 DNA topoisomerase II (topo II)

DNA topoisomerase II (topo II) is a nuclear phosphoprotein involved in DNA replication and chromosome dynamics. These enzymes catalyse the ATP-dependent passage of one DNA duplex (the transport or T-segment) through a transient, double-stranded break in another (the gate or G-segment), navigating DNA through the protein using a set of dissociable internal interfaces, or 'gates' [65,66]. The family of DNA topoisomerase II includes two related but genetically distinct isoforms isforms TOPII α and II β in mammalian cells.

The human topoisomerase II α gene (TOP2 α) is localized on chromosome 17q21-22 [67] whereas TOP2 β maps to chromosome 3p24 [68]. The cDNAs for the human α and β isoforms encode p170 and p180 proteins of 1531- and 1621-amino-acid [68], respectively. TOP2 α lies close to the epidermal growth factor-like receptor gene ERBB2 (HER2) and the retinoic acid receptor locus RAR in a region of chromosomal 17 which is amplified in some human breast cancer [69]. The two enzymes are closely similar in structure each comprising three functional domains defined by sites of cleavage by trypsin or staphylococcal V8 proteases: an N-terminal ATPase domain (approximately residues 1-400); a DNA breakage-reunion region (400-1220); and the C-terminal domain which carries a multitude of phosphorylation sites [70].

In addition to its role in cell division, TOP2 α is also found to be related to the MDR in tumor. It is the major molecular target for a large group of clinically relevant, structurally different cytotoxic agents known as TOP2 α inhibitors including the anthracycline class of antitumor cytotoxic agents [71]. These drugs all act by forming covalent bonds with TOP2 α , creating a complex that introduces permanent double-strand breaks in DNA leading to apoptosis. Reduced topoisomerase II expression or function can contribute to resistance to agents such as anthracyclines and epipodophyllotoxins [72]. In vitro studies of breast tumor cell lines have shown that amplification of the TOP2 α gene leads to protein overexpression and sensitivity to anthracyclines [73,74]. Similarly, deletion of TOP2 α genomic alterations in breast cancer leads to a marked decrease in TOP2 α protein expression, which results in chemoresistance to TOP2 α inhibitor anticancer drugs in cell culture.

The HER-2 gene is another gene on chromosome 17 and it encodes for a ligandless, transmembrane glycoprotein receptor with intrinsic tyrosine-kinase activity. HER-2 gene amplification or protein overexpression occurs in about 20% of patients with breast cancer and is a recognized poor prognostic marker, often associated with endocrine resistant, high grade disease [75]. Recently research reported that the expression of TOP2 α is closely related to the expression of HER2 gene and co-expression of them may be a useful tool in predicting benefit from chemotherapy. Top II α is reported to be either amplified or deleted in nearly 90% of HER-2 amplified primary breast cancers [76]. Recent review of the Canadian-MA.5 trial assessed TOP2 α alterations and HER-2 amplification by FISH on tissue microarrays in 438 patients [77]. Top II α alterations occurred in 18% patients (12% amplification, 6% deletions) and were more common in large tumors and in HER-2 positive tumors. In patients with Top II α alterations, relative benefits of therapy were seen with CEF having statistical superiority over CMF in terms of RFS (adjusted HR 0.35, 95% CI 0.17 - 0.73, $p = 0.005$) and OS (adjusted HR 0.33, 95% CI 0.15 - 0.75, $p = 0.008$). However, there are also diffusing evidents. An analysis displayed topo II α mRNA overexpression in 19% of HER-2 negative patients [78]. In conclusion, the relationship between TOP II, HER2 and chemosensitivity needs further investigation.

2.3.3 Glucosylceramide synthase

Sphingolipids, which include ceramides and sphingosine, were first isolated and characterized in the late 1800s. Recent years, many studies have shown that they are not only structural and insert components of cell membranes but also associated with myriad process of cells including the proliferation, survival and death of cells. As an important member of sphingolipid metabolism, ceramide have been proven to be a second messenger of apoptosis [79, 80]. Cellular stress is known to increase ceramide levels in cells. So it is easily

to understand that increased ceramide has been observed in response to many anti-cancer drugs, such as doxorubicin, vincristine, paclitaxel, etoposide, PSC 833 and fenretinide.

Many enzymes have been confirmed to be responsible for the regulation of ceramide levels, such as ceramide synthase and sphingomyelinase which are responsible for the ceramide generation, and sphingomyelin synthase and ceramidase which take part in the ceramide metabolism [81]. Glucosylceramide synthase (GCS) is one of them. As an enzyme which catalyzes the first step in glycosphingolipid synthesis, GCS transfers UDP-glucose to ceramide to form glucosylceramide, which have been found to involve in many cellular processes such as cell proliferation, oncogenic transformation, differentiation, and tumor metastasis [82]. In addition, many studies show that GC was related with MDR in many tumor cells. In 1996, Lavie Y et al first reported that chemotherapy resistant MCF-7-AdrR breast cancer cells accumulate GC in comparison to wild-type MCF-7 cells [83]. After that, GC was found to confer to MDR in many other cancers [84-86]. So some people guessed that elevated GCS activity may be a novel form of multidrug resistance.

Then, Liu et al found that increased competence to glycosylate ceramide conferred adriamycin resistance in MCF-7 breast cancer cells by transfection with GCS cDNA [8], while using GCS inhibitor 1-phenyl-2-palmitoylamino-3-morpholino-propanol (PPMP) or transfection of doxorubicin-resistant MCF-7-AdrR cells with GCS antisense both restored cell sensitivity to doxorubicin or vinblastine and paclitaxel [86,87]. Ladisch found that blocking GCS with D, L-threo-phenyl-2-hexadecanoylamino-3-pyrrolidino-1-propanol (PPPP), was able to elevate ceramide levels and enhance vincristine cytotoxicity via programmed cell death [88]. All the following works demonstrated that GCS was potentially one MDR-related drug resistance mechanism.

Recently, Yong-Yu Liu et al reported that glucosylceramide synthase upregulates MDR1 expression in the regulation of cancer drug resistance through cSrc and β -catenin signaling in the ovarian cell line NCI/ADR-RES which was ever named MCF-7/AdrR [89]. This study revealed the importance of GCS in the mechanism of cancer drug resistance.

Further studies demonstrated that a GC-rich/Sp1 promoter binding region was of importance in the regulation of GCS expression and doxorubicin could induce activation of Sp1 and up-regulation of GCS and apoptosis in Leukemia drug-resistance cell line HL-60/ADR and ovarian cell line NCI/ADR-RES [81,90].

In 2009, Eugen Ruckhäberle et al analyzed microarray data of GCS expression in 1,681 breast tumors and found that expression of GCS was associated with a positive estrogen receptor (ER) status, lower histological grading, low Ki67 levels and ErbB2 negativity ($P < 0.001$ for all) [91]. This study revealed the expression profile of GCS in breast cancer. But, the study also found that GCS has no clearly correlation with *mdr1*. So the relationship between GCS and *mdr1* in breast cancer is still a puzzle.

2.4 Cancer stem cells and MDR

Stem cells are defined as cells with both self-renewal capacity and the ability to produce multiple distinct differentiated cell types to form all the cell types that are found in the mature tissue [92]. Thus, these two characteristics of stem cells confer the unique property of asymmetric division. Stem cells are quiescent or slowly cycling cells maintained in an undifferentiated state until normal functioning of the organism needs their participation. Stem cells are classified into two principal types: embryonic and adult stem cells [93].

Recent studies have revealed that they play important role in cancer biology. Cancer stem cells (CSC) have been detected in many tumors, such as retinoblastoma and melanoma [94,95]. In breast cancer, a CD44+/CD24-or low/Lin- cell population was first identified as CSC [96]. Later, aldehyde dehydrogenase (ALDH) 1 activity was reported to be associated with stem/progenitor properties in breast cancer [97].

Although the origin of cancer stem cells has not yet been elucidated, researchers proposed that the malignant transformation of a normal stem cell, or a progenitor cell that has acquired self-renewal ability may be the reason. Up to now, Three major pathways have been identified to be related with the regulation and maintenance of stem cells in adult life: Wnt, Hedgehog, and Notch[92].

It is well known that cancer chemotherapy targets dividing cells. Because stem cells are quiescent or slowly cycling cells under normal situation, it is easy to understand that cancer stem cells could escape from the killing of anti-tumor drugs.

Besides, the side-population (SP) cells may be another reasons why stem cells become multi-drug resistant. The isolation of SP cells is based on the technique described by Goodell et al. in 1996 [98]. While experimenting with staining of murine bone marrow cells with the vital dye, Hoechst 33342, they discovered that the display of Hoechst fluorescence simultaneously at two emission wavelengths (red 675 nm and blue 450 nm) localizes a distinct, small, nonstained cell population (0.1% of all cells) that express stem cells markers (Sca1+linneg/low), which were named SP cells. At first, they thought the exclusion of Hoechst 33342 by SP cells is an active process involving multidrug resistance transporter 1 (MDR1). But the following study show that MDR1 can not be taken as a single marker to identify and isolate SP cells. Zhou et al. have demonstrated the breast cancer resistance protein (BCRP) may also attend the SP phenotype[28]. The SPs from breast cancer contain primitive stem cell-like cells that can differentiate into epithelial tumors in vitro and in vivo and express stemness genes [28,99]. The characterization of cells within SP demonstrates that they are immature, poorly differentiated, and highly tumorigenic. Gene expression profiles of SP show that these cells are less differentiated than non-SP cells [100].

The ABC transporters may play three functions in CSCs. First, the ABC transporters can protect the CSCs against exogenous products able to penetrate the cell membrane barrier by active exclusion. Second, there is mounting speculation that ABC transporters repress the maturation and differentiation of stem cells. For example, the overexpression of ABCG2 inhibits hematopoietic development. The last, protection from hypoxia appears to be another function of ABC transporters in CSCs[101].

In conclusion, although the mechanisms of cancer stem cell are still unclear, the cancer stem cells must become target of chemotherapy.

2.5 Sex hormones and MDR

2.5.1 ER

Estrogens play key roles in development and maintenance of normal sexual and reproductive function. The most potent estrogen produced in the body is 17 β -estradiol(E2). Two metabolites of E2, estrone and estriol, although they are high-affinity ligands are much weaker agonists on estrogen receptors (ERs)[102]. Up to now, two type of ERs have been found which named ER α (NR3A1) and ER β (NR3A2). At the regulation of some genes, particularly those involved in proliferation, ER α and ER β can have opposite actions [103], a

finding which suggests that the overall proliferative response to E2 is the result of a balance between ER α and ER β signaling.

The expression of ER α is closely associated with breast cancer biology, especially the development of tumors; estrogen hormones induce expression of c-myc and c-fos protooncogenes sufficient for cell division and breast cancer progression[104]. Many studies demonstrated that breast carcinomas which lack ER α expression often reveal more aggressive phenotypes. Furthermore, ER α expression in tumor tissues is a favorable predictor of prognosis in endocrine treatment[105]. ER α typically functions as a transcription factor to regulate specific gene expression which binds to estrogen response elements (ERE) upstream of the target genes. The study of Lisa D. Coles et al. demonstrated that E2 could up-regulate the expression of p-gp in P-gp Overexpressing Cells (NCI-ADR-RES) [106].

Anti-estrogens, designed to block ER α , are widely and effectively used clinically in the treatment of breast cancer. The most common drugs including tamoxifen and toremifene. Some researches show that antiestrogens such as tamoxifen, metabolites of tamoxifen (4-hydroxytamoxifen and N-desmethyltamoxifen), droloxifen, and toremifene stimulated the p-gp ATPase activity and are substrates of p-gp. These results suggest that the antiestrogens may be potent drugs that reverse the multidrug-resistant phenotype mediated by P-gp[107]. However, another study displayed that tamoxifen activates CYP3A4 and MDR-1 genes through steroid and xenobiotic receptor (SXR) in breast cancer cells [108]. But, some other anti-estrogens seem to be more effective on reversing MDR. A study shows that the pure anti-oestrogen ICI 164 could enhance doxorubicin and VBL toxicity to MCF-7/Adr cells 25- and 35-fold, respectively and the pure anti-oestrogens iodotamoxifens completely reversed VBL resistance in the mdrl transfected lung cancer cell line [109].

Besides affection on p-gp, there are many data showing the relationship between anti-estrogens and BCRP. Imai et al. demonstrated that BCRP mRNA expression was induced by 17 β -estradiol in T47D:A18 cells [110]. Our research indicated that BCRP expression is upregulated by 17 β -estradiol via a novel pretranscriptional mechanism which might be involved in 17 β -estradiol-ER complexes binding to the ERE of BCRP promoter via the classical pathway to activate transcription of the BCRP gene[111]. Besides, we also found that tamoxifen and toremifene could reverse MDR mediated by BCRP in breast cancer cells[104].

2.5.2 Progesterone receptor (PR)

Like estrogens, the physiological effects of progesterone are mediated by interaction of the hormone with the progesterone receptor. Up to now, two types of PRs were detected, named as PRA and PRB, respectively. The two PRs are expressed from a single gene as a result of transcription from two alternative promoters[112]. In general, PRB acts as a stronger transcriptional activator, whereas PRA functions as a transcriptional inhibitor of PRB and ER[113]. PR expression in breast cancer is also an important indicator of likely responsiveness to endocrine agents. It has been shown that PRA and PRB are expressed in similar amounts in most breast tumors[114]. Some data indicated that progesterone via PRs may be related to the regulation of MDR in breast cancer.

In 1994, Rao US et al found that at 50 microM, progesterone stimulated the P-gp ATPase activity as effectively as verapamil and is a potent drugs inducing p-gp mediated MDR[115]. Recently study displays that transcriptional regulation by E2 and progesterone (P4) likely contributes to the modulation of P-gp levels[116].

Besides that, the relationship between PR and BCRP has also been focus on. Wang et al found there were progesterone response elements on the upstream of BCRP promoter[114] and they note that the identified PRE is exactly the same as the estrogen response element published by Ee et al[117]. They found that PRB is a strong activator of transcription of the BCRP promoter, and PRA represses the PRB activity in the human placental choriocarcinoma BeWo cells. But the real situation in breast cancer may be complex. Because 17β -estradiol can induce PRB expression and down-regulate BCRP expression through posttranscriptional modification[118]; On the other hand, PRA can repress the estrogen receptor activity[113]. So the relationship between progesterone receptor and BCRP needs further data.

2.6 EMT and MDR

Tumor invasiveness, and metastasis, as well as MDR are still great puzzle in the development and treatment of tumors. The interconversion between epithelial and mesenchymal cells (designated as epithelial-mesenchymal or mesenchymal-epithelial transition, EMT or MET, respectively) has received special attention and emerging evidence suggests that epithelial-mesenchymal transitions (EMTs) may take part in the above processes. An epithelial-mesenchymal transition (EMT) is a biologic process that allows a polarized epithelial cell, which normally interacts with basement membrane via its basal surface, to undergo multiple biochemical changes that enable it to assume a mesenchymal cell phenotype, which includes enhanced migratory capacity, invasiveness, elevated resistance to apoptosis, and greatly increased production of ECM components[119].

Kalluri R and Weinberg RA divided EMT into three types[119]. Type 1 EMTs can generate mesenchymal cells (primary mesenchyme) that have the potential to subsequently undergo a MET to generate secondary epithelia during implantation, embryogenesis, and organ development. Type 2 EMTs, the program begins as part of a repair-associated event that normally generates fibroblasts and other related cells in order to reconstruct tissues following trauma and inflammatory injury.

Type 3 EMTs occur in neoplastic cells that have previously undergone genetic and epigenetic changes, specifically in genes that favor clonal outgrowth and the development of localized tumors. Evidents show that EMT is critically linked with up-regulated invasion, metastasis, and angiogenesis. Figure 1 displays the relationship between EMT and progression of tumors. During the acquisition of EMT characteristics, cells lose epithelial cell-cell junctions, undergo actin cytoskeleton reorganization and decrease in the expression of proteins that promote cell-cell contact such as E-cadherin and β -catenin, and gain in the expression of mesenchymal markers such as vimentin, fibronectin, γ -smooth muscle actin (SMA), N-cadherin as well as increased activity of matrix metalloproteinases (MMPs) like MMP-2, MMP-3 and MMP-9, associated with an invasive phenotype[120].

The modulation of EMT is complicated. Many genes or signal transduction pathways are confirmed to take part in the regulation, such as hepatocyte growth factor (HGF)[121], transforming growth factor beta (TGF- β)[122], epidermal growth factor (EGF)[123], MMP-3[124] and so on. In addition, some transcriptional factors including snail and twist also play important role in EMT[125,126].

Recent studies have shown an intimate relationship between the EMT phenotype and MDR. Kajiyama et al. found that paclitaxel-resistant ovarian cancer cells showed phenotypic changes consistent with EMT[127]. These results were confirmed in other types of tumors

like gemcitabine-resistant pancreatic cancer cells, oxaliplatin-resistant colorectal cancer cells, lapatinib-resistant breast cancer[120]. In addition, tamoxifen-resistant breast cancer cells undergone EMT with altered β -catenin phosphorylation[128]. It has been indicated that mesenchymal-like cancers might be more sensitive to DNA damaging agents such as doxorubicin, whereas epithelial-like cancers are more sensitive to targeted therapies, such as EGFR and HER2 antagonists [129]. That may be the reason why mesenchymal-like, basal breast cancers are initially more sensitive to chemotherapy than epithelial-like luminal breast cancers[130]. However, it was discussed that basal, mesenchymal-like breast cancers possibly would be more prone to develop drug resistance. So more works need to do to investigate the links of EMT and MDR.

Fig. 1. The relationship between EMT and progression of tumors. Normal epithelial cells transform to tumor cells. After EMT, tumor cells invade into surrounding normal tissues and distant organs. Then, MET reverse the cells into epithelial cells in the metastasis.

2.7 Methylation and MDR

Cancer is known as a genetic disease. Gain, loss, and mutation of genetic information have long been known to contribute to cancer development and progression. It is being increasingly recognized that epigenetic alterations in cancer often serve as potent surrogates for genetic mutations. Methylation of CpG dinucleotides is an important pattern of epigenetics.

Methylation can directly interfere with the binding of transcription factors to inhibit replication and/or methyl-CpG binding proteins that can bind to methylated DNA, as well as regulatory proteins to inhibit transcription[131]. The patterns of CpG methylation are specie and tissue specific. The biological machinery of this system comprises a variety of regulatory proteins including DNA methyltransferases, putative demethylases, methyl-CpG binding proteins, histones modifying enzymes and chromatin remodeling complexes. Alterations in DNA methylation participate in the development of some human diseases, including tumor [132].

Then whether DNA methylation takes part in the modulation of MDR? The answer is yes. El-Osta et al. used inhibitors of DNA methyltransferase (5-azacytidine [5aC]) and histone deacetylase (trichostatin A [TSA]) to examine gene transcription, promoter methylation status, and the chromatin determinants associated with the MDR1 promoter and their result displayed that 5aC and TSA induced DNA demethylation, leading to reactivation of methylated MDR1[133]. Nakayama et al. demonstrated the hypomethylation status of the MDR1 promoter region might be a necessary condition for MDR1 gene overexpression and establishment of P-glycoprotein-mediated multidrug resistance in AML patients[134]. Detailed mapping of MDR1 promoter showed that its promoter is always hypermethylated in drug-sensitive cells, while the drugresistant cells have hypomethylated MDR1

promoter[135]. Gayatri Sharma et al used methylation-specific PCR to investigate the promoter methylation status of MDR1 in tumor and serum of 100 patients with invasive ductal carcinomas of breast (IDCs) and MDR1 was hypomethylated in 47% tumors and 44% paired serum of IDC patients [136].

The methylation of BCRP has also been focus on. To et al. have shown an active CpG island within the proximal ABCG2 promoter region contributing to inactivation of ABCG2[137]. A follow-up research by Turner et al. demonstrated that ABCG2 expression in multiple myeloma patients and in cell lines is regulated in part by promoter methylation[138].

DNA methylation has been found to anticipate the regulation of other MDR-related genes. Chekhun VF et al. found that the promoter regions of MDR1, GST-pi, genes were highly methylated in MCF-7 cell line but not in its MCF-7/R drug resistant variant. The results suggests that acquirement of doxorubicin resistance of MCF-7 cells is associated with DNA hypomethylation of the promoter regions of the MDR1, GST-pi[139].

3. Strategies to reverse MDR

Since MDR phenomena have been recognised, the war fighting against it has been continuing. Many strategies have been devised to overcome it and mainly divided into three types: modulators, immunotherapy and genetic therapy.

3.1 Modulators of MDR

Because P-gp is the best characterized gene conferring MDR and its wide effects, most modulators target for it. So herein, we divide the modulators into two types: modulators targeting P-gp and targeting other genes.

3.1.1 Modulators of P-gp

Up to now, numerous compounds have been shown to inhibit the drug efflux function of P-gp and therefore, reverse cellular resistance. Since P-gp was first detected in 1976, three generation of modulators are found or synthesis. The process of chemosensitization involves the co-administration of a MDR modulator with an anticancer drug in order to cause enhanced intracellular accumulation via impairing the P-gp function[2].

3.1.1.1 First generation modulators

The first compounds documented to reverse MDR was verapamil (VRP), one of the calcium channel blocker[140]. Studies displayed that VRP enhanced intracellular accumulation of many anticancer drugs, including DOX in numerous cell lines. Subsequent studies revealed that this MDR reversing character is shared by many other calcium channel blockers, clinically available calcium antagonists, and calmodulin antagonists, such as felodipine and trifluoroperazine[141]. Indole alkaloids, the anti-malarial quinine and the anti-arrhythmic quinidine, have also been shown to reverse MDR in vitro in experimental cell lines [142]. Cyclosporin A, a commonly used immunosuppressant for organ transplantation, remains one of the most effective first generation of MDR modulators[2].

A number of these first generation MDR modulators, such as VRP and CsA, displayed excellent MDR reversal activities both preclinically and clinically. However, a unique property shared by most first generation modulators is that they are therapeutic agents and typically reverse MDR at concentrations much higher than those required for their

individual therapeutic activity and at these elevated doses, both compounds exhibited severe and sometimes life-threatening toxicities[2].

3.1.1.2 Second generation modulators

In order to solve the high toxicity of the first generation modulators, many newer analogs of the first generation are researched which were more potent and considerably less toxic.

Analogues of VRP, including dexverapamil (less cardiotoxic R-enantiomer of VRP), emopamil, gallopamil, and Ro11-2933 (a tiapamil analogue) which reversed MDR in vitro to a degree equivalent to VRP, but with marginal toxicity in animal models were documented[2]. The non-immunosuppressive analogue of CsA, PSC 833, has demonstrated superior MDR reversal efficacy in conjunction with daunorubicin, DOX, vincristine, vinblastine, taxol, or mitoxantrone in many cell lines in vitro at concentrations of 0.5–2 mM[143].

Although these agents circumvented many of the problems experienced with first generation MDR modulators, when these agents were co-administered with anticancer agents for modulating P-gp-based MDR, they influenced the pharmacokinetics and biodistribution properties of the anticancer drugs, which resulted in increased toxicity to normal organs such as liver and kidney[144].

3.1.1.3 Third generation modulators

The third generation modulators of p-gp have recently been developed using structure-activity relationships and combinatorial chemistry approaches. These agents required low doses (in the nanomolar range (20–100 nM)) to achieve effective reversing concentrations in vivo.

The cyclopropyldibenzosuberane LY 335979 is the representative and is currently under investigation in phase II clinical trials. This substance is highly effective on P-gp-mediated MDR at the concentration of 0.1–0.2 μ M and shows a very strong affinity for P-gp[145]. Compared to CsA, LY 335979 is characterized by a 10-fold increased potency, latent modulating activity and a blockade specific for P-gp. Another drug the acridonecarboxamide GF 120918 exhibits similar characteristics to LY 335979, but seems to be more effective than LY 335979[146]. The effective concentration of it is 20–100 nM and is one of the most potent and selective MDR modulators disclosed thus far. Both of them are specific for P-gp-mediated MDR since it does not modulate MRP-mediated resistance. In addition, some bispecific chemosensitizers that block both P-gp and MRP were found, such as VX-710 and VX-853[2].

In summary, all the modulators of p-gp can be divided into 10 classifications. Table 1 selected list the modulators of p-gp of each classification reported.

3.1.2 Modulators of other genes

Besides agents targeting P-gp, drugs that inhibit other genes have also been developing. Table 2 displays the selected list of modulators that inhibit other MDR-related genes. Although these agents appear to be well tolerated in combination with anticancer drugs such as DOX, the lack selectivity for the tumor tissue P-GP is still their deficiency which is the cause of adversely affect therapy.

3.2 Immunotherapy of MDR

Another method of MDR reversal is the use of monoclonal antibodies, several of which can inhibit P-gp-mediated drug efflux in vitro. The monoclonal antibody (mAb) MRK16 is the

Immunosuppressant	Anti-arrhythmic agent
Cyclosporin A	Quinidine
Valspodar (PSC833)	Antifungal agent
HIV protease inhibitors	Ketoconazole
Ritonavir	Sedative
Saquinavir	Midazolam
Nelfi navir	Acridone carboxamide
Calcium channel blocker	LY 335979(zosoquidar)
Verapamil	GG918 (GF120918)
Bepridil	Peptide chemosensitiser
Diltiazem	Reversin 121
Flunarizine	Reversin 205
Progesterone antagonist	Anti-oestrogen
Mifepristone (RU486)	Tamoxifen

Table 1. Selected list of P-gp modulators[149]

Name	inhibitors
MRP1[3]	MS-209 XR-9576 (tariquidar) VX-710 (biricodar) Isothiocyanates tRA 98006 Agosterol A Rifampicin NSAIDs
BCRP(ABCG2)[3]	GF-120918 (elacridar) tRA 98006 Flavonoids Phytoestrogens Imatinib mesylate Fumitremorgin C TAG- 139
GST-pi[150]	Clofibrate Ethacrynic acid GSH analogs Gossypol Indomethacin Misonidazole Piriprost Quinones Quercetin Sulfasalazine
GCS[91]	PDMP PPMP Miglustat

Table 2. Selected list of modulator targeting other MDR-related genes

first antibody used for reversing MDR by Hamada and Tsuruo[147]. The results found that MRK16 increased intracellular accumulation and cytotoxicity of vincristine and actinomycin D in some MDR cell lines, but had no effect on doxorubicin cytotoxicity. An increase in the accumulation of vincristine and actinomycin D was also observed with two other anti-Pgp mAbs, HYB-241 and HYB-612[148].

3.3 Genetic therapy of MDR

The genetic therapy of MDR mainly includes two methods. The first method was established by Gottesman et al. They produced multidrug resistant bone marrow cells by transfecting them with vectors carrying the MDR1 cDNA and this process allowed bone marrow cells to apply a chemotherapeutic regimen at otherwise unacceptable doses, and thus overcoming MDR[149].

The other method is inhibiting MDR proteins including transcriptional/translational inhibition through the introduction of antisense oligonucleotides or ribozymes or RNA interference. Recently, researchers have done many works targeting different genes, such as *mdr1*, *MRP1*, *BCRP*, *GCS* and so on. These techniques were proved to have considerable effects on overcoming MDR in vitro and in animal models. However, as many of these methods require gene targeting and transfer, they are unlikely to produce any really significant in vivo applications anytime soon[149].

In summary, although many approaches have been adopted to battle with MDR, it will be for a long time to overcome it completely.

4. References

- [1] Germano S, O'Driscoll L. Breast cancer: understanding sensitivity and resistance to chemotherapy and targeted therapies to aid in personalised medicine. *Curr Cancer Drug Targets*. 2009; 9(3):398-418.
- [2] Rajesh Krishna, Lawrence D. Mayer. Multidrug resistance (MDR) in cancer Mechanisms, reversal using modulators of MDR and the role of MDR modulators in influencing the pharmacokinetics of anticancer drugs. *European Journal of Pharmaceutical Sciences*. 2000; 11(4): 265-283.
- [3] Tomris Ozben. Mechanisms and strategies to overcome multiple drug resistance in cancer. *FEBS Letters*. 2006; 580 (12): 2903-2909
- [4] Gottesman MM, Fojo T and Bates SE. Multidrug resistance in cancer: role of ATP-dependent transporters. *Nat Rev Cancer*. 2002; 2(1), 48-58
- [5] Higgins CF, Hiles ID, Salmond GPC, Gill DR, Downie JA, Evans IJ, Holland IB, Gray L, Buckel SD, Bell AW, Hermodson MA. A family of related ATP-binding subunits coupled to many distinct biological processes in bacteria. *Nature*. 1986; 323(6087): 448-450.
- [6] Sarkadi B, Ozvegy-Laczka C, Németh K, Váradi A. ABCG2--a transporter for all seasons. *FEBS Lett*. 2004; 567(1):116-20.
- [7] Juliano, R L, Ling, V. A surface glycoprotein modulating drug permeability in mutants Chinese hamster ovary cells. *Biochim Biophys Acta*. 1976; 455(1): 152-162.
- [8] Chen CJ, Chin JE, Ueda K, Clark DP, Pastan I, Gottesman MM, Roninson IB. Internal duplication and homology with bacterial transport proteins in the *mdr1* (P-glycoprotein) gene from multidrug resistant human cells. *Cell*. 1986; 47(3):381-389.
- [9] Lee CH, Bradley G, Zhang JT, Ling V. Differential expression of P-glycoprotein genes in primary rat hepatocyte culture. *J Cell Physiol*. 1993; 157(2):392-402.

- [10] Krishna R, Mayer LD. Multidrug resistance (MDR) in cancer. Mechanisms, reversal using modulators of MDR and the role of MDR modulators in influencing the pharmacokinetics of anticancer drugs. *Eur J Pharm Sci.* 2000; 11(4):265-283.
- [11] Frézard F, Pereira-Maia E, Quidu P, Priebe W, Garnier-Suillerot A. Garnier Suillerot. P-glycoprotein preferentially effluxes compounds containing free basic versus charged amine. *Eur. J. Biochem.* 2001; 268 (6): 1561-1567.
- [12] Conseil G, Perez-Victoria JM, Jault JM, Gamarro F, Goffeau A, Hofmann J, Di Pietro A. Protein kinase C effectors bind to multidrug ABC transporters and inhibit their activity. *Biochemistry.* 2001; 40(8):2564-2571.
- [13] Stavrovskaya AA. Cellular mechanisms of multidrug resistance of tumor cells. *Biochemistry (Mosc).* 2000; 65(1): 95-106
- [14] Klappe K, Hinrichs JW, Kroesen BJ, Sietsma H, Kok JW. MRP1 and glucosylceramide are coordinately over expressed and enriched in rafts during multidrug resistance acquisition in colon cancer cells. *Int J Cancer.* 2004;110(4):511-522.
- [15] Rudas M, Filipits M, Taucher S, Stranzl T, Steger GG, Jakesz R, Pirker R, Pohl G. Expression of MRP1, LRP and Pgp in breast carcinoma patients treated with preoperative chemotherapy. *Breast Cancer Res Treat* 2003;81(2):149-157.
- [16] Clarke R, Leonessa F, Trock B. Multidrug resistance/P-glycoprotein and breast cancer: review and metaanalysis. *Semin Oncol.* 2005; 32(6 Suppl 7):S9-15.
- [17] Leonard GD, Fojo T, Bates SE. The role of ABC transporters in clinical practice. *Oncologist.* 2003; 8(5):411-24.
- [18] Teodori E, Dei S, Scapecchi S, Gualtieri F. The medicinal chemistry of multidrug resistance (MDR) reversing drugs. *Farmaco.* 2002; 57(5):385-415.
- [19] Jedlitschky G, Leier I, Buchholz U, Center M, Keppler D. ATP-dependent transport of glutathione S-conjugates by the multidrug resistance-associated protein. *Cancer Res.* 1994; 54(18):4833-4836.
- [20] Manciu L, Chang XB, Riordan JR, Ruysschaert JM. Multidrug resistance protein MRP1 reconstituted into lipid vesicles: secondary structure and nucleotide-induced tertiary structure changes. *Biochemistry.* 2000; 39(42):13026-13033.
- [21] Keppler D, Leier I, Jedlitschky G, König J. ATP-dependent transport of glutathione S-conjugates by the multidrug resistance protein MRP1 and its apical isoform MRP2. *Chem Biol Interact.* 1998; 111-112:153-161.
- [22] Doyle LA, Yang W, Abruzzo LV, Krogmann T, Gao Y, Rishi AK, Ross DD. A multidrug resistance transporter from human MCF-7 breast cancer cells. *Proc. Natl. Acad. Sci. USA.* 1998; 95(26): 15665-15670.
- [23] Bailey-Dell KJ, Hassel B, Doyle LA, Ross DD. Promoter characterization and genomic organization of the human breast cancer resistance protein (ATP-binding cassette transporter G2 gene). *Biochim Biophys Acta.* 2001; 1520(3):234-241
- [24] J.D. Allen, A.H. Schinkel. Multidrug resistance and pharmacological protection mediated by the breast cancer resistance protein (BCRP/ABCG2), *Mol. Cancer Ther.* 2002; 1 (6): 427-434.
- [25] Bhatia A, Schäfer HJ, Hrycyna CA. Oligomerization of the human ABC transporter ABCG2: evaluation of the native protein and chimeric dimers. *Biochemistry.* 2005; 44(32):10893-10904.
- [26] Nakatomi K, Yoshikawa M, Oka M, Ikegami Y, Hayasaka S, Sano K, Shiozawa K, Kawabata S, Soda H, Ishikawa T, Tanabe S, Kohno S. Transport of 7-ethyl-10-hydroxycamptothecin (SN-38) by breast cancer resistance protein ABCG2 in human lung cancer cells, *Biochem. Biophys Res Commun.* 2001; 288 (4): 827-832.

- [27] Ni Z, Bikadi Z, Rosenberg MF, Mao Q. Structure and function of the human breast cancer resistance protein (BCRP/ABCG2). *Drug Metab.* 2010; 11(7):603-617.
- [28] Zhou S, Schuetz JD, Bunting KD, Colapietro AM, Sampath J, Morris JJ, Lagutina I, Grosveld GC, Osawa M, Nakauchi H, Sorrentino BP. The ABC transporter Bcrp1/ABCG2 is expressed in a wide variety of stem cells and is a molecular determinant of the side-population phenotype. *Nat Med.* 2001; 7(9):1028-1034.
- [29] Maliepaard M, Scheffer GL, Faneyte IF, van Gastelen MA, Pijnenborg AC, Schinkel AH, van De Vijver MJ, Scheper RJ, Schellens JH. Subcellular localization and distribution of the breast cancer resistance protein transporter in normal human tissues. *Cancer Res.* 2001; 61(8): 3458-3564.
- [30] Hirano M, Maeda K, Matsushima S, Nozaki Y, Kusuhara H, Sugiyama Y. Involvement of BCRP (ABCG2) in the biliary excretion of pitavastatin. *Mol Pharmacol.* 2005; 68(3):800-807.
- [31] Furukawa T, Wakabayashi K, Tamura A, Nakagawa H, Morishima Y, Osawa Y, Ishikawa T. Major SNP (Q141K) variant of human ABC transporter ABCG2 undergoes lysosomal and proteasomal degradations. *Pharm Res.* 2009; 26(2): 469-479.
- [32] Noguchi K, Katayama K, Mitsunashi J, Sugimoto Y. Functions of the breast cancer resistance protein (BCRP/ABCG2) in chemotherapy. *Adv Drug Deliv Rev.* 2009; 61(1):26-33.
- [33] Honjo Y, Hrycyna CA, Yan QW, Medina-Perez WY, Robey RW, van de Laar A, Litman T, Dean M, Bates SE. Acquired mutations in the MXR/BCRP/ABCP gene alter substrate specificity in MXR/BCRP/ABCP-overexpressing cells. *Cancer Res.* 2001; 61(18):6635-6639.
- [34] Polgar O, Ediriwickrema LS, Robey RW, Sharma A, Hegde RS, Li Y, Xia D, Ward Y, Dean M, Ozvegy-Laczka C, Sarkadi B, Bates SE. Arginine 383 is a crucial residue in ABCG2 biogenesis. *Biochim Biophys Acta* 2009; 1788(7):1434-1443.
- [35] Honjo Y, Hrycyna CA, Yan QW, Medina-Pérez WY, Robey RW, van de Laar A, Litman T, Dean M, Bates SE. Acquired mutations in the MXR/BCRP/ABCP gene alter substrate specificity in MXR/BCRP/ABCP-overexpressing cells. *Cancer Res.* 2001; 61(18):6635-6639.
- [36] Ross DD, Karp JE, Chen TT, Doyle LA. Expression of breast cancer resistance protein in blast cells from patients with acute leukemia. *Blood.* 2000; 96(1):365-368.
- [37] Benderra Z, Faussat AM, Sayada L, Perrot JY, Chaoui D, Marie JP, Legrand O. Breast cancer resistance protein and P-glycoprotein in 149 adult acute myeloid leukemias. *Clin Cancer Res.* 2004 Dec 1;10(23):7896-902.
- [38] Usuda J, Ohira T, Suga Y, Oikawa T, Ichinose S, Inoue T, Ohtani K, Maehara S, Imai K, Kubota M, Tsunoda Y, Tsutsui H, Furukawa K, Okunaka T, Sugimoto Y, Kato H. Breast cancer resistance protein (BCRP) affected acquired resistance to gefitinib in a "never-smoked" female patient with advanced non-small cell lung cancer. *Lung Cancer.* 2007;58(2):296-299.
- [39] O'Connor PM, Jackman J, Bae I, Myers TG, Fan S, Mutoh M, Scudiero DA, Monks A, Sausville EA, Weinstein JN, Friend S, Fornace AJ Jr, Kohn KW. Characterization of the p53 tumor suppressor pathway in cell lines of the National Cancer Institute anticancer drug screen and correlations with the growth-inhibitory potency of 123 anticancer agents. *Cancer Res.* 1997; 57(19):4285-4300.
- [40] Gasco M, Yulug IG, Crook T. TP53 mutations in familial breast cancer: functional aspects. *Hum Mutat.* 2003; 21(3): 301-306.

- [41] Aas T, Børresen AL, Geisler S, Smith-Sørensen B, Johnsen H, Varhaug JE, Akslen LA, Lønning PE. Specific P53 mutations are associated with de novo resistance to doxorubicin in breast cancer patients. *Nat Med*. 1996; 2(7):811-814.
- [42] Chin KV, Ueda K, Pastan I, Gottesman MM. Modulation of activity of the promoter of the human MDR1 gene by Ras and p53. *Science*. 1992; 255(5043): 459-462.
- [43] Zhou G, Kuo MT. Wild-type p53-mediated induction of rat *mdr1b* expression by the anticancer drug daunorubicin. *J Biol Chem*. 1998; 273(25): 15387-15394.
- [44] Bush JA, Li G. Cancer chemoresistance: the relationship between p53 and multidrug transporters. *Int J Cancer*. 2002; 98: 323-330.
- [45] Linn SC, Honkoop AH, Hoekman K, van der Valk P, Pinedo HM, Giaccone G. p53 and P-glycoprotein are often co-expressed and are associated with poor prognosis in breast cancer. *Br J Cancer*. 1996; 74: 63-68.
- [46] Linn SC, Pinedo HM, van Ark-Otte J, van der Valk P, Hoekman K, Honkoop AH, Vermorken JB, Giaccone G.. Expression of drug resistance proteins in breast cancer, in relation to chemotherapy. *Int J Cancer*. 1997; 71(5): 787-795.
- [47] Wang X, Wu X, Wang C, Zhang W, Ouyang Y, Yu Y, He Z.. Transcriptional suppression of breast cancer resistance protein (BCRP) by wild-type p53 through the NF-kappaB pathway in MCF-7 cells. *FEBS Lett*. 584(15): 3392-3397.
- [48] Coley HM. Mechanisms and strategies to overcome chemotherapy resistance in metastatic breast cancer. *Cancer Treat Rev*. 2008; 34(4): 378-390.
- [49] Stavrovskaya AA. Cellular mechanisms of multidrug resistance of tumor cells. *Biochemistry (Mosc)*. 2000; 65(1): 95-106.
- [50] Shitashige M, Toi M, Yano T, Shibata M, Matsuo Y, Shibasaki F. Dissociation of Bax from a Bcl-2/Bax heterodimer triggered by phosphorylation of serine 70 of Bcl-2. *J Biochem*. 2001; 130(6): 741-748.
- [51] Liu F, Xie ZH, Cai GP, Jiang YY. The effect of survivin on multidrug resistance mediated by P-glycoprotein in MCF-7 and its adriamycin resistant cells. *Biol Pharm Bull*. 2007; 30(12): 2279-2283.
- [52] Zhang M, Latham DE, Delaney MA, Chakravarti A. Survivin mediates resistance to antiandrogen therapy in prostate cancer. *Oncogene*. 2005; 24(15): 2474-2482.
- [53] Devling TW, Lindsay CD, McLellan LI, McMahon M, Hayes JD. Utility of siRNA against Keap1 as a strategy to stimulate a cancer chemopreventive phenotype. *Proc Natl Acad Sci USA*. 2005; 102:7280-5A.
- [54] Meijerman I, Beijnen JH, Schellens JH. Combined action and regulation of phase II enzymes and multidrug resistance proteins in multidrug resistance in cancer. *Cancer Treat Rev*. 2008; 34(6):505-520.
- [55] Su F, Hu X, Jia W, Gong C, Song E, Hamar P. Glutathion S transferase pi indicates chemotherapy resistance in breast cancer. *J Surg Res*. 2003 ;113(1):102-108.
- [56] Ban N, Takahashi Y, Takayama T, Kura T, Katahira T, Sakamaki S, et al. Transfection of glutathione S-transferase (GST)-pi antisense complementary DNA increases the sensitivity of a colon cancer cell line to adriamycin, cisplatin, melphalan, and etoposide. *Cancer Res* 1996; 56: 3577-3582.
- [57] Akan I, Akan S, Akca H, Savas B, Ozben T. Multidrug resistance-associated protein 1 (MRP1) mediated vincristine resistance: effects of N-acetylcysteine and buthionine sulfoximine. *Cancer Cell Int* 2005;5:22.
- [58] Adler V, Yin Z, Fuchs SY, Benezra M, Rosario L, Tew KD, Pincus MR, Sardana M, Henderson CJ, Wolf CR, Davis RJ, Ronai Z. Regulation of JNK signaling by GSTp. *EMBO J*. 1999; 18(5):1321-1334.

- [59] Benet LZ, Cummins CL, Wu CY. Unmasking the dynamic interplay between efflux transporters and metabolic enzymes. *Int J Pharm.* 2004; 277:3–9.
- [60] Tew KD, Monks A, Barone L, Rosser D, Akerman G, Montali JA. Glutathione-associated enzymes in the human cell lines of the National Cancer Institute Drug Screening Program. *Mol Pharmacol.* 1996; 50:149–159.
- [61] Chanas SA, Jiang Q, McMahon M, McWalter GK, McLellan LI, Elcombe CR, et al. Loss of the Nrf2 transcription factor causes a marked reduction in constitutive and inducible expression of the glutathione S-transferase *Gsta1*, *Gsta2*, *Gstm1*, *Gstm2*, *Gstm3* and *Gstm4* genes in the livers of male and female mice. *Biochem J.* 2002;365:405–416.
- [62] Hayashi A, Suzuki H, Itoh K, Yamamoto M, Sugiyama Y. Transcription factor Nrf2 is required for the constitutive and inducible expression of multidrug resistance-associated protein 1 in mouse embryo fibroblasts. *Biochem Biophys Res Commun.* 2003;310:824–829.
- [63] Moscow JA, Townsend AJ, Goldsmith ME, Whang-Peng J, Vickers PJ, Poisson R, Legault-Poisson S, Myers CE, Cowan KH. Isolation of the human anionic glutathione S-transferase cDNA and the relation of its gene expression to estrogen-receptor content in primary breast cancer. *Proc Natl Acad Sci U S A.* 1988;85(17):6518–6522.
- [64] Jhaveri MS, Morrow CS. Methylation-mediated regulation of the glutathione S-transferase P1 gene in human breast cancer cells. *Gene.* 1998; 210(1):1–7.
- [65] Berger JM, Gamblin SJ, Harrison SC, Wang JC. Structure and mechanism of DNA topoisomerase II. *Nature.* 1996; 379(6562):225–232.
- [66] Schmidt BH, Burgin AB, Dewese JE, Osheroff N, Berger JM. A novel and unified two-metal mechanism for DNA cleavage by type II and IA topoisomerases. *Nature.* 2010 Jun 3;465(7298):641–644.
- [67] Tsai-Pflugfelder M, Liu LF, Liu AA, Tewey KM, Whang-Peng J, Knutsen T, Huebner K, Croce CM, Wang JC. Cloning and sequencing of cDNA encoding human DNA topoisomerase II and localization of the gene to chromosome region 17q21–22. *Proc Natl Acad Sci U S A.* 1988 Oct;85(19):7177–7781.
- [68] Jenkins JR, Ayton P, Jones T, Davies SL, Simmons DL, Harris AL, Sheer D, Hickson ID. Isolation of cDNA clones encoding the L isozyme of human DNA topoisomerase II and localization of the gene to chromosome 3p24. *Nucleic Acids Res.* 1992; 20(21):5587–5592.
- [69] Keith WN, Douglas F, Wishart GC, McCallum HM, George WD, Kaye SB, Brown R. Co-amplification of *erbB2*, topoisomerase II α and the retinoic acid receptor α genes in breast cancer and allelic loss at topoisomerase I on chromosome 20. *Eur J Cancer* 1993(29A): 1469–1473.
- [70] Sng JH, Heaton VJ, Bell M, Maini P, Austin CA, Fisher LM. Molecular cloning and characterization of the human topoisomerase II α and II β genes: evidence for isoform evolution through gene duplication. *Biochim Biophys Acta.* 1999; 1444(3): 395–406.
- [71] Jarvinen TA, Liu ET. HER-2/neu and topoisomerase II α as simultaneous drug targets in cancer. *Comb Chem High Throughput Screen.* 2003;6:455–470.
- [72] Valkov NI, Sullivan DM. Drug resistance to DNA topoisomerase I and II inhibitors in human leukemia, lymphoma, and multiple myeloma. *Semin Hematol.* 1997; 34:48–62.

- [73] Järvinen TA, Tanner M, Rantanen V, Bärlund M, Borg A, Grénman S, Isola J.. Amplification and deletion of topoisomerase IIalpha associate with ErbB-2 amplification and affect sensitivity to topoisomerase II inhibitor doxorubicin in breast cancer. *Am J Pathol* 2000;156(3):839- 847.
- [74] Withoff S, Keith WN, Knol AJ, Coutts JC, Hoare SF, Mulder NH, de Vries EG.. Selection of a subpopulation with fewer DNA topoisomerase II alpha gene copies in a doxorubicin-resistant cell line panel. *Br J Cancer* 1996;74(4):502- 507.
- [75] Oakman C, Moretti E, Galardi F, Santarpia L, Di Leo A. The role of topoisomerase IIa and HER-2 in predicting sensitivity to anthracyclines in breast cancer patients. *Cancer Treat Rev.* 2009 Dec;35(8):662-7.
- [76] Jarvinen TA, Liu ET. HER-2/neu and topoisomerase II alpha in breast cancer. *Breast Cancer Res Treat* 2003;78(3): 299-311.
- [77] O'Malley FP, Chia S, Tu D, Shepherd LE, Levine MN, Bramwell VH, Andrulis IL, Pritchard KI.. Topoisomerase II alpha and responsiveness of breast cancer to adjuvant chemotherapy. *J Natl Cancer Inst* 2009;101(9): 644–650.
- [78] Gennari A, Sormani M, Pfeffer U. TOP mRNA expression in HER2 negative breast cancer. *San Antonio Breast Cancer Sympos.* 2008:6036.
- [79] Hannun YA, Luberto C, Argraves KM. Enzymes of sphingolipid metabolism: from modular to integrative signaling. *Biochemistry.* 2001;40(16):4893-4903.
- [80] Kolesnick R. The therapeutic potential of modulating the ceramide/sphingomyelin pathway. *J Clin Invest.* 2002; 110(1):3-8.
- [81] Uchida Y, Itoh M, Taguchi Y, Yamaoka S, Umehara H, Ichikawa S, Hirabayashi Y, Holleran WM, Okazaki T. Ceramide reduction and transcriptional up-regulation of glucosylceramide synthase through doxorubicin-activated Sp1 in drug-resistant HL-60/ADR cells. *Cancer Res.* 2004; 64(17):6271-6279.
- [82] Shukla A, Shukla GS, Radin NS. Control of kidney size by sex hormones: possible involvement of glucosylceramide. *Am J Physiol.* 1992; 262(1 pt 2):F24-29.
- [83] Lavie Y, Cao H, Bursten SL, Giuliano AE, Cabot MC. Accumulation of glucosylceramides in multidrug-resistant cancer cells. *J Biol Chem* 1996;271(32):19530-19536.
- [84] Lucci A, Cho WI, Han TY, Giuliano AE, Morton DL, Cabot MC. Glucosylceramide: a marker for multiple-drug resistant cancers. *Anticancer Res.* 1998;18(1B):475-480.
- [85] Yamashita T, Wada R, Sasaki T, Deng C, Bierfreund U, Sandhoff K, Proia RL. A vital role for glycosphingolipid synthesis during development and differentiation. *Proc Natl Acad Sci U S A.* 1999; 96(16):9142-9147.
- [86] Liu YY, Han TY, Giuliano AE, Cabot MC. Expression of glucosylceramide synthase, converting ceramide to glucosylceramide, confers adriamycin resistance in human breast cancer cells. *J Biol Chem* 1999;274(2):1140-1146.
- [87] Liu YY, Han TY, Giuliano AE, Hansen N, Cabot MC. Uncoupling ceramide glycosylation by transfection of glucosylceramide synthase antisense reverses adriamycin resistance. *J Biol Chem.* 2000;275(10):7138-7143.
- [88] Olshefski RS, Ladisch S. Glucosylceramide synthase inhibition enhances vincristine-induced cytotoxicity. *Int J Cancer.* 2001; 93(1):131-138.
- [89] Liu YY, Gupta V, Patwardhan GA, Bhinge K, Zhao Y, Bao J, Mehendale H, Cabot MC, Li YT, Jazwinski SM. Glucosylceramide synthase upregulates MDR1 expression in the regulation of cancer drug resistance through cSrc and beta-catenin signaling. *Mol Cancer.* 2010;9:145.

- [90] Liu YY, Yu JY, Yin D, Patwardhan GA, Gupta V, Hirabayashi Y, Holleran WM, Giuliano AE, Jazwinski SM, Gouaze-Andersson V, Consoli DP, Cabot MC...A role for ceramide in driving cancer cell resistance to doxorubicin. *FASEB J*. 2008;22(7):2541-2551.
- [91] Ruckhäberle E, Karn T, Hanker L, Gätje R, Metzler D, Holtrich U, Kaufmann M, Rody A. Prognostic relevance of glucosylceramide synthase (GCS) expression in breast cancer. *J Cancer Res Clin Oncol*. 2009;135(1):81-90.
- [92] Nguyen NP, Almeida FS, Chi A, Nguyen LM, Cohen D, Karlsson U, Vinh-Hung V. Molecular biology of breast cancer stem cells: potential clinical applications. *Cancer Treat Rev*. 2010; 36(6):485-91. Review.
- [93] Fuchs E, Segre JA. Stem cells: a new lease on life, *Cell*. 2000; 100(1):143-55. Review.
- [94] Seigel GM, Campbell LM, Narayan M, Gonzalez-Fernandez F. Gonzalez-Fernandez. Cancer stem cell characteristics in retinoblastoma, *Mol Vis*. 2005 Sep 12;11:729-37.
- [95] Grichnik JM, Burch JA, Schulteis RD, Shan S, Liu J, Darrow TL, Vervaert CE, Seigler HF. Melanoma, a tumor based on a mutant stem cell? *J. Invest. Dermatol*. 2006; 126 (1) 142-153.
- [96] Al-Hajj M, Wicha MS, Benito-Hernandez A, Morrison SJ, Clarke MF. Prospective identification of tumorigenic breast cancer cells. *Proc Natl Acad Sci USA*. 2003; 100(7): 3983-3988
- [97] Ginestier C, Hee Hur M, Charafe-Jauffret E, Monville F, Dutcher J, Brown M, Jacquemier J, Viens P, Kleer CG, Liu S, Schott A, Hayes D, Birnbaum D, Wicha MS, Dontu G. ALDH1 is a marker of normal and malignant human mammary stem cells and a predictor of poor clinical outcome. *Cell Stem Cell*. 2007; 1(5): 555-567
- [98] Goodell MA, Brose K, Paradis G, et al. Isolation and functional properties of murine hematopoietic stem cells that are replicating in vivo. *J Exp Med*, 1996; 183(4):1797-806.
- [99] Patrawala L, Calhoun T, Schneider-Broussard R, Zhou J, Claypool K, Tang DG. Side population is enriched in tumorigenic, stem-like cancer cells, whereas ABCG2+ and ABCG2- cancer cells are similarly tumorigenic. *Cancer Res*, 2005; 65(14):6207-6219.
- [100] Decraene C, Benchaouir R, Dillies MA, Israeli D, Bortoli S, Rochon C, Rameau P, Pitaval A, Tronik-Le Roux D, Danos O, Gidrol X, Garcia L, Piétu G. Global transcriptional characterization of SP and MP cells from the myogenic C2C12 cell line: effect of FGF6, *Physiol. Physiol Genomics*. 2005 Oct 17;23(2):132-149.
- [101] Hadnagy A, Gaboury L, Beaulieu R, Balicki D. SP analysis may be used to identify cancer stem cell populations. *Exp Cell Res*. 2006; 312(19):3701-3710.
- [102] Heldring N, Pike A, Andersson S, Matthews J, Cheng G, Hartman J, Tujague M, Ström A, Treuter E, Warner M, Gustafsson JA. Estrogen Receptors: How Do They Signal and What Are Their Targets. *Physiol Rev*. 2007; 87(3):905-31. Review
- [103] Liu MM, Albanese C, Anderson CM, Hilty K, Webb P, Uht RM, Price RH Jr, Pestell RG, Kushner PJ. Opposing action of estrogen receptors alpha and beta on cyclin D1 gene expression. *J Biol Chem*. 2002; 277: 24353-24360.
- [104] Zhang Y, Wang H, Wei L, Li G, Yu J, Gao Y, Gao P, Zhang X, Wei F, Yin D, Zhou G. Transcriptional modulation of BCRP gene to reverse multidrug resistance by toremifene in breast adenocarcinoma cells. *Breast Cancer Res Treat*. 2010;123(3):679-689.
- [105] Hayashi SI, Eguchi H, Tanimoto K, Yoshida T, Omoto Y, Inoue A, Yoshida N, Yamaguchi Y. The expression and function of estrogen receptor alpha and beta in

- human breast cancer and its clinical application. *Endocr Relat Cancer*. 2003; 10(2):193-202.
- [106] Coles LD, Lee IJ, Voulalas PJ, Eddington ND. Estradiol and progesterone-mediated regulation of P-gp in P-gp overexpressing cells (NCI-ADR-RES) and placental cells (JAR). *Mol Pharm*. 2009; 6(6):1816-1825.
- [107] Rao US, Fine RL, Scarborough GA. Antiestrogens and steroid hormones: substrates of the human P-glycoprotein. *Biochem Pharmacol*. 1994; 48(2):287-292.
- [108] Nagaoka R, Iwasaki T, Rokutanda N, Takeshita A, Koibuchi Y, Horiguchi J, Shimokawa N, Iino Y, Morishita Y, Koibuchi N. Tamoxifen activates CYP3A4 and MDR1 genes through steroid and xenobiotic receptor in breast cancer cells. *Endocrine*. 2006; 30(3):261-268.
- [109] Kirk J, Syed SK, Harris AL, Jarman M, Roufogalis BD, Stratford IJ, Carmichael. . Reversal of P-glycoprotein-mediated multidrug resistance by pure anti-oestrogens and novel tamoxifen derivatives. *J Biochem Pharmacol*. 1994; 48(2):277-85.
- [110] Imai Y, Ishikawa E, Asada S, Sugimoto Y. Estrogen-mediated post transcriptional down-regulation of breast cancer resistance protein/ABCG2. *Cancer Res*. 2005; 65:596-604.
- [111] Zhang Y, Zhou G, Wang H, Zhang X, Wei F, Cai Y, Yin D. Transcriptional upregulation of breast cancer resistance protein by 17beta-estradiol in ERalpha-positive MCF-7 breast cancer cells. *Oncology*. 2006; 71(5-6):446-455.
- [112] Kastner P, Krust A, Turcotte B, Stropp U, Tora L, Gronemeyer H, Chambon P. Two distinct estrogen-regulated promoters generate transcripts encoding the two functionally different human progesterone receptor forms A and B. *EMBO J*. 1990; 9(5): 1603-1614.
- [113] Giangrande PH, McDonnell DP. The A and B isoforms of the human progesterone receptor: two functionally different transcription factors encoded by a single gene. *Recent Prog Horm Res*. 1999; 54: 291-313.
- [114] Wang H, Lee EW, Zhou L, Leung PC, Ross DD, Unadkat JD, Mao Q. Progesterone receptor (PR) isoforms PRA and PRB differentially regulate expression of the breast cancer resistance protein in human placental choriocarcinoma BeWo cells. *Mol Pharmacol*. 2008; 73: 845-854.
- [115] Rao US, Fine RL, Scarborough GA. Antiestrogens and steroid hormones: substrates of the human P-glycoprotein. *Biochem Pharmacol*. 1994; 48: 287-292.
- [116] Coles LD, Lee IJ, Voulalas PJ, Eddington ND. Estradiol and progesterone-mediated regulation of P-gp in P-gp overexpressing cells (NCI-ADR-RES) and placental cells (JAR). *Mol Pharm*. 2009; 6: 1816-1825.
- [117] Ee PL, Kamalakaran S, Tonetti D, He X, Ross DD, Beck WT. Identification of a novel estrogen response element in the breast cancer resistance protein (ABCG2) gene. *Cancer Res*. 2004; 64: 1247-1251.
- [118] Wang H, Zhou L, Gupta A, Vethanayagam RR, Zhang Y, Unadkat JD, Mao Q. Regulation of BCRP/ABCG2 expression by progesterone and 17beta-estradiol in human placental BeWo cells. *Am J Physiol Endocrinol Metab*. 2006; 290: E798-807.
- [119] Kalluri R, Weinberg RA. The basics of epithelial-mesenchymal transition. *J Clin Invest*. 2009; 119: 1420-1428.
- [120] Wang Z, Li Y, Ahmad A, Azmi AS, Kong D, Banerjee S, Sarkar FH. Targeting miRNAs involved in cancer stem cell and EMT regulation: An emerging concept in overcoming drug resistance. *Drug Resist Updat*. 2010; 13(4-5): 109-118.

- [121] Balkovetz DF, Pollack AL, Mostov KE. Hepatocyte growth factor alters the polarity of Madin-Darby canine kidney cell monolayers. *J Biol Chem.* 1997; 272: 3471-3477.
- [122] Janda E, Lehmann K, Killisch I, Jechlinger M, Herzig M, Downward J, Beug H, Grünert S. Janda E, Lehmann K, Killisch I, et al. Ras and TGF[β] cooperatively regulate epithelial cell plasticity and metastasis: dissection of Ras signaling pathways. *The Journal of cell biology.* 2002; 156(2): 299-313.
- [123] Lu Z, Ghosh S, Wang Z, Hunter T. Downregulation of caveolin-1 function by EGF leads to the loss of E-cadherin, increased transcriptional activity of β -catenin, and enhanced tumor cell invasion. *Cancer Cell.* 2003; 4: 499-515.
- [124] Radisky DC, Levy DD, Littlepage LE, Liu H, Nelson CM, Fata JE, Leake D, Godden EL, Albertson DG, Nieto MA, Werb Z, Bissell MJ. Rac1b and reactive oxygen species mediate MMP-3-induced EMT and genomic instability. *Nature.* 2005; 436(7047): 123-127.
- [125] Li QQ, Xu JD, Wang WJ, Cao XX, Chen Q, Tang F, Chen ZQ, Liu XP, Xu ZD. Twist1-mediated adriamycin-induced epithelial-mesenchymal transition relates to multidrug resistance and invasive potential in breast cancer cells. *Clin Cancer Res.* 2009; 15(8): 2657-2665.
- [126] de Herreros AG, Peiro S, Nassour M, Savagner P. Snail family regulation and epithelial mesenchymal transitions in breast cancer progression. *J Mammary Gland Biol Neoplasia.* 15: 135-147.
- [127] Kajiyama H, Shibata K, Terauchi M, Yamashita M, Ino K, Nawa A, Kikkawa F. Chemoresistance to paclitaxel induces epithelial-mesenchymal transition and enhances metastatic potential for epithelial ovarian carcinoma cells. *Int J Oncol.* 2007 ;31(2):277-283
- [128] Kim MR, Choi HK, Cho KB, Kim HS, Kang KW. Involvement of Pin1 induction in epithelial-mesenchymal transition of tamoxifen-resistant breast cancer cells. *Cancer Sci.* 2009 Oct;100(10):1834-1841.
- [129] Singh, A., Settleman, J., 2010. EMT, cancer stem cells and drug resistance: an emerging axis of evil in the war on cancer. *Oncogene*, doi:10.1038/onc.2010.215.
- [130] Carey, L.A., Dees, E.C., Sawyer, L., Gatti, L., Moore, D.T., Collichio, F., Ollila, D.W., Sartor, C.I., Graham, M.L., Perou, C.M., 2007. The triple negative paradox: primary tumor chemosensitivity of breast cancer subtypes. *Clin. Cancer Res.* 13, 2329-2334.
- [131] Segura-Pacheco B, Perez-Cardenas E, Taja-Chayeb L, Chavez-Blanco A, Revilla-Vazquez A, Benitez-Bribiesca L, Duenas-González A. Global DNA hypermethylation-associated cancer chemotherapy resistance and its reversion with the demethylating agent hydralazine. *J Transl Med.* 2006; 4: 32.
- [132] Rodriguez-Dorantes M, Tellez-Ascencio N, Cerbon MA, Lopez M, Cervantes A. DNA methylation: an epigenetic process of medical importance. *Rev Invest Clin.* 2004; 56: 56-71.
- [133] El-Osta A, Kantharidis P, Zalcberg JR, Wolffe AP. Precipitous release of methyl-CpG binding protein 2 and histone deacetylase 1 from the methylated human multidrug resistance gene (MDR1) on activation. *Mol Cell Biol.* 2002; 22: 1844-1857.
- [134] Nakayama M, Wada M, Harada T, Nagayama J, Kusaba H, Ohshima K, Kozuru M, Komatsu H, Ueda R, Kuwano M. Hypomethylation status of CpG sites at the promoter region and overexpression of the human MDR1 gene in acute myeloid leukemias. *Blood.* 1998; 92(11): 4296-4307.

- [135] Sharma D, Vertino PM. Epigenetic regulation of MDR1 gene in breast cancer: CpG methylation status dominates the stable maintenance of a silent gene. *Cancer Biol Ther.* 2004; 3: 549-550.
- [136] Sharma G, Mirza S, Parshad R, Srivastava A, Datta Gupta S, Pandya P, Ralhan R. CpG hypomethylation of MDR1 gene in tumor and serum of invasive ductal breast carcinoma patients. *Clin Biochem.* 43: 373-379.
- [137] To KK, Zhan Z, Bates SE. Aberrant promoter methylation of the ABCG2 gene in renal carcinoma. *Mol Cell Biol.* 2006; 26: 8572-8585.
- [138] Turner JG, Gump JL, Zhang C, Cook JM, Marchion D, Hazlehurst L, Munster P, Schell MJ, Dalton WS, Sullivan DM.. ABCG2 expression, function, and promoter methylation in human multiple myeloma. *Blood.* 2006; 108(12): 3881-3889.
- [139] Chekhun VF, Kulik GI, Yurchenko OV, Tryndyak VP, Todor IN, Luniv LS, Tregubova NA, Pryzimirska TV, Montgomery B, Rusetskaya NV, Pogribny IP. Role of DNA hypomethylation in the development of the resistance to doxorubicin in human MCF-7 breast adenocarcinoma cells. *Cancer Lett.* 2006; 231(1): 87-93.
- [140] Tsuruo T, Iida H, Tsukagoshi S, Sakurai Y. Overcoming of vincristine resistance in P388 leukemia in vivo and in vitro through enhanced cytotoxicity of vincristine and vinblastine by verapamil. *Cancer Res.* 1981; 41: 1967-1972.
- [141] Holtt V, Kouba M, Dietel M, Vogt G. Stereoisomers of calcium antagonists which differ markedly in their potencies as calcium blockers are equally effective in modulating drug transport by P-glycoprotein. *Biochem Pharmacol.* 1992; 43: 2601-8.
- [142] Sehested M, Jensen PB, Skovsgaard T, Bindslev N, Demant EJ, Friche E, Vindeløv L. Inhibition of vincristine binding to plasma membrane vesicles from daunorubicin-resistant Ehrlich ascites cells by multidrug resistance modulators. *Br J Cancer.* 1989; 60(6): 809-814.
- [143] Watanabe T, Naito M, Oh-hara T, Itoh Y, Cohen D, Tsuruo T. Modulation of multidrug resistance by SDZ PSC 833 in leukemic and solid-tumor-bearing mouse models. *Jpn J Cancer Res.* 1996; 87: 184-193.
- [144] Lum BL, Gosland MP. MDR expression in normal tissues. Pharmacologic implications for the clinical use of P-glycoprotein inhibitors. *Hematol Oncol Clin North Am.* 1995; 9: 319-336.
- [145] Dantzig AH, Law KL, Cao J, Starling JJ. Reversal of multidrug resistance by the P-glycoprotein modulator, LY335979, from the bench to the clinic. *Curr Med Chem.* 2001; 8: 39-50.
- [146] Wallstab A, Koester M, Bohme M, Keppler D. Selective inhibition of MDR1 P-glycoprotein-mediated transport by the acridone carboxamide derivative GG918. *Br J Cancer.* 1999; 79: 1053-1060.
- [147] Hamada H, Tsuruo T. Functional role for the 170- to 180-kDa glycoprotein specific to drug-resistant tumor cells as revealed by monoclonal antibodies. *Proc Natl Acad Sci U S A.* 1986 Oct;83(20):7785-7789.
- [148] Mechetner EB, Roninson IB. Efficient inhibition of P-glycoprotein-mediated multidrug resistance with a monoclonal antibody. *Proc Natl Acad Sci U S A.* 1992 Jul 1;89(13):5824-8.
- [149] Luqmani YA. Mechanisms of drug resistance in cancer chemotherapy. *Med Princ Pract.* 2005;14 Suppl 1:35-48.
- [150] Tew KD, Dutta S, Schultz M. Inhibitors of glutathione S-transferases as therapeutic agents. *Adv Drug Deliv Rev.* 1997; 26: 91-104.

Targeting New Pathways and Cell Death in Breast Cancer

Edited by Dr. Rebecca Aft

ISBN 978-953-51-0145-1

Hard cover, 190 pages

Publisher InTech

Published online 29, February, 2012

Published in print edition February, 2012

This book presents novel in interesting find by multiple accomplished investigators in breast cancer. These chapters elucidate new mechanisms of breast cancer cell death as well as discuss new pathways for therapeutic targeting.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Gengyin Zhou and Xiaofang Zhang (2012). Multidrug Resistance and Breast Cancer, Targeting New Pathways and Cell Death in Breast Cancer, Dr. Rebecca Aft (Ed.), ISBN: 978-953-51-0145-1, InTech, Available from: <http://www.intechopen.com/books/targeting-new-pathways-and-cell-death-in-breast-cancer/multidrug-resistance-and-breast-cancer>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](https://creativecommons.org/licenses/by/3.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen