

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

185,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Bariatric Surgery on Obese Type 2 Diabetes Patients

Junichirou Mori, Yoshihiko Sato and Mitsuhsa Komatsu
*Shinshu University,
Japan*

1. Introduction

Amidst a worldwide epidemic of diabetes, the World Health Organization estimates that 346 million people have diabetes and an estimated 3.4 million people died from consequences of high blood sugar in 2004¹⁾. Over time, diabetes can damage the heart, blood vessels, eyes, kidneys, and nerves. Especially diabetes increases the risk of heart disease and stroke.

Obesity carries with it significant risks of diabetes. Improvement in obesity is attendant with improvements in this ailment²⁾³⁾, and obese people consequently have been treated through pharmacotherapy, and intervention in life habits, including diet and exercise. Even with such treatment, however, it is very difficult to have satisfactory body weight loss. In the last few years, many studies have performed to compare intensive glucose control therapy with standard therapy. Most of the results show that body weight did not change with both intensive glucose control therapy and standard therapy⁴⁾⁵⁾. Moreover, many patients who are initially successful at weight loss then go on to rebound⁶⁾. Thus, promoting weight loss is a major issue in treatment, especially in severely obese patients. Recently, there has been an increase in patients with a BMI > 35 undergoing bariatric surgery and there has been a notable increase in studies describing effect of bariatric surgery on type 2 diabetes patients⁶⁾.

2. Effect of bariatric surgery on obesity and diabetes

Although an average of 55.9% loss in excess body weight was observed in bariatric surgery⁶⁾, the extent of operation-induced weight loss varies depending on the surgical method⁷⁾. While loss in body weight from intervention in life habits is insufficient, it was 46.2% for gastric binding and 59.5% for gastric bypass, respectively, thus showing that bariatric surgery leads to more efficient weight-loss results⁶⁾. However, although bariatric surgery generally leads to a great improvement in diabetes⁸⁾ (Table 1), there is a gradation of results depending on the procedure⁹⁾. Additionally for type 2 diabetes patients, studies on gastric bypass have shown that improvements in fasting plasma glucose and insulin sensitivity are evident prior to weight loss¹⁰⁾¹¹⁾. These kinds of changes are not observed in gastric binding⁶⁾¹²⁾. From these results, apart from improvements in insulin sensitivity induced through weight loss, gastric bypass is also thought to improve glucose metabolism.

	Total (n)	Gastric Binding (n)	Gastric Bypass (n)
Absolute Weight Loss (Kg)	-41.9 (266)	-26.0 (56)	-50.54 (129)
BMI Decrease (kg/m ²)	-14.0 (306)	-9.1 (56)	-18.0 (166)
Excess Loss (%)	-57.3 (267)	-41.0 (83)	-65.7 (184)
Fasting Insulin (pmol/L)	-123.9 (160)	-49.5 (56)	-153.7 (90)
HbA1c (%)	-2.4 (171)	-1.2 (83)	-3.0 (88)
Fasting Glucose (mmol/L)	-4.0 (296)	-3.2 (56)	-3.4 (164)
BMI: body mass index			

Table 1. Efficacy for Improvement in Diabetes-Related Outcomes for Diabetic and Glucose-Intolerant Patients ⁸⁾.

Incretin	Secretion	Function	Change after bariatric surgery
GLP-1	Distal ileum , Colon	Increase insulin release Slowing gastric emptying Controlling Glucagon secretion Induce satiety by working on the central nervous system	Increase
Ghrelin	Stomach	Stimulate appetite Increasing activity in the stomach Suppression of insulin secretion Increase growth hormone secretion	Decrease (Roux-en-Y Gastric Bypass)

Table 2. Summary of Intestin.

3. Effects of bariatric surgery on intestine hormone

One hypothesis to explain this phenomenon is the influence of gastrointestinal hormones. Glucagon-like peptide-1 (GLP-1), an intestinal hormone secreted from the distal ileum and colon in response to nutrient ingestion⁹⁾. GLP-1 acts on the beta cells to increase the level of cyclic AMP, leading to replenishment of the readily releasable pool of insulin granules during glucose-stimulated insulin secretion¹³⁾. Not only improving beta-cell function, GLP-1 is also involved in the proliferation and regeneration of pancreatic β -cells¹⁴⁾¹⁵⁾. Outside pancreatic effects, GLP-1 decreases dietary intake by slowing gastric emptying¹⁶⁾, controlling secretion of gastric acid¹⁷⁾ and glucagon¹⁸⁾, and induce satiety by working on the central nervous system¹⁹⁾²⁰⁾. There have been numerous studies detailing a post-operative increase in GLP-1 secretion from gastric bypass, and this increase occurs prior to post-gastric bypass weight loss²¹⁻²⁴⁾. Studies show that the post-gastric bypass GLP-1 level is significantly higher when compared to the post-gastric binding GLP-1 level^{25) 26)}. When considering the effect of GLP-1, it is possible that the increase in endogenous GLP-1 secretion plays an important role in the improvement of glucose metabolism by the gastric bypass surgery.

Although ghrelin is similar to GLP-1 in that it is related to the appetite, it is actually an appetite-stimulating hormone²⁷⁻²⁹⁾. It is likely that the appetite stimulating from ghrelin is due to its increasing activity in the stomach²⁸⁾³⁰⁾ and suppressing of insulin secretion³⁰⁾. Ghrelin levels increase in dietary restriction-induced weight loss and when there is a

negative energy balance, and conversely, decrease when eating or in the case of the obese³¹⁾³²⁾. However in the case of the obese, ghrelin levels become unchanged even when eating, and therefore, ghrelin level is a potential factor in obesity³³⁾. There are many reports demonstrating that fasting ghrelin levels decrease after Roux-en-Y Gastric Bypass compared to pre-operation³⁴⁾³⁵⁾. It has been reported that decrease in ghrelin levels occurs immediately following surgery and lasts for more than a year^{36) 37)}. Through Roux-en-Y Gastric Bypass, food bypasses the distal stomach in which ghrelin is released, and this may account for the post-bypass decrease in ghrelin levels³⁴⁾.

4. Incretin-therapy on type 2 diabetes patients

Recently GLP-1 analog/receptor agonists and GLP-1 degradation inhibitors are in clinical use. The beneficial points of these antidiabetes agents are glucose-dependent insulinitropism which may be reduced the risk of hypoglycemia³⁸⁾. Liraglutide is a GLP-1 analog with 97% sequence identity to human hormone. The mean reduction in HbA1c by liraglutide (1.8mg) was 1.14%³⁹⁾. In addition to effective glucose lowering, liraglutide produced beneficial effects on body weight. The mean reduction in body weight after 16weeks treatment of 1.8mg liraglutide was 3.6kg and was sustained throughout the 52 week study³⁹⁾. These results support the hypothesis that the bariatric surgery leads to a great improvement of diabetes not only by the reduction of the storage capacity of the stomach but also by the change in intestinal hormone.

5. Clinical application of bariatric surgery in Asia

As previously stated, there are reports that bariatric surgery leads to dramatic improvement in type 2 diabetes compared to pharmacotherapy and lifestyle intervention-based treatment. Will bariatric surgery replace conventional medication and/or life style intervention-based treatment in Asia? At present, however, most of these reports are not necessarily targeting regular subjects, given the subjects' extremely high average BMI of 47.9 kg/m² and relatively young average age of 40.2 years old⁶⁾.

Obesity in the Asian population is much less than in Western populations. The Ministry of Health, Labour and Welfare, Japan reported that only 3.7% of the population is obese (BMI>30)⁴⁰⁾. The rate of obesity in diabetes is reported to be similar to that in the rest of the Japanese population⁴¹⁾, and at present bariatric surgery has only a limited application in Japan. To increase the application of bariatric surgery in Asia, there is a need for a high-evidence level cohort study based on previous research that varies by age and obesity level in order to further the discussion on whether bariatric surgery should be given precedence over conventional medication and life style intervention-based treatment in patients with a BMI > 35.

6. Summary

Recently, in certain countries, there has been an increase in obese patients undergoing bariatric surgery which leads to more efficient weight-loss results. Bariatric surgery is an effective treatment option for severely obese patients for whom weight loss has been problematic with conventional pharmacotherapy and/or life style intervention-based

treatment. Gastric bypass has been shown not only to decrease body weight but to have an effect on incretin (Table 2). Change in incretin, especially GLP-1, could support the improvement in body weight. GLP-1 also has beneficially affect on pancreatic β -cells function, proliferation and regeneration. Thus even if diabetes did not cure by bariatric surgery, change in incretin have beneficially effect on diabetes. Therefore to choose the operative procedure of bariatric surgery, especially on patients who have basic disease, it is necessary to think about dynamic state of incretin.


7. References

- [1] World Health Organization: Diabetes. Fact sheet No312, Geneva, 2011
- [2] Pi-Sunyer X, Blackburn G, Yonavski SZ et al: Reduction in weight and cardiovascular disease risk factors in individuals with type 2 Diabetes: one-year results of the look AHEAD trial. *Diabetes Care* 30: 1374-1383, 2007
- [3] Espeland MA, Bray GA, Wing R et al: Describing patterns of weight changes using principal components analysis: results from the Action for Health in Diabetes (Look AHEAD) research group. *Ann Epidemiol* 19: 701-710, 2009
- [4] Patel A, MacMahon S, Travert F et al: Intensive blood glucose control and vascular outcomes in patients with type 2 diabetes. *N Engl J Med* 358: 2560-2572, 2008
- [5] Gerstein HC, Miller ME, Friedewald WT et al: Long-term effects of intensive glucose lowering on cardiovascular outcomes. *N Engl J Med* 364: 818-828, 2011
- [6] Buchwald H, Estok R, Sledge I et al: Weight and type 2 Diabetes after bariatric surgery: systematic review and meta-analysis. *Am J Med* 122: 248-256, 2009
- [7] Garb J, Welch G, Zagarins S, Kuhn J, Romanelli J: Bariatric surgery for the treatment of morbid obesity: a meta-analysis of weight loss outcomes for laparoscopic adjustable gastric banding and laparoscopic gastric bypass. *Obes Surg* 19: 1447-1455, 2009
- [8] Buchwald H, Avidor Y, Schoelles K et al: Bariatric surgery: a systematic review and meta-analysis. *JAMA* 292: 1724-1737, 2004
- [9] Pournaras DJ, LeRoux CW: The effect of bariatric surgery on gut hormones that alter appetite: *Diabetes Metab.* 35: 508-512, 2009
- [10] Bose M, Teixeira J, Laferrère B et al: Weight loss and incretin responsiveness improve glucose control independently after gastric bypass surgery. *J Diabetes* 2: 47-55, 2009
- [11] Bikman BT, Zheng D, Dohm GL et al: Mechanism for improved insulin sensitivity after gastric bypass surgery. *J Clin Endocrinol Metab* 93: 4656-4663, 2008
- [12] Vincent RP, leRoux CW: Changes in gut hormones after bariatric surgery. *Clin Endocrinol* 69: 173-179, 2008
- [13] Yajima H, Komatsu M, Shermerhorn T et al: cAMP enhances insulin secretion by an action on the ATP-sensitive K⁺ channel-independent pathway of glucose signaling in rat pancreatic islets. *Diabetes* 48:1006-1012, 1999
- [14] Holst JJ: The physiology of glucagon-like peptide 1. *Physiol Rev* 87: 1409-1439, 2007
- [15] Buteau J, Roduit R, Susini S, Prentki M: Glucagon-like peptide-1 promotes DNA synthesis, activates phosphatidylinositol 3-kinase and increases transcription factor pancreatic and duodenal homeobox gene 1 (PDX-1) DNA binding activity in beta (INS-1)-cells. *Diabetologia* 42: 856-864, 1999
- [16] Edholm T, Degerblad M, Schmidt PT, Hellström PM et al: Differential incretin effects of GIP and GLP-1 on gastric emptying, appetite and insulin-glucose homeostasis. *Neurogastroenterol Motil* 22:1191-1200, 2010

- [17] Nauck MA, Niedereichholz U, Schmiegeler WH et al: Glucagon-like peptide 1 inhibition of gastric emptying outweighs its insulinotropic effects in healthy humans. *Am J Physiol* 273: E981-E988, 1997
- [18] Asmar M, Bache M, Knop FK, Madsbad S, Holst JJ: Do the actions of glucagon-like peptide-1 on gastric emptying, appetite, and food intake involve release of amylin in humans? *J Clin Endocrinol Metab* 95: 2367-2375, 2010.
- [19] MaX, BruningJ, AshcroftFM: Glucagon-like peptide 1 stimulates hypothalamic proopiomelanocortin neurons. *J Neurosci* 27: 7125-7129, 2007
- [20] Pannacciulli N, Le DS, Krakoff J et al: Postprandial glucagon-like peptide-1 (GLP-1) response is positively associated with changes in neuronal activity of brain areas implicated in satiety and food intake regulation in humans. *NeuroImage* 35: 511-517, 2007
- [21] MacDonald PE, El-Kholy W, Wheeler MB et al: The multiple actions of GLP-1 on the process of glucose-stimulated insulin secretion. *Diabetes* 51: S434-S442, 2002
- [22] Goldfine AB, Mun EC, Patti ME et al: Patients with neuroglycopenia after gastric bypass surgery have exaggerated incretin and insulin secretory responses to a mixed meal. *J Clin Endocrinol Metab* 92: 4678-4685, 2007
- [23] LeRoux CW, Aylwin SJ, Bloom SR et al: Gut hormone profiles following bariatric surgery favor an anorectic state, facilitate weight loss, and improve metabolic parameters. *Ann Surg* 243: 108-114, 2006
- [24] Morínigo R, Moizé V, Vidal J et al: Glucagon-like peptide-1, peptide YY, hunger, and satiety after gastric bypass surgery in morbidly obese subjects. *J Clin Endocrinol Metab* 91: 1735-1740, 2006
- [25] Korner J, Bessler M, Inabnet W, Taveras C, Holst JJ: Exaggerated glucagon-like peptide-1 and blunted glucose-dependent insulinotropic peptide secretion are associated with Roux-en-Y gastric bypass but not adjustable gastric banding. *Surg Obes Relat Dis* 3: 597-601, 2007
- [26] Rodieux F, Giusti V, D'Alessio DA, Suter M, Tappy L: Effects of gastric bypass and gastric banding on glucose kinetics and gut hormone release. *Obesity (Silver Spring)* 16: 298-305, 2008
- [27] Murphy KG, Bloom SR: Gut hormones and the regulation of energy homeostasis. *Nature* 444: 854-859, 2006
- [28] Cummings DE, Overduin J: Gastrointestinal regulation of food intake. *J Clin Invest* 117: 13-23, 2007
- [29] Nakazato M, Murakami N, Date Y, Kojima M, Matsuo H, Kangawa K, Matsukura S: A role for ghrelin in the central regulation of feeding. *Nature* 409:194-198, 2001
- [30] Zwirska-Korczala K, Konturek SJ, Brzozowski T et al: Basal and postprandial plasma levels of PYY, ghrelin, cholecystokinin, gastrin and insulin in women with moderate and morbid obesity and metabolic syndrome. *J Physiol Pharmacol* 58:13-35, 2007
- [31] HansenTK, Dall R, Jørgensen JO et al: Weight loss increases circulating levels of ghrelin in human obesity. *Clin Endocrinol* 56: 203-206, 2002
- [32] Cummings DE, Foster-Schubert KE, Overduin J: Ghrelin and energy balance: focus on current controversies. *Curr Drug Targets* 6: 153-169, 2005
- [33] English PJ, Ghatei MA, Malik IA, Bloom SR, Wilding JP: Food fails to suppress ghrelin levels in obese humans. *J Clin Endocrinol Metab* 87: 2984, 2002

- [34] Lin E, Gletsu N, Smith CD et al: The effects of gastric surgery on systemic ghrelin levels in the morbidly obese. *Arch Surg* 139: 780-784, 2004
- [35] N. A. Tritos, E. Mun, A. Bertkau, R. Grayson, E. Maratos-Flier, and A. Goldfine: Serum ghrelin levels in response to glucose load in obese subjects post-gastric bypass surgery. *Obes Res* 11: 919-924, 2003
- [36] Leonetti F, Silecchia G, DiMario U et al: Different plasma ghrelin levels after laparoscopic gastric bypass and adjustable gastric banding in morbid obese subjects. *J Clin Endocrinol Metab* 88: 4227-4231, 2003
- [37] LeRoux CW, Welbourn R, Olber T et al: Gut hormones as mediators of appetite and weight loss after Roux-en-Y gastric bypass. *Ann Surg.* 246: 780-785, 2007
- [38] Ross SA and Ekoe JM: incretin agents in type 2 diabetes. *Can Fam Physician* 56(7): 639-648, 2010
- [39] Garber A, Henry R, Ratner R, Garcia-Hernandez PA Rodriguez-Pattzi H, Olvera-Alvarez I: Liraglutide versus glimepiride monotherapy for type 2 diabetes (LEAD-3 Mono): a randomized, 52-week, phase III, double-blind, parallel-treatment trial. *Lancet* 373(9662): 473-481. 2008
- [40] Ministry of Health, Labour and Welfare, Japan: Outline of the diabetes mellitus field study 2002. Tokyo, 2004
- [41] Sone H, Mizuno S, Fujii H, Yoshimura Y, Yamasaki Y, Ishibashi S, Katayama S, Saito Y, Ito H, Ohashi Y, Akanuma Y, Yamada N: Is the diagnosis of metabolic syndrome useful for predicting cardiovascular disease in Asian diabetic patients? Analysis from the Japan Diabetes Complications Study. *Diabetes Care.* 28:1463-1471, 2005

IntechOpen



Advanced Bariatric and Metabolic Surgery

Edited by Dr. Chih-Kun Huang

ISBN 978-953-307-926-4

Hard cover, 326 pages

Publisher InTech

Published online 29, February, 2012

Published in print edition February, 2012

Bariatric surgery has gained importance in the last 20 years because of the high prevalence of global obesity, and the vast understating of the physiological and pathological aspects of obesity and associated metabolic syndromes. This book has been written by a number of highly outstanding authors and pioneering bariatric surgeons from all over the world. The intended audience for this book includes all medical professionals involved in caring for bariatric patients. The chapters cover the choice of operation, preoperative preparation including psychological aspect, postoperative care and management of complication. It also extends to concept and result of metabolic surgery and scarless bariatric surgery.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Junichirou Mori, Yoshihiko Sato and Mitsuhsa Komatsu (2012). Bariatric Surgery on Obese Type 2 Diabetes Patients, Advanced Bariatric and Metabolic Surgery, Dr. Chih-Kun Huang (Ed.), ISBN: 978-953-307-926-4, InTech, Available from: <http://www.intechopen.com/books/advanced-bariatric-and-metabolic-surgery/bariatric-surgery-on-obese-type-2-diabetes-patients->

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](https://creativecommons.org/licenses/by/3.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen