We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Physiology of Astigmatism

Seyed-Farzad Mohammadi, Maryam Tahvildari and Hadi Z-Mehrjardi
*Eye Research Centre, Farabi Eye Hospital,
 Tehran University of Medical Sciences
 Iran*

1. Introduction

We know that the expression emmetropia is a conventional one and that in fact all normal human eyes have mild degrees of spherocylindrical errors (Shilo, 1997) or consist of a bitoric optical system, i.e. have principal meridians of relatively higher and lower powers at right angles. It is generally accepted that genetic factors have a significant role in determining ocular refractive status as well as astigmatism (Hammond et al., 2001) but many conditions and procedures such as surgery, suturing, wound healing, and ocular comorbidities modify the cylindrical status of the eye. Induced astigmatism and surgical correction of astigmatism are extensively addressed in other chapters of this book.

Manifest astigmatism is the vectorial sum of anterior corneal toricity and internal astigmatism. A variety of factors change the magnitude and shift the meridians of these cylindrical components and the perceived subjective astigmatism throughout life. Astigmatism is an extremely dynamic phenomenon, and changes in the shape of optical interfaces, refractive index, optical aperture, eyeball-extraocular structures (eyelids and extraocular muscles) interaction, visual tasks, accommodation, binocularity, tear film status, and even body position induce and modify baseline ocular astigmatism. In this chapter we shall focus on factors that determine baseline, diurnal, functional, and dynamic aspects of 'physiological astigmatism'.

2. Natural course of astigmatism in normal eyes

2.1 Age

Age-related evolution of ocular astigmatism in terms of power and axis has been observed in epidemiologic studies (Abrahamsson et al., 1988; Atkinson et al., 1980; Attebo et al., 1999; Baldwin & Mills, 1981; Ehrlich et al., 1997; Gwiazda et al., 1984; Hirsch, 1963; Kame et al., 1993; Sawada et al., 2008; Stirling, 1920). It is well documented that a high degree of astigmatism is present in neonates and infants; however, the reported amounts show discrepancies (Abrahamsson, et al. 1988; Howland & Sayles, 1985; Isenberg et al., 2004; Kohl & Samek, 1988; Varughese et al., 2005; Wood et al., 1995). The degree of astigmatism is even higher in preterm newborns and has an inverse association with postconceptional age and birth weight (Friling et al., 2004; Varghese et al., 2009). In near retinoscopy without cycloplegia, Gwiazda and colleagues found astigmatism of at least 1 D in about 55% of infants younger than 5 months, 10% of whom displayed a cylinder power of 3 D or more

(Gwiazda et al., 1984). In another study, photorefractive techniques showed that almost all infants at the age of 3 months had at least 1 D of astigmatism, which had decreased to adult levels by the age of 18 months (Atkinson et al., 1980). Likewise, a longitudinal study found astigmatism of at least 1 D in about 40% of infants at 3 months of age with a significant decrease to 4% by the age of 36 months. This reduction appears to be caused by the decrease in toricity of the cornea and the anterior lens (Mutti et al., 2004). Several studies have suggested that corneal shape changes throughout life. The linear reduction of the astigmatism to lower values with age is apparently a part of normal eye maturation (Friling et al., 2004) and emmetropisation. It has been suggested that the high astigmatism in early life induces and activates accommodation (Campbell & Westheimer, 1959; Howland, 1982).

Reports on the axis of astigmatism in infants are contradictory. According to Gullstrand, the natural form of the cornea is against the rule (Gullstrand 1962). Several studies have found a plus cylinder axis at 180 ± 20 (i.e. against the rule) in the majority of infants (Abrahamsson et al., 1988; Baldwin & Mills, 1981; Dobson et al., 1984; Gwiazda et al., 1984; Saunders 1986, 1988). While recent studies have raised questions about the reliability of previously used techniques. These studies have shown that with-the-rule astigmatism is more frequent among infants (Ehrlich et al. 1997; Isenberg et al., 2004; Mutti et al., 2004; Varghese et al., 2009).

As the child grows older, much of the early astigmatism will gradually disappear and transform into with the rule owing to eyelid pressure (Gwiazda et al. 1984). Most of the changes occur at ages 1-3 years, when the vertical and horizontal diameters of the cornea and its elasticity attain adult size and amount (Karesh, 1994). This with-the-rule astigmatism in preschool children is stabilized towards adolescence (Goss, 1991; Huynh et al., 2007; Shankar & Bobier, 2004). However, this is not always true, and a role for myopia development has been attributed to ocular astigmatism, as it may degrade optical blur cues and disrupt emmetropisation, which can lead to axial myopia progression in school-aged children (Gwiazda et al., 2000).

In early adulthood, astigmatism of more than one diopter is infrequent and is still with the rule. Lin and colleagues found a slight increase in the amount of astigmatism in medical students after five years (Lin et al., 1996). Other cross-sectional studies have indicated that mean total astigmatism changes with age, varying from as much as 0.62 D with the rule during adolescence to as much as 0.37 D against the rule in older ages (Baldwin & Mills, 1981). Baldwin and Mills found that steepening of the cornea in the horizontal meridian accounts for a major proportion of the increase in against-the-rule total astigmatism among older patients (Baldwin & Mills 1981). In the Blue Mountains Eye Study (Attebo et al., 1999), mean total astigmatism increased with age from 0.6 D to 1.2 D in youngest (49-59) to oldest (80-97) age groups. From a related study, an average of 1.6 D rise in total corneal astigmatism is documented for each five years of increase in age (Ho et al., 2010). Nuclear sclerosis cataract and the change in refractive index of the crystalline lens at older ages may contribute to myopic astigmatism (Fotadar et al., 2008).

Anterior corneal (and total) astigmatism shows flattening in the vertical meridian with aging, in contrast to a trend towards with-the-rule astigmatism on the posterior corneal surface (Ho et al., 2010). Against-the-rule astigmatism is the most common type of astigmatism in adults over 40 years of age. Interestingly, men are significantly more likely to develop against-the-rule astigmatism (Goto et al., 2001). In general, corneal toricity accounts

for the major component of total astigmatism (Asano et al., 2005; Ho et al., 2010); it is suggested that, with aging, upper eyelid pressure on the cornea and the tone of orbicularis muscle decrease (Marin-Amat, 1956). It has also been demonstrated that with-the-rule astigmatism decreases when eyelids are retracted from the cornea (Wilson et al., 1982). When relative steepening in the vertical meridian is abated, the intrinsic lenticular against-the-rule astigmatism will manifest. Decreases in action of extraocular muscles, especially the medial rectus (Marin-Amat, 1956), and vitreous syneresis and liquefaction may also contribute (Mehdizadeh, 2008).

The contribution of the lens to the ocular astigmatism is relatively constant throughout life (Hofstetter & Baldwin, 1957). Development of this lenticular astigmatism may be due to an emmetropisation phenomenon, as it effectively decreases manifest astigmatism in the early decades of life. But in older ages, lenticular astigmatism is manifested as an against-the-rule astigmatism when the corneal astigmatism is decreased (Artal et al., 2000, 2001; Ehrlich et al., 1997).

2.2 Diurnal changes of astigmatism in the normal eye

The magnitude and axis of astigmatism vary during the day; this variation can be described with regard to changes in eyelid pressure, extraocular muscle tension, pupil size and accommodation. From previous reports, it is postulated that generally the cornea has its flattest shape on awakening and steepens slightly until the evening (Manchester, 1970). Kiely et al. reported fluctuations in corneal asphericity during the day without recognizing a specific pattern (Kiely et al., 1982). Recently, diurnal variations in corneal topography have been studied by Read et al.: corneal wavefront error analysis revealed significant changes in astigmatism during the day (Read et al., 2005); see below.

2.2.1 Lid pressure and muscle tension in near tasks

Changes in corneal contour exerted through eyelid pressure have been widely discussed since the mid-1960s, and transient bilateral monocular diplopia after near work due to temporarily induced toricity in the cornea has been reported by a number of investigators (Bowman et al., 1978; Golnik & Eggenberger, 2001; Knoll, 1975; Mandell, 1966). It is agreed that visual tasks with significant downward gaze, such as reading, can alter corneal curvature owing to eyelid pressure (Collins et al., 2006; Read et al., 2007a). This will lead to horizontal bands on red reflex during retinoscopy (Ford et al., 1997) with concomitant topographic changes and corresponding distortions in Zernike wavefront analysis. Buehren et al. have reported changes towards against-the-rule astigmatism (Buehren et al., 2003).

In a recent study by Shaw et al., the average trend in the astigmatism axis due to near work was said to be against the rule, with approximately 0.25 D change within 15 minutes of 40° downward gaze, where both the upper and lower eyelids are in contact with the central 6 mm of the cornea. They also reported that eyelid tilt, curvature and position are important in the magnitude of corneal changes (Shaw et al., 2008).

Collins et al. demonstrated greater topographical changes in astigmatism during downward gaze with a larger angle (45° versus 25°) and with lateral eye movements (Collins et al., 2006). Studies on the time course of astigmatism regression have revealed slower recovery after longer periods of reading. Moreover, patterns of regression are similar among individuals, with a rapid recovery within the first 10 minutes after reading, and resolution takes between 30 to 60 minutes (Collins et al., 2005).

Regarding the role of extraocular muscles on corneal astigmatism, Lopping has mentioned that continuous use of the medial rectus muscle, especially during near tasks, imposes a force on the cornea which increases its radius of curvature in the horizontal meridian resulting in a shift towards against-the-rule astigmatism (Lopping & Weale, 1965).

These observations have implications for clinical testing, and it would be prudent that examinees avoid near tasks at least 30 minutes prior to refractive and topographic assessments.

2.2.2 Eyelid slant and tension

Apart from temporary changes of corneal curvature due to eyelid pressure, the cumulative effect of the eyelids contributes to naturally occurring astigmatism in healthy adults (see 2.1 above).

Slanting of the palpebral fissure is an important factor affecting corneal toricity (Read et al., 2007b). The magnitude of astigmatism increases as the palpebral fissure diverges from the horizontal plane. Male subjects show more downward fissure slanting, whereas female subjects show more upward fissure slanting (Garcia et al., 2003). People with Down's syndrome (Akinci et al., 2009; Little et al., 2009) or Treacher Collins syndrome (Wang et al., 1990) will show oblique astigmatism partly due to upward or downward slanting of the palpebral fissure.

Thicker or tighter eyelids tend to correspond with higher degrees of astigmatism as well. Asians and Native Americans show higher degrees of corneal astigmatism than other races (Osuobeni & Al Mijalli, 1997).

Corneal rigidity can also contribute to the amount of astigmatism caused by eyelid pressure. For instance, nutritional deficiencies are presumed to decrease corneal rigidity and flatten the horizontal meridian while steepening the vertical one (Lyle et al., 1972).

2.2.3 Pupil dynamics

We know that the optical system of the eye is not coaxial and at least three important axes have been described: optical axis (corneal optical center to lens's optical center), visual axis (object of regard to fovea; line of sight), and pupillary axis. There is a mild physiological pupil decentration in the nasal direction. Such physiological asymmetries, which have long been described (Walsh, 1988), induce coma (Wilson et al., 1992).

The pupil is the aperture for light entrance into the eye; excluding pharmacologic changes, pupil size and its (centroid) lateral position around the optical axis of the eye change according to ambient light (Walsh, 1988; Wilson et al., 1992), accommodative effort, and emotional status (Wilson et al., 1992).

Pupil size correlates with both the magnitude and orientation of astigmatism. Larger mesopic pupil sizes are detected with higher cylinder powers and are also associated with with-the-rule astigmatism rather than against-the-rule and oblique astigmatism (Cakmak et al., 2010). Larger pupil sizes—in low lighting conditions— increase the amount of higher order aberrations such as coma and may intensify the cylinder power in subjective/manifest refraction. Coma has been shown to be correlated with greater amounts of astigmatism (Hu et al. 2004). On the contrary, pupillary accommodative constriction reduces higher order aberrations including lenticular astigmatism (Sakai et al., 2007).

About 0.4 mm temporal pupil centroid shift in darkness was first reported by Walsh (Walsh, 1988); Wilson and Campbell (Wilson et al., 1992) then found shifts of up to 0.6 mm with decreased illumination, in nasal or temporal directions.

2.2.4 Accommodation and convergence

Three decades ago, Brzezinski introduced the expression 'accommodative astigmatism' and claimed that changes in lenticular astigmatism can neutralize corneal astigmatism and reduce the eye's overall toricity (Brzezinski, 1982). Other investigators have suggested that astigmatism increases as the accommodative response becomes larger (Denieul, 1982; Ukai & Ichihashi, 1991). According to Brzezinski, accommodative astigmatism is related to lens distortion due to inhomogeneous lens elasticity, variable constriction in ciliary muscles (which itself changes the lens power), and nonhomogeneous tension of the extraocular muscles during convergence (which causes corneal distortion). These may explain 'lag of accommodation', the phenomenon of less accommodative response than the accommodative stimulus in the horizontal meridian and the resultant with-the-rule astigmatism (Tsukamoto et al., 2001). Pupillary constriction may contribute to such changes as well (see above).

In a more recent study, Tsukamoto et al. found that all emmetropic subjects became astigmatic during accommodation, 93% with the rule (mean -1.96 D). Corneal astigmatism of with-the-rule orientation with mean values of 0.84 D and 0.91 D, respectively, for right and left eyes was detected without a direct association with the amount of accommodation. The eyes became emmetropic just after relaxation (Tsukamoto et al., 2000). Cheng et al. examined wavefront aberrations in a large adult population and found changes in astigmatism towards with the rule with an average of -0.1 D during maximum accommodation (Cheng et al., 2004). The mentioned pupillary and accommodative effects interact with the factors considered above during near tasks (see above).

Accommodation always accompanies convergence during near-vision tasks (Tait, 1933; Rosenfield & Gilmartin, 1988), and it is known that slight changes in cylinder power and axis (towards with the rule) occur during convergence alone (Beau Seigneur, 1946; Lopping & Weale, 1965; Tsukamoto, Nakajima et al., 2000). Seigneur has mentioned that this change is seen in a small percentage of eyes (Beau Seigneur, 1946), and many of the individuals do not experience any discomfort when using the same spectacles for far and near activities; nevertheless, for those who experience such an alteration, separate spectacle prescriptions for near and far distance vision might be beneficial.

2.2.5 Cyclotorsion and binocularity

Eye rotation around the Z axis (rolling or cyclotorsion) modifies the axis of ocular astigmatism in relation to the outside world. There is a complex interaction between accommodation, baseline astigmatism, and torsional alignment (Buehren et al., 2003; Read et al., 2007). These features contribute to eye fusional potential, depth perception, and depth of field (Regan & Spekreijse, 1970).

A number of reasons are implicated for physiological ocular torsion including unmasking of cyclophoria during monocular fixation and fusion loss (Tjon-Fo-Sang et al., 2002; Borish & Benjamin, 2006) and changing of body position from upright to supine (Park et al., 2009; Hori-Komai et al., 2007; Fea et al., 2006; Chernyak, 2004; Swami et al., 2002); these changes gain remarkable clinical significance when an individual is examined in the seating position but undergoes laser ablation in the supine position. Binocularity is also disturbed during corneal topography and wavefront aberrometry; binocular viewing is not normal during laser ablation in the supine position either.

Although several studies have shown significant incyclotorsion or excyclotorsion of about 2-4 degrees (maximum 9 to 14 degrees) as a result of changing the body position from seated

to supine (Swami et al., 2002; Chernyak, 2004; Fea et al., 2006; Neuhanh et al., 2010), a number of investigations have reported insignificant axis changes of less than 2 degrees, which can hardly affect astigmatic correction (Tjon-Fo-Sang et al., 2002; Becker et al., 2004). It has been suggested that axis misalignment of about 4 degrees will lead to 14% cylinder undercorrection during laser ablation (Swami et al., 2002; Neuhanh et al., 2010).

As mentioned above, cyclotorsion is frequently seen when switching from binocular to monocular vision, especially in those who have significant cyclophoria (Borish & Benjamin, 2006). Although it has been believed that an occluded eye shows excyclophoria under monocular occlusion of several hours (Graf et al., 2002), it is indeterminate whether ocular torsion resulting from monocular occlusion for a short period during refractive surgery or retinoscopy and monocular subjective refraction is clockwise or counter-clockwise; Hori-Komai et al. and Chang et al. both demonstrated that the magnitude and direction of cyclotorsion is different for each individual (Chang, 2008; Hori-Komai et al., 2007). This has significance in subjective refraction refinement and binocular balancing as well; in fact, a novel position in the phoropter allows maintenance of binocularity (fusion) while clarity of the images of the eyes are independently assessed (Borish & Benjamin, 2006).

Apart from body position and monocular vision, which account for static eye rotational alignment, dynamic cyclotorsion also occurs during laser ablation and may result in astigmatic undercorrection and/or induced astigmatism (Neuhanh et al., 2010; Chang, 2008; Hori-Komai et al., 2007). Fea et al. showed that blurring of the fixation target happens during ablation (after epithelium removal in surface ablation and following flap lifting in LASIK) and is an important factor for dynamic cyclotorsion, the magnitude of which seems to be significantly higher in the supine position (Fea et al., 2006). Modern eye trackers now are designed to dynamically follow the eye during laser ablation.

2.2.6 Tear film

The tear layer has a refractive index near to that of the cornea (1.33 versus 1.376) and refraction at the air-tear film interface accounts for the majority of refractive power of the anterior ocular surface (Oldenburg et al., 1990). Use of hard contact lenses to correct refractive errors creates a 'tear lens' in the contact lens-cornea interface which resolves the keratometric cylinder (Astin, 1989). This decouples anterior corneal astigmatism from internal astigmatism and manifests as 'residual astigmatism'. The nature of this astigmatism is frequently against the rule and at times can cause eye strain (see above).

The superior eyelid exerts pressure on the cornea, and tear accumulates over the lower eyelid margin due to gravity; this combination induces a vertical coma which may manifest as a cylinder (Montés-Micó et al., 2004a). Localized aggregation of lacrimal fluid is also caused by peripheral corneal lesions such as pterygium (Oldenburg et al., 1990; Walland et al., 1994; Yasar et al., 2003), limbal conjunctival carcinoma (Leccisotti, 2005), or nodules (Das et al., 2005). Such changes cause corneal astigmatism and are largely resolved after excision of the lesion or drying of the tear pool (Leccisotti, 2005; Yasar et al., 2003).

The tear film effect can also be discussed with regard to ocular wavefront changes during blinking. It has been agreed that higher order corneal aberrations show micro-fluctuations during the inter-blink interval. These dynamic variations of ocular surface topography have been widely investigated using high speed videokeratoscopes (E. Goto et al., 2003; T. Goto et al., 2004; Koh et al., 2002; Kojima et al., 2004; Montés-Micó et al., 2004a; Németh et al., 2002). Zhu et al. found that the height of the ocular surface increased about 2 mm within 0.5 s after blinking at the upper edge of the topography map. They also declared that absolute values

in horizontal coma and secondary astigmatism at 45° significantly increased during the inter-blink interval, while secondary astigmatism at 0° decreased considerably (Zhu et al., 2006). In another study, irregular astigmatism induced by tear film breakup was measured and significant increases were observed in coma, spherical aberration and total higher order aberrations (Koh et al., 2002).

It is therefore suggested that measurement of corneal wavefront aberrations for refractive surgery purposes should be done at a fixed interval after each blink (Montés-Micó et al., 2004b). Based on a number of studies that evaluated the variability of topography maps (Buehren et al., 2001; Iskander et al., 2005; Montés-Micó et al., 2004b), an interval of 1 to 4 seconds after blinking is suggested as the optimal time (Zhu et al., 2006).

2.2.7 Retinal astigmatism

From a historical point of view, directional variability in photoreceptor arrangement was proposed as a source of astigmatism (Mitchell et al., 1967); in other words, functional retinal elements may be more abundant or thicker in one axis than the other (Shlaer, 1937). More recently, a 'tilted' retina was simulated and it was observed to manifest as some degree of cylindrical error (Flüeler & Guyton, 1995). This could be the result of unequal lengthening of the sclera in different meridians during axial growth.

3. Conclusion

Although most of the materials presented in this chapter are of investigational interest, there is a resurging interest in these physiological issues owing to refractive surgery. Variations in tear film status, torsional alignment, and pupil features are sources of error in ocular refractive assessment and laser photoablation. Our objective should be firstly not to spoil the innate versatility and optical quality of the virgin eye; and secondly, to avoid inconsistencies in the outcome. The available optical models do not simulate optical performance of the eye perfectly, and the refractive surgery technology –in terms of diagnosis and treatment– does not fully follow our optical models either.

On the positive side, if we intend to make 'super vision' a reality (Applegate et al., 2004), we have to better understand the mentioned dynamics and interaction and 'personalize' treatments. Iris registration for example, can be used to avoid the negative effects of pupil centroid and astigmatism axis shifts during excimer laser ablation (Porter et al., 2006; Jing et al., 2008; Khalifa et al., 2009; Park et al., 2009). But this is just the beginning and we need dynamic optical models and advanced simulations to fulfil the mentioned objectives.

Additionally, an in-depth understanding of physiology of ocular astigmatism may throw light on the pathobiology of refractive errors and lead to new avenues for the prevention of clinically significant astigmatism.

4. References

- Abrahamsson, M., Fabian, G. & Sjostrand, J. (1988), 'Changes in astigmatism between the ages of 1 and 4 years: a longitudinal study', *Br J Ophthalmol*, 72 (2), 145-9.
- Akinci, A., et al. (2009), 'Refractive errors and strabismus in children with down syndrome: a controlled study', *J Pediatr Ophthalmol Strabismus*, 46 (2), 83-6.
- Applegate, R.A., Hilmantel, G. & Thibos, L.N. (2004). Assessment of visual performance, In: *Wavefront customized visual correction: The quest for supervision II*, Krueger, R.R., Applegate, R.A., Macrae, S.M., pp. (66), SLACK incorporated, ISBN: 1-55642-625-9, Thorofare.

- Artal, P., Guirao, A., Berrio, E. & Williams, D.R. (2001), 'Compensation of corneal aberrations by the internal optics in the human eye', *J Vis*, 1 (1), 1-8.
- (2002), 'Contribution of the cornea and internal surfaces to the change of ocular aberrations with age', *J Opt Soc Am A Opt Image Sci Vis*, 19 (1), 137-43.
- Asano, K., et al. (2005), 'Relationship between astigmatism and aging in middle-aged and elderly Japanese', *Jpn J Ophthalmol*, 49 (2), 127-33.
- Astin, C.L.K. (1989), 'Alternatives to toric contact lens fitting—for regular and irregular astigmatism', *Ophthalmic Physiol Opt*, 9 (3), 243-46.
- Atkinson, J., Braddick, O. & French, J. (1980), 'Infant astigmatism: its disappearance with age', *Vision Res*, 20 (11), 891-3.
- Attebo, K., Ivers, R.Q. & Mitchell, P. (1999), 'Refractive errors in an older population: the Blue Mountains Eye Study', *Ophthalmology*, 106 (6), 1066-72.
- Baldwin, W.R. & Mills, D. (1981), 'A longitudinal study of corneal astigmatism and total astigmatism', *Am J Optom Physiol Opt*, 58 (3), 206-11.
- Beau Seigneur, W. (1946), 'Changes in power and axis of cylindrical errors after convergence', *Aust J Optom*, 29 (6), 258-69.
- Becker, R. et al. (2004), 'Use of preoperative assessment of positionally induced cyclotorsion: a video-oculographic study', *Br J Ophthalmol*, 88 (3), 417-421.
- Bharti, S. & Bains, H.S. (2007), 'Active cyclotorsion error correction during LASIK for myopia and myopic astigmatism with the NIDEK EC-5000 CX III laser', *J Refract Surg*, 23 (9 Suppl), S1041-5.
- Borish, I.M. & Benjamin, W.J. (2006). Monocular and Binocular Subjective Refraction, In: *Borish's Clinical Refraction*, Benjamin, W.J., pp. (857-858), Butterworth-Heinemann, ISBN-13:978-0-7506-7524-6, Missouri.
- Bowman, K.J., Smith, G. & Carney, L.G. (1978), 'Corneal topography and monocular diplopia following near work', *J AAOPS*, 55 (12), 818-23.
- Brzezinski, M.A. (1982), 'Review: Astigmatic Accommodation (Sectional Accommodation)—a form of Dynamic Astigmatism', *Aust J Optom*, 65 (1), 5-11.
- Buehren, T., Collins, M.J. & Carney, L. (2003), 'Corneal aberrations and reading', *Optom Vis Sci*, 80 (2), 159-66.
- Buehren, T., et al. (2001), 'The stability of corneal topography in the post-blink interval', *Cornea*, 20 (8), 826-33.
- Buehren, T. et al. (2003), 'Corneal topography and accommodation', *Cornea*, 22 (4), 311-6.
- Cakmak, H.B., Cagil N, Simavli H, Duzen B & Simsek S. (2010), 'Refractive Error May Influence Mesopic Pupil Size', *Curr Eye Res*, 35 (2), 130-36.
- Campbell, F.W., Westheimer G. (1959), 'Factors influencing accommodation responses of the human eye', *J Opt Soc Am*, 49 (6), 568-71.
- Chang, J. (2008), 'Cyclotorsion during laser in situ keratomileusis', *J Cataract Refract Surg*, 34(10), 1720-6.
- Cheng, H., et al. (2004), 'A population study on changes in wave aberrations with accommodation', *J Vis*, 4 (4) 272-80.
- Chernyak, D.A. (2004), 'Cyclotorsional eye motion occurring between wavefront measurement and refractive surgery', *J Cataract Refract Surg*, 30 (3), 633-8.
- Collins, M.J., et al. (2005), 'Regression of lid-induced corneal topography changes after reading', *Optom Vis Sci*, 82 (9), 843-9.
- (2006), 'Factors influencing lid pressure on the cornea', *Eye & Contact Lens*, 32 (4), 168-73.

- Das, S., Link, B. & Seitz, B. (2005), 'Salzmann's nodular degeneration of the cornea: a review and case series', *Cornea*, 24 (7), 772-7.
- Denieul, P. (1982), 'Effects of stimulus vergence on mean accommodation response, microfluctuations of accommodation and optical quality of the human eye', *Vision Res*, 22 (5), 561-69.
- Dobson, V., Fulton, A. B. & Sebris, S. L. (1984), 'Cycloplegic refractions of infants and young children: the axis of astigmatism', *Invest Ophthalmol Vis Sci*, 25 (1), 83-7.
- Ehrlich, D. L., et al. (1997), 'Infant emmetropization: longitudinal changes in refraction components from nine to twenty months of age', *Optom Vis Sci*, 74 (10), 822-43.
- Fea, A. M., et al. (2006), 'Cyclotorsional eye movements during a simulated PRK procedure', *Eye (Lond)*, 20(7), 764-8.
- Ford, J.G., et al. (1997), 'Bilateral monocular diplopia associated with lid position during near work', *Cornea*, 16 (5), 525-30.
- Fotedar, R., Mitchell, P., Burlutsky, G. & Wang, J.J. (2008), 'Relationship of 10-year change in refraction to nuclear cataract and axial length findings from an older population', *Ophthalmology*, 115 (8), 1273-8.
- Friling, R., et al. (2004), 'Keratometry measurements in preterm and full term newborn infants', *Br J Ophthalmol*, 88 (1), 8-10.
- Flüeler, U. R. & D. L. Guyton. (1995), 'Does a tilted retina cause astigmatism? The ocular imagery and the retinoscopic reflex resulting from a tilted retina', *Surv Ophthalmol*, 40 (1), 45-50.
- Garcia, M.L., et al. (2003), 'Relationship between the axis and degree of high astigmatism and obliquity of palpebral fissure', *J AAPOS*, 7 (1), 14-22.
- Golnik, K.C. & Eggenberger, E. (2001), 'Symptomatic corneal topographic change induced by reading in downgaze', *J Neuroophthalmol*, 21 (3), 199-204.
- Goss, DA (1991), 'Childhood myopia', In: *Refractive Anomalies: Research and Clinical Applications*, Grosvenor, T.P., Flom, M.C., pp. (81-100), Butterworth-Heinmann, Boston.
- Goto, E., et al. (2003), 'Tear evaporation dynamics in normal subjects and subjects with obstructive meibomian gland dysfunction', *Invest Ophthalmol Vis Sci*, 44 (2), 533-9.
- Goto, T., et al. (2001), 'Gender- and age-related differences in corneal topography', *Cornea*, 20 (3), 270-6.
- (2004), 'Evaluation of the tear film stability after laser in situ keratomileusis using the tear film stability analysis system', *Am J Ophthalmol*, 137 (1), 116-20.
- Graf, E. W., J. S. Maxwell, et al. (2002), 'Changes in cyclotorsion and vertical eye alignment during prolonged monocular occlusion', *Vision Res*, 42 (9), 1185-94.
- Gullstrand, A (1962), The cornea, In: *Helmoltz's Treatise on Physiological Optics*, Helmholtz, H.V., Southal, J.P.C., Wade, N., Vol, 1, Dover, New York.
- Gwiazda, J., et al. (1984), 'Astigmatism in children: changes in axis and amount from birth to six years', *Invest Ophthalmol Vis Sci*, 25 (1), 88-92.
- Gwiazda, J., et al. (2000), 'Astigmatism and the development of myopia in children', *Vision Res*, 40 (1), 1019-26.
- Hammond, C. J., et al. (2001), 'Genes and environment in refractive error: the twin eye study', *Invest Ophthalmol Vis Sci*, 42 (6), 1232-6.
- Hirsch, M. J. (1963), 'Changes in astigmatism during the first eight years of school--an interim report from the Ojai longitudinal study', *Am J Optom Arch Am Acad Optom*, 40, 127-32.
- Ho, J. D., et al. (2010), 'Effects of aging on anterior and posterior corneal astigmatism', *Cornea*, 29 (6), 632-7.

- Hofstetter, H. W. & Baldwin, W. (1957), 'Bilateral correlation of residual astigmatism', *Am J Optom Arch Am Acad Optom*, 34 (7), 388-91.
- Hori-Komai, Y. et al. (2007), 'Detection of cyclotorsional rotation during excimer laser ablation in LASIK', *J Refract Surg*, 23(9), 911-15.
- Howland HC. (1982), 'Infant eyes: optics and accommodation', *Curr Eye Res*, 2(3), 217-24.
- Howland, H. C. & Sayles, N. (1985), 'Photokeratometric and photorefractive measurements of astigmatism in infants and young children', *Vision Res*, 25 (1), 73-81.
- Hu, J. R., et al. (2004), '[Higher-order aberrations in myopic and astigmatism eyes]', *Zhonghua Yan Ke Za Zhi*, 40 (1), 13-6.
- Huynh, S. C., et al. (2007), 'Astigmatism in 12-year-old Australian children: comparisons with a 6-year-old population', *Invest Ophthalmol Vis Sci*, 48 (1), 73-82.
- Isenberg, S. J., et al. (2004), 'Corneal topography of neonates and infants', *Arch Ophthalmol*, 122 (12), 1767-71.
- Iskander, D.R., Collins, M.J. & Davis, B. (2005), 'Evaluating tear film stability in the human eye with high-speed videokeratoscopy', *IEEE Trans Biomed Eng*, 52 (11), 1939-49.
- Jing, Z. et al. (2008), 'Comparison of visual performance between conventional LASIK and wavefront-guided LASIK with iris-registration', *Chin Med*, 121 (2), 137-142.
- Kame, R. T., Jue, T. S. & Shigekuni, D. M. (1993), 'A longitudinal study of corneal astigmatism changes in Asian eyes', *J Am Optom Assoc*, 64 (3), 215-9.
- Karesh, J.W. (1994), "Topographic Anatomy of the eye: an overview", In: *Duane's Foundations of Clinical Ophthalmology*, Jeager, E.A., Tasman, W., pp.(1-30), JB Lippincott, Philadelphia.
- Khalifa, M. et al. (2009). 'Iris registration in wavefront-guided LASIK to correct mixed astigmatism', *Journal Cataract Refract Surg*, 35 (3), 433-37.
- Kiely, P.M., Carney, L.G. & Smith, G. (1982), 'Diurnal variations of corneal topography and thickness', *Am Journal Optom Physiol Opt*, 59 (12), 976-82.
- Knoll, HA (1975), 'Letter: bilateral monocular diplopia after near work', *Am Journal Optom Physiol Opt*, 52 (2), 139.
- Koh, S., et al. (2002), 'Effect of tear film break-up on higher-order aberrations measured with wavefront sensor', *Am J Ophthalmol*, 134 (1), 115-17.
- Kohl, P. & Samek, M. (1988), 'Refractive error and preferential looking visual acuity in infants 12-24 months of age: year 2 of a longitudinal study', *J Am Optom Assoc*, 59 (9), 686-90.
- Kojima, T., et al. (2004), 'A new noninvasive tear stability analysis system for the assessment of dry eyes', *Invest ophthalmolvisci*, 45 (5), 1369-74.
- Leccisotti, A. (2005), 'Corneal topographic changes in a case of limbal conjunctival carcinoma', *Cornea*, 24 (8), 1021-3.
- Lin, L. L., et al. (1996), 'Changes in ocular refraction and its components among medical students--a 5-year longitudinal study', *Optom Vis Sci*, 73 (7), 495-8.
- Little, J. A., Woodhouse, J. M. & Saunders, K. J. (2009), 'Corneal power and astigmatism in Down syndrome', *Optom Vis Sci*, 86 (6), 748-54.
- Lopping, B. & Weale, RA (1965), 'Changes in corneal curvature following ocular convergence', *Vision Res*, 5 (4-5), 207-15.
- Lyle, WM, Grosvenor, T. & Dean, KC (1972), 'Corneal astigmatism in Amerind children', *Am J Optom Arch Am Acad Optom*, 49 (6), 517-24.
- Manchester, P. T., Jr. (1970), 'Hydration of the cornea', *Trans Am Ophthalmol Soc*, 68, 425-61.
- Mandell, R.B. (1966), 'Bilateral monocular diplopia following near work', *Am J Optom Arch Am Acad Optom*, 43 (8), 500-4.

- Marin-Amat, M. (1956), '[Physiological variations of corneal curvature during life time; their importance and transcendence into ocular refraction]', *Bull SocBelgeOphthalmol*, 113, 251-93.
- Mehdizadeh, M. (2008), 'Age and refraction', *Ophthalmology*, 115 (11), 2097; author reply 2097-8.
- Mitchell, D. E. et al. (1967), 'Effect of orientation on the modulation sensitivity for interference fringes on the retina', *J Opt Soc Am*, 57(2), 246-49.
- Montés-Micó, R., et al. (2004a), 'Temporal changes in optical quality of air-tear film interface at anterior cornea after blink', *Invest ophthalmolvisci*, 45 (6), 1752-57.
- (2004b), 'Postblink changes in total and corneal ocular aberrations', *Ophthalmology*, 111 (4), 758-67.
- Morgan, I. G., Rose, K. A. & Ellwein, L. B. (2010), 'Is emmetropia the natural endpoint for human refractive development? An analysis of population-based data from the refractive error study in children (RESC)', *Acta Ophthalmol*, 88 (8), 877-84.
- Mutti, D. O., et al. (2004), 'Refractive astigmatism and the toricity of ocular components in human infants', *Optom Vis Sci*, 81 (10), 753-61.
- Németh, J., et al. (2002), 'High-speed videotopographic measurement of tear film build-up time', *Invest ophthalmolvisci*, 43 (6), 1783-90.
- Neuhann, I. M. et al. (2010), 'Static and dynamic rotational eye tracking during LASIK treatment of myopic astigmatism with the Zyoptix laser platform and Advanced Control Eye Tracker', *J Refract Surg*, 26(1), 17-27.
- Oldenburg, J.B., et al. (1990), 'Mechanism of Corneal Topographic Changes', *Cornea*, 9 (3), 200-4.
- Osuoben, E.P. & Al Mijalli, M.H. (1997), 'Association between eyelid thickness and corneal astigmatism', *Clin Ex Optom*, 80 (1), 35-9.
- Park, S. H., M. Kim, et al. (2009). 'Measurement of pupil centroid shift and cyclotorsional displacement using iris registration', *Ophthalmologica*, 223 (3), 166-71.
- Porter, J. et al. (2006). 'Aberrations induced in wavefront-guided laser refractive surgery due to shifts between natural and dilated pupil center locations', *J Cataract Refract Surg*, 32 (1), 21-32.
- Read, S. A., Collins, M. J. & Carney, L. G. (2005), 'The diurnal variation of corneal topography and aberrations', *Cornea*, 24 (6), 678-87.
- Read, S.A., Collins, M.J. & Carney, L.G. (2007a), 'The influence of eyelid morphology on normal corneal shape', *Invest ophthalmolvisci*, 48 (1), 112-9.
- (2007b), 'A review of astigmatism and its possible genesis', *Clinical and Experimental Optometry*, 90 (1), 5-19.
- (2007c), 'Influence of accommodation on the anterior and posterior cornea', *J Cataract Refract Surg* 33 (11), 1877-85.
- Regan, D. & H. Spekreijse (1970), 'Electrophysiological correlate of binocular depth perception in man', *Nature*, 3 (5227), 92-4.
- Rosenfield, M. & B. Gilmartin (1988), 'The effect of vergence adaptation on convergent accommodation', *Ophthalmic Physiol Opt*, 8 (2), 172-77.
- Sakai, H., Hirata, Y. & Usui, S. (2007), 'Relationship between residual aberration and light-adapted pupil size', *Optom Vis Sci*, 84 (6), 517-21.
- Saunders, H. (1986), 'Changes in the orientation of the axis of astigmatism associated with age', *Ophthalmic Physiol Opt*, 6 (3), 343-4.
- (1988), 'Changes in the axis of astigmatism: a longitudinal study', *Ophthalmic Physiol Opt*, 8 (1), 37-42.
- Sawada, A., et al. (2008), 'Refractive errors in an elderly Japanese population: the Tajimi study', *Ophthalmology*, 115 (2), 363-70 e3.

- Shankar, S. & Bobier, W. R. (2004), 'Corneal and lenticular components of total astigmatism in a preschool sample', *Optom Vis Sci*, 81 (7), 536-42.
- Shaw, A.J., et al. (2008), 'Corneal refractive changes due to short-term eyelid pressure in downward gaze', *Journal Cataract Refract Surg*, 34 (9), 1546-53.
- Shen, E.P. et al. (2010), 'Manual limbal markings versus iris-registration software for correction of myopic astigmatism by laser in situ keratomileusis', *J Cataract Refract Surg*, 36(3), 431-6.
- Shilo, S. (1977), "Astigmatism of the mammalian cornea: evolutionary and perceptive significance", *Documenta Ophthalmologica*, 44(2), 403-419.
- Shlaer, S. (1937), 'The relation between visual acuity and illumination', *J Gen Physiol*, 21 (2), 165-88.
- Stirling, A. W. (1920), 'The Influence of Age Upon the Axis of Astigmatism', *Br J Ophthalmol*, 4 (11), 508-10.
- Swami, A.U. et al. (2002), 'Rotational malposition during laser in situ keratomileusis', *Am J Ophthalmol*, 133(4), 561-2.
- Tait, E.F. (1933), 'A reciprocal reflex system in the accommodation-convergence relationships', *Am J Psychol*, 45 (4), 647-62.
- Tjon-Fo-Sang, M. J. et al. (2002), 'Cyclotorsion: a possible cause of residual astigmatism in refractive surgery', *J Cataract Refract Surg*, 28 (4), 599-602.
- Tsukamoto, M., et al. (2000), 'Accommodation causes with-the-rule astigmatism in emmetropes', *Optom Vis Sci*, 77 (3), 150-5.
- (2001), 'The binocular accommodative response in uncorrected ametropia', *Optom Vis Sci*, 78 (10), 763-8.
- Ukai, K. & Ichihashi, Y. (1991), 'Changes in ocular astigmatism over the whole range of accommodation', *Optom Vis Sci*, 68 (10), 813-8.
- Varghese, R. M., Sreenivas, V., Puliye, J.M., & Varughese, S. (2009), 'Refractive status at birth: its relation to newborn physical parameters at birth and gestational age', *PLoS One*, 4 (2), e4469.
- Varughese, S., Varghese, R.M., Gupta, N., Ojha, R., Sreenivas V. & Puliye J.M., (2005), 'Refractive error at birth and its relation to gestational age', *Curr Eye Res*, 30 (6), 423-8.
- Walland, M.J., Stevens, J.D. & Steele, A.D.M. (1994), 'The effect of recurrent pterygium on corneal topography', *Cornea*, 13 (5), 463-4.
- Walsh, G. (1988). 'The effect of mydriasis on the pupillary centration of the human eye', *Ophthalmic Physiol Opt*, 8 (2), 178-82.
- Wang, F. M., Millman, A. L., Sidoti, P. A. & Goldberg, R. B. (1990), 'Ocular findings in Treacher Collins syndrome', *Am J Ophthalmol*, 110 (3), 280-6.
- Wilson, G., Bell, C. & Chotai, S. (1982), 'The effect of lifting the lids on corneal astigmatism', *Am J OptomPhysiol Opt*, 59 (8), 670-4.
- Wilson, M. A., Campbell, M. C. & Simonet, P. (1992). 'Change of pupil centration with change of illumination and pupil size', *Optom Vis Sci*, 69 (2), 129-36.
- Wood, I. C., Hodi, S. & Morgan, L. (1995), 'Longitudinal change of refractive error in infants during the first year of life', *Eye*, 9 (5), 551-7.
- Yasar, T., Ozdemir, M., Cinal, A., Demirok, A., Ilhan, B. & Durmus, A. (2003), 'Effects of fibrovascular traction and pooling of tears on corneal topographic changes induced by pterygium', *Eye*, 17 (4), 492-96.
- Zhu, M., Collins, M.J. & Iskander, D.R. (2006), 'Dynamics of ocular surface topography', *Eye*, 21 (5), 624-32.

Astigmatism - Optics, Physiology and Management

Edited by Dr. Michael Goggin

ISBN 978-953-51-0230-4

Hard cover, 308 pages

Publisher InTech

Published online 29, February, 2012

Published in print edition February, 2012

This book explores the development, optics and physiology of astigmatism and places this knowledge in the context of modern management of this aspect of refractive error. It is written by, and aimed at, the astigmatism practitioner to assist in understanding astigmatism and its amelioration by optical and surgical techniques. It also addresses the integration of astigmatism management into the surgical approach to cataract and corneal disease including corneal transplantation.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Seyed-Farzad Mohammadi, Maryam Tahvildari and Hadi Z-Mehrjardi (2012). Physiology of Astigmatism, Astigmatism - Optics, Physiology and Management, Dr. Michael Goggin (Ed.), ISBN: 978-953-51-0230-4, InTech, Available from: <http://www.intechopen.com/books/astigmatism-optics-physiology-and-management/astigmatism-physiology>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](https://creativecommons.org/licenses/by/3.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen