

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

186,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Photo- and Free Radical-Mediated Oxidation of Lipid Components During the Senescence of Phototrophic Organisms

Jean-François Rontani

*Laboratory of Microbiology, Geochemistry and Marine Ecology (UMR 6117),
Center of Oceanology of Marseille, Aix-Marseille University, Campus of Luminy, Marseille,
France*

1. Introduction

Recently, the role played by photochemical and free radical-mediated processes in the degradation of lipid components during the senescence of phototrophic organisms was investigated. The present paper reviews the results obtained in the course of these studies.

In a first part, visible and UV light-induced photooxidation of the main lipid cell components (chlorophylls, carotenoids, sterols, unsaturated fatty acids, highly branched isoprenoid and linear alkenes, alkenones, cuticular waxes ...) in senescent phototrophic organisms (phytoplankton, cyanobacteria, higher plants, purple sulfur bacteria and aerobic anoxygenic phototrophic bacteria) is examined. Probably due to its long lifetime in hydrophobic micro-environments and thus in senescent cells, singlet oxygen plays a key role in the photodegradation of most of the lipid components.

The second part of this paper describes the free radical oxidation (autoxidation) of lipid components during the senescence of phototrophic organisms, which have been virtually ignored until now in the literature. In senescent phototrophic organisms, the mechanism of initiation of free-radical oxidation seems to be the homolytic cleavage (catalyzed by some metal ions) of photochemically produced hydroperoxides. It was also demonstrated recently that viral infection and autocatalytic programmed cell death could also lead to elevated production of reactive oxygen species (ROS) able to induce the degradation of cell components.

2. Photodegradation processes in phototrophic organisms

Several works suggested photo-oxidation as an important sink of organic matter in the photic layer of oceans (Zafiriou, 1977; Zafiriou et al., 1984). However, due to the lack of suitable markers this phenomenon has never been fully addressed. Owing to the problem of stratospheric ozone depletion, some studies have recently examined the degradative effects of enhanced UV-B doses on phytoplanktonic lipids (He and Häder, 2002). However, photochemical damages in phytoplanktonic cells are not a monopoly of UV radiation. In fact, due to the presence of chlorophylls (which are very efficient photosensitizers (Foote, 1976; Knox and Dodge, 1985)), numerous organic components of phytoplankton are susceptible to being photodegraded during senescence by photosynthetically active radiation (PAR).

2.1 Photodegradation of the main lipidic components of phytoplankton during senescence

When a chlorophyll molecule absorbs a quantum of light energy, an excited singlet state (^1Chl) is formed which, in healthy cells, leads predominantly to the characteristic fast reactions of photosynthesis (Foote, 1976). However, a small proportion (<0.1%) undergoes intersystem crossing (ISC) to form the longer lived triplet state (^3Chl ; Knox and Dodge, 1985). ^3Chl is not only itself potentially damaging in type I reactions (hydrogen atom or electron abstraction) (Knox and Dodge, 1985), but can also generate highly reactive oxygen species (ROS) and, in particular, singlet oxygen ($^1\text{O}_2$), by reaction with ground state oxygen ($^3\text{O}_2$) via Type II processes. In order to avoid oxidative damage, there are many antioxidant protective mechanisms in chloroplasts. Carotenoids quench ^3Chl and $^1\text{O}_2$ by energy transfer mechanisms at very high rates (Foote, 1976) and tocopherols can remove $^1\text{O}_2$, $\text{O}_2^{\bullet-}$, HOO^{\bullet} and HO^{\bullet} by acting as sacrificial scavengers (Halliwell, 1987). Superoxide dismutase enzyme (SOD) and ascorbic acid may also scavenge $\text{O}_2^{\bullet-}$ (Halliwell, 1987), while catalase activity decreases H_2O_2 levels.

In senescent phototrophic organisms, the fast reactions of photosynthesis clearly do not operate, so an accelerated rate of formation of ^3Chl and $^1\text{O}_2$ would be expected (Nelson, 1993). The rate of formation of these potentially damaging species can then exceed the quenching capacity of the photoprotective system and photodegradation can occur (photodynamic effect; Merzlyak and Hendry, 1994). In phytodetritus, when the ordered structure of the thylakoid membranes has been disrupted, pigments tend to remain associated with other hydrophobic cellular components such as membrane lipids (Nelson, 1993). As a result, the photooxidative effect of chlorophyll sensitization might be strongly amplified within such a hydrophobic micro-environment. Moreover, the lifetime of $^1\text{O}_2$ produced from sensitizers in a lipid-rich hydrophobic environment could be longer, and its potential diffusive distance greater, than its behaviour in aqueous solution (Suwa et al., 1977). It is not surprising, therefore, that photodegradation processes act on the majority of unsaturated lipid components of senescent phytoplankton.

2.1.1 Chlorophylls

Irradiation of dead phytoplankton cells by PAR and UVR radiations results in rapid degradation of chlorophylls (Nelson, 1993; Rontani et al., 1995; Christodoulou et al., 2010). Photodegradation of chlorophyll-*a* and -*c* in killed cells of *E. huxleyi* appeared to be induced by both PAR and UVR (Christodoulou et al., 2010). The photochemical degradation of chlorophylls has so far been studied almost exclusively with respect to the macrocycle moiety of the molecule, which is the more reactive. Despite some progress regarding intermediary photoproducts (Engel et al., 1991; Iturraspe et al., 1994), no stable and specific markers for the chlorophyll macrocycle photodegradation have been characterised.

The isoprenoid phytyl side-chain of chlorophylls is also sensitive to photochemical processes. In fact, in phytodetritus, the photodegradation rates were only 3 to 5 times higher for the chlorophyll tetrapyrrolic structure than for the phytyl side-chain (Cuny et al., 1999; Christodoulou et al., 2010). Analysis of isoprenoid photoproducts of chlorophylls after irradiation of different dead phytoplanktonic cells by visible light clearly established that the photodegradation of the chlorophyll phytyl side-chain in phytodetritus involved mainly

Irradiation with UVR resulted in the additional production of small amounts of *Z*-phytol and *Z* and *E*-3,7,11,15-tetramethylhexadec-3-en-1,2-diols (**3,4**) (Christodoulou et al., 2010). The detection of *Z*-phytol allowed to demonstrate the induction of *cis-trans* photosensitized isomerization by UVR. These reactions probably involve triplet states of ketones as sensitizers. Type II photosensitized oxidation of the *Z* configuration of phytol, which should lead to the production of photoproducts of structures **a**, **b** and **c** (Fig. 1) (Schulte-Elte et al., 1979), explains the detection of small amounts of *Z* and *E*-3,7,11,15-tetramethylhexadec-3-en-1,2-diols (**3,4**) after irradiation with UVR. Irradiation with UVR also resulted in a faster degradation of chlorophyll phytyl side-chain oxidation products (Christodoulou et al., 2010). This higher reactivity was attributed to UVR-induced homolysis of the peroxy group of photoproducts of structures **a**, **b** and **c** (Fig. 1).

Phytyldiol (**2**) is ubiquitous in the marine environment and has been proposed as tracer for photodegradation of chlorophyll's phytyl side chain (Rontani et al. 1994; 1996a; Cuny and Rontani 1999). Further, the molar ratio phytyldiol:phytol (Chlorophyll Phytyl side-chain Photodegradation Index, CPPI) was employed to estimate the extent of chlorophyll photodegraded in natural marine samples by the empirical equation: chlorophyll photodegradation % = $(1 - (\text{CPPI} + 1)^{-18.5}) \times 100$ (Cuny et al. 2002).

2.1.2 Carotenoids

In phytodetritus, chlorophylls and carotenoids remain in a close molecular-scale association at relatively high localized concentrations, even though the structure of the thylakoid membrane has been disrupted (Nelson, 1993). Thus, the sensitized photooxidation of carotenoids is enhanced. The photosensitized oxidation (involving $^1\text{O}_2$) of carotenoids in solvents has been studied (Iseo et al., 1972) and loliolide (**5**), *iso*-loliolide (**6**) and dihydroactinidiolide (**7**) (Fig. 2) were identified as major photoproducts, depending on the functionality of carotenoids at C-3. Loliolide (**5**) and *iso*-loliolide (**6**) have been detected in killed cells of *Dunaliella* sp. irradiated by visible light (Rontani et al., 1998). However, due to their apparent production by anaerobic bacteria (Repeta, 1989) and during dark incubations of killed phytoplanktonic cells (Rontani et al., 1998), these compounds cannot constitute unequivocal indicators of photooxidative processes.




Fig. 2. Structure of the main carotenoid oxidation products.

2.1.3 Δ^5 -sterols

As important unsaturated components of biological membranes, Δ^5 -sterols are highly susceptible to photooxidative degradation during the senescence of phytoplankton. Irradiation by visible light of killed cells of *Skeletonema costatum*, *Dunaliella* sp.,

Phaeodactylum tricornutum and *Emiliania huxleyi* (Rontani et al., 1997a; 1997b; 1998) resulted in a quick photodegradation of the sterol components of these algae. The results obtained clearly established that the photooxidation of sterols in senescent cells of phytoplankton involves type II photoprocesses. These processes mainly produce Δ^6 -5 α -hydroperoxides (**8**) and to a lesser extent Δ^4 -6 α /6 β -hydroperoxides (**9** and **10**) (Fig. 3) (Nickon and Bagli, 1961; Kulig and Smith, 1973). Δ^6 -5 α -hydroperoxysterols (**8**) are relatively unstable and may undergo allylic rearrangement to Δ^5 -7 α -hydroperoxysterols (**11**), which in turn epimerize to the corresponding 7 β -hydroperoxides (**12**) (Fig. 3) (Smith, 1981). It was previously demonstrated that during singlet oxygen-mediated photooxidation of sterols in biological membranes (Korytowski *et al.*, 1992) and senescent phytoplanktonic cells (Rontani *et al.*, 1997a) the photogeneration of Δ^4 -6 α /6 β -hydroperoxides (**9** and **10**) was more favourable than in homogeneous solution (ratio Δ^4 -6 α /6 β -hydroperoxides/ Δ^6 -5 α -hydroperoxysterols ranging from 0.30 to 0.35 instead of 0.1).




Fig. 3. Type II photosensitized oxidation of Δ^5 sterols.

Allylic rearrangement of Δ^6 -5 α -hydroperoxides (**8**) appeared to take place very weakly in senescent phytoplanktonic cells (Rontani et al., 1997a; 1997b; 1998). This surprising stability was attributed by Korytowski et al. (1992) either to hydrogen bonding between the unsaturated fatty acyl chain of phospholipids and Δ^6 -5 α -hydroperoxides (**8**) which could hinder the allylic rearrangement, or to differences of polarity in the carbon 7-10 zone of the fatty acyl chain (where sterols tend to localize in phospholipid/sterol bilayers (MacIntosh, 1978)). It is also interesting to note that the reduction of hydroperoxysterols to the corresponding diols weakly operates in killed phytoplanktonic cells (Rontani et al., 1997a).

Δ^6 -5 α -Hydroperoxysterols (**8**) are potential type II photodegradation markers, not only because they are the major products of singlet oxygen attack on the steroidal Δ^5 -3 β - system, but also because biological functionalization of steroids at C-5 is rare. Unfortunately, if these compounds are particularly stable in phytodetritus, they decay slowly in the sediment to their corresponding Δ^5 -7 α / β -derivatives (**11** and **12**) (Rontani and Marchand, 2000), which are not selective markers (see chapter 3.3). Moreover, according to the stability of the alkyl radicals formed during β -scission of the corresponding alkoxy radicals, the following order of stability was proposed: Δ^4 -6-hydroperoxysterols (**9** and **10**) > Δ^5 -7-hydroperoxysterols (**11** and **12**) > Δ^6 -5-hydroperoxy-sterols (**8**) (Christodoulou et al., 2009). Consequently, Δ^4 -6 α / β -hydroperoxysterols (**9** and **10**) (or their degradative products Δ^4 -6 α / β -hydroxysterols and Δ^4 -6 α / β -oxosterols) may be considered as more reliable *in situ* markers of type II photodegradation processes than Δ^6 -5 α -hydroperoxides (**8**).

2.1.4 Unsaturated fatty acids

Chloroplast membrane components are particularly susceptible to type II photooxidation (Heath and Packer, 1968). This is the case notably for unsaturated fatty acids, which generally predominate in algal lipids, particularly in the photosynthetic membranes (Woods, 1974). In killed phytoplanktonic cells, the photodegradation rates of unsaturated fatty acids logically increase with their unsaturation degree (Rontani et al., 1998). Singlet oxygen-mediated photooxidation of monounsaturated fatty acids involves a direct reaction of $^1\text{O}_2$ with the carbon-carbon double bond by a concerted 'ene' addition (Frimer 1979) and leads to formation of hydroperoxides at each carbon of the original double bond. Thus, photooxidation of oleic acid produces a mixture of 9- and 10-hydroperoxides with an allylic *trans*-double bond (Frankel et al. 1979; Frankel, 1998), which can subsequently undergo stereoselective radical allylic rearrangement to 11-*trans* and 8-*trans* hydroperoxides, respectively (Porter et al. 1995) (Fig. 4).

The free radical nature of the allylic hydroperoxide rearrangement is supported by the observation that the rearrangement is catalysed by free radical initiators or light and inhibited by phenolic antioxidants (Porter et al., 1995). This allylic rearrangement weakly intervenes in most of the killed phytoplanktonic cells examined (Rontani et al., 1998). This was attributed to the relatively high localized fatty acid concentrations present in phytodetritus (Nelson, 1993), which favoured the dimerisation of hydroperoxides. Hydrogen atom abstraction to form allylperoxyl radicals does indeed occur readily from hydroperoxide monomers but not from hydroperoxide dimers (Porter et al., 1995).

During early diagenesis, isomeric hydroperoxyacids undergo heterolytic cleavage to aldehydes and ω -oxocarboxylic acids (Frimer, 1979) or homolytic cleavage and subsequent transformation to the corresponding alcohols or ketones (Fig. 5).

Taking into account the high amounts of photoproducts of mono-unsaturated fatty acids detected in the particulate matter samples (Marchand and Rontani, 2001; Christodoulou et al., 2009; Rontani et al., 2011a), and the well known increasing photooxidation rates of fatty acids with their degree of unsaturation (Frankel., 1998), it can be concluded that considerable amounts of poly-unsaturated fatty acids must be photooxidized during the senescence of phytoplankton in the marine environment. However, at this time photooxidation products of this kind of fatty acids could not be detected in natural samples.




Fig. 4. Type II photosensitized oxidation of oleic acid.




Fig. 5. Degradation of allylic hydroperoxides resulting from Type II photosensitized oxidation of monounsaturated fatty acids (the example given is this of 9-hydroperoxyoctadec-10-enoic acid) (RH = hydrogen donors, e.g. lipids or reduced sensitizers).

This is possibly due to: (i) the instability of the hydroperoxides formed, or (ii) the involvement of cross-linking reactions leading to the formation of macromolecular structures (Neff et al., 1988) non-amenable by gas chromatography.

2.1.5 Alkenones

Alkenones are a class of mono-, di-, tri-, tetra- and penta-unsaturated C₃₅-C₄₀ methyl and ethyl ketones (Boon et al., 1978; Volkman et al., 1980; de Leeuw et al., 1980; Marlowe et al., 1984; Prahl et al., 2006; Jaraula et al., 2010), which are produced by certain marine haptophytes. *Emiliana huxleyi* and *Gephyrocapsa oceanica* are the major sources of alkenones in the open ocean (Volkman et al., 1980; 1995; Conte et al., 1994). The unsaturation ratio of C₃₇ alkenones, defined as $U_{37}^{K'} = [C_{37:2}] / ([C_{37:2}] + [C_{37:3}])$ where [C_{37:2}] and [C_{37:3}] are the concentrations of di- and tri-unsaturated C₃₇ alkenones respectively, varies positively with the growth temperature of the alga (Prahl and Wakeham, 1987; Prahl et al., 1988). The $U_{37}^{K'}$ - growth temperature relationship in haptophyte algae and transferred to sinking marine particulate matter leads to a linear relationship between sedimentary C₃₇ alkenone composition and mean annual SST records throughout the oceans (Rosell-Melé et al., 1995; Müller et al., 1998). The $U_{37}^{K'}$ index is now routinely used for paleotemperature reconstruction.

For alkenones to be useful as measures of sea surface temperature in the geological record, it is essential that any effects of degradation in the water column and in sediments either do not affect the temperature signal established during their initial biosynthesis by the alga (Harvey, 2000; Grimalt et al., 2000), or if there is a change its extent can be reasonably estimated.

Visible light-induced photodegradation of these compounds was thus previously investigated in order to determine if photochemical processes could appreciably modify $U_{37}^{K'}$ ratios during algal senescence (Rontani et al., 1997b; Mouzdahir et al., 2001; Christodoulou et al., 2010). Though potentially selective, photochemical degradation of alkenones is not fast enough in killed cells of *E. huxleyi* to induce strong modifications of the $U_{37}^{K'}$ ratio before the photodestruction of the photosensitizing substances (Rontani et al., 1997b; Mouzdahir et al., 2001). UVR also appeared to be inefficient to alter the $U_{37}^{K'}$ ratio (Christodoulou et al., 2010).

This stability was attributed to the *trans* configuration of alkenone double bonds (Rechka and Maxwell, 1988) that is 7 to 10 times less sensitive against singlet oxygen-mediated oxidation than the classical *cis* configuration of fatty acids (Hurst et al., 1985). This may explain the difference of photoreactivity observed between the alkenones and fatty acids with the same number of unsaturations. We also previously attributed the poor photoreactivity of alkenones to a localisation of these compounds elsewhere than in cell membranes (Rontani et al., 1997b; Mouzdahir et al., 2001), which could significantly decrease the likelihood of interaction between singlet oxygen and alkenones. Although this hypothesis is well supported by the recent results of Eltgroth et al. (2005), who demonstrated that alkenones are mainly localized into cytoplasmic vesicles, the migration of singlet oxygen from phytodetritus to attached heterotrophic bacteria previously observed (Rontani et al., 2003a; Christodoulou et al., 2010) strongly suggests a diffusion of this excited form of oxygen also in these cytoplasmic vesicles.

2.1.6 n-Alkenes

The visible light-induced degradation of *n*-alkenes was previously investigated in killed cells of the Prymnesiophyceae *E. huxleyi* and the Eustigmatophyceae *Nannochloropsis salina* (Mouzdahir et al., 2001).

In *E. huxleyi* killed cells, minor C₃₁ and C₃₃ *n*-alkenes were strongly photodegraded, while the major C₃₇ and C₃₈ *n*-alkenes appeared particularly recalcitrant towards photochemical processes. These strong differences of photoreactivity imply distinct biological syntheses and/or functions for these two groups of hydrocarbons in *E. huxleyi* cells. Interestingly, the stereochemistry of the internal double bonds in C₃₁ and C₃₃ *n*-alkenes has been established to be *cis*, while C₃₇ and C₃₈ alkenes internal double bonds exhibit a *trans* geometry (Rieley et al., 1998; Grossi et al., 2000). The photochemical recalcitrance of C₃₇ and C₃₈ *n*-alkenes could thus be partly attributed to the *trans* geometry of their internal double bonds.

Irradiation of dead cells of *N. salina* resulted in a strong modification of the hydrocarbon fraction. It did not provide evidence of a significant light-dependent degradation of monounsaturated hydrocarbons; this result was attributed to the terminal position of the double bond in these compounds (Gelin et al., 1997), which is poorly reactive towards singlet oxygen (Hurst et al., 1985). In contrast, di-, tri-, and tetraenes were strongly photodegraded during irradiation. The visible light-dependent degradation of phytoplanktonic *n*-alkenes showed apparent second-order kinetics with respect to light exposure and the half-life doses obtained logically decrease with increasing number of double bonds in these compounds (Mouzdahir et al., 2001).

2.1.7 Highly branched isoprenoid (HBI) alkenes

HBI alkenes are widely distributed in aquatic environments (Rowland and Robson, 1990; Sinninghe-Damsté et al., 2004), although they appear to originate from a relatively small number of diatomaceous algae including *Haslea* spp., *Rhizosolenia* spp., *Pleurosigma* spp. and *Navicula* spp. (Volkman et al., 1994; Sinninghe-Damsté et al., 2004; Belt et al., 2000, 2001; Allard et al., 2001; Grossi et al., 2004). Despite this, they have been commonly reported in marine sediments worldwide and provide some insight into the deposition of organic matter from the water column. One HBI alkene, a mono-unsaturated isomer termed IP₂₅, has been used as a proxy for the occurrence of spring sea ice in the Arctic (e.g. Belt et al., 2007, 2010; Massé et al., 2008).

Examination of the photoreactivity of several mono-, di-, tri- and tetra-unsaturated HBI alkenes in the presence of a photosensitizer solution and in dead cells of *H. ostrearia* allowed to show that HBI alkenes possessing at least one tri-substituted double bond may be photo-oxidized at similar or higher rates compared to other highly reactive lipids (e.g. PUFAs, vitamin E and chlorophyll *a*) during the senescence of diatom cells (Rontani et al., 2011b). As a consequence, it is proposed that HBI alkenes possessing trisubstituted double bonds are likely to be susceptible to photodegradation within the euphotic zone. In contrast, HBIs containing only mono- and di-substituted double bonds were found to be significantly less reactive towards ¹O₂ and should, therefore, be relatively preserved during sedimentation through the water column (Rontani et al., 2011b). The kinetic experiments are supported by product analysis, which revealed that the main reaction with ¹O₂ primarily occurs with the trisubstituted double bonds of HBI alkenes affording tertiary and secondary allylic hydroperoxides (Fig. 6). In contrast, the extremely low photoreactivity of the HBI monoene

IP₂₅, can be attributed to its containing only the least photochemically reactive double bond. This lack of reactivity supports (in part) the good preservation of IP₂₅ generally observed in sediments (Belt et al., 2007, 2010; Massé et al., 2008).




Fig. 6. Type II photosensitized oxidation of HBI alkenes (RH = hydrogen donors)

2.2 Photodegradation processes in other phototrophic organisms

Visible light-dependent degradation processes have been also studied in senescent cells of two purple sulfur bacteria (*Thiohalocapsa halophila* and *Halochromatium salexigens*) isolated from microbial mats from Camargue (France) (Marchand and Rontani, 2003). These reactions act intensively on the phytyl side chain of bacteriochlorophyll-*a* and lead to the production of phytone (1) and phytyldiol (2) as in the case of chlorophylls (Fig. 1). Palmitoleic and *cis*-vaccenic acids also undergo strong photodegradation, affording mainly isomeric allylic oxo-, hydroxy- and hydroperoxyacids.

These processes were also investigated in aerobic anoxygenic phototrophic bacteria (AAPs) (Rontani et al., 2003a). These organisms constitute a relatively recently discovered bacterial group (Yurkov and Beatty, 1998) and seem to be widespread in the open ocean (Kolber et al., 2000). They perform photoheterotrophic metabolism, requiring organic carbon for growth, but they are capable to use photosynthesis as an auxiliary source of energy (Kolber et al., 2001). Though sensitive to photochemical processes in senescent purple sulfur bacteria (Marchand and Rontani, 2003), the isoprenoid phytyl side-chain of bacteriochlorophyll -*a* is not significantly photodegraded in senescent cells of AAPs (Rontani et al., 2003a). In contrast, significant amounts of allylic hydroxyacids arising from the photo-oxidation of the

major unsaturated fatty acid of these organisms (*cis*-vaccenic acid) could be detected after irradiation (Rontani et al., 2003a).

As in the case of phytoplankton and cyanobacteria, visible light-dependent degradation processes act significantly on the chlorophyll phytyl side-chain (Rontani et al., 1996b), unsaturated fatty acids and sterols (Rontani, Unpublished results) during terrestrial higher plant senescence affording similar photoproducts. 9-Hydroperoxy-18-hydroxyoctadec-10(*trans*)-enoic (**13**) and 10-hydroperoxy-18-hydroxyoctadec-8(*trans*)-enoic (**14**) acids deriving from type II photooxidation of 18-hydroxyoleic acid (**15**) (Fig. 7) were detected after visible light-induced senescence experiments carried out with *Petroselinum sativum* and subsequent cutin depolymerisation (Rontani et al., 2005a).




Fig. 7. Type II photosensitized oxidation of 18-hydroxyoleic acid in cutin polymers.

These results showed that in senescent plants, where the 1O_2 formation rate exceeds the quenching capacity of the photoprotective system, 1O_2 can migrate outside the chloroplasts and affect the unsaturated components of cutins. Significant amounts of 9,18-dihydroxyoctadec-10(*trans*)-enoic (**16**) and 10,18-dihydroxyoctadec-8(*trans*)-enoic (**17**) acids resulting from the reduction of these photoproducts of 18-hydroxyoleic acid were also detected in different natural samples (Rontani et al., 2005a). These results well support the significance of the photooxidation of the unsaturated components of higher plant cutins in the natural environment.

3. Free radical degradation (autoxidation) processes in phototrophic organisms

Autoxidation is the direct reaction of molecular oxygen with organic compounds under mild conditions. The autoxidation of organic compounds (in particular, lipids) involves free

radical reaction chains and thus includes an initiation, a propagation and a termination phase. Mechanisms of initiation for the free radical processes have been the subject of many studies. In senescent phytoplanktonic cells, initiation seems to result from the decomposition of hydroperoxides produced during photodegradation of cellular organic matter (Rontani et al., 2003b). Until now, autoxidative degradation in the marine environment has been largely ignored. Specific markers of these reactions have been highlighted by *in vitro* studies (Frankel, 1998; Rontani et al., 2003b; Rontani and Aubert, 2005). Using these markers, it was demonstrated *in situ* that autoxidation plays a very significant role in the degradation of particulate organic matter (Marchand et al., 2005; Rontani et al., 2006; Christodoulou et al., 2009; Rontani et al., 2011a).

Although the occurrence of autoxidation processes was clearly demonstrated *in situ*, it is not easy to induce these processes in laboratory cultures. Indeed, the mechanism of initiation of lipid radical oxidation, which has been debated for many years, seems to be the homolytic cleavage of photochemically produced hydroperoxides in phytodetritus (Rontani et al., 2003b). Redox-active metal ions are generally considered as the initiators of perhaps greatest importance for lipid oxidation in biological systems (Pokorny, 1987; Schaich, 1992). They may direct the cleavage of hydroperoxides either through alkoxy or peroxy radicals. In classical culture media (such as f/2) the metal chelator EDTA, which is present in high amounts, tightly binds free catalytic metal ions and thus renders them unavailable. EDTA thus acts in the culture media as an antioxidant and strongly limits radical oxidation processes.

Recently, autoxidative damages in cells of *E. huxleyi* strain CS-57 could be induced after incubation of this strain under an atmosphere of air + 0.5% CO₂ (Rontani et al., 2007a). The presence of additional CO₂ allowed: (i) to induce a stress that favoured oxidative damage and (ii) to decrease the pH of the culture medium releasing metal ions from EDTA complexes, which can act as catalysts of hydroperoxide homolysis.

It was also demonstrated recently that viral infection (Evans et al., 2006) and autocatalytic programmed cell death (Bidle and Falkowski, 2004) of phytoplanktonic cells could also lead to elevated production of reactive oxygen species (ROS) able to induce the degradation of cell components.

3.1 Chlorophyll phytyl side-chain

Autoxidation of the esterified chlorophyll phytyl chain involves either addition of peroxy radicals to the double bond or hydrogen abstraction at the allylic carbon 4 (Rontani and Aubert, 1994; Rontani and Aubert, 2005). Classical addition of peroxy radical to the double bond gives a tertiary radical (Fig. 8). This radical can then: (i) lead to *Z* and *E* epoxides (**18** and **19**) by fast intramolecular homolytic substitution (Fossey et al., 1995), or (ii) react with molecular oxygen affording (after hydrogen abstraction on another molecule of substrate) a diperoxide (**20**) (Fig. 8). Subsequent NaBH₄-reduction and alkaline hydrolysis of these compounds gives 3,7,11,15-tetramethylhexadecan-1,2,3-triol (**21**) (Fig. 8). In contrast, abstraction (by photochemically-produced peroxy radicals) of a hydrogen atom at the allylic carbon 4 of the phytyl chain and subsequent oxidation of the allylic radicals thus formed affords (after NaBH₄-reduction and alkaline hydrolysis) *Z* and *E* 3,7,11,15-tetramethylhexadec-3-en-1,2-diols (**3** and **4**) and *Z* and *E* 3,7,11,15-tetramethyl-hexadec-2-en-1,4-diols (**22** and **23**) (Fig. 8). Compounds **22** and **23** (which are well specific markers of free radical oxidation) could




Fig. 9. Free radical-mediated oxidation of oleic acid.

Free radical oxidative processes can be easily characterised based on the presence of *cis* allylic hydroperoxyacids, which cannot be produced photochemically (see Fig. 4) and are specific products of these degradation processes (Porter et al., 1995; Frankel, 1998).

Large amounts of oxidation products of oleic acid could be detected in cells of *E. huxleyi* grown under an atmosphere of air + 0.5% CO₂ for 10 days (Rontani et al., 2007a). The presence (after NaBH₄-reduction) of a high proportion of 11-hydroxyoctadec-*cis*-9-enoic (**27**) and 8-hydroxyoctadec-*cis*-9-enoic (**29**) acids (Fig. 10) showed that under these conditions the degradation of oleic acid mainly involved free radical oxidation processes.

3.3 Δ^5 -sterols

Free radical autoxidation of Δ^5 -sterols yields mainly 7 α - and 7 β -hydroperoxides and, to a lesser extent, 5 α / β ,6 α / β -epoxysterols and 3 β ,5 α ,6 β -trihydroxysterols (Smith, 1981; Morrissey and Kiely, 2006) (Fig. 11).




Fig. 10. Partial mass chromatogram of m/z 227, 329, 241 and 343 revealing the presence of oxidation products of oleic acid in the saponified fraction of *E. huxleyi* strain CS-57 grown under an atmosphere of air + 0.5% CO₂.

Owing to: their lack of specificity (possible formation by allylic rearrangement of photochemically-produced 5-hydroperoxides (see chapter 2.1.3), 7-hydroperoxides cannot be employed as tracers of autoxidation processes in phytodetritus. In contrast, it is generally considered that $5\alpha/\beta,6\alpha/\beta$ -epoxysterols arise mainly from peroxidation processes (Breuer and Björkhem, 1995; Giuffrida et al., 2004). Unfortunately, these compounds are not very stable and may be easily hydrolysed to the corresponding triol in seawater and during the treatment of the samples. $5\alpha/\beta,6\alpha/\beta$ -Epoxysterols and the corresponding $3\beta,5\alpha,6\beta$ -trihydroxysterols were thus finally selected as tracers of sterol autoxidation..

$5\alpha/\beta,6\alpha/\beta$ -Epoxysterols and $3\beta,5\alpha,6\beta$ -trihydroxysterols corresponding to sitosterol, stigmasterol and campesterol were previously detected in young and old cell cultures of *Chenopodium rubrum* (Meyer and Spiteller, 1997). The results showed that the increase of these oxidation products well correlated with the age of the culture.




Fig. 11. Free radical-mediated oxidation of Δ^5 sterols.

3.4 Vitamin E

Vitamin E is relatively abundant in most photosynthetic organisms, such as higher plants (Rise et al., 1988; Schultz, 1990), cyanobacteria (Dasilva and Jensen, 1971), microalgae (Brown et al., 1999) and macroalgae (Sanchez-Machado et al., 2002), where it plays an essential role in the removal of toxic forms of oxygen (singlet oxygen, superoxide anion, hydroxyl and peroxy radicals), by acting as sacrificial chemical scavenger (Halliwell, 1987); the process results in the irreversible oxidation of the tocopherol molecule. Vitamin E reacts rapidly with peroxy radicals, affording small amounts of phytone (1), 4,8,12,16-tetramethylheptadecan-4-olide, α -tocopherylquinone and epoxy- α -tocopherylquinones, and dimers and trimers as major oxidation products (Liebler, 1994; Frankel, 1998; Rontani et al., 2007b) (Fig. 12).

Isomeric trimers have been previously observed as products in numerous oxidations of vitamin E (e.g. Suarna et al., 1988; Krol et al., 2001). Such compounds cannot be easily detected since they are too heavy to be amenable by gas chromatography. However, methanolysis of the residues obtained resulted to the formation of high amounts of 5a-methoxytocopherol (**30**) arising from the methanolysis of the ketal group of trimers (Fig. 12) (Yamauchi et al., 1988). ESI-TOF MS analyses of oxidation products were also carried out in order to confirm the presence of high proportions of trimers (Nassiry et al., 2009).

Despite the intensive study of vitamin E oxidation since several decades, trimeric oxidation products could be detected in plants only very recently by Row et al. (2007). These authors detected these trimers in seeds of *Euryale ferox* containing extraordinarily high content of tocopherols. It is interesting to note that trimers were previously obtained as the major reaction products of vitamin E autoxidized under mild conditions in solution (1%) in methyl linoleate (Yamauchi et al., 1988). In plastoglobules, which are lipid monolayer subcompartments of the thylakoid membranes of chloroplasts (Maeda and Dellapenna, 2007), the concentration of tocopherols can reach 10% of the total fatty acids (Vidi et al., 2006). At such a concentration, the formation of a high proportion of trimers during photodynamic damages is thus very likely. In order to check this hypothesis, we searched for the presence of 5a-methoxytocopherol (**30**) after methanolysis of NaBH₄-reduced and non-reduced lipid extracts obtained from cells of *Emiliania huxleyi* strain TWP1 and *Chrysolida lamellosa* strain HAP17. The detection of significant amounts of this methanolysis product of trimers (Yamauchi et al., 1988) in these extracts (Nassiry et al., 2009) well supported the presence of such trimeric oxidation products of vitamin E in these algae.

3.5 Alkenones

The autoxidative reactivity of alkenones was studied in the laboratory in the presence of a radical initiator (di-*tert*-butyl nitroxide) and a radical enhancer (*tert*-butyl hydroperoxide) (Rontani et al., 2006). Alkenones appeared to be more sensitive towards oxidative free radical processes than analogues of other common marine lipids such as phytyl acetate, methyl oleate and cholesteryl acetate, and their oxidation rates increase in proportion with their number of double bonds. As the result of this increasing reactivity with degree of unsaturation, the $U_{37}^{K'}$ ratio increased significantly (up to 0.20) during the incubation.

Autoxidation of alkenones appears to mainly involve allylic hydrogen abstraction and subsequent oxidation of the allylic radical thus formed (Fig. 13). According to these processes, oxidation of each double bond of alkenones and subsequent NaBH₄ reduction affords four positional isomeric alkenediols. These compounds could be very useful indicators of autoxidation of alkenones but, unfortunately, they did not accumulate during the incubation. Indeed, due to the presence of additional reactive double bonds, hydroperoxyalkenones may undergo subsequent oxidation reactions affording, di-, tri- and tetrahydroperoxyalkenones according to the degree of unsaturation of the starting alkenone. In seawater, these different hydroperoxides may undergo two main degradative processes: (i) homolysis of the O-O bond leading to carbonyl (dehydration), alcoholic (reduction) and fragmentation (β -scission) products (Rontani et al., 2007c) and (ii) heterolysis of the O-O bond leading to the formation of two carbonyl fragments (Hock cleavage), this proton-catalysed cleavage being initiated by migration of groups to positive oxygen (Frimer, 1979). Dimeric and oligomeric compounds cross-linked through either peroxide or ether linkages (Frankel, 1998) may also be formed during autoxidation of alkenones.




Fig. 13. Characterization of oxidation products derived from the autoxidation of the ω 22 double bond of the $C_{37:3}$ alkenone (TMS = trimethylsilyl).

These results were corroborated by the further finding of significant amounts of alkenediols arising from NaBH_4 -reduction of the corresponding hydroperoxyalkenones in cultures of *E. huxleyi* strain CS-57 grown under an atmosphere of air + 0.5% CO_2 (Rontani et al., 2007a) and more recently after incubation of a culture of the strain *E. huxleyi* TWP1 under darkness (Rontani, Unpublished results) (Fig. 14) both exhibiting an anomalously high unsaturation ratio. It seems thus that autoxidation processes have the potential to affect alkenone distributions leading to a warm bias in estimates of palaeotemperatures derived from alkenone ratios in sediments.




Fig. 14. Partial mass fragmentograms of m/z 311 and 325 revealing the presence of silylated C_{37} and C_{38} alkenediols after NaBH_4 -reduction and silylation of the total lipid extract of *E. huxleyi* cells incubated under darkness (A) and standard autoxidation products of alkenones (B).

4. Conclusions

Due to the lack of adequate tracers, the role played by light-induced photochemical and free radical-mediated (autoxidative) processes during the degradation of lipid components of phototrophic organisms has been virtually ignored until now.

It was recently demonstrated that most of the unsaturated lipid components of these organisms (chlorophylls, carotenoids, unsaturated fatty acids, sterols, *n*-alkenes and HBI alkenes) could be photodegraded by visible and UV radiations during the senescence. This degradation mainly involves type II (i.e. involving $^1\text{O}_2$) photoprocesses. Singlet oxygen appeared to be sufficiently stable in this hydrophobic micro-environment to migrate outside the chloroplasts and affect the unsaturated components of cutins of higher plants.

Free radical-mediated oxidation (autoxidation) processes also intervene intensively during the senescence of phototrophic organisms. Induction of these processes seems to mainly result from the homolytic cleavage (catalyzed by some metal ions) of photochemically produced hydroperoxides. Unsaturated fatty acids, chlorophyll phytyl side-chain, vitamin E, sterols and alkenones appeared to be strongly affected by these degradative processes. In the case of alkenones, it is very important to note that autoxidative degradation processes may alter significantly their unsaturation ratio and thus constitute a potential source of biases during paleotemperature reconstruction.

5. Acknowledgements

Financial support over many years from the Centre National de la Recherche Scientifique (CNRS) and the Université de la Méditerranée is gratefully acknowledged.

6. References

- Allard, W. G., Belt, S.T., Massé, G., Naumann, R., Robert, J.-M. & Rowland, S.J. (2001). Tetra-unsaturated sesterterpenoids (Haslenes) from *Haslea ostrearia* and related species. *Phytochemistry* 56, 795-800.
- Belt, S.T., Allard, W.G., Massé, G., Robert, J.-M. & Rowland, S.J. (2000). Highly branched isoprenoids (HBIs): Identification of the most common and abundant sedimentary isomers. *Geochimica & Cosmochimica Acta* 64, 3839-3851.
- Belt, S.T., Massé, G., Rowland, S.J., Poulin, M., Michel, C. & Leblanc, B. (2007). A novel chemical fossil of palaeo sea ice : IP₂₅. *Organic Geochemistry* 38, 16-27.
- Belt, S.T., Vare, L.L., Massé, G., Manners, H., Price, J., MacLachlan, S., Andrews, J.T. & Schmidt, S. (2010). Striking similarities in temporal changes to seasonal sea ice conditions across the central Canadian Arctic Archipelago during the last 7,000 years. *Quaternary Science Reviews* 29, 3489-3504.
- Bidle, K.D. & Falkowski, P.G. (2004). Cell death in planktonic, photosynthetic microorganisms. *Nature Review* 2, 643-655.
- Boon, J.J., van der Meer, F.W., Schuyf, P.J., De Leeuw, J.W., Schenck, P.A. & Burlingame, A.L. (1978). Organic geochemical analyses of core samples from site 362 Walvis Ridge. DSDP Leg 40, *Initial Reports. DSDP Leg 38, 39, 40 and 41, Suppl.*, pp. 627-637.
- Breuer, O. & Blörkhem, I. (1995). Use of $^{18}\text{O}_2$ inhalation technique and mass isotopomer distribution analysis to study oxygenation of cholesterol in rat: evidence for in vitro formation of 7-oxo-, 7 β -hydroxy-, and 25-hydroxycholesterol. *Journal of Biological Chemistry* 270, 20278-20284.
- Brown, M.R., Mular, M., Miller, I., Farmer, C. & Trenerry, C. (1999). The vitamin content of microalgae used in aquaculture. *Journal of Applied Phycology* 11, 247-255.

- Christodoulou S., Marty J.-C., Miquel J.-C., Volkman J.K. & Rontani J.-F. (2009). Use of lipids and their degradation products as biomarkers for carbon cycling in the northwestern Mediterranean Sea. *Marine Chemistry* 113, 25-40.
- Christodoulou S., Joux F., Marty J.-C., Sempéré R. & Rontani J.-F. (2010). Comparative study of UV and visible light induced degradation of lipids in non-axenic senescent cells of *Emiliana huxleyi*. *Marine Chemistry* 119, 139-152.
- Conte, M.H., Volkman, J.K. & Eglinton, G. (1994). Lipid biomarkers of the Haptophyta. In: Green, J.C., Leadbeater, B.S.C. (Eds.), *The Haptophyte Algae. Systematics Association Special Volume No. 51*, Clarendon Press, Oxford, pp. 351-377.
- Cuny, P. & Rontani, J.-F. (1999). On the widespread occurrence of 3-methylidene-7,11,15-trimethylhexadecan-1,2-diol in the marine environment: a specific isoprenoid marker of chlorophyll photodegradation. *Marine Chemistry* 65, 155-165.
- Cuny, P., Romano, J.-C., Beker, B. & Rontani, J.-F. (1999). Comparison of the photodegradation rates of chlorophyll chlorin ring and phytol side chain in phytodetritus: is the phytyldiol versus phytol ratio (CPPI) a new biogeochemical index? *Journal of Experimental Marine Biology and Ecology* 237, 271-290.
- Cuny, P., Marty, J.-C., Chiaverini, J., Vescovali, I., Raphel, D. & Rontani, J.-F. (2002). One-year seasonal survey of the chlorophyll photodegradation process in the Northwestern Mediterranean Sea. *Deep Sea Research II* 49, 1987-2005.
- DaSilva, E.J. & Jensen, A. (1971). Content of α -tocopherol in some blue-green algae. *Biochimica et Biophysica Acta* 239, 345-347.
- De Leeuw, J.W., van der Meer, J.W., Rijpstra, W.I.C. & Schenck, P.A. (1980). On the occurrence and structural identification of long chain ketones and hydrocarbons in sediments. In: Douglas, A.G., Maxwell, J.R. (Eds.), *Advances in Organic Geochemistry 1979*. Pergamon Press, Oxford, pp. 211-7.
- Eltgroth, M.L., Watwood, R.L. & Wolfe, G.V. (2005). Production and cellular localization of neutral long-chain lipids in the Haptophyte algae *Isochrysis galbana* and *Emiliana huxleyi*. *Journal of Phycology* 41, 1000-1009.
- Engel, N., Jenny, T. A., Mooser, V. & Gossauer, A. (1991). Chlorophyll catabolism in *Chlorella protothecoides*. Isolation and structure elucidation of a red biline derivative. *FEBS Letters* 293, 131-133.
- Evans, C., Malin, G. & Mills, G.P. (2006). Viral infection of *Emiliana huxleyi* (Prymnesiophyceae) leads to elevated production of reactive oxygen species. *Journal of Phycology* 42, 1040-1047.
- Foote, C.S. (1976). Photosensitized oxidation and singlet oxygen: consequences in biological systems. In: Pryor, W.A. (Ed.), *Free Radicals in Biology*. Academic Press, New York, pp. 85-133.
- Fossey, J., Lefort, D. & Sorba, J. (1995). *Free radicals in organic chemistry*. Masson. Paris, pp 1-307.
- Frankel, E.N., Neff, W.E. & Bessler, T.R. (1979). Analysis of autoxidized fats by gas chromatography-mass spectrometry: V. photosensitized oxidation. *Lipids* 14, 961-967.
- Frankel, E.N. (1998). *Lipid Oxidation*. The Oily Press, Dundee.
- Frimer, A.A. (1983). Singlet oxygen in peroxide chemistry. In: Patai, S. (Ed.), *The Chemistry of Functional Groups, Peroxides*. John Wiley & sons Ltd., pp. 202-229.

- Gelin, F., Boogers, I., Noordeloos, A. A. M., Sinninghe Damsté, J. S., Riegman, R. & de Leeuw, J. W. (1997). Resistant biomacromolecules in marine microalgae of the classes Eustigmatophyceae and chlorophyceae: Geochemical implications. *Organic Geochemistry* 26, 659-675.
- Giuffrida, F., Destailats, F., Robert, F., Skibsted, L.H. & Dionisi F. (2004) Formation and hydrolysis of triacylglycerol and sterol epoxides : role of unsaturated triacylglycerol peroxy radicals. *Free Radical Biology & medicine* 37, 104-114.
- Grimalt, J.O., Rullkötter, J., Sicre, M.-A., Summons, R., Farrington, J., Harvey, H.R., Goñi, M. & Sawada, K. (2000). Modifications of the C₃₇ alkenone and alkenoate composition in the water column and sediment: possible implications for sea surface temperature estimates in paleoceanography. *Geochemistry Geophysics and Geosystems* 1, doi: 10.1029/2000G000053.
- Grossi, V., Raphel, D., Aubert, C. & Rontani, J.-F. (2000). The effect of growth temperature on the long-chain alkene composition in the marine coccolithophorid *Emiliania huxleyi*. *Phytochemistry* 54, 393-399.
- Grossi, V., Beker, B., Geenevasen, J.A.J., Schouten, S., Raphel, D., Fontaine, M.-F. & Sinninghe-Damsté, J.S. (2004). C₂₅ highly branched isoprenoid alkenes from the marine benthic diatom *Pleurosigma strigosum*. *Phytochemistry* 65, 3049-3055.
- Halliwell, B. (1987). Oxidative damage, lipid peroxidation and antioxidant protection in chloroplasts. *Chemistry and Physics of lipids* 44, 327-340.
- Harvey, H.R. (2000). Alteration processes of alkenones and related lipids in water columns and sediments. *Geochemistry Geophysics and Geosystems* 1, doi: 2000GC000054.
- He, Y.Y. & Häder, D.P. (2002) Reactive oxygen species and UV-B effect on cyanobacteria. *Photochemical and Photobiological Sciences* 1, 729-736.
- Heath, R.L. & Packer, L. (1968). Photoperoxidation in isolated chloroplasts. II. Role of electron transfer. *Archives of Biochemistry and Biophysics* 125, 850-857.
- Hurst, J.R., Wilson, S.L. & Schuster, G.B. (1985). The ene reaction of singlet oxygen: kinetic and product evidence in support of a peroxide intermediate. *Tetrahedron* 41, 2191-2197.
- Huysen, E.S. & Johnson, K.L. (1968). Concerning the nature of the polar effect in hydrogen atom abstractions from alcohols, ethers and esters. *Journal of Organic Chemistry* 33, 3972-3974.
- Iseo, S., Hyeon, S.B., Katsumura, S. & Sakan, T. (1972). Photooxygenation of carotenoids. II- The absolute configuration of loliolide and dihydroactinidiolide. *Tetrahedron Letters* 25, 2517-2520.
- Iturraspe, J., Engel, N. & Gossauer, A. (1994). Chlorophyll catabolism. Isolation and structure elucidation of chlorophyll b catabolites in *Chlorella protothecoides*. *Phytochemistry* 35, 1387-1390.
- Jaraula, C.M.B., Brassell, S.C, Morgan-Kiss, R, Doran, P.T. & Kenig, F. (2010) Origin and distribution of tri to pentaunsaturated alkenones in Lake Fryxell, East Antarctica. *Organic Geochemistry* 41, 386-397.
- Jeffrey, S.W. & Hallegraeff, G.M. (1987). Chlorophyllase distribution in ten classes of phytoplankton: a problem for chlorophyll analysis. *Marine Ecology Progress Series* 35, 293-304.
- Knox, J.P. & Dodge, A.D. (1985). Singlet oxygen and plants. *Phytochemistry* 24, 889-896.

- Kolber, Z.S., Van Dover, C.L., Niederman, R.A. & Falkowski, P.G. (2000) Bacterial photosynthesis in surface waters of the open ocean. *Nature* 407, 177-179.
- Kolber, Z.S., Plumley, F.G., Lang, A.S., Beatty, J.T., Blankenship, R.E., VanDover, C.L., Vetriani, C., Koblizek, M., Rathgeber, C. & Falkowski, P.G. (2001) Contribution of aerobic photoheterotrophic bacteria to the carbon cycle in the ocean. *Science* 292, 2492-2495.
- Korytowski, W., Bachowski, G.J. & Girotti, A.W. (1992). Photoperoxidation of cholesterol in homogeneous solution, isolated membranes, and cells: comparison of the 5 α - and 6 β -hydroperoxides as indicators of singlet oxygen intermediacy. *Photochemistry and Photobiology* 56, 1-8.
- Krol, E.S., Escalante, D.D.J. & Liebler, D.C. (2001). Mechanisms of dimer and trimer formation from ultraviolet-irradiated α -tocopherol. *Lipids* 36, 49-55.
- Kulig, M.J. & Smith, L.L. (1973). Sterol metabolism. XXV. Cholesterol oxidation by singlet molecular oxygen. *Journal of Organic Chemistry* 38, 3639-3642.
- Liebler, D.C. (1994). Tocopherone and epoxytocopherone products of vitamin E oxidation. *Methods in Enzymology* 234, 310-316.
- MacIntosh, T.J. (1978). The effect of cholesterol on the structure of phosphatidylcholine bilayers. *Biochimica et Biophysica Acta* 513, 43-58.
- Maeda, H. & DellaPenna, D. (2007). Tocopherol functions in photosynthetic organisms. *Current Opinion in Plant Biology* 10, 260-265.
- Marchand, D. & Rontani, J.-F. (2001). Characterization of photooxidation and autoxidation products of phytoplanktonic monounsaturated fatty acids in marine particulate matter and recent sediments. *Organic Geochemistry* 32, 287-304.
- Marchand, D. & Rontani, J.-F. (2003). Visible light-induced oxidation of lipid components of purple sulphur bacteria: A significant process in microbial mats, *Organic Geochemistry* 34, 61-79.
- Marchand, D., Marty, J.-C., Miquel, J.-C. & Rontani, J.-F. (2005). Lipids and their oxidation products as biomarkers for carbon cycling in the northwestern Mediterranean Sea: results from a sediment trap study. *Marine Chemistry* 95, 129-147.
- Marlowe, I.T., Green, J.C., Neal, A.C., Brassell, S.C., Eglinton, G. & Course, P.A. (1984). Long chain (n-C₃₇-C₃₉) alkenones in the Prymnesiophyceae. Distribution of alkenones and other lipids and their taxonomic significance. *British Phycology Journal* 19, 203-216.
- Massé, G., Rowland, S.J., Sicre, M.-A., Jacob, J., Jansen, E. & Belt, S.T. (2008). Abrupt climate changes for Iceland during the last millennium: Evidence from high resolution sea ice reconstructions. *Earth and Planetary Science Letters* 269, 565-569.
- Merzlyak, M.N. & Hendry, G.A.F. (1994). Free radical metabolism, pigment degradation and lipid peroxidation in leaves during senescence. *Proceedings of the Royal Society of Edinburgh* 102B, 459-471.
- Meyer, W. & Spitelner, G. (1997). Oxidized phytosterols increase by ageing in photoautotrophic cell cultures of *Chenopodium rubrum*. *Phytochemistry* 45, 297-302.
- Morrissey, P. A. & Kiely, M. (2006). Oxysterols: formation and biological function. In. *Advanced Dairy Chemistry 3rd edition*, Vol. 2 Lipids, Fox, P.F. & McSweeney, P.L.H., eds., Springer, New-York, pp. 641-674.

- Mouzdahir A., Grossi, V., Bakkas, S. & Rontani, J.-F. (2001). Photodegradation of long-chain alkenes in senescent cells of *Emiliana huxleyi* and *Nannochloropsis salina*. *Phytochemistry* 56, 677-684.
- Müller, P.J., Kirst, G., Ruhland, G., von Storch, I. & Rosell-Melé, A. (1998). Calibration of the alkenone paleotemperature index $U_{37}^{K'}$ based on core-tops from the eastern South Atlantic and global ocean (60°N-60°S). *Geochimica & Cosmochimica Acta* 62, 1757-1772.
- Nassiry, M., Aubert, C., Mouzdahir, A. & Rontani, J.-F. (2009) Generation of isoprenoid compounds and notably of prist-1-ene through photo- and autoxidative degradation of vitamin E. *Organic Geochemistry* 40, 38-50.
- Neff, W.E., Frankel, E.N. & Fujimoto, K. (1988). Autoxidative dimerization of methyl linolenate and its monohydroperoxides, hydroperoxy epidioxides and dihydroperoxides. *Journal of the American Oil Chemical Society* 65, 616-623.
- Nelson, J.R. (1993). Rates and possible mechanism of light-dependent degradation of pigments in detritus derived from phytoplankton. *Journal of Marine Research* 51, 155-179.
- Nickon, A. & Bagli, J.F. (1961). Reactivity and geochemistry in allylic systems. I. Stereochemistry of photosensitized oxygenation of monoolefins. *Journal of the American Chemical Society* 83, 1498-1508.
- Pokorny, J. (1987). Major factors affecting the autoxidation of lipids. In: Chan, H.W.-S. (Ed.), *Autoxidation of Unsaturated Lipids*. Academic Press, London, pp. 141-206.
- Porter, N.A., Caldwell, S.E. & Mills, K.A. (1995). Mechanisms of free radical oxidation of unsaturated lipids. *Lipids* 30, 277-290.
- Prahl, F.G. & Wakeham, S.G. (1987). Calibration of unsaturation patterns in long-chain ketone compositions for palaeotemperature assessment. *Nature* 330, 367-369.
- Prahl, F.G., Muehlhausen, L.A. & Zahnle, D.L. (1988). Further evaluation of long-chain alkenones as indicators of paleoceanographic conditions. *Geochimica & Cosmochimica Acta* 52, 2303-2310.
- Prahl F. G., Rontani J.-F., Volkman J. K., Sparrow M. A. & Royer I. M. (2006). Unusual C₃₅ and C₃₆ alkenones in a paleoceanographic benchmark strain of *Emiliana huxleyi*. *Geochimica & Cosmochimica Acta* 70, 2856-2867.
- Rechka, J.A. & Maxwell, J.R. (1988). Unusual long chain ketones of algal origin. *Tetrahedron Letters*. 29, 2599-600.
- Repeta, D.J. (1989). Carotenoid diagenesis in recent marine sediments. II- Degradation of fucoxanthin to loliolide. *Geochimica & Cosmochimica Acta* 53, 699-707.
- Rieley, G., Teece, M. A., Peakman, T. M., Raven, A. M., Greene, K. J., Clarke, T. P., Murray, M., Leftley, J. W., Campbell, C. N., Harris, R. P., Parkes, R. J. & Maxwell, J. R. (1998). Long-chain alkenes of the Haptophytes *Isochrysis galbana* and *Emiliana huxleyi*. *Lipids* 33, 617-625.
- Rise, M., Cojocar, M., Gottlieb, H.E. & Goldschmidt, E.E. (1988). Accumulation of α -tocopherol in senescing organs as related to chlorophyll degradation. *Plant Physiology* 89, 1028-1030.
- Rontani, J.-F. & Aubert, C. (1994) Effect of oxy-free radicals upon the phytyl chain during chlorophyll-a photodegradation. *Journal of Photochemistry and Photobiology*, A79, 167-172.

- Rontani, J.-F., Grossi, V., Faure, R. & Aubert, C. (1994). "Bound" 3-methylidene-7,11,15-trimethylhexadecan-1,2-diol: a new isoprenoid marker for the photodegradation of chlorophyll-a in seawater. *Organic Geochemistry* 21, 135-142.
- Rontani, J.-F., Beker, B., Raphel, D. & Baillet, G. (1995). Photodegradation of chlorophyll phytyl chain in dead phytoplanktonic cells. *Journal of Photochemistry and Photobiology* 85A, 137-142.
- Rontani, J.-F., Raphel, D. & Cuny, P. (1996a). Early diagenesis of the intact and photooxidized chlorophyll phytyl chain in a recent temperate sediment. *Organic Geochemistry* 24, 825-832.
- Rontani, J.-F., Cuny, P. & Grossi, V. (1996b). Photodegradation of chlorophyll phytyl chain in senescent leaves of higher plants. *Phytochemistry* 42, 347-351.
- Rontani, J.-F., Cuny, P. & Aubert, C. (1997a). Rates and mechanism of light-dependent degradation of sterols in senescing cells of phytoplankton. *Journal of Photochemistry and Photobiology* 111A, 139-144.
- Rontani, J.-F., Cuny, P., Grossi, V. & Beker, B. (1997b). Stability of long-chain alkenones in senescing cells of *Emiliana huxleyi*: effect of photochemical and aerobic microbial degradation on the alkenone unsaturation ratio ($U_{37}^{K'}$). *Organic Geochemistry* 26, 503-509.
- Rontani, J.-F., Cuny, P. & Grossi, V. (1998). Identification of a pool of lipid photoproducts in senescent phytoplanktonic cells. *Organic Geochemistry* 29, 1215-1225.
- Rontani, J.-F. & Marchand, D. (2000) Δ^5 -Stenol photoproducts of phytoplanktonic origin: a potential source of hydroperoxides in marine sediments? *Organic Geochemistry* 31, 169-180.
- Rontani J.-F., Koblizek M., Beker B., Bonin P. & Kolber Z. (2003a). On the origin of *cis*-vaccenic photodegradation products in the marine environment. *Lipids* 38, 1085-1092.
- Rontani, J.-F., Rabourdin, A., Marchand, D. & Aubert, C. (2003b). Photochemical oxidation and autoxidation of chlorophyll phytyl side chain in senescent phytoplanktonic cells: potential sources of several acyclic isoprenoid compounds in the marine environment. *Lipids* 38, 241-253.
- Rontani, J.-F. & Aubert, C. (2005). Characterization of isomeric allylic diols resulting from chlorophyll phytyl side-chain photo- and autoxidation by electron ionization gas chromatography/mass spectrometry. *Rapid Communications in Mass Spectrometry* 19, 637-646.
- Rontani, J.-F., Rabourdin, A., Pinot, F., Kandel, S. & Aubert, C. (2005) Visible light-induced oxidation of unsaturated components of cutins: a significant process during the senescence of higher plants. *Phytochemistry* 66, 313-321.
- Rontani, J.-F., Marty, J.-C., Miquel, J.-C. & Volkman, J.K. (2006). Free radical oxidation (autoxidation) of alkenones and other microalgal lipids in seawater. *Organic Geochemistry* 37, 354-368.
- Rontani, J.-F., Jameson, I., Christodoulou, S. & Volkman, J.K. (2007a). Free radical oxidation (autoxidation) of alkenones and other lipids in cells of *Emiliana huxleyi*. *Phytochemistry* 68, 913-924.
- Rontani, J.-F., Nassiry, M. & Mouzdahir, A. (2007b). Free radical oxidation (autoxidation) of α -tocopherol (vitamin E): A potential source of 4,8,12,16-tetramethylheptadecan-4-olide in the environment. *Organic Geochemistry* 38, 37-47.

- Rontani, J.-F., Harji, R. & Volkman, J.K. (2007c) Biomarkers derived from heterolytic and homolytic cleavage of allylic hydroperoxides resulting from alkenone autoxidation. *Marine Chemistry* 107, 230-243.
- Rontani, J.-F., Zabeti, N. & Wakeham, S.G. (2011a) Degradation of particulate organic matter in the equatorial Pacific Ocean: biotic or abiotic? *Limnology and Oceanography* 56, 333-349.
- Rontani, J.-F., Belt, S.T., Vaultier, F. & Brown, T.A. (2011b) Visible light-induced photo-oxidation of highly branched isoprenoid (HBI) alkenes: a significant dependence on the number and nature of the double bonds. *Organic Geochemistry* 42, 812-822
- Rosell-Melé, A., Carter, J.F., Parry, A.T. & Eglinton, G. (1995). Determination of the $U_{37}^{K'}$ index in geological samples. *Analytical Chemistry* 67, 1283-1289.
- Row, L.-C., Ho, J.-C. & Chen, C.-M. (2007). Cerebrosides and tocopherol trimers from the seeds of *Euryale ferox*. *Journal of Natural products* 70, 1214-1217.
- Rowland, S.J. & Robson, J.N. (1990). The widespread occurrence of highly branched isoprenoid acyclic C_{20} , C_{25} and C_{30} hydrocarbons in recent sediments and biota.- a review. *Marine Environmental Research* 30, 191-216.
- Sanchez-Machado, D.I., Lopez-Hernandez, J. & Paseiro-Losada, P. (2002). High-performance liquid chromatographic determination of α -tocopherol in macroalgae. *Journal of Chromatography* 976A, 277-284.
- Schaich, K.M. (1992). Metals and lipid oxidation. Contemporary issues. *Lipids* 27, 209-218.
- Schaich, K.M. (2005) Lipid oxidation: theoretical aspects. In: *Bailey's Industrial Oil and Fat Products*, Sixth edition, Shahidi, F. (Ed.), John Wiley & Sons, pp. 269-355.
- Schulte-Elte, K.H., Muller, B.L. & Pamingle, H. (1979). Photooxygenation of 3,3-dialkylsubstituted allyl alcohols. Occurrence of syn preference in the ene addition of 1O_2 at E/Z-isomeric alcohols. *Helvetica Chimica Acta* 62, 816-828.
- Schultz, G. (1990). Biosynthesis of α -tocopherol in chloroplasts of higher plants. *Fett Wissenschaft Technologie* 92, 86-91.
- Sinninghe-Damté, J.S., Muyzer, G., Abbas, B., Rampen, S.W., Massé, G., Allard, W.G., Belt, S.T., Robert, J.-M., Rowland, S.J., Moldovan, J.M., Barbanti, S.M., Fago, F.J., Denisevich, P., Dahl, J., Trindade, L.A.F. & Schouten, S. (2004). The rise of the Rhizosolenoid diatoms. *Science* 304, 584-587.
- Smith, L.L. (1981). *The Autoxidation of Cholesterol*. Plenum Press, New York.
- Suarna, C., Craig, D.C., Cross, K.J. & Southwell-Keely, P.T. (1988). Oxidations of vitamin E (α -tocopherol) and its model compound 2,2,5,7,8-pentamethyl-6-hydroxychroman. A new dimer. *Journal of Organic Chemistry* 53, 1281-1284.
- Suwa, K., Kimura, T. & Schaap, A.P. (1977). Reactivity of singlet molecular oxygen with cholesterol in a phospholipidic membrane matrix: a model for oxidative damage of membranes. *Biochemistry and Biophysical Research Communications* 75, 785-792.
- Vidi, P.-A., Kanwischer, M., Bagninsky, S., Austin, J.R., Csucs, G., Dörmann, P., Kessler, F. & Bréhélin, C. (2006). Tocopherol cyclase (VTE1) localization and vitamin E accumulation in chloroplast plastoglobule lipoprotein particles. *Journal of Biological Chemistry* 281, 11225-11234.
- Volkman, J.K., Eglinton, G., Corner, E.D.S. & Forsberg, T.E.V. (1980). Long chain alkenes and alkenones in the marine coccolithophorid *Emiliana huxleyi*. *Phytochemistry* 19, 2619-2622.

- Volkman, J. K., Barrett, S. M. & Dunstan, G. A. (1994). C₂₅ and C₃₀ highly branched isoprenoid alkenes in laboratory cultures of two marine diatoms. *Organic Geochemistry* 21, 407-414.
- Volkman, J.K., Barrett, S.M., Blackburn, S.I. & Sikes, E.L. (1995). Alkenones in *Gephyrocapsa oceanica* - implications for studies of paleoclimate. *Geochimica & Cosmochimica Acta* 59, 513–520.
- Wood, B.J.B. (1974). Fatty acids and saponifiable lipids. In: Steward, W.D. (Ed.), *Algal Physiology and Biochemistry*. University of California Press, Berkeley, pp. 236-265.
- Yamauchi, R., Kato, K. & Ueno, Y. (1988). Formation of trimers of α -tocopherol and its model compound, 2,2,5,7,8-pentamethylchroman-6-ol, in autoxidizing methyl linoleate. *Lipids* 23, 779-783.
- Yurkov, V.V. & Beatty, J.T. (1998) Aerobic anoxygenic phototrophic bacteria, *Microbiology and Molecular Biology Review* 62, 695-724.
- Zafiriou, O.C. (1977) Marine organic photochemistry previewed. *Marine Chemistry* 5, 497-522.
- Zafiriou, O.C., Jousset-Dubien, J., Zepp, R.G. & Zika, R.G. (1984) Photochemistry in natural waters. *Environmental Science and Technology* 18, 358A-370A.

IntechOpen



Senescence

Edited by Dr. Tetsuji Nagata

ISBN 978-953-51-0144-4

Hard cover, 850 pages

Publisher InTech

Published online 29, February, 2012

Published in print edition February, 2012

The book "Senescence" is aimed to describe all the phenomena related to aging and senescence of all forms of life on Earth, i.e. plants, animals and the human beings. The book contains 36 carefully reviewed chapters written by different authors, aiming to describe the aging and senescent changes of living creatures, i.e. plants and animals.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Jean-François Rontani (2012). Photo- and Free Radical-Mediated Oxidation of Lipid Components During the Senescence of Phototrophic Organisms, *Senescence*, Dr. Tetsuji Nagata (Ed.), ISBN: 978-953-51-0144-4, InTech, Available from: <http://www.intechopen.com/books/senescence/photo-and-free-radical-mediated-oxidation-of-lipid-components-during-the-senescence-of-phototrophic->

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen