We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Enzyme-Mediated Preparation of Flavonoid Esters and Their Applications

Jana Viskupicova^{1,2}, Miroslav Ondrejovic^{3,4} and Tibor Maliar³

¹*Institute of Experimental Pharmacology and Toxicology, Slovak Academy of Sciences*

²*Department of Biochemistry and Microbiology
Slovak University of Technology in Bratislava*

³*Department of Biotechnology, University of SS. Cyril and Methodius in Trnava*

⁴*Department of Biocentrum, Food Research Institute in Bratislava
Slovakia*

1. Introduction

Flavonoids comprise a group of plant polyphenols with a broad spectrum of biological activities. They have been shown to exert beneficial effects on human health and play an important role in prevention and/or treatment of several serious diseases, such as cancer, inflammation and cardiovascular disease (Middleton et al., 2000; Rice-Evans, 2001). Flavonoids are important beneficial components of food, pharmaceuticals, cosmetics and various commodity preparations due to their antimutagenic, hepatoprotective (Stefani et al., 1999), antiallergic (Berg & Daniel, 1988), antiviral (Middleton & Chithan, 1993) and antibacterial activity (Tarle & Dvorzak, 1990; Tereschuk et al., 1997; Singh & Nath, 1999; Quarenghi et al., 2000; Rauha et al., 2000). They are known to inhibit nucleic acid synthesis (Plaper et al., 2003; Cushnie & Lamb, 2006), cause disturbance in membranes (Stepanovic et al., 2003; Stapleton et al., 2004; Cushnie & Lamb, 2005) and affect energy metabolism (Haraguchi et al., 1998). But the most studied activity is their antioxidant action since they can readily eliminate reactive oxygen and nitrogen species or degradation products of lipid peroxidation and are thus effective inhibitors of oxidation (Ross & Kasum, 2002).

However, their commercial applications are limited due to low solubility in lipophilic environment and low availability for a living organism. Although aglycons, prenylated and methoxylated flavonoid derivatives may be implemented into such systems, they are rarely found in nature and are often unstable. In some plant species, the last step in the flavonoid biosynthesis is terminated by acylation which is known to increase solubility and stability of glycosylated flavonoids in lipophilic systems. Selectively acylated flavonoids with different aliphatic or aromatic acids may not only improve physicochemical properties of these molecules (Ishihara & Nakajima, 2003) but also introduce various beneficial properties to the maternal compound. These include penetration through the cell membrane (Suda et al., 2002; Kodelia et al., 1994) enhanced antioxidant activity (Viskupicova et al., 2010; Katsoura et al., 2006; Mellou et al., 2005), antimicrobial (Mellou et al., 2005), anti-proliferative (Mellou et al., 2006) and cytogenic (Kodelia et al., 1994) effect and improvement of thermostability and light-resistivity of certain flavonoids.

In nature, flavonoid acylation is catalyzed by various acyltransferases which are responsible for the transfer of aromatic or aliphatic acyl groups from a CoA-donor molecule to hydroxyl residues of flavonoid sugar moieties (Davies & Schwinn, 2006). Acylation is widespread especially among anthocyanins; more than 65% are reported to be acylated (Andersen & Jordheim, 2006). While the exact role of plant acylation is not yet fully understood, it is known that these modifications modulate the physiological activity of the resulting flavonoid ester by altering solubility, stability, reactivity and interaction with cellular targets (Ferrer et al., 2008). Acylation might be a prerequisite molecular tag for efficient vacuolar uptake of flavonoids (Kitamura, 2006; Nakayama et al., 2003). Some acylated flavonoids have been found to be involved in plant-insect interactions; they act as phytoalexins, oviposition stimulants, pollinator attractants (Iwashina, 2003), and insect antifeedants (Harborne & Williams, 1998). With respect to novel biological activities, acylation of flavonoids can result in changes in pigmentation (Bloor, 2001), insect antifeedant activity (Harborne & Williams, 1998) and antioxidant properties (Alluis & Dangles, 1999).

Over the past 15 years, there has been a substantial effort to take advantage of this naturally occurring phenomenon and to implement acylation methods into laboratories. However, the use of acyltransferases as modifying agents is rather inconvenient, as they require corresponding acylcoenzyme A, which must be either in stoichiometric amounts or regenerated *in situ*. Natural acyltransferases and cell extracts from *Ipomoea batatas* and *Perilla frutescens* containing acyltransferases were applied for selective flavonoid modification with aromatic acids (Tab.1) (Nakajima et al., 2000; Fujiwara et al., 1998).

Acyltransferase	Plant source	References
hydroxycinnamoyl-CoA:anthocyanin 3-O-glucosid-6"-O-acyltransferase	<i>Perilla frutescens</i>	Yonekura-Sakakibara et al., 2000
malonyl-CoA:anthocyanin 3-O-glucosid-6"-O-malonyltransferase	<i>Dahlia variabilis</i>	Wimmer et al., 1998
hydroxycinnamoyl-CoA:anthocyanin 5-O-glucosid-6"-O-acyltransferase	<i>Gentiana triflora</i>	Tanaka et al., 1996
hydroxycinnamoyl-CoA:anthocyanidin 3-rutinosid acyltransferase	<i>Petunia hybrida</i>	Brugliera & Koes, 2003
malonyl-CoA:anthocyanidin 5-O-glucosid-6"-O-malonyltransferase	<i>Salvia splendens</i>	Suzuki et al., 2001

Table 1. Acyltransferase catalysis of flavonoid acylation and their nature sources.

To solve this problem, the chemical approach was first investigated. It possessed a low degree of regioselectivity of esterification and drastic reaction conditions had to be applied (Patti et al., 2000). Later on, hydrolytic enzymes (lipases, esterases and proteases) have been recognized as useful agents due to their large availability, low cost, chemo-, regio- and enantioselectivity, mild condition processing and no need of cofactors (Collins & Kennedy, 1999; Nagasawa & Yamada, 1995).

Since the enzymatic preparation of flavonoid derivatives is a matter of several years, commercial applications have just been emerging. There are several patented inventions available to date, oriented on the flavonoid ester production and their use for the manufacture of pharmaceutical, dermopharmaceutical, cosmetic, nutritional or agri-foodstuff compositions

(Fukami et al., 2007; Moussou et al., 2004, 2007; Ghoul et al., 2006; Bok et al., 2001; Perrier et al., 2001; Otto et al., 2001; Nicolosi et al., 1999; Sakai et al., 1994).

This review presents available information on enzyme-mediated flavonoid acylation *in vitro*, emphasizing reaction parameters which influence performance and regioselectivity of the enzymatic reaction. In the second part, the paper focuses on biological effects of synthesized flavonoid esters as well as of those isolated from nature. Finally, the paper ends with application prospects of acylated flavonoids in the food, pharmaceutical and cosmetic industry.

2. Flavonoid esterification

Presently, the enzyme-catalyzed flavonoid esterification in organic media is a well-mastered technique for synthesis of selectively modified flavonoids. Results in this field suggest that a high degree of conversion to desired esters can be achieved when optimal reaction conditions are applied. The key factors, which influence regioselectivity and the performance of the enzymatic acylation of flavonoids, include type and concentration of enzyme, structure and concentration of the substrates (acyl donor, acyl acceptor and their ratio), nature of the reaction media, water content in the media, reaction temperature and nature of the reaction as reviewed in Chebil et al., 2006, 2007.

2.1 Enzymes

To date, the use of proteases, esterases, acyltransferases and lipases has been investigated in order to find the most potent biocatalyst for selective flavonoid acylation. These enzymes are often in the immobilized form which improves enzyme stability, facilitates product isolation, and enables enzyme reuse (Adamczak & Krishna, 2004).

2.1.1 Proteases

Proteases represent a class of enzymes which occupy a pivotal position with respect to their physiological roles as well as their commercial applications. They represent the first group of hydrolytic enzymes used for flavonoid modification. They perform both hydrolytic and synthetic functions. Since they are physiologically necessary for living organisms, proteases occur ubiquitously in diverse sources, such as plants, animals, and microorganisms. They are also classified as serine proteases, aspartic proteases, cysteine proteases, threonine proteases and metalloproteases, depending on the nature of the functional group at the active site.

Proteases have a large variety of applications, mainly in the detergent and food industries. In view of the recent trend of developing environmentally friendly technologies, proteases are envisaged to have extensive applications in leather treatment and in several bioremediation processes. Proteases are also extensively used in the pharmaceutical industry (Rao et al., 1998). Protease subtilisin was the first enzyme used for flavonoid ester synthesis conducted by Danieli et al. (1989, 1990). Later on, subtilisin was used for selective rutin acylation in organic solvents (Xiao et al., 2005; Kodelia et al., 1994). However, it has been reported that reactions catalyzed by subtilisin led to low conversion yields and a low degree of regioselectivity was observed (Danieli et al., 1990). These authors reported that the structure of the sugar moiety affected the regioselectivity. For flavonoid acylation, especially

serine proteases (subtilisin) have been used in ester synthesis (Danieli et al., 1989, 1990; Kodelia et al., 1994).

2.1.2 Esterases

Esterases (carboxyl esterases, EC 3.1.1.1) represent a diverse group of hydrolases catalyzing the cleavage and formation of ester bonds with wide distribution in animals, plants and microorganisms. A classification scheme for esterases is based on the specificity of the enzymes for the acid moiety of the substrate, such as the carboxyl esterases, aryl esterases, acetyl esterases, cholin esterases, cholesterol esterases, etc. (Jeager et al., 1999). Esterases show high regio- and stereospecificity, which makes them attractive biocatalysts for the production of optically pure compounds in fine-chemicals synthesis (reviewed in Bornscheuer, 2002).

They have the same reaction mechanism as lipases, but differ from them by their substrate specificity, since they prefer short-chain fatty acids, whereas lipases usually prefer long-chain fatty acids. Another difference lies in the interfacial activation (Hidalgo & Bornscheuer, 2006). In contrast to lipases, only a few esterases have commercial applications in organic synthesis because lipases are generally more enantioselective and resistant to organic solvents. The most widely used esterase is the preparation isolated from pig liver (Hidalgo & Bornscheuer, 2006). The practical applications of esterases in enzymatic transformation of flavonoids are not very attractive as it enables the implementation only of the molecule of a short aliphatic chain length, such as acetate, propionate and butyrate (Sakai et al., 1994).

2.1.3 Lipases

Today lipases stand amongst the most important biocatalysts in industry. Among them, microbial lipases find the biggest application use. They can be classified according to sequence alignment into three major groups: mammalian lipases (e.g. porcine pancreatic lipase), fungal lipases (*Candida rugosa* and *Rhizomucor* family) and bacterial lipases (*Staphylococcus* and *Pseudomonas* family) (Hidalgo & Bornscheuer, 2006). More than 50% of the reported lipases are produced by yeast in the forms of various isozymes (Vakhlu & Kour, 2006).

Lipases (triacylglycerol acylhydrolases, EC 3.1.1.3) belong to the class of serine hydrolases. They catalyze a wide range of reactions, including hydrolysis, inter-esterification, alcoholysis, acidolysis, esterification and aminolysis (Vakhlu & Kour, 2006). Under natural conditions, they catalyze the hydrolysis of ester bonds at the hydrophilic-hydrophobic interface. At this interface, lipases exhibit a phenomenon termed interfacial activation, which causes a remarkable increase in activity upon contact with a hydrophobic surface. The catalytic process involves a series of differentiated stages: contact with the interface, conformational change, penetration in the interface, and finally the catalysis itself (Hidalgo & Bornscheuer, 2006). Under certain experimental conditions, such as in the absence of water, they are capable of reversing the reaction. The reverse reaction leads to esterification and formation of glycerides from fatty acids and glycerol (Saxena et al., 1999). This synthetic activity of lipases is being successfully utilized also in flavonoid ester production.

Candida antarctica lipase B (CALB) is one of the most widely used biocatalysts in organic synthesis on both the laboratory and the commercial scale (Anderson et al., 1998; Uppenberg et al., 1995) due to its ability to accept a wide range of substrates, its non-aqueous medium tolerance and thermal deactivation resistance (Degn et al., 1999; Anderson et al., 1998; Cordova et al., 1998; Drouin et al., 1997). CALB belongs to the α/β hydrolase-fold superfamily with a conserved catalytic triad consisting of Ser105-His224-Asp187 (Uppenberg et al., 1995). It comprises 317 amino acid residues. The active site contains an oxyanion hole which stabilizes the transition state and the oxyanion in the reaction intermediate (Haefner et al., 1998). Reaction mechanism of CALB follows the bi-bi ping-pong mechanism, illustrated in Fig.1 (Kwon et al., 2007). The substrate molecule reacts with serine of the active site forming a tetrahedral intermediate which is stabilized by catalytic residues of His and Asp. In the next step alcohol is released and the acyl-enzyme complex is created. A nucleophilic attack (water in hydrolysis, alcohol in transesterification) causes another tetrahedral intermediate formation. In the last step, the intermediate is split into product and enzyme and is recovered for the next catalytic cycle (Patel, 2006).

Fig. 1. Reaction mechanism catalyzed by *Candida antarctica* lipase (Kwon et al., 2007).

The active site of CALB consists of a substrate-nonspecific acyl-binding site and a substrate specific alcohol-binding site (Cygler & Schrag, 1997; Uppenberg et al., 1995). It is selective for secondary alcohols (Uppenberg et al., 1995), as reflected by the geometry of the alcohol-binding site (Lutz, 2004). In contrast to most lipases, CALB has no lid covering the entrance to the active site and shows no interfacial activation (Martinelle et al., 1995). CALB is being frequently used in acylation of various natural compounds such as saccharides, steroids and natural glycosides, including flavonoids (Riva, 2002; Davis & Boyer, 2001). The proper enzyme selection plays multiple roles in flavonoid acylation. The biocatalyst significantly influences the regioselectivity of the reaction. Information is available mainly on the use of lipases for flavonoid ester synthesis; especially the use of lipase B from *Candida antarctica*,

which is preferred due to its acceptance of a wide range of substrates, good catalytic activity and a high degree of regioselectivity (Viskupicova et al., 2010; Katsoura et al., 2006, 2007; Ghoul et al., 2006; Mellou et al., 2005, 2006; Stevenson et al., 2006; Ardhaoui et al., 2004a, 2004b, 2004c; Passicos et al., 2004; Moussou et al., 2004; Gayot et al., 2003; Ishihara & Nakajima, 2003; Ishihara et al., 2002; Kontogianni et al., 2001, 2003; Nakajima et al., 1999, 2003; Gao et al., 2001; Otto et al., 2001; Danieli et al., 1997).

As for flavonoid aglycons, only two enzymes have been reported to be capable of acylating this skeleton – lipase from *Pseudomonas cepacia* and carboxyl esterase. Lambusta et al. (1993) investigated the use of *P. cepacia* lipase for catechin modification. They discovered that the acylation took place on the C5 and C7 hydroxyls. Sakai et al. (1994) observed that carboxyl esterase showed regioselectivity towards C3-OH of catechin. Sakai et al. (1994) explored the use of carboxyl esterase from *Streptomyces rochei* and *Aspergillus niger* for the 3-O-acylated catechin production.

2.2 Reaction conditions

The performance and regioselectivity of the enzyme-catalyzed flavonoid transformation is affected by several factors, including the type of enzyme, the nature of medium, reaction conditions, water content in the media, structure and concentration of substrates and their molar ratio. By varying these factors, significant changes in ester production and regioselectivity can be achieved.

2.2.1 Reaction media

Reaction media play an important role in enzymatic transformations. Methodologies for enzymatic flavonoid acylation have focused on searching a reaction medium which allows appropriate solubility of polar acyl acceptor (flavonoid glycoside) and nonpolar acyl donor as well as the highest possible enzymatic activity. Moreover, the medium has often been required to be nontoxic and harmless to biocatalyst. In order to meet the above-mentioned requirements, several scientific teams have dealt with proper medium selection (Viskupicova et al., 2006; Mellou et al., 2005; Kontogianni et al., 2001, 2003; Gao et al., 2001; Nakajima et al., 1999; Danieli et al., 1997).

Non-aqueous biocatalysis has several advantages over conventional aqueous catalysis: the suppression of hydrolytic activity of the biocatalyst which is carried out in water (Fossati & Riva, 2006), the enhanced solubility of hydrophobic substrates, the improvement of enzyme enantioselectivity, the exclusion of unwanted side reactions, the easy removal of some products, the enhanced enzyme thermostability and the elimination of microbial contamination (Rubin-Pitel & Zhao, 2006; Torres & Castro, 2004). Laane (1987) pointed out that log P, as a solvent parameter, correlated best with enzyme activity. Zaks & Klivanov (1988) reported that the activity of lipases was higher in hydrophobic solvents than in hydrophilic ones. Narayan & Klivanov (1993) claimed that it was hydrophobicity and not polarity or water miscibility which was important, whereas the log P parameter could be called a measure of solvent hydrophobicity. Trodler & Pleiss (2008), using multiple molecular dynamics simulations, showed that the structure of CALB possessed a high stability in solvents. In contrast to structure, flexibility is solvent-dependent; a lower dielectric constant led to decreased protein flexibility. This reduced flexibility of CALB in

non-polar solvents is not only a consequence of the interaction between organic solvent molecules and the protein, but it is also due to the interaction with the enzyme-bound water and its exchange on the surface (Trodler & Pleiss, 2008). In organic solvents, the surface area has been suggested to be reduced, leading to improved packing and increased stability of the enzyme (Toba & Merz, 1997).

Polar aprotic solvents such as dimethyl sulfoxid (DMSO), dimethylformamide (DMF), tetrahydrofuran (THF) and pyridine were first investigated (Nakajima et al., 1999; Danieli et al., 1997). However, it was observed that enzyme activity was readily deactivated in these solvents. To date enzymatic acylation of flavonoids has been successfully carried out in various organic solvents (Tab.2), while the most frequently used are 2-methylbutan-2-ol and acetone because of their low toxicity, their polarity allowing proper solubilization of substrates and high conversion yields.

Solvent	Reference
2-Methylbutan-2-ol	Ghoul et al., 2006; Ardhaoui et al., 2004a, 2004b, 2004c; Passicos et al., 2004; Gayot et al., 2003
Acetone	Ghoul et al., 2006; Mellou et al., 2005, 2006; Kontogianni et al., 2001, 2003; Ishihara et al., 2002, Ishihara & Nakajima, 2003; Nakajima et al., 1999, 2003; Danieli et al., 1997
Acetonitrile	Ghoul et al., 2006; Ishihara & Nakajima, 2003; Nakajima et al., 1997, 1999
2-Methylpropan-2-ol	Ghoul et al., 2006; Stevenson et al., 2006; Mellou et al., 2005; Moussou et al., 2004; Kontogianni et al., 2001, 2003; Otto et al., 2001
Dioxane	Ghoul et al., 2006; Danieli et al., 1997
Pyridine	Danieli et al., 1990, 1997
THF, DMSO, DMF	Kontogianni et al., 2001, 2003; Danieli et al., 1997
Binaric mixtures of solvents	Ghoul et al., 2006; Gao et al., 2001; Nakajima et al., 1999; Danieli et al., 1997

Table 2. Organic solvents used in flavonoid acylation.

The effect of the solvent on conversion yield depends on the nature of both the acyl donor and the flavonoid (Chebil et al., 2006). Although much has been done in this area, it is quite difficult to deduce any general conclusion on solvent choice because the available data are controversial and sometimes even contrary.

Recently, ionic liquids have received growing attention as an alternative to organic solvents used for the enzymatic transformation of various compounds (Katsoura et al., 2006; Kragl et al., 2006; Jain et al., 2005; Lozano et al., 2004; Reetz et al., 2003; Van Rantwick et al., 2003). The potential of these “green solvents” lies in their unique physicochemical properties, such as non-volatility, nonflammability, thermal stability and good solubility for many polar and less polar organic compounds (Jain et al., 2005; Wilkes, 2004; Itoh et al., 2003; Van Rantwick et al., 2003). Probably the most promising advantage of the use of ionic liquids is their potential application in food, pharmaceutical and cosmetic preparations due to their reduced toxicity (Jarstoff et al., 2003). Due to the many above-mentioned advantages of ionic liquids for enzyme-mediated transformations, several flavonoid esters have been recently

prepared in such media (Katsoura et al., 2006, 2007; Kragl et al., 2006). The biocatalytic process showed significantly higher reaction rates, regioselectivity and yield conversions compared to those achieved in organic solvents. Thus ionic liquid use seems to be a challenging approach to conventional solvent catalysis.

The solvent-free approach for elimination of the co-solvent of the reaction has been recently introduced as an alternative for conventional solvents (Enaud et al., 2004; Kontogianni et al., 2001, 2003). It is based on the use of one reactant in the role of the solvent. The authors reported rapid reaction rates; however, the conversion yields were slightly decreased. In spite of the attractiveness, the use of solvent-free systems is characterized by a serious drawback due to the necessity to eliminate the excess of the acyl donor for the recovery of the synthesized products (Chebil et al., 2006).

2.2.2 Water content

Water content in reaction media is a crucial parameter in lipase-catalyzed synthesis as it alters the thermodynamic equilibrium of the reaction towards hydrolysis or synthesis. Moreover, it is involved in noncovalent interactions which keep the right conformation of an enzyme catalytic site (Foresti et al., 2007). The amount of water required for the catalytic process depends on the enzyme, its form (native or immobilized), the enzyme support, and on the solvent nature (Arroyo et al., 1999; Zaks & Klivanov, 1988). The influence of water content in the reaction system on enzyme activity is variable with various enzymes (lipase from *Rhizomucor miehei*, *Rhizomucor niveus*, *Humicola lanuginosa*, *Candida rugosa*, *Pseudomonas cepacia*).

In general, the water amount which is considered to be optimal for esterifications in organic solvents is 0.2 – 3% (Rocha et al., 1999; Yadav & Piyush, 2003; Iso et al., 2001). The enzymatic esterification of flavonoids in non-aqueous media is greatly influenced by the water content of the reaction system (Ardhaoui et al., 2004b; Gayot et al., 2003; Kontogianni et al., 2003). Ardhaoui et al. (2004b) observed the best enzyme activity when water content was maintained at 200 ppm. Gayot et al. (2003) found that the optimal value of water in an organic reaction medium equaled 0.05% (v/v). Kontogianni et al. (2003) reported that highest flavonoid conversion was reached when initial water activity was 0.11 or less.

2.2.3 Temperature

Temperature represents a significant physical factor in enzyme-catalyzed reactions. It affects viscosity of the reaction medium, enzyme stability, and substrate and product solubility.

Since lipase from *C. antarctica* belongs to thermostable enzymes, improved catalytic activity was observed at higher temperatures (Arroyo et al., 1999). To date, flavonoid transformation has been carried out in the temperature range 30 – 100°C (Ghoul et al., 2006; Katsoura et al., 2006; Stevenson et al., 2006; Mellou et al., 2005; Ardhaoui et al., 2004a, 2004b, 2004c; Moussou et al., 2004; Passicos et al., 2004; Enaud et al., 2004; Gayot et al., 2003; Kontogianni et al., 2003; Ishihara et al., 2002; Gao et al., 2001; Otto et al., 2001; Nakajima et al., 1999; Danieli et al., 1990). The choice of temperature depends on the enzyme and solvent used. The majority of authors performed flavonoid acylation at 60°C due to the best enzyme activity, good solubility of substrates and highest yields of resulting esters reached (Viskupicova et al., 2006, 2010; Ghoul et al., 2006; Katsoura et al., 2006; Stevenson et al., 2006;

Ardhaoui et al., 2004a, 2004b, 2004c; Moussou et al., 2004; Passicos et al., 2004; Enaud et al., 2004; Gayot et al., 2003; Otto et al., 2001). Our results on the effect of temperature on naringin conversion are presented in Fig.2 and are in accordance with other authors (Viskupicova et al., 2006).

Fig. 2. Effect of temperature on naringin conversion to naringinpalmitate in 2-methylbutan-2-ol catalyzed by *C. antarctica* lipase after 24 h (Viskupicova et al., 2006).

2.3 Acyl donors and acceptors

2.3.1 Acyl donor

Since lipase-catalyzed acylation takes place through the formation of an acyl-enzyme intermediate, the nature of the acyl donor has a notable effect on reactivity. The ideal acyl donor should be inexpensive, fast acylating, and completely non-reactive in the absence of the enzyme (Ballesteros et al., 2006). Many acylating agents have been tested in flavonoid esterification, such as aromatic or aliphatic organic acids, substituted or not (Tab.3). Special attention was attributed to fatty acid ester production (Katsoura et al., 2006; Mellou et al., 2005, 2006; Ardhaoui et al., 2004a, 2004b, 2004c; Enaud et al., 2004; Gayot et al., 2003; Kontogianni et al., 2003). This approach enables to improve flavonoid solubility and stability in lipophilic systems. The proper acyl donor selection may significantly influence not only the physicochemical but also biological properties of the resulting esters.

A simple way to increase the reaction rate and conversion yield in acylation is to use an excess of acyl donor (Patti et al., 2000). Many authors have tried to determine the optimal molar ratio of flavonoid/acyl donor in order to achieve the highest possible yields. The molar ratios 1:1 to 1:15 (acyl acceptor/acyl donor) have been investigated, whereas the majority agreed on the ratio 1:5 to be the most suitable for the best reaction performance (Mellou et al., 2006; Gayot et al., 2003; Ishihara & Nakajima, 2003; Ishihara et al., 2002; Kontogianni et al., 2001). A better solution is offered by the use of special acyl donors which ensure a more or less irreversible reaction. This can be achieved by the introduction of electron-withdrawing substituents (esters), resulting in higher conversion yields and reaction rates. The use of vinyl esters allows a several times faster reaction progress than do other activated esters (Ballesteros et al., 2006). Enzymatic synthesis of flavonoid esters can be realized by two basic approaches, i.e. esterification and transesterification (Fig.3).

Aliphatic acids	Aromatic acids
Acetic*	Caffeic*
Malic*	p-Coumaric*
Malonic*	Ferulic*
Succinic*	Gallic*
Tartaric*	p-Hydroxybenzoic*
Butyric	Sinapic*
Crotonic	Benzoic
n-Butanoic	Cinnamic
Isobutyric	Isoferulic
Isovaleric	Methylsinapic
Lactic	
3-methylbutyric	
Quinic	
Vinylpropionic	
Tiglic	

*acyl donors found in anthocyanins

Table 3. Acyl donors found in flavonols, flavones (Williams, 2006) and anthocyanins (Andersen & Jordheim, 2006).

Fig. 3. Mechanism of isoquercitrin esterification and transesterification (Chebil et al., 2006).

Pleiss et al. (1998) studied the acyl binding site of CALB and found the enzyme to be selective for short and medium fatty acid chain length. This fact may be attributed to the structure of the lipase acyl binding pocket, which is an elliptical, narrow cleft of 9.5 × 4.5 Å. With increasing carbon number of a fatty acid or molecule size, the steric hindrance is involved resulting in low efficiency of the enzymatic reaction (Riva et al., 1988; Wang et al., 1988; Carrea et al., 1989). This fact was experimentally confirmed by Katsoura et al. (2006) and by Viskupicova & Ondrejovic (2007) whose results showed higher performance of the naringin and rutin esterification when fatty acids up to C10 were introduced. On the other hand, Ardhaoui et al. (2004b) and Kontogianni et al. (2003) reported that the fatty acid chain length had no significant effect on conversion yield when fatty acids of a medium and high chain length were used.

Thus, the effect of fatty acid chain length on flavonoid acylation still remains a matter of discussion. Our team conducted a series of experiments with both saturated and unsaturated fatty acids and found a correlation between log P of the acids tested and conversion yields (Viskupicova et al., 2010). It would be interesting to take this parameter into consideration when assessing the influence of an acyl donor on the reaction progress.

Only little progress has been achieved in flavonoid esterification with aromatic acids (Stevenson et al., 2006; Enaud et al., 2004; Gao et al., 2001; Nakajima et al., 2000). It has been observed that the performance of the process depends mainly on the nature of the substitutions, the position of the hydroxyls and the length of the spacers.

2.3.2 Acyl acceptor

The structure of acyl acceptor (flavonoid), especially stereochemistry of glycosidic bonds, plays an important role in flavonoid acylation. The structural differences, such as the number and position of hydroxyl groups, the nature of saccharidic moiety, as well as the position of glycosidic bonds, influence the flavonoid solubility, and thus affect the overall conversion yield.

Available studies are concerned mainly with acylation on flavonoid glycosides. Among polyphenolic compounds, naringin and rutin are the most widely used substrates. For the naringin molecule, which possesses a primary hydroxyl group on glucose, the acylation takes place on the 6''-OH (Katsoura et al., 2006; Konntogianni et al., 2001, 2003; Ishihara et al., 2002; Gao et al., 2001; Otto et al., 2001; Danieli et al., 1990) since the primary hydroxyl is favored by CALB (Fig.4). However, in rutin, which has no primary hydroxyl available, either the 3''-OH of glucose (Ishihara et al., 2002; Danieli & Riva, 1994) or the 4'''-OH of rhamnose (Fig.4) (Viskupicova et al., 2010; Mellou et al., 2006; Ardhaoui et al., 2004a, 2004b, 2004c) can be acylated. Danieli et al. (1997) observed the rutin-3'',4'''-O-diester formation. When subtilisin was used as biocatalyst, naringin-3''-O-ester and rutin-3''-O-ester were synthesized (Danieli et al., 1990).

Fig. 4. Acylation sites of naringin (left) and rutin (right) molecule.

The concentration of the flavonoid also affects the performance of the acylation reaction. The conversion yield and the initial rate rise with increasing flavonoid concentration. However, the amount of flavonoid is limited by its solubility in a reaction medium (Chebil et al., 2006, 2007).

3. Influence of flavonoid derivatization on biological activities

3.1 Esters with aromatic acids

Aromatic acids, along with flavonoids, belong to the group of phenols of secondary metabolism of living organisms. The described secondary metabolites represent a store of biologically active compounds, displaying various biological activities. We can therefore assume that physicochemical and biological properties of the initial flavonoids may be improved by acylation of flavonoids with aromatic acids. However, by this reaction a new compound can also gain novel activities provided by the aromatic acids.

Flavonoid acylation with aromatic acids was reported to improve physiological activities, such as UV-absorbing capacity, radical scavenging ability (Delazar et al., 2005; Ishihara & Nakajima, 2003; Harborne & Williams, 2000; Alluis & Dangles 1999; Jungblut et al., 1995) pigment stabilization (especially anthocyanins) (Ishihara & Nakajima 2003), and interaction with cellular targets (Ferrer et al., 2008).

Flavonoid esters acylated with *p*-coumaric acid were found to increase antioxidant (Pajero et al., 2005) and anti-inflammatory activities (Harborne & Williams, 2000), as well as antiproliferative and cytotoxic effects on various cancer cell lines (Mitrokotsa et al., 1993). Moreover, *p*-coumaroyl esters of quercetin and kaempferol were reported to have positive effects on cerebrovascular disorders (Calis et al., 1995). Similarly, flavonoid esters esterified with cinnamic acid were shown to exhibit antiproliferative activity against several human cancer cell lines (Duarte-Almeida et al., 2007). Flavonoid acylation with caffeic acid contributes to the enhancement of antioxidant properties (Pajero et al., 2005). Flavonolignans acylated with truxinic acid were shown to possess hepatoprotective as well as anticancer activity (Sharma et al., 2003).

3.2 Esters with aliphatic acids

Biological activities of aliphatic acids are not of a big importance in comparison with aromatic acids. These compounds are mainly accepted as energy storage and components of several compartments of cells, such as membranes, enzymes, surfactants, etc. In the literature, more studies can be found describing changes in biological activities of flavonoids after their acylation with aliphatic acids.

The aliphatic acylation of anthocyanins with malonic acid is important for enhancing the pigment solubility in water, protecting glycosides from enzymatic degradation and stabilizing anthocyanin structures (Nakayama et al., 2003). Several *in vitro* observations suggest that acylation with malonic acid or sinapic acid is crucial for efficient flavonoid accumulation in plants.

Fatty acid esters of catechins were reported to display antitumor, antibacterial and 5- α reductase inhibiting activity (Fukami et al., 2007) as well as antioxidant properties (Sakai et al., 1994). Lee et al. (2003) reported anti-atherogenic activity of two naringenin derivatives, 7-*O*-oleic ester and 7-*O*-cetyl ether.

Acylation of the flavonoid molecule with polyunsaturated fatty acids introduces potential antitumor and antiangiogenic properties (Mellou et al., 2006). Anticarcinogenic effects were observed also in silybin esters acylated with butyric and lauric acid (Xanthakis et al., 2010). Recently, we found that acylation of rutin with unsaturated fatty acids, such as oleic, α -linoleic and linolenic, increased the antioxidant potential of the initial compound (Viskupicova et al., 2010). This observation is in accordance with the results of Mellou et al. (2006) and Katsoura et al. (2006).

In the field of fatty acid ester synthesis, information on the photoprotective effectiveness of new quercetin derivatives acylated with acetic, propionic and palmitic acids, has been reported. The authors found that esterification with a short side-chain (such as acetate or propionate) may improve migration through the aqueous environment and interaction with or penetration into phospholipid membranes (Sajja et al., 2003).

Recent experimental findings indicate that acylation of flavonoid may increase enzyme inhibitory activity. Lin et al. (2010) observed increased 5 α -reductase inhibition after acylation of (-)-epigallocatechins. Salem et al. (2011) showed that the acylation of isorhamnetin-3-*O*-glucoside with different aliphatic acids enhanced its capacity to inhibit xanthine oxidase. Our recent investigations showed that lipophilic rutin and naringin esters were strong inhibitors of transport enzymes such as sarcoplasmic reticulum Ca²⁺-ATPase and plasma membrane Ca²⁺-ATPase (Augustyniak et al., 2010; Viskupicova et al., 2009), and thus might be useful in calcium regulation. We presume that there might be a general mechanism involved in the enhanced inhibitory activity of the acylated flavonoids on structurally diverse classes of enzymes which seems to be donated by the medium to long fatty acid chains.

4. Application perspectives

The following section provides a summary of patented inventions available in the commercial sphere. These include practical applications in food, pharmaceuticals and cosmetics.

4.1 Food

The major contribution of acylated flavonoids in the food industry lies in the improvement of stability and solubility of initial molecules, e.g. by reducing lipid oxidation in oil/fat based food systems, desirable modification of unwanted sensory properties of certain flavonoids, taking advantage of pigment stabilization by the means of flavonoid acylation, or other food characteristics. Furthermore, selectively acylated flavonoids may cause significant changes in their bioavailability and bioactivity, and when consumed, may thus play a role in preventing diseases.

Flavonoid acylation is a useful tool for modification of sensory properties of food. While flavonoids provide a variety of health benefits, flavonoid-containing food often suffers from bitter and astringent taste. Degenhardt et al. (2007) found that certain glycosylation and acylation patterns can effectively modulate these negative taste factors in edible preparations, pharmaceutical preparations and cosmetics with mouth contact (i.e. tooth paste, mouth wash). Both the taste intensity and the taste profile perception are improved by the novel compounds. Ghoul et al. (2006) introduced a process for the selective preparation of acylated flavonoid glycosides with improved stability and solubility in various preparations with their antioxidant effect remaining intact or being improved.

Another particular advantage obtained by these modified flavonoids is the bifunctional character of their molecule with higher biological activity. Free unsaturated fatty acids represent a potential risk because they are highly reactive and by creating free radicals they cause undesirable damage in food. Enzymatic synthesis of flavonoids with unsaturated fatty acids was found to be a useful solution for the stabilization of these highly oxidizable acids (Viskupicova et al., 2010; Mellou et al., 2006).

Another important benefit of acylated anthocyanins lies in the use as food colorants which can serve as a useful alternative to synthetic additives (Giusti & Wrolstad, 2003; Fox, 2000; Asen et al., 1979). The discovery of acylated anthocyanins with increased stability has shown that these pigments may provide food products with the desirable color and stability at a wide pH range. Examples of suitable acylated anthocyanin sources may be radishes, red

potatoes, red cabbage, black carrots, and purple sweet potatoes (reviewed in Giusti & Wrolstad, 2003). The invention of Asen et al. (1979) refers to a stable food colorant from a natural source. It relates to an anthocyanin isolated from the Heavenly Blue Morning Glory (*Ipomoea tricolor* Cav cv), peonidin 3-(dicaFFEylsophoroside)-5-glucoside, which is characterized by the stability of colors ranging from purplish-red to blue produced in food and beverage products at pH values from about 2.0 to about 8.0. Fox (2000) reported the invention referring to a stable, ruby red natural colorant (anthocyanins acylated with chlorogenic acid) derived from purple sunflower hulls, useful as a coloring agent in food products, cosmetics, pharmaceuticals and other materials.

4.2 Pharmaceuticals

In recent years, coronary artery diseases, such as atherosclerosis and hypercholesterolemia, represent a major cause of death, exceeding even oncological causes or infectious diseases. Novel acylated flavanone derivatives are effective in the treatment or prevention of elevated blood lipid level-related diseases, e.g. hyperlipidemia, arteriosclerosis, angina pectoris, stroke and hepatic diseases since they exert inhibitory effects on acylcolicholesterol acyl transferase activity and HMG-CoA reductase activity. In spite of their potent efficacies, the flavanone derivatives exhibited no toxicity or mitogenicity in tests using mice (Bok et al., 2001).

Mellou et al. (2005) carried out enzymatic acylation on Greek endemic plants and reported that this modification increased both their antioxidant activity towards isolated low-density lipoproteins (LDL) and serum model and antimicrobial activity against two Gram-positive bacteria, *Staphylococcus aureus* and *Bacillus cereus*. Katsoura et al. (2006) also found that biocatalytic acylation of rutin with various acyl donors affected its antioxidant potential towards both isolated LDL and total serum model *in vitro*. A significant increase in antioxidant activity was observed for rutin-4'''-oleate.

The 6''-O-esterification of kaempferol-3-O-glucoside (astragalin) with *p*-coumaric acid was found to increase its anti-inflammatory activity eight times compared to the initial flavonoid, while addition of another *p*-coumaroyl group at 2'' position gave an activity 30 times greater than that of astragalin (Harborne & Williams, 2000). Another kaempferol derivative, kaempferol 3-(2'',3''-di-E-*p*-coumaroyl)rhannoside, was found to possess a cytotoxic effect. It significantly modulated the proliferation of promyelocytic cell line HL60 and MOLT3 (a T-ALL with phenotypic characteristics of cortical thymocytes) (Mitrokotsa et al., 1993). Also Demetzos et al. (1997) synthesized novel flavonoid esters with cytotoxic activity. These acetylated esters of tiliroside exhibited a strong cytotoxic effect against four leukemic cell lines (HL60, DAUDI, HUT78 and MOLT3), whilst the maternal compound had no effect (Demetzos et al., 1997). Tricin-7-O-β-(6''-methoxycinnamic)-glucoside, a flavone from sugarcane, was found to exhibit antiproliferative activity against several human cancer cell lines, with higher selectivity toward cells of the breast resistant NIC/ADR line (Duarte-Almeida et al., 2007). Mellou et al. (2006) provided evidence that flavonoid derivatives esterified with polyunsaturated fatty acids were able to decrease the production of vascular endothelial growth factor by K562 human leukemia cells unlike the initial flavonoids, indicating that these novel compounds might possess improved anti-angiogenic and anti-tumor properties. Anticancer activity was established also in two O-acylated flavonoids, daglesiosides I and II, which were isolated from the leaves of *Pseudotsuga menziesii* (Sharma et al., 2003).

Parejo et al. (2005) examined quercetagen glycosides acylated with caffeic and *p*-coumaric acid for antioxidant activity. They found that these compounds exhibited a high radical scavenging activity in comparison with reference compounds. Fatty acid derivatives of catechins are described as having antitumorigenesis promoting activity or 5- α reductase inhibiting activity, as well as antibacterial activity (Fukami et al., 2007). Since these acylated catechin compounds have a greatly superior solubility in fats and oils than any catechins previously known, they may be used as a highly effective antioxidative agents (Sakai et al., 1994).

A different catechin derivative, 3-*O*-octanoyl-(+)-catechin, was synthesized by Aoshima et al. (2005) by incorporation of an octanoyl chain into (+)-catechin. This ester was found to be more efficient than catechin in inhibiting the response of ionotropic gamma-aminobutyric acid receptors and Na⁺/glucose cotransporters expressed in *Xenopus oocytes* in a noncompetitive manner. Moreover, it induced a nonspecific membrane current and decreased the membrane potential of the oocyte. This newly synthesized catechin derivative possibly binds to the lipid membrane more strongly than do catechin, (-)-epicatechin gallate, or (-)-epigallocatechin-3-gallate, and as a result it perturbs the membrane structure (Aoshima et al., 2005).

4.3 Cosmetics

The majority of cosmetic or dermatopharmaceutical compositions consist of a fatty phase, the oily products of which have a certain tendency to oxidize, even at room temperature. The consequence of this oxidation is to profoundly modify the properties, which makes them unusable after a variable time period. In order to protect the compositions with respect to these oxidation phenomena, it is common practice to incorporate protective agents which act as antioxidizing agents (N'guyen, 1995). By virtue of the skin-protecting and skin-cleansing properties of flavonoids and their effects against aging, against skin discoloration and on the appearance of the skin, they have been used as constituents of cosmetic or dermatopharmaceutical compositions. They also act on the mechanical properties of hair (Ghoul et al., 2006).

Moussou et al. (2007) found that the esters of flavonoids with omega-substituted C6 to C22 fatty acids have the property to protect the skin cells against damage caused by UV radiation. According to the invention, these esters of flavonoids protect skin cells against UVA and UVB radiation in a more effective manner than flavonoids alone. Moreover, these esters demonstrated their property to stimulate glutathione metabolism of human skin cells after UVA irradiation, i.e. to stimulate their cellular defenses. They have also anti-inflammatory and soothing properties, as demonstrated by the inhibition of released protein kinase PGE2 after UVB irradiation. Thus these flavonoid esters may be used to protect the skin and scalp and/or to fight against UV and sun damage, erythema, sunburn, mitochondrial or nuclear DNA damage, to prevent or fight photo-aging, providing improvement for signs of aging as skin wrinkles, elasticity lost and decrease in skin thickness (Moussou et al., 2007).

Perrier et al. (2001) discovered that specific flavonoid esters can be stabilized while preserving their initial properties, particularly free radical inhibition and enzyme inhibition, and for applications associated with these properties: venous tonics, agents for increasing the strength of blood capillaries, inhibitors of blotchiness, inhibitors of chemical, physical or actinic erythema, agents for treating sensitive skin, decongestants, draining agents, slimming agents,

anti-wrinkle agents, stimulators of the synthesis of the components of the extracellular matrix, toners for making the skin more elastic and anti-ageing agents (Perrier et al., 2001).

5. Conclusions

Flavonoids, having a wide spectrum of health-beneficial activities, seem to be applicable in various areas of national management from food additivization to pharmaceutical preparations with the purpose of prevention and/or treatment of important civilization diseases. Their chemical structure determines not only biological effects on human health but also their solubility, stability and bioavailability. Recently, selective enzyme-mediated acylation of flavonoids has been introduced to confer improved biological properties to the novel compounds including both biological activity of initial flavonoid and other parameters determined by the chemical structure of an acyl donor. In the past, proteases, esterases and acyltransferases were used for the preparation of acylated flavonoids. In light of our review, immobilized lipases, especially *Candida antarctica* B lipase, are suitable for this purpose. Not only the given enzyme but also the reaction conditions have a distinct influence on the performance of acylation. This aspect must be considered when producing acylated flavonoids in technology scale for potential uses in the food, pharmaceutical and cosmetic industry.

6. Acknowledgment

The work was supported by The Slovak Research and Development Agency in the frame of the Project APVV-VMSP-II-0021-09 and by The Agency of the Ministry of Education, Science, Research and Sport of the Slovak Republic for the Structural Funds of EU, OP R&D of ERDF in the frame of the Project „Evaluation of natural substances and their selection for prevention and treatment of lifestyle diseases (ITMS 26240220040).

7. References

- Adamczak, M. & Krishna, S.H. (2004). Strategies for improving enzymes for efficient biocatalysis. *Food Technology and Biotechnology*, Vol.42, pp. 251-264.
- Alluis, B. & Dangles, O. (1999). Acylated flavone glucosides: synthesis, conformational investigation, and complexation properties. *Helvetica Chimica Acta*, Vol.82, pp. 2201-2212.
- Andersen, Ø.M. & Jordheim, M. (2006). The anthocyanins. In: Andersen, Ø.M.; Markham K.R.: *Flavonoids: chemistry, biochemistry and applications*. New York: Taylor & Francis Group, CRC Press, 471 – 551. ISBN: 0-8493-2021-6.
- Anderson, E.M.; Larsson, K.M. & Kirk, O. (1998). One biocatalyst – many applications: the use of *Candida antarctica* B-lipase in organic synthesis. *Biocatalysis and Biotransformation*, Vol.16, pp. 181-204.
- Aoshima, H.; Okita, Y.; Hossain, S.J.; Fukue, K.; Mito, M.; Orihara, Y.; Yokoyama, T.; Yamada, M.; Kumagai, A.; Nagaoka, Y.; Uesatos, S. & Hara, Y. (2005). Effect of 3-O-octanoyl-(+)-catechin on the responses of GABA_A receptors and Na⁺/glucose cotransporters expressed in *Xenopus oocytes* and on the oocyte membrane potential. *Journal of Agricultural and Food Chemistry*, Vol.53, pp. 1955-1959.
- Ardhaoui, M.; Falcimaigne, A.; Engasser, J.-M.; Moussou, P.; Pauly, G. & Ghoul, M. (2004a). Acylation of natural flavonoids using lipase of *Candida antarctica* as biocatalyst. *Journal of Molecular Catalysis B: Enzymatic*, Vol.29, pp. 63-67.

- Ardhaoui, M.; Falcimaigne, A.; Ognier, S.; Engasser, J.-M.; Moussou, P.; Pauly, G. & Ghoul, M. (2004b). Effect of acyl donor chain length and substitutions pattern on the enzymatic acylation of flavonoids. *Journal of Biotechnology*, Vol.110, pp. 265-272.
- Ardhaoui, M.; Falcimaigne, A.; Engasser, J.-M.; Moussou, P.; Pauly, G. & Ghoul, M. (2004c). Enzymatic synthesis of new aromatic and aliphatic esters of flavonoids using *Candida antarctica* as biocatalyst. *Biocatalysis and Biotransformation*, Vol.22, pp. 253-259.
- Arroyo, M.; Sánchez-Montero, J.M. & Sinisterra, J.V. (1999). Thermal stabilization of immobilized lipase B from *Candida antarctica* on different supports: Effect of water activity on enzymatic activity in organic media. *Enzyme & Microbial Technology*, Vol.24, pp. 3-12.
- Asen, S.; Stewart, R.N. & Norris, K.H. (1979). Stable foods and beverages containing the anthocyanin, peonidin-3-(dicaffeoylsophoroside)-5-glucosid. Patent US 4172902.
- Augustyniak, A.; Bartosz, G.; Cipak, A.; Duburs, G.; Horakova, L.; Luczaj, W.; Majekova, M.; Odysseos, A.D.; Rackova, L.; Skrzydlewska, E.; Stefek, M.; Strosova, M.; Tirzitis, G.; Viskupicova, J.; Vraha, P.S. & Zarkovic, N. (2010). Natural and synthetic antioxidants: an updated overview. *Free Radical Research*, Vol.44, pp. 1216-1262.
- Ballesteros, A.; Plou, F.J.; Alcade, M.; Ferrer, M.; Garcia-Arellano, H.; Reyes-Duarte, D. & Ghazi, I. (2006). Enzymatic synthesis of sugar esters and oligosaccharides from renewable resources. In: Patel, R.N.: *Biocatalysis in the pharmaceutical and biotechnology industries*. New York: CRC Press, Taylor & Francis Group, 463-488. ISBN: 0-8493-3732-1.
- Berg, P.A. & Daniel, P.T. (1988). Plant flavonoids in biology and medicine II. *Progress in Clinical and Biological Research*, Vol.280, pp. 157-171.
- Bloor, S.J. (2001). Deep blue anthocyanins from blue *Dianella* berries. *Phytochemistry*, Vol.58, pp. 923-927.
- Bok, S.-H.; Jeong, T.-S.; Lee, S.-K.; Kim, J.-R.; Moon, S.-S.; Choi, M.-S.; Hyun, B.-H.; Lee, C.-H. & Choi, Y.-K. (2001). Flavanone derivatives and composition for preventing or treating blood lipid level-related diseases comprising same. Patent US 6455577.
- Bornscheuer, U.T. (2002). Microbial carboxyl esterases: classification, properties and application in biocatalysis. *FEMS Microbiology Reviews*, Vol.26, pp. 73-81.
- Calis, I.; Ozipek, M. & Ruedi, P. (1995). Enzyme-mediated regioselective acylation of flavonoid glycosides. *FABAD Journal of Pharmaceutical Sciences*, Vol.20, pp. 55-59.
- Carrea, G.; Riva, S.; Secundo, F. & Danieli, B. (1989). Enzymatic synthesis of various 1-O-sucrose and 1-O-fructose esters. *Journal of the Chemical Society Perkin Transactions*, Vol.1, pp. 1057-1061.
- Chebil, L.; Humeau, C.; Falcimaigne, A.; Engasser, J.-M. & Ghoul, M. (2006). Enzymatic acylation of flavonoids. *Process Biochemistry*, Vol.41, pp. 2237-2251.
- Chebil, L.; Anthoni, J.; Humeau, C.; Gerardin, C.; Engasser, J.-M. & Ghoul, M. (2007). Enzymatic acylation of flavonoids: Effect of the nature of the substrate, origin of lipase, and operating conditions on conversion yield and regioselectivity. *Journal of Agricultural and Food Chemistry*, Vol.55, pp. 9496-9502.
- Collins, A.M. & Kennedy, M.J. (1999). Biotransformations and bioconversions in New Zealand: Past endeavours and future potential. *Australasian Biotechnology*, Vol.9, pp. 86-94.
- Cordova A.; Iverson T. & Hult K. (1998). Lipase catalysed formation of macrocycles by the ring opening polymerisation of ϵ -caprolactone. *Polymer*, Vol.39, pp. 6519-6524.
- Cushnie, T.P.T. & Lamb, A.J. (2005). Detection of galangin-induced cytoplasmic membrane damage in *Staphylococcus aureus* by measuring potassium loss. *Journal of Ethnopharmacology*, Vol.101, pp. 243-248.

- Cushnie, T.P.T. & Lamb, A.J. (2006). Assessment of the antibacterial activity of galangin against 4-quinolone resistant strains of *Staphylococcus aureus*. *Phytomedicine*, Vol.13, pp. 191-197.
- Cygler, M., & Schrag, J. (1997). Structure as a basis for understanding interfacial properties of lipases. *Methods in Enzymology*, Vol.284, pp. 3-27.
- Danieli, B.; De Bellis, P.; Carrea, G. & Riva, S. (1989). Enzyme-mediated acylation of flavonoid monoglycosides. *Heterocycles*, Vol.29, pp. 2061-2064.
- Danieli, B.; De Bellis, P.; Carrea, G. & Riva, S. (1990). Enzyme-mediated regioselective acylations of flavonoid disaccharide monoglycosides. *Helvetica Chimica Acta*, Vol.73, pp. 1837-1844.
- Danieli, B.; Luisetti, M.; Sampognaro, G.; Carrea, G. & Riva, S. (1997). Regioselective acylation of polyhydroxylated natural compounds catalyzed by *Candida antarctica* lipase B (Novozym 435) in organic solvents. *Journal of Molecular Catalysis B: Enzymatic*, Vol.3, pp. 193-201.
- Davis, B. G. & Boyer, V. (2001). Biocatalysis and enzymes in organic synthesis. *Natural Product Reports*, Vol.18, pp. 618-640.
- Davies, K.M. & Schwinn, K.E. (2006). Molecular biology and biotechnology of flavonoid biosynthesis. In: Andersen, Ø.M.; Markham, K.R.: Flavonoids: chemistry, biochemistry and applications. New York: Taylor & Francis Group, CRC Press, 143-218. ISBN: 0-8493-2021-6.
- Degenhardt, A.; Ullrich, F.; Hofmann, T. & Stark, T. (2007). Flavonoid sugar addition products, method for manufacture and use thereof. Patent US 20070269570.
- Degn P.; Pedersen L.H.; Duus J. & Zimmerman, W. (1999). Lipase catalysed synthesis of glucose fatty acid esters in *t*-butanol. *Biotechnology Letters*, Vol.21, pp. 275-280.
- Delazar, A.; Celik, S.; Gokturk, R.S.; Nahar, L. & Sarker, S.D. (2005). Two acylated flavonoid glycosides from *Stachys bombycina*, and their free radical scavenging activity. *Pharmazie*, Vol.60, pp. 878-880.
- Demetzos, C.; Magiatis, P.; Typas, M.A.; Dimas, K.; Sotiriadou, R.; Perez, S. & Kokkinopoulos, D. (1997). Biotransformation of the flavonoid tiliroside to 7-methylether tiliroside: bioactivity of this metabolite and of its acetylated derivative. *Cellular and Molecular Life Sciences*, Vol.53, pp. 587-592.
- Drouin, J.; Costante, J. & Guibé-Jampel, E. (1997). A thermostable microbial enzyme for fast preparative organic chemistry: Preparation of R-(+)-1-phenylethanol from (±)-1-phenylethyl pentanoate and *n*-butanol. *Journal of Chemical Education*, Vol.74, pp. 992-995.
- Duarte-Almeida, J.M.; Negri, G.; Salatino, A.; De Carvalho, J.E. & Lajolo, F.M. (2007). Antiproliferative and antioxidant activities of a tricin acylated glycoside from sugarcane (*Saccharum officinarum*) juice. *Phytochemistry*, Vol.68, pp. 1165-1171.
- Enaud, E.; Humeau, C.; Piffaut, B. & Girardin, M. (2004). Enzymatic synthesis of new aromatic esters of phloridzin. *Journal of Molecular Catalysis B: Enzymatic*, Vol.27, pp. 1-6.
- Ferrer, J.-L.; Austin, M.B.; Stewart, C. & Noel, J.P. (2008). Structure and function of enzymes involved in the biosynthesis of phenylpropanoids. *Plant Physiology and Biochemistry*, Vol.46, pp. 356-370.
- Foresti, M.L.; Pedernera, M.; Bucalá, V. & Ferreira, M.L. (2007). Multiple effects of water on solvent-free enzymatic esterifications. *Enzyme & Microbial Technology*, Vol.41, pp. 62-70.
- Fossati, E. & Riva, S. (2006). Stereoselective modifications of polyhydroxylated steroids. In: Patel, R.N.: Biocatalysis in the pharmaceutical and biotechnology industries. New York: CRC Press, Taylor & Francis Group, 591-604. ISBN: 0-8493-3732-1.

- Fox, G.J. (2000). Natural red sunflower anthocyanin colorant with naturally stabilized color qualities, and the process of making. Patent US 6132791.
- Fujiwara, H.; Tanaka, Y.; Yonekura-Sakakibara, K.; Fukuchi-Mizutani, M.; Nakao, M.; Fukui, Y.; Yamaguchi, M.; Ashikari, T. & Kusumi, T. (1998). cDNA cloning, gene expression and subcellular localization of anthocyanin 5-aromatic acyltransferase from *Gentiana triflora*. *The Plant Journal*, Vol.16, pp. 421-431.
- Fukami, H.; Nakao, M.; Namikawa, K. & Maeda, M. (2007). Esterified catechin, process for producing the same, food and drink or cosmetic containing the same. Patent EP 1849779.
- Gao, C.; Mayon, P.; Macmanus, D.A. & Vulfson, E. N. (2001). Novel enzymatic approach to the synthesis of flavonoid glycosides and their esters. *Biotechnology & Bioengineering*, Vol.71, pp. 235-243.
- Gayot, S.; Santarelli, X. & Coulon, D. (2003). Modification of flavonoid using lipase in non-conventional media: effect of the water content. *Journal of Biotechnology*, Vol.101, pp. 29-36.
- Ghoul, M.; Engasser, J.-M.; Moussou, P.; Pauly, G.; Ardhaoui, M. & Falcimaigne, A. (2006). Enzymatic production of acyl flavonoid derivatives. Patent US 20060115880.
- Giusti, M.M. & Wrolstad, R.E. (2003). Acylated anthocyanins from edible sources and their applications in food systems. *Biochemical Engineering Journal*, Vol.14, pp. 217-225.
- Haeflner, F.; Norin, T. & Hult, K. (1998). Molecular modeling of the enantioselectivity in lipase-catalyzed transesterification reactions. *Biophysical Journal*, Vol.74, pp. 1251-1262.
- Haraguchi, H.; Tanimoto, K.; Tamura, Y.; Mizutani, K. & Kinoshita, T. (1998). Mode of antibacterial action of retrochalcones from *Glycyrrhiza inflata*. *Phytochemistry*, Vol.48, pp. 125-129.
- Harborne, J.B. & Williams, C.A. (1998). Anthocyanins and other flavonoids. *Natural Product Reports*, Vol.15, 631-652.
- Harborne, J.B. & Williams, C.A. (2000). Advances in flavonoid research since 1992. *Phytochemistry*, Vol.55, pp. 481-504.
- Hidalgo, A. & Bornscheuer, U.T. (2006). Direct evolution of lipases and esterases for organic synthesis. In: Patel, R.N.: Biocatalysis in the pharmaceutical and biotechnology industries. New York: CRC Press, Taylor & Francis Group, 159-179. ISBN: 0-8493-3732-1.
- Ishihara, K. & Nakajima, N. (2003). Structural aspects of acylated plant pigments: stabilization of flavonoid glucosides and interpretation of their functions. *Journal of Molecular Catalysis B: Enzymatic*, Vol.23, No.2-6, pp. 411-417.
- Ishihara, K.; Nishimura, Y.; Kubo, T.; Okada, C.; Hamada, H. & Nakajima, N. (2002). Enzyme-catalyzed acylation of plant polyphenols for interpretation of their functions. *Plant Biotechnology*, Vol.19, pp. 211-214.
- Iso, M.; Chen, B.; Eguchi, M.; Kudo, T. & Shrestha, S. (2001). Production of biodiesel fuel from triglycerides and alcohol using immobilized lipase. *Journal of Molecular Catalysis B: Enzymatic*, Vol.16, pp. 53-58.
- Itoh, T.; Nishimura, Y.; Ouchi, N. & Hayase, S. (2003). 1-Butyl-2,3-dimethylimidazolium tetrafluoroborate: the most desirable ionic liquid solvent for recycling use of enzyme in lipase-catalyzed transesterification using vinyl acetate as acyl donor. *Journal of Molecular Catalysis B: Enzymatic*, Vol.26, pp. 41-45.
- Iwashina, T. (2003). Flavonoid function and activity to plants and other organisms. *Biological Science in Space*, Vol.17, pp. 24-44.

- Jain, N.; Kumar, A.; Chauhan, S. & Chauhan, S.M.S. (2005). Chemical and biochemical transformations in ionic liquids. *Tetrahedron*, Vol.61, pp. 1015-1060.
- Jarstoff, B.; Störmann, R.; Ranke, J.; Mölter, K.; Stock, F.; Oberheitmann, B.; Hoffmann, J.; Nüchter, M.; Ondruschka, B. & Filser, J. (2003). How hazardous are ionic liquids? Structure-activity relationships and biological testing as important elements for sustainability evaluation. *Green Chemistry*, Vol.5, pp. 136-142.
- Jeager, K.E.; Dijkstra, B.W. & Reetz, M.T. (1999). Bacterial biocatalysts: molecular biology, three-dimensional structures, and biotechnological applications. *Annual Review of Microbiology*, Vol.53, pp. 315-351.
- Jungblut, T.P.; Schnitzler, J.P.; Hertkorn, N.; Metzger, J.W.; Heller, W. & Sandermann, H. (1995). Acylated flavonoids as plant defense compounds against environmentally relevant ultraviolet-B radiation in *Scots pine* seedlings. *Phytochemicals and health. Proceedings, tenth annual Penn State Symposium in Plant Physiology*, Rockville, Md.: American Society of Plant Physiologists, 266 - 267.
- Katsoura, M.H.; Polydera, A.C.; Tsironis, L.; Tselepis, A.D. & Stamatis, H. (2006). Use of ionic liquids as media for the biocatalytic preparation of flavonoid derivatives with antioxidant potency. *Journal of Biotechnology*, Vol.123, pp. 491-503.
- Katsoura, M.H.; Polydera, A.C.; Katapodis, P.; Kolisis, F.N. & Stamatis, H. (2007). Effect of different reaction parameters on the lipase-catalyzed selective acylation of polyhydroxylated natural compounds in ionic liquids. *Process Biochemistry*, Vol.42, pp. 1326-1334.
- Kitamura, S. (2006). Transport of flavonoids: from cytosolic synthesis to vacuolar accumulation. In Grotewold, E.: *The science of flavonoids*. Springer Science Business Media, New York, 123-146. ISBN: 0-387-28821-X
- Kodelia, G.; Athanasiou, K. & Kolisis, F.N. (1994). Enzymatic synthesis of butyryl-rutin ester in organic solvents and its cytogenetic effects in mammalian cells in culture. *Applied Biochemistry and Biotechnology*, Vol.44, pp. 205-212.
- Kontogianni, A.; Skouridou, V.; Sereti, V.; Stamatis, H. & Kolisis, F.N. (2003). Lipase-catalyzed esterification of rutin and naringin with fatty acids of medium carbon chain. *Journal of Molecular Catalysis B: Enzymatic*, Vol.21, pp. 59-62.
- Kontogianni, A.; Skouridou, V.; Sereti, V.; Stamatis, H. & Kolisis, F.N. (2001). Regioselective acylation of flavonoids catalyzed by lipase in low toxicity media. *European Journal of Lipid Science and Technology*, Vol.103, pp. 655-660.
- Kragl, U.; Kaftzik, N.; Schofer, S. & Wasserscheid, P. (2006). U.S. Patent No. 20060211096.
- Kwon, C.H.; Dae, Y.S.; Jong, H.L. & Seung, W.K. (2007). Molecular modeling and its experimental verification for the catalytic mechanism of *Candida antarctica* lipase B. *Journal of Microbiology and Biotechnology*, Vol.17, No.7, pp. 1098-1105.
- Laane, C. (1987). Medium engineering for bioorganic synthesis. *Biocatalysis*, Vol.30, pp. 80-87.
- Lambusta, D.; Nicolosi, G.; Patti, A. & Piattelli, M. (1993). Enzyme-mediated regioprotection-deprotection of hydroxyl groups in (+)-catechin. *Synthesis*, Vol.11, pp. 1155-1158.
- Lee, S.; Lee, C.-H.; Moon, S.-S.; Kim, E.; Kim, C.-T.; Kim, B.-H.; Bok, S.-H.; Jeong, T.-S. (2003). Naringenin derivatives as anti-atherogenic agents. *Bioorganic & Medicinal Chemistry Letters*, Vol.13, pp. 3901-3903.
- Lin, S.F.; Lin, Y.-H.; Lin, M.; Kao, Y.-F.; Wang, R.-W.; Teng, L.-W.; Chuang, S.-H.; Chang, J.-M. Yuan, T.-T.; Fu, K.C.; Huang, K.P.; Lee, Y.-S.; Chiang, C.-C.; Yang, S.-C.; Lai, C.-L.; Liao, C.-B.; Chen, P.; Lin, Y.-S.; Lai, K.-T.; Huang, H.-J.; Yang, J.-Y.; Liu, C.-W.; Wei, W.-Y.; Chen, C.-K.; Hiipakka, R.A.; Liao, S. & Huang, J.-J. (2010). Synthesis and

- structure-activity relationship of 3-O-acylated (-)-epigallocatechins as 5 α -reductase inhibitors. *European Journal of Medicinal Chemistry*, Vol.45, pp. 6068-6076.
- Lozano, P.; De Diego, T.; Carrie, D.; Vaultier, M. & Iborra, J. L. (2004). Synthesis of glycidyl esters catalyzed by lipases in ionic liquids and supercritical carbon dioxide. *Journal of Molecular Catalysis A: Chemical*, Vol.214, pp. 113-119.
- Lutz, S. (2004). Engineering lipase B from *Candida antarctica*. *Tetrahedron: Asymmetry*, Vol.15, pp. 2743-2748.
- Martinelle, M.; Holmquist, M. & Hult, K. (1995). On the interfacial activation of *Candida antarctica* lipase A and B as compared with *Humicola lanuginosa* lipase. *Biochimica et Biophysica Acta*, Vol.1258, pp. 272-276.
- Mellou, F.; Lazari, D.; Skaltsa, H.; Tselepis, A.D.; Kolisis, F.N. & Stamatis, H. (2005). Biocatalytic preparation of acylated derivatives of flavonoid glycosides enhances their antioxidant and antimicrobial activity. *Journal of Biotechnology*, Vol.116, pp. 295-304.
- Mellou, F.; Loutrari, H.; Stamatis, H.; Roussos, C. & Kolisis, F.N. (2006). Enzymatic esterification of flavonoids with unsaturated fatty acids: Effect of novel esters on vascular endothelial growth factor release from K562 cells. *Process Biochemistry*, Vol.41, pp. 2029-2034.
- Middleton, E.Jr. & Chithan, K. (1993). The impact of plant flavonoids on mammalian biology: implications for immunity, inflammation and cancer. In.: Harborne, J.B.: The flavonoids: advances in research since 1986. London, Chapman and Hall, 619-652, ISBN 0-412-48070-0.
- Middleton, E.Jr.; Kandaswami, C. & Theoharides, T.C. (2000). The effects of plant flavonoids on mammalian cells: Implications for inflammation, heart disease, and cancer. *Pharmacological Reviews*, Vol.52, pp. 673-751.
- Mitrokotsa, D.; Mitaku, S.; Demetzos, C.; Harvala, C.; Mentis, A.; Perez, S. & Kokkinopoulos, D. (1993). Bioactive compounds from the buds of *Platinus orientalis* and isolation of a new kaempferol glycoside. *Planta Medica*, Vol.59, pp. 517-520.
- Moussou, P.; Falcimaigne, A.; Ghoul, M.; Danoux, L. & Pauly, G. (2007). Esters of flavonoids with ω -substituted C6-C22 fatty acids. Patent US 20070184098.
- Moussou, P.; Falcimaigne, A.; Pauly, G.; Ghoul, M.; Engasser, J.-M. & Ardhaoui, M. (2004). Preparation of flavonoid derivatives. Patent EP1426445.
- Nagasawa, T. & Yamada, H. (1995). Microbial production of commodity chemicals. *Pure & Applied Chemistry*, Vol.67, pp. 1241-1256.
- Nakajima, N.; Ishihara, K.; Hamada, H.; Kawabe, S.-I. & Furuya, T. (2000). Regioselective acylation of flavonoid glucoside with aromatic acid by an enzymatic reaction system from cultured cells of *Ipomoea batatas*. *Journal of Bioscience & Bioengineering*, Vol.90, pp. 347-349.
- Nakajima, N.; Ishihara, K.; Itoh, T.; Furuya, T. & Hamada, H. (1999). Lipase catalysed direct and regioselective acylation of flavonoid glucoside for mechanistic investigation of stable plant pigments. *Journal of Bioscience and Bioengineering*, Vol.61, pp. 1926-1928.
- Nakajima, N.; Sugimoto, M.; Yokoi, H.; Tsuji, H. & Ishihara, K. (2003). Comparison of acylated plant pigments: Light-resistance and radical-scavenging ability. *Bioscience, Biotechnology, and Biochemistry*, Vol.67, pp. 1828-1831.
- Nakayama, T.; Suzuki, H. & Nishino, T. (2003). Anthocyanin acyltransferases: specificities, mechanism, phylogenetics, and applications. *Journal of Molecular Catalysis B: Enzymatic*, Vol.23, pp. 117-132.

- Narayan, V.S. & Klivanov, A.M. (1993). Are water-immiscibility and apolarity of the solvent relevant to enzyme efficiency? *Biotechnology & Bioengineering*, Vol.41, pp. 390-393.
- N'guyen, Q.L. (1995). Cosmetic or dermopharmaceutical composition containing, in combination, a lauroylmethionate of a basic amino acid and at least one polyphenol. Patent US 5431912.
- Nicolosi, G.; Piattelli, M.; Lambusta, D. & Patti, A. (1999). Biocatalytic process for the preparation of 3-O-acyl-flavonoids. Patent WO/1999/066062.
- Otto, R.; Geers, B.; Weiss, A.; Petersohn, D.; Schlotmann, K. & Schroeder, K.R. (2001). Novel flavone glycoside derivatives for use in cosmetics, pharmaceuticals and nutrition. Patent EP 1274712.
- Parejo, I.; Bastida, J.; Viladomat, F. & Codina, C. (2005). Acylated quercetagenin glycosides with antioxidant activity from *Tagetes maxima*. *Phytochemistry*, Vol.66, pp. 2356-2362.
- Patel, R.N. (2006). Biocatalysis for synthesis for chiral pharmaceutical intermediates. In: Patel, R.N.: Biocatalysis in the pharmaceutical and biotechnology industries. New York: CRC Press, Taylor & Francis Group, 103-158. ISBN: 0-8493-3732-1.
- Passicos, E.; Santarelli, X. & Coulon, D. (2004). Regioselective acylation of flavonoids catalyzed by immobilized *Candida antarctica* lipase under reduced pressure. *Biotechnology Letters*, Vol.26, pp. 1073-1076.
- Patti, A.; Piattelli, M. & Nicolosi, G. (2000). Use of *Mucor miehei* lipase in the preparation of long chain 3-O-acylcatechins. *Journal of Molecular Catalysis B: Enzymatic*, Vol.10, pp. 577-582.
- Perrier, E.; Mariotte, A.-M.; Boumendjel, A. & Bresson-Rival, D. (2001). Flavonoide esters and their use notably in cosmetics. Patent US 6235294.
- Plaper, A.; Golob, M.; Hafner, I.; Oblak, M.; Solmajer, T. & Jerala, R. (2003). Characterization of quercetin binding site on DNA gyrase. *Biochemical and Biophysical Research Communications*, Vol.306, pp. 530-536.
- Pleiss, J.; Fischer, M. & Schmid, R. D. (1998). Anatomy of lipase binding sites: the scissile fatty acid binding site. *Chemistry and Physics of Lipids*, Vol.93, pp. 67-80.
- Quarenghi, M.V.; Tereschuk, M.L.; Baigori, M.D. & Abdala, L.R. (2000). Antimicrobial activity of flowers from *Anthemis cotula*. *Fitoterapia*, Vol.71, pp. 710-712.
- Rao, M.B.; Tanksale, A.M.; Ghatge, M.S. & Deshpande, V.V. (1998). Molecular and biotechnological aspects of microbial proteases. *Microbiology and Molecular Biology Reviews*, Vol.62, pp. 597-635.
- Rauha, J.P.; Remes, S.; Heinonen, M.; Hopia, A.; Kähkönen, M.; Kujala, T.; Pihlaja, K.; Vuorela, H. & Vuorela, P. (2000). Antimicrobial effect of Finnish plant extracts containing flavonoids and other phenolic compounds. *International Journal of Food Microbiology*, Vol.56, pp. 3-12.
- Reetz, M.T.; Wiesenhöffer, W.; Francio, G. & Leitner, W. (2003). Continuous flow enzymatic kinetic resolution and enantiomer separation using ionic liquid/supercritical carbon dioxide media. *Advanced Synthesis & Catalysis*, Vol.345, pp. 01221-01228.
- Rice-Evans, C.A. (2001). Flavonoid Antioxidants. *Current Medicinal Chemistry*, Vol.8, pp. 797-807.
- Riva, S. (2002). Enzymatic modification of the sugar moieties of natural glycosides. *Journal of Molecular Catalysis B: Enzymatic*, Vol.19-20, pp. 43-54.
- Riva, S.; Chopineau, J.; Kieboom, A.P.G. & Klivanov, A.M. (1988). Protease-catalyzed regioselective esterification of sugars and related compounds in anhydrous dimethylformamide. *Journal of the American Chemical Society*, Vol.110, pp. 584-589.

- Rocha, J.; Gil, M. & Garcia, F. (1999). Optimisation of the enzymatic synthesis of *n*-octyl oleate with immobilized lipase in the absence of solvents. *Journal of Chemical Technology & Biotechnology*, Vol.74, pp. 607-612.
- Ross, J.A. & Kasum, C.M. (2002). Dietary flavonoids: bioavailability, metabolic effects, and safety. *Annul Reviews in Nutrition*, Vol.22, pp. 19-34.
- Rubin-Pitel, S.B. & Zhao, H. (2006). Recent Advances in biocatalysis by directed enzyme evolution. *Combinatorial Chemistry & High Throughput Screening*, Vol.9, pp. 247-257.
- Saija, A.; Tomaino, A.; Trombetta, D.; Pellegrino, M.L.; Tita, B.; Messina, C.; Bonina, F.P.; Rocco, C.; Nicolosi, G. & Castelli, F. (2003). 'In vitro' antioxidant and photoprotective properties and interaction with model membranes of three new quercetin esters. *European Journal of Pharmaceutics and Biopharmaceutics*, Vol.56, pp. 167-174.
- Sakai, M.; Suzuki, M.; Nanjo, F. & Hara, Y. (1994). 3-O-acylated catechins and methods of producing same. Patent EP 0618203.
- Salem, J.H.; Chevalot, I.; Harscoat-Schiavo, C.; Paris, C.; Fick, M. & Humeau, C. (2011). Biological activities of flavonoids from *N. retusa* and their acylated derivatives. *Food Chemistry*, Vol.124, pp. 486-494.
- Saxena, R.K.; Ghosh, P.K.; Gupta, R.; Davidson, W.S.; Bradoo, S. & Gulati, R. (1999). Microbial lipases: potential biocatalysts for the future industry. *Current Science*, Vol.77, pp. 101-115.
- Sharma, G.; Singh, R.P.; Chan, D.C. & Agarwal, R. (2003) Silibinin induces growth inhibition and apoptotic cell death in human lung carcinoma cells. *Anticancer Research*, Vol.23, pp. 2649-2655.
- Singh, R.K. & Nath, G. (1999). Antimicrobial activity of *Elaeocarpus sphaericus*. *Phytotherapy Research*, Vol.13, pp. 448-450.
- Stapleton, P.D.; Shah, S.; Anderson, J.C.; Hara, Y.; Hamilton-Miller, J.M.T. & Taylor, P.W. (2004). Modulation of beta-lactam resistance in *Staphylococcus aureus* by catechins and gallates. *International Journal of Antimicrobial Agents*, Vol.23, pp. 462-467.
- Stefani, E.D.; Boffetta, P.; Deneo-Pellegrini, H.; Mendilaharsu, M.; Carzoglio, J.C.; Ronco, A. & Olivera, L. (1999). Dietary antioxidants and lung cancer risk: a case-control study in Uruguay. *Nutrition and Cancer*, Vol.34, pp. 100-110.
- Stepanovic, S.; Antic, N.; Dakic, I. & Svabic-Vlahovic, M. (2003). *In vitro* antimicrobial activity of propolis and synergism between propolis and antimicrobial drugs. *Research in Microbiology*, Vol.158, pp. 353-357.
- Stevenson, D.E.; Wibisono, R.; Jensen, D.J.; Stanley, R.A. & Cooney, J.M. (2006). Direct acylation of flavonoid glycosides with phenolic acids catalyzed by *Candida antarctica* lipase B (Novozym 435®). *Enzyme and Microbial Technology*, Vol.39, pp. 1236-1241.
- Suda, I.; Oki, T.; Masuda, M.; Nishiba, Y.; Furuta, S.; Matsugano, K.; Sugita, K. & Terahara, N. (2002). Direct absorption of acylated anthocyanin in purple-fleshed sweet potato into rats. *Journal of Agricultural and Food Chemistry*, Vol.50, pp. 1672-1676.
- Tarle, D. & Dvorzak, I. (1990). Antimicrobial activity of the plant *Cirsium oleraceum* (L.). Scop. *Acta Pharmaceutica Jugoslavica*, Vol.40, pp. 567-571.
- Tereschuk, M.L.; Riera, M.V.; Castro, G.R. & Abdala, L.R. (1997). Antimicrobial activity of flavonoids from leaves of *Tagetes minuta*. *Journal of Ethnopharmacology*, Vol.56, pp. 227-232.
- Toba, S. & Merz, K. M. (1997). The concept of solvent compatibility and its impact on protein stability and activity enhancement in non-aqueous solvents. *Journal of the American Chemical Society*, Vol.119, pp. 9939-9948.

- Torres, S. & Castro, G.R. (2004). Non-aqueous biocatalysis in homogeneous solvent systems. *Food Technology & Biotechnology*, Vol.42, pp. 271-277.
- Trodler, P. & Pleiss, J. (2008). Modeling structure and flexibility of *Candida antarctica* lipase B in organic solvents. *BMC Structural Biology* [online].
- Uppenberg, J.; Öhrner, N.; Norin, M.; Hult, K.; Kleywegt, G.J.; Patkar, S.; Waagen, V.; Anthonsen, T. & Jones, T.A. (1995). Crystallographic and molecular-modelling studies of lipase B from *Candida antarctica* reveal a stereospecificity pocket for secondary alcohols. *Biochemistry*, Vol.33, pp. 16838-16851.
- Van Rantwick, F.; Lau, R.M. & Sheldon, R.A. (2003). Biocatalytic transformations in ionic liquids. *Trends in Biotechnology*, Vol.21, pp. 131-138.
- Vakhlu, J. & Kour, A. (2006). Yeast lipases: enzyme purification, biochemical properties and gene cloning. *Electronic Journal of Biotechnology* [online].
- Viskupicova, J.; Danihelova, M.; Ondrejovic, M.; Liptaj, T. & Sturdik, E. (2010). Lipophilic rutin derivatives for antioxidant protection of oil-based foods. *Food Chemistry*, Vol.123, pp. 45-50.
- Viskupicova, J.; Maliar, T.; Psenakova, I. & Sturdik, E. (2006). Enzymatic acylation of naringin. *Nova Biotechnologica*, Vol.6, pp. 149-159 [In Slovak].
- Viskupicova, J. & Ondrejovic, M. (2007). Effect of fatty acid chain length on enzymatic esterification of rutin. In: Book of abstracts of the 1st International Conference of Applied Natural Sciences, November 7 – 9, 2007 (pp. 59), Trnava: UCM.
- Viskupicova, J.; Strosova, M.; Sturdik, E. & Horakova, L. (2009). Modulating effect of flavonoids and their derivatives on sarcoplasmic reticulum Ca^{2+} -ATPase oxidized by hypochloric acid and peroxynitrite. *Neuroendocrinology Letters*, Vol.30, pp. 148-151.
- Wang, Y.F.; Lalonde, J.J.; Momongan, M.; Bergbreiter, D.E. & Wong, C.H. (1988). Lipase-catalyzed irreversible transesterifications using enol esters as acylating reagents: preparative enantio and regioselective synthesis of alcohols, glycerol derivatives, sugars, and organometallics. *Journal of the American Chemical Society*, Vol.110, pp. 7200-7205.
- Williams, C.A. (2006). Flavone and flavonol O-glycosides. In: Andersen, Ø.M.; Markham, K.R.: Flavonoids: chemistry, biochemistry and applications. New York: Taylor & Francis Group, CRC Press, 397-441. ISBN: 0-8493-2021-6.
- Wilkes, J.S. (2004). Properties of ionic liquids solvents for catalysis. *Journal of Molecular Catalysis A: Chemical*, Vol.214, pp. 11-17.
- Xanthakis, E.; Theodosiou, E.; Magkouta, S.; Stamatis, H. & Loutrari, H. (2010). Enzymatic transformation of flavonoids and terpenoids: structural and functional diversity of the novel derivatives. *Pure & Applied Chemistry*, Vol.82, pp. 1-16.
- Xiao, Y.-M.; Wu, Q.; Wu, W.-B.; Zhang, Q.-Y. & Lin, X.-F. (2005). Controllable regioselective acylation of rutin catalyzed by enzymes in non-aqueous solvents. *Biotechnology Letters*, Vol.27, pp. 1591-1595.
- Yadav, G.D. & Piyush, S.L. (2003). Kinetics and mechanism of synthesis of butyl isobutyrate over immobilised lipases. *Biochemical Engineering Journal*, Vol.16, pp. 245-252.
- Zaks, A. & Klivanov, A.M. (1988). Enzyme catalysis in nonaqueous solvents. *Journal of Biological Chemistry*, Vol.263, pp. 3194-3201.

Biochemistry

Edited by Prof. Deniz Ekinçi

ISBN 978-953-51-0076-8

Hard cover, 452 pages

Publisher InTech

Published online 02, March, 2012

Published in print edition March, 2012

Over the recent years, biochemistry has become responsible for explaining living processes such that many scientists in the life sciences from agronomy to medicine are engaged in biochemical research. This book contains an overview focusing on the research area of proteins, enzymes, cellular mechanisms and chemical compounds used in relevant approaches. The book deals with basic issues and some of the recent developments in biochemistry. Particular emphasis is devoted to both theoretical and experimental aspect of modern biochemistry. The primary target audience for the book includes students, researchers, biologists, chemists, chemical engineers and professionals who are interested in biochemistry, molecular biology and associated areas. The book is written by international scientists with expertise in protein biochemistry, enzymology, molecular biology and genetics many of which are active in biochemical and biomedical research. We hope that the book will enhance the knowledge of scientists in the complexities of some biochemical approaches; it will stimulate both professionals and students to dedicate part of their future research in understanding relevant mechanisms and applications of biochemistry.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Jana Viskupicova, Miroslav Ondrejovic and Tibor Maliar (2012). Enzyme-Mediated Preparation of Flavonoid Esters and Their Applications, *Biochemistry*, Prof. Deniz Ekinçi (Ed.), ISBN: 978-953-51-0076-8, InTech, Available from: <http://www.intechopen.com/books/biochemistry/enzyme-mediated-preparation-of-flavonoid-esters-and-their-applications>

INTech
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](https://creativecommons.org/licenses/by/3.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen