

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

185,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Photosynthetic Inhibitors

Róbson Ricardo Teixeira, Jorge Luiz Pereira and Wagner Luiz Pereira
*Federal University of Viçosa, Viçosa, Minas Gerais State
 Brazil*

1. Introduction

Life on Earth is dependent on sunlight. In the process known as photosynthesis, plants, algae and certain bacterias are capable of using this source of energy to drive the synthesis of organic compounds. The oxygenic photosynthesis results in the release of molecular oxygen and the removal of carbon dioxide from the atmosphere that is used to synthesize carbohydrates. This process, which involves a complex series of electron transfer reactions, provides the energy and reduced carbon required for the survival of virtually all life on our planet, as well as the molecular oxygen necessary for the survival of oxygen consuming organisms (Nelson, 2011; Nelson & Yocum, 2006; Rutherford & Faller, 2001).

In modern agriculture, farmers continuously face a battle to achieve products in high yields and better quality to feed an ever increasing world population (Stetter & Lieb, 2000). The optimization of agriculture techniques demands, along with other requirements, the application of crop protection agents to control a variety of diseases and pests, among which are weeds. Weeds compete with crops for nutrients, water, and physical space, may harbor insect and disease pests, and are thus capable of greatly undermining both crop quality and yield. In view of the problems caused by weed species, their control is highly desirable.

Among the methods to control weeds, the use of herbicides or weed killers has become the most reliable and least expensive tool for weed control in places where highly mechanized agriculture is practiced. Since the introduction of 2,4-dichlorophenoxyacetic acid (2,4-D) in 1946, several classes of herbicides were developed that are effective for broad-spectrum of weed control (Böger et al, 2002; Cobb, 1992; Ware, 2000).

It is well know that various compounds can interfere with photosynthetic electron transport. This fact has been explored by agrochemical companies to develop an assortment of herbicides to control weeds. Some representative members of commercial photosynthetic inhibitors are diuron (1), atrazine (2), paraquat (3) e diquat (4) (Figure 1).

The photosynthetic inhibitors can be divided into two distinct groups, the inhibitors of photosystem II exemplified by diuron (1) and atrazine (2) and the inhibitors of photosystem I such as paraquat (3) and diquat (4). It is worth to mention that compounds that inhibit photosystem II account for 30% of the sales in the herbicide market (Draber, 1992).

Although there are a variety of herbicides that can control a broad spectrum of weeds, there is still a necessity for the development of new active ingredients. Several reasons can be mentioned to support this statement. Herbicides should have a favorable combination of properties, such as high specific activity, low application rates, crop tolerance, and low mammalian toxicity. Increasing public concern for environmental pollution derived from

agricultural practices also requires that herbicides be rapidly degraded by soilborne microorganisms. Moreover, during the past years, intensive and repeated applications of the same active ingredients cause the selection for and development of herbicide resistance (Devine & Shukla, 2000; Beckie, 2006; Gressel, 2009; Preston, 2004). Starting from 1960s, hundreds of weed biotypes have been reported as surviving herbicide application (Heap, 2011).

Intensive efforts have thus been undertaken to discover new compounds with favorable environmental and safety features to selectively control weeds. In this regard, the photosynthetic system has been target aiming to find new weed killers. In this book chapter, it will be covered the developments concerning the search for new photosynthetic inhibitors during the last ten years.




Fig. 1. Examples of commercial photosynthetic inhibitors.

2. Natural products as source of photosynthetic inhibitors

The strategies used to identify new chemical agents to control weeds can no longer be distinguished from pharmaceutical research and development (Short, 2005). Three major different approaches have been employed. The first one refers to the systematic screening of large numbers of synthetic compounds. Subsequently, lead compounds are optimized. This has been the most widely used strategy by agrochemical companies (Böger et al, 2002; Cobb, 1992; Rüegg et al, 2006; Ware, 2000). The second one is the rational design of specific inhibitors of key metabolic processes (Lein et al, 2004). However, to date such an approach has not been fully successful to produce commercial herbicides.

A third strategy is related to the exploitation of natural products either directly as herbicides (Copping & Duke, 2007) or as leads for the development of new herbicides (Barbosa et al, 2008; Dayan et al, 1999; Dayan et al, 2009; Duke et al, 2000a, 2002; Hütter, 2011; Macías et al, 2007; Pillmoor et al, 1993; Scharader et al, 2010; Strange, 2007; Vyvyan, 2002). This strategy can be considered attractive for several reasons. A vast array of compounds has been isolated from nature and most of them have not been evaluated for agrochemical purposes. Different from what happens to the majority of synthetic agrochemicals, most natural products are water soluble. Moreover, due to natural selection these compounds can present bioactivity in very low concentration. The great variety of chemical structures found in nature can afford chemical agents for weed control that are toxicologically and environmentally benign. Furthermore, the molecular sites where natural products exert their action can be quite different from known molecular targets (Duke et al, 2005; Duke,

Duke et al, 2000b). This is particularly important to overcome the resistance problem. It is worth to mention that the few natural products based herbicides mesotrione, sulcotrione, cinmethylin, bialaphos and glyphosate (vide infra) act on molecular sites that were not known before they were introduced.

Even though the great variety of natural products has been relatively little explored, several active principles were discovered in this way (Figure 2).




Fig. 2. Structures of compounds 5-11 mentioned in the text.

Bialaphos (5) and phosphinotricin (6) were originally isolated from various strains of the bacteria *Streptomyces* spp. (Saxena & Pandey, 2001). Bialaphos is a proherbicide that is metabolized into the active ingredient phosphinotricin in the treated plant. Currently, bialaphos is commercialized in Japan with the name of Herbiace®. It should be mentioned that phosphinotricin (6) is also produced synthetically as a racemic mixture and commercialized as glyphosate. Leptospermone (7), a major component of the essential oil of *Leptospermum scoparium* (Myrtaceae) (van Klink et al, 1999), was chemically modified to make mesotrione (Mitchell et al, 2001). Mesotrione (8) is the active principle of the commercial herbicide Callisto®, which is commercialized by Syngenta and suitable for use in corn fields. Another example is sulcotrione (Chaabane et al, 2005), an herbicide marketed in Europe by Bayer Crop Science under the trade name Mikado®. Sulcotrione (9) is used to control a broad range of annual and perennial broadleaf weeds in maize and sugar cane crops. The commercial herbicide cinmethylin (11) (Figure 2) is a 2-benzyl ether derivative of the natural product 1,4-cineole (10) that was developed to control annual grasses (Romagni et al, 2000a,b; Duke & Oliva, 2004). Moreover, an increasing number of other natural products have been described in the literature as potential leads for the development of chemical agents for weed control among which are coumarins, benzoquinones, flavonoids, terpenoids and lactones (Barbosa et al, 2008). The natural product pool has been explored in the search of new photosynthetic inhibitors and the advances in this regard will be described below.

The diterpene labdane-8 α ,15-diol (**12**) and its acetyl derivative (**13**) (Figure 3) were isolated from the hexan extract of the stems of *Croton clatoglanduliferus*. Biological assays carried out with intact spinach chloroplasts showed that these compounds are capable of interfere with ATP synthesis (Morales-Flores et al., 2007).





Fig. 3. Structures of labdane-8 α ,15-diol and its acetyl derivative.

It is well established that in the oxygenic photosynthesis there is a transfer of electron from water to NADP⁺ affording the reduced form NADPH. In this biochemical process, two different reaction centers, known as photosystem II and photosystem I, work concurrently but in series. In the presence of light photosystem II feeds electrons to photosystem I. The electrons are transferred from photosystem II to the photosystem I by intermediate carriers as described by the well known Z scheme of photosynthesis (Figure 4). The net reaction is the transfer of electrons from water to NADP⁺.

It has been proposed that electron transport is indirectly coupled to phosphorylation (ATP synthesis) through an electrochemical potential of hydrogen ions (protons) build up across the biological membranes involved in the electron transport (Mitchel, 1961). The electrochemical gradient, in turn, is consumed in the formation of ATP from ADP and inorganic phosphate. There is proportionality between changes in pH and changes in hydrogen ion concentrations. The electron transport gives two protons translocated for each electron transferred from photosystem II to photosystem I (Allen, 2003). At pH 8.0 one proton ion is consumed irreversibly in the synthesis of ATP:



As can be noticed in the previous equation, ATP formation can be measured in vitro, using intact spinach chloroplast, by determining the basicity of the medium. In the presence of an artificial electron receptor such as methylviologen and under continuous actinic illumination, the pH rise continues to increase linearly with time corresponding to the steady state rate of ATP formation. Under appropriate conditions (Morales-Flores et al., 2007), the rate of ATP formation can be determined by back titration with hydrochloric acid using a microelectrode attached to a potentiometer.

Diterpenes (**12**) and (**13**) (Figure 3) inhibited ATP synthesis coupled to electron transport from water to methylviologen in freshly lysed intact spinach chloroplasts in a concentration dependent manner. The IC₅₀ (the concentrations causing 50% inhibition in vitro) were determined being equal to 72 $\mu\text{mol L}^{-1}$ for compound (**12**) and 10 $\mu\text{mol L}^{-1}$ for compound (**13**) (Morales-Flores et al., 2007). As mentioned above, the light-dependent phosphorylation is coupled to electron transport flow. Thus, ATP formation can be inhibited by (a) blockage of the electron transport within thylakoid chain, (b) by dissipation of the H⁺ gradient, that is, uncoupling of the ATP synthesis process from the electron transport, and (c) by direct inhibition of the H⁺-ATPase complex. Reagents that block electron transport avoid ATP synthesis because the generation of the transmembrane electrochemical gradient is not formed; as previously stated, the driving force for ATP synthesis is dependent upon electron flow. Chemicals that increase proton permeability of thylakoid membranes uncouple

phosphorylation from electron flow. Uncoupling agents inhibit ATP synthesis by decreasing the proton gradient but allow electron transport to occur at high rates. In contrast, direct inhibitors of photophosphorylation block both phosphorylation and that portion of electron transport that is a consequence of proton efflux linked to phosphorylation (Veiga et al, 2007a).




Fig. 4. The Z scheme and the electron transfer processes involved in oxygenic photosynthesis.

In order to elucidate the mechanism of action of (12) and (13) on photosynthesis, their effect on non-cyclic electron transport from water to methylviologen (basal, phosphorylating, and uncoupled) was investigated. It was found that as concentration of these compounds increased up to $300 \mu\text{mol L}^{-1}$, the rates of basal, phosphorylation, and uncoupled electron transport were inhibited being basal and uncoupled electron rates the more affected. The IC_{50} found for compound (12) were $200 \mu\text{mol L}^{-1}$ (basal) and $76 \mu\text{mol L}^{-1}$ (uncoupled) while for (13) IC_{50} for these rates were 31 and $71.5 \mu\text{mol L}^{-1}$ for basal and uncoupled, respectively. The partial reactions involved in the electron transport flow can be characterized in great detail by using highly specific artificial donors and acceptors (Trebst, 2007; Morales-Florest et al, 2007; as cited in Allen & Holmes, 1986). The effect of the natural products (12) and (13) (Figure 3) on photosystem I and II and their partial reactions were evaluated. It was determined that terpene (12) inhibited the partial reaction from water to DCPIP (2,6-dichlorophenol indophenol). Inhibition of this partial reaction indicates that the compounds inhibits the range of reactions of photosystem II electron transport chain from water to Q_B (Figure 4), because oxidized DCPIP accepts electrons at this level. It was also found that (12) inhibits the photosystem II partial reaction from water to sodium silicomolybdate (SiMo).

Inhibition of this partial reaction means that the (12) affects the partial reaction from water to quinone Q_A since SiMo is used as electron acceptor at or before Q_A site (Morales-Flores et al, 2007, as cited in Giaquinta & Dilley, 1975). The effect of compound (12) on the electron flow from diphenylcarbazide (DPC) to DCPIP was also evaluated. In this case, DCPD donates electrons at P_{680} level and the evaluation is carried out with Tris treated chloroplast. Such a treatment inhibits the oxygen evolving complex activity. Compound (12) did not inhibit this last partial reaction (electron flow from DPC to DCPIP). The results found for the activities of compound (12) on the partial reactions of photosystem II led to the indication that for this natural product the first site of interaction and inhibition is at the oxygen evolving complex. Further investigations carried out with terpene (12) showed that it is able to inhibit the partial reaction from reduced phenylmetasulfate (PMS) and methylviologen. While PMS donates electrons at the P_{680} level, methylviologen accepts electrons at F_x level in photosystem I. Therefore, it was established that the second site of interaction of compound (12) is located between P_{700} and F_x of photosystem I electron transport chain (Morales-Flores et al, 2007).

The effect of (13) on the partial reactions mentioned above was also evaluated being found the following results. It inhibited the partial reactions from water to DCPIP, from water to SiMo, and from DPC to DCPIP. These results indicate that the compound inhibits the electron flow in photosystem II in the evolving complex system as well as in the path between P_{680} and Q_A or at least at P_{680} site. The labdane 13 also inhibited the partial reaction from reduced PMS to oxidized methylviologen (Morales-Flores et al, 2007).

The survey of literature revealed that various other terpenes (Figure 5) have been evaluated as potential inhibitors of photosynthesis in a similar manner described for compounds (12) and (13).




Fig. 5. Structures of photosynthetic inhibitor terpenes 14-18.

The sesquiterpenoids (14) and (15) were isolated from the dichloromethane extract of the leaves of *Celastrus vulcanicola* (Torres-Romero et al, 2008). Similar to what happens to other terpenes, both compounds are capable of inhibiting the photosynthetic photophosphorylation in isolated spinach chloroplasts from water to methylviologen in a concentration-dependent manner ($IC_{50} = 25 \mu\text{mol L}^{-1}$ for 14 and $IC_{50} = 44 \mu\text{mol L}^{-1}$ for 15). At lower concentrations (up to $50 \mu\text{mol L}^{-1}$) compound (14) inhibited basal and phosphorylating electrons rates (approximately 83% and 79%, respectively); however, at

higher concentrations this terpenes reversed both electron transport rates (at 300 $\mu\text{mol L}^{-1}$, the electron flow rates were 32% and 68% for basal and phosphorylating, respectively). In addition, this compound also inhibited the uncoupled electron transport rate. Further investigations showed that (14) has two targets of interaction: one is located at the oxygen-evolving complex, and the other located at the H^+ -ATPase complex. Compound (15) behaves likewise to (14) albeit being less active (Torres-Romero et al, 2008).

The fractioning of the hexane extract from air-dried leaves of *Maytenus imbricata* led to the isolation of seco-3,4-triterpenoid (16). It inhibits the formation of ATP coupled to electron transport from water to methylviologen and its IC_{50} was 148 $\mu\text{mol L}^{-1}$. Evaluation of the effect of increasing concentration of (16) on basal, phosphorylating, and uncoupled electron flow rates showed that the phosphorylating and uncoupled electron flow are inhibited up to 300 $\mu\text{mol L}^{-1}$; after this concentration both activities are enhanced. Experimental results support the fact that this triterpenoid acts on the oxygen evolving complex since it inhibits the electro flow from water to sodium silicomolybdate (SiMo). Moreover, it behaves as uncoupler activating the Mg^{+2} -ATPase complex (Souza & Silva et al, 2007).

Epifriedelinol (17) and canophyllol (18) (Figure 5) are friedelane triterpenoids isolated from the stems of *Celastrus vulcanicola*. Their activities on photosynthesis as determined in vitro can be summarized as follows. Epifriedelinol (17) and canophyllol (18) inhibits the ATP synthesis coupled to electron transport from water to methylviologen as concentration increases presenting $\text{IC}_{50} = 82 \mu\text{mol L}^{-1}$ for compound (17) and 124 $\mu\text{mol L}^{-1}$ for compound (18). The evaluation of their effects on noncyclic electron transport from water to methyl viologen under basal, phosphorylating, and uncoupled conditions showed that epifriedelinol (17) did not affect these processes. On the other hand, compound 18 partially inhibited the electron transport rates at different degrees. Both compounds moderately enhanced the light-activated Mg^{+2} -ATPase activity indicating that they affect the variation of pH and thus avoiding the ATP formation. Thus, the friedelane triterpenoids (17) and (18) has two targest of interaction: one is a decrease of pH variation blocking the ATP formation and the second by interacting and inhibiting the Mg^{+2} -ATPase activity in thylakoids. It was also found that compound (18) is capable of inhibiting the electron flow from water to the electron acceptor 2,5-dichloro-1,4-benzoquinone (DCBQ) acting, therefore, as Hill reaction inhibitor (Torres-Romero et al, 2010).

Besides terpenes, the effects of several other natural products (Figure 6) as photosynthetic inhibitors have been investigated.

Demuner and co-workers reported the isolation of 4-methoxy-5-methyl-6-(3-methylbut-2-enyloxy)isobenzofuran-1(3H)-one (19) from the phytopathogenic fungus *Nympha alternantherae* (Demuner et al, 2006). This isobenzofuranone acts as Hill reaction inhibitor and uncoupler of photosynthesis, displaying inhibitory activity on ATP synthesis ($\text{IC}_{50} = 66 \mu\text{mol L}^{-1}$).

Siderin (20) is a secondary metabolite produce by *Toona ciliate* (Meliaceae). Veiga and co-workers showed that this coumarin inhibited ATP synthesis presenting $\text{IC}_{50} = 27.0 \mu\text{mol L}^{-1}$ and did not inhibit photosystem I electron transport. Siderin does inhibit partial reactions of photosystem electron flow frow water to DCPIP, from water do sodium silicomolybdate (SiMo), and partially inhibits electron flow from DPC to DCPIP. All these results support the fact that the site of inhibition of (20) is the donor and acceptor sites of photosystem II, between P_{680} and Q_A (Veiga et al, 2007b).

The lactone lasiodiplodin (21) was isolated from the ethanolic extract from the fungus *Botryosphaeria rhodina*. As the concentration of (21) is raised, ATP synthesis is inhibited

($IC_{50} = 35.6 \mu\text{mol L}^{-1}$). The investigation of the phytotoxic effect of this compound on photosynthesis resulted in the identification of three new different sites of interaction and inhibition: one at CF_1 ATPase complex, the second in the oxygen evolving complex, and the third at the electron transfer path between P_{680} and Q_A . These targets are different from that of displayed by synthetic herbicides. This finding corroborates the fact that the exploitation of the natural product pool can afford compounds presenting different molecular targets compared to well known herbicides (Veiga et al, 2007a).




Fig. 6. Structures of compounds **19-28** mentioned in the text.

The bioactivity-guided chemical analysis of *Selaginella lepidophylla* resulted in the isolation of the biflavonoids robustaflavone (**22**), 2,3-dihydrorobustaflavone (**23**), and 2,3-dihydrorobustaflavone-5-methyl ether (**24**) (Figure 6). The in vitro assays revealed that all the isolated flavonoids inhibited the ATP synthesis coupled to electron flow from water to methylviologen in isolated freshly lysed intact spinach chloroplasts. The IC_{50} values for this activity were 44, 39, and $79 \mu\text{mol L}^{-1}$ for compounds (**22**), (**23**), and (**24**), respectively. Considering the three modes of electron transport (basal, uncoupling and phosphorylating) from water to methylviologen, all of them are partially inhibited by the three flavonoids being compound (**22**) the most active. The mode of action of these compounds were further investigated. Compound (**22**) did not affect the photosystem I. However, it interacts with photosystem II in the span of oxygen evolving complex to P_{680} . The interaction and inhibition target of (**23**) was located at $Cytb_6f$ complex to plastocyanin (PC) (Figure 4). Flavonoid (**24**) had no effect on photosystem I or II. However, it acts as energy transfer inhibitors since increasing concentrations of (**22**), (**23**), and (**24**) inhibited the Mg^{+2} -ATPase activity (Aguilar et al, 2008).

A set of fifteen substances belonging to two groups of organic compounds, nonenolides and cytochalasins, had their phytotoxic effects evaluated against the perennial weed species

Cirsium arvense and *Sonchus arvensis* (Berestetskiy et al, 2008). Among the nonelides evaluate by leaf disc-puncture bioassay, stagonolide A (25) displayed the highest phytotoxic effect (necrosis diameter > 6 mm) on leaves of *C. arvense*. Considering the cytochalasins, only cytochalasyn A (26) showed high phytotoxic effect on this weed species (necrosis diameter ~3 mm) (Berestetskiy et al, 2008).

The most phytotoxic compound against *S. arvensis* was the cytochalasyn deoxaphomin (27) (necrosis diameter ~7 mm). High phytotoxic effects were also observed for stagonolide A (25), cytochalasin A (26) and cytochalasin B (28) (necrosis diameter ~ 4.5, 5.5 and 4 mm, respectively) (Berestetskiy et al, 2008).


Photometric assays in the range of 450-950 nm showed that stagonolide A (25) and cytochalasin B (28) presented effects on photosynthesis. Twenty four hours after the observation of the first necrosis on leaf discs, both compounds caused significant decrease of the light absorption at the wave length 450 nm by *C. arvense*. This experimental observation was associated with the reduction of β -carotene or/and chlorophyll *b* content in leaf tissue of *C. arvense* since these pigments have a peak of resonant absorption near the 450 nm. At the wavelength of 530 nm and 550 nm, it was noticed increase in the light absorption for compounds (25) and (28). However, stagonolide A (25) had significantly stronger effect at 550 nm compared to cytochalasin B (28). Since cytochromes have a peak absorption in the range of 530-550 nm the substances (25) and (28) probably increased the concentration of these proteins, and did not affect electron transport. Treatment of leaf discs by stagonolide (25) also led to reduction of light absorption in the wavelength region between 630-690 nm by *C. arvense* leaves (Berestetsky et al, 2008). The peaks of light absorption in this region are characteristic of chlorophyll intermediates, phytychlorophyllide and chlorophyllide (Berestetskiy et al, 2008, as cited by Duke et al, 1991).

3. Synthetic studies towards the discovery of new photosynthetic inhibitors

In the research and development of new agrochemicals, organic synthesis plays an important role. After the discovery of a lead structure, an optimization process of it is carried out in order to improve the biological activity as well physico-chemical properties of the lead. Within this context, organic chemists utilize a vast array of chemical reactions to prepare derivatives of a lead compound. In this section it will be discussed recent advances concerning synthetic studies aimed to discover new photosynthetic inhibitors. Details about the preparation of the mentioned compounds can be found in the cited references and will not be covered herein since this is not the focus of this book chapter.

Several papers has been published in the last few years dealing with the synthesis of various aromatic nitrogenated compounds and their biological evaluation as photosynthetic inhibitors (Jampilek et al, 2009a,b; Musiol et al, 2007; Musiol et al, 2008; Musiol et al, 2010; Otevrel et al, 2010). The general structures and their associated most active compounds are presented in Figure 7 along with the IC₅₀ data. For all of these substances, the in vitro evaluation of their inhibitory activity on the electron transport in isolated intact spinach chloroplasts was determined spectrophotometrically using the artificial electron acceptor 2,6-dichlorophenol indophenol (DCPIP). The rate of photosynthetic electron transport was monitored as photoreduction of DCPIP. The biological assays utilized diuron (1) (Figure 1) as positive control which the IC₅₀ determined under the conditions used in the experiments was 1.9 $\mu\text{mol L}^{-1}$.


Quinoline is a structural motif found in various classes of bioactive compound. Musiol and co-workers prepared seventeen quinoline derivatives presenting the general structure (29). Some of the synthesized compound could not be biologically evaluated due to poor water solubility. The ten studied quinolines presented low inhibitory activity on photosynthetic electron transport with IC₅₀ values ranging from 26 to 487 μmol L⁻¹. Compound (29a) was the most active presenting IC₅₀ = 26 μmol L⁻¹ (Musiol et al, 2007).



Compound	IC ₅₀ (μmol L ⁻¹)
29a	26.0
30a	32.0
30b	95.0
30c	8.7
30d	107.0
30e	137.0
31a	16.0
31b	7.2
32a	157
32b	126
33a	33.5
33b	54.4
34a	2.7
34b	1.6
34c	1.6
34d	1.0
Diuron	1,9

Fig. 7. Structures of aromatic nitrogenated compounds 29-34.

For the two groups of amides (structures **30** and **31**) based on the quinoline scaffold system, it was found that the for the seventeen compounds of general structure (**30**) the most active ones were those possessing the nitro group (-NO₂) at the R¹ position (**30a-30e**). Compounds lacking this functionality at the specified position were completely inactive. Considering the quinolines (**31**), no simple structure-activity relationship explaining the observed activity was found being the two most active derivatives (**31a**) and (**31b**) (Musiol et al, 2008). In a series of twelve ring-substituted 4-hydroxy-1*H*-quinolin-2-one derivatives (general structure **32**), all of the compounds displayed very low inhibitory activity (IC₅₀ ranging from 126 to 925 µmol L⁻¹) on electron transport in photosynthesis from water to DCPIP (Jampilek et al, 2009a). The best activities were observed with derivatives (**32a**) and (**32b**). In another series of hydroxilated derivatives, among the fourteen ring-substituted 8-hydroxyquinoline derivatives (**33**), two compounds (**33a**, **33b**) showed moderate inhibitory activity (Musiol et al, 2010).



Compound	IC ₅₀ (µmol L ⁻¹)
35a	169±86
35b	0.32±0.19
35c	42±8
35d	32±8
35e	6.6±0.8
35f	2.6±0.6
36a	0.21±0.03
36b	18±4
36c	2.9±0.6
36d	0.57±0.11
Diuron	0.27±0.02
Hexazinone	0.11±0.01
Lenacil	0.08±0.02

Fig. 8. Structures of pyrazoles **35** and **36** and the corresponding IC₅₀ data.

The investigation of the influence of twelve ring substituted salicylanilides and carbamoylpehnylcarbamates, represented by the general structure (**34**), on photosynthetic apparatus afforded compounds (**34a-34d**) (Figure 7) with biological activities compared do diuron. The remaining evaluated compounds displayed low to moderate activity. In addition, it was also suggested based on experimental results that the site of action of the

compounds could be in Q_B , which is the second quinone acceptor on the oxidizing site of photosystem II.

The ability of ten pyrazole derivatives presenting the general structures (35) and (36) to act as photosystem II inhibitors was evaluated (Vicentini et al, 2004). In the presence of increasing concentrations of the compounds, it was observed that all of them are capable of interfering with the light-driven reduction of ferricyanide by isolated spinach chloroplasts. As can be seen in Figure 8, with the only exception of compound (35a), the IC_{50} ranged from 0.2 to 42 $\mu\text{mol L}^{-1}$. It is important to mention that the effectiveness of the most active compounds was comparable to three commercial herbicides (diuron, hexazinone and lenacil) used as positive controls in the experiments.

The pyrazoles (35) and (36) were also evaluated in vivo against the blue-green alga *Spirulina platensis* as well as the eukaryotic alga belonging to the genus *Chlorella*. Once again, for some derivatives the observed activities are remarkable and comparable to commercial herbicides used as positive controls (Vicentini et al, 2004).

Even more remarkable biological activity was displayed by another series of pyrazoles presenting the general structures (37) and (38) (Figura 9). All of these compounds inhibited the Hill reaction from water to ferricyanide in the presence of illuminated chloroplasts, with IC_{50} ranging from 10^{-6} mol L^{-1} to 10^{-4} mol L^{-1} . The efficacy of (37j) ($IC_{50} = 0.64 \mu\text{mol L}^{-1}$) is comparable to commercial herbicides such as diuron, lenacil and hexazinone ($IC_{50} = 0.27$, 0.08 and 0.11 $\mu\text{mol L}^{-1}$) (Vicentini et al, 2005).




Fig. 9. Structures of pyrazoles 37 and 38 mentioned in the text.

One strategy that could be used to discover novel herbicides is to synthesize analogues of a natural products which is known to present phytotoxic activity. It is also possible to take a structure of a phytotoxic natural product itself and carry out structural modifications on it. Such an approach can result in compounds with improved biological activity as well better physico-chemical properties. The natural product (39) (Figure 10) was isolated from *Pterodon polygalaeflorus*. Biological evaluation of (39) as potential inhibitor of ATP synthesis in isolated chloroplasts from spinach revealed that this substance is completely inactive. However, its β -lactone derivative (40) inhibited this process as concentration increases. For this compound, the IC_{50} was 90 $\mu\text{mol L}^{-1}$. Similar to other terpenes previously described in this chapter, lactone (40) did not affect photosystem I but it did inhibit electron transport through photosystem II by targeting the oxygen evolving complex as well as the redox

enzymes of the electron transport chain in the span between P_{680} and Q_A (King-Díaz et al 2006).





Fig. 10. Structure of diterpene (39) and its synthetic derivative (40).

Other derivatives of compound (39) have been recently evaluated and present phytotoxic properties on photosynthesis (King-Díaz et al, 2010).

The nostoclides (41) correspond to a pair of naturally-occurring lactones produced by a cyanobacterium (*Nostoc* sp.) symbiont of *Peltigera canina* (L.). It has been suggested that these chlorinated compounds may be allelopathic agents since *P. canina* cultures are usually not contaminated with microorganisms (Yang et al, 1993). In view of that, it was decided to investigate the potential phytotoxicity of nostoclides analogues (Barbosa et. al. 2006). In this context, several derivatives (general structures 42 and 43, Figure 11) were prepared, and their ability to interfere with the light-driven reduction of ferricyanide by isolated spinach chloroplasts thylakoid membranes (Hill reaction) was subsequently evaluated (Barbosa et al., 2007; Teixeira et al, 2008).

A number of nostoclides derivatives, at various degrees, exhibited inhibitory properties in the micromolar range against the basal electron flow from water to ferricyanide. As a general trend, the non-brominated derivatives (43) presented higher effectiveness than their brominated counterparts. The most active compounds (43a-43e) derivatives along with their IC_{50} values are presented in Figure 11.



Compound	Benzylidene group	IC_{50} ($\mu\text{mol L}^{-1}$)
43a	Z-4-nitrobenzylidene	1.7 ± 0.7
43b	Z-2-fluorobenzylidene	10.1 ± 3.2
43c	Z-4-trifluoromethylbenzylidene	8.3 ± 2.3
43d	Z-2-trifluoromethylbenzylidene	11.8 ± 4.5
43e	Z-4-ethylbenzylidene	9.1 ± 3.2

Fig. 11. Structure of natural product nostoclides and some synthetic analogues.

More recently, a QSAR (Quantitative Structure Activity Relationship) investigation was carried out on a series of nostoclide analogues presenting the general structure (43) to correlate molecular descriptions with their in vitro biological activity (the ability to interfere with light-driven reduction of ferricyanide by isolated spinach chloroplasts thylakoid membranes). The results of this investigation suggested that the degree of inhibition efficiency of this class of compounds is intimately associated with their polarity (Teixeira et al, 2009). Thus, it is likely that new nostoclide analogues with higher polarity could display improved biological activity. At the moment that this book chapter is written, there is no report on the literature concerning the synthesis of nostoclide analogues with higher polarity than the previous ones already prepared.

4. Conclusions

The ongoing need for new agents to control weeds has stimulated the search for new photosynthetic inhibitors. We described in this chapter a variety of compounds presenting this type of activity. The natural products have been explored toward this end resulting in the identification of compounds with various structural motifs. Such an approach has resulted in the discovery of photosynthetic inhibitors with new modes of action. This, in turn, can be helpful in dealing with resistance a problem to be faced in weed management. It is possible to anticipate that promising inhibitors of photosynthesis will certainly be found by exploring the natural product pool. From nature, it is also possible that more active compounds with low toxicity and improved selectivity will be found. Promising photosynthetic inhibitors has also been revealed by the synthetic studies. One important challenge in the field of weed management is related to selectivity. In other words, chemicals should exert their action only on weeds. As can be noticed in the discussion of the studies published in the literature during the last years, this issue has not been addressed. Future works should also be concerned with this important matter.

5. Acknowledgment

We are grateful to Fundação de Amparo à Pesquisa de Minas Gerais (FAPEMIG) and Fundação Arthur Bernardes (FUNARBE) for financial support.

6. References

- Aguilar, M. I.; Romero, M. G.; Chávez, M. I.; King-Díaz, B.; Lotina-Hennsen, B. (2008). Biflavonoids isolated from *Selaginella lepidophylla* inhibit photosynthesis in spinach chloroplasts. *Journal of Agricultural and Food Chemistry*, Vol.56, No.16, (August 2008), pp. 6994-7000, ISSN 0021-8561.
- Allen, J. F. (2003). Cyclic, pseudocyclic and noncyclic photophosphorylation: new links in the chain. *TRENDS in Plant Science*, Vol.8, No.1, (January 2003), pp. 15-19, ISSN 1360-1383.

- Allen, J. F.; Holmes, N. G. (1986). Electron transport partial reactions. In *Photosynthesis, energy transduction. A practical approach*, M. F. Hipkinns, N. R. Baker (Eds), 103-141, Chap. 5., IRL Press, ISBN 09479467519, Oxford, United Kingdom.
- Barbosa, L. C. A.; Demuner, A. J.; de Alvarenga, E. S.; Oliveira, Alberto.; King-Diaz, B.; Lotina-Hennsen, Blas. (2006). Phytogrowth- and photosynthesis-inhibiting properties of nostocclides analogues. *Pest Management Science*, Vol.62, No.3 (March 2006), pp. 214-222, ISSN 1526-498X.
- Barbosa, L. C. A.; Rocha, M. E.; Teixeira, R. R.; Maltha, C. R. A.; Forlani, G. (2007). Synthesis of 3-(4-bromobenzyl)-5-(arylmethylene)-5H-furan-2-ones and their activity as inhibitors of the photosynthetic electron transport chain. *Journal of Agricultural and Food Chemistry*, Vol.55, No.21 (October 2007), pp. 8562-8569, ISSN 0021-8561.
- Barbosa, L. C. A.; Teixeira, R. R.; Montanari, R. M. (2008). Phytotoxic natural products as models for the development of crop protection agents. In: *Current Trends in Phytochemistry*, F. Epifano (Ed.), 21-59, Research Signpost, ISBN 978-81-308-0277-0, Trivandrum, Kerala, India.
- Beckie, H. J. (2006). Herbicide-Resistant Weeds: Management Tactics and Practices. *Weed Technology*, Vol.20, No.3, (July 2006), pp. 793-814, ISSN 0890-037X.
- Berestestskiy, A.; Dmitriev, A.; Mitina, G.; Lisker, I.; Andolfi, A.; Evidente, A. (2008). Nonelides and cytochalasins with phytotoxic activity against *Cirsium arvense* and *Sonchus arvensis*: a structure-activity relationships study. *Phytochemistry*, Vol.69, No.4, (February 2008), pp. 953-960, ISSN 0031-9422.
- Böger, P.; Wakabayashi, K.; Hirai, K. (2002). *Herbicide Classes in Development. Mode of Action, Targets, Genetic Engineering, Chemistry*, Springer-Verlag, ISBN 9783540431473, Berlin, Germany.
- Chaabane, H.; Cooper, J. F.; Azouzi, L.; Coste, C. M. (2005). Influence of soil properties on the adsorption-desorption of sulcotrione and its hydrolysis metabolites on various soils. *Journal of Agricultural and Food Chemistry*, Vol.53, No.10, (April 2005), pp. 4091-4095, ISSN 0021-8561.
- Cobb, A. (1992). *Herbicides and Plant Physiology*, Chapman and Hall, ISBN 0 412 43860 7, London, United Kingdom.
- Copping, L. G.; Duke, S. O. (2007). Natural products that have been used commercially as crop protection agents. *Pest Management Science*, Vol.63, No.6, (June 2007), ISSN 1526-498X.
- Dayan, F. E.; Cantrell, C. L.; Duke, S. O. (2009). Natural products in crop protection. *Bioorganic and Medicinal Chemistry*, Vol.17, No.12, (June 2009), pp. 4022-4034, ISSN 0968-0896.
- Dayan F.; Romagnì, J.; Tellez, M.; Rimando A.; Duke, S. (1999). Managing weeds with natural products. *Pesticide Outlook*, Vol.10, No.5, (October 1999), pp. 185-188, ISSN 0956-1250.
- Demuner, A. J.; Barbosa, L. C. A.; Veiga, T. A. M.; Barreto, R. W.; King-Diaz, B.; Lotina-Hennsen, B. (2006). Phytotoxic constituents from *Nimbya alternantherae*. *Biochemical Systematics and Ecology*, Vol.34, No.11, (November 2006), pp. 790-795, ISSN 0305-1978.

- Devine, M. D.; Shukla, A. (2000) Altered target sites as a mechanism for herbicide resistance. *Crop Protection*, Vol.19, No.8-10, (September 2000), pp. 881-889, ISSN 0261-2194
- Draber, W. (1992). *Rational Approaches to Structure, Activity, and Ecotoxicology of Agrochemicals*. W. Draber; T. Fujita (Eds), p. 278, CRC, ISBN 0849358590, Boca Raton, Florida, United States.
- Duke, S. O.; Dayan, F. E.; Rimando, A. M.; Schrader, K. K.; Aliotta, G.; Oliva, A.; Romagni, J. G. (2002). Chemicals from nature for weed management. *Weed Science*, Vol.50, No.2, (March 2002), pp. 138-151, ISSN 0043-1745.
- Duke, S. O.; Dayan, F. E.; Kagan, I. A.; Baerson, S. R. (2005). New herbicide target sites from natural compounds. *ACS Symposium Series 892 (New discoveries in agrochemicals)*, pp. 151-160, ISSN 0097-6156.
- Duke, S. O.; Dayan, F. E.; Romagni, J. G.; Rimando, A. M. (2000a). Natural products as sources of herbicides: current status and future trends. *Weed research*, Vol.40, No.1, (February 2000), pp. 99-111, ISSN 0043-1737.
- Duke, S. O.; Duke, M. V.; Sherman, T. D.; Nandihalli, U. B. (1991). Spectrophotometric and spectrofluorometric methods in weed science. *Weed Science*, Vol. 39, No.3, pp. 505-13, ISSN 0043-1745.
- Duke, S. O.; Romagni, J. G.; Dayan, F. E. (2000b). Natural products as sources of new mechanisms of herbicidal action. *Crop Protection*, Vol.19, No.8-10, (September 2000), pp. 583-589, ISSN 0261-2194.
- Duke, S. O.; Oliva, A. (2004). In: *Allelopathy – Chemistry and mode of action of allelochemicals*, F. A. Macías, J. C. G. Galindo, J. M. G. Molinillo, H. G. Cuttler (Eds.), 201-216, CRC Press, Boca Raton, ISBN 0-8493-1964-1.
- Giaquinta, R. T.; Dilley, R. A. A partial reaction in photosystem II: reduction of silicomolybdate prior to the site of dichlorophenyldimethylurea inhibition. *Biophysica Acta*, Vol. 387, No.2, pp. 288-305, ISSN 0006-3002.
- Gressel, J. Evolving understanding of the evolution of herbicide resistance. (2009). *Pest Management Science*, Vol.65, No.11, (November, 2009), pp. 1164-1173, ISSN 1526-498X.
- Heap, I. M. (2011). International Survey of Herbicide Resistant Weeds, In: *WeedScience.com*, May 30th 2011, Available from: www.weedscience.org.
- Hüter, O. F. (2011). Use of natural products in the crop protection industry. *Phytochemistry review*, Vol.10, No.2, (June 2011), pp. 185-194, ISSN 1568-7767.
- Jampilek, J.; Musiol, R.; Finster, J.; Pesko, M.; Carroll, J.; Kralova, K.; Vejsova, M.; O'Mahony, J.; Coffey, A.; Dohnal, J.; Polanski, J. (2009b). Investigating biological activity spectrum for novel styrylquinazoline analogues. *Molecules*, Vol.14, No.10, (October 2009), pp. 4246-4265, ISSN 1420-3049.
- Jampilek, J.; Musiol, R.; Pesko, M.; Kralova, K.; Vejsova, M.; Carroll, J.; Coffey, A.; Finster, J.; Tabak, D.; Niedbala, H.; Kozik, V.; Polanski, J.; Csollei, J.; Dohnal, J. (2009a). Ring-substituted 4-hydroxy-1H-quinolin-2-ones: preparation and biological activity. (2009a). *Molecules*, Vol.14, No.3, (March 2009), pp. 1145-1159, ISSN 1420-3049.

- King-Díaz, B.; Castelo-Branco, P. A.; dos Santos, F. J. L.; Rubinger, M. M. M.; Ferreira-Alves, D. L.; Veloso, D. P. (2010). Furanoditerpene Ester and thiocarbonyldioxy derivatives inhibit photosynthesis. *Pesticide Biochemistry and Physiology*, Vol.96, No.3, (March, 2010), pp. 119-126, ISSN 0048-3575.
- King-Díaz, B.; Pérez-Reyes, A.; Santos, F. J. L.; Ferreira-Alves, D. L.; Veloso, D. P.; Carvajal, S. U.; Lotina-Hennsen, B. (2006). Natural diterpene β -lactone derivative as photosystem II inhibitor on spinach chloroplasts. *Pesticide Biochemistry and Physiology*, Vol.84, No.2, (February, 2006), pp. 109-115, ISSN 0048-3575.
- Lein, W.; Börnke, F.; Reindl, A.; Ehrhardt, T.; Stitt, M.; Sonnewald, U. (2004). Target-based discovery of novel herbicides. *Current Opinion in Plant Biology*, Vol.7, No.2, (April 2004), pp. 219-225, ISSN 1369-5266
- Macías, F. A.; Molinillo, J. M. G.; Varela, R. M.; Galindo, J. C. G. (2007). Allelopathy – a natural alternative for weed control. *Pest Management Science*, Vol.63, No.4, (April 2007), pp. 327-348, ISSN 1526-498X.
- Mitchell, G.; Barlett, D. W.; Fraser, T. E. M.; Hawkes, T. R.; Holt, D. C.; Townson, J. K.; Wichert, R. A. (2001). Mesotrione: a new selective herbicide for use in maize. *Pest Management Science*, Vol.57, No.2, (February 2001), pp. 120-128, ISSN 1526-498X.
- Mitchell, P. (1961). Coupling of phosphorylation to electron and proton transfer by a chemiosmotic type of mechanism. *Nature* Vol.19, (July 1961), pp. 144-148, ISSN 0028-0836.
- Morales-Flores, F.; Aguilar, M. I.; King-Díaz, B.; de Santiago-Gómez, J.-R.; Lotina-Hennsen, B. (2007). Natural diterpenes from *Croton ciliatoglanduliferus* as photosystem II and photosystem I inhibitors in spinach chloroplasts. *Photosynthesis Research*, Vol.91, No.1 (January 2007), pp. 71-80, ISSN 0166-8595.
- Musiol, R.; Jampilek, J.; Nycz, J. E.; Pesko, M.; Carroll, J.; Kralova, K.; Vejsova, M.; O'Mahony, J.; Coffey, A.; Mrozek, A.; Polanski, J. (2010). Investigating the activity spectrum for ring-substituted 8-hydroxyquinolines. *Molecules*, Vol.15, No.1 (January 2010), pp. 288-304, ISSN 1420-3049.
- Musiol, R.; Jampilek, J.; Kralova, K.; Richardson, D. R.; Kalinowski, D.; Podeszwa, B.; Finster, J.; Niedbala, H.; Palka, A.; Polanski, J. (2007). Investigating biological activity spectrum for novel quinoline analogues. *Bioorganic and Medicinal Chemistry*, Vol.15, No.3, (February 2007), pp. 1280-1288, ISSN 0968-0896.
- Musiol, R.; Tabak, D.; Niedbala, H.; Podeszwa, B.; Jampilek, J.; Kralova, K.; Dohnal, J.; Finster, J.; Mencil, A.; Polanski, J. (2008). Investigating biological activity spectrum of novel quinoline analogues 2: hydroxyquinolinecarboxamides with photosynthesis-inhibiting activity. *Bioorganic and Medicinal Chemistry*, Vol.16, No.8, (April 2008), pp. 4490-4499, ISSN 0968-0896.
- Nelson, N. (2011). Photosystems and global effects of oxygenic photosynthesis. *Biochimica et Biophysica Acta*, Vol.1807, No.8, (August 2011), pp. 856-863, ISSN 0005-2728.
- Nelson, N. ; Yocum, C. F. Structure and function of photosystems I and II. (2006). *Annual Review of Plant Biology*, Vol.57, (June 2006), 521-565, ISSN 1543-5008.


- Otevrel, J. ; Mandelova, Z. ; Pesko, M. ; Guo, J. ; Kralova, K. ; Sersen, F. ; Vejsova, M. ; Kalinowski, D. S. ; Kovacevic, Z. ; Coffey, A. ; Csollei, J. ; Richardson, D. R. ; Jampilek, J. (2010). Investigating the spectrum of biological activity of ring-substituted salicylanilides and carbamoylphenylcarbamates. *Molecules*, Vol.15, No.11, (November 2010), pp. 8122-8142, ISSN 1420-3049.
- Pillmoor, J. B.; Wright K.; Terry A. S. (1993). Natural products as a source of agrochemicals and leads for chemical synthesis. *Pesticide Science*, Vol.39, No.2, (June 1993), pp. 131-140 ISSN 0031-613X.
- Preston, C. (2004). Herbicide resistance in weeds endowed by enhanced detoxification : complications for management. *Weed Science* Vol.52, No.3 (May 2004), pp. 448-453, ISSN 0043-1745.
- Romagni, J. G. ; Allen, S. N. ; Dayan, F. E. (2000a). Allelopathic effects of volatile cineoles on two weedy plant species. *Journal of Chemical Ecology*, Vol.26, No.1, (January 2000), pp 303-313, ISSN 0098-0331.
- Romagni, J. G. ; Duke, S. O. ; Dayan, F. E. (2000b). Inhibition of plant asparagine synthetase by monoterpenes cineoles. *Plant Physiology*, Vol.123, No.2, (June, 2000), pp. 725-732, ISSN 0032-0889.
- Rüegg, W. T. ; Quadranti M. ; Zoschke, A. Herbicide research and development challenges and opportunities. (2007). *Weed Research*, Vol.47, No.4, (August 2007), pp. 271-275, ISSN 0043-1737.
- Rutherford, A. W. ; Faller, P. (2001). The heart of photosynthesis in glorious 3D. *TRENDS in Biochemical Sciences*, Vol.26, No.6, (June 2001), pp. 341-344, ISSN 0968-0004.
- Saxena, S. ; Pandey, A. K. (2001). Microbial metabolites as eco-friendly agrochemicals for the next millennium. *Applied Microbiology Biotechnology*, Vol.55, No.4, (May 2001), pp. 395-403, ISSN 0175-7598.
- Schrader, K. K. ; Andolfi, A. ; Cantrell, C. L. ; Cimmino, A. ; Duke, S. O. ; Osbrink, W. ; Wedge, D. E ; Evidente, A. (2010). A survey of phytotoxic microbial and plant metabolites as potential products for pest management. *Chemistry and Biodiversity*, Vol.7, No.9, (September 2010), pp. 2261-2280, ISSN 1612-1872.
- Short, P. L. (2005). Growing agrochem r & d. *Chemical and Engineering News*, Vol.8, No.38, (September 2005), pp. 19-22, ISSN 0009-2347.
- Stetter, J. ; Lieb. F. (2000). Innovation in crop protection : trends in research. *Angewandte Chemie International Edition*, Vol.39, No.10, (May 2000), pp. 1725-1744, ISSN 1433-7851.
- Strange, R. N. Phytotoxins produced by microbial plant pathogens. (2007). *Natural Product Reports*, Vol.24, No.1, (January 2007), pp. 127-144, ISSN 0265-0568.
- Souza e Silva, S. R. ; Silva, G. D. F. ; Barbosa, L. C. A.; Duarte, L. P. ; King-Diaz, B. ; Archundia-Camacho, F. ; Lotina-Hennsen, B. (2007). Uncoupling and inhibition properties of 3,4-seco-friedelan-3-oic acid isolated from *Maytenus imbricata*. *Pesticide Biochemistry and Physiology*, Vol.87, No.2 (February 2007), pp. 109-114, ISSN 0048-3575.
- Teixeira, R. R. ; Barbosa, L. C. A. ; Forlani, G. ; Veloso, D. P. ; Carneiro, J. W. M. (2008). Synthesis of photosynthesis-inhibiting nostoclide analogues. *Journal of*

- Agricultural and Food Chemistry*, Vol.56, No.7 (April 2008), pp. 2321-2329, ISSN 0021-8561.
- Teixeira, R. R. ; Pinheiro, P. F. ; Barbosa, L. C. A. ; Carneiro, J. W. M. ; Forlani, G. (2009). QSAR modeling of photosynthesis-inhibiting nostoclide derivatives. *Pest Management Science*, Vol.66, No.2 (February 2009), pp. 196-202, ISSN 1526-498x.
- Torres-Romero, D. ; King-Díaz, B. ; Jiménez, I. A. ; Lotina-Hennsen, B. L. ; Bazzocchi, I. L. (2008). Sesquiterpenes from *Celastrus vulcaniola*. *Journal of Natural Products*, Vol.71, No.8, (August 2008), pp.1331-1335, ISSN 0163-3864.
- Torres-Romero, D. ; King-Díaz, B. ; Strasser, R. J. ; Jiménez, I. A. ; Lotina-Hennsen, B. ; Bazzocchi, I. L. (2010). Friedelane triterpenes from *Celastrus vulcanicola* as photosynthetic inhibitors. *Journal of Agricultural and Food Chemistry*, Vol.58, No.20, (October 2010), pp. 10847-10854, ISSN 0021-8561.
- Trebst, A. (2007). Inhibitors in the functional dissection of the photosynthetic electron transport system. *Photosynthesis research*, Vol.92, No.2 (May 2007), pp. 217-224, ISSN 0166-8595.
- van Klink, J. W. ; Brophy, J. J. ; Perry, N. B. ; Weavers, R. T. (1999). β -triketones from Myrtaceae: Isoleptospermone from *Leptospermum socparium* and papuanone from *Corymbia dallachiana*. *Journal of Natural Products*, Vol. 62, No.3 (February 1999), pp. 487-489, ISSN 0163-3864.
- Veiga, T. A. M. ; González-Vázquez, R. ; Neto, J. O. ; Silva, M. F. G. F. ; King-Díaz, B. ; Lotina-Hennsen, B. (2007b). Siderin from *Toona ciliata* (Meliaceae) as photosystem II inhibitor on spinach thylakoids. *Archives of Biochemistry and Biophysics*, Vol. 465, No.1, (September 2007), pp. 38-43, ISSN 0003-9861.
- Veiga, T. A. M. ; Silva, S. C. ; Francisco, A. C. ; Filho, E. R. ; Vieira, P. C. ; Fernandes, J. B. ; Silva, M. F. G. ; Müller, M. W. ; Lotina-Hennsen, B. (2007a). Inhibition of photophosphorylation and electron transport chain in thylakoids by lasiodiplodin, a natural product from *Botryosphaeria rhodina*. *Journal of Agricultural and Food Chemistry*, Vol.55, No.10, (May 2007), pp. 4217-4221, ISSN 0021-8561.
- Vicentini, C. B. ; Guccione, S. ; Giurato, Laura. Ciacchio, R. ; Mares, D. ; Forlani, G. (2005). Pyrazole derivatives as photosynthetic electron transport inhibitors : new leads and structure-activity relationship. *Journal of Agricultural and Food Chemistry*, Vol.53, No.10, (May 2005), pp. 3848-3855, ISSN 0021-8561.
- Vicentini, C. B. ; Mares, D. ; Tartari, A. ; Manfrini, M. ; Forlani, G. (2004). Synthesis of pyrazole derivatives and their evaluation as photosynthetic electron transport inhibitors. *Journal of Agricultural and Food Chemistry*, Vol.52, No.7, (April 2004), pp. 1898-1906, ISSN 0021-8561.
- Vyvyan, J. R. (2002). Allelochemicals as leads for new herbicides and agrochemicals. *Tetrahedron*, Vol.58, No.9, (February 2002), pp. 1631-1646, ISSN 0040-4020.
- Ware, G. W. (2000). *The Pesticide Book* (5th Edition), Thomson Publications, ISBN 0-913702-63-3, Fresno, California, United States.

Yang, X. ; Shimizu, Y. ; Steiner, J. R. ; Clardy, J. Nostoclides I and II, extracellular metabolites from a symbiotic cyanobacterium *Nostoc* sp., from the lichen *Peltigera canina*. (1993). *Tetrahedron*, Vol. 34, No.5 (January 1993), pp. 761-764, ISSN 0040-4039.

IntechOpen

IntechOpen



Applied Photosynthesis

Edited by Dr Mohammad Najafpour

ISBN 978-953-51-0061-4

Hard cover, 422 pages

Publisher InTech

Published online 02, March, 2012

Published in print edition March, 2012

Photosynthesis is one of the most important reactions on Earth, and it is a scientific field that is intrinsically interdisciplinary, with many research groups examining it. This book is aimed at providing applied aspects of photosynthesis. Different research groups have collected their valuable results from the study of this interesting process. In this book, there are two sections: Fundamental and Applied aspects. All sections have been written by experts in their fields. The book chapters present different and new subjects, from photosynthetic inhibitors, to interaction between flowering initiation and photosynthesis.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Róbson Ricardo Teixeira, Jorge Luiz Pereira and Wagner Luiz Pereira (2012). Photosynthetic Inhibitors, Applied Photosynthesis, Dr Mohammad Najafpour (Ed.), ISBN: 978-953-51-0061-4, InTech, Available from: <http://www.intechopen.com/books/applied-photosynthesis/photosynthetic-inhibitors>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](https://creativecommons.org/licenses/by/3.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen