

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

186,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Molecular Breeding of Grasses by Transgenic Approaches for Biofuel Production

Wataru Takahashi and Tadashi Takamizo

*Forage Crop Research Division, NARO Institute of Livestock and Grassland Science
Japan*

1. Introduction

Since the Industrial Revolution, fossil fuels have been consumed in massive amounts. However, little fossil fuels are estimated to remain, and the number of new oil fields in the world continues to decline. Thus, the future supply of fossil fuels will be tight and will not continue to support our levels of consumption for long. In addition, rapid consumption of fossil fuels for energy production continues to elevate the atmospheric carbon dioxide concentration, which was 280 ppm before the Industrial Revolution and is 380 ppm today. The emissions, including carbon dioxide, derived from the combustion of fossil fuels act as greenhouse gases and cause the serious problem of global warming.

In this context, society must develop beyond the consumption of our limited fossil fuels. Plant biomass, which is organic matter derived from the photosynthetic fixation of atmospheric carbon dioxide, holds much promise as a future energy source. The use of plant biomass for energy is called carbon neutral, because the combustion of plant biomass only releases carbon dioxide that was originally fixed from the atmosphere by plants. Because a sustainable society based on carbon-neutral energy must be constructed to preserve the global environment, we must develop related practical technology as soon as possible.

However, recent hikes in the global price of grain, caused by various political and economic events, has made us consider shifting toward using non-grain plant biomass to improve both energy and food-supply security. Thus, research into the production of biofuel from lignocellulose, which constitutes the bodies of woody and grass plant species, is a major field of study in modern plant science.

Needless to say, this research includes biotechnological approaches. In particular, genetic engineering is expected to be a key technology in herbaceous plant breeding, because most of the bioenergy crop species are not yet domesticated well. This chapter focuses on recent progress in the molecular breeding of herbaceous plants.

2. Herbaceous perennials for dedicated lignocellulosic biofuel crops

High biomass yields are indispensable to successful cultivate energy crops that can compete with and replace fossil fuels (Karpenstein-Machan, 2001; Lewandowski et al., 2003). Considering food-supply security, net energy yield, and environmental protection,

large and fast-growing non-food grass species are among the most promising candidates for lignocellulose production. Perennial species are more attractive than annuals because they have minimal input requirements, commonly yield more aboveground biomass in the form of stems and leaves, and their well-developed root systems can serve as carbon sinks.

Name (Scientific name) ^a	Photosystem	Yield (t DM ha ⁻¹ yr ⁻¹) ^b	Location	Citation
Switchgrass (<i>Panicum virgatum</i> L.)	C4	15.9 [†]	U.S.A. (Oklahoma)	Aravindhakshan et al. (2010)
Sugarcane (<i>Saccharum</i> spp. hybrids)	C4	41.3 ^c	Japan (South western islands)	Terajima et al. (2010)
<i>Miscanthus</i> species (<i>Miscanthus</i> × <i>giganteus</i>)	C4	28.7 [†]	Central Italy (Pisa)	Angelini et al. (2009)
		12.4 [†]	U.S.A. (Oklahoma)	Aravindhakshan et al. (2010)
<i>Erianthus</i> species (<i>Erianthus</i> <i>arundinaceus</i> Retz.)	C4	58.4	Japan (Northern Kanto region)	Ando et al. (2011)
Giant reed (<i>Arundo donax</i> L.)	C3	37.7 [†]	Central Italy (Pisa)	Angelini et al. (2009)

^a Actual scientific names of experimental cultivars or lines used in the study cited are given for *Erianthus* species, *Miscanthus* species, and sugarcane.

^b Values followed by the same obelisk were obtained by the same study.

^c An average value calculated with the data from three different locations.

Table 1. Summary information for perennial grasses described in this review (modified and updated from Lewandowski et al., 2003)

C4 perennial grasses show promise as lignocellulosic biofuel crops because their highly efficient C4 photosynthesis often yields more biomass than C3 grasses (Jakob et al., 2009). There are many available candidate perennial grasses, which differ in their potential productivity, chemical and physical biomass properties, environmental demands and crop management requirements (Lewandowski et al., 2003). This review will focus on noteworthy C4 perennial grasses as dedicated lignocellulosic biofuel crops, including switchgrass (*Panicum virgatum* L.), *Miscanthus* species, *Erianthus* species, and sugarcane (*Saccharum* species). In addition, the high biomass C3 grass species giant reed (*Arundo donax* L.), which has a photosynthetic capacity comparable to or higher than that of C4 grasses (Angelini et al., 2009; Rossa et al., 1998), will be discussed as well. These perennial grasses are summarized in Table 1 and will be briefly described in sections 2.1 through 2.3.

2.1 Switchgrass

Switchgrass (*Panicum virgatum* L.) is an outcrossing perennial warm-season forage grass originally from the prairies of North America. In the 1990’s, the United States Department of Energy sponsored a 10-year research project to evaluate and develop this grass as a dedicated herbaceous lignocellulosic energy crop because of its potential for high fuel

yields, environmental enhancement, and ability to grow well on marginal cropland without heavy fertilizing or intensive management (Bouton, 2008; McLaughlin & Kszos, 2005; Wright & Turhollow, 2010).

Switchgrasses are divided into two ecotypes: lowland and upland (Bouton, 2008; Lewandowski et al., 2003). The lowland ecotypes are tall and robust plants adapted to wetter sites; they mature later and have longer growth periods and higher biomass yields than upland ecotypes. Although somatic chromosome numbers vary in switchgrass ($2n$ from $2x = 18$ to $12x = 108$) (Hopkins et al., 1996), the lowland ecotypes are predominantly tetraploid ($2n = 4x = 36$) (Bouton, 2008; Lewandowski et al., 2003). The upland ecotypes have shorter and fine-stemmed morphology, and are adapted to drier habitats. They are commonly octoploid ($2n = 8x = 72$), or occasionally tetraploid ($2n = 4x = 36$) (Bouton, 2008) or hexaploid ($2n = 6x = 54$) (Lewandowski et al., 2003). Irrespective of ecotype, switchgrasses with the same ploidy level can be intercrossed (Lewandowski et al., 2003).

2.2 *Saccharum* complex

The term “*Saccharum* complex” was first coined by Mukherjee (1957) and originally encompassed four closely-related interfertile genera, *Saccharum*, *Erianthus*, *Sclerostachya*, and *Narenga*. Based on species richness and the geographic distributions of endemic species, India was considered the center of maximum variation of the *Saccharum* complex (Mukherjee, 1957). Eventually, the genus *Miscanthus* was added to the *Saccharum* complex, because it was thought to be involved in the origin of *Saccharum* (Daniels et al., 1975, as cited in Alwala et al., 2006; and Amalraj & Balasundaram, 2006).

Modern sugarcane varieties are mostly derived from interspecific hybridization within the genus *Saccharum* (Amalraj & Balasundaram, 2006), and intergeneric hybridization between *Saccharum* and other genera in the *Saccharum* complex is thought to be the primary gene pool for sugarcane breeding (Cheavegatti-Gianotto et al., 2011). Thus, the *Saccharum* complex can also be considered as the primary gene pool for breeding non-domesticated *Miscanthus* and *Erianthus* as well.

In sections 2.2.1 through 2.2.3, we will address individual genera within the *Saccharum* complex. However, a comprehensive molecular breeding system with intercrossing across the complex should ultimately be undertaken in the development of novel hybrid biofuel crops.

2.2.1 Sugarcane (*Saccharum* species)

Sugarcane is a tall perennial C4 grass that is cultivated in tropical and subtropical regions of the world. Notably, this grass stores high concentrations of sucrose in the stem. Approximately, 65 to 70% of global sugar production in the form of sucrose is derived from sugarcane (FAO, 2003). The potential of sugarcane as an important energy crop was argued because of the advent of large-scale sugarcane-based ethanol production in Brazil (Tew & Cobill, 2008).

Sugarcane belongs to the genus *Saccharum*. Although six polyploid species are recognized within *Saccharum* (Table 2), modern cultivars for sugar production are mostly derived from interspecific hybridization between *S. officinarum* ($2n = 8x = 80$) and *S. spontaneum* ($2n$ from $5x = 40$ to $16x = 128$) and are thus complex polyploids with variable chromosome numbers

(2n approx. from 100 to 120) (Henry, 2010; Piperidis et al., 2010). Of the other four species, *S. robustum*, *S. barberi*, and *S. sinense* have also provided minor contributions to the breeding of some modern sugarcane cultivars (Cheavegatti-Gianotto et al., 2011).

Species	Chromosome number (2n)	Genomic contribution to modern interspecific hybrid cultivars and elite lines (%) ^a	Note
<i>S. barberi</i>	From 111 to 120	-	Semi-sweet Indian cane
<i>S. edule</i>	From 60 to 80	-	Cultivated for edible inflorescence
<i>S. officinarum</i>	80	From 70 to 80	Domesticated sweet cane
<i>S. robustum</i>	60 or 80	-	Putative ancestor of <i>S. officinarum</i>
<i>S. sinense</i>	From 81 to 124	-	Semi-sweet Chinese cane
<i>S. spontaneum</i>	From 40 to 128	From 10 to 23	Wild cane found throughout Asia

^a Values are the proportions of total chromosome complement reported by Piperidis et al. (2010) where the proportion of chromosomes derived from interspecific exchanges between *S. officinarum* and *S. spontaneum* are shown to be 8-13%.

Table 2. Species of *Saccharum* and their characteristics (modified from Henry, 2010)

The breeding of sugarcane as a dedicated biomass crop called "energy cane" can be categorized into three strategies with different objectives: the “sugar model”, the “sugar-and-fiber model”, and the “fiber-only model”. The fiber yield of energy cane is important because of its use for electricity generation, cellulosic ethanol production, and so forth; fiber is considered a by-product in the sugar and sugar-and-fiber models or the main product in the fiber-only model (Tew & Cobill, 2008). In the sugar model, improved sugar yield and sugar content are the main foci, so traditional sugarcane cultivars can be used. In the sugar-and-fiber and fiber-only models, Type I and Type II energy canes, respectively, would be used. Tew & Cobill (2008) defined Type I and Type II energy canes as follows:

- Type I energy cane is bred and cultivated to maximize both its sugar and fiber content.
- Type II energy cane is bred and cultivated primarily or solely for its fiber content.

Recently, Japanese breeders succeeded in developing a high-quality Type I energy cane cultivar ‘KY01-2044’ with 1.5 times the total biomass yield and 1.3 times the total sugar yield than the major Japanese sugar-producing cultivar (Asia Biomass Office, 2010; Terajima et al., 2010). The new cultivar allowed the establishment of an experimental system for simultaneous production of ethanol and sucrose from total sugar with residual fiber as a heat source. In addition to ethanol production, this system is designed to produce an amount of sucrose comparable to conventional sugar production systems.

2.2.2 *Miscanthus* species

Miscanthus is a genus of C4 perennial rhizomatous grasses widely distributed in Asia and the Pacific Islands. *Miscanthus* was thought to consist of 17 species divided into four sections (Deuter, 2000). However, recent taxonomic analyses using molecular markers revealed that the genus can be reduced to approximately 11-12 species. The species *M. sinensis* ssp. *condensatus* is sometimes recognized at specific rank as *M. condensatus* (Clifton-Brown et al., 2008) (Table 3).

Species	Chromosome number and ploidy level ^a
<i>M. floridulus</i> (Labill.) Warb.	2n = 2x = 38
<i>M. intermedius</i> (Honda) Honda	2n = 6x = 114
<i>M. longiberbis</i> Nakai	-
<i>M. lutarioparius</i>	-
<i>M. oligostachyus</i> Stapf.	2n = 2x = 38
<i>M. paniculatus</i> (B. S. Sun) Renvoize & S. L. Chen	-
<i>M. sacchariflorus</i> (Maxim.) Hack.	2n = 2x = 38 (in China) 2n = 4x = 76 (in Japan)
<i>M. sinensis</i> Anderss.	2n = 2x = 38
<i>M. sinensis</i> ssp. <i>condensatus</i> (Hackel) T. Koyama	2n = 2x = 38
<i>M. tinctorius</i> (Steud.) Hack.	2n = 2x = 38
<i>M. transmorrisonensis</i> Hayata	2n = 2x = 38
<i>M. × giganteus</i> Greef & Deuter ex Hodkinson and Renvoize (Hybrid species between <i>M. sacchariflorus</i> and <i>M. sinensis</i>)	2n = 3x = 57

^a Major cytotypes from Deuter (2000) are shown.

Table 3. Species in the genus *Miscanthus* and their chromosome numbers

The basic chromosome number of *Miscanthus* species is 19, and polyploids and aneuploids are observed (Deuter, 2000). Of the species in Table 3, *M. sacchariflorus*, *M. sinensis*, *M. sinensis* ssp. *condensatus*, *M. floridulus*, and *M. × giganteus* are of interest for biomass production (Deuter, 2000). In particular, the triploid hybrid species *M. × giganteus* shows superior characteristics, such as high biomass yield, and is thus thought to be the most practical *Miscanthus* species for bioenergy production, especially in Europe and North America (Lewandowski et al., 2000; Pyter et al., 2007). The existence of this promising hybrid will further encourage interspecific hybridization within *Miscanthus*, because the genus has a lot of genetic diversity within and between species. In addition, the frost tolerance, growth at low temperature, and robustness against pests and diseases of *Miscanthus* make it a potential gene pool for developing widely-adaptable stress-tolerant cultivars of sugarcane through intergeneric hybridization (Clifton-Brown et al., 2008).

2.2.3 *Erianthus* species

Erianthus is a tall C4 perennial rhizomatous grass. The genus *Erianthus* was erected by André Michaux in *Flora Boreali-Americana* in 1803 for the New World species *E. saccharoides* Michaux (Tagane et al., 2011). Old World species have distinct morphology from New World species (Grassl, 1972, as cited in Tagane et al., 2011) and are characterized by the

presence of a distinctive luteolin, di-C-glycoside, that is not present in the New World species (Williams et al., 1974). The genus is widely distributed in the Americas (New World species) and in the Mediterranean, India, China, South East Asia, and New Guinea (Old World species) (Amalraj & Balasundaram, 2006).

The seven Old World species comprise the section *Ripidium* (Table 4). The basic chromosome number of these *Erianthus* species is $x = 10$, the same as *Saccharum officinarum*. Polyploids are also observed, as in other genera in the *Saccharum* complex.

Species	Chromosome number ^a
<i>E. arundinaceus</i> (Retz.) Jesw.	2n = 30, 40, 60
<i>E. bengalense</i> (Retz.)	2n = 20, 30, 40, 60
<i>E. elephantinus</i> Hook. f.	2n = 20
<i>E. hostii</i> Griseb.	2n = 20
<i>E. kanashiroi</i> Ohwi	2n = 60
<i>E. procerus</i> (Roxb.) Raizada	2n = 40
<i>E. ravennae</i> (L.)	2n = 20

^a Chromosome numbers shown by Amalraj & Balasundaram (2006) are shown.

Table 4. Species in the genus *Erianthus* sect. *Ripidium* and their chromosome numbers

The Old World species are increasingly used in sugarcane breeding because of their high biomass yields, drought tolerance, and resistance to pests and diseases. Two species in particular, *E. arundinaceus* and *E. procerus*, are considered most useful due to their disease resistance, ratooning ability, vigor, and environmental stress tolerance (Tagane et al., 2011). Recently, Ando et al. (2011) reported the wintering ability and high biomass yields of the *E. arundinaceus* clone JW630 grown experimentally in Nasushiobara, Japan, where the monthly mean minimum air temperatures from December to March are below freezing, indicating its potential as breeding stock for cold-tolerant sugarcane and as a dedicated lignocellulosic biofuel crop (Table 1).

2.3 Giant reed

Giant reed (*Arundo donax* L.) is a rhizomatous C3 perennial grass that evolved in Asia but is also considered a native to the Mediterranean region (Lewandowski et al., 2003). The grass is used to make reeds for woodwind musical instruments, in construction, and as a source of cellulose for rayon manufacture and paper pulp production (Perdue, 1958).

Recently, giant reed has gained attention as a source of lignocellulose for bioenergy production because of its high yield and strong pest resistance (Lewandowski et al., 2003). Studies of the giant reed in Central Italy demonstrated biomass production equivalent to or higher than *Miscanthus × giganteus* (Angelini et al., 2009) (Table 1). Because giant reed exhibits heavy metal tolerance, it could be used simultaneously for both phytoremediation and bioenergy production (Papazoglou, 2007; Papazoglou et al., 2005).

Giant reed is highly polyploid, with a possible base chromosome number $x = 12$ [(2n = 10x = 120 - 8 or 2n = 9x = 108 + 4) (Gorenflot et al., 1972) or (2n = 110) (Lewandowski et al., 2003)]. Although isozyme and DNA analyses have revealed genetic diversity in giant reed (Khudamrongsawat et al., 2004; Lewandowski et al., 2003), the plant is thought to produce

no viable seed, owing to aberrant division of the megaspore mother cell (Bhanwra et al., 1982). This behavior suggests that conventional breeding through sexual hybridization cannot be performed.

3. Plant regeneration systems

In most case, effective plant regeneration systems with *in vitro* cell cultures are required to establish reliable transformation systems in plants. In this section, we summarize the regeneration systems of switchgrass, sugarcane, *Miscanthus*, *Erianthus*, and giant reed.

3.1 Switchgrass regeneration systems

Biotechnology research on the molecular breeding of switchgrass began in 1992 at the University of Tennessee with the support of Oak Ridge National Laboratory (Vogel & Jung, 2001). This work yielded the first reported plant regeneration system through embryogenesis and organogenesis (Denchev & Conger, 1994). In the experiments, mature caryopses and young leaf segments were used as explants for callus induction. Then, the influences of the synthetic auxins 2,4-dichlorophenoxyacetic acid (2,4-D) and 4-amino-3,5,6-trichloropyridine-2-carboxylic acid (picloram) in combination with 6-benzyladenine (BA) were examined on the plant regeneration system. The combination of 2,4-D and BA led to best results in experiments with mature caryopses (Denchev & Conger, 1995). Alexandrova et al. (1996a) increased the efficiency of switchgrass regeneration by using immature inflorescences obtained from node cultures in aseptic conditions; one immature inflorescence can produce hundreds of spikelets of the same genotype in a single Petri dish that can be easily used as explants for callus induction. Most recently, a new medium, LP9, for the production, maintenance, and regeneration of switchgrass callus was reported (Burris et al., 2009). Callus produced on LP9 can be easily propagated, maintains its regenerability for at least six months, and is adaptable to *Agrobacterium*-mediated transformation (Burris et al., 2009).

Embryogenic suspension cultures initiated from embryogenic callus were also established in switchgrass (Gupta & Conger, 1999), and osmotic pre-treatment with 0.3 M each of sorbitol and mannitol was effective for plant regeneration from suspension cultures (Odjakova & Conger, 1999). Another type of culture system that is highly attractive for gene transfer experiments was also established: multiple shoot clumps were induced from intact seedlings with various combinations of 2,4-D and 1-phenyl-3-(1,2,3-thiadiazol-5-yl) urea (TDZ) (Gupta & Conger, 1998).

Most of the research mentioned above used the lowland cultivar 'Alamo'. However, because switchgrass is outcrossing and self-infertile, genetic variation exists not only among cultivars but also within a cultivar, indicating that the establishment of a totipotent tissue culture is not highly dependent on the cultivar itself but rather on screening highly regenerable genotypes as in other outcrossing plants, like ryegrasses (Takahashi et al., 2004). Therefore, Odjakova & Conger (1999) and Burris et al. (2009) used specific single genotypes Alamo 2702 and Alamo 2, respectively, in their experiments, and Alexandrova et al. (1996b) developed a micropropagation procedure with node culture for the multiplication of such selected genotypes. Interestingly, the selected line HR8, having high somatic embryogenic capacity, was recently bred using recurrent tissue culture selection to allow easy induction of embryogenic callus from mature seeds (Xu et al., 2011).

3.2 Sugarcane regeneration systems

The first sugarcane regeneration system via callus culture was reported in 1964 (Nickell, 1964, as cited in Lakshmanan, 2006) and was followed by numerous reports resulting from a great deal of research activity. These early reports have been well-reviewed elsewhere (Lakshmanan, 2006; Suprasanna et al., 2008a). Although many papers on plant regeneration from sugarcane calli are still being published, they generally vary only in reporting specific culture conditions, which must be optimized for each cultivar and genotype because of the outcrossing nature of sugarcane. Most callus cultures are supplemented with the auxin 2,4-D, but sometimes other synthetic auxins, such as picloram and 3,6-dichloro-2-methoxybenzoic acid (dicamba) are used, with or without the cytokinins BA, TDZ, or kinetin. Research on sugarcane regeneration systems may be stalled. To spark innovation, here we update the previous reviews (Lakshmanan, 2006; Suprasanna et al., 2008a) by describing innovative reports published since 2007.

In addition to plant hormones, exogenous amino acids may positively or negatively influence somatic embryogenesis in plants. In sugarcane tissue culture, the addition of arginine to culture media was found to significantly induce somatic embryogenesis and to promote plant regeneration (Nieves et al., 2008). Similarly, glycine, arginine, and cysteine positively affected somatic embryogenesis and subsequent plant regeneration; 0.75 mM glycine was the most effective treatment studied (Asad et al., 2009).

The moisture status of *in vitro* cultures influences plant regeneration in sugarcane. Partial desiccation has been reported to enhance plant regeneration from calli (Garcia et al., 2007; Kaur & Gosal, 2009). A similar phenomenon was observed even in irradiated embryogenic cultures (Suprasanna et al., 2008b).

A fundamental but important aspect of sugarcane tissue culture is the selection of the initial explants. In most previous reports, explants were obtained from field-grown plants. However, Garcia et al. (2007) recommended using sugarcane plants grown *in vitro* because they provided a year-round source of physiologically uniform explants and could be prevented from releasing large amounts of phenolic compounds, which can hamper tissue culture (Garcia et al., 2007). Basnayake et al. (2011) recommended using explants obtained from donor plants with good water supplies, especially when working with genotypes that are recalcitrant to tissue culture due to phenolic compounds.

3.3 *Miscanthus* regeneration systems

To the best of our knowledge, the first preliminary regeneration experiments in *Miscanthus* were conducted with immature inflorescences, mature leaves, immature leaves, nodal segments, internodal segments, meristematic regions, and ovules as explants, which were cultured on media containing a wide range of auxin types (Gawel et al., 1987, as cited in Gawel et al., 1990). In these experiments, immature inflorescences were the only explants that produced calli, and only media containing 2,4-D or picloram with no cytokinin produced a regeneration response (Gawel et al., 1987, as cited in Gawel et al., 1990). The research group thus focused on immature inflorescences as explants and published that 9.0 μ M of 2,4-D was effective for callus induction and subsequent plant regeneration in *M. sinensis* cultivars 'Gracillimus', 'Variegatus', and 'Zebrinus' (Gawel et al., 1990). In these experiments, callus induction and plant regeneration were not distinguished from one

another; shoot formation occurred during successive culture on the same culture medium as callus induction (Gawel et al., 1990).

Many researchers have focused on the high-biomass species *M. × giganteus*. The auxins 2,4-D or 2,4,5-trichlorophenoxyacetic acid in combination with the cytokinin 6-furfurylaminopurine were best for inducing regenerable embryogenic callus from immature inflorescences (Lewandowski & Kahnt, 1992). Although immature inflorescences emit fewer browning substances than shoot tips or leaf primordia (Lewandowski & Kahnt, 1993), browning substances from the explants were detrimental to tissue culture (Lewandowski & Kahnt, 1992). Ascorbic acid, cysteine, and watering the explants were not effective treatments, but liquid culture gave better results because the browning substances did not accumulate around the explants or calli (Lewandowski & Kahnt, 1992).

The effects of different explants, such as shoot apices, leaves and root sections of *in vitro*-propagated plants and leaf and immature inflorescence sections from greenhouse-grown plants, were examined in *M. × giganteus*. Shoot apices had the highest percentage of embryogenic callus formation, while immature inflorescence-derived callus had the highest regenerability (Holme & Petersen, 1996). The growth conditions of donor plants influence tissue culture (Creemers-Molenaar et al., 1988), as has been observed in *M. × giganteus* (Holme & Petersen, 1996). Leaf explants from *in vitro*-propagated shoots and from greenhouse-grown plants showed differences in embryogenic callus formation; the best results were obtained from leaves of *in vitro*-propagated shoots and older leaves of greenhouse-grown plants (Holme & Petersen, 1996). However, supplying BA to the callus induction medium led to different results: a higher percentage of regenerable shoot-forming callus was formed on shoot apices compared with leaf sections of *in vitro*-grown shoots when 0.4 μM BA was supplied (Petersen, 1997). Also, small immature inflorescences of *M. × giganteus*, between 2.5 and 8 mm, were more suitable for embryogenic callus formation than explants from shorter or longer inflorescences, shoot apices or leaf explants, indicating that the size and type of explant influence culture responses (Petersen et al., 1999). Furthermore, different carbon sources, and their sterilization methods, influenced *M. × giganteus* tissue culture. Significant differences were reported for carbon sources and their sterilization methods in tissue cultures of various explant-derived calli. Leaf explants were more affected by the carbon sources than were shoot apices or immature inflorescences, and both callus proliferation and plant regeneration were generally improved by the use of filter-sterilized carbon sources (Petersen et al., 1999).

As mentioned above, severe browning is a major problem in *Miscanthus* tissue culture, but supplying proline to callus induction and suspension cultures effectively prevented browning in *M. × giganteus* (Holme et al., 1997). Proline is thought to inhibit polyphenol oxidase, which causes enzymatic browning of cultured tissues (Öztürk & Demir, 2002). The addition of proline to callus induction media increased embryogenic callus formation on shoot apices and leaf explants. However, results varied with proline concentration and the basal salts in the medium. Specifically, 12.5 to 50 mM proline in callus induction media with Murashige & Skoog (MS) salts (Murashige & Skoog, 1962) increased embryogenic callus formation more than media with N6 salts (Chu et al., 1975). The inhibitory effects of proline on tissue browning in *Miscanthus* were confirmed by another group (Głowacka et al., 2010). Conversely, adding honey instead of sucrose to callus induction media inhibited browning of cultured immature inflorescences, and a combination of the honey and banana pulp was best for inducing regenerable callus in *M. × giganteus* (Płażek & Dubert, 2009, 2010).

Media component (mg L ⁻¹ final conc.)	Media		
	HB ^a	MS	N6
Macroelements			
KNO ₃	2000	1900	2830
NH ₄ NO ₃	1500	1650	
(NH ₄) ₂ SO ₄			463
CaCl ₂ •2H ₂ O	200	440	166
MgSO ₄ •7H ₂ O	300	370	185
KH ₂ PO ₄	300	170	400
Microelements			
Na ₂ -EDTA	37.25	37.3	37.25
FeSO ₄ •7H ₂ O	27.85	27.8	27.85
MnSO ₄ •4H ₂ O	11.25	22.3	4.4
ZnSO ₄ •7H ₂ O	4.3	8.6	1.5
CuSO ₄ •5H ₂ O	0.0125	0.025	
CoCl ₂ •6H ₂ O	0.0125	0.025	
H ₃ BO ₃	3.1	6.2	1.6
KI	0.415	0.83	0.8
Vitamins			
Inositol	100	100	
Nicotinic acid	0.5	0.5	0.5
Pyridoxine HCl	0.5	0.5	0.5
Thiamine HCl	0.4	0.1	1
Another organic			
Glycine	2	2	2

^a Composition presented in Sun et al. (1999).

Table 5. Composition of HB, MS, and N6 basal media

Recently, studies involving tissue cultures of *M. sinensis* have been reported. Głowacka & Jeżowski (2009a, 2009b) reported culturing anthers of *M. sinensis* and subsequently demonstrated the effects of inflorescence developmental stage on callus induction and plant regeneration (Głowacka et al., 2010). Mature seeds can also serve as explants for tissue culture in *M. sinensis*. Mature seeds of 18 accessions from various sites in Japan were subjected to callus induction, and a combination of a relatively high 2,4-D concentration (5 mg L⁻¹) and a relatively low BA concentration (0.01 mg L⁻¹) efficiently induced regenerable calli that could be used to produce transgenic *Miscanthus* plants (Wang et al., 2011). Interestingly, there was a correlation between the average annual air temperature at accession collecting sites and the frequency of embryogenic callus induction; seeds collected from warmer regions formed a higher percentage of embryogenic calli (Wang et al., 2011). Most recently, Zhang et al. (2011) germinated seeds collected in China and compared explants from the epicotyls, young leaves, and radicles. The epicotyl was best for embryogenic callus formation and plant regeneration from the callus. In the experiments, Holley & Baker (HB) medium (Holley & Baker, 1963, as cited in Zhang et al., 2011) was better for callus induction than other common basal media, such as MS medium, N6 medium (Chu et al., 1975), or half strength of the MS medium. Zhang et al. surmised that HB medium would work well for callus induction in other grass species, because the medium had been effective for callus culture of wheat (Sun et al., 1999). The

components of HB medium will be of interest to many researchers, so we compare HB, MS, and N6 media in Table 5. We were unable to obtain the original publication (Holley & Baker, 1963, as reported in Zhang et al., 2011), so we give the composition of HB basal medium presented in Sun et al. (1999).

3.4 *Erianthus* regeneration systems

To date, only one published report concerns the plant regeneration system of *Erianthus* species. Callus induction and plant regeneration from calli was achieved in *E. elephantinus* (Jalaja & Sreenivasan, 1999). Calli were induced from expanding leaves, leaf sheaths, and immature inflorescences as explants on MS basal medium containing 2 mg L⁻¹ 2,4-D. As in *Miscanthus* species, tissue browning occurred, especially in explants from leaf sheaths (Jalaja & Sreenivasan, 1999). Unfortunately, detailed data on the frequencies of callus induction and plan regeneration from the callus were not reported because the study focused mainly on somaclonal variation, such as morphology, pollen fertility, and chromosome aberrations, of the regenerants (Jalaja & Sreenivasan, 1999).

3.4.1 Protocol for plant regeneration system of *Erianthus arundinaceus*

We have recently succeeded in establishing a plant regeneration system in *E. arundinaceus*. Here, we present the simplified protocol. All culture media used were based on MS medium containing 3% (w v⁻¹) sucrose, adjusted to pH 5.8, and solidified with 0.25% (w v⁻¹) Gelrite (Wako, Osaka, Japan). Components of the culture media are shown in Table 6.

Media	Components ^a
Plant maintenance medium	0.3% (w v ⁻¹) activated charcoal 3%(w v ⁻¹) sucrose
Callus induction medium	5 mg L ⁻¹ 2,4-D 25 mM L-proline 750 mg ⁻¹ MgCl ₂ • 6H ₂ O 3%(w v ⁻¹) sucrose
Multiple shoot formation medium	1 mg L ⁻¹ BA 3 mg L ⁻¹ NAA 3%(w v ⁻¹) sucrose
Shoot elongation medium	0.3 mg L ⁻¹ GA 3%(w v ⁻¹) sucrose

^a Additives added to MS basal medium are shown for each medium.

Table 6. Culture media used in our plant regeneration system for *Erianthus arundinaceus*

- Establishment of *in vitro*-grown plants

We obtained *in vitro*-grown plants of the *E. arundinaceus* clone JW630 (Ando et al., 2011). Shoot tips, each containing an apical meristem, of greenhouse-grown plants (approx. 3 cm long) were used as explant sources. After removing the mature outer leaves, the shoot tips were submerged in 70% ethanol for 1 min, surface-sterilized in 10% (v v⁻¹) sodium hypochlorite solution (1% available chlorine) for 20 min, rinsed twice in sterile distilled

water under aseptic conditions, aseptically stripped of more outer leaves, and cultured on plant maintenance medium under long-day conditions [16 h light ($80 \mu\text{mol m}^{-2} \text{s}^{-1}$)/8 h dark; 28°C]. The explants were subcultured every two weeks until rooting.

- Callus induction from axillary buds of *in vitro*-grown plants

Axillary buds at the proximal end of each leaf were isolated from *in vitro*-grown plants under a stereomicroscope. The isolated buds were placed on callus induction medium, and cultured under continuous fluorescent light ($40 \mu\text{mol m}^{-2} \text{s}^{-1}$) at 25°C . They were subcultured every two weeks.

- Plant regeneration from calli

For plant regeneration, calli were transferred to the plant maintenance medium and cultured under long-day conditions [16 h light ($80 \mu\text{mol m}^{-2} \text{s}^{-1}$)/8 h dark; 28°C].

- Results and tips

As with other C_4 plants, such as switchgrass, sugarcane, and *Miscanthus*, explant browning was severe at first. Rooting of the explants took about two months, but once rooted, the *in vitro* plants grew vigorously on the plant maintenance medium (Figure 1A). Calli were easily induced from axillary buds of *in vitro*-grown plants (Figure 1B), and plant regeneration was observed from the calli (Figure 1C). However, calli induced from explants from field-grown plants were rare due to severe bacterial contamination and tissue browning, even when we used apical meristems, immature inflorescences, and young leaf rolls (data not shown). The *in vitro*-grown plants produced multiple shoot clumps when cultured on multiple shoot formation medium (Figure 2A), and the newly formed short shoots maintained their morphological states and could elongate to normal morphology after being transferred to shoot elongation medium containing gibberellic acid (GA) (Figure 2B).



Fig. 1. Plant regeneration system for *Erianthus arundinaceus*. An *in vitro*-grown plant cultured for three months (A), a callus induced from an axially bud of the *in vitro*-grown plants (B), and plant regeneration from calli (C).




Fig. 2. Multiple shoot formation in *Erianthus arundinaceus*. A multiple shoot clump derived from an *in vitro*-grown plant (A), and elongated shoots from the multiple shoot clumps (B).

3.5 Giant reed regeneration system

To the best of our knowledge, only one scientific article on the regeneration of giant reed has been published so far (Takahashi et al., 2010). In the system, *in vitro* propagation was first optimized for the year-round production of explants (Figure 3).




Fig. 3. An *in vitro*-grown giant reed plant. The photo was taken two months after subculture (adapted from Takahashi et al., 2010).

Calli were induced from axillary buds isolated from the *in vitro*-grown plants. Several combinations of 2,4-D and BA were used for callus induction. They influenced both callus induction frequency and later plant regeneration on hormone-free media because of possible carryover effects of plant hormones in the original callus induction media. Media supplemented with 3 mg L⁻¹ 2,4-D appeared to stabilize callus formation and subsequent shoot formation on hormone-free medium (Figure 4).




Fig. 4. Effect of original callus induction media on subsequent shoot formation on hormone-free media in giant reed. The concentrations of 2,4-D and BA are those in original callus induction media (modified and adapted from Takahashi et al., 2010).

In total, 11 genotypes were treated under optimized conditions, and data suggest genotypic effects in tissue culture response, although the giant reed is thought to propagate asexually because its seeds are non-viable. Ex-callus induction frequencies and shoot formation frequencies ranged from 82.9 to 100% and from 0 to 37.5%, respectively.

4. Transformation systems

We summarize here several earlier reports on transformation systems in switchgrass, sugarcane, *Miscanthus*, and giant reed. We know of no such reports for *Erianthus*.

4.1 Switchgrass transformation systems

There are some reports on the establishment of transformation systems in switchgrass. Gene transfer methods and names of transgenes used so far are shown in Table 7.

Transgenes ^a	Gene transfer methods	Reference
<i>sgfp</i> , [<i>bar</i>]	Gene gun	Richards et al. (2001)
<i>uidA</i> , [<i>bar</i>]	<i>Agrobacterium</i>	Somleva et al. (2002)
<i>pporRFP</i> , [<i>hph</i>]	<i>Agrobacterium</i>	Burris et al. (2009)
<i>uidA</i> , <i>GUSPlus</i> , [<i>hph</i>]	<i>Agrobacterium</i>	Xi et al. (2009)
<i>sgfp</i> , [<i>hph</i>]	<i>Agrobacterium</i>	Li & Qu (2011)
<i>sgfp</i> , <i>GUSPlus</i> , [<i>bar</i>] ^b , [<i>hph</i>]	<i>Agrobacterium</i>	Xu et al. (2011)
<i>uidA</i> , [<i>bar</i>], [<i>hph</i>], [<i>nph</i>]	<i>Agrobacterium</i>	Song et al. (2011)

^aSelectable marker genes in brackets.
^bNot used as a selectable marker.

Table 7. A list of transgenes and gene transfer methods in used in reported switchgrass transformation systems

The first transgenic switchgrass plants were produced by the gene gun method (Richards et al., 2001). Tungsten particles coated with a plasmid harboring the *sgfp* and *bar* genes were introduced into immature inflorescence-derived embryogenic calli. In total, 2,430 calli were subjected to the gene transfer, and 97 plants eventually showed tolerance, conferred by the *bar* gene, to 0.1% Basta herbicide (Richards et al., 2001). Basta tolerance was also observed in T₁ transgenic progeny resulting from crosses between transgenic and nontransgenic control plants, indicating inheritance of the *bar* gene (Richards et al., 2001).

All subsequent reports on switchgrass transformation have used *Agrobacterium*-mediated transformation (Table 7). The first such report involved sophisticated experiments that examined the effects of various explants of different genotypes for callus induction and of various concentrations of acetosyringone during inoculation and cocultivation on gene transfer efficiency (Somleva et al., 2002). The gene transformation efficiencies ranged from 0 to 97.3%, but the efficiencies were clearly depended on genotype and kind of explant. Improvements in basal media should lead to efficient transformation in switchgrass. Recently, switchgrass Type II calli were induced from newly-developed LP9 medium (see section 3.1) and were subjected to the *Agrobacterium*-mediated transformation with a final transformation efficiency of 4.4% (Burris et al., 2009).

Because *Agrobacterium* is the causal agent of crown gall, interactions between *Agrobacterium* and plant cells during gene transfer often trigger an undesired plant defense mechanism, called hypersensitive cell death, that leads to lower gene transfer efficiency. Thus, highly compatible *Agrobacterium* strains must be selected for plant transformation. Indeed, different strains of *Agrobacterium*-strain varied in their transient transgene expression efficiencies (Chen et al., 2010) and in stable transformation rates in switchgrass (Song et al., 2011; Xi et al., 2009). For stable transformation of switchgrass, the most noteworthy *Agrobacterium* strain so far has been EHA105; two different research groups found it to be more effective than other strains, such as AGL1, GV3101, and LBA4404 (Song et al., 2011; Xi et al., 2009).

Li & Qu (2011) recently developed the most high-through-put system to date, with a stable transformation efficiency of over 90%. Although most previous transformation systems employed the cultivar 'Alamo', the authors found higher transformation efficiency with the cultivar 'Performer'. Their modifications included infection under vacuum, co-cultivation in desiccation conditions, resting between co-cultivation and selection, and L-proline supplementation in the callus culture and selection media (Li & Qu, 2011). Interestingly, this system also employed *Agrobacterium* strain EHA105, suggesting that it may be highly compatible with 'Performer'. Switchgrass transformation using a EHA105 with the switchgrass selected line HR8, which has high somatic embryogenic capacity (Xu et al., 2011) (see section 3.1), may be a promising avenue for future research.

Transgene inheritance and phenotypic expression in progeny derived from *Agrobacterium*-mediated transformation were observed in some of these reports (Li & Qu, 2011; Somleva et al., 2002; Xi et al., 2009).

4.2 Sugarcane transformation systems

Numerous reports on transformation systems in sugarcane have been previously reviewed (Hotta et al., 2010; Lakshmanan et al., 2005; Suprasanna et al., 2008a). Here, we summarize two key studies published in 2011.

Sophisticated adjustment of culture conditions for several cultivars have resulted in transformation systems widely-adaptable to different genotypes. Basnayake et al. (2011) investigated the effects of 2,4-D levels during callus proliferation, geneticin concentrations during selection, and/or light intensity during regeneration in 16 Australian sugarcane cultivars destined for microprojectile-mediated transformation. This study will be a useful guide for the rapid optimization of key tissue culture variables for efficient genetic transformation of other sugarcane cultivars.

Microprojectile-mediated gene transfer is the most common gene transfer method in sugarcane (Hotta et al., 2010), but the method often causes complex transgene integration that may result in gene silencing. However, minimal expression cassettes lacking vector backbone sequences have been reported to support simple transgene integration in plants. Most recently, a linear minimal expression cassette for the *npt* gene was introduced into embryogenic callus by microprojectile-mediated gene transfer of different amounts of the expression cassette DNA (Kim et al., 2011). Genomic DNA from transgenic sugarcane plants displayed two to 13 *npt* hybridization signals on Southern blots, and the authors observed a trend toward reduced transgene integration complexity and reduced transgene expression levels when lower amounts of the minimal expression cassette were used per shot. This

suggests that backbone free minimal expression cassettes might be efficiently integrated and expressed in sugarcane and other plant species.

4.3 *Miscanthus* transformation systems

The first transgenic *Miscanthus* plants produced via particle bombardment were reported in *Miscanthus sacchariflorus* (Zili et al., 2004, as cited in Jakob et al., 2009), although we were unable to obtain the original reference and cannot report the details. Later, transgenic *M. sinensis* plants were produced by particle bombardment methods (Wang et al., 2011); four transgenic plants containing a foreign *hph* gene with or without the *gfp* were recovered from 120 bombarded calli, and foreign gene expression was confirmed by reverse transcription-PCR analyses. So far, there are no other reports on *Miscanthus* transformation.

4.4 Giant reed transformation systems

An optimized particle bombardment protocol for gene transfer with embryogenic calli was recently reported in giant reed (Dhir et al., 2010). In the study, embryogenic calli were induced from inflorescence segments collected from field-grown mature plants, and several physical parameters, such as helium pressure, bombardment distance to target tissue, and vacuum pressure, together with other factors such as gold microparticle size, DNA concentration, and the number of bombardments, were examined with transient expression of beta-glucuronidase (GUS) and green fluorescent protein (GFP) genes (Dhir et al., 2010). Unfortunately, however, no transgenic plants were obtained due to the lack of regeneration potential of the tissue culture system (Dhir et al., 2010).

We also tried to produce transgenic giant reed plants by using particle bombardment-mediated gene transfer in combination with the plant regeneration system described above (see section 3.5). We obtained more than 100 hygromycin-resistant calli from ca. 5,000 bombarded calli (Figure 5A). Transformation of these calli with foreign *hph* was confirmed by PCR analyses (data not shown). However, we could not obtain transgenic plants from the resistant calli, because the calli had lost their shoot formation potential and produced only adventitious roots on regeneration medium (Figure 5B). Thus, improvements in these tissue culture systems are needed to produce transgenic giant reed plants.




Fig. 5. Transgenic calli of giant reed. A hygromycin-resistant callus propagated during selective culture with the antibiotic hygromycin (arrow) (A), and adventitious root formation from the calli on plant regeneration medium (B).

5. Transgenic plants with improved traits

Here, we summarize reports of transgenic switchgrass and sugarcane plants having distinctive improved traits that will be of significance for biofuel production. In switchgrass, we know of only four such reports, as shown in Table 8. However, in sugarcane, several reports exist on the metabolic engineering of value-added sugarcane via carbohydrate biosynthesis and increased sucrose accumulation; the target genes and the resulting transgenic plants have been discussed in previous reviews (Hotta et al., 2010; Lakshmanan et al., 2005; Suprasanna et al., 2008a; Waclawovsky et al., 2010). Thus, for transgenic sugarcane, we focus here on reports published from 2007 to 2011 (Table 8). No studies have yet been reported for improved traits in *Miscanthus*, *Erianthus*, or giant reed.

Transgenes ^a	Gene transfer methods	Reference
Switchgrass		
<i>phaA</i> , <i>phaB</i> , <i>phaC</i> , [<i>bar</i>]	<i>Agrobacterium</i>	Somleva et al. (2008)
<i>PvCAD</i> ^b [<i>hph</i>]	<i>Agrobacterium</i>	Fu et al. (2011b)
<i>PviCAD2</i> ^b [<i>hph</i>]	<i>Agrobacterium</i>	Saathoff et al. (2011)
<i>COMT</i> ^b [<i>hph</i>]	<i>Agrobacterium</i>	Fu et al. (2011a)
Sugarcane		
<i>phaA</i> , <i>phaB</i> , <i>phaC</i> , [<i>npt</i>]	Gene gun	Petrasovits et al. (2007)
<i>mds6pdh</i> , <i>zmglk</i> , [<i>npt</i>]	Gene gun	Chong et al. (2007)
<i>HIS</i> CaneCPI-1, [<i>npt</i>]	Gene gun	Ribeiro et al. (2008)
<i>avidin</i> ^c , [<i>adhA</i>]	Gene gun	Jackson et al. (2010)
<i>mutB</i> , [<i>npt</i>]	Gene gun	Hamerli & Birch (2011)

^a Selectable marker genes in brackets.
^b Partial inverted repeat sequences for RNAi technology.
^c Several signal peptides were fused for subcellular localization.

Table 8. A list of transgenic plants with improved traits

5.1 Transgenic switchgrass plants

A pioneering study of transgenic switchgrass involved the successful production of value-added transgenic plants that could synthesize polyhydroxybutyrate (PHB), a biodegradable alternative to standard consumer plastic (Somleva et al., 2008). Primary transgenic plants containing up to 3.72% dry weight of PHB in leaf tissues and 1.23% dry weight of PHB in whole tillers were obtained. Polymer accumulation was also analyzed in the T1 generation. This achievement resulted from the successful expression of a functional multigene pathway involving complex metabolic engineering (Somleva et al., 2008).

In the first half of 2011, three reports were published on the downregulation of lignin biosynthesis genes using RNAi technology to improve ethanol production from lignocellulosic biomass (Fu et al., 2011a; Fu et al., 2011b; Saathoff et al., 2011). The first two reports were published at almost the same time and both targeted the gene encoding cinnamyl alcohol dehydrogenase (CAD), a key enzyme for catalyzing the last step of lignin

monomer biosynthesis. The resulting transgenic plants showed significantly fewer transcripts of the target gene, reduced CAD activity, lower lignin content (Fu et al., 2011b; Saathoff et al., 2011), and altered lignin composition (Fu et al., 2011b). Furthermore, these modified lignin biosynthesizers had improved sugar release from cell walls with or without chemical pre-treatment (Fu et al., 2011b; Saathoff et al., 2011). In another study, the expression of the lignin biosynthesis-related *O*-methyltransferase gene was similarly downregulated, and the resultant transgenic plants had reduced lignin content, digestibility by cellulase, and up to 38% more efficient ethanol yield using conventional biomass fermentation processes (Fu et al., 2011a).

5.2 Transgenic sugarcane plants

PHB production is also targeted in sugarcane molecular breeding. The same gene set and the same strategy used with transgenic switchgrass (Somleva et al., 2008) were also employed for producing transgenic sugarcane (Petrasovits et al., 2007). In the study, PHB accumulated in leaves of transgenic plants to a maximum of 1.88% of dry weight without obvious deleterious effects. The PHB concentration in culm internodes of the transgenic plants was much lower (0.0033%) than in leaves (Petrasovits et al., 2007).

The same research group also examined sugar manipulation in sugarcane by engineering a new carbon sink for the six-carbon sugar alcohol sorbitol; sorbitol has intrinsic value as a non-caloric sweetener and is also used to manufacture ascorbic acid (Chong et al., 2007). Transgenic sugarcane plants expressing the *mds6pdh* gene accumulated the sorbitol. The average amounts of sorbitol detected in the most productive line were 120 mg g⁻¹ dry weight (equivalent to 61% of the soluble sugars) in the leaf lamina and 10 mg g⁻¹ dry weight in the stalk pith, but the accumulation caused evident necrosis in expanding leaves and reduced growth (Chong et al., 2007). More recently, another group focused on production of the sucrose isomer trehalulose (Hamerli & Birch, 2011). The transgenic sugarcane plants accumulated the trehalulose up to 600 mM in juice from mature nodes. Contrary to the case of the sorbitol-accumulating sugarcane, the trehalulose-accumulating transgenic plants were vigorous, and trehalulose production in selected lines was retained over multiple vegetative generations under glasshouse and field conditions (Hamerli & Birch, 2011).

A high value-added dedicated lignocellulosic biofuel crop could also be developed by accumulating heterologous functional proteins that could be used for therapeutic, industrial, or other purposes. This concept is adaptable to the production of cystatin, a natural inhibitor of cysteine proteinases, that can be used to protect against insect attacks (Ribeiro et al., 2008). A transformed sugarcane plant expressing high levels of the His-tagged cystatin gene *HIS*CaneCPI-1 was reported to be useful for production and purification of functional *HIS*CaneCPI-1 protein (Ribeiro et al., 2008). The *HIS*CaneCPI-1 protein purified through affinity chromatography in a nickel column was able to inhibit the catalytic activity of midgut cysteine proteinase purified from the sugarcane weevil *Sphenophorus levis* and human cathepsin L in nanomolar amounts, indicating that this system can be used for the production of functional recombinant proteins (Ribeiro et al., 2008). The accumulated recombinant proteins may disturb cell metabolism. If so, subcellular targeting of the recombinant proteins would be necessary to optimize protein yield and avoid detrimental effects of the accumulated protein. Subcellular targeting peptides are thus considered a useful tool for compartmentalization of recombinant proteins. To find ideal subcellular

targeting for recombinant protein accumulation in sugar cane, Jackson et al. (2010) used the glycoprotein avidin, which is potentially toxic to cells, as a test protein fused with several subcellular targeting peptides. Accumulation of avidin was directed to the apoplast, endoplasmic reticulum, and lytic and delta type vacuoles. The study identified the delta type vacuole as a promising target, but the efficiency may depend on tissue type. If the protein is resistant or can be protected by proteolytic attack, the lytic vacuole would be a preferable target (Jackson et al., 2010).

6. Conclusions

We described recent progress in the identification of candidate herbaceous perennials for dedicated lignocellulosic biofuel crops. The biotechnological approaches for molecular breeding of these plants as dedicated biofuel crops are still in immature stages. Biotic and abiotic stress tolerance and herbicide resistance are the minimum required traits. Other traits and related technologies will be important and will make large impacts in the molecular breeding of biofuel crop; these include low-fertilizer needs, sophisticated manipulation technology of secondary cell wall biosynthesis, high-efficiency photosynthesis for high productivity, promoted and synchronized flowering for hybridization, accelerated generation and, especially, organellar transformation for the effective accumulation of high-value chemicals and proteins. These traits and candidate target genes are reviewed in other works (Jakob et al., 2009; Vega-Sánchez & Ronald, 2010; Vogel & Jung, 2001).

7. Acknowledgments

This work was supported by the Japan Society for the Promotion of Science (JSPS) Grant-in-Aid for Scientific Research (C) (23580027).

8. References

- Alexandrova, K. S., Denchev, P. D. & Conger, B. V. (1996a). In vitro development of inflorescences from switchgrass nodal segments. *Crop Science*, Vol. 36, No. 1, pp. 175-178, ISSN 0011-183X
- Alexandrova, K. S., Denchev, P. D. & Conger, B. V. (1996b). Micropropagation of switchgrass by node culture. *Crop Science*, Vol. 36, No. 6, pp. 1709-1711, ISSN 0011-183X
- Alwala, S., Suman, A., Arro, J. A., Veremis, J. C. & Kimbeng, C. A. (2006). Target region amplification polymorphism (TRAP) for assessing genetic diversity in sugarcane germplasm collections. *Crop Science*, Vol. 46, pp. 448-455, ISSN 0011-183X
- Amalraj, V. A. & Balasundaram, N. (2006). On the taxonomy of the members of 'Saccharum complex'. *Genetic Resources and Crop Evolution*, Vol. 53, No. 1, pp. 35-41, ISSN 0925-9864
- Ando, S., Sugiura, M., Yamada, T., Katsuta, M., Ishikawa, S., Terashima, Y., Sugimoto, A. & Matsuoka, M. (2011). Overwintering ability and dry matter production of sugarcane hybrids and relatives in the Kanto region of Japan. *Japan Agricultural Research Quarterly*, Vol. 45, No. 3, pp. 259-267, ISSN 0021-3551
- Angelini, L. G., Ceccarini, L., Nasso, N. & Bonari, E. (2009). Comparison of *Arundo donax* L. and *Miscanthus x giganteus* in a long-term field experiment in

- Central Italy: Analysis of productive characteristics and energy balance. *Biomass and Bioenergy*, Vol. 33, No. 4, pp. 635-643, ISSN 0961-9534
- Aravindhakshan, S. C., Epplin, F. M. & Taliaferro, C. M. (2010). Economics of switchgrass and miscanthus relative to coal as feedstock for generating electricity. *Biomass and Bioenergy*, Vol. 34, No. 9, pp. 1375-1383, ISSN 0961-9534
- Asad, S., Arshad, M., Mansoor, S. & Zafar, Y. (2009). Effect of various amino acids on shoot regeneration of sugarcane (*Saccharum officinarum* L.). *African Journal of Biotechnology*, Vol. 8, No. 7, pp. 1214-1218, ISSN 1684-5315
- Asia Biomass Office (2010). To expand domestic biofuel production in Japan. In: *Biomass Topics*, 17.09.2011, Available from:
<http://www.asiabiomass.jp/english/topics/1005_02.html>
- Basnayake, S. W. V., Moyle, R. & Birch, R. G. (2011). Embryogenic callus proliferation and regeneration conditions for genetic transformation of diverse sugarcane cultivars. *Plant Cell Reports*, Vol. 30, No. 3, pp. 439-448, ISSN 0721-7714
- Bhanwra, R. K., Choda, S. P. & Kumar, S. (1982). Comparative embryology of some grasses. *Proceedings of the Indian National Science Academy*, Vol. 48, pp. 152-162
- Bouton, J. (2008). Improvement of switchgrass as a bioenergy crop, In: *Genetic Improvement of Bioenergy Crops*, W. Vermerris (Ed.), pp. 295-308, Springer, ISBN 978-0-387-70804-1, New York, USA
- Burris, J. N., Mann, D. G. J., Joyce, B. L. & Stewart, C. N., Jr. (2009). An improved tissue culture system for embryogenic callus production and plant regeneration in switchgrass (*Panicum virgatum* L.). *Bioenergy Research*, Vol. 2, No. 4, pp. 267-274, ISSN 1939-1234
- Cheavegatti-Gianotto, A., de Abreu, H. M. C., Arruda, P., Bessalho Filho, J. C., Burnquist, W. L., Creste, S., di Ciero, L., Ferro, J. A., de Oliveira Figueira, A. V., de Sousa Filgueiras, T., Grossi-de-Sá, M. d. F., Guzzo, E. C., Hoffmann, H. P., de Andrade Landell, M. G., Macedo, N., Matsuoka, S., de Castro Reinach, F., Romano, E., da Silva, W. J., de Castro Silva Filho, M. & César Ulian, E. (2011). Sugarcane (*Saccharum X officinarum*): A reference study for the regulation of genetically modified cultivars in Brazil *Tropical Plant Biology*, Vol. 4, No. 1, pp. 62-89, ISSN 1935-9756
- Chen, X., Equi, R., Baxter, H., Berk, K., Han, J., Agarwal, S. & Zale, J. (2010). A high-throughput transient gene expression system for switchgrass (*Panicum virgatum* L.) seedlings. *Biotechnology for Biofuels*, Vol. 3, No. 9, Retrieved from:
<<http://www.biotechnologyforbiofuels.com/content/3/1/9>>
- Chong, B. F., Bonnett, G. D., Glassop, D., O'Shea, M. G. & Brumbley, S. M. (2007). Growth and metabolism in sugarcane are altered by the creation of a new hexose-phosphate sink. *Plant Biotechnology Journal*, Vol. 5, No. 2, pp. 240-253, ISSN 1467-7644
- Chu, C. C., Wang, C. C., Sun, C. S., Hsu, C., Yin, K. C., Chu, C. Y. & Bi, F. Y. (1975). Establishment of an efficient medium for anther culture of rice through comparative experiments on nitrogen-sources. *Scientia Sinica*, Vol. 18, No. 5, pp. 659-668
- Clifton-Brown, J., Chiang, Y.-C. & Hodkinson, T. R. (2008). *Miscanthus*: Genetic resources and breeding potential to enhance bioenergy production, In: *Genetic Improvement of Bioenergy Crops*, W. Vermerris (Ed.), pp. 273-294, Springer, ISBN 978-0-387-70804-1, New York, USA

- Creemers-Molenaar, J., Loeffen, J. P. M. & Van der Valk, P. (1988). The effect of 2,4-dichlorophenoxyacetic acid and donor plant environment on plant regeneration from immature inflorescence-derived callus of *Lolium perenne* L. and *Lolium multiflorum* L. *Plant Science*, Vol. 57, No. 2, pp. 165-172, ISSN 0168-9452
- Denchev, P. D. & Conger, B. V. (1994). Plant-regeneration from callus-cultures of switchgrass. *Crop Science*, Vol. 34, No. 6, pp. 1623-1627, ISSN 0011-183X
- Denchev, P. D. & Conger, B. V. (1995). In-vitro culture of switchgrass - influence of 2,4-D and picloram in combination with benzyladenine on callus initiation and regeneration. *Plant Cell, Tissue and Organ Culture*, Vol. 40, No. 1, pp. 43-48, ISSN 0167-6857
- Deuter, M. (2000). Breeding approaches to improvement of yield and quality in *Miscanthus* grown in Europe, In: *European miscanthus improvement – final Report September 2000*, I. Lewandowski & J. C. Clifton-Brown (Eds.), pp. 28-52, Institute of Crop Production and Grassland Research, University of Hohenheim, Stuttgart, Germany
- Dhir, S., Knowles, K., Pagan, C. L., Mann, J. & Dhir, S. (2010). Optimization and transformation of *Arundo donax* L. using particle bombardment. *African Journal of Biotechnology*, Vol. 9, No. 39, pp. 6460-6469, ISSN 1684-5315
- FAO (2003). Important commodities in agricultural trade: sugar. In: *FAO fact sheets: Input for the WTO ministerial meeting in Cancún*, 16.09.2011, Available from: <<http://www.fao.org/docrep/005/y4852e/y4852e11.htm>>
- Fu, C. X., Mielenz, J. R., Xiao, X. R., Ge, Y. X., Hamilton, C. Y., Rodriguez, M., Chen, F., Foston, M., Ragauskas, A., Bouton, J., Dixon, R. A. & Wang, Z. Y. (2011a). Genetic manipulation of lignin reduces recalcitrance and improves ethanol production from switchgrass. *Proceedings of the National Academy of Sciences of the United States of America*, Vol. 108, No. 9, pp. 3803-3808, ISSN 0027-8424
- Fu, C. X., Xiao, X. R., Xi, Y. J., Ge, Y. X., Chen, F., Bouton, J., Dixon, R. A. & Wang, Z. Y. (2011b). Downregulation of cinnamyl alcohol dehydrogenase (CAD) leads to improved saccharification efficiency in switchgrass. *Bioenergy Research*, Vol. 4, No. 3, pp. 153-164, ISSN 1939-1234
- Garcia, R., Cidade, D., Castellar, A., Lips, A., Magioli, C., Callado, C. & Mansur, E. (2007). In vitro morphogenesis patterns from shoot apices of sugar cane are determined by light and type of growth regulator. *Plant Cell, Tissue and Organ Culture*, Vol. 90, No. 2, pp. 181-190, ISSN 0167-6857
- Gawel, N. J., Robacker, C. D. & Corley, W. L. (1990). In vitro propagation of *Miscanthus sinensis*. *Hortscience*, Vol. 25, No. 10, pp. 1291-1293, ISSN 0018-5345
- Głowacka, K. & Jezowski, S. (2009a). Genetic and nongenetic factors influencing callus induction in *Miscanthus sinensis* (Anderss.) anther cultures. *Journal of Applied Genetics*, Vol. 50, No. 4, pp. 341-345, ISSN 1234-1983
- Głowacka, K. & Jezowski, S. (2009b). In vitro culture of *Miscanthus sinensis* anthers. *Acta Biologica Cracoviensia Series Botanica*, Vol. 51, Suppl. 1, p. 17, ISSN 0001-5296
- Głowacka, K., Jezowski, S. & Kaczmarek, Z. (2010). The effects of genotype, inflorescence developmental stage and induction medium on callus induction and plant regeneration in two *Miscanthus* species. *Plant Cell, Tissue and Organ Culture*, Vol. 102, No. 1, pp. 79-86, ISSN 0167-6857

- Gorenflot, R., Raicu, P., Cartier, D. & Plantefol, L. (1972). Caryologie de la canne de Provence (*Arundo donax* L.). *Comptes rendus de l'Académie des Sciences*, Vol. 274, No. 3, pp. 391-393, (in French)
- Gupta, S. D. & Conger, B. V. (1998). *In vitro* differentiation of multiple shoot clumps from intact seedlings of switchgrass. *In Vitro Cellular & Developmental Biology-Plant*, Vol. 34, No. 3, pp. 196-202, ISSN 1054-5476
- Gupta, S. D. & Conger, B. V. (1999). Somatic embryogenesis and plant regeneration from suspension cultures of switchgrass. *Crop Science*, Vol. 39, No. 1, pp. 243-247, ISSN 0011-183X
- Hamerli, D. & Birch, R. G. (2011). Transgenic expression of trehalulose synthase results in high concentrations of the sucrose isomer trehalulose in mature stems of field-grown sugarcane. *Plant Biotechnology Journal*, Vol. 9, No. 1, pp. 32-37, ISSN 1467-7644
- Henry, R. J. (2010). Basic information on the sugarcane plant, In: *Genetics, Genomics and Breeding of Sugarcane*, R. Henry & C. Kole (Eds.), pp. 1-7, CRC Press, ISBN 978-1-57808-684-9, Boca Raton, Florida, USA
- Holme, I. B., Krogstrup, P. & Hansen, J. (1997). Embryogenic callus formation, growth and regeneration in callus and suspension cultures of *Miscanthus x ogiformis* Honda 'Giganteus' as affected by proline. *Plant Cell, Tissue and Organ Culture*, Vol. 50, No. 3, pp. 203-210, ISSN 0167-6857
- Holme, I. B. & Petersen, K. K. (1996). Callus induction and plant regeneration from different explant types of *Miscanthus x ogiformis* Honda 'Giganteus'. *Plant Cell, Tissue and Organ Culture*, Vol. 45, No. 1, pp. 43-52, ISSN 0167-6857
- Hopkins, A. A., Taliaferro, C. M., Murphy, C. D. & Christian, D. (1996). Chromosome number and nuclear DNA content of several switchgrass populations. *Crop Science*, Vol. 36, No. 5, pp. 1192-1195, ISSN 0011-183X
- Hotta, C. T., Lembke, C. G., Domingues, D. S., Ochoa, E. A., Cruz, G. M. Q., Melotto-Passarin, D. M., Marconi, T. G., Santos, M. O., Mollinari, M., Margarido, G. R. A., Crivellari, A. C., dos Santos, W. D., de Souza, A. P., Hoshino, A. A., Carrer, H., Souza, A. P., Garcia, A. A. F., Buckeridge, M. S., Menossi, M., Van Sluys, M.-A. & Souza, G. M. (2010). The biotechnology roadmap for sugarcane improvement. *Tropical Plant Biology*, Vol. 3, No. 2, pp. 75-87, ISSN 1935-9756
- Jackson, M. A., Nutt, K. A., Hassall, R. & Rae, A. L. (2010). Comparative efficiency of subcellular targeting signals for expression of a toxic protein in sugarcane. *Functional Plant Biology*, Vol. 37, No. 8, pp. 785-793, ISSN 1445-4408
- Jakob, K., Zhou, F. S. & Paterson, A. (2009). Genetic improvement of C4 grasses as cellulosic biofuel feedstocks. *In Vitro Cellular & Developmental Biology-Plant*, Vol. 45, No. 3, pp. 291-305, ISSN 1054-5476
- Jalaja, N. & Sreenivasan, T. (1999). *In vitro* regeneration and cytological studies on somaclones of *Erianthus elephantinus* Hook. F. *Sugar Tech*, Vol. 1, No. 4, pp. 132-138, ISSN 0972-1525
- Karpenstein-Machan, M. (2001). Sustainable cultivation concepts for domestic energy production from biomass. *Critical Reviews in Plant Sciences*, Vol. 20, No. 1, pp. 1-14, ISSN 0735-2689
- Kaur, A. & Gosal, S. S. (2009). Desiccation of callus enhances somatic embryogenesis and subsequent shoot regeneration in sugarcane. *Indian Journal of Biotechnology*, Vol. 8, No. 3, pp. 332-334, ISSN 0972-5849


- Khudamrongsawat, J., Tayyar, R. & Holt, J. S. (2004). Genetic diversity of giant reed (*Arundo donax*) in the Santa Ana River, California. *Weed Science*, Vol. 52, No. 3, pp. 395-405, ISSN 0043-1745
- Kim, J. Y., Gallo, M. & Altpeter, F. (2011). Analysis of transgene integration and expression following biolistic transfer of different quantities of minimal expression cassette into sugarcane (*Saccharum* spp. hybrids). *Plant Cell, Tissue and Organ Culture*, ISSN 0167-6857, (Published online: September 21, 2011)
- Lakshmanan, P. (2006). Somatic embryogenesis in sugarcane - An addendum to the invited review 'Sugarcane biotechnology: The challenges and opportunities,' *In Vitro Cell. Dev. Biol. Plant* 41(4): 345-363; 2005. *In Vitro Cellular & Developmental Biology-Plant*, Vol. 42, No. 3, pp. 201-205, ISSN 1054-5476
- Lakshmanan, P., Geijskes, R. J., Aitken, K. S., Grof, C. L. P., Bonnett, G. D. & Smith, G. R. (2005). Sugarcane biotechnology: The challenges and opportunities. *In Vitro Cellular & Developmental Biology-Plant*, Vol. 41, No. 4, pp. 345-363, ISSN 1054-5476
- Lewandowski, I., Clifton-Brown, J. C., Scurlock, J. M. O. & Huisman, W. (2000). Miscanthus: European experience with a novel energy crop. *Biomass and Bioenergy*, Vol. 19, No. 4, pp. 209-227, ISSN 0961-9534
- Lewandowski, I. & Kahnt, G. (1992). Development of a tissue culture system with unemerged inflorescences of *Miscanthus* 'Giganteus' for the induction and regeneration of somatic embryoids. *Beiträge zur Biologie der Pflanzen*, Vol. 67, No. 3, pp. 439-451, ISSN 0005-8041
- Lewandowski, I. & Kahnt, G. (1993). Possibilities for the establishment of an *In-vitro*-propagationsystem for *Miscanthus* "Giganteus" by using different parts of the plant as donor-tissue. *Bodenkultur*, Vol. 44, No. 3, pp. 243-252, ISSN 0006-5471, (in German)
- Lewandowski, I., Scurlock, J. M. O., Lindvall, E. & Christou, M. (2003). The development and current status of perennial rhizomatous grasses as energy crops in the US and Europe. *Biomass and Bioenergy*, Vol. 25, No. 4, pp. 335-361, ISSN 0961-9534
- Li, R. & Qu, R. (2011). High throughput *Agrobacterium*-mediated switchgrass transformation. *Biomass and Bioenergy*, Vol. 35, No. 3, pp. 1046-1054, ISSN 0961-9534
- McLaughlin, S. B. & Kszos, L. A. (2005). Development of switchgrass (*Panicum virgatum*) as a bioenergy feedstock in the United States. *Biomass and Bioenergy*, Vol. 28, No. 6, pp. 515-535, ISSN 0961-9534
- Mukherjee, S. K. (1957). Origin and distribution of *Saccharum*. *Botanical Gazette*, Vol. 119, pp. 55-61, ISSN 0006-8071
- Murashige, T. & Skoog, F. (1962). A revised medium for rapid growth and bioassays with tobacco tissue cultures. *Physiologia plantarum*, Vol. 15, No. 3, pp. 473-497, ISSN 0031-9317
- Nieves, N., Sagarra, F., González, R., Lezcano, Y., Cid, M., Blanco, M. A. & Castillo, R. (2008). Effect of exogenous arginine on sugarcane (*Saccharum* sp.) somatic embryogenesis, free polyamines and the contents of the soluble proteins and proline. *Plant Cell, Tissue and Organ Culture*, Vol. 95, No. 3, pp. 313-320, ISSN 0167-6857
- Odjakova, M. K. & Conger, B. V. (1999). The influence of osmotic pretreatment and inoculum age on the initiation and regenerability of switchgrass suspension

- cultures. *In Vitro Cellular & Developmental Biology-Plant*, Vol. 35, No. 6, pp. 442-444, ISSN 1054-5476
- Öztürk, L. & Demir, Y. (2002). *In vivo* and *in vitro* protective role of proline. *Plant Growth Regulation*, Vol. 38, No. 3, pp. 259-264, ISSN 0167-6903
- Papazoglou, E. G. (2007). *Arundo donax* L. stress tolerance under irrigation with heavy metal aqueous solutions. *Desalination*, Vol. 211, No. 1-3, pp. 304-313, ISSN 0011-9164
- Papazoglou, E. G., Karantounias, G. A., Vemmos, S. N. & Bouranis, D. L. (2005). Photosynthesis and growth responses of giant reed (*Arundo donax* L.) to the heavy metals Cd and Ni. *Environment International*, Vol. 31, No. 2, pp. 243-249, ISSN 0160-4120
- Perdue, R. E. (1958). *Arundo donax*—Source of musical reeds and industrial cellulose. *Economic Botany*, Vol. 12, No. 4, pp. 368-404, ISSN 0013-0001
- Petersen, K. K. (1997). Callus induction and plant regeneration in *Miscanthus x ogiformis* Honda 'Giganteus' as influenced by benzyladenine. *Plant Cell, Tissue and Organ Culture*, Vol. 49, No. 2, pp. 137-140, ISSN 0167-6857
- Petersen, K. K., Hansen, J. & Krogstrup, P. (1999). Significance of different carbon sources and sterilization methods on callus induction and plant regeneration of *Miscanthus x ogiformis* Honda 'Giganteus'. *Plant Cell, Tissue and Organ Culture*, Vol. 58, No. 3, pp. 189-197, ISSN 0167-6857
- Petrasovits, L. A., Purnell, M. P., Nielsen, L. K. & Brumbley, S. M. (2007). Production of polyhydroxybutyrate in sugarcane. *Plant Biotechnology Journal*, Vol. 5, No. 1, pp. 162-172, ISSN 1467-7652
- Piperidis, G., Piperidis, N. & D'Hont, A. (2010). Molecular cytogenetic investigation of chromosome composition and transmission in sugarcane. *Molecular Genetics and Genomics*, Vol. 284, No. 1, pp. 65-73, ISSN 1617-4615
- Plažek, A. & Dubert, F. (2009). Optimization of medium for callus induction and plant regeneration of *Miscanthus x giganteus*. *Acta Biologica Cracoviensia Series Botanica*, Vol. 51, Suppl. 1, pp. 56-56, ISSN 0001-5296
- Plažek, A. & Dubert, F. (2010). Improvement of medium for *Miscanthus x giganteus* callus induction and plant regeneration. *Acta Biologica Cracoviensia Series Botanica*, Vol. 52, No. 1, pp. 105-110, ISSN 0001-5296
- Pyter, R., Voigt, T., Heaton, E., Dohleman, F. & Long, S. (2007). Giant miscanthus: biomass crop for Illinois. *Proceeding of the 6th National New Crops Symposium*, pp. 39-42, San Diego, California, October 14-18, 2007
- Ribeiro, C. W., Soares-Costa, A., Falco, M. C., Chabregas, S. M., Ulian, E. C., Cotrin, S. S., Carmona, A. K., Santana, L. A., Oliva, M. L. V. & Henrique-Silva, F. (2008). Production of a his-tagged canecystatin in transgenic sugarcane and subsequent purification. *Biotechnology Progress*, Vol. 24, No. 5, pp. 1060-1066, ISSN 8756-7938
- Richards, H. A., Rudas, V. A., Sun, H., McDaniel, J. K., Tomaszewski, Z. & Conger, B. V. (2001). Construction of a GFP-BAR plasmid and its use for switchgrass transformation. *Plant Cell Reports*, Vol. 20, No. 1, pp. 48-54, ISSN 0721-7714
- Rossa, B., Tüffers, A. V., Naidoo, G. & von Willert, D. J. (1998). *Arundo donax* L. (Poaceae) - a C-3 species with unusually high photosynthetic capacity. *Botanica Acta*, Vol. 111, No. 3, pp. 216-221, ISSN 0932-8629
- Saathoff, A. J., Sarath, G., Chow, E. K., Dien, B. S. & Tobias, C. M. (2011). Downregulation of cinnamyl-alcohol dehydrogenase in switchgrass by RNA silencing results in

- enhanced glucose release after cellulase treatment. *PLoS ONE*, Vol. 6, No. e16416, Retrieved from:
<http://www.plosone.org/article/info%3Adoi%2F10.1371%2Fjournal.pone.0016416>
- Somleva, M. N., Snell, K. D., Beaulieu, J. J., Peoples, O. P., Garrison, B. R. & Patterson, N. A. (2008). Production of polyhydroxybutyrate in switchgrass, a value-added co-product in an important lignocellulosic biomass crop. *Plant Biotechnology Journal*, Vol. 6, No. 7, pp. 663-678, ISSN 1467-7652
- Somleva, M. N., Tomaszewski, Z. & Conger, B. V. (2002). *Agrobacterium*-mediated genetic transformation of switchgrass. *Crop Science*, Vol. 42, No. 6, pp. 2080-2087, ISSN 0011-183X
- Song, G., Walworth, A. & Hancock, J. (2011). Factors influencing *Agrobacterium*-mediated transformation of switchgrass cultivars. *Plant Cell, Tissue and Organ Culture*, ISSN 0167-6857, (Published online: September 24, 2011)
- Sun, G. Z., Ma, M. Q., Zhang, Y. Q., Xian, X. L., Cai, X. L. & Li, X. P. (1999). A medium adapted to embryogenic callus induction and subcultures of wheat. *Journal of Hebei Agricultural Sciences*, Vol. 3, No. 2, pp. 24-26, ISSN 1008-1631, (in Chinese)
- Suprasanna, P., Patade, V. Y. & Bapat, V. A. (2008a). Sugarcane biotechnology - A perspective on recent developments and emerging opportunities, In: *Advances in plant biotechnology*, G. P. Rao, Y. Zhao, V. V. Radchuk & S. K. Bhatnagar (Eds.), pp. 313-342, Studium Press LLC, ISBN 19-3369-938-8, Houston, Texas, USA
- Suprasanna, P., Rupali, C., Desai, N. S. & Bapat, V. A. (2008b). Partial desiccation augments plant regeneration from irradiated embryogenic cultures of sugarcane. *Plant Cell, Tissue and Organ Culture*, Vol. 92, No. 1, pp. 101-105, ISSN 0167-6857
- Tagane, S., Ponragdee, W., Sansayawichai, T., Sugimoto, A. & Terajima, Y. (2011). Characterization and taxonomical note about Thai *Erianthus* germplasm collection: the morphology, flowering phenology and biogeography among *E. procerus* and three types of *E. arundinaceus*. *Genetic Resources and Crop Evolution*, ISSN 0925-9864
- Takahashi, W., Komatsu, T., Fujimori, M. & Takamizo, T. (2004). Screening of regenerable genotypes of Italian ryegrass (*Lolium multiflorum* Lam.). *Plant Production Science*, Vol. 7, No. 1, pp. 55-61, ISSN 1343-943X
- Takahashi, W., Takamizo, T., Kobayashi, M. & Ebina, M. (2010). Plant regeneration from calli in giant reed (*Arundo donax* L.). *Grassland Science*, Vol. 56, No. 4, pp. 224-229, ISSN 1744-6961
- Terajima, Y., Terauchi, T., Sakaigaichi, T., Hattori, T., Fujisaki, N., Teruya, H., Naitou, T., Daiku, M., Matsuoka, M., Ohara, S., Irei, S., Ujihara, K. & Sugimoto, A. (2010). Productivities of a large biomass sugarcane line KY01-2044 in Nansei islands. *Proceeding of the 229th Meeting of the Crop Science Society of Japan*, pp. 124-125, Utsunomiya, Japan, March 30-31, 2010, (in Japanese)
- Tew, T. L. & Cobill, R. M. (2008). Genetic improvement of suagarcane (*Saccharum* spp.) as an energy crop, In: *Genetic Improvement of Bioenergy Crops*, W. Vermerris (Ed.), pp. 249-272, Springer, ISBN 978-0-387-70804-1, New York, USA
- Vega-Sánchez, M. E. & Ronald, P. C. (2010). Genetic and biotechnological approaches for biofuel crop improvement. *Current Opinion in Biotechnology*, Vol. 21, No. 2, pp. 218-224, ISSN 0958-1669
- Vogel, K. P. & Jung, H.-J. G. (2001). Genetic modification of herbaceous plants for feed and fuel. *Critical Reviews in Plant Sciences*, Vol. 20, No. 1, pp. 15-49, ISSN 0735-2689

- Waclawovsky, A. J., Sato, P. M., Lembke, C. G., Moore, P. H. & Souza, G. M. (2010). Sugarcane for bioenergy production: an assessment of yield and regulation of sucrose content. *Plant Biotechnology Journal*, Vol. 8, No. 3, pp. 263-276, ISSN 1467-7644
- Wang, X., Yamada, T., Kong, F. J., Abe, Y., Hoshino, Y., Sato, H., Takamizo, T., Kanazawa, A. & Yamada, T. (2011). Establishment of an efficient *in vitro* culture and particle bombardment-mediated transformation systems in *Miscanthus sinensis* Anderss., a potential bioenergy crop. *Global Change Biology Bioenergy*, Vol. 3, No. 4, pp. 322-332, ISSN 1757-1693
- Williams, C. A., Harborne, J. B. & Smith, P. (1974). Taxonomic significance of leaf flavonoids in *Saccharum* and related genera. *Phytochemistry*, Vol. 13, No. 7, pp. 1141-1149, ISSN 0031-9422
- Wright, L. & Turhollow, A. (2010). Switchgrass selection as a "model" bioenergy crop: A history of the process. *Biomass and Bioenergy*, Vol. 34, No. 6, pp. 851-868, ISSN 0961-9534
- Xi, Y., Fu, C., Ge, Y., Nandakumar, R., Hisano, H., Bouton, J. & Wang, Z.-Y. (2009). *Agrobacterium*-mediated transformation of switchgrass and inheritance of the transgenes. *Bioenergy Research*, Vol. 2, No. 4, pp. 275-283, ISSN 1939-1234
- Xu, B., Huang, L., Shen, Z., Welbaum, G. E., Zhang, X. & Zhao, B. (2011). Selection and characterization of a new switchgrass (*Panicum virgatum* L.) line with high somatic embryogenic capacity for genetic transformation. *Scientia Horticulturae*, Vol. 129, No. 4, pp. 854-861, ISSN 0304-4238
- Zhang, Q. X., Sun, Y., Hu, H. K., Chen, B., Hong, C. T., Guo, H. P., Pan, Y. H. & Zheng, B. S. (2011). Micropropagation and plant regeneration from embryogenic callus of *Miscanthus sinensis*. *In Vitro Cellular & Developmental Biology - Plant*, ISSN 1054-5476, (Published online: Augst 20, 2011)

IntechOpen



Transgenic Plants - Advances and Limitations

Edited by PhD. Yelda Ozden Çiftçi

ISBN 978-953-51-0181-9

Hard cover, 478 pages

Publisher InTech

Published online 07, March, 2012

Published in print edition March, 2012

Development of efficient transformation protocols is becoming a complementary strategy to conventional breeding techniques for the improvement of crops. Thus, Transgenic Plants - Advances and Limitations covers the recent advances carried on improvement of transformation methods together with assessment of the impact of genetically transformed crops on biosafety. Each chapter has been written by one or more experienced researchers in the field and then carefully edited to ensure thoroughness and consistency.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Wataru Takahashi and Tadashi Takamizo (2012). Molecular Breeding of Grasses by Transgenic Approaches for Biofuel Production, Transgenic Plants - Advances and Limitations, PhD. Yelda Ozden Çiftçi (Ed.), ISBN: 978-953-51-0181-9, InTech, Available from: <http://www.intechopen.com/books/transgenic-plants-advances-and-limitations/molecular-breeding-of-grasses-by-transgenic-approaches-for-biofuel-production>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen