We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Nasopharyngeal Carcinoma: The Role for Chemotherapeutics and Targeted Agents

Jared Knol¹, Tiffany King² and Mark Agulnik³

¹*Department of Medicine, Feinberg School of Medicine, Northwestern University, Chicago, Illinois*

²*Division of Hematology/Oncology, Northwestern Medical Faculty Foundation, Chicago, Illinois*

³*Division of Hematology/Oncology, Department of Medicine, Feinberg School of Medicine, Northwestern University, Robert H. Lurie Comprehensive Cancer Center, Chicago, Illinois USA*

1. Introduction

Nasopharyngeal carcinoma (NPC) is a distinguishing and rare form of head and neck cancer whose predominant tumor type arises in the nasopharynx, the narrow tubular passage behind the nasal cavity. The disease is classified into three histopathological types by The World Health Organization (WHO): Keratinizing squamous cell carcinoma (SCC, WHO Type I), Nonkeratinizing carcinoma: differentiated (WHO Type II) and undifferentiated (WHO Type III) and basaloid squamous cell carcinoma¹. Worldwide, there are 80,000 incident cases and 50,000 deaths annually,² however the disease is vastly more common in certain regions in Asia and Africa than anywhere else in the world. In the United States, the incidence ranges from 0.2 and 0.5 cases per 100,000 population, which constitutes roughly 0.02% of all cancers, compared with endemic areas such as areas in Asia, where NPC might represent 25% of all cancers³. In fact, it is sometimes referred to as Cantonese cancer because it occurs in about 25 cases per 100,000 people in this region, 25 times higher than the rest of the world⁴.

Making it distinct is the fact that it differs from other head and neck squamous cell carcinomas (HNSCCs) in epidemiology, histology, natural history, and response to treatment. The remarkable geographic and demographic variation of NPC incidence suggests a multifactorial etiology of NPC. In endemic populations, the risk of NPC appears to be related to an interaction of these factors, namely Epstein-Barr virus (EBV) infection; genetic predisposition; and environmental factors such as the traditionally high intake of preserved foods⁵⁻⁷.

2. RT and chemotherapy for locally advanced disease

The radiosensitizing properties of systemic chemotherapy are firmly established for head and neck tumors⁸, and NPC is inherently more chemosensitive than other head and neck malignancies⁹. Therefore, it follows intuitively that adding systemic chemotherapy or targeted therapies to radiotherapy could provide NPC patients with additional clinical benefit. This hypothesis was tested in the late 1990s, and the utility of platinum-based chemotherapy in addition to radiation (RT) in locally advanced NPC was clearly

demonstrated in 1998 when the Intergroup 0099 trial was terminated prematurely due to a clear survival benefit in the chemotherapy with radiotherapy arm¹⁰. Since that time several randomized trials have been published, all varying in their chemotherapy regimens and timing of chemotherapy in relation to RT (Tables 1 and 2).

Trial	No. of pts	Treatment Arms
Neoadjuvant + RT vs RT alone		
VUMCA (1996)	a: 171 b: 168	a: Bleomycin/Epirubicin/Cisplatin → RT b: RT
Chua et al (1998)	a: 167 b: 167	a: Cisplatin/Epirubicin → RT b: RT
Ma et al (2001)	a: 224 b: 225	a: Cisplatin/Bleomycin/5-FU → RT b: RT
Hareyama et al (2002)	a: 40 b: 40	a: Cisplatin/5-FU → RT b: RT
Concurrent + RT vs RT alone		
Lin et al (2003)	a: 141 b: 143	a: Cisplatin/5-FU + RT b: RT
Chan et al (2002, 2004)	a: 174 b: 176	a: Cisplatin + RT b: RT
Zhang et al (2005)	a: 59 b: 56	a: Oxaliplatin + RT b: RT
Adjuvant + RT vs RT alone		
Rossi et al (1988)	a: 113 b: 116	a: RT → Vincristine/Cyclophosphamide/Adriamycin b: RT
Chi et al (2002)	a: 77	a: RT → Cisplatin/Fluorouracil/Leucovorin b: RT
Neoadjuvant + Adjuvant + RT vs RT alone		
Chan et al (1995)	a: 37 b: 40	a: Cisplatin/Fluorouracil → RT → Cisplatin/5-FU b: RT
Concurrent + Adjuvant + RT vs RT alone		
Al-Sarraf et al (1998, 2001)	a: 93 b: 92	a: Cisplatin + RT → Cisplatin/5-FU b: RT
Wee et al (2005)	a: 111 b: 110	a: Cisplatin + RT → Cisplatin/5-FU b: RT
Lee et al (2005, 2010)	a: 172 b: 176	a: Cisplatin + RT → Cisplatin/5-FU b: RT
Lee et al (2006)	a: 51 b: 44 c: 52 d: 42	a: Cisplatin + RT → Cisplatin/5-FU b: Cisplatin + AF RT → Cisplatin/5-FU c: AF RT d: RT
Kwong et al (2004)	a1: 57 a2: 53 a3: 54 b: 55	a1: Uracil-Tegafur + RT → Cisplatin/5-FU,Vincristine/Bleomycin/MTX a2: Uracil-Tegafur + RT a3: RT → Cisplatin/5-FU, Vincristine/Bleomycin/MTX b: RT
Chen et al (2008)	a: 158 b: 158	a: Cisplatin + RT → Cisplatin/5-FU b: RT

Table 1. Randomized Trials of Chemotherapy with RT vs RT alone in locally advanced NPC

Trial	Treatment Arm	OS	DFS	Median F/U (mo)
Neoadjuvant + RT vs RT alone				
VUMCA (1996)	a	3yr – 60%	3yr – 52%*	49
	b	54%	32%	
Chua et al (1998)	a	3yr – 78%	3yr – 48%	30
	b	71%	42%	
Ma et al (2001)	a	5yr – 63%	5yr – 59%*	62
	b	56%	49%	
Hareyama et al (2002)	a	5yr – 60%	5yr – 55%	49
	b	48%	43%	
Chua (2005)	a	5yr – 62%	5yr – 51%*	67
	b	58%	43%	
Concurrent + RT vs RT alone				
Lin et al (2003)	a	5yr – 72%*	5yr – 72%*	65
	b	54%	53%	
Chan et al (2002, 2004)	a	NR	2yr – 76%	33
	b	NR	69%	
	a	5yr – 70%	5yr – 60%	65
	b	59%	52%	
Zhang et al (2005)	a	2yr – 100%*	2yr – 96%*	24
	b	77%	83%	
Adjuvant + RT vs RT alone				
Rossi et al (1998)	a	5yr – 55%	5yr – 54%	49.5
	b	61%	50%	
Chi et al (2002)	a	4yr – 59%	4yr – 58%	43
	b	67%	56%	
Neoadjuvant + Adjuvant + RT vs RT alone				
Chan et al (1995)	a	2yr – 80%	2yr – 68%	28.5
	b	81%	72%	
Concurrent + Adjuvant + RT vs RT alone				
Al-Sarraf et al (1998, 2001)	a	3yr – 76%*	3yr – 66%*	32.4
	b	46%	26%	
	a	5yr – 67%*	5yr – 58%*	60
	b	37%	29%	
Wee et al (2005)	a	2yr – 85%*	3yr – 80%*	38.4
	b	78%	65%	
Lee et al (2005, 2010)	a	3yr – 77%	3yr – 69%	25
	b	76%	61%	
	a	5yr – 68%	5yr – 62%*	70.8
	b	64%	53%	
	a	8yr – 61%	NR	
	b	54%	NR	
Lee et al (2006)	a	3yr – 87%	3yr – 73%	33
	b	88%	88%	
	c	73%	63%	
	d	83%	65%	
Kwong et al (2004)	a1	3yr – 89%	3yr – 70%	32.5
	a2	84%	69%	
	a3	71%	54%	
	b	83%	61%	
Chen et al (2008)	a	2yr – 90%*	2yr – 85%*	29
	b	80%	73%	

* statistically significant result

Table 2. OS and DFS of randomized trials of chemotherapy with RT vs. RT alone in locally advanced NPC

2.1 Neoadjuvant chemotherapy

Four trials have assessed the role of neoadjuvant chemotherapy followed by RT versus RT alone. The VUMCA study¹¹, active from 1989-1993, was a multi-center phase II trial of 339 patients, in which 171 patients were randomized to receive neoadjuvant bleomycin, epirubicin, and cisplatin (BEC). The remaining 168 patients received RT alone. After a median followup of 49 months the authors noted increased disease free survival (DFS) in the chemotherapy with RT arm (52% vs 32%). However, no difference in overall survival (OS) was seen, and the trial was notable for an 8% rate of treatment-related death in the experimental arm versus 1% among patients treated with RT alone.

A small study of 80 patients, active from 1991-1998, was published by Hareyama et al¹². Forty patients were randomized to cisplatin and 5-fluorouracil prior to RT, and the control arm received RT alone. A trend toward both OS and DFS at three years was seen in the neoadjuvant arm (60% and 55% vs 48% and 43%, respectively), though this did not reach statistical significance. In 1998, Chua et al published the results of a trial active from 1989-1993, in which 334 patients were randomized to either cisplatin and epirubicin with RT or RT alone¹³. Intention to treat analysis at three years revealed a trend toward improved DFS and OS in the chemoradiotherapy arm (58% and 80% vs 46% and 72%, respectively), however the results did not reach statistical significance.

In 2001 Ma et al reported a trial, active from 1993-1994, in which 449 patients were randomized to two to three cycles of cisplatin, 5-fluorouracil, and bleomycin with RT versus RT alone¹⁴. Intention to treat analysis at five years revealed a statistically significant improvement in DFS (59% vs 49%), however, while a trend toward improved OS was noted this did not reach statistical significance. In 2005, Chua published an analysis of pooled data from the latter two trials, which together included 784 patients with a median follow up of 67 months¹⁵. This analysis revealed a statistically significant DFS benefit in the neoadjuvant arm (51% vs 43%). However, consistent with the other trials, a trend toward OS (62% vs 58%) did not reach statistical significance. In summary, four trials of neoadjuvant platinum-based chemotherapy in addition to RT have revealed, at best, modest improvement in DFS and a trend toward improved overall survival. However to date no trial has shown a clear and statistically significant survival benefit with the neoadjuvant strategy.

2.2 Concurrent chemotherapy

To date, three trials have assessed the strategy of concurrent chemotherapy and RT as compared to RT alone. In 2002, Chan et al reported a trial, active from 1994-1997, of 350 patients randomized to weekly low-dose cisplatin plus RT versus RT alone¹⁶. Notably, the study population in this trial had relatively advanced disease: 90% of patients were Ho's stage III or IV, and over 70% were AJCC stage III or IV. A trend toward improved DFS at two years was seen (76% vs 69%), however this did not reach statistical significance. Updated data published in 2005 revealed modestly improved DFS and OS for the chemotherapy arm at five years: DFS was 60% and 52% ($p = 0.06$) and OS was 70% and 59% ($p = 0.05$) for the chemotherapy + RT and RT alone arms, respectively¹⁷. A subgroup analysis of this data revealed that patients with more advanced disease (T3 and T4) derived the most benefit. Lin et al reported data in 2003 from a 1993-1999 study of concurrent cisplatin and 5-fluorouracil plus

RT vs RT alone¹⁸. All patients in this trial had AJCC stage III or IV disease. The study revealed statistically significant improvement in both DFS and OS with concurrent chemotherapy: 5-year OS rates for the chemotherapy arm were 72% compared with 54% in the control arm, and the 5-year DFS rates were 72 versus 53%, respectively. Of note, significantly more toxicity was noted in the chemotherapy arm. In 2005, Zhang et al published a trial, active from 2001-2003, of 115 patients randomized to six doses of weekly oxaliplatin plus concurrent RT versus RT alone¹⁹. Consistent with the earlier cisplatin-based trials, this study revealed a significant DFS and OS benefit for patients treated with concurrent platinum-based chemotherapy. For the chemotherapy + RT and RT alone arms, respectively, two year OS was 100% versus 77% and two year DFS was 96% versus 83%. In summary, three trials of concurrent platinum-based chemotherapy plus radiotherapy have revealed statistically significant improvements in DFS and OS as compared to radiotherapy alone. In these trials, the benefit was greatest in patients with advanced disease.

2.3 Adjuvant chemotherapy

The first trial that addressed the question of adding chemotherapy to RT for patients with NPC used an adjuvant strategy. In 1988, Rossi et al published the results of a 4 year multi-center trial, active from 1979-1983, in which 229 patients were randomized to either adjuvant vincristine, cyclophosphamide, and adriamycin following RT or RT alone²⁰. Follow up at four years did not reveal a statistically significant difference in DFS or OS between the two groups. Notably, no platinum agent was used in this trial, and therefore the lack of benefit must be interpreted in light of subsequent studies which have established platinum-based chemotherapy as the cornerstone of chemotherapy plus RT strategies in this disease. However, a later study of adjuvant chemotherapy, which did include cisplatin, was published in 2002 by Chi et al and revealed no significant DFS or OS benefit²¹. Between 1994 and 1999, 157 patients were randomized to either adjuvant cisplatin with 5-fluorouracil and leucovorin plus RT versus RT alone. At five years, the DFS rates were 54.4% vs 49.5% and the OS rates were 60.5% and 54.5%, respectively, for the chemotherapy and RT arms. Neither of these differences reached statistical significance. In summary, two trials have addressed the strategy of adjuvant chemotherapy following RT, one of which was published prior to the Intergroup 0099 trial and as such did not include a platinum agent. Neither study revealed a statistically significant improvement in survival with adjuvant chemotherapy following RT as compared to RT alone.

2.4 Neoadjuvant plus adjuvant chemotherapy

To date, only one study has assessed the effect of combined neoadjuvant and adjuvant chemotherapy in addition to RT. In 1995, Chan et al reported a study of 82 patients randomized to two cycles of neoadjuvant and four cycles of adjuvant chemotherapy (both with cisplatin) plus RT vs RT alone²². No difference in either DFS or OS was noted after two years follow up. In summary, the single published trial of combined neoadjuvant and adjuvant chemotherapy (cisplatin) plus RT revealed no benefit of this strategy as compared to RT alone.

2.5 Concurrent plus adjuvant chemotherapy

To date, the most influential study to address the question of chemotherapy plus RT in locally advanced NPC utilized a combined concurrent and adjuvant platinum-based

chemotherapy strategy, and, consequently, this approach has subsequently received the most research attention. In 1998, Al-Sarraf et al published the results of the Intergroup 0099 study, active between 1989 and 1995, in which 193 patients were randomized to either RT alone or a chemotherapy arm¹⁰. Patients in the chemotherapy group were treated with cisplatin on days 1, 22 and 43 of the concurrent RT, and three adjuvant cycles of cisplatin with 5-fluorouracil were given monthly after completion of chemoradiotherapy. At 3 years, DFS was 69% in the chemotherapy group and 24% in the RT alone arm. Overall survival at 3 years was 78 *vs* 47%, favoring chemotherapy. These results prompted early closure of the study, given the clear benefit demonstrated with concurrent plus adjuvant cisplatin. Updated analysis at 5 years confirmed the benefit of treatment, with 5-year DFS rates of 58 *vs* 29% and 5-year OS rates of 67 *vs* 37%, both favoring the combined therapy arm²³. Analysis of the National Cancer database since the first published results of the Intergroup 0099 study data in 1998 has demonstrated that this study has led to a demonstrable change in NPC management. Of all patients enrolled in the database and matching the eligibility criteria of the Intergroup 0099 study, only 38% received chemotherapy along with RT prior to 1997, while since the publication of the data, 65% of these patients have received concurrent and adjuvant chemotherapy²⁴.

One-quarter of all patients treated on the Intergroup 0099 protocol had World Health Organization (WHO) stage I histology (keratinizing squamous cell carcinoma). A phase III randomized trial active from 1997-2003, using a similar chemotherapy and RT plan but restricted to patients with WHO type IIa (nonkeratinising squamous cell carcinoma) and IIb (undifferentiated carcinoma) histologies was published by Wee et al in 2005²⁵. The two and three year DFS and OS rates were statistically significant and favored the use of chemotherapy, confirming the findings of the Intergroup 0099 trial. Of note, the chemotherapy regimen used in the Wee et al study differed slightly from the Intergroup 0099 trial in that the dose of cisplatin was given in divided doses rather than one dose, however, the total dose remained the same. Lee et al have published data from the Hong Kong NPC Study Group in which the Intergroup 0099 regimen was applied to patients with nonkeratinizing or undifferentiated NPC²⁶. Preliminary results published in 2004 suggested a trend toward improved DFS in the chemotherapy arm. Long term follow up data published in 2010 confirmed the trend toward improved DFS seen on the initial analysis (62% *vs* 53% favoring chemotherapy). However, while five-year analysis showed a clear reduction death due to disease progression in the chemotherapy arm (28% *vs* 38% favoring chemotherapy, $p = 0.08$), the five-year overall survival data revealed only a modest trend favoring chemotherapy (68% *vs* 64%, $p = 0.22$)²⁷. The discrepancy between a clear decreased in disease-associated death and nearly identical five-year overall survival was attributed to an increase in non-cancer deaths among patients in the chemotherapy arm.

A similar trial, also published by Lee et al, utilized the identical chemotherapy as IG-0099, but assessed both the therapeutic gain with concurrent and adjuvant chemotherapy and/or accelerated RT²⁸. Four arms were evaluated, conventional RT *vs.* accelerated fraction RT *vs.* conventional RT with concurrent and adjuvant chemotherapy *vs.* accelerated fraction RT with concurrent and adjuvant chemotherapy. After 189 patients were randomized, the study was closed due to poor accrual. Preliminary data after a median follow-up of 2.9 years, showed no statistically significant change in either OS or DFS at 3-years.

In 2008, Chen et al published the results of a study of 316 patients performed between 2002 and 2005, in which subjects randomized to the chemotherapy arm received weekly cisplatin concurrent with RT, followed by cisplatin and 5-fluorouracil every four weeks for three cycles following completion of RT²⁹. Preliminary analysis after two years revealed a statistically significant difference in both overall (89.8% vs 79.7%) and disease free (84.6% vs 72.5%) survival, favoring chemotherapy, at the expense of increased toxicity in the chemotherapy arm.

Finally, a factorial study of four different regimens was published by Kwong et al in 2004³⁰. This study assessed the combination of RT alone *vs* three other schemas: RT with adjuvant chemotherapy, concurrent chemoradiotherapy and lastly, concurrent chemoradiotherapy followed by adjuvant chemotherapy. UFT (uracil and tegafur in a 4:1 molar ratio) was given concurrent with RT, while the adjuvant therapy consisted of alternating cycles of cisplatin/5-fluorouracil and vincristine/bleomycin/methotrexate.

Although a trend towards improved DFS and OS was noted with the addition of concurrent chemotherapy, it did not reach statistical significance at 3 years. In assessing distant metastases rates, a significant reduction was attributable to concurrent chemotherapy. In this study, adjuvant chemotherapy did not improve outcome.

3. Chemotherapy for recurrent and metastatic disease

The management of recurrent and metastatic NPC remains challenging. Several studies have evaluated the use of platinum drugs in combination with various other chemotherapeutic agents including gemcitabine^{31,32}, bleomycin-5FU³³, 5FU³⁴, capecitabine³⁵, bleomycin/epirubicin/5FU³⁶, paclitaxel³⁷, and docetaxel³⁸. Given the known chemosensitivity of NPC, it is not surprising that these regimens are associated with good overall response rates, ranging from 56% - 79%. However, the duration of response is short, and median overall survival in these trials was on average approximately one year (range: 11 - 25 mo). Two trials of non-platinum based monotherapy (gemcitabine³⁹ and capecitabine⁴⁰) were associated with worse overall response rates (28% and 37%, respectively) and similar median overall survival (7.2 and 14 months, respectively). Adding vinorelbine to gemcitabine for patients with platinum-resistant disease yielded an overall response rate of 36% with a median overall survival of 11.9 months⁴¹. Irinotecan used as monotherapy in heavily pretreated patients has been associated with an overall response rate of 14% (all partial responses) and median overall survival of 11.4 months⁴². The poor clinical outcomes associated with traditional chemotherapeutic agents underscores the urgent need for new and more effective therapeutic options in the locally advanced and metastatic setting.

4. Targeted therapy

Given the poor clinical outcomes associated with chemotherapeutics in the metastatic setting, evaluation of targeted therapies become of utmost importance (Table 3).

Epidermal growth factor receptor (EGFR) expression in NPC has been correlated with decreased survival, increased rates of locoregional failure, and more aggressive disease^{43,44}. Drawing on those findings, Chan et al published a study in 2005 evaluating the anti-EGFR

Table 3. Targeted therapy in NPC

Reference	Targeted Tx	Other Tx	# of patients/ # evaluable	Disease Status	Overall Response
Chan (2005)	Cetuximab	Carboplatin	60/59	Recurrent/Metastatic	11.7% (1)
Ma (2007)^	Cetuximab	Cisplatin + IMRT	20/12	Locally advanced	83%
Elser (2007) (3)	Sorafenib	None	7/NR	NR	N/A (4)
Chua (2008)	Gefitinib	None	19/NR	Recurrent/Metastatic	0%
Ma (2008)	Gefitinib	None	16/15	Recurrent/Metastatic	0%
You (2009)^	Erlotinib	Cisplatin/Carboplatin + Gemcitabine (5)	20/11	Recurrent/Metastatic	0%

(1)All partial responses
(2)No composite grade 3/4 toxicity data reported. Reports that 85% of patients experienced grade 3/4 leukopenia, 85% grade 3/4 mucositis.
(3)This study included both SCCHN and NPC patients. Data is presented here only for the 7 NPC patients
(4)One of 26 evaluable SCCHN/NPC patients had a partial response. However, it is not reported whether
(5)Erlotinib was given as maintenance after completion of 6 cycles platinum + gemcitabine or as 2nd line therapy
(6)Reports grade 3/4 toxicity as “rare”, most common was rash in (33%)
^ Abstract

monoclonal antibody, cetuximab, in combination with carboplatin⁴⁵. This multi-center phase II trial enrolled 60 patients, all of who had evidence of disease progression within 12 months of platinum-based chemotherapy. Notably, 93.3% of patients in this trial had stage IV disease at the time of enrollment, and 85% had distant metastases. The treatment protocol consisted of carboplatin infused every three weeks for a maximum of eight cycles, along with cetuximab 400mg/m² followed by 250mg/m² weekly. The median number of cetuximab infusions completed was 10, with a range of 1-30. Only 53.3% of the patients received at least 3 carboplatin infusions. Of patients enrolled, 11.7% had a partial response to therapy; no complete responses were observed in this study and 48.3% of patients had stable disease. Median time to progression was 81 days, and median overall survival was 233 days. Grade 3 or 4 toxicity was observed in 51.7% of patients, 31.7% of whom had toxicity that was attributed to cetuximab. A subsequent trial of cetuximab in combination with cisplatin-IMRT was reported in abstract form by Ma et al in 2008⁴⁶. The 20 patients in this trial differed from those in Chan et al in that all subjects had untreated, non-metastatic disease. The cetuximab dosing was the same (400mg/m² initially, followed by 250mg/m² weekly). Cisplatin was administered at 30mg/m² weekly, along with weekly IMRT. Ninety percent of patients received at least 5 doses of cetuximab, and 80% received at least 5 cisplatin infusions. Preliminary analysis at 3 months revealed promising results: of 12 patients evaluable for response, 10 had a complete response and the remaining two had stable disease.

Two studies have evaluated the use of the small molecule EGFR tyrosine kinase inhibitor, gefitinib, as monotherapy in advanced platinum-resistant NPC. In 2008 Chua et al reported a single-center phase II study of 19 patients with relapsed or progressive despite at least two prior chemotherapy regimens (one of which included a platinum drug)⁴⁷. In contrast to the previously mentioned trials, this study protocol assessed response to anti-EGFR therapy alone without any concurrent chemotherapeutic agent. Patients in the study received gefitinib 250mg daily until disease progression, unacceptable toxicity, or patient refusal. Median treatment duration was 10 weeks; 63% of patients were taken off study due to disease progression, and the remaining patients expressed a preference not to continue treatment. The regimen was well tolerated, with no grade 3 or 4 toxicity reported. Unfortunately, no patients in this study achieved either partial or complete response. Similar disappointing results were reported by Ma et al in 2008⁴⁸. In that trial, 16 patients with progressive disease after prior platinum-based therapy were treated with oral gefitinib 500 mg/day. Three patients achieved stable disease (range: 2.8 – 8.5 months), and the drug was generally well tolerated. However, consistent with the results reported by Chua et al, no patient achieved a partial or complete response. In 2009, You et al presented data in abstract form reporting their study of the EGFR tyrosine kinase inhibitor, erlotinib⁴⁹. In that small trial of 20 patients, subjects were first treated with up to 6 cycles of gemcitabine + a platinum agent (either cisplatin or carboplatin), followed by erlotinib 150 mg/day as maintenance (if stable disease) or 2nd line therapy (if progressive disease prior to 6 chemotherapy cycles). Of the 11 patients evaluable for response to erlotinib, three had stable disease that was maintained for 3, 4, and 7 months.

Sorafenib, a multi-kinase inhibitor with activity against VEGFR-2, VEGFR-3, PDGFR, FLT-3, c-kit, and the Raf isoforms c-Raf and b-Raf, was evaluated in a 2007 study published by Elser et al.⁵⁰ Of particular theoretical interest is the anti-angiogenesis activity of sorafenib

mediated by its activity against the vascular endothelium growth factor receptor (VEGFR), given the known VEGF overexpression in NPC⁵¹⁻⁵³. Notably, this study was not limited to NPC and also enrolled patients with SCCHN; of 27 patients enrolled, only 7 had NPC. Patients received sorafenib 400 mg twice daily as monotherapy, and therapy was generally well tolerated with few grade 3 or 4 toxicities. Although the reported response data for the entire patient cohort was promising (9 of 26 evaluable patients had stable disease, and one had a partial response), response data specific to the NPC subgroup was not separately reported. Median time to progression for the 7 patients with NPC was 3.2 months, and median overall survival was 7.7 months.

In addition to monoclonal antibody and small molecule targeted therapy, immunotherapy may play a role in the future management of NPC. This approach has great theoretical appeal in NPC, given the strong association with Epstein-Barr virus (EBV). NPC cells express two distinct EBV latent membrane proteins, LMP-1 and LMP-2, and these proteins represent targets for adoptive immunotherapy⁵⁴. In a phase I study published by Straathof et al in 2005, 10 patients with advanced NPC were treated with EBV-specific cytotoxic T lymphocytes (CTL)⁵⁵. Four patients were in remission at the time of enrollment, and all remained disease-free 19-27 months after infusion. Of the remaining 6 patients, all with relapsed or refractory disease, 2 had complete responses, 1 had a partial response, 1 had stable disease and 2 had no response. CTL was well tolerated in this study. Follow up data from the same group was published in 2010, and this again demonstrated clinical benefit following CTL infusion⁵⁶. Of 23 patients in this study (all with a history of relapsed/refractory NPC), 8 were in remission and 15 were not. Of the 8 patients in remission, 5 remained disease free at the time of publication (range: 17-75 months post-infusion). Furthermore, CTL infusion was associated with a 48.7% overall response rate in patients with active disease, though for the subset of patients with metastatic disease the overall response rate was only 10%.

5. Conclusion

In summary, the anti-EGFR monoclonal antibody, cetuximab, has been associated with a small but significant overall response in patients with recurrent/metastatic disease who progressed despite platinum-based therapy. Furthermore, nearly half of the patients in that study achieved stable disease. Preliminary data in the locally advanced setting are more promising, with one small trial reporting an 83% overall response. In both of these trials, cetuximab was given in combination with platinum-based chemotherapy. Treatment with small molecule inhibitors has been less successful: two studies of the tyrosine kinase inhibitor, gefitinib, have shown no benefit. Inhibitors of angiogenesis are theoretically attractive subjects of investigation given the documented VEGF overexpression in NPC, and these agents warrant further investigation. Finally, the clear association between NPC and EBV represents a disease-specific biologic property that may be exploited via immunotherapy, and early work with EBV-specific cytotoxic T lymphocytes has produced promising results.

6. References

- [1] Thompson L: World Health Organization classification of tumours: pathology and genetics of head and neck tumours. *Ear Nose Throat J* 85:74, 2006

- [2] Parkin DM, Bray F, Ferlay J, et al: Global cancer statistics, 2002. *CA Cancer J Clin* 55:74-108, 2005
- [3] Phillips CA, Hendershot DM: Functional electrical stimulation and reciprocating gait orthosis for ambulation exercise in a tetraplegic patient: a case study. *Paraplegia* 29:268-76, 1991
- [4] Chang ET, Adami HO: The enigmatic epidemiology of nasopharyngeal carcinoma. *Cancer Epidemiol Biomarkers Prev* 15:1765-77, 2006
- [5] Yuan JM, Wang XL, Xiang YB, et al: Preserved foods in relation to risk of nasopharyngeal carcinoma in Shanghai, China. *Int J Cancer* 85:358-63, 2000
- [6] Farrow DC, Vaughan TL, Berwick M, et al: Diet and nasopharyngeal cancer in a low-risk population. *Int J Cancer* 78:675-9, 1998
- [7] Liebowitz D: Nasopharyngeal carcinoma: the Epstein-Barr virus association. *Semin Oncol* 21:376-81, 1994
- [8] Seiwert TY, Salama JK, Vokes EE: The chemoradiation paradigm in head and neck cancer. *Nat Clin Pract Oncol* 4:156-71, 2007
- [9] Chan AT, Teo PM, Leung TW, et al: The role of chemotherapy in the management of nasopharyngeal carcinoma. *Cancer* 82:1003-12, 1998
- [10] Al-Sarraf M, LeBlanc M, Giri PG, et al: Chemoradiotherapy versus radiotherapy in patients with advanced nasopharyngeal cancer: phase III randomized Intergroup study 0099. *J Clin Oncol* 16:1310-7, 1998
- [11] Preliminary results of a randomized trial comparing neoadjuvant chemotherapy (cisplatin, epirubicin, bleomycin) plus radiotherapy vs. radiotherapy alone in stage IV(> or = N2, M0) undifferentiated nasopharyngeal carcinoma: a positive effect on progression-free survival. International Nasopharynx Cancer Study Group. VUMCA I trial. *Int J Radiat Oncol Biol Phys* 35:463-9, 1996
- [12] Hareyama M, Sakata K, Shirato H, et al: A prospective, randomized trial comparing neoadjuvant chemotherapy with radiotherapy alone in patients with advanced nasopharyngeal carcinoma. *Cancer* 94:2217-23, 2002
- [13] Chua DT, Sham JS, Choy D, et al: Preliminary report of the Asian-Oceanian Clinical Oncology Association randomized trial comparing cisplatin and epirubicin followed by radiotherapy versus radiotherapy alone in the treatment of patients with locoregionally advanced nasopharyngeal carcinoma. Asian-Oceanian Clinical Oncology Association Nasopharynx Cancer Study Group. *Cancer* 83:2270-83, 1998
- [14] Ma J, Mai HQ, Hong MH, et al: Results of a prospective randomized trial comparing neoadjuvant chemotherapy plus radiotherapy with radiotherapy alone in patients with locoregionally advanced nasopharyngeal carcinoma. *J Clin Oncol* 19:1350-7, 2001
- [15] Chua DT, Ma J, Sham JS, et al: Long-term survival after cisplatin-based induction chemotherapy and radiotherapy for nasopharyngeal carcinoma: a pooled data analysis of two phase III trials. *J Clin Oncol* 23:1118-24, 2005
- [16] Chan AT, Teo PM, Ngan RK, et al: Concurrent chemotherapy-radiotherapy compared with radiotherapy alone in locoregionally advanced nasopharyngeal carcinoma: progression-free survival analysis of a phase III randomized trial. *J Clin Oncol* 20:2038-44, 2002

- [17] Chan AT, Leung SF, Ngan RK, et al: Overall survival after concurrent cisplatin-radiotherapy compared with radiotherapy alone in locoregionally advanced nasopharyngeal carcinoma. *J Natl Cancer Inst* 97:536-9, 2005
- [18] Lin JC, Jan JS, Hsu CY, et al: Phase III study of concurrent chemoradiotherapy versus radiotherapy alone for advanced nasopharyngeal carcinoma: positive effect on overall and progression-free survival. *J Clin Oncol* 21:631-7, 2003
- [19] Zhang L, Zhao C, Peng PJ, et al: Phase III study comparing standard radiotherapy with or without weekly oxaliplatin in treatment of locoregionally advanced nasopharyngeal carcinoma: preliminary results. *J Clin Oncol* 23:8461-8, 2005
- [20] Rossi A, Molinari R, Boracchi P, et al: Adjuvant chemotherapy with vincristine, cyclophosphamide, and doxorubicin after radiotherapy in local-regional nasopharyngeal cancer: results of a 4-year multicenter randomized study. *J Clin Oncol* 6:1401-10, 1988
- [21] Chi KH, Chang YC, Guo WY, et al: A phase III study of adjuvant chemotherapy in advanced nasopharyngeal carcinoma patients. *Int J Radiat Oncol Biol Phys* 52:1238-44, 2002
- [22] Chan AT, Teo PM, Leung TW, et al: A prospective randomized study of chemotherapy adjunctive to definitive radiotherapy in advanced nasopharyngeal carcinoma. *Int J Radiat Oncol Biol Phys* 33:569-77, 1995
- [23] Al-Sarraf M: Superiority of Five Year Survival with Chemo-Radiotherapy (CT-RT) vs Radiotherapy in Patients (Pts) with Locally Advanced Nasopharyngeal Cancer (NPC). Intergroup (0099) (SWOG 8892, RTOG 8817, ECOG 2388) Phase III Study: Final Report. *Proc Am Soc Clin Oncol* 20:abstract 905, 2001
- [24] Hoffman H CJ, Weber R, Ang K, Porter K, Langer C.: Changing patterns of practice in the management of nasopharynx carcinoma (NPC): Analysis of the National Cancer Database (NCDB). *ASCO Meeting Abstracts* Sept 3:5501, 2004
- [25] Wee J, Tan EH, Tai BC, et al: Randomized trial of radiotherapy versus concurrent chemoradiotherapy followed by adjuvant chemotherapy in patients with American Joint Committee on Cancer/International Union against cancer stage III and IV nasopharyngeal cancer of the endemic variety. *J Clin Oncol* 23:6730-8, 2005
- [26] Lee AW, Lau WH, Tung SY, et al: Preliminary results of a randomized study on therapeutic gain by concurrent chemotherapy for regionally-advanced nasopharyngeal carcinoma: NPC-9901 Trial by the Hong Kong Nasopharyngeal Cancer Study Group. *J Clin Oncol* 23:6966-75, 2005
- [27] Lee AW, Tung SY, Chua DT, et al: Randomized trial of radiotherapy plus concurrent-adjuvant chemotherapy vs radiotherapy alone for regionally advanced nasopharyngeal carcinoma. *J Natl Cancer Inst* 102:1188-98, 2010
- [28] Lee AW, Tung SY, Chan AT, et al: Preliminary results of a randomized study (NPC-9902 Trial) on therapeutic gain by concurrent chemotherapy and/or accelerated fractionation for locally advanced nasopharyngeal carcinoma. *Int J Radiat Oncol Biol Phys* 66:142-51, 2006
- [29] Chen Y, Liu MZ, Liang SB, et al: Preliminary results of a prospective randomized trial comparing concurrent chemoradiotherapy plus adjuvant chemotherapy with radiotherapy alone in patients with locoregionally advanced nasopharyngeal carcinoma in endemic regions of china. *Int J Radiat Oncol Biol Phys* 71:1356-64, 2008

- [30] Kwong DL, Sham JS, Au GK, et al: Concurrent and adjuvant chemotherapy for nasopharyngeal carcinoma: a factorial study. *J Clin Oncol* 22:2643-53, 2004
- [31] Ngan RK, Yiu HH, Lau WH, et al: Combination gemcitabine and cisplatin chemotherapy for metastatic or recurrent nasopharyngeal carcinoma: report of a phase II study. *Ann Oncol* 13:1252-8, 2002
- [32] Ma BB, Hui EP, Wong SC, et al: Multicenter phase II study of gemcitabine and oxaliplatin in advanced nasopharyngeal carcinoma--correlation with excision repair cross-complementing-1 polymorphisms. *Ann Oncol* 20:1854-9, 2009
- [33] Boussen H, Cvitkovic E, Wendling JL, et al: Chemotherapy of metastatic and/or recurrent undifferentiated nasopharyngeal carcinoma with cisplatin, bleomycin, and fluorouracil. *J Clin Oncol* 9:1675-81, 1991
- [34] Au E, Ang PT: A phase II trial of 5-fluorouracil and cisplatin in recurrent or metastatic nasopharyngeal carcinoma. *Ann Oncol* 5:87-9, 1994
- [35] Li YH, Wang FH, Jiang WQ, et al: Phase II study of capecitabine and cisplatin combination as first-line chemotherapy in Chinese patients with metastatic nasopharyngeal carcinoma. *Cancer Chemother Pharmacol* 62:539-44, 2008
- [36] Taamma A, Fandi A, Azli N, et al: Phase II trial of chemotherapy with 5-fluorouracil, bleomycin, epirubicin, and cisplatin for patients with locally advanced, metastatic, or recurrent undifferentiated carcinoma of the nasopharyngeal type. *Cancer* 86:1101-8, 1999
- [37] Yeo W, Leung TW, Chan AT, et al: A phase II study of combination paclitaxel and carboplatin in advanced nasopharyngeal carcinoma. *Eur J Cancer* 34:2027-31, 1998
- [38] Chua DT, Sham JS, Au GK: A phase II study of docetaxel and cisplatin as first-line chemotherapy in patients with metastatic nasopharyngeal carcinoma. *Oral Oncol* 41:589-95, 2005
- [39] Foo KF, Tan EH, Leong SS, et al: Gemcitabine in metastatic nasopharyngeal carcinoma of the undifferentiated type. *Ann Oncol* 13:150-6, 2002
- [40] Chua D, Wei WI, Sham JS, et al: Capecitabine monotherapy for recurrent and metastatic nasopharyngeal cancer. *Jpn J Clin Oncol* 38:244-9, 2008
- [41] Wang CC, Chang JY, Liu TW, et al: Phase II study of gemcitabine plus vinorelbine in the treatment of cisplatin-resistant nasopharyngeal carcinoma. *Head Neck* 28:74-80, 2006
- [42] Poon D, Chowbay B, Cheung YB, et al: Phase II study of irinotecan (CPT-11) as salvage therapy for advanced nasopharyngeal carcinoma. *Cancer* 103:576-81, 2005
- [43] Chua DT, Nicholls JM, Sham JS, et al: Prognostic value of epidermal growth factor receptor expression in patients with advanced stage nasopharyngeal carcinoma treated with induction chemotherapy and radiotherapy. *Int J Radiat Oncol Biol Phys* 59:11-20, 2004
- [44] Leong JL, Loh KS, Putti TC, et al: Epidermal growth factor receptor in undifferentiated carcinoma of the nasopharynx. *Laryngoscope* 114:153-7, 2004
- [45] Chan AT, Hsu MM, Goh BC, et al: Multicenter, phase II study of cetuximab in combination with carboplatin in patients with recurrent or metastatic nasopharyngeal carcinoma. *J Clin Oncol* 23:3568-76, 2005
- [46] Ma BB KM, Hui EP, King AD, Chan SL, Yu BK, Chiu SK, Lee FH, Chan AT.: A phase II study of concurrent cetuximab-cisplatin and intensity-modulated radiotherapy (IMRT) in locoregionally advanced nasopharyngeal carcinoma (NPC) with

- correlation using dynamic contrast-enhanced magnetic resonance imaging (DCE-MRI). JCO Abstract, 2008
- [47] Chua DT, Wei WI, Wong MP, et al: Phase II study of gefitinib for the treatment of recurrent and metastatic nasopharyngeal carcinoma. *Head Neck* 30:863-7, 2008
- [48] Ma B, Hui EP, King A, et al: A phase II study of patients with metastatic or locoregionally recurrent nasopharyngeal carcinoma and evaluation of plasma Epstein-Barr virus DNA as a biomarker of efficacy. *Cancer Chemother Pharmacol* 62:59-64, 2008
- [49] You B CE, Chin SF, Wang L, Jarvi A, Bharadwaj RR, Kamel-Reid S, Perez-Ordóñez B, Siu LL: A phase II trial of erlotinib after gemcitabine plus platinum-based chemotherapy in patients (pts) with recurrent and/or metastatic nasopharyngeal carcinoma (NPC) *European Journal of Cancer Supplements*, 2009
- [50] Elser C, Siu LL, Winquist E, et al: Phase II trial of sorafenib in patients with recurrent or metastatic squamous cell carcinoma of the head and neck or nasopharyngeal carcinoma. *J Clin Oncol* 25:3766-73, 2007
- [51] Krishna SM, James S, Balaram P: Expression of VEGF as prognosticator in primary nasopharyngeal cancer and its relation to EBV status. *Virus Res* 115:85-90, 2006
- [52] Montag M, Dyckhoff G, Lohr J, et al: Angiogenic growth factors in tissue homogenates of HNSCC: expression pattern, prognostic relevance, and interrelationships. *Cancer Sci* 100:1210-8, 2009
- [53] Segawa Y, Oda Y, Yamamoto H, et al: Close correlation between CXCR4 and VEGF expression and their prognostic implications in nasopharyngeal carcinoma. *Oncol Rep* 21:1197-202, 2009
- [54] Lee SP, Tierney RJ, Thomas WA, et al: Conserved CTL epitopes within EBV latent membrane protein 2: a potential target for CTL-based tumor therapy. *J Immunol* 158:3325-34, 1997
- [55] Straathof KC, Bollard CM, Popat U, et al: Treatment of nasopharyngeal carcinoma with Epstein-Barr virus--specific T lymphocytes. *Blood* 105:1898-904, 2005
- [56] Louis CU, Straathof K, Bollard CM, et al: Adoptive transfer of EBV-specific T cells results in sustained clinical responses in patients with locoregional nasopharyngeal carcinoma. *J Immunother* 33:983-90, 2010

IntechOpen

Head and Neck Cancer

Edited by Dr. Mark Agulnik

ISBN 978-953-51-0236-6

Hard cover, 440 pages

Publisher InTech

Published online 14, March, 2012

Published in print edition March, 2012

Head and Neck Cancer provides an interesting and comprehensive overview of all aspects of head and neck cancer including overviews of the disease, basic science aspects pertaining to the disease, diagnosis, treatment and outcomes for patients with this disease. The chapters written by world renowned experts cover the entire discipline of head and neck oncology and include discussions of regional disparity is, advances in basic science understanding, advances in her radiotherapy, chemotherapy and targeted agents as well as a focus on reconstruction, prostheses, and aspects of quality of life and health outcomes. The book is designed to be both practical and comprehensive for every physician treating his complex disease.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Jared Knol, Tiffany King and Mark Agulnik (2012). Nasopharyngeal Carcinoma: The Role for Chemotherapeutics and Targeted Agents, Head and Neck Cancer, Dr. Mark Agulnik (Ed.), ISBN: 978-953-51-0236-6, InTech, Available from: <http://www.intechopen.com/books/head-and-neck-cancer/nasopharyngeal-carcinoma-the-role-for-chemotherapeutics-and-targeted-agents>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](https://creativecommons.org/licenses/by/3.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen