We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Wildlife Cats Reproductive Biotechnology

Regina Celia Rodrigues da Paz

*Departamento de Ciências Básicas e Produção Animal,
Faculdade de Agronomia, Medicina Veterinária e Zootecnia,
Universidade Federal de Mato Grosso – DCBPA/FAMEV/UFMT
Brazil*

1. Introduction

The techniques used in assisted reproduction for wildlife or domestic animals are similar, and consist in the collection, evaluation, and cryopreservation of semen; artificial insemination (AI), *in vitro* fertilization (IVF), and embryo transfer (ET).

Considering that the maintenance of genetic diversity is reproduction dependent, reproduction technologies are important tools for the preservation of endangered species, through the development of methods to increase the fertility in these animals (Howard, 1993).

Several studies show that hormone therapies may induce ovulation and/or superovulation in wild animals (Pope, 2000). The artificial induction of ovulation and superovulation are important components in assisted reproduction techniques, especially in cats, because their ovulation is induced by external stimulations (Hamner et al., 1970).

According to The World Zoo Conservation Strategy ([IUDZG/CBSG], 1993), some of the many benefits from using assisted reproduction techniques in conservation management programs are:

1. To enable the exchange of genetic material between two or more zoos collaborating in the same program, including animals in captivity and in the wild. Semen transportation is economical and reduces the risks involved in animal transfers and consequently diseases transmission;
2. To enable reproduction in animals with physical and reproductive behavioral disabilities. This is important for animals that are representative of genetic lines that cannot be lost;
3. To enable rapid population growth when only a small founder population is available;
4. To assist in maintaining the ratio between males and females by selectively transplanting embryos of one sex;
5. To determine the number of offspring per individual;
6. To promote the formation of databases for gametes and embryos from species of interest.

The reproductive techniques used for domestic animals are gradually being used for zoo animals (Comizzoli et al., 2000; Dresser et al., 1986).

Artificial insemination has been conducted in different species of carnivores such as cougar (*Felis concolor*), leopard (*Panthera pardus saxicolor*), cheetah (*Acinonyx jubatus*), tiger (*Panthera tigris altaica*), ocelot (*Leopardus pardalis*), tigrina (*Leopardus tigrinus*), and jaguar (*Panthera onca*) (Dresser et al., 1982; Donoghue et al., 1993; Howard et al., 1992a; Jimenez et al., 1999; Moore et al., 1981; Moraes et al., 1997; Silva et al., 2000; Swanson et al., 1996a).

In vitro fertilization (IVF) has also been performed in captive wild cats such as tiger (*Panthera tigris altaica*), jaguar (*Panthera onca*), ocelot (*Leopardus pardalis*) and tigrina (*Leopardus tigrinus*) (Donoghue et al., 1990; Morato et al., 2000; Swanson & Brown, 2004).

The application of artificial reproductive methods in wild animals has not been successful, showing low reproductive rates. Some of the several reasons for these low reproductive rates are lack of knowledge on the species' physiology, poor sperm or oocytes quality, and difficulties in adapting from the methodologies used in experimental models.

Considering the zoos' limitation to maintain genetically viable populations of threatened species, the establishment of genetic banks containing semen, oocytes, embryos and cells emerge as a strategy to ensure the genetic diversity of populations (Lasley et al., 1994). The potential of assisted reproduction for endangered species should be emphasized by the possibility of semen and embryos cryopreservation, which are genetically valuable for the future populations.

Although there are some records related to the female reproductive physiology (Table 1) and semen characteristics for various Neotropical felids species (Table 2), there is a little knowledge about the fertilization ability using artificial methods (Table 3) in these animals. Studies about the application of assisted reproduction techniques in Neotropical felids showed limitations because of the lack of basic knowledge on the physiology of the species.

2. General reproductive characteristics in wildlife cats

The large number of felines and their wide geographical distribution determine many of the species' particularities, mainly in the reproductive aspects. The reproductive seasonality is one of the most variable aspect between species.

In domestic cats (*Felis catus*), the reproductive seasonality is related to photoperiod (Johnston et al., 1996; Shille et al., 1979; Tsuitsui & Stabenfeldt, 1996), whereas in wild animals, it is also related to high or low food supply during the seasons (Ewer, 1975).

The ovulation mechanism is another variable aspect among cat species. Studies evaluating serum hormone levels confirmed that some wild cats like tiger (*Panthera tigris*) (Seal et al., 1985), snow leopard (*Panthera uncia*) (Schmidt et al., 1993), jaguars (*Panthera onca*) (Wildt et al., 1979) and cougars (*Puma concolor*) (Bonney et al., 1981), have induced reflex ovulation, similar to the domestic cat (Johnston et al., 1996; Shille et al., 1979; Tsuitsui & Stabenfeldt, 1996).

However, female leopards (*Panthera pardus*) presented two ovulation mechanisms in two different situations. When kept isolated, they showed typical hormonal profile for ovulation reflex mechanism; but, when housed in pairs with another female, the ovulation was probably stimulated by physical contact (Schmidt et al., 1988).

Lionesses (*Panthera leo*), when isolated from the males, showed an ovulation pattern distinct from other cats. Schmidt et al. (1979), using serum hormone levels and corpus luteum

visualization, demonstrated that this species presents spontaneous ovulation in a higher frequency compared to what is described for other felines.

The genus *Leopardus* is polyestral and can cycle all year round (Morais et al., 1996; Moreira et al., 2001); the margay (*Leopardus wiedii*) is the only species in this genus presenting spontaneous ovulation (Moreira et al., 2001).

According to Tebet (1999), the estrous cycle in ocelots is characterized by the presence of serum estradiol peaks associated with relatively low levels of serum progesterone (<2.61 ng/mL). This demonstrates the polyestral characteristic of this species, similar to previously observed for other species such as cats (Shille al., 1979; Tsuitsui & Stabenfeldt, 1996; Verhage et al., 1976), tigers (Seal et al., 1985), and cheetahs (Brown et al., 1996).

Ocelot females that ovulated and were not fertilized showed a period of increased progesterone serum concentration and estrus inhibition (Tebet, 1999), called pseudopregnancy or diestrus, similar to domestic cats (Feldman & Nelson, 1996) and leopards (Schmidt et al., 1988).

Under an evolutionary analysis and among other factors, the process of spontaneous or induced ovulation may be related to the sociability of the species. Thus, solitary cats would require a longer estrus period, extended viability of oocytes, and extended time for the ovulation to occur, after the couple meet in the wild (Ewer, 1975).

The detection of fecal estrogens and progestins, through the analyses of fecal metabolites in domestic and wild cats such as the leopard cat (*Felis bengalensis*), cheetah (*Acinonyx jubatus*), clouded leopard (*Neofelis nebulosa*), and snow leopard (*Panthera uncia*) was successfully performed by Brown et al. (1994). Likewise, this methodology has been widely used to monitor ovarian function in Neotropical felines such as the ocelot and margay (Morais et al, 1996; Moreira et al., 2001).

In this context, noninvasive methods such as the quantification of fecal hormonal metabolites are increasingly being used in wildlife animals.

The estrus cycle, gestational time, and number of pups observed in different species of Neotropical wild cats are presented in Table 1.

Common Name	Scientific Name	Estrus (days)	Estrous Cycle (days)	Gestation (days)	Pups (n°)
Ocelot	<i>Leopardus pardalis</i>	4.63±0.63 ¹	16.5 ± 1.5 ² 18.4±1.6 ⁴	70-85 ⁶	1-4 ⁶
Tigrinus	<i>Leopardus tigrinus</i>	3.0-9.0 ⁵	15.8 ± 1.5 ² 16.7±1.3 ⁴	73-78 ⁶	1-4 ⁶
Margay	<i>Leopardus wiedii</i>	4.0-10.0 ⁵	19.5 ± 2.1 ² 17.6±1.5 ⁴	81-84 ⁶	1 ⁶
Geofroy's cat	<i>Oncifelis geoffroyi</i>	2.5±0.5 ¹	20.0 ⁵	72-76 ⁶	1-3 ⁶

Common Name	Scientific Name	Estrus (days)	Estrous Cycle (days)	Gestation (days)	Pups (n°)
Pallas cat	<i>Oncifelis colocolo</i>	-	-	80-85 ⁶	1-3 ⁶
Jaguarondi	<i>Herpaelurus yaguarundi</i>	3.17±0.75 ¹	53.63±2.41 ¹	72-75 ⁶	1-4 ⁶
Puma	<i>Puma concolor</i>	8.0 ⁵	23.0 ⁵	84-98 ⁶	1-6 ⁶
Jaguar	<i>Panthera onca</i>	12.0±1.0 ³	47.2±5.4 ³	90-111 ⁶	1-2 ⁶

Mellen 1989¹; Morais et al., 1996²; Morato & Paz, 2001³; Moreira et al., 2001⁴; Oliveira, 1994⁵; Oliveira & Cassaro, 1997⁶.

Table 1. Neotropical wildlife cats reproductive characteristics.

3. Sperm collection

Two methods are used for the collection of semen from wild animals: the first is the epididymis' semen collection from dead animals or after castration or vasectomy. The second is through an artificial vagina, electroejaculation, or digital manipulation.

3.1 Post mortem sperm collection

In the post mortem sperm collection, immediately after death or castration, the testes should be kept cold at 5°C (Howard, 1993). Generally, postmortem epididymidal spermatozoa remain viable for several days after the animal's death. However, this period depends on the species, the storage conditions of the testes, and methods used for sperm collection. Several techniques could be used including: totally cutting (homogenization) the cauda epididymis; washing out or aspiration (flushing) through sperm duct; making cuts (mincing) in the cauda epididymis and squeezing of content; or just squeezing (pressing). The first two techniques are the most frequently used (Maksudov et al., 2008).

For cats, the epididymis is washed and homogenized in HEPES medium plus Ham's F10, and the sperm is recovered after centrifugation. Live sperm can be recovered within the first twelve hours of the epididymis isolation using this technique (Howard, 1993). In vasectomized animals, the semen is collected by aspiration from the epididymis' tail using syringe and needle. The syringe should contain HEPES medium plus HAM'S F10 and the sperm is recovered after centrifugation.

Sperm recovered from the epididymis of cats are known to be motile, viable, and capable of penetrating oocytes (Goodrowe & Hay, 1993; Hay & Goodrowe, 1993). However, epididymal sperm have naturally more cytoplasmic droplets, which normally are lost during transport through the duct (Briz et al., 1995).

According to Tebet et al. (2006) there were no significant differences between the fresh or frozen-thawed domestic cat spermatozoa for the variables: sperm motility, plasma membrane integrity and morphology of electroejaculated and epididymal spermatozoa analyzed immediately after collection, and after freezing and thawing. Jewgenow (Jewgenow et al., 1997) reported motility of frozen-thawed epididymal sperm of lions (*Panthera leo*), tigers (*Panthera tigris*), leopards (*Panthera pardus*), pumas (*Felis concolor*) and jaguar (*Panthera onca*).

Epididymal spermatozoa were collected from immobilized adult male lion by caudal epididymectomy, cryopreserved, and used for IVF of *in vitro* matured lionesses' ova. The post-thawed motility ranged from 55-65%, and the percentages of fertilized ova were 12.7% and 11.5% for 30 and 36 hours of *in vitro* maturation, respectively (Bartels et al., 2000).

It has been shown, for several species, that the methods developed for ejaculated sperm are effective for freezing epididymal spermatozoa. However, the physiology of ejaculated and epididymal spermatozoa are different thus, it can be assumed that optimum methods of freezing and thawing may be different.

Postmortem material that can be retrieved in zoos usually belonged to the aged animals with different diseases or that died because of accidents (stress, fights). Moreover, some species display seasonal breeding. All these factors can influence on the spermatozoa's presence and quality in epididymis, at the collection time (Maksudov et al., 2008).

Analysis of different factors showed that the concurrency of death with breeding season has the strongest affect on the spermatozoa content in testicles of dead animals. The spermatozoa content and quality were almost equal in males that died of sickness (cancer, chronic cardiovascular or excretory system disorders) and in males that died in accidents (stress, fights). However, the quality of postmortem semen of animals that died of natural death is worse than that of animals that died accidentally (Maksudov et al., 2008).

The energy invested by a male with copulation is minimal compared to that of the female, for whom the costs continue throughout pregnancy and lactation. Therefore, it is a significant genetic advantage for cats to be reproductively active throughout the year, or at least to remain active well outside of the usual female breeding season (Spindler & Wildt, 1999).

This strategy is apparent in wild felids. The female Siberian tiger exhibits more estrual activity in January-June than at any other time of the year (Seal et al., 1985), however, the sperm quality is fairly consistent throughout the year (Byers et al., 1990). The quantity and quality of the sperm from some felids can vary when a distinctive female reproductive seasonality is known, as in the snow leopard (Johnston et al., 1994) and Pallas' cat (Swanson et al., 1996), the males remain reproductively active for longer periods than the females of the same species.

The collection of the postmortem sperm of recently dead animals belonging to endangered species can be of substantial importance, and therefore, the method of choice for the preservation of reproductive cells, in the wild, at zoos, and in national parks. The preservation and utilization of postmortem represent the last chance to obtain offspring from the dead males in cases of unexpected loss of valuable animals (Maksudov et al., 2008).

3.2 In vivo sperm collection

The collection of semen, through digital manipulation or using an artificial vagina are indicated because they promote a natural and normal ejaculation, however, these methods require intensive animal training. These techniques are effective for wild dogs, but not routinely used in wild cats.

The electroejaculation is the most used method in wild cats because it can be performed in anesthetized animals. However, one of the disadvantages of using this method in cats is the urine contamination of the semen. The urine contamination occurs when the voltage exceeds the minimum necessary level for ejaculation or when the electrode is positioned cranially. One alternative to minimize this problem would be the catheterization or cystocentesis before start procedure.

The Tiletamine-Zolazepam combination has been the most used anesthetic protocol for this procedure because it produces insignificant changes in the ejaculate. The ketamine hydrochloride and xylazine association in the same syringe has been used for semen collection in small cats, but according to Dooley et al. (1991), these drugs, in combination, seems to be related to retrograde ejaculation.

The electroejaculator should indicate both the voltage and amperage. The voltage should reach up to 12V, controlled by the command button, which should provide smooth control and gradual increase in power output.

The rectal bipolar electrode used for electroejaculation should have three longitudinal strips of copper. The copper strips must have a 0.4 cm apart and protruding approximately 0.2 cm. The electrode, previously lubricated with mineral oil, should be introduced into the rectum with the longitudinal strips ventrally positioned, applying light pressure to increase contact with the pelvic plexus region. The diameter of the electrode should be specific to each species (Table 2).

The electrical series follows a specific protocol with 80 stimuli divided into three series: 30 (series 1: 10 stimuli at 2, 3 and 4V), 30 (Series 2: 10 stimulations in 3, 4 and 5V) and 20 (series 3: 10 stimulations in 4 and 5V) (Howard 1993). In jaguars, the last series of stimuli can reach up to 6V to achieve ejaculation (Paz et al., 2000).

The stimulation cycle starts at 1 second from 0 voltage to the desired voltage, 2 to 3 seconds at the desired voltage, and 3 seconds returning to 0 voltage. An interval of 10 minutes should be used for resting between sets.

Before the start of the series, the penis must be exposed, examined, and washed with saline solution and gauze. The semen collection should be performed in plastic tubes that are maintained warm at 37° C water bath. For each series, the tubes should be replaced in order to avoid urine contamination. All ejaculates should be used; the total volume of semen is the sum of each ejaculate's volume (Figure 1).

4. Sperm evaluation

The appearance is the first evaluation of the semen: changes in color may be associated with diseases in the accessory organs and testes. The ejaculate's volume is the second aspect to be evaluated and must be determined immediately after collection. The volume provides information about the semen production in different species. The pH determination is important because it may indicate urine (acidic pH) or bacteria contamination (basic pH).

For sperm evaluation, an aliquot of the ejaculate is placed on a microscope slide warmed at 37°C, covered with a warmed glass coverslip, and examined at 400 X magnification. The

semen should be evaluated for motility and progressive sperm motility. Motility is expressed in percentages, with 0% being the value for immobile spermatozoa and 100% for maximum spermatozoa performance. The sperm type of movement is evaluated by the progressive sperm motility in scale from 0 to 5 (0 - no motility, 1 - poor lateral movement with some progression, 2 - moderate lateral movement with occasional progression, 3 - slow progression, 4 - progression, 5 - rapid progression) (Howard, 1993).

Fig. 1. Ocelot's penis exposed before semen collection; the penis' spines can be observed (left). Ocelot's electroejaculation procedure (right). Pictures: Regina Paz.

The sperm morphology and concentration can be evaluated by fixing an aliquot of semen (1:3 dilution) in a 10% formaldehyde saline solution or in a 2.5% glutaraldehyde solution after the preparation of samples in a humidified chamber.

For the determination of the sperm morphology, 200 cells per slide were counted at 1000 X magnification under light microscopy; the abnormalities were classified as primary or secondary defects expressed as percentages. According to the primary defects presented in the sample, the sperm can be classified as macrocephalic, microcephalic, bicephalic, pyriform head, rounded head, abnormal acrosome, abnormal midpiece, no midpiece, tightly coiled tail and biflagelate. According to the secondary defects presented in the sample, the sperm can be classified as bent midpiece with or without droplet, bent tail with or without droplet, and proximal or distal droplet.

The concentration can be evaluated using a Neubauer chamber at 400 X magnification under light microscopy. The volume, concentration, motility, vigor and abnormal sperm data in different species are presented in Table 2.

Species	Probe (cm)	N ^o Ejaculates	Volume (mL)	Concentr. (x10 ⁶ /mL)	Motility (%)	Vigour (0-5)	Normals (%)
<i>Ocelot</i> (<i>L. pardalis</i>)	1.0 ¹	5 ¹ 38 ²	0.3±0.1 ¹ 0.62±0.08 ²	28.0±17.0 ¹ 53.8±17.8 ²	72.0±12.5 ¹ 70.4±2.3 ²	4.0±0.5 ¹ -	80.8±0.9 ¹ 58.4±5.8 ²
<i>Tigrinus</i> (<i>L. tigrinus</i>)	1.0 ¹	18 ²	0.11±0.02 ²	78.5±33.8 ²	62.1±5.7 ²	-	35.6±6.0 ²
<i>Margay</i> (<i>L. wiedii</i>)	1.0 ¹	11 ¹ 27 ²	0.2±0.1 ¹ 0.31±0.05 ²	79.9±28.1 ¹ 14.2±5.3 ²	86.0±3.3 ¹ 62.8±5.3 ²	4.6±0.2 ¹ -	48.5±6.1 ¹ 39.5±7.7 ²
<i>Jaguarondi</i> (<i>H. yagouarundi</i>)	1.0 ¹	3 ¹ 21 ²	0.1±0.1 ¹ 0.08±0.02 ²	12.5±9.4 ¹ 7.2±4.0 ²	50.0±9.9 ¹ 57.8±2.5 ²	3.5±0.4 ¹ -	35.4±14.3 ¹ 25.7±4.6 ²
<i>Pampas cat</i> (<i>O. colocolo</i>)	1.0 ¹	5 ¹ 2 ²	0.3±0.1 ¹ 0.08±0.01 ²	10.8±5.7 ¹ 364.0±326.0 ²	36.7±6.6 ¹ 81.3±6.3 ²	2.8±0.2 ¹ -	65.9±23.8 ¹ 56.5±0.5 ²
<i>Geofroy's cat</i> (<i>O. geofroy</i>)	1.0 ¹	8 ¹ 24 ²	0.2±0.1 ¹ 0.21±0.03 ²	300.0±233.2 ¹ 66.5±24.4 ²	73.0±4.4 ¹ 64.0±4.7 ²	4.0±0.3 ¹ -	29.0±11.5 ¹ 46.9±5.0 ²
<i>Puma</i> (<i>P. concolor</i>)	1.6 ¹	12 ¹	2.8±0.5 ¹	20.2±4.7 ¹	52.0±8.0 ¹	3.5±0.2 ¹	23.4±3.7 ¹
<i>Jaguar</i> (<i>P. onca</i>)	3.0 ¹ 2.3 ³	5 ¹ 38 ³	2.7±0.6 ¹ 5.7±1.71 ³	12.0±1.9 ¹ 13.16±10.76 ³	82.0±5.8 ¹ 56.9±9.35 ³	4.1±0.3 ¹ 3.02±0.77 ³	58.2±11.1 ¹ 65.73±6.7 ³

Howard, 1993¹; Morais, 2001²; Paz et al., 2000³;

Table 2. Neotropical wildlife cats seminal characteristics.

The poor semen quality in carnivores may be related to the nutritional status of the animals. Rodrigues da Paz et al. (2006), studying the reproduction of jaguars, observed a positive correlation between the improvement in the semen quality and the decrease of primary defects, after diet supplementation with vitamins and minerals. Ocelots, tigrinus and margays showed an increase in the number of ejaculates and 20-30% improvement related to sperm defects after receiving vitamin and mineral supplementation (Morais, 2001).

The seminal plasma constituents compromise the sperm viability in some species. Thus, washing the ejaculates in culture media (HEPES, HAM'SF-10) by centrifugation is efficient in removing the seminal plasma, which could contain bacteria and other undesirable microorganisms, especially when the semen will be used for intrauterine artificial insemination.

5. Sperm cryopreservation

The semen cryopreservation procedures should be initiated only after the centrifugation at 300g for 10 minutes in culture medium or HEPES HAM'SF-10. This procedure is essential for eliminating microorganisms and seminal plasma remove.

The "Double Step" cryopreservation method, using glycerol as cryoprotectant, is in general, used for the semen of most carnivores. This method use two fractions: fraction A containing nutritional constituents and antibiotics; and fraction B containing nutritional constituents, antibiotics, and the cryoprotectant. The PDV medium is used for the semen cryopreservation of the majority of wildlife cats. The fraction A contains 20% egg yolk, 11% lactose, 1000 IU penicillin/mL, 1000 mg streptomycin/mL; the fraction B contains 20% egg yolk, 11% lactose, 8% glycerol, 1000 IU penicillin/mL and 1000 mg streptomycin/mL.

After being removed from liquid nitrogen, the semen straws should be immediately thawed for 1 min in waterbath at 37°C, evaluated for total motility (%) and progressive sperm motility (scale, 0-5) before use.

The first step in the process of freezing semen is the removal of the supernatant after centrifugation of the semen collected in HAM'S F-10 or HEPES culture media, and the subsequent resuspension of the pellet in PDV fraction A at 37 °C. This mixture should be kept in the refrigerator for 2 hours followed by a subsequent slow addition of the PDV fraction B. The material is then transferred to cooled 0.25 mL straws and kept in the refrigerator for 30 minutes. Afterwards, each straw is placed in liquid nitrogen vapor for 20 min, immersed in liquid nitrogen, transferred to the racks, and loaded into the canisters for long-term liquid nitrogen storage at -196°C.

The straws, racks and canister identification are extremely important, being the determining factor for the germplasm bank establishment and successful operation. The material collected might be extremely valuable for populations in the future and the safety use of this material depends on the correct identification.

The straws identification must contain the animal species (scientific name), tattoo or microchip number, the institution to which the animal belongs, and the date. For free-living animals, the straws must contain the species, the location where the animal was captured, and the date. The racks may be identified by numbers or if applicable, by species. A registry, which can be computerized, is essential to record all of the straws, racks, and canisters, thereby facilitating the location of the material.

The reasons for the poor quality of wildlife semen after thawing are still unknown and involve a range of information and specific characteristics for each species, which are also not yet clearly understood. New tests with different protocols and different cryoprotectors for each species of interest are required in order to maximize the spermatozoa viability after cryopreservation procedures.

6. Ovarian activity induction and superovulation

The currently used ovarian stimulation and superovulation protocols require injections of exogenous gonadotropins, which consist of large complexes of glycoproteins. Equine Chorionic Gonadotropin (eCG) and Human Chorionic Gonadotrophin (hCG) are frequently used due to their long half-life in circulation (24-48h) and good ovarian response with a single application. Other hormones used are the porcine Follicle Stimulating Hormone (pFSH) and porcine Luteinizing Hormone (pLH), these hormones are characterized as short half-life (~ 2h) gonadotropins, therefore, they present the disadvantage of requiring multiple applications to produce a good ovarian response (Crichton et al., 2003; Dresser et al., 1988; Pope, 2000; Wildt et al., 1981).

Studies on wild cats report the use of eCG/hCG in combination, mainly to avoid the stress associated with multiple injections of FSH (Roth et al., 1997). However, the use of porcine FSH/LH determined equivalent number of oocytes compared to the established protocol for eCG/hCG used in tigers (*Panthera tigris*) (Crichton et al., 2000), ocelot (*Leopardus pardalis*) and tigrinas (*Leopardus tigrinus*) (Paz et al., 2005, 2006), demonstrating that the stress caused by daily injections did not influence the ovarian response.

The eCG/hCG combination has been used successfully in tigers (*Panthera tigris*), cheetahs (*Acinonyx jubatus*), clouded leopards (*Neofelis nebulosa*), pumas (*Puma concolor*), ocelots (*Leopardus pardalis*) and tigrinas (*Leopardus tigrinus*) (Barone et al., 1994; Donoghue, 1993; Donoghue et al., 1990; Howard, 1992b; Moore et al., 1981; Moraes et al., 1997; Morato et al., 2000; Swanson, 1996a). The FSH and hCG combination was used successfully in the Indian desert leopard (*Felis sylvestris ornata*) (Pope et al., 1989), and the pFSH and pLH combination in tigers (Crichton et al., 2000, 2003) jaguars (*Panthera onca*) (Morato et al., 2000), ocelots (*Leopardus pardalis*) and tigrinas (*Leopardus tigrinus*) (Paz et al., 2005, 2006).

Swanson et al. (1995, 1996a) suggest that the repeated administration of exogenous gonadotropins, within short time intervals is a problem because it causes a reduction in the ovarian stimulation which is immunologically mediated. The repeated administration of exogenous gonadotropins has been associated with the production of neutralizing immunoglobulin, which prevents the ovarian response to superovulation protocols.

Alternating gonadotropins regimens in sequential treatments are indicated because of variable immunoglobulin affinities to different exogenous gonadotropins (Maurer et al., 1968; Swanson et al., 1995).

Ocelots and tigrinas treated four to six times, at 4-month intervals, with alternating exogenous gonadotropin regimens (eCG/hCG and pFSH/pLH) did not show a reduction in ovarian response (total follicles and Corpora Lutea), oocyte maturation or exogenous gonadotropins antibodies production over time (Paz et al., 2005, 2006). The findings suggest that, these endangered cat species may be managed intensively with the use of alternating exogenous gonadotropin regimens in assisted reproduction procedures without compromising ovarian responsiveness to these hormones.

Specie	Procedure	Treatment 1		Treatment 2	
		ECG (UI)	hCG (UI)	pFSH (UI)	pLH (UI)
Ocelot (<i>Leopardus pardalis</i>)	AI ⁶	400 ⁶	200 ⁶	-	-
	IVF ^{4,5}	500 ^{4,5}	225 ^{4,5}	50 ^{4,5}	20 ^{4,5}
Tigrinus (<i>Leopardus tigrinus</i>)	AI ^{4,5}	75 ^{4,5}	100 ^{4,5}	30 ^{4,5}	10 ^{4,5}
	IVF ⁶	200 ⁶	150 ⁶	-	-
Gato mourisco (<i>H. yagouarundi</i>)	AI ⁷	100 ⁷	75 ⁷	-	-
	IVF ⁷	200 ⁷	150 ⁷	-	-
Puma (<i>Puma concolor</i>)	AI ¹	200 ¹	100 ¹	-	-
	IVF	-	-	-	-
Jaguar (<i>Panthera onca</i>)	AI ²	200 ²	150 ²	-	-
	IVF ³	-	-	50 ³	25 ³

Barone et al., 1994¹; Jimenez et al., 1999²; Morato et al., 2000³; Paz et al., 2005⁴; Paz et al., 2006⁵; Swanson et al., 1996a⁶; Swanson (Personal Communication)⁷.

Table 3. Ovarian stimulation with exogenous gonadotropins used in wildlife cats reproduction (AI= Artificial Insemination and IVF= *In vitro* fertilization).

The exogenous gonadotropins dosage used for ovarian stimulation is another important factor to the fertilization rate and subsequent embryonic development (Donoghue et al., 1993). Species with similar size and weight may require varying dosages, possibly because

they have different sensitivities to exogenous gonadotropins (Roth et al., 1997; Swanson et al., 1996b). The nutritional status of the animal also influences the fertilization success rate. Swanson et al. (2002a), studying ocelots and tigrinas in Brazil, observed better quality of oocytes and increase fertilization rates after supplementing the diet with vitamins and minerals.

7. Artificial insemination and oocytes collection

The rate of artificial insemination success in carnivores is influenced by the localization of the semen deposition. Non-surgical methods of semen deposition in the vagina has shown inferior results compared with the surgical method, with semen deposition directly into the uterus. This can be explained by the chemical restraint need in wild animals, with the anesthesia compromising the sperm transportation in non-surgically insemination (Howard, 1993).

The artificial insemination success with semen deposition in the uterine horn is described in several species of wild cats such as puma (*Puma concolor*) (Moore et al., 1981); leopard (*Panthera pardus saxicolor*) (Dresser et al., 1982); cheetah (*Acinonyx jubatus*) (Howard, 1992); tiger (*Panthera tigris altaica*) (Donoghue et al., 1993); ocelot (*Leopardus pardalis*) (Swanson, 1996a) and tigrina (*Leopardus tigrinus*) (Moraes et al., 1997).

Artificial insemination using video-laparoscopy technique has been developed for the semen deposition directly into the uterine horn, close to the oviduct where fertilization occurs, in addition to being a less invasive method. In this procedure, the ovaries and uterine horns can be accessed and evaluated for thickness, consistency and color in all species. In cats, the ovaries are easily observed, facilitating the counting and characterization of pre-ovulatory follicles (brighter small elevated areas) and post-ovulatory corpus luteum (yellow-red area).

According to ovarian stimulation protocols, the animals should be inseminated within 24 to 48 hours after hCG or pLH administration, or after the ovulation process. Inhalatory anesthesia is necessary to perform this procedure (Figure 2).

Fig. 2. Anesthesia with isoflurane gas mask (left) and intubation (right). Pictures: Regina Paz.

The anesthetized cats should be secured in dorsal recumbency with the use of leg ties on a tilting surgical table, and the abdominal region of each female should be clipped and prepped

with alternating applications of Betadine scrub and alcohol. A pneumoperitoneum should be created by means of CO₂ gas introduced through a *Verres* needle inserted transcutaneously into the central abdominal cavity. A 7-mm-diameter laparoscope should be inserted through a 1cm skin incision slightly cranial to the umbilicus. The ovaries could be manipulated with the *Verres* needle probe, and each ovary should be closely examined to determine the number of mature follicles (≥ 2 mm diameter), recently-formed corpora lutea, and corpora albicans.

To stabilize the uterine horn, where the cannula will be introduced to deposit the semen, grasping forceps is inserted laterally, 4 to 5 cm of the umbilicus. This procedure maintains the uterus close to the abdominal wall. The horn to be inseminated is the ovary that shows the corpus luteum after ovulation. The procedure should be performed in both uterine horns if both ovaries present the corpus luteum.

For the semen deposition, the uterine horn is cannulated using a 20G sterile needle catheter inserted through the abdominal cavity, near the uterine lumen. As soon as the needle pierces the uterine horn, it is removed, keeping the catheter in place. Inside the catheter, a sterile polypropylene tube must be inserted, which will be connected to a syringe containing the semen. (Figure 3).

The intrauterine artificial insemination by laparoscopy is less invasive because the semen deposition occurs directly in the uterine horn without laparotomy. This methodology resulted in a 46.2% increase in cheetah's pregnancy rates (Howard, 1992).

Similarly, Donoghue et al. (1993) reported the first birth of a tiger cub (*Panthera tigris altaica*) in Siberian Tiger Species Survival Plan (SSP Program) after intrauterine insemination by video-laparoscopy in females stimulated with eCG and hCG. This result demonstrates the importance in using assisted reproduction methods in the production of genetically viable population with recommended breeding by a management program.

Fig. 3. Intrauterine insemination set: 1mL syringe, polypropylene tube and 22G needle (left). Connected set (right). Pictures: Regina Paz.

The procedures for oocyte retrieval in *in vitro* fertilization are performed using the laparoscopy technique. The ovaries' visualization and the oocyte retrieval in carnivores are species-specific. Cats' ovaries are easily accessed and the follicles easily aspirated.

General anesthesia is needed to perform this procedure; isoflurane inhalation anesthesia is generally used. After anesthetized, the animals are placed in the supine position (45 degrees) and pneumoperitoneum is created with the *Verres* needle, which can be coupled to a CO₂ gas automatic insufflator or a manual pump.

The endoscope is placed near the umbilicus for the evaluation of ovaries and observation of the follicles. Only mature follicles, larger than 2 mm, should be aspirated. The *Verres* needle is used for the follicles' size determination and for the maintenance of the ovary in an adequate position, near to the abdominal wall.

The mature follicles are aspirated with a 22G needle attached to polypropylene tubing connected to a sterile collection tube (15mL) containing M199 culture medium and heparin, which is attached to a vacuum aspiration pump (Figure 4). After the collection, the oocytes are placed in petri dishes with culture medium and observed by stereomicroscopy at 400 X magnification for their classification.

The oocytes from carnivores are dark and contain lipidic drops. The maturation status is characterized in I, II and III according morphological aspects. Oocytes I are of excellent quality, characterized by a uniformly dark cytoplasm, nucleus with a distinct corona radiata, and an expansive cumulus cell mass. Oocytes II are of regular quality, characterized by a non-uniformly cytoplasm, nucleus with an indistinct corona radiata, and a non-expansive cumulus cell mass. Oocytes III are degenerated and characterized by an abnormal cytoplasm, nucleus without corona radiata or cumulus cell mass (Goodrowe et al., 1988; Johnston et al., 1989).

Fig. 4. Follicular aspiration using video-laparoscopy and *Verres* needle in ocelot (left). Aspiration follicular system (right). Pictures: Regina Paz.

8. *In vitro* fertilização and embryo transfer

The *in vitro* fertilization technique has been applied in wild animals after follicular aspiration using laparoscopy technique, oocyte retrieval post-mortem, or after ovariectomy. According Swanson (1998), oocytes collected from ovaries can be refrigerated at 5°C for 24 hours without maturation and changes in the fertilization potential.

Oocytes recovery from refrigerated ovaries can be achieved using the follicular aspiration technique with a syringe and needle, or through ovary laceration and oocyte harvest using stereomicroscopy. The second technique is used in small animals, which present ovaries with small diameters because the follicle aspiration would be difficult. The M199 culture medium is used for follicular aspiration and ovaries laceration.

However, laparoscopic follicular aspiration is the most used oocyte retrieval technique for IVF, which should be preceded by hormonal treatment. According Howard (1999), the

treatment with exogenous gonadotropins and laparoscopy are the basic requirements for ovarian stimulation and oocyte retrieval for IVF procedures.

The immature oocytes collected ($\cong 60\%$) in cats become mature in 24-32 hours in culture media. These about 70% are fertilized, however, only a small percentage, from 20 to 30% develops in blastocysts (Johnston et al., 1989).

Donoghue et al. (1990) reported the birth of the first wild cat from IVF after an embryo transfer. Tiger cubs (*Panthera tigris*) were produced *in vitro* using excellent quality embryos, containing two to four cells, and surgically transferred to the oviducts of two females. Pregnancy was successful in one of these females and three kittens were born after 107 days.

The first embryos produced by IVF in Brazil were jaguar (*Panthera onca*) embryos. The ovarian stimulation with pFSH/LH produced $\cong 25$ follicles/female ($> 80\%$), however, despite the recovery of high quality oocytes, the fertilization percentages were low ($> 25\%$) (Morato et al., 2000).

In a Project involving the São Paulo University/Brazil (USP), the Mata Ciliar Association/Brazil (AMC) and the Cincinnati Zoo/USA (CREW), 128 ocelot (*Leopardus pardalis*) and tigrina (*Leopardus tigrinus*) embryos were produced. The animals were treated with eCG/hCG and produced $\cong 10$ follicles/female. Follicles were aspirated by laparoscopy and 7-9 of excellent quality oocytes/female were recovered. Of these, 60% were fertilized *in vitro*, resulting in 76 ocelot and 52 tigrina embryos (Swanson & Brown, 2004). Two ocelots became pregnant with normal pregnancy development and birth, one from the Cincinnati Zoo/USA and another from the Sao Bernardo do Campo Zoo/Brazil. However, only the offspring born to the American female survived after birth (Swanson, 2002b).

Among all the Neotropical cats, the ocelot is the only species that produced offspring after the transfer of frozen embryos (Swanson, 2001, 2002b). These births were the result of a cooperative effort between Brazil and the USA for the development of adequate management programs for this species in captivity, and for the establishment of genetically viable populations between the Brazilian and American populations.

Based on the percentage of successful embryonic cleavage after thawing domestic cat embryos, which is less than 70% (Pope, 2000), the difficulties to develop feline embryos used post-freezing is recognized. The difficulties may be related to inappropriate timing for the embryo implantation or fetal survival (Swanson & Brown, 2004) and not only related to the quality of the embryos. Thus, it is necessary to detect the female natural receptor estrus in order to perform the transfer of thawed embryos. This determination is achieved through measurement of fecal steroid levels according to the enzyme-immunoassay (EIA) technique for the fecal estrogens metabolites.

After the determination of the natural estrus, the ovulation should be induced with GnRH (Gonadotropin Releasing Hormone) and the embryos should be transferred by video-laparoscopy directly in the ostium of the oviduct.

9. Transmission of reproductive diseases

The methods used for assisted reproduction in wild animals should be free of contamination or diseases, therefore, some measures must be taken to avoid compromising the procedures. The semen centrifugation and seminal plasma removal are essential for artificial insemination with fresh or frozen semen and *in vitro* fertilization procedures.

The methods for processing semen without centrifugation and removal of the seminal plasma were responsible for the development of pyometra in 40% of inseminated domestic cats, regardless of the semen being diluted in culture medium containing antibiotics penicillin and streptomycin (Howard, 1993). It is believed that the donors might carry bacteria in their normal flora, eg *E. coli*, which would cause infection in the females.

In a study conducted by Paz et al. (1999), aiming at determining preputial microbiota in nine adult male jaguars (*Panthera onca*), the most frequently observed microorganism were *Staphylococcus sp* (40%), followed by *Streptococcus sp* (30%), *Escherichia coli* (20%) and *Corynebacterium sp* (10%).

The preputial microbiota in the genus *Leopardus* was assessed by Guido et al. (2000), and the results were *Escherichia coli* (40%), *Proteus rettgeri* (40%) and *Yersinia pseudotuberculosis* (20%) in tigrina (*Leopardus tigrinus*) (n = 5), *Staphylococcus sp* (42.9%), *Escherichia coli* (28.5%), *Streptococcus sp* (14.3%), *Staphylococcus sp* + *Streptococcus sp* (14.3%) in ocelots (*Leopardus pardalis*) (n = 6) and only *Staphylococcus sp* in margay (*Leopardus wieddi*) (n = 1).

The feline immunodeficiency virus is present in the semen of domestic cats and can be transmitted to females by AI (Jordan et al., 1995, 1996). This aspect should be taken into consideration during assisted reproduction procedures performed in wild cats.

In addition, a reproductive evaluation and clinical examination should be performed in wild cats before the animal inclusion in the management programs using assisted reproduction.

10. References

- Bartels, P.; Lubbe, K.; Killian, L.; Friedmann, Y.; Van Dyk, G. & Mortimer, D. (2000). In vitro maturation and fertilization of lion (*Panthera onca*) oocytes using frozen-thawed epididymal spermatozoa recovered by cauda epididymectomy of an immobilized lion. *Theriogenology*, Vol. 53, pp. 325.
- Barone, M. A.; Wildt, D. E.; Byers, A. P.; Roelke, M. E.; Glass, C. M. & Howard, J. G. (1994). Gonadotrophin dose and timing of anaesthesia for laparoscopic artificial insemination in the puma (*Felis concolor*). *Journal of Reproduction and Fertility*, Vol.101, pp. 103-108.
- Bonney, R. C.; Moore, H. D. M. & Jones, D. M. (1981). Plasma concentrations of oestradiol-17 β and progesterone, and laparoscopic observations of the ovary in the puma (*Felis concolor*) during oestrus, pseudopregnancy and pregnancy. *Journal of Reproduction and Fertility*, Vol.63, pp. 523-532.
- Briz, M. D.; Bonet, S.; Pinart, B.; Egozcue, J. & Campos, R. (1995). Comparative study of boar sperm coming from the caput, corpus, and cauda regions of the epididymis. *Journal of Andrology*, Vol. 16, pp. 175-188.
- Brown, J. L.; Wasser, S. K.; Wildt, D.E. & Graham, L. H. (1994). Comparative aspects of steroid hormone metabolism and ovarian activity in felids, measured noninvasively in feces. *Biology of Reproduction*, Vol.51, pp. 766-86.
- Brown, J. L.; Terio, K. A. & Graham, L. H. (1996). Fecal androgen metabolite analysis for non invasive monitoring of testicular steroidogenic activity in felids. *Zoo Biology*, Vol.15, pp. 425-434.
- Byers, A. P.; Hunter, A. G.; Seal, E. S.; Graham, E.F. & Tilson, R.L. (1990). Effect of season on seminal traits and serum hormone concentration in male Siberian tiger (*Panthera tigris*). *Journal of Reproduction and Fertility*, Vol. 90, pp. 119-125.

- Comizzoli, P.; Mermillod, P. & Mauget, R. (2000). Reproductive biotechnologies for endangered mammalian species. *Reproduction Nutrition Development*, Vol.40, pp. 493-504.
- Crichton, E. G.; Armstrong, D. L.; Vajta, G.; Pope, C. E. & Loskutoff, N. M. (2000). Development competence in vitro of embryo produce from siberian tigers (*Panthera tigris altaica*) cryopreserved by controlled rate freezing versus vitrification. *Theriogenology*, Vol.53, No.1, pp.328. Abstract.
- Crichton, E. G.; Bedows, E.; Miller-Lindholm, A. K.; Baldwin, D. M.; Armstrong, D. L.; Graham, L. H.; Ford, J. J.; Gjorret, J. O.; Hyttel, P.; Pope, C. E.; Vajta, G. & Loskutoff, N. M. (2003). Efficacy of porcine gonadotropins for repeated stimulation of ovarian activity for oocyte retrieval and in vitro embryo production and cryopreservation in siberian tigers (*Panthera tigris altaica*). *Biology of Reproduction*, Vol.68, pp. 105-113.
- Donoghue, A. M.; Johnston, L. A.; Seal, U. S.; Armstrong, D. L.; Tilson, R. L.; Wolf, P.; Petrini, L. G.; Simmons, T.; Groos, T. & Wildt, D. E. (1990). In vitro fertilization and embryo development in vitro and in vivo in the tiger (*Panthera tigris*). *Biology of Reproduction*, Vol.43, pp. 733-44.
- Donoghue, A. M.; Johnston, L. A.; Armstrong, D. L.; Simmons, L. G. & Wildt, D. E. (1993). Birth of a siberian tiger cub (*Panthera tigris altaica*) following laparoscopic intrauterine artificial insemination. *Journal of Zoo and Wildlife Medicine*, Vol.24, No.2, pp. 185-89.
- Dooley, M. P.; Pineda, M. H; Hopper, J. G. & Hsu, W. H. (1991). Retrograde flow of spermatozoa into the urinary bladder of cats during eletroejaculation, collection of semen with an artificial vagina, and mating. *American Journal Veterinary Research*, Vol.52, No.5, pp. 687-91.
- Dresser, B. L.; Kramer, L.; Reece, B. & Russel, P. T. (1982). Induction of ovulation and successful artificial insemination in a Persian Leopard (*Panthera pardus saxicolor*). *Zoo Biology*, Vol.1, pp. 55-57.
- Dresser, B. L.; Sehlhorst, C. S.; Keller, G.; Kramer, L. W. & Reece, R. W. (1986). Artificial insemination and embryo transfer in the felidae. *Proceedings of International Symposium for World Conservation Strategies for Tigers*, pp. 4-18, Cincinnati, Ohio, USA.
- Dresser, B. L.; Gelwicks, E. J.; Wachs, K. B. & Keller, G. L. (1988). First successful transfer of cryopreserved feline (*Felis catus*) embryos resulting in live offspring. *The Journal of Experimental Zoology*, Vol.246, pp. 180-186.
- Ewer, R. F. (1975). The evolution of mating system in the felidae. In: *The world's cats: biology, behavior and management of reproduction*. Eaton, R. L. pp. 11-20, London: Unimark Publishing.
- Feldman, E. C. & Nelson, R. W. (1996). *Canine and Feline Endocrinology and Reproduction*. pp. 529-46, 2 ed, Philadelphia: W.B. Saunders.
- Goodrowe, K. L.; Hay, M. (1993). Characteristics and zona binding ability of fresh and cooled domestic cat epididymal spermatozoa. *Theriogenology*, Vol. 40, pp. 967-75.
- Goodrowe, K. L.; Wall, R. J.; O'Brien, S. J.; Schmidt, P. M. & Wildt, D. E. (1988). Developmental competence of domestic cat follicular oocytes after fertilization in vitro. *Biology of Reproduction*, Vol.39, pp. 355-372.
- Guido, M. C.; Paz, R. C. R.; Costa, E. O.; Zuge, R. M.; Benites, N. R.; Barnabe, V. H. & Barnabe, R. C. , (2000). Microbiota prepucial e vaginal de felinos neotropicais mantidos em cativeiro. *Proceedings of Combravet XVII Congresso Brasileiro de Medicina Veterinária*, pp.03, Águas de Lindóia, São Paulo, Brasil, 2000.

- Hamner, C. E.; Jennings, L. L. & Sojka, N. J. (1970). Cat (*Felis catus* L.) spermatozoa require capacitation. *Journal Reproduction and Fertility*, Vol.23, pp. 477-480.
- Hay, M. A.; Goodrowe, K. L. (1993). Comparative cryopreservation and capacitation of spermatozoa from epididymides and vasa deferentia of the domestic cat. *Journal Reproduction and Fertility Supplement*, Vol. 47, p. 297-30.
- Howard, J.; Donoghue, A. M.; Barone, M. A.; Goodrowe, K. L.; Blumer, E. S.; Snodgrass, B. S.; Sarnes, D.; Michael, T.; Bush, M. & Wildt, D. E. (1992). Successful induction of ovarian activity and laparoscopic intrauterine artificial insemination in the cheetah (*Acinonyx jubatus*). *Journal of Zoo and Wildlife Medicine*, Vol.23, No.3, pp. 288-300a.
- Howard, J. (1992). Embryogenesis in conservation biology or how to make an endangered species embryo. *Theriogenology*, Vol.37, No.1 b.
- Howard, J. G. (1993). Semen collection and analysis in carnivores. In: *Zoo & Wild Animal Medicine Current Therapy*, Fowler, M. E, pp.390-399, 3 ed. Philadelphia: W.B. Saunders.
- Howard J. G. (1999). Assisted reproductive techniques in nondomestic carnivores. In: *Zoo & Wild Animal Medicine*, Fowler, M.E. & Miller, R.E.. pp.449-457, 4 ed., Philadelphia: W.B. Saunders.
- IUDZG/CBSG. (1993). *The word zoo conservation strategy: the role of the zoos and aquaria of the world in global conservation*. Chicago: Zoological Society, 76 p.
- Jewgenow, K.; Blottner, S.; Lengwinat, T. & Meyer, H.H.D. (1997). New methods for gametes rescue from gonads of nondomestic felids. *Journal of Reproduction and Fertility Supplement*, Vol. 51, pp. 33-39.
- Jimenez, T. G.; Zuge, R.; Paz, R. C. R.; López, J. E. & Crudeli, G. A. (1999). Sincronización de celo e inseminación artificial por video laparoscopia en yaguararé (*Panthera onca*) en cautiverio. *Comunicaciones Científicas y Tecnológicas*, Vol.4, pp. 67-70.
- Johnston, L.A.; O'Brien, S. J. & Wildt, D. E. (1989). In vitro maturation and fertilization of domestic cat follicular oocytes. *Gamete Research*, Vol. 24, No.3, pp. 343-56.
- Johnston, L. A.; Armstrong, D. L. & Brown, J. L. (1994). Seasonal effects on seminal and endocrine traits in the captive snow leopard (*Panthera uncia*). *Journal of Reproduction and Fertility*, Vol. 102, pp. 229-236.
- Johnston, S. D.; Root, M. V. & Olson, P. N. S. (1996). Ovarian and testicular functions in the domestic cat: clinical management of spontaneous reproductive disease. *Animal Reproduction Science*, Vol.42, pp. 261-274.
- Jordan, H. L.; Howard, J. G. & Tompkins, W. A. (1995). Detection of feline immunodeficiency virus in semen from seropositive domestic cats (*Felis catus*). *Journal Virology*, Vol.69, pp. 7328-7333.
- Jordan, H. L.; Howard, J. G & Sellon, R. K. (1996). Transmission of feline immunodeficiency virus in domestic cats via artificial insemination. *Journal Virology*, Vol.70, pp. 8224-8228.
- Lasley, B. L.; Loskutoff, N. M. & Anderson, G. B. (1994). The limitation of conventional breeding programs and the need and promise of assisted reproduction in non-domestic species. *Theriogenology*, Vol.41, pp. 119-132.
- Maksudov GYu, Shishova NV, & Katkov II. 2009. In the Cycle of Life: Cryopreservation of Post-Mortem Sperm as a Valuable Source in Restoration of Rare and Endangered Species. In: *Endangered Species: New Research*. Eds.: A.M. Columbus and L. Kuznetsov. NOVA Science Publishers, Hauppauge NY, Ch. 8, pp 189-240.
- Maurer, R. R.; Hunt, W. L. & Foote, R. H. (1968). Repeated superovulation following administration of exogenous gonadotrophins in dutch-belted rabbits. *Journal Reproduction and Fertility*, Vol.15, pp.93-102.

- Mellen, J.D. 1989. *Reproductive behavior of small captive exotic cats (Felis sp)*. (Doctoral thesis) University of California. Davis/CA.
- Moore, H. D. M.; Bonney, R. C. & Jones, D. M. (1981). Successful induced ovulation and artificial insemination in the puma (*Felis concolor*). *Veterinary Record*, Vol.108, pp.282-83.
- Moraes, W.; Morais, R.; Moreira, N.; Lacerda, O.; Gomes, M. L. F.; Mucciolo, R. G. & Swanson, W. F. (1997). Successful artificial insemination after exogenous gonadotropin treatment in the ocelot (*Leopardus pardalis*) and tigrina (*Leopardus tigrina*). *Proceedings of American Association of Zoo Veterinarians Congress*, p. 334-335.
- Morais, R. N.; Moreira, W.; Mucciolo, R. G.; Lacerda, O.; Gomes, M. L. F.; Swanson, W. F.; Graham, L. H. & Brown, J. L. (1996). Testicular and ovarian function in south american small felids assessed by fecal steroids. *Proceedings of American Association of Zoo Veterinarians Congress*, pp. 561-565.
- Morais, R. N. (2001). Reproduction in small felid males. In: *Biology, Medicine and Surgery of South American Wild Animals*, Fowler, M. E. & Cubas, Z. S. (ed). 1 ed. Ames: Iowa State University Press. pp.312-316.
- Morato, R. G.; Crichton, E. G.; Paz, R. C. R.; Zúge, R. M.; Moura, C. A.; Nunes, A. V. L.; Teixeira, R. H.; Porto-Filho, L.; Guimarães, M. A. B. V.; Correa, S. H. R.; Barnabe, R. C.; Armstrong, D. L. & Loskutoff, N. M. (2000). Ovarian stimulation and successful in vitro fertilization in the jaguar (*Panthera onca*). *Theriogenology*, Vol.53, No.1, pp.339. Abstract.
- Morato, R. G. & Paz, R. C. R. (2001). *Reproduction in jaguars*. In: *Biology, Medicine and Surgery of South American Wild Animals*, Fowler, M. E. & Cubas, Z. S. (ed). 1 ed. Ames: Iowa State University Press. pp.308-312.
- Moreira, N.; Monteiro-Filho, E. L. A.; Moraes, W.; Swanson, W. F.; Graham, L. H.; Pasqual, O. L.; Gomes, K. L. F.; Morais, R. N.; Wildt, D. E. & Brown, J. L. (2001). Reproductive steroid hormones and ovarian activity in felids of the leopardus genus. *Zoo Biology*, Vol.20, pp.103-116.
- Oliveira, T. G. (1994). *Neotropical Cats Ecology and Conservation*. São Luis: EDUFMA, 220 p.
- Oliveira, T. G. & CASSARO, K. , (1997). *Guia de Identificação dos Felinos Brasileiros*. São Paulo: Fundação Parque Zoológico de São Paulo, 60p.
- Paz, R. C. R.; Guido, M. C.; Costa, E. O.; Zúge, R. M.; Morato, R. G.; Guimarães, M. A. B. V.; Nunes, A. L. V.; Felipe, P. A. N.; Jimenez, G. T.; Crudeli, G. A.; Barnabe, V. H. & Barnabe, R. C. (1999). Microbiota Prepuccial de onças pintadas (*Panthera onca*) mantidas em cativeiro. *Proceedings of Congresso Brasileiro para Conservação de Felinos Neotropicais*, Jundiaí, São Paulo, Brasil.
- Paz, R. C. R.; Zúge, R. M.; Morato, R. G.; Barnabe, V. H.; Barnabe, R. C. & Felipe, P. A. N. (2000). Capacidade de Penetração de sêmen congelado de onça pintada (*Panthera onca*) em oócitos heterólogos. *Brazilian Journal of Veterinary Research and Animal Science*, Vol.37, No.6, pp. 462-466.
- Paz, R. C. R.; Swanson, W. F.; Dias, E. A.; Adania, C. H.; Barnabe, V. H. & Barnabe, R. C. (2005). Ovarian and immunological responses to alternating exogenous gonadotropin regimens in the ocelots (*Leopardus pardalis*) and tigrina (*Leopardus tigrinus*). *Zoo Biology*, Vol.24, pp.247-260.
- Paz, R. C. R.; Dias, E. A.; Adania, C. H.; Barnabe, V. H. & Barnabe, R. C. (2006). Ovarian response to repeated administration of alternating exogenous gonadotropin regimens in the ocelot (*Leopardus pardalis*) and tigrinus (*Leopardus tigrinus*). *Theriogenology*, Vol. 66, pp. 1787-1789.

- Pope, C. E.; Gelwicks, E. J.; Wachs, K. B.; Keller, G. L.; Maruska, E. J. & Dresser, P. L. (1989). Successful interspecies transfer of embryos from the Indian desert cat (*Felis silvestris ornata*) to the domestic cat (*Felis catus*) following in vitro fertilization. *Biology of Reproduction*, Vol.40, pp. 61. Supplementum 40.
- Pope, C. E. (2000). Embryo technology in conservation efforts for endangered felids. *Theriogenology*, Vol.53, pp. 163-174.
- Rodrigues da Paz, R. C.; Morato, R. G.; Carciofi, A. C.; Guimarães, M.A.B.V.; Pessuti, C.; Ferraz, E. S.; Ferreira, F. & Barnabe, R.C. (2006). Influence of nutrition on the quality of sêmen in Jaguars (*Panthera onca*) in Brazilian zoos. *Internationa Zoo Yearbook*, Vol.40, p 351-359.
- Roth, T. L.; Wolfe, B. A.; Long, J. A.; Howard, J. & Wildt, D. E. (1997). Effects of equine chorionic gonadotropin, human chorionic gonadotropin, and laparoscopic artificial insemination on embryo, endocrine, and luteal characteristics in the domestic cat. *Biology of Reproduction*, Vol.57, pp.165-71.
- Seal, U. S.; Plotka, E. D.; Smith, J. D.; Wright, F. H.; Reindl, N. J.; Taylor, R. S. & Seal, M. F. (1985). Immunoreactive luteinizing hormone estradiol, progesterone, testosterone, and androstenedione levels during the breeding season and anestrus in siberian tigers. *Biology of Reproduction*, Vol. 32, pp. 361-368.
- Schmidt, A. M.; Nadal, L. A.; Schmith, M. J. & Beamer, N. B. (1979). Serum concentrations of oestradiol and progesterone during the normal oestrus cycle and early pregnancy in the lion (*Panthera leo*). *Journal of Reproduction and Fertility*, Vol. 57, pp. 267-272.
- Schmidt, A. M.; Hess, D. L.; Schmith, M. J.; Smith, R. C. & Lewis, C. R. (1988). Serum concentration of oestradiol and progesterone, and sexual behavior during the normal oestrus cycle in the leopard (*Panthera pardus*). *Journal Reproduction and Fertility*, Vol. 82, pp. 43-49.
- Schmidt, A. M.; Hess, D. L.; Schmith, M. J. & Lewis, C. R. (1993). Serum concentration of oestradiol and progesterone and frequency os sexual behavior during the normal oestrous cycle in the snow leopard (*Panthera uncia*). *Journal Reproduction and Fertility*, Vol. 98, pp. 91-95.
- Seal, U. S.; Plotka, E. D.; Smith, J. D.; Wright, F. H.; Reindl, N. J.; Taylor, R. S. & Seal, M. F. (1985). Immunoreactive luteinizing hormone estradiol, progesterone, testosterone, and androstenedione levels during the breeding season and anestrus in siberian tigers. *Biology of Reproduction*, Vol.32, pp.361-368.
- Shille, V. M.; Lundström, K. E. & Stabenfeldt, G. H. (1979). Follicular function in the domestic cat as determined by estradiol-17 β concentration in plasma: relation to estrous behavior and cornification of exfoliated vaginal epithelium. *Biology of Reproduction*, Vol.21, pp. 953-963.
- Silva, J. C.; Leitão, R. M.; Lapão, N. E.; Cunha, M. B.; Cunha, T. P.; Silva, J. P. & Paisana, F. C. (2000). Birth of siberian tiger (*Panthera tigris altaica*) cubs after transvaginal artificial insemination. *Journal of Zoo and Wildlife Medicine*, Vol.34, pp. 566-569.
- Spindler, R. E.; Wildt, D. E. (1999). Circannual variations in intraovarian oocyte but not epididymal sperm quality in the domestic cat. *Biology of Reproduction*, Vol. 61, pp. 188-194.
- Swanson, W. F.; Brown, J. L.; Wildt, D. E. (1996). Influence of season on reproductive traits of the male Pallas'cat (*Felis Manul*) and implications for captive management. *Journal of Zoo and Wildlife Medicine*, Vol. 27, pp. 234-240.
- Swanson, W. F.; Horohov, D. W. & Godke, R. A. (1995). Production of exogenous gonadotrophin-neutralizing immunoglobulins in cats after repeated eCG-hCG

- treatment and relevance for assisted reproduction in felids. *Journal of Reproduction and Fertility*, Vol.105, pp. 35-41.
- Swanson, W. F.; Howard, J. G.; Roth, T. L.; Brown, J. L.; Alvarado, T.; Burton, M.; Starnes, D. & Wildt, D. E. (1996). Responsiveness of ovaries to exogenous gonadotrophins and laparoscopic artificial insemination with frozen-thawed spermatozoa in ocelots (*Felis pardalis*). *Journal of Reproduction and Fertility*, Vol.106, pp. 87-94 a.
- Swanson, W. F.; Roth, T. L.; Graham, K.; Horohov, D. W. & Godke, R. A. (1996). Kinetics of the humoral immune response to multiple treatments with exogenous gonadotropins and relation to ovarian responsiveness in domestic cats. *American Journal Veterinary Research*, Vol.57, No.3, pp. 302-307 b.
- Swanson, W. F. (1998). Reproduction in domestic cats: model for endangered nondomestic cat species. In: *Curso de extensão - Felinos selvagens: biotécnicas reprodutivas e conservação*, Morais, R. N. & Gomes, M. L. F. Curitiba: Departamento de Fisiologia da Universidade Federal do Paraná, 100 p.
- Swanson, W. F. (2001). Reproductive biotechnology and conservation of the forgotten felids - the small cats. *Proceedings of 1st International Symposium on Assisted Reproductive Technologies: Conservation & Genetic Management Wildlife*, Omaha, Nebraska, USA, pp. 100-120.
- Swanson, W. F.; Paz, R. C. R.; Morais, R. N.; Gomes, M. L. G.; Moraes, W. & Adania, C. H. (2002). Influence of species and diet on efficiency of in vitro fertilization in two endangered Brazilian felids - The ocelot (*Leopardus pardalis*) and tigrina (*Leopardus tigrinus*). *Theriogenology*, Vol.57, No.1, pp. 593. Abstract.a
- Swanson, W. F. (2002). The role of science and reproductive biotechnology in establishing and managing the Brazilian ocelot population in U.S. and Brazilian Zoos. *Proceedings of Annual Conference of the American Zoo & Aquarium Association*, Fort Worth, Texas, USA. pp.75-78.b
- Swanson, W. F. & Brown, J. L. (2004). International training programs in reproductive sciences for conservation of Latin American felids. *Animal Reproduction Science*, Vol.82-83, pp. 21-34.
- Tebet, J. M. (1999). *Aspectos clínicos e fisiológicos do ciclo estral da jaguatirica (Leopardus pardalis, L.1758)*. (Dissertação Mestrado em Reprodução Animal) - Faculdade de Medicina Veterinária e Zootecnia, Universidade Estadual Paulista "Júlio de Mesquita Filho", Botucatu, São Paulo. pp. 102.
- Tebet, J. M.; Martins, M. I. M.; Chirinea, V. H.; Souza, F. F.; Campagnol, D.; Lopes, M. D. (2006). Cryopreservation effects on domestic cat epididymal versus electroejaculated spermatozoa. *Theriogenology*, Vol. 66, pp. 1629-1632.
- Tsutsui, T. & Stabenfeldt, G. H. (1996). Biology of ovarian cycles, pregnancy and pseudopregnancy in the domestic cat. *Journal of Reproduction and Fertility*, pp. 29-35. Supplementum 47.
- Verhage, H. G.; Beamer, N. B. & Brenner, R. M. (1976). Plasma levels of estradiol and progesterone in the cat during polyestrus, pregnancy and pseudopregnancy. *Biology of Reproduction*, Vol.14, pp. 579-585.
- Wildt, D. E.; Platz, C. C.; Chakraborty, P. K. & Seager, S. W. J. (1979). Oestrus and ovarian activity in a female jaguar (*Panthera onca*). *Journal of Reproduction and Fertility*, Vol.56, pp. 555-58.
- Wildt, D. E.; Platz, C. G.; Seager, S. W. J. & Bush, M. (1981). Induction of ovarian activity in the cheetah (*Acinonyx jubatus*). *Biology of Reproduction*, Vol.24, pp. 217-222.

Current Frontiers in Cryobiology

Edited by Prof. Igor Katkov

ISBN 978-953-51-0191-8

Hard cover, 574 pages

Publisher InTech

Published online 09, March, 2012

Published in print edition March, 2012

Almost a decade has passed since the last textbook on the science of cryobiology, *Life in the Frozen State*, was published. Recently, there have been some serious tectonic shifts in cryobiology which were perhaps not seen on the surface but will have a profound effect on both the future of cryobiology and the development of new cryopreservation methods. We feel that it is time to revise the previous paradigms and dogmas, discuss the conceptually new cryobiological ideas, and introduce the recently emerged practical protocols for cryopreservation. The present books, "Current Frontiers in Cryobiology" and "Current Frontiers in Cryopreservation" will serve the purpose. This is a global effort by scientists from 27 countries from all continents and we hope it will be interesting to a wide audience.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Regina Celia Rodrigues da Paz (2012). *Wildlife Cats Reproductive Biotechnology*, Current Frontiers in Cryobiology, Prof. Igor Katkov (Ed.), ISBN: 978-953-51-0191-8, InTech, Available from:
<http://www.intechopen.com/books/current-frontiers-in-cryobiology/wildlife-cats-reproductive-biotechnology>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](https://creativecommons.org/licenses/by/3.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen