We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Effect of Antisense Squalene Synthase Gene Expression on the Increase of Artemisinin Content in *Artemisia annua*

Hong Wang et al.*

Graduate University of the Chinese Academy of Sciences, Beijing, China

1. Introduction

Artemisinin, a sesquiterpene lactone endoperoxide, is a valuable and powerful antimalarial drug obtained from the aerial parts of a Chinese herb, *Artemisia annua* (Liu et al., 1979). Artemisinin and its derivatives show few or no side effects with the existing antimalarial drugs, it has consequently been regarded as the next generation of antimalarial drugs (Looaresuwan, 1994). Currently, commercial production of artemisinin mainly based on its extraction and purification from plant material, however, the endogenous production of artemisinin is very low (0.01%-0.8% dry weight) (Wallaart et al., 1999). In view of the limited availability of artemisinin and the increased demand, the synthetic preparation of artemisinin becomes an attractive proposition. However due to its complex structure, the complete chemical synthesis is very difficult (Schmid & Hofheinz, 1983). *Artemisia annua* as the only valid source, many research groups have directed their investigations toward the enhancement of artemisinin production in *A. annua* cell cultures or whole plants by biotechnological approaches. However these approaches were still proved to be not successful (Ghingra et al., 2000). Recently, several genes in artemisinin biosynthesis have been cloned, and important advances in artemisinin biosynthesis have been achieved, which makes it possible to regulate artemisinin biosynthesis in a direct way, for example, by metabolic engineering (Abdin et al., 2003).

Artemisinin is synthesized by the isoprenoid pathway. In the cytosol, isoprenoids are synthesized via the classical acetate/mevalonate pathway (Fig. 1). In this pathway, farnesyl diphosphate (FDP) occupies a central position and serves as a common substrate for the first committed reactions of sterols and sesquiterpenes, such as artemisinin. Therefore, this point represents a potentially important controlling point for balancing sterol synthesis and sesquiterpenes synthesis. From metabolic engineering point of view (Fig. 1), there are two

*Yugang Song¹, Haiyan Shen¹, Yan Liu², Zhenqiu Li³, Huahong Wang², Jianlin Chen², Benye Liu² and Hechun Ye²

¹Graduate University of the Chinese Academy of Sciences, Beijing, China

²Key Laboratory of Photosynthesis and Environmental Molecular Physiology, Institute of Botany, the Chinese Academy of Sciences, Beijing, China

³Hebei University, Baoding, China

ways to increase the flux to artemisinin biosynthesis, on the one hand, we can overexpress the key genes involved in the biosynthesis of artemisinin; on the other hand, we can inhibit the genes involved in other pathways competing for its precursors.

ADS: amorpha-4,11-diene synthase; FPP: farnesyl diphosphate; FPS: farnesyl diphosphate synthase; HMG-CoA: 3-hydroxy-3-methylglutaryl CoA; HMGR: HMG-CoA reductase; HMGS: HMG-CoA synthase; IPP: isopentenyl diphosphate; MVA: mevalonate; SES: sesquiterpene synthase; SQS: squalene synthase.

Fig. 1. Diagram of the mevalonate pathway leading to the biosynthesis of sesquiterpenes and sterols.

Squalene synthase (SQS) catalyzes the condensation of two molecules of farnesyl diphosphate (FDP) to form the linear 30 carbon compound squalene, the first committed precursor for sterol biosynthesis (Goldstein & Brown, 1990). SQS is generally described as a crucial branch point enzyme for synthesizing sterol intriguing as a potential regulatory point that controls carbon flux into either sterol or into non-sterol isoprenoids (such as sesquiterpenes). So if the SQS gene expression is inhibited by genetic manipulation, the carbon flux into sterol may be diverted to sesquiterpenes, and the biosynthesis of sesquiterpenes may be increased. With the purpose to increase artemisinin production, we have cloned squalene synthase cDNA (SQS) (Liu et al., 2003). In this chapter, we report the construction of the antisense SQS plant expression vector, and its effects on inhibition of SQS gene expression on squalene and artemisinin biosynthesis.

2. Materials and methods

2.1 Plant materials

A high artemisinin producing *Artemisia annua* L. strain 001 was collected from Sichuan Province of China. The seeds were surface sterilized and cultured on the Murashige &

2.2 Construction of antisense plant gene expression vector

2.3 *Agrobacterium* preparation and genetic transformation

Plant transformation was basically performed according to the method of Han et al (2005). The leaves of 2-week-old aseptic seedlings were immersed in 50 mL *A. tumefaciens*, which was at the log phase of growth and was diluted 10-fold with MS medium. After 20 min, the

infected leaves were taken out, blotted with sterile paper and co-cultured on solid MS medium at 26 °C for 2 or 3 d. After co-cultivation, the leaves were transferred to shoot-inducing medium (MS medium supplemented 1.0 mg/L 6-BA and 0.05 mg/L NAA) containing 20 mg/L Kan to induce shoot and 500 mg/L cefotaxime (Cefo) to kill residual *Agrobacterium*. The medium without Kan was used as a control. Here the key difference with Han's method is that a sheet of sterile filter paper was placed on the shoot-inducing medium, since we found by such a simple action, the shoot induction frequency can be noticeably increased. One week later, the leaves were transferred to fresh MS selection medium with 400 mg/L Cefo. After 4-6 weeks selection Kan resistant shoots were obtained and transferred to 100 mL Erlenmeyer flask with 40 mL regeneration medium (MS basal medium supplemented with 0.05 mg/L NAA) to induce roots. All the rooted seedlings were kept in the greenhouse at 25 °C under the fluorescent lamps (with a light intensity of 3000 lx, 16h), with a relative humidity of 40%.

2.4 PCR detection

The integration of SQS gene into *A. annua* genome was confirmed by the polymerase chain reaction (PCR). The CTAB method was used to purify the genomic DNA from transgenic *A. annua* leaves. The forward primer was 5'-CCA CGT CTT CAA AGC AAG TGG ATT -3', designed according to the sequence of CaMV 35S promoter, and the reverse primer was 5'-GCA GAG CTC GGA TTT GGA TCT TGA AGA A G -3', designed according to the sequence at the 3'-terminal region of SQS gene. 30-cycle reactions, each consisted of heated denaturation (94 °C for 40 s), annealing (55 °C for 30 s) and extension (72 °C for 2 min), were carried out and the reaction mixtures were subjected to agarose gel electrophoresis.

2.5 RNA isolation and RT-PCR detection

For reverse transcription-polymerase chain reactions (RT-PCR), 1 g of total RNA isolated from the leaves of the transformed and non-transformed plants was used. The RNA extraction protocol was done as described in Sambrook et al (1989). The first strand cDNA was synthesized by using a first-strand cDNA synthesis kit (TaKaRa), according to the manufacturer's instructions. The resultant first-strand cDNA was used as a template, and the PCR primers, P1: 5'-GGA ACC ATG GGT AGT TTG AAA GCA GTA TTG-3, and P2: 5'-GCC TGG ATC CCT TGA CTC TCT CTT AAC TAT-3', were designed according to the SQS gene sequence of *A. annua*. The PCR was performed in the same way as described in the Section of PCR detection. At the same time, to normalize the amount of mRNA in each PCR reaction, a PCR product of actin in *A. annua* was amplified, and the primers were: 5'-AAC TGG GAT GAC ATG GAG AAG ATA T-3', and 5'-TCA CAC TTC ATG ATG GAG TTG TAG G-3'.

2.6 Squalene analysis

Squalene analysis was carried out according to the method described by Wentzinger et al (2002). For all the plants analyzed, young leaves was collected. Extraction and purification of the samples were in accordance with Wentzinger et al (2002), and the samples were analyzed by GC, the GC injection port was operated at 120 °C. The oven temperature programmed from 120 °C to 180 °C at 15 °C min⁻¹ and from 180 °C to 260 °C at rate of 25 °C min⁻¹. The final temperature was maintained for 25 min. The results were compared to standards.

2.7 Determination of artemisinin

The detection of artemisinin was performed according to the method of Zhao & Zeng (1986). Fresh leaves of the transformed and non-transformed plants were collected and dried to constant weight in an oven at 50 °C. Then the dried leaves were ground to fine powder. Exactly 0.05 g powder was added to an extraction bottle containing 40 mL petroleum ether (30-60 °C) and treated in a supersonic bath for 2 min. The extraction mixture was filtered and the petroleum ether was evaporated. The residue was dissolved in 1 mL methanol and centrifuged at 12000 r/min to precipitate the undissolved components. The supernatant was used for detection of artemisinin by HPLC.

200 µL of the above prepared methanol solution was placed in a 10 mL tube and 800 µL methanol and 4 mL 0.2% sodium hydroxide were added, mixed and maintained in a 50 °C water bath for 30 min, then the reaction mixture was cooled to room temperature. 0.5 mL of the reaction mixture was placed in a 1.5 mL Eppendorf tube, and 100 µL methanol and 400 µL 0.05 M acetic acid were added, mixed and the sample purified by filtering on a NC filter (40 µm). The artemisinin standard (Sigma, MO) solutions with concentrations of 3, 6, 12, 24 and 48 µg/mL were prepared in the same way as the sample.

C18 reverse column was 4.6×250 mm, 5 µm. The mobile phase was 0.01 M phosphate buffer (pH 7.0) : methanol (55:45), with flow rate 1 mL/min. The wavelength of the UV detector was 288.6 nm and the injection volume was 20 µL. Artemisinin standard appeared at 4 min 30 s under the above mentioned conditions.

3. Results

3.1 Regeneration of transgenic *A. annua*

Leaf discs, which were infected with *A. tumefaciens* strain EHA105 harboring the binary vector pBISQS, were co-cultured for 36-48 h at 26 °C in dark, then transferred to shoot-inducing medium. Shoots usually start to appear within 2-3 weeks on this medium. In order to obtain higher frequency of shoot induction, a sheet of filter paper was placed on the shoot-inducing medium during the transformant selection step, which was proved to be a very effective means for shoot induction (Song et al., 2006). Regenerated shoots are rooted on MS medium with 0.05 mg/L NAA, followed by transferring to greenhouse (16 h light at 25 °C).

3.2 Molecular analysis of transgenic plants

To investigate the presence of antisense SQS gene in the putatively transformed plants, genomic DNA of 4 Kan-resistant plants regenerated from the leaves inoculated with EHA105 was isolated, and PCR analysis was performed. The antisense SQS gene was detected as 1660-bp fragments in all 4 analyzed plants (Fig. 3). The amplified fragments were of the same size as the predicted one. The fragment in the nontransformed plant was not amplified.

RT-PCR was performed using specifically designed primers according to the squalene synthase cDNA sequence in *Artemisia annua*. These primers allow specific amplification of *A. annua* squalene synthase cDNA. The results showed that the suppressed expression of the

endogenous *A. annua* squalene synthase gene in lines SQS3 and SQS5, but the transcriptional level in line SQS2 had no noticeable difference to that of the control (Fig. 4).

1. positive control; 2-5. transformed lines; 6. negative control; 7. DNA ladder.

Fig. 3. PCR amplification of the transformed plants and the control.

1. 001 line (non-transgenic control); 2. SQS2; 3. SQS3; 4. SQS5.

Fig. 4. RT-PCR analysis of the transformed plants and the control.

3.3 Detection of squalene content

In order to determine the effects of inhibiting squalene synthase gene expression on sterol biosynthesis, the leaves of transgenic lines SQS3, SQS5, and these of the control 001 were selected to detect squalene content. The results of GC-MS showed that in SQS3 and SQS5 transgenic lines, squalene content is decreased by 19.4% and 21.6% respectively in comparison with the control (Fig. 5).

1. 001 line (control); 2. SQS3; 3. SQS5.

Fig. 5. Analysis of squalene content of the transgenic plants and the control.

3.4 Determination of artemisinin

Artemisinin was detected by HPLC. The results indicated that artemisinin content of SQS3 and SQS5 transgenic lines was increased by 23.2% and 21.5%, respectively compared with that of the control (Fig. 6) and in SQS2 transgenic line, the artemisinin content manifested no obvious variation compared with the control.

The above results demonstrated a clear negative correlation between squalene content and artemisinin content, which implies that the inhibiting of squalene synthase gene expression caused part of the flux for squalene biosynthesis diverting to artemisinin biosynthesis.

1. 001 line (control); 2. SQS2; 3. SQS3; 4. SQS5

Fig. 6. Analysis of artemisinin content of the transgenic plants and the control.

4. Discussion

Since squalene synthase is commonly depicted as the incipient and crucial branch point enzyme of the isoprenoid pathway to sterol biosynthesis, it has attracted considerable interest as a potential regulatory point that controls carbon flux into sterols. Several researchers reported the induction of sesquiterpene phytoalexins biosynthesis had been correlated with suppression of sterol biosynthesis in elicitor-treated tobacco cell cultures (Chappell et al., 1989; McGarvey & Croteau, 1995; Yin et al., 1997). The induction of one enzyme and suppression of the other are thought to be one mechanism that regulates the production of squalene and sesquiterpene (Devarenne et al., 1998). The biosynthesis of artemisinin belongs to the isoprenoid pathway, in this pathway, squalene synthase and amorph-4,11-diene synthase are positioned at putative branch points in isoprenoid metabolism, these two enzymes catalyze the common farnesyl diphosphate to form squalene and amorph-4,11-diene, respectively. Furthermore, amorph-4,11-diene synthase is considered as a key enzyme in artemisinin biosynthesis (Bouwmeester et al., 1999), so squalene synthase can be considered as a competitive enzyme of artemisinin biosynthesis. Therefore, the inhibiting of SQS gene expression may increase the biosynthesis of artemisinin.

In order to increase artemisinin content, we introduced antisense squalene synthase gene into *A. annua* via *Agrobacterium*-mediated transformation and transgenic plants were obtained. It has been shown that the transgenic plants had an increase of artemisinin content in lines SQS3 and SQS5 and a reduction in squalene content. This may be due to the suppression of squalene synthase gene expression caused part of the carbon flux to squalene biosynthesis diverting to artemisinin biosynthesis. At the same time, in correspondence with the decline in squalene content, the endogenous squalene synthase transcript level of the SQS3 and SQS5 transgenic lines were reduced compared with the control. These results strongly supported that the overexpression of squalene synthase in antisense orientation had a relevant effect on endogenous squalene metabolism in transgenic tobacco plants (Zhang et al., 2005). Though the SQS gene expression of the transgenic plants was inhibited, both the growth and the phenotype of the transgenic plants showed no obvious difference to those of the control, and this was in consistent with the results of transgenic tobacco (Zhang et al., 2005).

Our results as well as other related reports all indicated that it is possible to increase artemisinin content of *A. annua* by inhibiting the expression of genes competing precursors with artemisinin in transgenic plants (Yang et al., 2008; Zhang et al., 2009; Chen et al., 2011). But it has caused only a limited increase in artemisinin content, perhaps this is because of the biosynthesis of artemisinin is controlled by multi-genes and the contribution of manipulating one gene to artemisinin biosynthesis is limited. Recently, the study of artemisinin has made great progress, and more and more genes of artemisinin biosynthesis, such as cytochrome P450 monooxygenase (CYP71AV1), double bond reductase 2 (DBR2) and aldehyde dehydrogenase 1 (ALDH1) (Teoh et al., 2006; Covello et al., 2007; Zhang et al., 2008; Teoh et al., 2009), have been cloned, this makes it possible to regulate multi-genes of artemisinin biosynthesis in the future.

5. Conclusion

The antisense squalene synthase (SQS) gene was transferred into *A. annua* via *Agrobacterium*-mediated transformation, and the artemisinin content of one of the transgenic lines showed an increase of 23.2% in comparison to the wild-type control. The results demonstrated that inhibiting pathway competing for precursor of artemisinin by anti-sense technology is an effective means of increasing the artemisinin content of *A. annua* plants.

6. Acknowledgements

This work was supported by the National Natural Science Foundation of China (NSFC) (No. 61173098) and the Knowledge Innovation Program of the Chinese Academy of Sciences (KSCX2-EW-J-29).

7. References

- Abdin, M., Israr, M., Rehman, R. & Jain, S. (2003). Artemisinin, a novel antimalarial drug: biochemical and molecular approaches for enhanced production. *Planta Medica*, Vol.69, No.4 (April 2003), pp. 289-299, ISSN 0032-0943

- Bouwmeester, H., Wallaart, T., Janssen, M., van Loo, B., Jansen, B., Posthumus, M., Schmidt, C., De Kraker, J., Konig, W. & Franssen, M. (1999). Amorpha-4,11-diene synthase catalyses the first probable step in artemisinin biosynthesis. *Phytochemistry*, Vol.52, No.5 (November 1999), pp. 843-854, ISSN 0031-9422
- Chappell, J., Von Lanken, C., Vögeli, U. & Bhatt, P. (1989). Sterol and sesquiterpenoid biosynthesis during a growth cycle of tobacco cell suspension cultures. *Plant Cell Reports*, Vol.8, No.1 (January 1989), pp. 48–52, ISSN 0721-7714
- Chen, J., Fang, H., Ji, Y., Pu, G., Guo, Y., Huang, L., Du, Z., Liu, B., Ye, H., Li, G. & Wang, H. Artemisinin biosynthesis enhancement in transgenic *Artemisia annua* plants by downregulation of the β -caryophyllene synthase gene. *Planta Medica*, DOI: 10.1055/s-0030-1271038, Epub ahead of print, ISSN 0032-0943
- Covello, P., Teoh, K., Polichuk, D., Reed, D. & Nowak, G. (2007). Functional genomics and the biosynthesis of artemisinin. *Phytochemistry*, Vol.68, No.14 (July 2007), pp. 1864-1871, ISSN 0031-9422
- Devarenne, T., Shin, D., Back, K., Yin, S. & Chappell, J. (1998). Molecular characterization of tobacco squalene synthase and regulation in response to fungal elicitor. *Archives of Biochemistry and Biophysics*, Vol.349, No.2 (January 1998), pp. 205–215, ISSN 0003-9861
- Ghingra, V., Vishweshwar Rao, K. & Lakshmi Narasu, M. (2000). Current status of aremisinin and its derivatives as antimalarial drugs. *Life Sciences*, Vol.66, No.4 (December 1999), pp. 279-300, ISSN 0024-3205
- Goldstein, J. & Brown, M. (1990). Regulation of the mevalonate pathway. *Nature*, Vol.343, No.6257 (February 1990), pp. 425–430, ISSN 0028-0836
- Han, J., Wang, H., Ye, H., Liu, Y., Li, Z., Zhang, Y., Zhang, Y., Yan, F. & Li, G. (2005). High efficiency of genetic transformation and regeneration of *Artemisia annua* L. via *Agrobacterium tumefaciens*-mediated procedure. *Plant Science*, Vol.168, No.1 (January 2005), pp. 73-80, ISSN 0168-9452
- Liu, J., Ni, M., Fan, J., Tu, Y., Wu, Z., Wu, Y. & Chou, W. (1979). Structure and reaction of arteannuin. *Acta Chimica Sinica*, Vol.37, No.2 (April 1979), pp. 129-143, ISSN0567-7351
- Liu, Y., Ye, H., Wang, H. & Li, G. (2003). Molecular cloning, Escherichia coli expression and genomic organization of squalene synthase gene from *Artemisia annua*. *Acta Botanica Sinica*, Vol. 45, No.5 (May 2003), pp. 608-613, ISSN 0577-7496
- Looaresuwan, S. (1994). Overview of clinical studies on artemisinin derivatives in Thailand. *Transactions of the Royal Society of Tropical Medicine and Hygiene*, Vol.88, No.suppl (June 1994), pp. s9-11, ISSN 0035-9203.
- McGarvey, D., & Croteau, R. (1995). Terpenoid metabolism. *The Plant Cell*, Vol.7, No.7 (July 1995), pp. 1015–1026, ISSN 1040-4651
- Murashige, T. & Skoog, F. (1962). A revised medium for rapid growth and bioassays with tobacco tissue cultures. *Physiologia Plantarum*, Vol.15, No.3 (September 1962), pp. 473-497, ISSN 0031-9317
- Sambrook, J., Fritsch, E. & Maniatis, T. (1989). *Molecular cloning: A laboratory manual* (second edition), Cold Spring Harbor Laboratory Press, ISBN 0-87969-309-6, New York
- Schmid, G. & Hofheinz, W. (1983). Total synthesis of qinghaosu. *Journal of the American Chemical Society*, Vol.105, No.3 (February 1983), pp. 624-625, ISSN 0002-7863

- Song, Y., Li, Z., Wang, H., Liu, B., Li, G. & Ye, H. (2006). Effect of filter paper on shoot induction and genetic transformation of *Artemisia annua* L. *Chinese Journal of Applied and Environmental Biology*, Vol.12, No.6 (December 2006), pp. 788-791, ISSN 1006-687X
- Teoh, K., Polichuk, D., Reed, D., Nowak, G. & Covello, P. (2006). *Artemisia annua* L. (Asteraceae) trichome-specific cDNAs reveal CYP71AV1, a cytochrome P450 with a key role in the biosynthesis of the antimalarial sesquiterpene lactone artemisinin. *FEBS Letters*, Vol.580, No.5 (February 2006), pp. 1411-1416, ISSN 0014-5793
- Teoh, K., Polichuk, D., Reed, D. & Covello, P. (2009). Molecular cloning of an aldehyde dehydrogenase implicated in artemisinin biosynthesis in *Artemisia annua*. *Botany*, Vol.87, No.6 (June 2009), pp. 635-642, ISSN 1916-2790
- Wallaart, T., Pras, N. & Quax, W. (1999). Seasonal variations of artemisinin and its biosynthetic precursors in tetraploid *Artemisia annua* plants compared with the diploid wild-type. *Planta Medica*, Vol.65, No.8 (December 1999), pp. 723-728, ISSN 0032-0943
- Wentzinger, L., Bach, T. & Hartmann, M. (2002). Inhibition of squalene synthase and squalene epoxidase in tobacco cells triggers an up-regulation of 3-hydroxy-3-methylglutaryl coenzyme a reductase. *Plant Physiology*, Vol.130, No.1 (September 2002), pp. 334-346, ISSN 0032-0889
- Yang, R., Feng, L., Yang, X., Yin, L., Xu, X. & Zeng, Q. (2008). Quantitative transcript profiling reveals down-regulation of a sterol pathway relevant gene and overexpression of artemisinin biogenetic genes in transgenic *Artemisia annua* plants. *Planta Medica*, Vol.74, No.12 (October 2008), pp. 1510-1516, ISSN 0032-0943
- Yin, S., Mei, L., Newman, J., Back, K. & Chappell, J. (1997). Regulation of sesquiterpene cyclase gene expression. Characterization of an elicitor- and pathogen-inducible promoter. *Plant Physiology*, Vol.115, No.2 (October 1997), pp. 437-451, ISSN 0032-0889
- Zhang, L., Jing, F., Li, F., Li, M., Wang, Y., Wang, G., Sun, X. & Tang, K. (2009). Development of transgenic *Artemisia annua* (Chinese wormwood) plants with an enhanced content of artemisinin, an effective anti-malarial drug, by hairpin-RNA-mediated gene silencing. *Biotechnology and Applied Biochemistry*, Vol.52, No.3 (March 2009), pp. 199-207, ISSN 0885-4513
- Zhang, Y., Liu, Y., Wang, H., Ye, H. & Li, G. (2005). Regulation of squalene synthase gene expression in tobacco by antisense transformation with an *Artemisia annua* squalene synthase gene. *Journal of Agricultural Biotechnology*, Vol.13, No.4 (August 2005), pp. 416-422, ISSN 1674-7968
- Zhang, Y., Teoh, K., Reed, D., Maes, L., Goossens, A., Olson, D., Ross, A. & Covello, P. (2008). The molecular cloning of artemisinic aldehyde Delta 11(13) reductase and its role in glandular trichome-dependent biosynthesis of artemisinin in *Artemisia annua*. *The Journal of Biological Chemistry*, Vol.283, No.31 (August 2008), pp. 21501-21508, ISSN 0021-9258
- Zhao, S., & Zeng, M. (1986). Determination of qinghaosu in *Artemisia annua* L. by high performance liquid chromatography. *Chinese Journal of Pharmaceutical Analysis*, Vol.6, No.1 (January 1986), pp. 3-5, ISSN 0254-1793

Transgenic Plants - Advances and Limitations

Edited by PhD. Yelda Ozden Çiftçi

ISBN 978-953-51-0181-9

Hard cover, 478 pages

Publisher InTech

Published online 07, March, 2012

Published in print edition March, 2012

Development of efficient transformation protocols is becoming a complementary strategy to conventional breeding techniques for the improvement of crops. Thus, Transgenic Plants - Advances and Limitations covers the recent advances carried on improvement of transformation methods together with assessment of the impact of genetically transformed crops on biosafety. Each chapter has been written by one or more experienced researchers in the field and then carefully edited to ensure thoroughness and consistency.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Hong Wang, Yugang Song, Haiyan Shen, Yan Liu, Zhenqiu Li, Huahong Wang, Jianlin Chen, Benye Liu and Hechun Ye (2012). Effect of Antisense Squalene Synthase Gene Expression on the Increase of Artemisinin Content in Artemisia annua, Transgenic Plants - Advances and Limitations, PhD. Yelda Ozden Çiftçi (Ed.), ISBN: 978-953-51-0181-9, InTech, Available from: <http://www.intechopen.com/books/transgenic-plants-advances-and-limitations/effect-of-antisense-squalene-synthase-gene-expression-on-the-increase-of-artemisinin-content-in-arte>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](https://creativecommons.org/licenses/by/3.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen