

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

185,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Potato Peel as a Source of Important Phytochemical Antioxidant Nutraceuticals and Their Role in Human Health – A Review

A. Al-Weshahy and V.A. Rao*

*Department of Nutritional Sciences, Faculty of Medicine,
University of Toronto, Toronto, Ontario,
Canada*

1. Introduction

Food processing as one of the most important industries over the globe, however, by-products of such industrial activity that are mainly organic material must be handling in appropriate manner to avoid any environmental violence. Sanitation and disposal problem of food processing by-products is priority hence many approaches were suggested including recycling of such ingredients and utilize in several food and / or non-food applications (1). By-products of food processing is an inexpensive, affordable, and valuable starting material for the extraction of value added products such as dietary fiber, natural antioxidants, biopolymers, and natural food additives(2-4). However, the central dogma is still the stability, and economic feasibility of the processing development (5,6).

Potato (*Solanum tuberosum* L.) is one of the major world crops with world annual production of 180 million tones on 2009 according to the Food and Agriculture Organization (FAO) (7). Potato vegetation is dated back to South America around 500 B.C (8) and over centuries, it becomes a cornerstone in human nutrition in which nutritional quality was well established and documented and considers a source for many nutrients as shown in (table 1).

Of over 30 MT of annual production, tenth goes to industry to produce variable products including French fries and potato chips while the rest consumes freshly (7). By-products from potato processing, as like as any other food processing industry, are principally organic materials thus management and disposal are crucial toward a clean industry.

Potato peel (PP) has been studied as feed for pigs (9).However, potato peel is not suitable, without further treatments, as feed for non-ruminants because it is too fibrous to be digested (10). On the other hand, potato peel has been successfully used in feeding of multi-gastric animals. Milk fat from cows fed with PP was reported to be 3.3 g/kg higher than that of control (11).

* Corresponding Author

Water (g)	79.3	Calcium (mg)	12	Vitamin A, RAE (mcg)	0	Monounsaturated fatty acids, total (g)	0
Energy (kcal)	77	Copper (mg)	0.11	Vitamin C (mg)	19.7	16:1 (g)	0
Protein (g)	2.02	Iron (mg)	0.78	Vitamin B-6 (mg)	0.3	18:1 (g)	0
Fat, total (g)	0.09	Magnesium (mg)	23	Choline, total (mg)	12.1	20:1 (g)	0
Carbohydrate (g)	17.5	Phosphorus (mg)	57	Vitamin B-12 (mcg)	0	22:1 (g)	0
Sugars, total (g)	0.78	Potassium (mg)	421	Vitamin B-12, added (mcg)	0	Polyunsaturated fatty acids, total (g)	0.04
Fiber, total dietary (g)	2.2	Selenium (mcg)	0.3	Vitamin E, alpha tocopherol (mg)	0.01	18:2 (g)	0.03
Alcohol (g)	0	Sodium (mg)	6	Vitamin E, added (mg)	0	18:3 (g)	0.01
Cholesterol (mg)	0	Zinc (mg)	0.29	Vitamin D (D2 + D3) (mcg)	0	18:4 (g)	0
Saturated fatty acids, total (g)	0.03	Carotene, beta (mcg)	1	Folate, DFE (mcg)	16	20:4 (g)	0
4:0 (g)	0	Carotene, alpha (mcg)	0	Folate, food (mcg)	16	20:5 n-3 (g)	0
6:0 (g)	0	Cryptoxanthin, beta (mcg)	0	Folate, total (mcg)	16	22:5 n-3 (g)	0
8:0 (g)	0	Lutein + zeaxanthin (mcg)	8	Folic acid (mcg)	0	22:6 n-3 (g)	0
10:0 (g)	0	Lycopene (mcg)	0	Vitamin K (mcg)	1.9		
12:0 (g)	0	Caffeine (mg)	0	Niacin (mg)	1.05		
14:0 (g)	0	Theobromine (mg)	0	Retinol (mcg)	0		
16:0 (g)	0.02			Riboflavin (mg)	0.03		
18:0 (g)	0			Thiamin (mg)	0.08		

Values calculated per 100 gm fresh weight
Source: USDA Food and Nutrient Database for Dietary Studies (FNDDS)

Table 1. Chemical composition and nutritional value of potato.

The present review summarizes most attempts of potato peels utilization in food and non-food applications that include the extraction and verification of bioactive ingredients and nutritional quality of potato peels and their applications.

Although PP is being used for feeding livestock, by-products from potato processing industry still outpace such limited utilization. Another aspect was its utilization as a source for natural antimicrobial compounds (12).

2. Potato peels as a source of dietary fibers in food applications

Dietary fiber is well known as a bulking agent, increasing the intestinal mobility and hydration of the feces (13). Several authors have reviewed the importance of consumption of moderate amounts of dietary fibers for human health (14).

Scientifically speaking, dietary fiber is a broad term that includes several carbohydrates; cellulose, hemicelluloses, lignins, pectins, gums etc. (15). (16) reported that PP fibers are primarily insoluble, and can bind bile acids in-vitro. It is believed that binding of bile acids is one of the mechanisms whereby certain sources of dietary fibers lower plasma cholesterol. (17) studied the hypocholesterolemic effect of dietary fiber from PP and found that after four weeks of feeding on potato peels, rats showed 40 % reduction in plasma cholesterol content and 30% of hepatic fat cholesterol levels were reduced as compared with animals fed only with cellulose supplemented diet. Defects of dietary fiber on lipid-profile influence several health related issues. High concentrations of low-density lipoprotein (LDL) cholesterol, other dyslipidemia (high

concentration of triglycerides and low concentration of high-density lipoprotein [HDL] cholesterol), leads to blood platelets aggregation (18), risk factors for cardiovascular diseases (CVD) (19), and hypertension (20). Moreover, high intake of dietary fibers has a positive influence on blood glucose profile and it is related health complications, in healthy and diabetic individuals of both types. By altering the gastric emptying time, dietary fibers are able to affect the absorption of other simple sugars.

The effect of dietary fibers on blood glucose and insulin response has been demonstrated by many other authors as well (21,22).

Different sources of dietary fibers have been used to replace wheat flour in the preparation of bakery products. Potato peel was introduced as a promising source of dietary fiber. Since approximately 50% of potato peels (w/w) is dietary fibers (23). (24) studied the physical and chemical characteristics of PP and reported PP as being superior to wheat bran in its content of total dietary fiber, water holding capacity, and low quantities of starchy components. Wheat flour was also substituted with PP in the production of white bread, but it increased crumb darkening and reduced the loaf volume (24).

(25) found that PP caused a musty odor in breads, but the extrusion of potato peel before its utilization can diminish this aroma from the final product. (26) mentioned the effect of using PP in biscuit processing. The resulted biscuits produced using 5 and 10 % (w/w) of PP replaced wheat flour were smaller in terms of their stack weight and sensory score (color and appearance) in proportion to the levels used. In addition, supplementary biscuits were harder to bite than control biscuits.

(27) found that muffins with 25 % PP were darker, lower in height and more resistant to compression. Similarly, cookies with 10 and 15 % PP (w/w) were darker, harder, and smaller in diameter than control cookies. From a different prospect, (23) suggested the possible effect of potato peel as a good source of dietary fiber to be acting as anticarcinogenic material. Dietary fibers are known as protective materials against mutagenesis and carcinogenesis via several well established mechanisms that include binding of carcinogenic and mutagenic substances, reduce intestines-transit time, increase water absorption and fecal bulk, and lower fecal pH through the fermentation by intestinal

microflora (28). The probable role of extruded vs. unextruded potato peel, in comparison with wheat bran and cellulose as other sources of dietary fiber, in binding of the carcinogenic benzopyrene in designed in-vitro digestion model were investigated(16).

Un-extruded steam peels bound more benzopyrene than did peels extruded at temperature of 110 °C and feed moisture of 30%. Moreover, binding of benzopyrene by abrasion potato peels was lower than that by steam potato peels. In addition, abrasion peels contain approximately 25% dietary fiber and 50% starch, which may have maintained benzopyrene in suspension and / or higher amount of polyphenols remaining in abrasion peel, in contrast with steam peel, which may also have role in lowering of levels of benzopyrene by interaction and binding (16).

The method of peeling was found to be a key factor influencing the chemical composition of peels and its suitability for further utilization. (29) compared the influence of peeling method on composition of PP. Abrasion, method of peeling used by potato chip manufacturers, results in more starch and less dietary fiber than the steam peeling method used in the production of dehydrated potatoes. Potato peels with either abrasion or steam peeling methods were extruded; at barrel temperatures of either 110 °C or 150 °C and fed moistures of either 30 or 35%. Extrusion was associated with an increase in total dietary fiber and lignin contents and a decrease in starch content in steam peels. Lignin content was found to decrease but total dietary fiber content was unaffected in extruded abrasion peels. Soluble nonstarch polysaccharides increased in both types of peeling because of extrusion.

The higher quantity of glucose recovered from the insoluble fiber fraction of extruded steam peels was reported to be the possible reason in the formation of resistant starch. On the other hand, manual peeling of potato produced peels with approximately 63%, on a dry weight basis of alcohol-insoluble fibers, which was separated into pectic substances, hemicellulose, cellulose, and lignin. These fractions consisted of 3.4% pectin, 2.2% cellulose, 14.7% protein, 66.8% starch, and 7.7% ash. The sugars in the alcohol-soluble fraction consisted of 1.4% total soluble sugars and 0.9% reducing sugars (30).

3. Potato peels contribution in biotechnology

Several agro-industrial byproducts such as wheat bran, rice bran, molasses bran, barley bran, maize meal, soybean meal, potato peel and coconut oil cake have been screened as low-cost solid substrates for microbial production of enzymes to be use either in food applications or in other industrial sectors. Potato peel was reported to be an excellent substrate for the production of thermostable alpha-amylase, a starch hydrolyzing enzyme extensively used in different food industries, under solid-state controlled growth conditions (31) and was successfully used in some applications (32). Moreover, (33) and (34) used potato peel as a low-cost agro-industrial medium in production of both alpha-amylase and alkaline protease enzymes, respectively, to be used as detergents.

The antibacterial activity of the potato peel was found to be species dependent and the water extract of potato peel was effective only at high concentration against gram negative and one gram positive bacteria (35). (12) also investigated the extract of potato peel as antimicrobial agent against different types of bacteria and fungi. The results indicated the antibacterial and antifungal activities of potato peel to be species independent and when

compared to one of antibiotics, streptomycin, potato peel had significant effect against *Pseudomonas aeruginosa* and *Clavibacter michiganensis*. In addition to its antimicrobial properties, potato peel can also function as a metal binder, probably due to its contents of polyphenolic compounds.

Potato peel also was able to support the economical growth and production of several extracellular hydrolytic enzymes, using *Bacillus subtilis* (30). (36) achieved the highest productivity, in terms of amount and activity, using *Bacillus amyloliquefaciens* grown on solid media of potato peels, compare to peel samples of several fruits and vegetables, to produce β -mannanase (Figure 1). The growing interest in mannanase production for Industrial applications is due to its importance in the bioconversion of agro-industrial residues, which randomly hydrolyzes the axial chain of hetero-mannans, the major softwood hemicellulose (37).




Fig. 1. Specific activity of mannanase produced by *B. amyloliquefaciens* grown on different lignocellulosic materials. Source {Mabrouk, 2008 #87932}

Mannanases have been tested in several industrial processes, such as extraction of vegetable oils from leguminous seeds, viscosity reduction of extracts during the manufacturing of instant coffee and oligosaccharides as well as its importance in textile and paper industries. (38,39).

4. Potato peels in food applications

Potato peel also has acquired attention as a natural antioxidant in food system due to its high content of polyphenols, which was reported to be 10 times higher than their levels in the flesh (40) accounting for approximately 50 % of all polyphenols in potato tuber (41). Therefore, the effective utilization of potato peel as an antioxidant in food has been investigated extensively. Synthetic antioxidants, especially butylated hydroxytoluene (BHT) and butylated hydroxyanisole (BHA) are commonly used to prevent the oxidation process (42).

Concerns have been raised regarding the use of these synthetic antioxidants as being toxic and carcinogenic (43). Synthetic antioxidants may cause liver swelling and disturb the normal levels of liver enzymes (44). There is a strong need for alternative and effective antioxidants, from natural sources, to prevent deterioration of foods.

One of the first attempts to re-utilize PP was using it as a natural antioxidant in food systems. The oxidative deterioration of fats and oils is responsible for rancid odors and flavors, with a consequent decrease in nutritional quality and safety caused by the formation of secondary, potentially toxic, compounds. Thus, addition of antioxidants is an important step in processing to preserve flavor and color deterioration and to avoid nutrient loss. Due to the concerns of synthetic antioxidants in humans' foods, potato peel and extracts was proposed as a possible potential antioxidant in food systems (45). The same authors also measured the composition of fatty acids and polyphenolic content in peel of different potato varieties they also measured the antioxidant capacity of the petroleum-ether extracts in comparison with commercial antioxidants, tertiary butylhydroquinone (TBHQ), butylated hydroxyanisole, and butylated hydroxytoluene. Based on the results of their work, they suggested potato peel extract as a possible effective natural antioxidant retarding the oxidation process in vegetable oils; in particular colored skin-potato varieties which have been shown to contain the highest antioxidant activity reflecting high content of polyphenols (anthocyanins) that are responsible for their appearance (45).

(46) measured both amounts and type of phenolic compounds in freeze-dried, aqueous extract of potato peel. Chlorogenic (50.31%), gallic (41.67%), protocatechuic (7.815%), and caffeic (0.21%) acids were the major phenolic compounds detected in this study. They also measured the stability of sunflower oil heated at 63 °C when spiked with individual phenolic compounds that were detected in extract and with potato peel-water extract, respectively. When compared with control, sunflower oil in presence of aqueous potato peel extract was superior to BHT but not ascorbic acid derivative in preventing rancidity-related deteriorations in heated oil. On the other hand, (47) qualitatively and quantitatively, assessed polyphenols in either methanolic extract at 4 °C or water extract at 25 and 100 °C, respectively, of potato peel. Methanol extract of PP was shown to contain highest amounts of total polyphenolic compounds than water extract. (46). (47) reported that the main polyphenol detected in potato peel extract was chlorogenic acid. In terms of its safety as natural antioxidants in foods, potato peel freeze-dried extract was found not to be mutagenic when tested with *Salmonella typhimurium*-*Escherichia coli* microsome assay (35).

Moreover, the extract of potato peel had both bactericidal and bacteriostatic effects but only at a high concentration. Thus, potato peel was suggested as a possible safe, natural antioxidant to preserve foods.

More recently, potato peels, as well as some other agro-industrial by products, were examined for their content of total polyphenols and related antioxidant capacities (48). When extracted with mixture of 0.1 % of HCL in methanol: acetone: water (60:30:10, v/v/v), potato peel was found to contain approximately 177mg of total polyphenols /100 gm of fresh weight of peel.

In addition, (48) measured the total antioxidant capacity of potato peel extracts by measuring their antiradical scavenging activities and reducing power. A strong positive correlation was observed between total polyphenolic compounds in potato peel extracts and their antioxidant potency.

Lipid oxidation is a major cause of muscle food deterioration, affecting color, flavor, texture, and nutritional value ((49) and ((50). This oxidative deterioration of muscle involves the oxidation of the unsaturated fatty acids, catalyzed by hemoproteins as well as non-heme iron. (51) used freeze-dried extract of potato peel at two different levels of 500 and 1000 ppm as a natural antioxidant in ground beef patties to measure the efficacy of PP extract against the commercial antioxidant, control.

Freeze-dried extract of potato peel showed maximum antioxidant capacity at pH range of 5-6 while lost this activity at neutral and alkaline pH. In addition, antioxidant capacity, as measured with β -carotene linolate method, was significantly decreased when freeze-dried extract was boiled for more than 30 minutes and at temperature higher than 80 °C. (51) also studied the effect of heat, time, and pH on stability of potato peel extract. Further development in terms of amount being use and optimum storage conditions of food product was suggested.

(52) explored if the potato peel extract possess antioxidant potentiality in retarding lipid oxidation in lamb meat preserved with radiation. Radiation processing is one of the most effective technologies that can extend shelf life and eliminate pathogenic bacteria in raw meat and meat products. However, meat on irradiation may undergo pronounced oxidative changes that influence the sensory quality of meat (53). When added to lamb meat in ratio of 0.04%, before irradiation process, potato peel extract was able to retard lipid oxidation in a similar manner as butylated hydroxytoluene (BHT). More recently, potato peel extract were studied as a natural antioxidant to prevent deteriorations of food lipids (54). Using six different solvents and two other food processing by-products.

Compared to sesame cake and sugar beet pulp, potato peel was superior in its content of polyphenolic compounds also showed the strongest antioxidant capacity using three different confirming assays (54).

(55) and his colleagues also evaluated the effectiveness of potato peel extract, as natural antioxidant during 60 days storage of refined soy-bean oil at 25 and 45 °C. Free fatty acids, peroxide values, and iodine values were the criteria to assess the antioxidant activity of potato peels extract. Different organic solvents, including ethanol, methanol, acetone, hexane, petroleum-ether, and diethyl ether, were used to prepare extracts of potato peels.

The maximum efficiency of extraction was achieved by petroleum ether meanwhile diethyl ether, methanol, hexane, ethanol, and acetone, respectively, had had lower amounts of polyphenolic compounds in the extract. Potato peel extracted with petroleum ether exhibited strongest antioxidant activity in soybean oil during storage, equal almost to the antioxidant activity of synthetic antioxidants (BHA and BHT). However, the level of potato peel extract needed was 8–12 times higher than that of the synthetic antioxidants to control the development of rancidity during storage of cooking oils at high temperature. They suggested that potato peel extract in oils, fats and other food products can safely be used as natural antioxidant to suppress lipid oxidation (55).

Polyphenols in plants are present in either free form or bound to cell wall polysaccharides. Not all polyphenols are, therefore, readily extracted resulting in their underestimation by many researches.

(56) and (57) mention the incompleteness of data related to polyphenols in potato peel. They discussed the presence of other polyphenols in potato peel that were ignored by previous works. Principally they were interested to figure out the amount of ferulic acid-sugar esters in potato peels since many established works showed its ability to suppress the LDL-cholesterol oxidation in human plasma (58).

In their work (59) and his colleagues studied the thermal stability of ferulic acid which affects adhesion and texture of plant foods. (57) measured both free and bound forms of polyphenols in potato peel and their related free radical scavenging activities. Their results showed the superiority of bound- form of polyphenols, ferulic acid, in potato peels to quench free radicals, as measured by free radical scavenging activity technique, and strongly recommended it as a strong natural antioxidant.

5. Potentiality of potato peels antioxidants in biological systems

In light of above investigations, the potentiality of potato peel or potato peel extracts to act as antioxidant in biological systems have draw great scientific attention. The antioxidant potency of freeze-dried aqueous extract of potato peel was investigated, employing various established antioxidant measurement techniques in vitro; lipid peroxidation in rat liver homogenate, 1,1-diphenyl-2-picrylhydrazyl (DPPH), as well as superoxide/hydroxyl radical scavenging, reducing power, and iron ion chelation (60).

The freeze-dried aqueous extract of potato peel powder showed strong inhibitory activity toward lipid peroxidation in rats' liver homogenate induced by the $\text{FeCl}_2\text{-H}_2\text{O}_2$ system. Furthermore, the water extract of PP exhibited a strong concentration-dependent inhibition of deoxyribose (DNA) oxidation.

The antioxidant activity measurements of potato peel extract did emphasis its strong reducing power activity, superoxide scavenging ability as well as its ion chelating potency. The in vitro results suggest the possibility that potato peel waste could be effectively employed as an ingredient in health or functional food, to alleviate oxidative stress (60).

In patients with diabetes mellitus, either type I or type II, it is well known that the production of ROS and lipid peroxides (LPO) increase (61). It has been suggested that oxidative stress (OS) is responsible for the pathophysiology of diabetes (62). The OS was reported to be related to hyperglycemia (63).

Hyperglycemia causes nonenzymatic glycation of protein through the Maillard reaction and alters energy metabolism, which results in an elevation of ROS levels and further development of diabetic complications (64). Potato peel was found to influence both glycemic index and antioxidant status in streptozotocin (STZ)-induced diabetic male Wistar rats (65). In that study, diabetic rats fed potato peel-powder-supplemented diet for 4 weeks showed significant decrease in blood glucose levels. In addition, Incorporation of PP powder into the diet reduced significantly the hypertrophy of both liver and kidney in STZ-diabetic rats and normalized the activities of serum alanine-aminotransferase (ALT) and aspartate-aminotransferase (AST).

Serum enzymes such as aspartate aminotransferase and alanine aminotransferase are employed in the evaluation of hepatic disorders and the increase in their levels reflects acute

liver damage and inflammatory hepatocellular disorders (66). In addition, (65) measured both hepatic and renal-lipids oxidation, glutathione as well as activities of various antioxidant enzymes in liver and kidney of STZ-diabetic rats. It is suggested that PP powder in the diet may also be able to attenuate the eye lens damage associated with the diabetic conditions.

The strong antioxidant properties of potato peel extract (PP), which have been attributed to high content of polyphenolic compounds, was tested as a protective agent against erythrocytes-oxidative induced-damage in vitro (67) and by the measurement of morphological alterations and the structural alterations in the cell membrane. The total polyphenolic content in aqueous PP in this experiment was found to be 3.93 mg/g, dry weight based.

The major phenolic acids present were: gallic acid, caffeic acid, chlorogenic acid and protocatechuic acid. An experimental pro-oxidant system was used to induce lipid peroxidation in rat red blood cells (RBCs) and human RBC membranes. PP was found to inhibit lipid peroxidation with similar effectiveness in both the systems (about 80-85% inhibition by PP at 2.5 mg/ml).

While PP *per se* did not cause any morphological alteration in the erythrocytes, under the experimental conditions used, PP significantly inhibited the H₂O₂-induced morphological alterations in rat RBCs as revealed by scanning electron microscopy. (67) also suggested PP extract as a significant protector of human erythrocyte membrane proteins from oxidative damage probably by acting as a strong antioxidant. By virtue of its unique vascular and metabolic features, the liver is exposed to absorbed drugs and xenobiotics in concentrated form.

Detoxification reactions (phase I and phase II) metabolize xenobiotics aiming to increase substrate hydrophilicity for excretion. Drug-metabolizing enzymes detoxify many xenobiotics but bioactivate or increase the toxicity of others (68). In case of bioactivation, the liver is the first organ exposed to the damaging effects of the newly formed toxic substance.

Therefore, protective mechanisms relevant to the liver are of particular interest. Because free radicals and reactive oxygen species play a central role in liver diseases pathology and progression, dietary antioxidants have been proposed as therapeutic agents counteracting liver damage (69). (70) predicted a possible protective role of the potato peel extract against induced liver injury in rats.

Prior to the treatment with tetrachlorocarbon (CCl₄), the animals received doses of 100 mg, freeze-dried aqueous potato peel extract, / kg of body weight for seven consecutive days. The animals receiving an acute dose of carbon tetrachloride (CCl₄) to induce liver injury followed this. Liver enzymes as well as biomarkers of oxidative stress either in liver or in blood serum were measured. The administration of PP to rats orally at 100 mg/kg body weight/day reduced the risk of hepatic damage induced as a result of CCl₄ administration, as observed by evaluating of serum lactate dehydrogenase (LDH), aspartate aminotransferase (AST), and alanine aminotransferase (ALT) levels in CCl₄-treated rats. Also, the deteriorations of malondialdehyde (MDA); which is one of the end products of lipid peroxidation in the liver tissue, and depletion of hepatic glutathione (GSH) level; which is an important indicator of oxidative stress (71), were significantly higher in CCl₄-treated rats in comparison with animals that were pretreated with potato peel extract.

Super oxidase dismutase (SOD), catalase (CAT), and glutathione peroxidase (GPX) constitute a mutually supportive team of antioxidant defense against reactive oxygen species. SOD is the first line of antioxidant defense and it hastens the dismutation of superoxide radical to H₂O₂, while CAT/GPX converts H₂O₂ to water (72).

According to this scenario, a significant increase in SOD activity as a defense against the presence of free radicals can, generate overwhelmingly large amounts of H₂O₂. This in turn would disable the organism to handle the excessive H₂O₂, which would again result in oxidative stress.

Thus, increase in SOD accompanied by a decrease in CAT/GPX activity indicates a state of oxidative stress due to free radical generation (73). (70) elucidated the significant positive correlation between dosing of potato peel extract and SOD levels in rats treated with CCl₄. Additionally, histopathological studies showed that CCl₄ induced a remarkable degeneration in hepatocytes and focal necrosis. However, in general, lesions were markedly diminished in rats that were pre-administered with potato peel extract prior CCl₄ administration.

Study	Results	Reference
Solubility of PP fibres and its ability to bind bile salts	Are primarily insoluble, and can bind bile acids in-vitro. It is believed that binding of bile acids is one of the mechanisms whereby certain sources of dietary fibers lower plasma cholesterol	{Camire, 1993 #88855}
The hypocholesterolemic effect of dietary fiber from PP	40 % reduction in plasma cholesterol content and 30% of hepatic fat cholesterol levels were reduced in rats fed on potato peels compared to normal diet	{Lazarov, 1996 #88875}
Physical and chemical characteristics of PP and substitution of PP wheat flour in the production of white bread	Being superior to wheat bran in its content of total dietary fiber, water holding capacity, and low quantities of starchy components. Potato peels increased crumb darkening and reduced the loaf volume	{Toma, 1979 #88853}
The probable role of extruded vs. unextruded potato peel, in comparison with wheat bran and cellulose as other sources of dietary fiber, in binding of the carcinogenic benzopyrene in designed in-vitro digestion model	1. Un-extruded steam peels bounds more benzopyrene than did peels extruded at temperature of 110 °C and feed moisture of 30%. 2. Binding of benzopyrene by abrasion potato peels was lower than that by steam potato peels. 3. Abrasion peels contain approximately 25% dietary fiber and 50% starch, which may have maintained benzopyrene in suspension and / or higher amount of polyphenols remaining in abrasion peel, in contrast with steam peel, which may also have role in lowering of levels of benzopyrene by interaction and binding	{Camire, 1993 #88855}.

Raw and extruded PP fibres in the production of bread	PP caused a musty odor in breads, but the extrusion of potato peel before its utilization can diminish this aroma from the final product.	{Orr, 1982 #88797}
The effect of using PP in biscuit processing.	The resulted biscuits produced using 5 and 10 % (w/w) of PP replaced wheat flour were smaller in terms of their stack weight and sensory score (color and appearance) in proportion to the levels used. In addition, supplementary biscuits were harder to bite than control biscuits.	{Abdel-Magied, 1991 #88876}
Potato peels in the production of muffins and cookies	Muffins with 25 % PP were darker, lower in height and more resistant to compression. Similarly, cookies with 10 and 15 % PP (w/w) were darker, harder, and smaller in diameter than control cookies.	{Arora, 1994 #86730}
The influence of peeling method on chemical composition of PP.	1. Abrasion method used by potato chip manufacturers, results in more starch and less dietary fiber than the steam peeling method used in the production of dehydrated potatoes. 2. Extrusion was associated with an increase in total dietary fiber and lignin contents and a decrease in starch content in steam peels. 3. Lignin content was found to decrease but total dietary fiber content was unaffected in extruded abrasion peels. 4.Soluble nonstarch polysaccharides increased in both types of peeling because of extrusion.	{Camire, 1997 #88802}
Manual peeling of potato peels	Approximately 63%, on a dry weight basis of alcohol-insoluble fibers was separated into pectic substances, hemicellulose, cellulose, and lignin. These fractions consisted of 3.4% pectin, 2.2% cellulose, 14.7% protein, 66.8% starch, and 7.7% ash. The sugars in the alcohol-soluble fraction consisted of 1.4% total soluble sugars and 0.9% reducing sugars	{Mahmood, 1998 #88884}.

Potato peel used as a low-cost agro-industrial medium in production of both alpha-amylase and alkaline protease enzymes and several extracellular hydrolytic enzymes	High yield and high activity of the produced enzymes	{Mukherjee, 2008 #88896}, {Mukherjee, 2009 #88895}, {Fadel, 1999 #88873} and {Fadel, 2002 #88874} {Mahmood, 1998 #88884}. {Mabrouk, 2008 #88880} {McCleary, 1988 #88881} {Gubitz, 1996 #88882; Ademark, 1998 #88883}.
The extract of potato peel as antimicrobial.	Antibacterial and antifungal activities of potato peel to be species independent	{Prasad, 2007 #88515}
Potato peel extracts was proposed as a possible potential antioxidant in food systems	Extract was superior to BHT but not ascorbic acid derivative in preventing rancidity-related deteriorations in heated oil	{Onyeneko, 1993 #88898} {Rodriguez De Sotillo, 1994 #80192} ({Rhee, 1996 #88903} ({Yin, 1997 #88904}., {Makrisa, 2007 #88902} {Mansour, 2000 #80194} {Formanek, 2003 #88897}. {Kanatt, 2005 #89686} {Mohdaly, 2010 #88957}. {Mohdaly, 2010 #88957}. {Rehman, 2004 #86733} {Rehman, 2004 #86733}.
Amounts and type of phenolic compounds in freeze-dried, aqueous extract of potato peel.	1. Chlorogenic (50.31%), gallic (41.67%), protocatechuic (7.815%), and caffeic (0.21%) acids were the major phenolic compounds detected in this study. 2. Qualitatively and quantitatively, assessed polyphenols in either methanolic extract at 4 °C or water extract at 25 and 100 °C, respectively, of potato peel. Methanol extract of PP was shown to contain highest amounts of total polyphenolic compounds than water extract. reported that the main polyphenol detected in potato peel extract was chlorogenic acid.	{Rodriguez De Sotillo, 1994 #80192} {Ishii, 1997 #88910}. {Nara, 2006 #45746} {Rodriguez De Sotillo, 1994 #80191}

Safety of potato peels extract when used as natural antioxidants in foods,	Potato peel freeze-dried extract was found not to be mutagenic when tested with <i>Salmonella typhimurium</i> - <i>Escherichia coli</i> microsome assay	{Rodriguez De Sotillo, 1998 #80193}.
The potentiality of potato peel or potato peel extracts to act as antioxidant in biological systems	1. Inhibitory activity toward lipid peroxidation in rats' liver homogenate induced by the FeCl ₂ -H ₂ O ₂ system. the water extract of PP exhibited a strong concentration-dependent inhibition of deoxyribose (DNA) oxidation. 2. Potato peel was found to influence both glycemic index and antioxidant status in streptozotocin (STZ)-induced diabetic male Wistar rats reduced significantly the hypertrophy of both liver and kidney and normalized the activities of serum alanine-aminotransferase (ALT) and aspartate-aminotransferase (AST). 3. PP extract as a significant protector of human erythrocyte membrane proteins from oxidative damage probably by acting as a strong antioxidant.	{Singh, 2004 #79 {Singh, 2005 #88890}. 926}. {Singh, 2005 #88890}
Protective role of the potato peel extract against induced liver injury using CCL ₄ in rats.	1. The administration of PP to rats orally at 100 mg/kg body weight/day reduced the risk of hepatic damage induced as a result of CCL ₄ administration, as observed by evaluating of serum lactate dehydrogenase (LDH), aspartate aminotransferase (AST), and alanine aminotransferase (ALT) levels in CCL ₄ -treated rats. 2. Significant positive correlation between dosing of potato peel extract and SOD levels in rats treated with CCL ₄ 3. histopathological studies showed that CCL ₄ induced a remarkable degeneration in hepatocytes and focal necrosis. However, in general, lesions were markedly diminished in rats that were pre-administered with potato peel extract prior CCL ₄ administration.	{Singh, 2008 #87208}

Table 2. Summary of potato peels (PP) attempts of reutilization.

6. References

- [1] Schieber A, Stintzing FC, Carle R. By-products of plant food processing as a source of functional compounds-recent developments. *Trends in Food Science and Technology* 2001;12(11):401-413.
- [2] Chiellini E, Cinelli P, Chiellini F, Imam SH. Environmentally degradable bio-based polymeric blends and composites. *Macromolecular Bioscience* 2004;4:218-231.
- [3] Bildstein MO, García J, Faraldi M, Colvine S. Value-added for the european tomato processing industry. *Acta Horticulturae* 2009;823:195-198.
- [4] Di Mauro A, Arena E, Fallico B, Passerini A, Maccarone E. Recovery of anthocyanins from pulp wash of pigmented oranges by concentration on resins *Journal of Agricultural and Food Chemistry* 2002;50(21):5968-5974.
- [5] Bhushan S, Kalia K, Sharma M, Singh K, Ahuja PS. Processing of apple pomace for bioactive molecules. *Critical Reviews in Biotechnology* 2008;28:285-296.
- [6] Chiu SW, Chan SM. Production of pigments by *Monascus purpureus* using sugar-cane bagasse in roller bottle cultures. *World Journal of Microbiology and Biotechnology* 1992;8(1):68-70.
- [7] FAOSTAT. Volume 2009; 2009.
- [8] Bradshaw JE, Ramsay G, Jaspreet S, Lovedeep K. Potato Origin and Production. *Advances in Potato Chemistry and Technology*. San Diego, U.S.A.: Academic Press; 2009. p 1-26.
- [9] Natu RB, Mazza G, Jadhav SJ. Waste utilization. In: Salunkhe DK, Kadam SS, Jadhav SJ, editors. *Potato: production, processing, and products*: CRC Press, Boca Raton, U.S.A.; 1991. p 175-199.
- [10] Birch GG, Blakebrough N, Parker KJ. *Enzymes and Food Processing*. London: Applied Science Publishers Ltd; 1981. 296 p. p.
- [11] Jurjanz S, Colin-Schoellen O, Gardeur JN, Laurent F. Alteration of milk fat by variation in the source and amount of starch in a total mixed diet fed to dairy cows. *Journal of Dairy Science* 1998;81(11):2924-2933.
- [12] Prasad AGD, Pushpa HN. Antimicrobial activity of potato peel waste. *Asian Journal of Microbiology, Biotechnology and Environmental Sciences* 2007;9(3):559-561.
- [13] Forsythe WA, Chenoweth WL, Bennink MR. The effect of various dietary fibres on serum cholesterol and laxation in the rat. *Journal of Nutrition* 1976;106:26-32.
- [14] Ballesteros MN, Cabrera RM, Saucedo MS, Yepiz-Plascencia GM, Ortega MI, Valencia ME. Dietary fiber and lifestyle influence serum lipids in free living adult men. *Journal of the American College of Nutrition* 2001;20(6):649-655.
- [15] Gallaher D, Schneeman BO. Dietary fibre. In: Bowman AB, Russell MR, editors. *Present knowledge in nutrition*. Washington, DC, USA: ILSI; 2001. p 805.
- [16] Camire ME, Zhao J, Violette DA. In vitro binding of bile acids by extruded potato peels. *Journal of Agricultural and Food Chemistry* 1993;41(12):2391-2394.
- [17] Lazarov K, Werman MJ. Hypocholesterlaemic effect of potato peels as a dietary fiber source. *Journal of Medical Sciences Research* 1996;24:581-582.
- [18] Bagger M, Andersen O, Nielsen JB, Rytting KR. Dietary fibres reduce blood pressure, serum total cholesterol and platelet aggregation in rats. *British Journal of Nutrition* 1996;75(3):483-493.
- [19] Erkkila AT, Lichtenstein AH. Fiber and cardiovascular disease risk: how strong is the evidence? *Journal of Cardiovascular Nursing* 2006;21(1):3-8.


- [20] Alonso A, Beunza JJ, Bes-Rastrollo M, Pajares RM, Martinez-Gonzalez MA. Vegetable protein and fiber from cereal are inversely associated with the risk of hypertension in a Spanish cohort. *Archives of Medical Research* 2006;37(6):778-786.
- [21] Onyechi UA, Judd PA, Ellis PR. African plant foods rich in non-starch polysaccharides reduce postprandial blood glucose and insulin concentrations in healthy human subjects. *British Journal of Nutrition* 1998 80(5):419-428.
- [22] Chandalia M, Garg A, Lutjohann D, von Bergmann k, Grundy SM, Brinkley LJ. Beneficial effect of high dietary fiber intake in patients with type 2 diabetes mellitus. *The New England Journal of Medicine* 2000;342:1392-1398.
- [23] Camire ME, Zhao J, Dougherty MP, Bushway RJ. In vitro binding of benzo[a]pyrene by extruded potato peels. *Journal of Agricultural and Food Chemistry* 1995;43:970-973.
- [24] Toma RB, Orr PH, D'appolonia B, Dintzis FR, Tabekhia MM. Physical and chemical properties of potato peel as a source of dietary fiber in bread. *Journal of Food Science* 1979;44:1403-1407.
- [25] Orr PH, Toma RB, Munson ST, D'Appolonia B. Sensory evaluation of breads containing various levels of potato peel. *American Potato Journal* 1982;59(12):605-612.
- [26] Abdel-Magied MM. Effect of dietary fiber of potato peel on the rheological and organoleptic characteristics of biscuits. *Egyptian Journal of Food Science* 1991;19:293-300.
- [27] Arora A, Camire ME. Performance of potato peels in muffins and cookies. *Food Research International* 1994;27:15-22.
- [28] Harris PJ, Ferguson LR. Dietary fibre: Its composition and role in protection against colorectal cancer. *Mutation Research* 1993;290:97-110.
- [29] Camire ME, Violette D, Dougherty MP, McLaughlin MA. Potato peel dietary fiber composition: effects of peeling and extrusion cooking processes. *Journal of Agricultural and Food Chemistry* 1997;45(4):1404-1408.
- [30] Mahmood AU, Greenman J, Scragg AH. Orange and potato peel extracts : Analysis and use as *Bacillus* substrates for the production of extracellular enzymes in continuous culture. *Enzyme and Microbial Technology* 1998;22(2):130-137.
- [31] Fadel M. Utilization of potato chips industry by products for the production of thermostable bacterial alpha amylase using solid state fermentation system: 1. Effect of incubation period, temperature, moisture level and inoculum size. *Egyptian Journal of Microbiology* 1999;34:433-445.
- [32] Fadel M. Utilization of potato chips industry by products for production of thermostable bacterial alpha-amylase using solid state fermentation system. 2. Effect of moistening agent, supplementary substrates nitrogen source, and application of the solid fermented substrate for some starches digestion. *Egyptian Journal of Microbiology* 2002;34(4):533-546.
- [33] Mukherjee KA, Adhikari H, Rai KS. Production of alkaline protease by a thermophilic *Bacillus subtilis* under solid-state fermentation (SSF) condition using *Imperata cylindrica* grass and potato peel as low-cost medium: Characterization and application of enzyme in detergent formulation. *Biochemical Engineering Journal* 2008;39:353-361.

- [34] Mukherjee KA, Borahb M, Rai KS. To study the influence of different components of fermentable substrates on induction of extracellular α -amylase synthesis by *Bacillus subtilis* DM-03 in solid-state fermentation and exploration of feasibility for inclusion of α -amylase in laundry detergent formulations. *Biochemical Engineering Journal* 2009;43:149-156.
- [35] Rodriguez De Sotillo D, Hadley M, Wolf-Hall C. Potato peel extract a nonmutagenic antioxidant with potential antimicrobial activity. *Journal of Food Science* 1998;63(5):907-910.
- [36] Mabrouk EMM, El Ahwany MDA. Production of β -mannanase by *Bacillus amylolequifaciens* 10A1 cultured on potato peels. *African Journal of Biotechnology* 2008;7(8):1123-1128.
- [37] McCleary BV. β -Mananasa. *Methods in Enzymology* 1988;160:596-610.
- [38] Gubitz GM, Hayn M, Urbanz G, Steiner W. Purification and properties of an acid β -mannanase from *Sclerotium rolfii*. *Journal of Biotechnology* 1996;45:165-172.
- [39] Ademark P, Varga A, Medve J, Harjunpa AV, Drakenberg T, Tjerneld F. Softwood hemicellulose-degrading enzymes from *Aspergillus niger*: Purification and properties of a β -mannanase. *Journal of Biotechnology* 1998;63:199-210.
- [40] Malmberg A, Theander O. Free and conjugated phenolic acids and aldehydes in potato tubers. *Swedish Journal of Agricultural Research* 1984;14:119-125.
- [41] Friedman M. Chemistry, biochemistry, and dietary role of potato polyphenols. *Journal of Agricultural and Food Chemistry* 1997;45(5):1523-1540.
- [42] Sobedio JL, Kaitaramita J, Grandgirál A, Malkkl Y. Quality assessment of industrial pre-fried French fries. *American Oil Chemical Society* 1991;68:299-302.
- [43] Ito N, Hiroze M, Fukushima G, Tauda H, Shira T, Tatematsu M. Studies on antioxidant; their carcinogenic and modifying effects on chemical carcinogenesis. *Food Chemistry Toxicology* 1986;24:1071-1081.
- [44] Madhavi DL, Deshpande SS, Salunkhe DK. Butylated hydroxyanisole (BHA; tert-butyl-4-hydroxyanisole) and Butylated hydroxytoluene (BHT; 2,6-di-tert-butyl-p-cresol). In: Madhavi DL, Deshpande SS, Salunkhe DK, editors. *Food antioxidants: Technological, toxicological, and health perspectives*. New York, U.S.A.: Marcel Dekker; 1996. p 1-215.
- [45] Onyeneho NS, Hettiarachchy SN. Antioxidant activity, fatty acids and phenolic acids compositions of potato peels. *Journal of the Science of Food and Agriculture* 1993;62:345-350.
- [46] Rodriguez De Sotillo D, Hadley M, Holm ET. Potato peel waste: Stability and antioxidant activity of a freeze-dried extract. *Journal of Food Science* 1994;59(5):1031-1033.
- [47] Rodriguez De Sotillo D, Hadley M, Holm ET. Phenolics in aqueous potato peel extract: Extraction, identification and degradation. *Journal of Food Science* 1994;59(3):649-651.
- [48] Makrisa PM, Boskou G, Andrikopoulos KN. Polyphenolic content and in vitro antioxidant characteristics of wine industry and other agri-food solid waste extracts. *Journal of Food Composition and Analysis* 2007;20:125-132.
- [49] Rhee KS, Anderson LM, Sams AR. Lipid oxidation potential of beef, chicken and pork. *Journal of Food Science* 1996;61:8-12.

- [50] Yin MC, Cheng WS. Oxymyoglobin and lipid oxidation in phosphatidylcholine liposomes retarded by α -tocopherol and β -carotene. *Journal of Food Science* 1997;62:1095-1097.
- [51] Mansour EH, Khalil AH. Evaluation of antioxidant activity of some plant extracts and their application to ground beef patties. *Food Chemistry* 2000;69(2):135-141.
- [52] Kanatt SR, Chander R, Radhakrishna P, Sharma A. Potato peel extract: a natural antioxidant for retarding lipid peroxidation in radiation processed lamb meat. *Journal of Agricultural and Food Chemistry* 2005;53(5):1499-1504.
- [53] Formanek Z, Lynch A, Galvin K, Farkas J, Kerry JP. Combined effects of irradiation and the use of natural antioxidants on the shelf-life stability of overwrapped minced beef. *Meat Science* 2003;63:433-440.
- [54] Mohdaly AA, Sarhan MA, Smetanska I, Mahmoud A. Antioxidant properties of various solvent extracts of potato peel, sugar beet pulp and sesame cake. *Journal of the Science of Food and Agriculture* 2010;90(2):218-226.
- [55] Rehman Z, Habib F, Shah WH. Utilization of potato peels extract as a natural antioxidant in soy bean oil. *Food Chemistry* 2004;85:215-220.
- [56] Fry SC. Cross-linking of matrix polymers in the growing cell walls of angiosperms. *Annual Review of Plant Physiology* 1986;37:165-186.
- [57] Nara K, Miyoshi T, Honma T, Koga H. Antioxidative activity of bound-form phenolics in potato peel. *Bioscience, Biotechnology, and Biochemistry* 2006;70(6):1489-1491.
- [58] Ohta T, Semboku N, Kuchii A, Egashira Y, Sanada H. Antioxidant activity of corn bran cell-wall fragments in the LDL oxidation system. *Journal of Agricultural and Food Chemistry* 1997;45(5):1644-1648.
- [59] Ishii T. Structure and functions of feruloylated polysaccharides. *Plant Science* 1997;127(2):111-127.
- [60] Singh N, Rajini PS. Free radical scavenging activity of an aqueous extract of potato peel. *Food Chemistry* 2004;85(4):611-616.
- [61] Kuyvenhoven JP, Meinders AE. Oxidative stress and diabetes mellitus. Pathogenesis of long-term complications. *European Journal of Internal Medicine* 1999;10:9-19.
- [62] Dandona P, Thusu K, Cook S, Synder B, Makaowski J, Armstrong D, Nicotera T. Oxidative damage to DNA in diabetes mellitus. *Lancet* 1996;347:444-445.
- [63] Baynes JW, Thorpe SR. Role of oxidative stress in diabetic complications: A new perspective on an old paradigm. *Diabetes* 1999;48:1-9.
- [64] Shimoi K, Okitsu A, Green MHL, Lowe JE, Ohta T, Kaji K, Terato H, Ide H, Kinane N. Oxidative DNA damage induced by high glucose and its suppression in human umbilical vein endothelial cells. *Mutation Research* 2001;480-481:371-378.
- [65] Singh N, Kamath V, Rajini PS. Attenuation of hyperglycemia and associated biochemical parameters in STZ-induced diabetic rats by dietary supplementation of potato peel powder. *Clinica Chimica Acta* 2005;353(1-2):165-175.
- [66] El BK, Hashimoto Y, Muzandu K, Ikenaka Y, Ibrahim ZS, Kazusaka A, Fujita S, Ishizuka M. Protective effect of *Pleurotus cornucopiae* mushroom extract on carbon tetrachloride-induced hepatotoxicity. *Japanese Journal of Veterinary Research* 2009;57(2):109-118.

- [67] Singh N, Rajini PS. Antioxidant-mediated protective effect of potato peel extract in erythrocytes against oxidative damage. *Chemico-Biological Interactions* 2008;173(2):97-104.
- [68] Jaeschke H, Gores GJ, Cederbaum AI, Hinson JA, Pessayre D, Lemasters JJ. Mechanisms of hepatotoxicity. *Toxicological Sciences* 2002;65(2):166-176.
- [69] Higuchi H, Gores GJ. Mechanisms of liver injury: An overview. *Current Molecular Medicine* 2003;3(6):483-490.
- [70] Singh N, Kamath V, Narasimhamurthy K, Rajini PS. Protective effect of potato peel extract against carbon tetrachloride-induced liver injury in rats. *Environmental Toxicology and Pharmacology* 2008;26(2):241-246.
- [71] Souza MF, Rao VSN, Silveira ER. Inhibition of lipid peroxidation by ternatin, a tetramethoxyflavone from *Egletes viscosa* L. *Phytomedicine* 1997;4(1):27-31.
- [72] Halliwell B, Gutteridge JM. Role of free radicals and catalytic metal ions in human disease: An overview. *Methods in Enzymology* 1990;186:1-85.
- [73] Tasaduq SA, Singh K, Sethi S, Sharma SC, Bedi KL, Singh J, Jaggi BS. Hepatocurative and antioxidant profile of HP-1, a polyherbal phytomedicine *Human and Experimental Toxicology* 2003;22(12):639-645.

IntechOpen



Phytochemicals as Nutraceuticals - Global Approaches to Their Role in Nutrition and Health

Edited by Dr Venketeshwer Rao

ISBN 978-953-51-0203-8

Hard cover, 278 pages

Publisher InTech

Published online 23, March, 2012

Published in print edition March, 2012

Phytochemicals are biologically active compounds present in plants used for food and medicine. A great deal of interest has been generated recently in the isolation, characterization and biological activity of these phytochemicals. This book is in response to the need for more current and global scope of phytochemicals. It contains chapters written by internationally recognized authors. The topics covered in the book range from their occurrence, chemical and physical characteristics, analytical procedures, biological activity, safety and industrial applications. The book has been planned to meet the needs of the researchers, health professionals, government regulatory agencies and industries. This book will serve as a standard reference book in this important and fast growing area of phytochemicals, human nutrition and health.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

A. Al-Weshahy and V.A. Rao (2012). Potato Peel as a Source of Important Phytochemical Antioxidant Nutraceuticals and Their Role in Human Health - A Review, *Phytochemicals as Nutraceuticals - Global Approaches to Their Role in Nutrition and Health*, Dr Venketeshwer Rao (Ed.), ISBN: 978-953-51-0203-8, InTech, Available from: <http://www.intechopen.com/books/phytochemicals-as-nutraceuticals-global-approaches-to-their-role-in-nutrition-and-health/potato-peel-as-a-source-of-important-phytochemical-antioxidant-nutraceuticals-and-their-role-in-huma>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](https://creativecommons.org/licenses/by/3.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen