We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Advances in Genetics of Non Syndromic Neural Tube Defects

Patrizia De Marco
Neurosurgery Department
G. Gaslini Institute, Genoa,
Italy

1. Introduction

This chapter revisits established and emerging information on the causes of the human Neural Tube Defects (NTD). These congenital malformations, affecting 1 per 1000 births, have adverse consequences that afflict society as well as the affected individuals. The mechanisms by which they arise are still unknown.

Investigators have focused their efforts in many research fields (developmental and metabolic pathways and animal models) to understand the mechanisms underlying the etiology of these devastating defects. Nevertheless, few candidate genes have proved to have significant impact on the development of NTDs. Moreover, the existing body of literature is still fragmented and constrained by limited size, lack of power, and lack of replicated and independent studies.

The review attempts to describe the results obtained by the analysis of candidate genes from three types of evidences: developmental pathway and biochemical pathways (folate metabolism). The update of the literature includes the most recent findings.

This chapter aims to improve the knowledge of the readers on a such complex multifactorial disease outlining that, even though the remarkable recent progresses, further studies are needed to clarify the molecular basis of NTDs.

The knowledge of the etiology can help us in developing strategies for prevention of the NTDs, by the identification of parents who have a higher chance of having an affected child. In these families it may be possible to clarify the mode of inheritance and estimate the recurrence risk of the disorder. Actually, we are far to have immediate clinical benefits for NTDs, but providing clinicians with a better knowledge of the development of the disease, we could speed up the search for a focused prevention and a more complete familial counselling.

Neural tube defects (NTDs) are congenital malformations that involve failure of the neural tube closure during the early phases of development at any level of the rostro-caudal axis. The incidence of NTDs in European countries excluding UK and Ireland was 0.1-0.6 per 1000 births in the period 1989-2002. [1] Each year, more than 4500 pregnancies in the European Union are affected by NTDs. [2]

In order to gain insights into the embryonic basis of NTDs, it is important to understand the morphogenetic processes and the underlying molecular mechanisms involving the neural tube closure. Neurulation begins with formation of the neural plate, a thickening of the ectoderm on the dorsal surface of the post-gastrulation embryo. Neurulation occurs in a two-part process: i) primary neurulation (weeks 3-4) resulting in formation of the neural tube that will develop into brain and most of spinal cord; ii) secondary neurulation (weeks 5-6) that creates the neural tube caudal to the mid-sacral region. [3] During the primary neurulation, the neural plate is subject to shaping and folding, with fusion along the midline to form the tube. The secondary neural tube is derived from a population of mesenchymal cells, the tail bud, which undergo proliferation and condensation followed by cavitation and fusion with the primary neural tube. [4] Neurulation is driven by redundant mechanisms both at the tissue and cellular level. [5]. Active processes required for neural tube closure include convergent extension cell movements, expansion of the cranial mesenchyme, contraction of actin filaments, bending of the neural plate, and adhesion of the neural folds. Insight into these mechanisms are being provided by the many mouse models in which NTDs occur as a result of genetic mutations. [6] Neural tube closure is a discontinuous process and proceed bi-directionally in a zipper-like fashion. Five closure sites exist in human embryos and on the basis of the study of clinical cases, van Allen et al. [7] proposed a multisite model in which the initial site of closure starts at the rhombencephalic segment and the caudal end is the last to close.

There are several different types of NTDs. The distribution of different types may vary in different populations and can also vary by the birth prevalence of the defects. [8] The most common classification includes anencephaly, myelomeningocele (also called spina bifida) and encephalocele. Craniorachischisis and iniencephaly are considered to be rarer in most populations. NTDs group includes other conditions, broadly described as occult spinal dysraphisms (lipoma, tight filum terminale) and some complex dysraphic states as diastematomyelia, spinal segmental dysgenesis, and caudal regression syndrome. [9] Failure of the closure at any level of the body axis leads to open NTDs where the affected nervous tissues are exposed to the surface. Failure of the secondary neurulation leads to closed forms of NTDs, where the defect is covered by the skin and it is not exposed to the surface of the body.

The majority of NTD cases are non syndromic or isolated NTDs, meaning that the NTD is the only defect. A number of anomalies including clubfoot, deformation of the lower limbs, hydrocephalus and Arnold Chiari II malformation have been widely accepted as secondary abnormalities that may accompany NTDs. Aetiology of non syndromic NTDs is multifactorial with both environmental and genetic factors implicated. Non syndromic NTDs have an important genetic component and it is important that research is focused at determining the basis of this genetic aetiology. The aim of this chapter is provide evidences for genetic contribution to non syndromic NTDs along with the discussion of different approaches for identifying such genes. The identification of genetic factors for NTDs is a relatively a new area of research. In fact, there are many different mutations in mice resulting in NTDs, but only few well characterized genetic factors in humans. Here we provide an overview of the major causative genes and the more recent genetic advances emerging thanks to the powerful tool of mouse models.

2. Evidence for genetic contribution of non syndromic NTDs

There are strong evidence supporting a genetic component to NTDs. NTDs can be associated with various chromosome rearrangements, the most common include trisomy 13, trisomy 18, tetraploidy, and deletions involving chromosomes 2, 3, 11, 13, 22. Genetic syndromes with NTDs include Meckel-Gruber syndrome and Currarino syndrome. [10] Overall heritability in NTDs is estimated to be 60% with multiple genes involved. Recurrence risk for sibling is 2-5%, which represents up to 50-fold increased risk respect to that observed in the general population [11]. Positive family history families with many affected children have implied an autosomal recessive model of inheritance. [12] Inspection of multiplex families demonstrated that affected relative pairs are related at either the second- or third-degree, suggesting oligogenic inheritance. [13]. The recurrence risk and the inheritance pattern in families have been demonstrated to follow a multifactorial threshold model in which the liability to these malformations follows a normal distribution pattern: additive effects of several genetic and environmental factors cumulate to increase the liability until a threshold value beyond which the phenotype become expressed. Twin studies demonstrated that concordance rate is relatively low among cotwins of affected cases. [11]. Recurrences in families in which the case is affected with spina bifida tend to be spina bifida and families; recurrence in families in which the case is anencephaly tend to be anencephaly. [14-15] However, one third of recurrent cases involve a NTDs phenotype that is different from the cases phenotype. This intra-familial heterogeneity may be the consequence of the pleiotropic effect of a underlying causative gene. NTD occurrence vary among different ethnic groups and it is unknown if this population-specific incidence is due to differences in the frequency of risk-associated alleles or to specific dietary habits or exposures. The highest incidence is in Northern China (7/1000) and the lowest in Japan (0.1/1000). In Europe, the highest rate is reported in Ireland (3/1000). Furthermore, an Irish study demonstrated that predisposing genetic factors may be transmitted from the mother's side of the family. [16] Another important line of evidence for a genetic basis for NTDs comes from the numerous mouse models, both spontaneous and induced via a gene-driven process. [17].

3. Identification of susceptibility genes involved in NTDs

The number and identity of genes predisposing to NTDs remain unknown. Once genes implicated in the development of NTDs are identified, it will be essential to understand potential additive or multiplicative effects of genetic and environmental factors. Candidate genes for NTDs can be classified into two categories according to how they have been identified: biological and positional. Candidate genes based on biological plausibility have resulted from research done in animal models and from biological pathways important for NTDs. Candidate genes can also be identified through positional techniques via chromosomal rearrangements in NTD patients or genomic screen in linkage analysis.

A number of genes have been identified from research done in experimental animals, most notably from mouse models. The mouse model remains one of the most well studied animal models for NTDs. More than 190 genes in mouse produce varying NTDs phenotypes. [17]. There are both naturally occurring and induced strains of mice. The two most common mouse phenotypes are exencephaly, that is comparable to human anencephaly, and craniorachischisis, in which the neural tube remains open along the entire body axis. The mouse system is useful for the study of NTDs because the genetic map of mouse is noted

and many of the human homologues are known. Moreover, mouse has a short gestational period and does not require the same ethical considerations as the studies in humans. Another advantage is that both genetic and environmental factors may be studied. For example three mouse mutants have been demonstrated to have a reduced risk for NTDs after maternal folate supplementation: *crooked* (*Cd*), *splotch* (*Sp*) and *cartilage homeoprotein1* (*Cart1*). [17] Nevertheless, there are obvious differences between mouse and human, so that it is possible that genes contributing to NTDs in mouse not always do the same in humans. The majority of human NTDs are isolated and show a multifactorial inheritance. In contrast, NTDs in mouse are a lethal condition in the homozygous knock-out state and NTDs is a part of a larger syndrome. Mice have a four-sites model of neural tube closure. In humans controversies have been arisen about the number of closure sites, although there is a general agreement on the presence of rostral and caudal sites. The majority of candidate genes from mouse are involved in a wide range of functions, in crucial steps of neurulation and only few are implicated in cellular functions such as genome stability and DNA repair. Despite a such long list of candidate genes, studies failed to show positive association in humans. Another system that can be used is the zebrafish (*Brachydanio rerio*), that is a cost-effective model system for studying NTDs since fertilized eggs can develop rapidly into transparent embryos, allowing all stages of vertebrate development to be visualized easily. [18] Zebrafish has proven to be very useful for studying embryological processes, including morphogenesis, lineage specification, and mechanisms of acquired cell fate. Furthermore, assays have been developed to perform both gain and loss of function experiments. Partial loss of function phenotypes can be generated by microinjection of antisense morpholino. [19] A forward genetic approach (random mutagenesis) will lead to identification of potential NTDs genes. Such a screen would be useful in expanding the collection of potential NTDs risk genes.

The approach for identifying positional candidate genes is through a genomic screen in which genetic markers at known locations are tested in families with two or more affected individuals. Thanks to advances in the density and polymorphisms of genetic markers, highthroughput genotyping methods, and powered statistical genetic linkage analysis, it is possible to perform large-scale investigation of human complex traits such as NTDs. Regions with excess sharing of alleles among affected individuals are considered region of interest and may harbour genes that predispose to NTDs. Few studies have performed genome-wide linkage studies in families with multiple affected individuals, given the scarcity of these large families. In fact, high morbidity and mortality rates associated with the disorder, and pregnancy termination after case identification are factors that limit the ascertainment of multiplex families. [20] A recent genomic screen was performed on 44 multiplex American NTDs families identifying linkage to chromosome 7 and 10. Two candidate genes (*Meox2* and *Twist1*) were identified on chromosome 7 and three candidates (*FGFR2*, *GFRA1*, *Pax2*) were found on critical region of the chromosome 10. [21]

Statistical tests are employed to assess evidence of allelic association for NTDs. Infact allelic association is population-specific. There are two methods of testing allelic association in NTDs candidate genes: case-control and family-based studies. Traditionally, NTDs studies have employed case-control studies, where the allelic frequency in an unrelated group of patients are compared to a group of controls individuals. This study can be flawed, if the controls are not properly matched with patients for age, gender, and geographic origin. This population admixture may result in

biased results and spurious associations. In family based studies, the frequencies of alleles transmitted by a parent to the affected child are compared with that of the allele not transmitted. The family studies can also detect a paternal or maternal effect. The most used family-based test is the Transmission disequilibrium test (TDT). [22] When one parent is not available for the analysis unaffected sibling can be used as controls. Association studies in patients with NTDs were often underpowered by a small sample size, leading to inconsistent results. Larger cohorts of patients and controls are needed to reach definitive conclusion regarding the role of the most of studied genes. Meta-analysis for variants of genes are also warrented.

4. Genetic variations in folate metabolism

A large number of experimental and observational clinical research studies over the past half century demonstrated that folic acid supplementation decreases both the occurrence and recurrence risk of NTDs by approximately 70%. [23-24] The recommended intakes are 4 mg/day for high risk women and 0.4 mg/day for all others. Maternal periconceptional folate supplementation may act to overcome insufficient maternal folate levels. However, in most cases, mothers of affected foetuses have either normal folate status or are, at most, mildly folate-deficient, arguing against maternal folate deficiency as a major causative factor. [25-26] Alternatively, it is postulated that supplemental folic acid may act to overcome an underlying defect in folate metabolism that results from a genetic mutation in the mother or in the foetus. Although a protective effect of folate is recognized, the mechanism by which some women develop low folate levels and predispose their offspring to NTDs and other congenital malformation is still debated.

Fig. 1. Folate and homocysteine metabolism

Folic acid is a water soluble B vitamin that participates in transfer of single-carbon units in several critical pathways (Figure 1). These pathways include synthesis or catabolism of amino acids (serine, glycine, histidine, and methionine) and synthesis of nucleotides. Through their role in methionine synthesis, folates are also critical for the methylation cycle, since methionine is the methyl donor for many important methylation reactions, including DNA methylation. Folate-dependent conversion of homocysteine to methionine begins early in development in all tissues and provides a link between folate and homocysteine metabolism. An elevation of plasma homocysteine, mild hyperhomocysteinemia, has become recognized as a risk factor for NTDs [25, 27].

Proteins that mediate, or are functionally associated with, folate metabolism have provided candidates for genetic analysis in human NTDs. There are over 25 proteins involved in the folate and closed-related homocysteine pathways. Several of the corresponding genes have been examined as risk factors for NTDs, but few have showed a positive association (Table 1).

Variant	Association with NTDs risk
MTHFR C677T	+
MTHFR A298C	+/-
MTHFR C116T	+ (only in one study)
MTHFR G1793A	+ (only in one study)
MTHFD1 G19658A	+
MTHHD1L 781-6823ATT(7-9)	+
MTR A2756G	-
MTRR A66G	+ (in mothers with low levels of vitamin B12)
CBS 844ins68	+
BHMT G742A	+
RFC-1 A80G	+
FR alpha/Betas	+
TCII A67G; G280A; A701G; C776G; C1043T; G1196A	+
GCPII C1561T	-

Table 1. Variants of genes involved in folate and homocysteine metabolism studied for association with NTDs risk

Methylenetetrahydrofolate reductase (MTHFR) converts 5,10methylenETHF to 5-mthylTHF, the methyl donor for homocysteine conversion to methionine. The 677C→T variant converts an alanine residue to a valine residue. Biochemical studied confirmed that this variant is characterized by residual activity after heating. [28] The variant is extremely common, with homozygosity frequencies ranging between 5% and 15% in many populations. [29] The association of the mutant 677T/T genotype with hyperhomocysteinemia occurs only in individuals with low folate status. Folate has been shown to stabilize the mutant human enzyme and should prevent hyperhomocysteinemia in mutant individuals [30]. The MTHFR 677T/T Homozygous mutant genotype first emerged as a possible risk factor for NTDs in some populations [31-33], even if other studies failed to demonstrate an association. [34-35] Two meta-analysis were performed in 1997 [36] and 2000 [29]. The latter study found a pooled odds ratio for cases 677T/T homozygous cases of 1.7 (95%CI 1.4-2.2), with a pooled attributable fraction of NTDs cases of 6%. A second mutation in the *MTHFR* gene

(1298A→C) has also been positively associated with NTDs [37-39]. The 1298 variant is in strong linkage disequilibrium with the 677 variant, such that 1298C and 677T changes are rarely seen on the same allele. Therefore, the two mutations can occur on separate alleles in the same individual who may be predisposed to hyperhomocysteinemia, if folate status is low. Compound heterozygosity for the 677T and 1298C alleles (677C/T/1298A/C) has been suggested to increase the risk, although the results have been conflicting. [37, 40-41] Moreover, a very recent report identified other two polymorphisms of *MTHFR* gene a 116C→T (P39P) and 1793G→A (R594Q). A possible association between NTDs cases and these two new polymorphisms was found; further analysis demonstrated that this association is driven by the linkage disequilibrium with the 677C→T NTD risk factor [42]. The *MTHFR* 677 variant has been suggested to account for approximately 10% of the NTDs risk. [29]. Since up to 70% of NTD cases may be prevented by the folate, other variants in folate-related genes have to be involved.

Given its important role in folate metabolism, the trifunctional enzyme methylenetetrahydrofolate dehydrogenase/methenyltetrahydrofolate cyclohydrolase/formyltetrahydrofolate synthetase (*MTHFD1*) may play a role in NTD pathogenesis. Brody et al. demonstrated that the polymorphism 1958G→A (R653Q) within the *MTHFD1* gene resulting in the substitution of a conserved arginine by glutamine is a maternal risk factor for NTDs. [43] Mothers who possess two copies of the Q allele have an increased risk of an NTD-affected pregnancy. There was also a suggestion that the Q allele decreases fetal viability. Very recently, the impact of the *MTHFD1* 1958G>A polymorphism on NTD risk was evaluated in the Italian population both by case-control and family-based studies. An increased risk was found for the heterozygous 1958G/A (OR=1.69; P=0.04) and homozygous 1958A/A (OR=1.91; P=0.02) genotypes in the children. The risk of an NTD-affected pregnancy of the mothers was increased 1.67-fold (P=0.04) only when a dominant effect (1958G/A or 1958A/A vs 1958G/G) of the 1958A allele was analysed. Family-based tests also confirmed a significant excess of transmission of the 1958A allele to affected individuals, demonstrating that this variant is a genetic risk factor for Italian NTD cases. [44] A potential role of the mitochondrial paralogue *MTHFD1L* as candidate gene for NTDs association has been also investigated. In particular, a common triallelic deletion/insertion variant 781-6823ATT(7-9) which influences splicing efficiency was tested and a significant increased risk for allele 1 [ATT(7)] was identified. [45]

The methionine synthase (MTR), a vitamin B₁₂-dependent enzyme, utilizes 5-methyltetrahydrofolate, generated by *MTHFR*, for homocysteine remethylation to methionine. A polymorphism of the MTR gene, the 2756A→G (D919G) that lead to the substitution of an aspartic acid with a glycine residue, has been studied in several studies, but the majority have not reported a significant association [36, 40, 46-48].

A common variant in the gene methionine synthase reductases (*MTRR*), the 66A→G (I22M), that converts a isoleucine into a methionine residue, has been shown to confer a four-fold increased risk for NTDs in mothers having low vitamin B₁₂ levels [49]. Other studies have also suggested an effect on NTDs risk, both for mothers and children with 66GG mutant genotype, but interactions with nutrients has not been examined. [50-51]. The *MTRR* mutant genotype combined with *MTHFR* mutant genotype conferred a significant four-fold increased risk in children [49]. An interaction of MTR and *MTRR* genotypes has also recently reported. [50]

Cystathionine beta synthase (CBS) catalyzes the first step in the transsulfuration pathway of homocysteine. The 68-bp insertion in the exon 8 of the gene appeared to be associated with an increased risk in several studies. [46, 52-53]. Multiplicative interaction of the CBS 68-bp insertion with the homozygosity for the MTHFR 677C→T variant was showed to confer a five-fold increased risk for NTDs children, The MTHFR mutation alone had a two-fold increased risk, whereas there was no risk for CBS mutant genotype alone. [54].

Betaine homocysteine methyltransferase (BHMT) catalyzes homocysteine remethylation to methione, using betaine as the methyl donor. Homozygosity for the BHMT R239Q variant is associated with a decreased risk in mother and children [55]. The 80GG genotype for the 80A→G variant of the reduced folate carrier (RFC1), which transports 5-methyltetrahydrofolate across the plasma membrane, confers an increased risk for mothers with low red blood cell folate [55] and for children whose mothers do not use vitamins [56]. A recent study by Rothenberg et al. showed that some mothers with a NTDs-pregnancy produce autoantibodies that bind to folate receptors (FRs) on the placental membrane [57] The authors suggest that folate supplementation would bypass autoantibody formation that mediates the placental FRs blockage.

Several variants have been identified in transcobalamin II (*TCII*) gene (67A→G, 280G→A, 701A→G, 776C→G, 1043C→T, 1196G→A), but none of the SNPs were associated with a significant increased risk for NTD. Among them, three *TCII* SNPs (67A→G, 776C→G, 1196G→A) affected the transcobalamin concentration, even if these variants only partially accounted for the reduced proportion of vitamin B₁₂. [58].

The glutamate carboxypeptidase II (GCPII) is localized in the jejunum and catalyzes the conversion of folate polyglutamates to monoglutamates prior to absorption. The GCPII C1561T (H475Y) variant was identified and studied as risk factor for NTDs, but no study demonstrated an association. [59-60]

In general, there are many different polymorphism in genes invoved in folate transport or metabolism that have been identified and studied for association with NTDs risk, but data are not yet convincing for most of them. The only one well characterized genetic risk factor is the MTHFR C677T. Many studies are biased by small sample size and insufficient statistical power. The interaction between genes and non genetic factors is an intriguing area of investigation, but it requires larger cohort of patients and controls (Table 2).

MS A2756G/MTHFR C677T	(Morrison et al., 1998)
MS A2756G/MTRR A66G	(Zhu et al., 2003)
MTRR A66G/MTHFR C677T	(Relton et al., 2004)
MTRR A66G/FGCP C1561T	(Relton et al., 2004)
MTRR A66G/MTHFD1 G1958A	(van der Put et al., 1997; Hol et al., 1998)
MTHFR A1298C/RFC-1 A80G	(DeMarco et al., 2003)
RFC-1 A80G/CBS 844ins68	(Relton et al., 2004)
MTFR C677T/CBS 31 bp VNTR	(Afman et al., 2003)
MTHFR C677T/CBS 844ins68	(Botto and Mastroiacovo 1998; de Franchis et al. 2002;Relton et al., 2004)

Table 2. Gene-gene interactions between variants of genes involved in folate/homocysteine metabolism

5. From animal models to humans

Mouse mutant strains that exhibit NTDs have arisen by spontaneous mutations, or are the result of mutagen- and gene-trapped induced mutations. Other mutants have arisen through manipulation of an already identified and characterized gene that is manipulated to alter its function. The majority of the known mouse models are those generated by gene knock-out technology, that mainly develop cranial defects. [17, 61-62] The penetrance of NTDs in mouse models is frequently affected by both genetic background and environmental influences. Mouse mutant genes produce NTDs, for the most part, when they are homozygous. Mouse with double heterozygosity for two mutant genes also develop NTDs, demonstrating that gene-gene interaction may promote the development of NTDs. [63] A variety of exogenous agents when administered to embryos exert teratogenic effects with production of abnormalities that include NTDs. Drugs, chemicals, or other substances extraneous to pregnancy cause NTDs. In addition, molecules derived from maternal metabolism, such as those produced in maternal diseases condition, may induce NTDs. For example, elevated glucose and ketone body concentrations, as found in diabetes mellitus, increased the NTD risk. [64] Teratogens interact directly with genetic mutant effects in the causation of NTDs. The frequency of exencephaly in the *curly tail* strain is elevated by administration of all-trans retinoic acid. [65] This demonstrates that exogenous agents can exacerbating the severity of the phenotype interacting with a genetic mutation. An exogenous agent that ameliorates the genetic predisposition to NTDs in mouse models is a candidate therapeutic agent. Several NTDs mouse mutant, *Sp*, *Cart1*, *Cd*, *Splotch*, have shown to interact with folic acid, exhibiting a reduction of NTDs incidence. [66-68] But not all mouse models are folate responsive, demonstrating that there is a proportion of NTDs defects that require another preventive therapy. Inositol was shown preventive in folate-resistant mouse models, such as *curly tail* strain, an action that operates via stimulation of protein kinase C activity. [69]

Many of the genes that are mutated or disrupted in NTD mouse models are now being identified. Few genes have been identified as contributing to human NTD aetiology. Recently, animal models demonstrated an essential role of for the planar cell polarity (PCP) signaling pathway in the process of the convergent extension. Systematic analysis of human homologues PCP genes is giving promising results for our understanding of molecular basis of human NTDs.

6. Genes of PCP pathway and NTDs

Several studies showed that early, during embryogenesis of CNS, the major driving force essential to the shaping of the neural plate is a process which is referred to as convergent extension (CE). This is a morphogenetic process by which cells elongate medio-laterally and intercalate with other neighboring cells forming a longer and narrower array. This changes lead to the conversion of an initially wide and short neural plate into a narrow and elongated one. [70] A crucial role in CE has been assigned to the Planar Cell Polarity (PCP) pathway, a highly conserved, non-canonical Wnt-frizzled-dishevelled signaling cascade (Figure 2). PCP signaling plays a key role in establishing and maintaining polarity in the plane of epithelia in *Drosophila* and in epithelial and non-epithelial tissues in vertebrates.

[71-72] Genetic studies of mutants affecting complex structures in the fly have identified a group of proteins referred to as “core PCP” components, that include transmembrane proteins such as Frizzled (Fz), Strabismus/Van Gogh (Stbm/Vang) and Flamingo (Fmi), as well as cytoplasmic proteins, including Dishevelled (Dsh/Dvl), Prickle (Pk), and Diego (Dgo). [73-74] Downstream of the core PCP members, additional factors mediate the PCP signaling in different tissues, the so-called “PCP effectors”, that include the proteins Inturned, Fuzzy and Fritz. [75-77] Evidence for the involvement of the PCP pathway in CE process in vertebrate has emerged from studies of a wide range of mutants of orthologs of *Drosophila* PCP genes in several animal models such as zebrafish, *Xenopus* and mouse. [78-83] In mouse, *Loop-tail* (*Lp*) was the first mutant to implicate a role of PCP pathway and CE process in NTD pathogenesis. [84-85] *Lp* heterozygotes are characterized by a “looped” tail appearance, while homozygotes develop a severe NTD resembling human craniorachischisis. [86] NTD in *Lp* is caused by independent missense mutations S464N (*Lp*) and D255E (*Lpm1Jus*), localized in the proposed C-terminal cytoplasmic domain of a gene, now referred to as Van Gogh-like 2 (Vangl2). [84-85; 87] A novel experimentally induced allele, *Lp* (*m2Jus*), having a missense mutation, R259L, in Vangl2 has been recently reported. This mutation segregates in a recessive manner, with all heterozygotes appearing normal, and 47% of homozygotes showing a looped-tail. Homozygous *Lp* (*m2Jus*) embryos showed spina bifida in 12%. [88] Seven other mutant mice carrying mutations in some of PCP genes and fail to complete neural tube closure leading to craniorachischisis have been described; they include: *circle-tail* (*Crc*), *crash* (*Crsh*), *Ptk7*, and *dishevelled1*^{-/-}/*dishevelled2*^{-/-} (*Dvl1*^{-/-}/*Dvl2*^{-/-}), *Frizzled3*^{-/-}/*Frizzled6*^{-/-} (*Fzd3*^{-/-}/*Fzd6*^{-/-}), *dishevelled3/Lp* (*Dvl3*^{+/-}/*Lp*⁺) double-knockout mice, and *Fuzzy*. [89-94] The *Crsh* mouse harbors a mutation in *Celsr1* that encodes a protein orthologous to *Drosophila* Flamingo. [93] Like Vangl2, this gene functions in the PCP pathway. In the *Crc* mouse, a point mutation was identified introducing a stop codon into the apical cell polarity gene scribble (*Scrb1*), a PDZ domain-containing gene that is the ortholog of *Drosophila* scribble. [95] Scribble was not known, at a first time, to be a PCP component in *Drosophila*. However, a polarity defect is observed in the inner ear of the *Crc* mice, suggesting a role for *Scrb1* in establishment of polarity in vertebrates. [96] A mutation in the protein tyrosine kinase 7 (*Ptk7*) gene, which encodes a conserved transmembrane protein with tyrosine kinase homology, disrupts neural tube closure and stereociliary bundle orientation and shows genetic interaction with *Lp*. [93] Double null homozygous embryos for both *dishevelled1* and *dishevelled2* genes (*Dvl1*^{-/-}/*Dvl2*^{-/-}) as well as *Frizzled3* and *Frizzled6* genes (*Fzd3*^{-/-}/*Fzd6*^{-/-}), two members of highly homologous seven-pass receptor family, also exhibit NTDs that closely resemble the craniorachischisis observed in single knock-out mice. [91, 97] Although neurulation appeared normal in both *Dvl3*^{-/-} and *LtapLp*⁺/*Lp*⁺ mutants, combined mutants *Dvl3*^{+/-}/*LtapLp*⁺ displayed incomplete neural tube closure. [94] *Fuzzy* knockout mice exhibit both NTDs and defective primary cilia. [98] The exact mechanism by which the PCP pathway regulates CE cellular movements remains poorly understood. A recent study reported that PCP signalling was requested for re-establishing cell polarity that is transiently lost in dividing cells during neurulation. In fact, during mitosis, dividing cells lose their polarized features threatening the overall organization of the developing tissue; PCP signaling acts through quickly repolarization of the daughter cells and directs their integration into the neuroepithelium. [99] Evidence for an intrinsic role of ciliogenesis in PCP is derived from study of Bardet-Biedl syndrome

(BBS), a genetically heterogeneous human disorder with pleiotropic manifestations including obesity, polydactily, endocrine dysfunction, cystic renal disease, progressive photoreceptor degeneration and hearing loss. [100-101] BBS genes share the common features that the encoded proteins localized to the cilium or its cellular anchor, the basal body. [102] Targeted disruption of *Bbs1*, *Bbs4* or *Bbs6* in mouse lead to phenotype shared with PCP mutants, including NTDs (14% of *Bbs4*^{-/-} mice display exencephaly) [103]. A second link between PCP and cilia has come from the identification of the mouse *Inversin* (*Invs*) gene, which encodes a large adaptor-like protein with homology to *Drosophila* PCP protein Diego. Recent studies in zebrafish and *Xenopus* embryos revealed that knock-down of *Invs* disrupts CE process. [104] The subcellular localization of inversin is complex and dynamic, and includes the basal bodies, primary cilia and, during metaphase and anaphase, the spindle poles. [105-107] A recent study in *Xenopus* demonstrated a link between ciliogenesis, Hedgehog signalling (HH) and PCP [108]. Disruption of orthologs of two *Drosophila* PCP effectors, *Inturned* (*In*) and *Fuzzy* (*Fy*), leads to failure of ciliogenesis, resulting from incorrect orientation of ciliary microtubules. In *Xenopus*, the absence of *In* and *Fy* elicited prominent rostral NTDs in addition to more caudal NTDs, predicted to result from disruption of PCP signalling. These caudal defects were shown to arise from failure of CE in the neural plate, whereas the rostral defects were shown to stem from a failure of Hedgehog signalling. [108]. Finally, it is tempting to speculate that both PCP and Hedgehog phenotypes may be linked in humans to NTDs. Mutations in two genes, *MKS1* and *MKS3*, that were predicted to be involved in cilia formation have been recently found in patients with Meckel syndrome (MKS), characterized by bilateral renal cystic dysplasia, central nervous system malformations, bilateral upper and lower limb polydactyly and fibrocystic changes of the liver. [109-112]. The gene mutated in type 1 MKS encodes a protein associated with the base of the cilium in vertebrates and nematodes. Loss of function of mouse *Mks1* results in an accurate model of human MKS, with structural abnormalities in the neural tube, biliary duct, limb patterning, bone development and the kidney that mirror the human syndrome. Analysis of patterning in the neural tube and the limb demonstrates altered Hedgehog pathway signaling underlies some MKS defects.[113]

7. PCP pathway and human NTDs

In humans mutations in *VANGL1* and *VANGL2* genes cause NTDs and abrogate the physical interaction between VANGL proteins and other members of PCP pathway. In a first study three Italian patients who were heterozygotes for missense mutations of *VANGL1* (V239I, R274Q, and M328T) were identified. The V239I mutation was identified in a girl with a severe form of caudal regression The girl's mother also carried the mutation, but showed no clinical signs of NTDs. Her brother having a milder form of the disorder, a dermal sinus, also carried the V239I mutation. The V239I mutation was absent in the father, in the maternal aunt and in the maternal grandparents, indicating that it had arisen *de novo* in the germline of one of the maternal grandparents or somatically in the mother, with subsequent transmission through the mother's germline. Valine at position 239 is located in the fourth predicted transmembrane domain of the *VANGL1* protein. It is invariant and part of a 'VLLE' motif, which is conserved across all known VANGL proteins. In vitro studies demonstrated that V239I mutation abrogated interaction between *VANGL1* and their binding partners, *Dvl1*, *Dvl2*, and *Dvl3*. The R274Q mutation was found in a girl with

myelomeningocele. Her mother, who carried the R274Q mutation, and maternal aunt had vertebral schisis, a minimal sign of NTD. Arg274 is in the cytoplasmic domain of VANGL1 and is invariant in all known orthologs except in *C. elegans*, in which it is replaced by glutamate. The M328T mutation was carried by a child with myelomeningocele, hydrocephalus, and Chiari II malformation. M328T mutation occurs in the predicted cytoplasmic domain of VANGL1. We did not detect these variants in 150 ancestrally matched controls. Thus, these are the first genetic mutations clearly linked to NTDs such as spina bifida in humans. [114].

Fig. 2. Planar Cell Polarity (PCP) pathway.

Wnt proteins bind to its receptors, Frizzled (Fz), leading to recruitment and activation of Dishevelled, that is a multimodule protein PCP activation requires formation of a multimodule protein-complex including two transmembrane proteins, Vangl (1/2) and Flamingo/Celsr, and three cytoplasmic proteins, Dishevelled, Diego and Prickle. These proteins acquire an asymmetric localization at the plasma membrane that is crucial for proper PCP signaling. Scribble is a cytoplasmic protein binds to and genetically interacts with Vangl, suggesting a role in PCP signaling. Ptk7 receptor is a regulator and shows genetic interaction with Vangl. Downstream effectors include small GTPases Rho or Rac, which are active when bound to GTP. The Formin homology protein, Daam1, as an important link between Dishevelled and the Rho GTPase for cytoskeletal modulation. Alternatively, Dvl can initiate Rac signaling and its downstream effector c-Jun N-terminal kinase (JNK) that promote sites of actin polymerization modulating lamellipodia extension. Inturned (In) and Fuzzy that are down-stream regulator of PCP pathway do influence convergent extension and they also are broadly required for Hedgehog (HH) signaling

Validation of the potential pathogenic effect of VANGL1 V239I, M328T, and R274Q mutations *in vivo* was performed by investigating their effect on CE in zebrafish. Knocking down the expression of *tri*, the ortholog of *Vangl2*, using an antisense morpholino (MO), led to a defective CE manifested by a shortened body axis and widened somites. Co-injection of the human VANGL1 with the *tri*-MO was able to partially rescue the *tri*-MO induced phenotype in zebrafish. In contrast, co-injection of V239I and M328T, failed to rescue this phenotype. Overexpression studies evaluating the ability of the human VANGL1 alleles to induce a CE phenotype when injected at high doses in zebrafish embryos have been carried out. While overexpressing the wild-type allele led to a severely defective CE, overexpression of either V239I and M328T variants failed to do so. Thus, results from both *tri*-MO knockdown/rescue results and overexpression assays suggest that these two variants most likely represent "loss-of-function" alleles that affect protein function during embryonic development. [115].

Overall, human VANGL1 gene has been sequenced in a cohort of 810 NTD patients with various ethnic origin. Eight missense mutations both in familial (V239I, R274Q, S83L, and R181E) and sporadic (M238T, F153S, L202F, and A404S) cases have been identified. These mutations affect evolutionary conserved amino residues that are distributed along the entire length of the VANGL1 protein. Since these many of these variants do not represent obvious null mutations (like stop codon, deletions), functional experiments are needed to investigate their effect on the protein function. All mutations detected so far in VANGL1 in NTDs are heterozygous, leading to the speculation that these variants may act as partial loss of function alleles and interact with other environmental and genetic factors to cause the NTD phenotype. The finding of VANGL1 mutations in both open and closed NTDs support this hypothesis of common underlying molecular mechanisms. [116]

Sequencing of *VANGL2* in the same cohort of patients led to the identification of six novel heterozygous missense mutations in seven patients, that could be pathogenic based on genetic and initial validation data [117]. Four of these mutations, R135W, R177H, L242V, R270H, were predicted to be damaging to protein function using bioinformatics' tools, and two others, T247M and R482H, affect highly conserved residues across evolution. Five mutations were identified in patients affected with closed spinal NTDs, suggesting that *VANGL2* mutations may predispose to NTDs is approximately 2.5% of closed spinal NTDs. A Chinese study recently reported the identification of three other missense mutations in *VANGL2* in fetuses with a cranial NTDs: S84F, R353C, and F437S. [118]

Recently, three non-synonymous Fuzzy amino acid substitutions in some patients with NTDs have been identified, resulting in alteration of the length of primary cilia and cell movement. Since Fuzzy knockout mice exhibit both NTDs and defective primary cilia and Fuzzy is expressed in the emerging neural tube, mutations in Fuzzy may account for a subset of NTDs in humans. [98]

Thus, the evidence is accumulating for an important contribution of PCP genes to the pathogenesis of human NTDs, necessitating a detailed analysis of other not yet explored PCP genes in large cohorts of patients.

8. Conclusions

NTDs are a group of severe congenital malformations having a profound physical, emotional, and financial effects on families and communities. Despite their importance in

terms both of human suffering and cost to the health care system, the causes of most cases of NTDs are not known. There is growing evidence that many NTDs have a genetic background. Elucidation of genetic causes predisposing to NTDs are important challenge in light of preconception care program that aim to reduce reproductive risks before conception and to improve the chance for a healthy birth outcome. Further, the genetic screening done for research setting may become available for diagnostic purposes, allowing to identify the predisposition before conception and to provide reproductive options to minimize the chance of having affected children in the future.

9. References

- [1] EUROCAT Working Group: Surveillance of Congenital anomalies in Europe 1980-1999. 2002. EUROCAT, University of Ulster, Report No.: 8.
- [2] Busby A, Abramsky L, Dolk H, Armstrong B, Addor MC, Anneren G, Armstrong N, Baguette A, Barisic I, Berghold A, Bianca S, Braz P, Calzolari E, Christiansen M, Cocchi G, Daltveit AK, De Walle H, Edwards G, Gatt M, Gener B, Gillerot Y, Gjergja R, Goujard J, Haeusler M, Latos-Bielenska A, McDonnell R, Neville A, Olars B, Portillo I, Ritvanen A, Robert-Gnansia E, Rösch C, Scarano G, Steinbicker V. 2005. Preventing neural tube defects in Europe: a missed opportunity. *Repro Toxic* 20 (3): 393-402.
- [3] Schoenwolf GC, Smith JL. 2000. Mechanisms of neurulation. *Methods Mol Biol* 136 125-134.
- [4] Catala M. Genetic control of caudal development. 2002. *Clin Genet* 61: 89-96.
- [5] Colas J.F. and Schoenwolf. 2001. Towards a cellular and molecular understanding of neurulation. *Dev Dyn* 221, 117-145.
- [6] Copp AJ, Greene ND, Murdoch JN: 2003. The genetic basis of mammalian neurulation. *Nat Rev genet* 4: 784-793.
- [7] van Allen MI, Kalousek DK, Chernoff GF, Juroff D, Harris M, McGillivray BC, Young S-L, Langlois S, MacLeod PM, Chitayat D, et al.. 1993. Evidence for multi-site closure of the neural tube in humans. *Am J Med Genet* 47, 723-743.
- [8] Botto LD, Moore CA, Khoury MJ, Erickson JD. 1999. Neural-tube defects. *New England Journal of Medicine* 341: 1509-1519.
- [9] Rossi A, Cama A, Piatelli G, Ravegnani M, Biancheri R, Tortori-Donati P. 2004. Spinal dysraphism: MR imaging rationale. *J Neuroradiol* 31: 3-24.
- [10] O'Reilly GC, Shields LE. Karyotyping for isolated neural tube defects. 2000. A report of two cases. *J Reprod Med* 45:950-952.
- [11] Elwood JM, Little J, Elwood JH. 1992. *Epidemiology and control of neural tube defects*. Oxford:Oxford University Press.
- [12] Fuhrmann W, Seeger W, Böhm R. 1971. Apparently monogenic inheritance of anencephaly and spina bifida in a kindred. *Hum Genet* 13(3) :241-243.
- [13] Partington MD, McLone DG. 1995. Hereditary factors in the etiology of neural tube defects. Results of a survey. *Pediatr Neurosurg*. 23(6): 311-316.
- [14] Toriello HV, Higgins JV. 1985. Possible causal heterogeneity in spina bifida cystica. *Am J Med Genet*. 21(1): 13-20.

- [15] Garabedian BH, Fraser FC. 1993. Upper and lower neural tube defects: an alternate hypothesis. *J Med Genet* 30(10): 849-51.
- [16] Byrne J, Carolan S. 2006. Adverse reproductive outcomes among pregnancies of aunts and (spouses of) uncles in Irish families with neural tube defects. *Am J Med Genet A* 140(1):52-61.
- [17] Juriloff DM, Harris MJ. 2000. Mouse models for neural tube closure defects. *Hum Mol Genet.* 19(6): 993-1000.
- [18] Udvardi AJ, Linney E. 2003. Windows into development: historic, current, and future perspectives on transgenic zebrafish. *Dev Biol* 256(1):1-17.
- [19] Wienholds E, Schulte-Merker S, Walderich B, Plasterk RH. 2002. Target-selected inactivation of the zebrafish *rag1* gene. *Science* 297:99-102.
- [20] Sebold CD, Melvin EC, Siegel D, Mehlretter L, Enterline DS, Nye JS, Kessler J, Bassuk A, Speer MC, George TM; NTD Collaborative Group. 2005. Recurrence risks for neural tube defects in siblings of patients with lipomyelomeningocele. *Genet Med* 7(1):64-67.
- [21] Rampersaud E, Bassuk AG, Enterline DS, George TM, Siegel DG, Melvin EC, Aben J, Allen J, Aylsworth A, Brei T, Bodurtha J, Buran C, Floyd LE, Hammock P, Iskandar B, Ito J, Kessler JA, Lasarsky N, Mack P, Mackey J, McLone D, Meeropol E, Mehlretter L, Mitchell LE, Oakes WJ, Nye JS, Powell C, Sawin K, Stevenson R, Walker M, West SG, Worley G, Gilbert JR, Speer MC. 2005. Whole genomewide linkage screen for neural tube defects reveals regions of interest on chromosomes 7 and 10. *J Med Genet* 42(12):940-946.
- [22] Spielman RS, McGinnis RE, Ewens WJ. 1993. Transmission test for linkage disequilibrium: the insulin gene region and insulin-dependent diabetes mellitus (IDDM). *Am J Hum Genet* 52(3):506-16.
- [23] MRC Vitamin Study Group. 1991. Prevention of neural tube defects: results of the Medical Research Council Vitamin Study. MRC Vitamin Study research Group. *Lancet* 338, 131-137.
- [24] Czeizel AE, Dudas I. 1992. Prevention of the first occurrence of neural-tube defects by periconceptional vitamin supplementation. *N Engl J Med* 327, 1832-1835.
- [25] Mills JL, Scott JM, Kirke PN, McPartlin JM, Conley MR, Weir DG, Molloy AM, Lee YJ. 1996. Homocysteine and neural tube defects. *J Nutr* 126(3):756S-760S.
- [26] Scott JM. 1999. Folate and vitamin B12 *Proc Nutr Soc* 58(2):441-448.
- [27] Steegers-Theunissen RP, Boers GH, Trijbels FJ, Finkelstein JD, Blom HJ, Thomas CM, Borm GF, Wouters MG, Eskes TK. 1994. Maternal hyperhomocysteinemia: a risk factor for neural-tube defects? *Metabolism* ;43(12):1475-80.
- [28] Frost P, Blom HJ, Milos R et al. 1995. A candidate genetic risk factor for vascular disease: a common mutation in methylenetetrahydrofolate reductase. *Nature Genet* 10: 111-113.
- [29] Botto LD, Yang Q. 2000. 5,10-Methylenetetrahydrofolate reductase gene variants and congenital anomalies: a HuGE review. *Am J Epidemiol.* 151: 862-877.
- [30] Malinow MR, Nieto FJ, Kruger WD, Duell PB, Hess DL, Gluckman RA, Block PC, Holzgang CR, Anderson PH, Seltzer D, Upson B, Lin QR. 1997. The effects of folic acid supplementation on plasma total homocysteine are modulated by

- multivitamin use and methylenetetrahydrofolate reductase genotypes. *Arterioscler Thromb Vasc Biol* 17(6):1157-1162.
- [31] Van der Put NM, Steegers-Theunissen RP, Frosst P, Trijbels FJ, Eskes TK, van den Heuvel LP, Mariman EC, den Heyer M, Rozen R, Blom HJ. 1995. Mutated methylenetetrahydrofolate reductase as risk factor for spina bifida. *Lancet* 346:1070-1071.
- [32] Ou CY, Stevenson RE, Brown VK, Schwart CE, Allen WP, Khoury MJ, Rozen R, Oakley GP, Adams MJ. 1996. Methylenetetrahydrofolate reductase genetic polymorphism as a risk factor for neural tube defects. *Am J Med Genet* 63: 610-614.
- [33] Christensen B, Arbour L, Tran P, Leclerc D, Sabbaghian N, Platt R, Gilfix BM, Rosenblatt DS, Gravel RA, Forbes P, Rozen R. 1999. Genetic polymorphisms in methylenetetrahydrofolate reductase and methionine synthase, folate levels in red blood cells, and risk of neural tube defects. *Am J Med Genet* 84: 151-157.
- [34] Koch MC, Stegmann K, Ziegler A, Schroter B, Ermert A. 1998. Evaluation of the MTHFR C677T allele and the MTHFR gene locus in a German spina bifida population. *Eur J Pediatr* 157: 487-492.
- [35] Gonzales-Herrera L, Garcia-Escalante G, Castillo-Zapata I, Canto-Herrera J, Caballos-Quintal J, Pinto-Escalante D, Diaz-Rubio F, Del Angel RM, Orozco-Orozco L. 2002. Frequency of the thermolabile variant C677T in the MTHFR gene and lack of association with neural tube defects in the State of Yucatan, Mexico. *Clin Genet* 62: 394-398.
- [36] van der Put NM, Eskes TK, Blom HJ. 1997. Is the common 677C-->T mutation in the methylenetetrahydrofolate reductase gene a risk factor for neural tube defects? A meta-analysis. *QJM*. 90(2):111-115.
- [37] van der Put NM, Gabreels F, Stevens EM, Smeitink JA, Trijbels FJ, Eskes TK, van den Heuvel LP, Blom HJ. 1998. A second common mutation in the methylenetetrahydrofolate reductase gene: an additional risk factor for neural-tube defects? *Am J Hum Genet* 62: 1044-1051.
- [38] Weisberg I, Tran P, Christensen B, Sibani S, Rozen R. 1998. A second genetic polymorphism in methylenetetrahydrofolate reductase (MTHFR) associated with decreased enzyme activity. *Mol Genet Metab* 64: 169-172.
- [39] De Marco P, Calevo MG, Moroni A, Arata L, Merello E, Finnell RH, Zhu H, Andreussi L, Cama A, Capra V. 2002. Study of MTHFR and MS polymorphisms as risk factors for NTD in the Italian population. *J Hum Genet* 47: 319-324.
- [40] Trembath D, Sherbondy AL, Vandyke DC, Shaw GM, Todoroff K, Lammer EJ, Finnell RH, Marker S, Lerner G, Murray JC. 1999. Analysis of select folate pathway genes, PAX3, and human T in a Midwestern neural tube defect population. *Teratology* 59(5):331-341.
- [41] Richter B, Stegmann K, Röper B, Böddiker I, Ngo ET, Koch MC. 2001. Interaction of folate and homocysteine pathway genotypes evaluated in susceptibility to neural tube defects (NTD) in a German population. *J Hum Genet* 46(3):105-109.
- [42] O'Leary VB, Mills JL, Parle-McDermott A, Pangilinan F, Molly AM, Cox C, Weiler A, Conley M, Kirke PN, Scott JM, Brody LC. 2005. Screening for new MTHFR polymorphisms and NTD risk. *Am J Med Genet* 138: 99-106.

- [43] Brody LC, Conley M, Cox C, Kirke PN, McKeever MP, Mills JL, Molloy AM, O'Leary VB, Parle-McDermott A, Scott JM, Swanson DA. 2002. A polymorphism, R653Q, in the trifunctional enzyme methylenetetrahydrofolate dehydrogenase/methenyltetrahydrofolate cyclohydrolase/formyltetrahydrofolate synthetase is a maternal genetic risk factor for neural tube defects: report of the Birth Defects Research Group. *Am J Hum Genet* 71:1207-1215.
- [44] De Marco P, Merello E, Calevo MG, Mascelli S, Raso A, Cama A, Capra V. 2006. Evaluation of methylenetetrahydrofolate-dehydrogenase 1958G>A polymorphism for neural tube defect risk. *J Hum Genet* 51: 98-103.
- [45] Pearle-McDermott A, Pangilinan F, O'Brien KK, Mills JL, Magee AM, Troendle J, Sutton M, Scott JM, Kirke PN, Molloy AM, Brody LC. 2009. A common variant in MTHFD1L is associated with neural tube defects and mRNA splicing efficiency. *Hum Mutat* 30(12): 1650-1656.
- [46] Morrison K, Papapetrou C, Hol FA, Mariman EC, Lynch SA, Burn J, Edwards YH. 1998. Susceptibility to spina bifida; an association study of five candidate genes. *Ann Hum Genet* 62: 379-396.
- [47] Brody LC, Baker PJ, Chines PS, Musick A, Molloy AM, Swanson DA, Kirke PN, Ghosh S, Scott JM, Mills JL. 1999. Methionine synthase: high-resolution mapping of the human gene and evaluation as a candidate locus for neural tube defects. *Mol Genet Metab* 1999 67(4):324-333.
- [48] Johanning GL, Tamura T, Johnston KE, Wenstrom KD. Comorbidity of 5,10-methylenetetrahydrofolate reductase and methionine synthase gene polymorphisms and risk for neural tube defects. 2000. *J Med Genet* 37(12):949-951.
- [49] Wilson A, Platt R, Wu Q, Leclerc D, Christensen B, Yang H, Gravel RA, Rozen R. A common variant in methionine synthase reductase combined with low cobalamin (vitamin B12) increases risk for spina bifida. 1999. *Mol Genet Metab* 67(4):317-323.
- [50] Zhu H, Wicker NJ, Shaw GM, Lammer EJ, Hendricks K, Suarez L, Canfield M, Finnell RH. 2003. Homocysteine remethylation enzyme polymorphisms and increased risks for neural tube defects. *Mol Genet Metab* 78(3):216-221.
- [51] Guéant-Rodriguez RM, Rendeli C, Namour B, Venuti L, Romano A, Anello G, Bosco P, Debard R, Gérard P, Viola M, Salvaggio E, Guéant JL. 2003. Transcobalamin and methionine synthase reductase mutated polymorphisms aggravate the risk of neural tube defects in humans. *Neurosci Lett* 344(3):189-192.
- [52] Ramsbottom D, Scott JM, Molloy A, Weir DG, Kirke PN, Mills JL, Gallagher PM, Whitehead AS. 1997. Are common mutations of cystathionine beta-synthase involved in the aetiology of neural tube defects? *Clin Genet* 51(1):39-42.
- [53] Speer MC, Nye J, McLone D, Worley G, Melvin EC, Viles KD, Franklin A, Drake C, Mackey J, George TM. 1999. Possible interaction of genotypes at cystathionine beta-synthase and methylenetetrahydrofolate reductase (MTHFR) in neural tube defects. NTD Collaborative Group. *Clin Genet* 56(2):142-144.
- [54] Botto LD, Mastroiacovo P. 1998. Exploring gene-gene interactions in the etiology of neural tube defects. *Clin Genet* 53(6):456-459.

- [55] Morin I, Platt R, Weisberg I, Sabbaghian N, Wu Q, Garrow TA, Rozen R. 2003. Common variant in betaine-homocysteine methyltransferase (BHMT) and risk for spina bifida. *Am J Med Genet A* 119A(2):172-176.
- [56] Shaw GM, Lammer EJ, Zhu H, Baker MW, Neri E, Finnell RH. 2002. Maternal periconceptional vitamin use, genetic variation of infant reduced folate carrier (A80G), and risk of spina bifida. *Am J Med Genet* 108(1):1-6.
- [57] Rothenberg SP, da Costa MP, Sequeira JM, Cracco J, Roberts JL, Weedon J, Quadros EV. 2004. Autoantibodies against folate receptors in women with a pregnancy complicated by a neural-tube defect. *N Engl J Med*. 350(2):134-142.
- [58] Afman LA, Van Der Put NM, Thomas CM, Trijbels JM, Blom HJ. 2001. Reduced vitamin B12 binding by transcobalamin II increases the risk of neural tube defects. *QJM* 94(3):159-166.
- [59] Vieira AR, Trembath D, Vandyke DC, Murray JC, Marker S, Lerner G, Bonner E, Speer M. 2002. Studies with His475Tyr glutamate carboxipeptidase II polymorphism and neural tube defects. *Am J Med Genet* 111(2):218-219.
- [60] Afman LA, Trijbels FJ, Blom HJ. 2003. The H475Y polymorphism in the glutamate carboxypeptidase II gene increases plasma folate without affecting the risk for neural tube defects in humans. *J Nutr* 133(1):75-77.
- [61] Juriloff DM, Harris MJ. 2000. Mouse models for neural tube closure defects. *Hum Mol Genet* 9(6):993-1000.
- [62] Harris MJ, Juriloff DM. 2007. Mouse mutants with neural tube closure defects and their role in understanding human neural tube defects. *Birth Defects Res A Clin Mol Teratol* 79(3):187-210.
- [63] Estibeiro JP, Brook FA, Copp AJ. 1993. Interaction between splotch (Sp) and curly tail (ct) mouse mutants in the embryonic development of neural tube defects. *Development* 119(1):113-121.
- [64] Sadler TW, Hunter ES 3rd, Wynn RE, Phillips LS. 1989. Evidence for multifactorial origin of diabetes-induced embryopathies. *Diabetes* 38(1):70-74.
- [65] Tom C, Juriloff DM, Harris MJ. 1991. Studies of the effect of retinoic acid on anterior neural tube closure in mice genetically liable to exencephaly. *Teratology* 43(1):27-40.
- [66] Zhao Q, Behringer RR, de Crombrughe B. 1996. Prenatal folic acid treatment suppresses acrania and meroanencephaly in mice mutant for the *Cart1* homeobox gene. *Nat Genet* 13(3):275-283.
- [67] Fleming A, Copp AJ. 1998. Embryonic folate metabolism and mouse neural tube defects. *Science* 280(5372):2107-2109.
- [68] Gefrides LA, Bennett GD, Finnell RH. 2002. Effects of folate supplementation on the risk of spontaneous and induced neural tube defects in Splotch mice. *Teratology* 65(2):63-69.
- [69] Greene ND, Copp AJ. 1997. Inositol prevents folate-resistant neural tube defects in the mouse. *Nat Med* 3(1):60-66.
- [70] Wallingford JB, Harland RM. 2002. Neural tube closure requires Dishevelled-dependent convergent extension of the midline. *Development* 129, 5815-5825.

- [71] Strutt D. 2003. Frizzled signalling and cell polarisation in *Drosophila* and vertebrates. *Development* 130:4501-4513.
- [72] Klein TJ, Mlodzik M. 2005. Planar cell polarization: an emerging model points in the right direction. *Annu Cell Dev Biol* 21: 155-176.
- [73] Jenny A, Mlodzik M. 2006. Planar cell polarity signaling: a common mechanism for cellular polarization. *Mt Sinai J Med* 73:738-750.
- [74] Jones C, Chen P. 2007. Planar cell polarity signaling in vertebrates. *Bioessays* 29 (2):120-32.
- [75] Adler PN. 2002. Planar signaling and morphogenesis in *Drosophila*. *Dev Cell* 2:525-535.
- [76] Adler PN, Zhu CM, Stone D. 2004. Inturned localizes to the proximal side of wing cells under the instruction of upstream planar polarity proteins. *Current Biology* 14: 2046-2051.
- [77] Collier S, Lee H, Burgess R, Adler P. 2005. 40 repeat protein fritz links cytoskeletal planar polarity to frizzled subcellular localization in the *drosophila* epidermis. *Genetics* 169: 2035-2045.
- [78] Wallingford JB. 2005. Neural tube closure and neural tube defects: studies in animal models reveal known knowns and known unknowns. *Am J Med Genet C Semin Med Genet* 135:59-68.
- [79] Barrow JR. 2006. Wnt/PCP signaling: a veritable polar star in establishing patterns of polarity in embryonic tissues. *Semin cell Dev Biol* 17 (2): 185-93.
- [80] Montcouquiol M, Crenshaw EB III, Kelley MW. 2006. Noncanonical Wnt signaling and neural polarity. *Annu Rev Neurosci* 29:363-386.
- [81] Wallingford JB, Rowning BA, Vogeli KM. 2000. Dishevelled controls cell polarity during *Xenopus* gastrulation. *Nature* 405:81-85.
- [82] Tada M, Smith JC. 2000. Xwnt11 is a target of *Xenopus* Brachyury: regulation of gastrulation movements via Dishevelled, but not through the canonical Wnt pathway. *Development* 127: 2227-2238
- [83] Heisenberg CP, Tada M, Rauch GJ, Saude L, Concha ML, Geisler R, Stemple DL, Smith JC, and Wilson SW. 2000. Silberblick/Wnt11 mediates convergent extension movements during zebrafish gastrulation. *Nature* 405: 76-81
- [84] Kibar Z, Vogan KJ, Groulx N, Groulx N, Justice MJ, Underhill DA, Gros P. 2001. Ltap, a mammalian homolog of *Drosophila* Strabismus/Van Gogh, is altered in the mouse neural tube mutant Loop-tail. *Nat Genet* 28:251-255.
- [85] Murdoch JN, Doudney K, Paternotte C, Copp AJ, Stenier P. 2001. Severe neural tube defects in the loop-tail mouse result from mutation of Lpp1, a novel gene involved in floor plate specification. *Hum Mol Genet* 10: 2593-2601.
- [86] Stein K, Rudin IA. 1953. Development of mice homozygous for the gene for looped-tail. *J hered* 44: 59-69.
- [87] Kibar Z, Underhill DA, Canonne-Hergaux F, Gauthier S, Justice MJ, Gros P. 2001. Identification of a new chemically induced allele (Lp(m1Jus)) at the loop-tail locus: morphology, histology, and genetic mapping. *Genomics* 72:331-337.

- [88] Guyot MC, Bosoi CM, Kharfallah F, Reynolds A, Drapeau P, Justice M, Gros P, Kibar Z. 2011. A novel hypomorphic Looptail allele at the planar cell polarity Vangl2 gene. *Dev Dyn*. 2011 240(4):839-849
- [89] Smith LJ and Stein JK. 1962. Axial elongation in the mouse and its retardation in homozygous loop tail mice. *J mbryol Exp Morphol* 10: 73-87.
- [90] Murdoch JN, Rachel RA, Shah S, Beermann F, Stanier P, Mason Ca, Copp AJ. 2001. Circletail, a new mouse mutant with severe neural tube defects: chromosomal localization and interaction with the loop-tail mutation. *Genomics* 78:55-63.
- [91] Hamblet NS, Lijam N, Ruiz-Lozano P. 2002. Dishevelled 2 is essential for cardiac outflow tract development, somite segmentation and neural tube closure. *Development* 129:5827-5838
- [92] Curtin JA, Quint E, Tsipouri V, Arkell RM, Cattanach B, Copp AJ, Henderson DJ, Spurr N, Stanier P, Fisher EM, Nolan PM, Steel KP, Brown SD, Gray IC, Murdoch JN. 2003. Mutation of Celsr1 disrupts planar polarity of inner ear hair cells and causes severe neural tube defects in the mouse. *Curr Biol* 13:1129-1133.
- [93] Lu X, Borchers AG, Jolicœur C, Rayburn H, Baker JC, Tessier-Lavigne M. 2004. PTK7/CCK-4 is a novel regulator of planar cell polarity in vertebrates. *Nature* 430: 93-98.
- [94] Etheridge SL, Ray S, Li S, Hamblet NS, Lijam N, Tsang M, Greer J, Kardos N, Wang J, Sussman DJ, Chen P, Wynshaw-Boris A. 2008. Murine dishevelled 3 functions in redundant pathways with dishevelled 1 and 2 in normal cardiac outflow tract, cochlea, and neural tube development. *PLoS Genet* 4(11):e1000259.
- [95] Murdoch JN, Henderson DJ, Doudney K, Gaston-Massuet C, Philipps HM, Paternotte C, Arkell R, Stanier P, Copp AJ. 2003. Disruption of scribble (Scrb1) causes severe neural tube defects in the circletail mouse. *Hum Mol Genet* 12:87-98.
- [96] Montcouquiol M, Rachel RA, Lanford PJ, Copeland NG, Jenkins NA, Kelley MW. 2003. Identification of Vangl2 and Scrb1 as planar polarity genes in mammals. *Nature* 423:173-177.
- [97] Wang Y, Guo N, Nathans J. 2006. The role of Frizzled3 and Frizzled6 in neural tube closure and in the planar polarity of inner-ear sensory hair cells. *J Neurosci* 26:2147-2156.
- [98] Seo JH, Zilber Y, Babayeva S, Liu J, Kyriakopoulos P, De Marco P, Merello E, Capra V, Gros P, Torban E. 2011. Mutations in the planar cell polarity gene, Fuzzy, are associated with neural tube defects in humans. *Hum Mol Genet*. 2011 Aug 25.
- [99] Ciruna B, Jenny A, Lee D, Mlodzik M, Schier AF. 2006. Planar cell polarity signalling couples cell division and morphogenesis during neurulation. *Nature* 439:220-224.
- [100] Bisgrove BW, Yost HJ. 2006. The roles of cilia in developmental disorders and disease. *Development* 133: 4131-4143.
- [101] Davis EE, Brueckner M, Katsanis N. 2006. The emerging complexity of the vertebrate cilium: New functional roles for an ancient organelle. *Developmental Cell* 11: 9-19.
- [102] Ansley SJ, Badano JL, Blacque OE, Hill J, Hoskins BE, Leitch CC, Kim JC, Ross AJ, Eichers ER, Teslovich TM, Mah AK, Johnsen RC, Cavender JC, Lewis RA, Leroux MR, Beales PL, and Katsanis N. 2003. Basal body dysfunction is a likely cause of pleiotropic Bardet-Biedl syndrome. *Nature* 425: 628-633.

- [103] Ross AJ, May-Simera H, Eichers ER, Kai M, Hill J, Jagger D J, Leitch CC, Chapple JP, Munro PM, Fisher S, Tan PL, Phillips HM, Leroux MR, Henderson DJ, Murdoch JN, Copp AJ, Eliot MM, Lupski JR, Kemp DT, Dollfus H, Tada M, Katsanis N, forge A, Bales PL. 2005. Disruption of Bardet-Biedl syndrome ciliary proteins perturbs planar cell polarity in vertebrates. *Nature Genet* 37: 1135-1140.
- [104] Simons M, Gloy J, Ganner A, Bullerkotte A, Bashkurov M, Kronig C, Schermer B, Benzing T, Cabello OA, Jenny A, Mlodzik M, Polok B, Driever W, Obara T, Walz G. 2005. Inversin, the gene product mutated in nephronophthisis type II, functions as a molecular switch between Wnt signaling pathways. *Nat Genet* 37:537-543
- [105] Morgan D, Goodship J, Essner JJ, Vogan KJ, Turnpenny L, Yost HJ, Tabin CJ, Strachan T. 2002. The left-right determinant inversin has highly conserved ankyrin repeat and IQ domains and interacts with calmodulin. *Human Genetics* 110: 377-384.
- [106] Watanabe D, Saijoh Y, Nonaka S, Sasaki G, Ikawa Y, Yokoyama T, Hamada H. 2003. The left-right determinant Inversin is a component of node monocilia and other 9+0 cilia. *Development* 130: 1725-1734.
- [107] Eley L, Turnpenny L, Yates LM, Craighead AS, Morgan D, Whistler C, Goodship JA, Strachan T. 2004. A perspective on inversin. *Cell Biology International* 28: 119-124
- [108] Park TJ, Haigo SL, Wallingford JB. 2006. Ciliogenesis defects in embryos lacking inturned or fuzzy function are associated with failure of planar cell polarity and Hedgehog signaling. *Nat Genet* 38 (3): 303-11.
- [109] Ahdab-Barmada, M.; Claassen, D. 1990. A distinctive triad of malformations of the central nervous system in the Meckel-Gruber syndrome. *J Neuropath Exp Neurol* 49: 610-620.
- [110] Salonen R, Paavola P. 1998. Meckel syndrome. *Journal of Medical Genetics* 35: 497-501.
- [111] Smith U, Pasha S, Cox P, Attie-Bitach T, Maher ER, Johnson CA. 2005. Fine-mapping the MKS3 locus for Meckel-Gruber syndrome. *Journal of Medical Genetics* 42: S107
- [112] Kyttälä M, Tallila J, Salonen R, Kopra O, Kohlschmidt N, Pavola-Sakki P, Peltonen L, Kestila M. 2006. MKS1, encoding a component of the flagellar apparatus basal body proteome, is mutated in Meckel syndrome. *Nat Genet* 38:155-157.
- [113] Weatherbee SD, Niswander LA, Anderson KV. 2009. A mouse model for Meckel syndrome reveals Mks1 is required for ciliogenesis and Hedgehog signaling. *Hum Mol Genet* 18(23):4565-4575.
- [114] Kibar Z, Torban E, Mc Dearnid JR, Reynolds A, Mathieu M, kirillova I, De Marco P, Merello E, Hayes JM, Wallingford JB, Drapeau P, Capra V, Gros P. 2007. Mutations in VANGL1 associated in neural-tube defects. *N Engl J Med* 361(14): 1432-1437.
- [115] Reynolds A, McDearmid JR, Lachance S, De Marco P, Merello E, Capra V, Gros P, Drapeau P, Kibar Z. 2010. VANGL1 rare variants associated with neural tube defects affect convergent extension in zebrafish. *Mech Dev* 127:385-392.
- [116] Kibar Z, Bosoi CM, Kooistra M, Salem S, Finnell RH, De Marco P, Merello E, Bassuk AG, Capra V, Gros P. 2009. Novel Mutations in VANGL1 in Neural Tube Defects. *Hum Mutat* 30:E706-E715.

- [117] Kibar Z, Salem S, Bosoi CM, Pauwels E, De Marco P, Merello E, Bassuk AG, Capra V, Gros P. 2010. Contribution of VANGL2 mutations to isolated neural tube defects. *Clin Genet* 2011 80(1):76-82.
- [118] Lei YP, Zhang T, Li H, Wu BL, Jin L, Wang HY. 2010. VANGL2 mutations in human cranial neural-tube defects *N Engl J Med* 362:2232-2235.

IntechOpen

IntechOpen

Neural Tube Defects - Role of Folate, Prevention Strategies and Genetics

Edited by Dr. Kannan Laksmi Narasimhan

ISBN 978-953-51-0317-2

Hard cover, 200 pages

Publisher InTech

Published online 16, March, 2012

Published in print edition March, 2012

The book Neural Tube Defects - Role of Folate, Prevention Strategies and Genetics has several eminent international authors and the book is a resource for anybody who is interested in this very important subject. The authors are distinguished and the chapters are a product of their extensive research.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Patrizia De Marco (2012). Advances in Genetics of Non Syndromic Neural Tube Defects, Neural Tube Defects - Role of Folate, Prevention Strategies and Genetics, Dr. Kannan Laksmi Narasimhan (Ed.), ISBN: 978-953-51-0317-2, InTech, Available from: <http://www.intechopen.com/books/neural-tube-defects-role-of-folate-prevention-strategies-and-genetics/advances-in-genetics-of-non-syndromic-neural-tube-defects>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](https://creativecommons.org/licenses/by/3.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen