We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Molecular Electrophoretic Technique for Authentication of the Fish Genetic Diversity

Tsai-Hsin Chiu, Yi-Cheng Su, Hui-Chiu Lin and Chung-Kang Hsu

¹Department of Food Science, National PengHu University of Science and Technology

²Seafood Research and Education Center, Oregon State University

³Penghu Marine Biology Research Center, Fisheries Research Institute, COA, EY

^{1,3}Taiwan

²USA

1. Introduction

Cobia (*Rachycentron canadum*) is the sole representative of their family, the Rachycentridae. They are distributed worldwide in tropical and subtropical seas, as the Atlantic and Pacific Oceans (Miao, et al., 2009). There are several species, including cobia, seabream, red porgy, snappers, scads and groupers that are raised by cage culture in Taiwan. Among these cage-cultured fishes, cobia certainly takes a leading distribution in both annual total production (81.9%) and total value production (75.4%) as compared to the rest in Taiwan (Fisheries Agency 2006).

Giant grouper (*Epinephelus lanceolatus*) are also found in tropical and subtropical waters from the Indo-Western Pacific Ocean. It is one of the two largest species of groupers in the world. Due to its fast growth and high price, giant grouper currently is regarded as a favorite species for marine culture in Taiwan (Hseu, et al., 2004).

Red coral trout (*Plectropomus leopardus*) a reef-associated fish in Western Pacific, distributed from southern Japan to Australia and eastward to the Caroline Islands (Zhang, et al., 2010). Only few studies concerning population genetics of *Plectropomus leopardus* has been reported.

All of cobia, giant grouper, and red coral trout are high-valued fish market in Taiwan and neighboring countries, including China, Japan, and Vietnam. For the globalization of the seafood industry, seafood authentication and food safety are very important. We must know that the source of fish or accurately species of the fish. Traditional method to distinguish the fish species was observed the external traits. It can cause the error judgment. Today, DNA-based methods are also more frequently employed for food authentication (Lockley and Bardsley, 2000). It has proven to be reliable, sensitive and fast for many aspects of fish species and food authentication. Asensio et al. (2009) were suggesting that the species-specific PCR method could be potentially used by regulatory agencies as routine control assay for the commercial grouper fillets authentication. PCR-based methods commonly used for fish species identification include PCR-sequencing, random amplified polymorphic DNA (RAPD), inter simple sequence repeat (ISSR). Those methods are simplicity, specificity and sensitivity.

Recently, many researchers have reported for the assessment of genetic structure of aquaculture species such as red mullet (*Mullus barbatus*), Tropical abalone (*Haliotis asinina*), and suminoe oyster (*Crassostrea ariakensis*) using several kinds of molecular markers (Garoia et al., 2004; Tang et al., 2004; Zhang et al., 2005; Maltagliati, et al., 2006), including RAPD, ISSR, AFLP, and RFLP methods. Many molecular methods are available for studying various aspects of wild populations, captive broodstocks and interactions between wild cultured stocks of fish and other aquatic species (Okumus and Ciftci, 2003). Among those methods, RAPD and ISSR technology were cheaper, simple, and fast. And just only one primer could obtain the different profiles for genomic analysis (Welsh and McClelland, 1990). RAPD is simple, rapid and cheap, it have high polymorphism. RAPD analysis has been used to evaluate genetic diversity for species, subspecies and population identification in common carp (Bártfai, et al., 2003), Indian major carps (Barman, et al., 2003). The microsatellite method was already used to study of genetic diversity of other grouper (Antoro, et al., 2006; Ramirez, et al., 2006; Wang, et al., 2007). Zeng, et al. (2008) have report that genetic analysis of Malaysia and Taiwan wild populations of giant grouper by microsatellite method. Their results were shown polymorphic loci in those populations, but they didn't discriminate the wild and cultivated populations of giant grouper. Beside, genetic markers can be suitable for assessing the differences between culture stocks and wild population and monitoring the changes in the genetic variation (Okumus, et al., 2003). Monitoring the genetic diversity of natural populations and fish raised in fish hatcheries is fundamentally important for species conservation. Molecular markers can be very useful in this context (Povh et al., 2008).

In our study, we try to identify the seafood products, including cobia, giant grouper, and red coral trout from cultivated and wild populations by molecular markers, and provide the fish population genetic diversity for seafood management and good monitoring for brood stock management.

2. Methods and material

2.1 Fish sampling and genomic DNA extraction

14 giant grouper (*Epinephelus lanceolatus*) and 14 cobia (*Rachycentron canadum*) were collected from Southern Taiwan, as Penghu Island, Kaohsiung, and Pengtung during 2007-2009. Those were selected in different cultured farms and local markets. 14 of red coral trout (*Plectropomus leopardus*) were collected from Penghu Island during 2009-2011. All samples were described in Table 1. All specimens were confirmed in the laboratory.

Approximately 1g of fish muscle tissue samples were cut into small pieces and pulverized in liquid nitrogen. The powdered fish samples were obtained and extracted genomic DNA using the QIAGEN® DNeasy Blood kit (QIAGEN Inc., Valencia, California) according to manufacturer's instructions. The extracted DNA concentration in 200 µl of sterile water and then the quality of DNA were assessed by a Qubit™ Fluorometer (Invitrogen, USA). The DNAs were stored at -20°C until PCR amplifications.

2.2 PCR-RAPD method

A total of 95 RAPD primers were used for PCR, which were shown in Table 2. Those sequences were obtained from University of British Columbia Biotechnology Laboratory,

No.	Species	Sampling sources	No.	Species	Sampling sources	No.	Species	Sampling sources
1	<i>E. lanceolatus</i>	Wild	R1	<i>R. canadum</i>	Wild	71411	<i>P. leopardus</i>	Wild
E2	<i>E. lanceolatus</i>	Cultivated	R2	<i>R. canadum</i>	Wild	71412	<i>P. leopardus</i>	Wild
E3	<i>E. lanceolatus</i>	Cultivated	R3	<i>R. canadum</i>	Cultivated	71413	<i>P. leopardus</i>	Wild
E4	<i>E. lanceolatus</i>	Cultivated	R4	<i>R. canadum</i>	Wild	82011	<i>P. leopardus</i>	Wild
E5	<i>E. lanceolatus</i>	Wild	R5	<i>R. canadum</i>	Cultivated	91321	<i>P. leopardus</i>	Wild
E6	<i>E. lanceolatus</i>	Cultivated	R6	<i>R. canadum</i>	Wild	91711	<i>P. leopardus</i>	Wild
E7	<i>E. lanceolatus</i>	Wild	R7	<i>R. canadum</i>	Cultivated	91712	<i>P. leopardus</i>	Wild
E8	<i>E. lanceolatus</i>	Wild	R8	<i>R. canadum</i>	Cultivated	71421	<i>P. leopardus</i>	Cultivated
E9	<i>E. lanceolatus</i>	Wild	R9	<i>R. canadum</i>	Wild	71422	<i>P. leopardus</i>	Cultivated
E10	<i>E. lanceolatus</i>	Cultivated	R10	<i>R. canadum</i>	Wild	71423	<i>P. leopardus</i>	Cultivated
E11	<i>E. lanceolatus</i>	Cultivated	R11	<i>R. canadum</i>	Wild	81621	<i>P. leopardus</i>	Cultivated
E12	<i>E. lanceolatus</i>	Cultivated	R12	<i>R. canadum</i>	Wild	80622	<i>P. leopardus</i>	Cultivated
E13	<i>E. lanceolatus</i>	Wild	R13	<i>R. canadum</i>	Wild	81623	<i>P. leopardus</i>	Cultivated
E14	<i>E. lanceolatus</i>	Wild	R14	<i>R. canadum</i>	Wild	91322	<i>P. leopardus</i>	Cultivated

Table 1. Specimens of *E. lanceolatus*, *R. canadum*, and *P. leopardus* fish analyzed and locality where they were collected

RAPD Analysis Kit (Amersham Pharmacia Biotech, Piscataway, NJ), and Operon primer kit (Operon, Advanced Biotechnologies). DNA amplification was performed in a final volume of 25 µl in the “Gene Amp PCR System 2720” thermal cycler (Applied Biosystems Inc., USA). The reaction mix contained 20 mM Tris-HCl, pH8.0, 50 mM KCl, 2 mM MgCl₂, 10 mM dNTPs each (dATP, dCTP, dGTP, dTTP), 20 µM of primer, 2.5 U *Taq*-polymerase (Promega, Co., Wisconsin, USA) and 1 µl of the 10 ng extracted DNA. The pre-amplification PCR procedure was: treatment at 94°C for 5 min, followed by 35 cycles of denaturation at 94°C for 30s, annealing at a primer-specific annealing temperature as table 2 for 30s and extension at 72°C for 30s, and final extension at 72°C for 10 min. The annealing temperature of *Cobia* was 36°C. A 10 µl of the PCR product were analyzed in a 2 % agarose gel in 0.5 X TBE. The electrophoresis was performed at a constant voltage of 150 V for 150 min and 250 V for 1 min. The gel was stained with ethidium bromide and visualized under UV light.

2.3 PCR-ISSR method

ISSR primers of this study were listed in the Table 3. A total 59 primers were screened. Pre-amplification PCR reaction was conducted in 25 µl reaction containing 12.5 µl PCR master mix (Promega, Co., Wisconsin, USA), 1µl each primer, 1 µl of the 10 ng extracted DNA, and 10.5 µl dH₂O. Then, the mixtures were subjected to 94°C for 5 min, followed by 35 cycles of denaturation at 94°C for 30s, 30s at a primer-specific annealing temperature as table 3,

Primer	Sequence of primer (5'-3')	Tm(°C)	references	Primer	Sequence of primer (5'-3')	Tm (°C)	references
RAPD16	GGTGGCGGGA	56	RAPD Primer Set#1, University of British Columbia	RAPD540	CGGACCGCGT	56	Set#6
RAPD17	CCTGGGCCTC	56		RPAD542	CCCATGGCCC	56	
RAPD22	CCCTTGGGGG	56		RAPD563	CGCCGCTCCT	56	
RAPD23	CCCGCCTTCC	56		RAPD571	GCGCGGCACT	56	
RAPD31	CCGGCCTTCC	56		RAPD584	GCGGGCAGGA	56	
RAPD34	CCGGCCCCAA	56		RAPD585	CCCGCGAGTC	56	Set#7
RAPD50	TTCCCCGCGC	56		RAPD592	GGGCGAGTGC	56	
RAPD56	TGCCCCGAGC	56		RAPD595	GTACCGCGC	56	
RAPD63	TTCCCCGCCC	56		RAPD598	ACGGGCGCTC	56	
RAPD64	GAGGGCGGGA	56		RAPD601	CCGCCCCACTG	56	
RAPD65	AGGGGCGGGA	56		RAPD603	ACCCACCGCG	56	
RAPD67	GAGGGCGAGC	56		RAPD606	CGGTCGGCCA	56	
RAPD70	GGGCACGCGA	56		RAPD615	CGTCGAGCGG	56	
RAPD71	GAGGGCGAGG	56		RAPD620	TTGCGCCCCG	56	
RAPD73	GGGCACGCGA	56		RAPD625	CCGCTGCAGC	56	
RAPD81	GAGCACGGGG	56		RAPD626	CCAAGCCCCG	56	
RAPD83	GGGCTCGTGG	56		RAPD640	CGTGGGGCCT	56	
RAPD84	GGGCGCGAGT	56		RAPD647	CCTGTGGGGG	56	
RAPD86	GGGGGGAAGG	56		RAPD769	GGGTGGTGGG	56	Set#8
RAPD87	GGGGGGAAGC	56		RAPD770	GGGAGGAGGG	56	
RAPD88	CGGGGGATGG	56		RAPD771	CCCTCCTCCC	56	
RAPD89	GGGGGCTTGG	56		RAPD772	CCCACCACCC	56	
RAPD94	GGGGGGAACC	56	RAPD Primer Set#2	primer1	GGTGCGGGAA	36	Ready-To-Go. RAPD Analysis Kit (Amersham Pharmacia Biotech, Piscataway, N J)
RAPD95	GGGGGGTTGG	56		primer 2	GTTTCGCTCC	36	
RAPD96	GGCGGCATGG	56		primer 3	GTAGACCCGT	36	
RAPD105	CTCGGTGGG	56		primer 4	AAGAGCCCGT	36	
RAPD106	CGTCTGCCCC	56		primer 5	AACGCGCAAC	36	
RAPD115	TTCCGCGGGC	56		primer 6	CCCGTCAGCA	36	
RAPD149	AGCAGCGTGG	56	Set#3	OPA1	CAGGCCCTTC	56	(Operon, Advanced Biotechnologies)
RAPD157	CGTGGGCAGG	56		OPA2	TGCCGAGCTG	56	
RAPD158	TAGCCGTGGC	56		OPA3	AGTCAGCCAC	56	
RAPD173	CAGGCGGCGT	56		OPA4	AATCGGGCTG	56	
RAPD174	AACGGGCAGC	56		OPA5	AGGGGTCTTG	56	
RAPD190	AGAATCCGCC	56		OPA6	GGTCCCTGAC	56	
RAPD196	CTCCTCCCCC	56		OPA7	GAAACGGGTG	56	
RAPD198	GCAGGACTGC	56		OPA8	GTGACGTAGG	56	
RAPD210	GCACCGAGAG	56		OPA9	GGGTAACGCC	56	
RAPD211	GAAGCGCGAT	56		OPA10	GTGATCGCAG	56	
RAPD218	CTCAGCCCAG	56	Set#4	OPA11	CAATCGCCGT	56	Portman International (China) Limited, Hong Kong
RAPD241	GCCCGACGCG	56		OPA12	TCGGCGATAG	56	
RAPD245	CGCGTGCCAG	56		OPA13	CAGCACCCAC	56	
RAPD270	TGCGCGCGGG	56		OPA14	TCTGTGCTGG	56	
RAPD286	CGGAGCCGGC	56		S514	CAGGATTCCC	56	
RAPD287	CGAACGGCGG	56		S1036	AAGGCACGAC	56	
RAPD319	GTGGCCGCGC	56		S1040	CCTGTTCCT	56	
RAPD480	GGAGGGGGGA	56		S1042	TCGCACAGTC	56	
RAPD534	CACCCCTGTC	56	Set#6	S1201	CCATTCCGAG	56	
RAPD536	GCCCCTCGTC	56					

Table 2. RAPD primers of PCR amplification

extension at 72°C for 30s, and final extension at 72°C for 5 min before analysis by the electrophoresis as described previously.

Primer	Sequence of primer(5'-3')	Tm(°C)	References
ISSR1	(GGAC) ₃ A	48	Pazza et al. (2007)
ISSR2	(GGAC) ₃ C	48	
ISSR3	(GGAC) ₃ T	48	
ISSR4	(TGTC) ₄	48	
ISSR5	(GGAC) ₄	48	
ISSR6	(GGAT) ₄	48	
ISSR7	(TAGG) ₄	48	
ISSR8	(GACA) ₄	48	
ISSR801	(AT) ₈ T	50	Liu et al. (2006)
ISSR817	(CA) ₈ A	48	
ISSR825	(AC) ₈ T	48	
ISSR842	(CA) ₈ YG	52	
ISSR848	(CA) ₈ RG	51	
ISSR850	(GT) ₈ YC	51	
ISSR855	(AC) ₈ YT	50	
ISSR856	(AC) ₈ YA	49	
ISSR858	(TG) ₈ RT	48	
ISSR859	(TG) ₈ RC	52	
ISSR860	(TG) ₈ RA	51	
ISSR888	BDB(CA) ₇	52	
SAS1	(GTG) ₄ GC	55	Maltagliati et al.(2006)
SAS3	(GAG) ₄ G	55	
UBC809	(AG) ₈ G	55	
UBC811	(GA) ₈ C	55	
UBC827	(AC) ₈ G	55	
IT1	(CA) ₈ GT	55	
IT2	(CA) ₈ AC	55	
IT3	(GAG) ₄ AG	55	
PT1	(GT) ₈ C	55	
ISSR807	(AG) ₈ T	48	UBC Primer Set#9, University of British Columbia
ISSR819	(GT) ₈ A	48	
ISSR822	(TC) ₈ A	48	
ISSR831	(AT) ₈ YA	48	
ISSR834	(AG) ₈ YT	48	
ISSR843	(CT) ₈ RA	48	
ISSR852	(TC) ₈ RA	48	
ISSR861	(ACC) ₆	48	
ISSR862	(AGC) ₆	48	
ISSR868	(GAA) ₆	48	
ISSR871	(TA) ₈ RG	48	
ISSR873	(GACA) ₄	48	
ISSR877	(TGCA) ₄	48	
ISSR9	(GAG) ₅ RY	55	Hou et al., 2006
ISSR10	VBV(CA) ₈	54	
ISSR11	VDV(GT) ₈	51	
ISSR12	HVHT(GT) ₇	51	
ISSR13	(CT) ₈ A	49	
ISSR14	(TG) ₈ GT	48	
ISSR15	(AG) ₈ TG	54	
ISSR16	(TC) ₈ C	52	
ISSR17	(TG) ₈ G	53	
ISSR18	(TG) ₆ R	45	

ISSR19	(CA) ₆ RY	44
ISSR20	(GT) ₆ YR	44
ISSR21	(GT) ₆ AY	43
ISSR22	(ACTG) ₄	52
ISSR23	(GACA) ₄	48
ISSR24	(CAC) ₆	57

Table 3. ISSR primers of PCR amplification

2.4 Genetic distances and phylogenetic analysis

Patterns from RAPD and ISSR methods were scored for the presence (1) or absence (0) of clear bands to analyze genetic similarities using the Dice coefficient of similarity. Similarity matrix cluster and phylogenetic analysis was used to reveal association among strains based on the unweighted pair group method with arithmetic averages (UPGMA) using the NTSYSpc software (Numerical taxonomy and multivariate analysis system, version 2.01b, State University of New York, Stony Brook, NY, USA) according to Rohlf (1997).

3. Results

3.1 RAPD method of giant group

A total of 14 giant grouper including cultivate and wild were obtained. Analysis on species of giant grouper in cultivate and wild. For RAPD method amplification, the species, For RAPD method, total 95 of RAPD oligonucleotide primers were used to screening the genetic diversity of giant grouper. There are 21 RAPD primers (22.1%) have polymorphic bands. Total 279 bands were generated by those primers and 86 polymorphic bands (31%). The primer RAPD 115 (5'-TTCCGCGGGC-3') was got the more diversity than other primers, have 8 polymorphic bands (Fig 1). The RAPD 245 primer was generated the less bands, only

Fig. 1. RAPD profiles of the 14 *E. lanceolatus* fish obtained using the primer 115. Lanes M: Bio-100 bp DNA Ladder. Lanes 1-7: *E. lanceolatus* fish wild population (E1, E5, E7, E8, E9, E13, and E14); Lanes 8-14: *E. lanceolatus* fish cultivate population (E2, E3, E4, E6, E10 E11, and E12).

2 polymorphic bands. The sequence and PCR-RAPD condition were listed in Table 2. All primers were generated bands ranging in size from 100 to 3000 bp. The results shown that the ratio of polymorphic bands were between 13.3~66.7% by 21 RAPD primers. For dendrogram analysis, two groups were identified by RAPD 115 primer (Fig 2). E1, E5, E14, E7, E8, and E9 samples were clustered in group I, which were collected from wild population. Group II, which including E13, E10, E2, E3, E4, E6, E11, and E12 samples. For Group II, all samples were belonged to cultivated populations. For giant grouper, wild (seven samples) and cultivated (seven samples) populations of giant grouper can be discriminated by RAPD method.

Fig. 2. UPGMA consensus dendrogram of dissimilarity among individuals analyzed using the primer RAPD 115.

3.2 ISSR method of giant group

Results of ISSR analysis, 59 primers were used in this study. According the results of ISSR method, 17 primers (29%) have polymorphic patterns. Total 166 bands were generated, 58 polymorphic bands (34.9%). The primer ISSR IT3 was got the more diversity than other primers, have 20 bands. The ISSR 15 primer was generated the less bands, only 3 bands. All the polymorphic patterns were ranged between 100~3000 bp. ISSR primer 868 (5'-(GAA)₆-3') was better distinguished than other primers. The result was shown in Fig 3. For giant grouper, the patterns of ISSR primer868 could discriminate giant grouper between wild and cultivated populations. For dendrogram analysis, four groups were clustered by ISSR primer868 primer (Fig 4). Among those groups, Group I, Group III, and Group IV were collected from wild population. Samples were clustered in Group II were from cultivated populations. We also found that the results of ISSR method have the same tend to RAPD method. ISSR method was more discriminate ability than RAPD method.

Fig. 3. ISSR profiles of the 14 giant grouper fish obtained using the primer ISSR 868. Lanes M: Bio-100bp DNA Ladder. Lanes 1-7: giant grouper fish wild population (E1, E5, E7, E8, E9, E13, and E14); Lanes 8-14: giant grouper fish cultivated population (E2, E3, E4, E6, E10, E11, and E12).

Fig. 4. UPGMA consensus dendrogram of dissimilarity among individuals analysed using the primer ISSR 868.

3.3 RAPD and ISSR methods of cobia

Ninety-five RAPD primers and 59 ISSR primers were used for PCR amplification. The results were shown that all the cobia samples were the same patterns and no polymorphic

bands. The primer ISSR UBC809 and RAPD31 were generated 12 to 17 bands and in size from 100 to 2000 bp. The results were also shown in Fig 5 and Fig 6.

Fig. 5. ISSR profiles of the 14 *R. canadum* fish obtained using the primer UBC 809. Lanes M: Bio-100 bp DNA Ladder. Lanes: 1, 2, 4, 6, 9, 10, 11, 12, 13, and 14 (R1, R2, R4, R6, R9, R10, R11, R12, R13, and R14) *R. canadum* fish wild population; Lanes: 3, 5, 7, and 8 (R3, R5, R7, and R8) *R. canadum* fish cultivate population.

Fig. 6. RAPD profiles of the 14 *R. canadum* fish obtained using the primer 31. Lanes M: Bio-100 bp DNA Ladder. Lanes : 1, 2, 4, 6, 9, 10, 11, 12, 13, and 14 (R1, R2, R4, R6, R9, R10, R11, R12, R13, and R14) *R. canadum* fish wild population; Lanes: 3, 5, 7, and 8 (R3, R5, R7, and R8) *R. canadum* fish cultivate population.

Sequence variability of mitochondrial DNA regions was low between the six cobia (Garnet, et al., 2002). These results were also similar in our study. Both RAPD and ISSR methods have no different patterns. Hence, this could provide more useful information of molecular genetic data in population and stock enhancement studies.

3.4 ISSR methods of red coral trout

For screening ISSR primers, the ISSR primer15 (ISSR15: 5'-(AG)₈TG -3') was better distinguished than other primers. The result was shown in Fig 7. The primer ISSR 15 was generated 10 to 16 bands and in size from 200 to 2000 bp. For dendrogram analysis, three groups were identified. Group I, the 71412, 71413, 71422, and 82011 were clustered in the group. Group II, including 81621, 91321, 71411, and 91712 samples; group III including 71423, 80622, 91322, 71421, and 81623 samples. All the nodes of the dendrograms ranged from 90 to 100%. The result was shown in Fig 8.

Fig. 7. ISSR profiles of the 14 red coral trout obtained using the primer ISSR 15. Lanes M: Bio-100 bp DNA Ladder. Lanes: 4, 5, 6, 8, 9, 10, and 12 (71411, 71412, 71413, 82011, 91321, 91711, and 91712) red coral trout fish wild population; Lanes: 1, 2, 3, 7, 11, 13, and 14 (71421, 71422, 71423, 81621, 80622, 81623, and 91322) red coral trout fish cultivate population.

4. Discussions

RAPD and ISSR methods were generally used for genetic diversity and populations study; those methods also could be used to analyze the breeding relationship. For species identification and genetic resource/diversity analysis, RAPD and microsatellites method were recommended (Liu & Cordes, 2004). The RAPD techniques has been used for discrimination of populations of species of the genus *Barbus*, grouper, Nile perch and wreck fish, salmonids, among others (Partis & Wells, 1996; Callejas & Ochando, 2001; Asensio et al., 2002; Jin, Cho, Seong, Park, Kong & Hong, 2006). Genetic analysis with RAPD markers is relatively easy, fast, and efficient. RAPD analysis, however, may not be practical for identifying interbreed species (Martinez, Elvevoll & Haug, 1997). SSRs are inherited in a co-dominant fashion. This allows one to discriminate between homo- and heterozygous state, and increases the efficiency of genetic mapping and population genetic studies. ISSR markers have recently been used successfully for genetic analysis in hatchery and wild *Paralichthys olivaceus* strains. It indicates that molecular marker systems contribute greater levels of capability for the detection of polymorphism, and provide a better solution for the assessment of genetic variations (Shikano, 2005; Liu, Chen, Li & Li, 2006).

Fig. 8. UPGMA consensus dendrogram of dissimilarity among individuals analysed using the primer ISSR 15.

According to our results, RAPD and ISSR method can be effectively discriminate giant grouper from different sources, such as cultivate and wild, even if fish are from different cultivate farms. But cobia and red coral trout were less discriminate ability. The study found that ISSR and RAPD methods were positively high correlations. Giant grouper species have highly genetic diversity. A comparison of RAPD and ISSR patterns in 14 giant grouper samples, ISSR primers have higher polymorphism and fewer bands than those of RAPD primers. It could provide simple and convenient method to discriminate genetic variation of giant grouper samples. In this study, ISSR method could distinguish genetic variation within specie and different populations. Some reports also have suggested that ISSR may reveal a much higher numbers of polymorphic fragments per primer than those of RAPD (Esselman, et al., 1999). Among these markers, microsatellite DNAs have revolutionized the use of molecular genetic markers in the applications mentioned before, and the markers are destined to dominate this type of studies in the coming years (Asensio, 2007). It also has been revealed as important tools in studies regarding the genetic structure of populations, phylogeographic relations and phylogenetic reconstruction in fish (Antunes, et al., 2010).

5. Conclusion

We developed DNA molecular marker techniques which could be used to generate information for fish genetic diversity, species identification, trace genetic variation between different individuals in aquaculture, authenticate fish, fishery products and provide good reference resources for species sources and relationships.

6. Acknowledgment

This research project was supported by the Council of Agriculture (COA) (97AS-4.1.2-AI-I2).

7. References

- Antoro, S.; Na-Nakom, U. & Koedprang, W. (2006). Study of Genetic Diversity of Orange-Spotted, *Epinephelus coioides*, from Thailand and Indonesia Using Microsatellite Markers. *Marine Biotechnology*, Vol.8, pp. 17-26, ISSN 1432-1793
- Antunes, R. S. P.; Gomes, V. N.; Prioli, S. M. A. P.; Prioli, R. A.; Júlio Jr.; H. F. ; Prioli, L. M.; Agostinho, C. S. & Prioli, A. J. (2010). Molecular Characterization and Phylogenetic Relationships Among Species of the Genus *Brycon* (Characiformes: Characidae) from Four Hydrographic Basins in Brazil. *Genetics and Molecular Research*, Vol.9, pp. 674-684, ISSN 1676-5680
- Asensio, L.; González, I.; Fernández, A.; Rodríguez, M. A.; Lobo, E.; Hernández, P. E.; García, T. & Martín, R. (2002). Application of Random Amplified Polymorphic DNA (RAPD) Analysis for Identification of Grouper (*Epinephelus guaza*), Wreck fish (*Polyprion americanus*), and Nile perch (*Lates niloticus*) Fillets. *Journal of Food Protection*, Vol.65, pp.432-435, ISSN 0362-028x
- Asensio, L. (2007). PCR-Based Methods for Fish and Fishery Products Authentication. *Trends in Food Science and Technology*, Vol.18, pp. 558-566, ISSN 0924-2244
- Asensio, L.; González, I.; Rojas, M.; García, T. & Martín, R. (2009). PCR-Based Methodology for the Authentication of Grouper (*Epinephelus marginatus*) in Commercial Fish Fillets. *Food control*, Vol.20, pp. 618-622, ISSN 0956-7135
- Barman, H. K.; Barat, A.; Yadav, B. M.; Banerjee, S.; Meher P. K.; Reddy, P. V. G. K. & Jana, R. K. (2003). Genetic Variation Between Four Species of Indian Major Carps as Revealed by Random Amplified Polymorphic DNA Assay. *Aquaculture*, Vol. 217, pp. 115-123, ISSN 0044-8486
- Bártfai, R.; Egedi, S.; Yue, G. H.; Kovács, B.; Urbányi, B.; Tamás, G.; Horváth, L. & Orbán, L. (2003). Genetic Analysis of Two Common Carp Broodstocks by RAPD and Microsatellite Markers. *Aquaculture*, Vol. 219, pp. 157-167, ISSN 0044-8486
- Callejas, C. & Ochando, M. D. (2001). Molecular Identification (RAPD) of the Eight Species of the Genus *Barbus* (Cyprinidae) in the Iberian Peninsula. *Journal of Fish Biology*, Vol. 59, pp. 1589-1599, ISSN 1095-8649
- Esselman, E. J.; Jianqiang, L.; Crawford, D. J.; Windus, J. L. & Wolfe, A. D. (1999). Clonal Diversity in the Rare *Calamagrostis porteri* ssp. *insperata* (Poaceae): Comparative Results for Allozymes and Random Amplified Polymorphic DNA and Intersimple Sequence Repeat Markers. *Molecular Ecology*, Vol. 8, pp. 443-453, ISSN 1365-294X
- Garber, A. F.; Grater, W. D.; Stuck, K. C. & Franks, J. S. (2002). Characterization of the Mitochondrial DNA Control Region of Cobia, *Rachycentron vanadium*, from Mississippi Coastal Waters. *Gulf and Caribbean Fisheries Institute*, Vol. 53, pp. 570-578, ISSN 0072-9019
- Garoia, F.; Guarniero, I.; Piccinetti, C. & Tinti, F. (2004). First Microsatellite Loci of Red Mullet (*Mullus barbatus*) and Their Application to Genetic Structure Analysis of Adriatic Chared Stock. *Marine biotechnology*, Vol. 6, pp. 446-452, ISSN 1436-2236
- Hou, L.; Lü, H.; Zou, X.; Bi, X.; Yan, D. & He, C. (2006). Genetic Characterizations of *Macra Veneriformis* (Bivalve) Along the Chinese Coast Using ISSR-PCR Marker. *Aquaculture*, Vol. 261, pp. 865-871, ISSN 0044-8486
- Hseu, J.-R.; Hwang, P.-P. & Ting, Y.-Y. (2004). Morphometric Model and Laboratory Analysis of Intracohort Cannibalism in Giant Grouper *Epinephelus lanceolatus* Fry. *Fisheries Science*, Vol. 70, pp. 482-486, ISSN 1444-2906

- Jin, L. G.; Cho, J. G.; Seong, K. B.; Park, J. Y.; Kong, I. S. & Hong, Y. K. (2006). 18 rRNA Gene Sequences and Random Amplified Polymorphic DNA Used in Discriminating Manchurian Trout from Other Freshwater Salmonids. *Fisheries Science*, Vol. 72, pp. 903-905, ISSN 1444-2906
- Liu, Y.-G.; Chen, S.-L.; Li, J. & Li, B.-F. (2006). Genetic Diversity in Three Japanese Flounder (*Paralichthys olivaceus*) Populations Revealed by ISSR Markers. *Aquaculture*, Vol. 255, pp. 565-572, ISSN 0044-8486
- Liu, Z. J. & Cordes, J. F. (2004). DNA Marker Technologies and Their Applications in Aquaculture Genetics. *Aquaculture*, Vol. 238, pp. 1-37, ISSN 0044-8486
- Lockley, A. & Bardsley, R. (2000). DNA-Based Methods for Food Authentication. *Trends in Food Science & Technology*, Vol. 11, pp. 67-77, ISSN 0924-2244
- Maltagliati, F.; Lai, T.; Casu, M.; Valdesalici, S. & Castelli, A. (2006). Identification of Endangered Mediterranean Cyprinodontiform Fish by Means of DNA Inter-Simple Sequence repeats (ISSRs). *Biochemical systematics and Ecology*, Vol. 34, pp. 626-634, ISSN 0305-1978
- Martinez, I.; Elvevoll, E. O. & Haug, T. (1997). RAPD Typing of North-East Atlantic Minke Whale (*Balaenoptera acutorostrata*). *ICES Journal of Marine Science*, Vol. 54, pp. 478-484, ISSN 1095-9289
- Miao, S.; Jen, C. C.; Huang, C. T. & Hu S.-H. (2009). Ecological and Economic Analysis for Cobia *Rachycentron canadum* Commercial Cage Culture in Taiwan. *Aquaculture International*, Vol. 17, pp. 125-141, ISSN 0967-6120
- Okumus, İ. & Çiftçi, Y. (2003). Fish Population Genetics and Molecular Marker: II - Molecular Markers and Their Applications in Fisheries Aquaculture. *Turkish Journal of Fisheries and Aquatic Sciences*, Vol. 3, pp. 51-79, ISSN 1303-2712
- Partis, L. & Wells, R. J. (1996). Identification of Fish Species Using Random Amplified Polymorphic DNA (RAPD). *Molecular and Cellular Probes*, Vol. 10, pp. 435-441, ISSN 0890-8508
- Pazza, R.; Kavalco K. F.; Prioli Soˆnia M. A. P.; Prioli, A. J. & Bertollo, L. A. C. (2007). Chromosome Polymorphism in *Astyanax Fasciatus* (Teleostei, Characidae), Part 3: Analysis of the RAPD and ISSR Molecular Markers. *Biochemical Systematics and Ecology*, Vol. 35, pp. 843-851, ISSN 0305-1978
- Povh, J. A.; Lopera-Barrero, N. M.; Ribeiro, R. P.; Lupchinski Jr. E.; Gomes, P. C. & Lopes, T. S. (2008). Genetic Monitoring of Fish Repopulation Programs Using Molecular Markers. *Ciencia e Investigación Agraria*, Vol. 35, pp. 1-10, ISSN 0718-1620
- Ramirez, M. A.; Pareicia-Acevedo, J. & Planas, S. (2006). New Microsatellite Resources for Groupers (Serranidae). *Molecular Ecology Notes*, Vol. 6, pp. 813-817, ISSN 1755-0998
- Rohlf, F. J. (1997). "NTSYS-pc v. 2.01b Numerical Taxonomy and Multivariate Analysis System." Applied Biostatistics, Inc., Setauket, NY.
- Shikano, T. (2005). Marker-Based Estimation of Heritability for Body Color Variation in Japanese Flounder *Paralichthys Olivaceus*. *Aquaculture*, Vol. 249, pp. 95-105, ISSN 0044-8486
- Tang, S.; Tassanakajon, A.; Klinbunga, S.; Jarayabhand, P. & Menasveta, P. (2004). Population Structure of Tropical Abalone (*Haliotis asinina*) in Coastal Waters of Thailand Determined Using Microsatellite Markers. *Marine Biotechnology*, Vol. 6, pp. 604-611, ISSN 1432-1793

- Wang, S. F.; Du, J. Y. & Wang, J. (2007). Identification of *Epinephelus malabaticus* and *Epinephelus coioides* Using DNA Markers. *Acta Oceanology Sinica*, Vol. 26, pp. 122-129, ISSN 1869-1099
- Welsh, J. & McClelland, M. (1990). Fingerprinting Genomes Using PCR Arbitrary Primers. *Nucleic Acids Research*, Vol. 18, pp. 7213-7218, ISSN 1362-4962
- Zeng, H.-S.; Ding, S.-X.; Wang, J. & Su, Y.-Q. (2008). Characterization of Eight Polymorphic Microsatellite Loci for the Giant Grouper (*Epinephelus lanceolatus* Bloch). *Molecular Ecology Resources*, Vol. 8, pp. 805-807, ISSN 1755-0998
- Zhang, Q.; Allen Jr. S. K. & Reece, K. S. (2005). Genetic Variation in Wild and Hatchery Stocks of Suminoe Oyster (*Crassostrea ariakensis*) Assessed by PCR-RFLP and Microsatellite Markers. *Marine Biotechnology*, Vol. 7, pp. 588-599, ISSN 1432-1793
- Zhang, J.; Liu, H. & Song, Y. (2010). Development and Characterization of Polymorphic Microsatellite Loci for a Threatened Reef Fish *Plectropomus leopardus*. *Conservation Genetics Resources*, Vol. 2 (Supplement 1), pp. 101-103, ISSN 1877-7260

IntechOpen

Gel Electrophoresis - Advanced Techniques

Edited by Dr. Sameh Magdeldin

ISBN 978-953-51-0457-5

Hard cover, 500 pages

Publisher InTech

Published online 04, April, 2012

Published in print edition April, 2012

As a basic concept, gel electrophoresis is a biotechnology technique in which macromolecules such as DNA, RNA or protein are fractionated according to their physical properties such as molecular weight or charge. These molecules are forced through a porous gel matrix under electric field enabling uncounted applications and uses. Delivered between your hands, a second book of this Gel electrophoresis series (Gel Electrophoresis- Advanced Techniques) covers a part, but not all, applications of this versatile technique in both medical and life science fields. We try to keep the contents of the book crisp and comprehensive, and hope that it will receive overwhelming interest and deliver benefits and valuable information to the readers.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Tsai-Hsin Chiu, Yi-Cheng Su, Hui-Chiu Lin and Chung-Kang Hsu (2012). Molecular Electrophoretic Technique for Authentication of the Fish Genetic Diversity, Gel Electrophoresis - Advanced Techniques, Dr. Sameh Magdeldin (Ed.), ISBN: 978-953-51-0457-5, InTech, Available from: <http://www.intechopen.com/books/gel-electrophoresis-advanced-techniques/molecular-electrophoretic-technique-for-authentication-of-the-fish-genetic-diversity>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](https://creativecommons.org/licenses/by/3.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen