

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

186,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



The Contribution of Fibronectin ED-A Expression to Myofibroblast Transdifferentiation in Diabetic Renal Fibrosis

Keisuke Ina, Hirokazu Kitamura,
Shuji Tatsukawa and Yoshihisa Fujikura
Oita University
Japan

1. Introduction

The number of cases in which hemodialysis therapy for diabetic nephropathy is required has been increasing. Currently, the goal of treatment for renal fibrosis is not only to prevent the development and progression of the disease, but also to promote its remission and regression.

It is well known that glomerulosclerosis and tubulointerstitial fibrosis occur during the early stages of diabetic nephropathy. It has been demonstrated that the development of tubulointerstitial lesions is more closely correlated with a progressive decline in renal function compared to glomerular lesions (Bohle et al., 1991). Tubulointerstitial fibrosis ultimately leads to renal failure as a result of renal atrophy. The myofibroblasts emerging in tubulointerstitial fibrosis tissue have been indicated to play a crucial role in the development and progression of fibrosis (Simonson, 2007). They overproduce extracellular matrix (ECM) molecules, including type I collagen and fibronectin, and repress ECM degradation through the production of tissue inhibitor of metalloproteinase-1 (TIMP-1) (Edwards et al., 1987) and plasminogen activator inhibitor-1 (PAI-1) in response to transforming growth factor- β_1 (TGF- β_1), which is increased in DN (Laiho et al., 1987), followed by ECM accumulation (i.e. fibrosis). Furthermore, the myofibroblasts may induce fibrosis tissue contraction via increased cell contraction promoted by α -smooth muscle actin (α -SMA) expression, leading to renal atrophy and failure. Myofibroblasts are considered to be derived from various kinds of cells, e.g. fibroblasts (Strutz and Zeisberg, 2006), epithelial cells (Iwano et al., 2002), endothelial cells (Zeisberg et al., 2008), pericytes (Humphreys et al., 2010), and bone marrow-originated cells (Keeley et al., 2010), via stimulation by TGF- β_1 .

Although α -SMA expression, which is the most pronounced characteristic of myofibroblasts, has been considered to cause tissue contraction, the detailed mechanism has not yet been determined. Myofiboblasts were first described in wound-healing granulation tissue

(Gabbiani et al., 1971), where they were referred to as “modified fibroblasts” on the basis of their ultrastructural identification in transmission electron microscopic studies. These cells include a fibrillar system corresponding to stress fibers, nuclear indentation implying cell contraction, and cell-to-ECM and cell-to-cell junctions. Thus, modified fibroblasts (myofibroblasts) were described as an intermediate type of cell between fibroblasts and smooth muscle cells due to their ability to produce ECM and contract. Later investigators have noted that α -SMA expression in the stress fibers was the major characteristic of myofibroblasts.

At present, cells with α -SMA-positive stress fibers in the interstitium, except for vascular components, are recognized to be myofibroblasts. In vertebrate tissues, six actin isoforms have been identified. Four actin isoforms, including the α -skeletal, α -cardiac, α -vascular and γ -enteric isoforms, are tissue-restricted and involved in cell contraction. The other two actin isoforms, the β -cytoplasmic and γ -cytoplasmic isoforms, are ubiquitously expressed. These have been considered to constitute noncontractile cytoskeletons. The α -vascular actin is the SMA isoform, referred to as “ α -SMA”. During myofibroblast transdifferentiation, the actin isoform transitions, at least in part, from the cytosolic actin to α -SMA. It has been proposed that myofibroblasts have the ability to contract due to their expression of α -SMA.

Fibronectin, a 440 kDa dimeric glycoprotein, is one of the ECM molecules that exerts various functions, including adhesion, migration and differentiation. The fibronectin subunit is composed of three internally homologous repeats and contains binding sites for cell surface receptors and for other ECM components (Magnusson & Mosher, 1998; Peterson et al., 1983). Fibronectin polymorphisms arise from the alternative splicing of mRNA at three regions, ED-A, ED-B and IIICS (Norton & Hynes, 1987) (Fig. 1). The two former sequences are either omitted or included, while the latter one varies in length. Fibronectins are present in an insoluble form in the ECM, and in a soluble form in circulation, namely as cellular fibronectin and plasma fibronectin. Only cellular fibronectin contains ED-A and ED-B (Vartio et al., 1987). The fibronectin ED-A and fibronectin ED-B isoforms are predominantly expressed in embryonic tissue, and are seldom found in normal adult tissues. However, fibronectin ED-A has been demonstrated to be re-expressed in a variety of reactive and pathological processes, such as wound healing (Ffrench-Constant et al., 1989), arterial intimal thickening (Glukhova et al., 1989) and liver fibrosis (Jamagin et al., 1994). *In vitro*, TGF- β_1 has been shown to increase fibronectin production, and it regulates the splicing pattern of fibronectin pre-messenger RNA, followed by expression of the ED-A domain (Balza et al., 1988; Borsi et al., 1990). Moreover, it has been suggested that the fibronectin ED-A isoform modulates various cells to generate myofibroblasts which express α -SMA (Bochaton-Piallat et al., 2000; Ffrench-Constant et al., 1989; Jamagin et al., 1994). Gabbiani and coworkers demonstrated that the fibronectin ED-A isoform plays a crucial role in TGF- β_1 -induced myofibroblast transdifferentiation in an *in vitro* study using human subcutaneous fibroblasts (Serini et al., 1998).

In the present study, we examined whether TGF- β_1 induces the expression of the ED-A domain in fibronectin, and whether this, in turn, leads to α -SMA expression in renal fibroblasts by using an *in vitro* model of fibrosis due to diabetic nephropathy. Furthermore, whether the fibronectin ED-A isoform stimulates the contraction of the collagen gel through the expression of α -SMA and the accumulation of type I collagen was examined.




Fig. 1. The molecular structure of fibronectin ED-A and ED-B.

The fibronectin subunit is composed of three different repeating sequences: type I (rectangles), type II (ovals) and type III (circles). Sets of repeats constitute binding domains for fibrin, fibronectin (FN), collagen, cells and heparin as described. Sequence variations can occur in all three points of the fibronectin subunit: extra domains A and B (ED-A and ED-B, respectively) and the IIICS (Wierzbicka-Patynowski & Schwarzbauer, 2003). Arrows indicate approximate intron positions determined in the murine gene.

2. Materials and methods

2.1 Materials

Porcine type I collagen solution was purchased from Nitta Gelatin (Osaka, Japan), recombinant human TGF- β_1 was purchased from R and D Systems (MN, USA), and the mouse monoclonal antibodies against α -SMA and ED-A were obtained from Sigma-Aldrich (MO, USA) and Abcam (Cambridge, UK), respectively. The fluorescein isothiocyanate (FITC)-conjugated goat anti-mouse IgG secondary antibody against these monoclonal antibodies was purchased from Caltag Laboratories (CA, USA). Rat type I collagen ELISA kits were obtained from Cusabio (Wuhan, China).

2.2 Cell culture

Normal rat kidney fibroblasts (NRK 49F cells) were obtained from the RIKEN Cell Bank (Tsukuba, Japan). The cells were maintained in Dulbecco's minimal essential medium (D-MEM) (Nissui Pharmaceutical, Tokyo, Japan) containing 100 IU ml⁻¹ penicillin, 100 μ g ml⁻¹ streptomycin, and 10% fetal bovine serum (FBS) (JRH Biosciences, KS) at 37°C in a humidified, 5% CO₂ atmosphere. FBS had been heat-inactivated at 56°C for 30 min prior to use. Cells from passage 3 to passage 8 were used in the experiments described below.

Monolayer cultures of 2×10^4 cells ml^{-1} were grown on two-well Labtek Chamber Slides (AS ONE, Osaka, Japan). The cell cultures were divided into 4 groups: a control group, cultured with 0.5% FBS; a TGF- β_1 group, cultured with 0.5% FBS + 5 ng ml^{-1} TGF- β_1 ; an anti-ED-A antibody group, cultured with 0.5% FBS + 5 ng ml^{-1} TGF- β_1 + 10 $\mu\text{g ml}^{-1}$ anti-ED-A antibody; an IgG group, cultured with 0.5% FBS + 5 ng ml^{-1} TGF- β_1 + 10 $\mu\text{g ml}^{-1}$ mouse IgG (negative control, R & D Systems, Inc., MN, USA). Furthermore, each group was sub-divided into two groups: the 1D (1 day) group, which was cultured for 24h after the initiation of culture; the 2D (2 day) group, which was cultured for 24h after refeeding the cells the same medium 24 h after culture initiation. The cells and the supernatants of all groups were subjected to immunofluorescence studies for fibronectin ED-A or α -SMA, and to an ELISA for type I collagen, respectively.

2.3 Immunofluorescence microscopy for fibronectin ED-A and α -SMA

The cells from monolayer cultures were fixed in 4% paraformaldehyde in 0.1 M phosphate-buffered saline (pH 7.4) for 10 min. After rinsing, the cells were permeabilized with 0.5% Triton X-100 for 20 min in order to identify α -SMA in the cells, this process was omitted in studies for detecting fibronectin ED-A. After washing, the cells were incubated with the primary antibody against α -SMA or fibronectin ED-A for 2 h at room temperature. After being rinsed, they were incubated with FITC-conjugated goat anti-mouse IgG secondary antibody for 1 h at room temperature. After staining, the cells were rinsed, observed and photographed using an Olympus BX 60 microscope equipped with epifluorescence optics. The assessment of the percentage of cells with α -SMA-positive stress fibers was performed as follows: when there were one or more stress fibers with α -SMA-positive staining in a cell, the cell was considered to be α -SMA-positive and was referred to as a “myofibroblast”. The percentage of myofibroblasts was represented by counting the α -SMA-positive cells among 100 to 500 cells.

2.4 Gel contraction assay

A total of 7 ml of a collagen solution (3 mg/ml stock) was gently mixed with 2 ml of 5-fold-concentrated D-MEM on ice. The mixture was adjusted to pH 7.4 with 1 ml of 200 mM HEPES plus 2.2% NaHCO_3 in distilled water. This collagen mixture (0.3ml) was added to each well of a 24 well plate (Becton-Dickinson Labware, NJ). Then, the collagen gel was formed by the incubation of the mixture for 30 min at 37°C. After gelatinization, an NRK 49F cell suspension (2×10^4 cells ml^{-1}) was poured on the gel in each well. Preincubation of the cell-populated collagen lattice was performed in D-MEM + 10% FBS for 24 h. Next, the gels were rinsed with serum-free D-MEM three times for 30 min. Then, the gel cultures were divided into 4 groups similar to the immunofluorescence studies: a control group, cultured in 0.5% FBS; a TGF- β_1 group, cultured in 0.5% FBS + 5 ng ml^{-1} TGF- β_1 ; an anti-ED-A antibody group, cultured in 0.5% FBS + 5 ng ml^{-1} TGF- β_1 + 10 $\mu\text{g ml}^{-1}$ anti-ED-A antibody and an IgG group, cultured in 0.5% FBS + 5 ng ml^{-1} TGF- β_1 + 10 $\mu\text{g ml}^{-1}$ mouse IgG. The gels were then detached from the lateral wall and bottom of each well with a microspatula and cultured for 48 h. After 48 h in culture, the diameter of each gel was measured with a ruler to assess the gel contraction after treatment with the reagents. In addition, to confirm that the shrinkage of the cell-populated collagen lattice occurred due to gel contraction, rather than gel digestion, another study was performed. For this study, after gelatinization, the

NRK49F cell suspension (5000 cells per 5 μ l) was placed in a very small area near the margin of the gel, and incubated for 1h at 37°C. Then, preincubation was done as above. After rinsing, the gels were cultured in the control medium and the medium with 5ng ml⁻¹ TGF- β_1 for 48h at 37°C following gel detachment. After 48h, the shapes of the gels were observed.

2.5 Enzyme-linked immunosolvent assay for type I collagen accumulation in the supernatants of the cultures

The supernatants from monolayer cultures were added to the wells of 96 well plate, which were pre-coated with a mouse monoclonal antibody against rat type I collagen, and were incubated for 2 h at 37°C. After removing the liquid from each well, a biotin-conjugated anti-mouse IgG secondary antibody solution was added, and cells were incubated for 1 h at 37°C. After rinsing, each well was exposed to the horseradish peroxidase-avidin solution for 1 h at 37°C. After washing, the TMB substrate was added, and cells were incubated for 20 min at 37°C. Then, the stop solution was added to each well, while the optical density of each well was determined using a microplate reader, the Multiskan GO (Thermo Fisher Scientific, Yokohama, Japan), at 450 nm. The value of the optical density was converted to the type I collagen concentration in the supernatant of the monolayer culture.

2.6 Statistical analysis

The experimental data are presented as the means \pm SD. The statistical analyses were performed using Student's *t*-test. Differences were considered to be statistically significant at $P < 0.05$.

3. Results

3.1 TGF- β_1 - induced the expression of ED-A and α -SMA

ED-A expression was induced in a dot or short linear pattern on the cell surface by TGF- β_1 by 24 h after initiation of culture (the 1D • TGF- β_1 group), and was further increased after the 48 h culture (the 2D • TGF- β_1 group) (Figs. 2c, 2d). On the other hand, the expression of intracellular α -SMA was not observed 24 h after the initiation of culture (the 1D • TGF- β_1 group), but was induced in stress fiber formations after 48 h in culture (the 2D • TGF- β_1 group) (Figs. 2g, 2h). The expression of ED-A and α -SMA was not observed in the cells cultured in the medium without TGF- β_1 (the 1D and 2D control groups) (Figs. 2a, 2b, 2e, 2f).

3.2 The effects of the anti-ED-A antibody on TGF- β_1 - induced α -SMA expression

The TGF- β_1 - induced α -SMA expression was markedly decreased by adding an anti-ED-A antibody to the medium (Fig. 3A). The rates of α -SMA-positive cells in stress fiber formations were as follows: the 2D control group (0%), the 2D TGF- β_1 group (97%), the 2D anti-ED-A antibody group (36%), and the 2D IgG group (95%) (Fig. 3B).

3.3 The characteristics of TGF- β_1 -induced gel shrinkage

When the cells were ubiquitously poured on the gel, TGF- β_1 -induced gel shrinkage occurred in the shape of a circle after culture for 48h (Fig. 4Aa). The diameters of gels were $99.4 \pm 1.0\%$




Fig. 2. Immunofluorescent staining for ED-A and α -SMA induced by TGF- β_1 . a, c, e, g, ID groups; b, d, f, h, 2D groups; a, b, c, d, Immunostaining for ED-A; e, f, g, h, Immunostaining for α -SMA; a, b, e, f, control groups; c, d, g, h, TGF- β_1 groups. a, No immunostaining. b, No immunostaining. c, Immunostaining for ED-A is shown by the dot or short linear pattern on the cell surface. d, Immunostaining for ED-A was increased. e, No immunostaining. f, No immunostaining. g, No immunostaining. h, Immunostaining for α -SMA is seen in the stress fiber formation in the cells. Bar, 10 μ m.

in the control medium group and $86.4 \pm 1.7\%$ in the TGF- β_1 group ($p < 0.001$) (Fig. 4Ab). On the other hand, when the cells were placed in a small region close to the margin of the gel, the gel exhibited indentation toward the cell mass in response to TGF- β_1 (Fig. 4B). If gel shrinkage occurs due to gel digestion, the gel would reveal the features of digestion, such as the dissolution of the gel around the cell mass, and would form a thin layer or disappear. However, because the present study did not show these features, gel digestion was not considered to be responsible for the observed gel shrinkage. The observed indentation was thought to be caused by retraction of the gel resulting from cell contraction.

3.4 The effects of the anti-ED-A antibody on TGF- β_1 -induced gel contraction

Gel contraction did not occur in the control medium throughout the experimental period. In addition, no gel contraction was observed in any of the 1D groups. In the 2D groups, TGF- β_1 remarkably evoked gel contraction. The treatment with the anti-ED-A antibody was shown to repress TGF- β_1 -induced gel contraction. The diameters of the gels in the four groups were as follows: the 2D control group (100%), the 2D TGF- β_1 group ($84.0 \pm 1.9\%$), the 2D anti-ED-A antibody group ($89.5 \pm 1.9\%$, $p < 0.05$ vs the 2D TGF- β_1 group), and the 2D IgG group ($83.7 \pm 1.6\%$, $p < 0.05$ vs the 2D anti-ED-A antibody group) (Fig. 5).

3.5 The effects of the anti-ED-A antibody on TGF- β_1 -stimulated type I collagen accumulation in the supernatants of the cell cultures

There were no significant differences in the accumulation of type I collagen in the cell supernatants between the three 1D groups (the control group, 1420 ± 200 ; the TGF- β_1 group, 1470 ± 100 ; the anti-ED-A antibody group, 1510 ± 160 ; the IgG group, 1513 ± 26 pg ml $^{-1}$). In the 2D groups, the accumulation of type I collagen in the supernatant of the TGF- β_1 group was increased. The increased accumulation of collagen was repressed by addition of the anti-ED-A antibody to the medium with TGF- β_1 (Fig. 5). The concentrations of the supernatants in the four 2D groups (control, TGF- β_1 , anti-ED-A antibody, IgG) were 1980 ± 220 , 2730 ± 110 , 2490 ± 18 , 2570 ± 3 pg ml $^{-1}$, respectively ($p < 0.001$, the control group vs the TGF- β_1 group; $p < 0.05$, the TGF- β_1 group and the IgG group vs the anti-ED-A antibody group) (Fig. 6).

4. Discussion

Myofibroblasts expressing α -SMA have been shown to emerge in fibrotic diseases (Tomasek et al., 2002). In general, α -SMA is considered to be profoundly involved in the progression of fibrosis, but the detailed significance of α -SMA expression remains unresolved. For example, it is unclear whether myofibroblasts are really essential for fibrosis. In contrast to general considerations, there have been some reports indicating that myofibroblasts are not necessary for the progression of fibrosis. For example, Takeji et al. suggested that α -SMA expression provided benefits for fibrosis (Takeji et al., 2006). It was demonstrated that defects in α -SMA enhanced the progression of fibrosis by performing unilateral ureteral obstruction in a renal interstitial fibrosis model in an α -SMA knockout mouse. Furthermore, transfection of the α -SMA gene into α -SMA $^{-/-}$ myofibroblasts from these mice suppressed fibrosis. These results indicated that α -SMA expression suppresses, rather than accelerates, fibrosis. The α -SMA $^{-/-}$ myofibroblasts that were obtained possessed the ability to migrate, proliferate, and overproduced type I collagen. These cells did not express α -SMA, but did start to overexpress skeletal muscle α -actin, smooth muscle γ -actin, and related molecules.

These cells also overproduced type I collagen compared to conventional myofibroblasts expressing α -SMA. Because the cells which were established by transfection of the α -SMA gene into the α -SMA^{-/-} cells failed to produce type I collagen, these cells might resemble smooth muscle cells. It is likely that myofibroblasts are an intermediate type of cells between α -SMA^{-/-} cells and smooth muscle cells, given their ability to express α -SMA and produce type I collagen. Taken together, their report does not refute the generally accepted concept




Fig. 3. Immunofluorescent staining for α -SMA was influenced by the anti-ED-A antibody Aa, the 2D TGF- β ₁ group; b, the 2D anti-ED-A antibody group; c, the 2D IgG group; B, Counts of cells immunostained for α -SMA in the stress fiber formation. Aa and Ac, Immunostaining for α -SMA was shown in the stress fiber formation. Ab, Cells expressing α -SMA in the stress fiber formation were markedly decreased in number. Immunostaining for ED-A was seen in response to addition of the anti-ED-A antibody to the medium. B. Cells with α -SMA expression induced by TGF- β ₁ were decreased in number in the anti-ED-A antibody group. Bar, 10 μ m.




Fig. 4. Morphological alterations of gels in response to TGF- β ₁ according to differences in the pouring methods. A. Ubiquitous pouring of cells. a. The shape of the gel b. the diameters of the gels. B. Spot pouring of cells. Aa. The change in shape induced by TGF- β ₁ is indicated by the circle. Ab. TGF- β ₁ significantly induced gel shrinkage. The data represent the diameters of the gels compared to the well. B. The gel was deformed, as shown by the indentation toward the cell mass (arrows).




Fig. 5. Type I collagen gel contraction. No gel contraction occurred in any of the 1D groups. TGF-β₁ was shown to evoke gel contraction in the 2D TGF-β₁ and IgG groups. TGF-β₁-induced gel contraction was attenuated by adding the anti-ED-A antibody. The data represent the diameters of the gels compared to the well.




Fig. 6. Type I collagen accumulated in the supernatants of the monolayer cultures. The TGF-β₁ and anti-ED-A antibodies do not influence the type I collagen accumulation in 1D groups. TGF-β₁ increased the type I collagen accumulation in the 2D TGF-β₁ and IgG groups. The TGF-β₁- accelerated collagen accumulation was repressed by the anti-ED-A antibody.

that α -SMA contributes to the progression of fibrosis, but do add an additional role for α -SMA producing cells. There was another report (Leemans et al., 2009) that suggested that renal fibrogenesis persists despite decreasing numbers of α -SMA⁺ myofibroblasts. This also does not contradict the general consensus about the role of myofibroblasts, because it can be considered that after myofibroblasts induce the accumulation of collagen fibrils by forming rigid fibrils which are resistant to proteolysis by proteinases, they undergo apoptosis. Subsequently, the α -SMA⁻ fibroblasts might add collagen fibers to these rigid fibers. In this case, myofibroblasts would make up the nucleus of the accumulated collagen fibrils.

In the present study, TGF- β_1 induced the production of fibronectin ED-A, which was anchored to kidney fibroblasts, and subsequently induced α -SMA expression in the stress fiber formations in cells. Stress fiber formation is involved in cell contraction, which results in gel contraction via the consecutive binding of α -SMA, focal adhesion, integrins, and type I collagen. In addition, the α -SMA expression was strongly inhibited by exposing the cells to an anti-ED-A antibody. Therefore, the TGF- β_1 -induced expression of α -SMA was considered to be, at least to a large extent, dependent on ED-A expression. This is compatible with the fact that the fibronectin ED-A isoform contributes to the transdifferentiation of a variety of cells into myofibroblasts with α -SMA (Bochaton-Piallat et al., 2000; French-Constant et al., 1989; Jamagin et al., 1994).

It was shown in the renal fibroblasts that the intracellular signaling pathways of TGF- β_1 -induced fibronectin ED-A expression involves Smad 3 (Bondi et al., 2010). The pathways leading to α -SMA expression and myofibroblast transdifferentiation differed between reports. Our previous report suggested that the pathway of TGF- β_1 -induced α -SMA expression involved p38 MAPK (Ina et al., 2011). This finding is consistent with a study by Miura et al. (Miura et al., 2006), in which retinal pigment epithelial cells were used. On the other hand, Masszi et al. (Masszi et al., 2003) have found that the activation of the Rho kinase pathway was involved in the α -SMA expression in renal tubular epithelial cells. The contrast between these findings might be due to the differences in the cells used. How ED-A expression activated the p38 MAPK pathway in the present study remains to be determined. However, ED-A has been identified as an endogenous ligand for Toll-like receptor-4 (Gondokaryono et al., 2007; Lefebvre et al., 2011; Ricupero et al., 2001) which is well known to be expressed in inflammatory cells such as macrophages, mast cells, and leukocytes, and is associated with innate immunity. Recently, it was demonstrated that Toll-like receptor 4 is also expressed in renal interstitial fibroblasts (Sorensen et al., 2011). Furthermore, the activation of Toll-like receptor 4 was reported to promote renal tubulointerstitial fibrosis (Campbell et al., 2011; Sorensen et al., 2011). It is therefore possible that ED-A binds to Toll-like receptor 4 and activates it, leading to stimulation of the related signaling pathways, eventually promoting the signaling by TGF- β_1 and stimulating α -SMA expression.

Because the immunostaining for fibronectin ED-A was positive in both conventional immunocytochemistry and in that without the first antibody (the anti-ED-A antibody) of the cells exposed to the antibody in culture, the epitope appears to be present in a region other than the receptor-binding site. In this case, the binding of the antibody to the epitope might cause alterations in the conformation of the receptor binding site, and inhibit the binding of ED-A to the receptor. In the current study, since contraction of the NRK49F cell-populated collagen lattice was suppressed by the adding anti-ED-A antibody, the ED-A expression was considered to be involved in TGF- β_1 -induced gel contraction.




Fig. 7. The hypothesis: about the contribution of fibronectin ED-A expression to α-SMA expression and type I collagen accumulation. We hypothesize the mechanism as follows: TGF-β₁ induces ED-A domain expression through the pathway involving Smad 3 in the fibronectin molecule. The fibronectin ED-A is anchored to the cell via a combination of the cell-binding domain and the integrin on the cell. Subsequently, the ED-A domain binds to the receptor (the Toll-like receptor) and stimulates it. Furthermore, the signaling transfers to one of the TGF-β₁-activated pathways, which may be the p38 MAPK pathway. Finally, the ED-A-stimulated pathway induces α-SMA expression and type I collagen accumulation, which involves the increased synthesis of collagen and decreased degradation of collagen via the production of proteinase inhibitors, TIMPs and PAI-1.

The previous study demonstrated that TGF- β_1 caused the contraction of the fibroblast (without α -SMA expression)-populated collagen lattice in culture medium with 10% FBS. The TGF- β_1 -induced gel contraction was accelerated by expressing α -SMA, which was induced by a low concentration of FBS and the addition of TGF- β_1 (Ina et al., 2011). Therefore, TGF- β_1 was suggested to exert distinct functions on α -SMA expression and gel contraction. In addition, it was shown that TGF- β_1 -induced gel contraction occurred via the p38 MAPK pathway and the Rho kinase pathway (Ina et al., 2011). Taken together, since the p38 MARK pathway is involved in both α -SMA expression and gel contraction, it is likely that ED-A expression might stimulate the p38 MAPK pathway activated by TGF- β_1 , followed by myofibroblast transdifferentiation and gel contraction.

These findings still have not clarified what gel contraction means in the fibrosis of diabetic nephropathy. We believe that it corresponds to the renal atrophy following fibrosis. However, it is also possible that gel contraction might reflect the shrinkage of collagen fibrils, allowing collagen molecules to become nearer to each other, followed by increased crosslinking, which leads to the formation of rigid fibrils. These are resistant to proteolysis by proteinases, thus leading to the accumulation of fibrils, and therefore, fibrosis.

In the present study, we also demonstrated that accumulation of type I collagen was, at least in part, dependent on ED-A expression. To our knowledge, this is the first report that ED-A mediates TGF- β_1 -stimulated type I collagen accumulation, while there was a previous report that ED-A stimulated type I collagen mRNA expression (Serini et al., 1998). There are both Smad-dependent and independent pathways, including the p38 MAPK pathway, via which TGF- β_1 can stimulate the production of ECM containing type I collagen (Eickelberg, 2001). The observed α -SMA expression, gel contraction, and type I collagen accumulation are, at least in part, dependent on ED-A expression. The α -SMA expression was the most dependent of these events on ED-A. We therefore hypothesized that α -SMA expression is, to a large extent, dependent on ED-A expression via the p38 MAPK pathway. On the other hand, gel contraction and type I collagen accumulation were, to a lesser extent, dependent on ED-A expression, because they are comparatively mediated by pathways other than the p38 MAPK pathway. ED-A binds to Toll-like receptor-4 and activates it, followed by signaling via diverse pathways in inflammatory cells (Akira & Takeda, 2004). The p38 MAPK pathway is also included in these pathways. Thus, it is most likely that the α -SMA expression, gel contraction, and type I collagen accumulation induced by ED-A expression are mediated by activating the p38 MAPK pathway (Fig. 6). In other words, ED-A appears to enhance the activity of the p38 MARK pathway stimulated by TGF- β_1 .

5. Conclusion

The TGF- β_1 -induced α -SMA expression, followed by gel contraction and stimulated type I collagen accumulation, were, at least in part, dependent on fibronectin ED-A expression. These findings suggest that tubulointerstitial fibrosis associated with diabetic nephropathy and fibrotic tissue contraction leading to renal atrophy are partly mediated by TGF- β_1 -induced fibronectin ED-A expression. In addition, since the fibronectin ED-A isoform is seldom found in normal adult subjects, the negative effects induced by

inhibition of ED-A expression or activity would likely be minimal. Taken together, our findings and those of previous studies suggest that ED-A may be a target for treatment of diabetic nephropathy.

6. Acknowledgement

The authors thank Ms. Yukari Goto for her excellent secretarial assistance.

7. References

- Akira, S., & Takeda, K. (2004). Toll-like receptor signaling. *Nature Reviews*, Vol. 4, pp. 499-511, 1474-1733
- Balza, E.L., Borsi, G., Allemanni, G., & Zardi, L. (1988). Transforming growth factor β regulates the levels of different fibronectin isoforms in normal cultured fibroblasts. *FEBS Letters*, Vol. 228, pp. 42-44, 0014-5793
- Bohle, A., Wehrmann, M., Bogenschütz, O., Batz, C., & Müller, G.A. (1991). The pathogenesis of chronic renal failure in diabetic nephropathy. *Pathology, Research and Practice*, Vol. 187, pp. 251-259, 0344-0338
- Bochaton-Piallat, M.L., Kapetanios, A.D., Donati, G., Redard, M., Gabbiani, G., & Pournaras, C.J. (2000). TGF- β 1, TGF- β receptor II and ED-A fibronectin expression in myofibroblast of vitreoretinopathy. *Investigation Ophthalmology & Visual Science*, Vol. 41, No.8, pp. 2336-2342, 1552-5783
- Bondi, C.D., Manickam, N., Lee, D.Y., Block, K., Gorin, Y., Abboud, H.E., & Barnes, J.L. (2010). NAD(P)H oxidase mediates TGF- β 1-induced activation of kidney myofibroblasts. *Journal of the American Society of Nephrology*, Vol. 21, pp.93-102, 1046-6673
- Borsi, L., Castellani, P., Risso, A.M., Leprini, A., & Zardi, L. (1990). Transforming growth factor β regulates the splicing pattern of fibronectin messenger RNA precursor. *FEBS Letters*, Vol. 261, pp. 175-178, 0014-5793
- Campbell, M.T., Hile, K.L., Zhang, H., Asanuma, H., Vanderbrink, B.A., Rink, R.R., & Meldrum, K.K. (2011). Toll-like receptor 4: a novel signaling pathway during renal fibrogenesis. *Journal of Surgical Research*, Vol. 168, No. 1, pp. e61-e69, 0022-4804
- Edwards, D.R., Murphy, G., Reynolds, J.J., Whitham, S.E., Docherty, A.J.P., Angel, P., & Heath, J.K. (1987). Transforming growth factor beta modulates the expression of collagenase and metalloprotease inhibitor. *EMBO Journal*, Vol. 6, pp. 1899-1904, 0261-4189
- Eickelberg, O. (2001). Endless healing: TGF- β , SMADs, and fibrosis. *FEBS Letters*, Vol. 506, pp. 11-14, 0014-5793
- Ffrench-Constant, C., Van, W.L., Dvorak, H.F., & Hynes, R.O. (1989). Reappearance of an embryonic pattern of fibronectin splicing during wound healing in the adult rat. *The Journal of Cell Biology*, Vol. 109, pp. 903-914, 0021-9525


- Gabbiani, G. & Ryan, G., & Majno, G. (1971). Presence of modified fibroblasts in granulation tissue and their possible in wound contraction. *Experientia*, Vol. 27, pp. 549-550, 0014-4754
- Glukhova, M.A., Frid, M.G., Shekhomin, B.V., Vasilevskaya, T.D., Grunwald, J., Saginati, M., & Koteliensky, V.E. (1989). Expression of extra domain-A fibronectin sequence in vascular smooth muscle cells is phenotype dependent. *The Journal of Cell Biology*, Vol. 109, pp. 357-366, 0021-9525
- Gondokaryono, S.P., Ushio, H., Niyonsaba, F., Hara, M., Takenaka, H., Jayawardana, S.T.M., Ikeda, S., Okumura, K., & Ogawa, H. (2007). The extra domain A of fibronectin stimulates murine mast cells via Toll-like receptor 4. *Journal of Leukocyte Biology*, Vol. 82, pp. 657-665, 1938-3673
- Humphreys, B.D., Lin, S.L., Kobayashi, A., Hudson, T.E., Nowlin, B.T., Bonventre, J.V., Valerius, M.T., McMahon, A.P., & Duffield, J.S. (2010). Fate tracing reveals the pericyte and not epithelial origin of myofibroblasts in kidney fibrosis. *The American Journal of Pathology*, Vol. 176, pp. 85-97, 0002-9440
- Ina, K., Kitamura, H., Tatsukawa, S., & Fujikura, Y. (2011). Significance of α -SMA in myofibroblasts emerging in renal tubulointerstitial fibrosis. *Histology and Histopathology*, Vol. 26, pp. 855-866, 1699-5848
- Iwano, M., Plieth, D., Danoff, T.M., Xue, C., Okeda, H., & Neilson, E.G. (2002). Evidence that fibroblasts derive from epithelium during tissue fibrosis. *The Journal of Clinical Investigation*, Vol. 110, pp. 341-350, 0021-9738
- Jamagin, W.R., Rockey, D.C., Koteliensky, V.E., Wang, S.S., & Bissel, D.M. (1994). Expression of variant fibronectins in wound healing: Cellular source and biological activity of the EIIIA segment in rat hepatic fibrogenesis. *The Journal of Cell Biology*, Vol. 127, pp. 2037-2048, 0021-9525
- Keeley, E.C., Mehrad, B., & Strieter, R.M. (2010). Fibrocytes: bringing new insights into mechanisms of inflammation and fibrosis. *The International Journal of Biochemistry & Cell Biology*, Vol. 42, pp. 535-542, 1357-2725
- Laiho, M., Saksela, O., & Keski-Oja, J. (1987). Transforming growth factor- β induction of type-1 plasminogen activator inhibitor. *Journal of Biological Chemistry*, Vol. 262, pp. 17467-17474, 0021-9258
- Leemans, J.C., Butter, L.M., Pulskens, W.P., Teske, G.J., Claessen, N., V.d.Poll, T., & Florequin, S. (2009). The role of Toll-like receptor 2 in inflammation and fibrosis during progressive renal injury. *PLoS ONE*, Vol. 4, pp. e5704, 1932-6203
- Lefebvre, J.S., Lévesque, T., Picard, S., Paré, G., Gravel, A., Flamand, L., & Borgeat, P. (2011). Extra domain A of fibronectin primes leukotriene biosynthesis and stimulates neutrophil migration through activation of Toll-like receptor 4. *Arthritis & Rheumatism*, Vol. 63, No.6, pp. 1527-1533, 1529-0131
- Magnusson, M.K., & Mosher, D.F. (1998). Fibronectin: Structure, assembly, and cardiovascular implications. *Arteriosclerosis, Thrombosis, and Vascular Biology*, Vol. 18, pp. 1363-1370, 0276-5047
- Masszi, A., Ciano, C.D., Sirokmány, G., Arthur, W.T., Rotstein, O.D., Wang, J., McCulloch, C.A.G., Rosivall, L., Mucsi, I., & Kapus, A. (2003). Central role for Rho in TGF- β 1-induced α -smooth muscle actin expression during epithelial-mesenchymal

- transition. *American Journal of Physiology-Renal Physiology*, Vol. 284, No. 5, pp. F911-F924, 1522-1466
- Miura, M., Hata, H., Hirayama, K., Kita, T., Noda, Y., Fujisawa, K., Shimokawa, H., & Ishibashi, T. (2006). Critical role of the Rho-kinase pathway in TGF- β 2-dependent collagen gel contraction by retinal pigment epithelial cells. *Experimental Eye Research*, Vol. 82, pp. 849-859, 0014-4835
- Norton, P.A., & Hynes, R.O. (1987). Alternative splicing of chicken fibronectin in embryos and in normal and transformed cells. *Molecular and Cellular Biology*, Vol. 7, pp. 4297-4307
- Petersen, T.E., Thogersen, H.C., Skorstengaard, K., Vibe-Pedersen, K., Sahi, P., Sottrup-Jensen, L., & Magnusson, S. (1983). Partial primary structure of bovine plasma fibronectin: Three types of internal homology. *Proceedings of the National Academy of Sciences of the United States of America*, Vol. 80, pp. 137-141, 0027-8424
- Ricupero, D.A., Poliks, C.F., Rishikof, D.C., Kuang, P-P., & Goldstein, R.H. (2001). Apigenin decreases expression of the myofibroblast phenotype. *FEBS Letters*, Vol. 506, pp. 15-21, I0014-5793
- Serini, G., Bochaton-Piallat, M.L., Ropraz, P., Geinoz, A., Borsi, L., Zardi, L., & Gabbiani, G. (1998). The fibronectin domain ED-A is crucial for myofibroblastic phenotype induction by transforming growth factor- β 1. *The Journal of Cell Biology*, Vol. 142, No.3, pp. 873-881, 0021-9525
- Simonson, M.S. (2007). Phenotypic transitions and fibrosis in diabetic nephropathy. *Kidney International*, Vol. 71, pp. 846-854, 0085-2538
- Sörensen, I., Susnik, N., Inhester, T., Degen, J.L., Melk, A., Haller, H., & Schmitt, R (2011). Fibrinogen, acting as a mitogen for tubulointerstitial fibroblasts, promotes renal fibrosis. *Kidney International*, [epub ahead of print], 0085-2539
- Strutz, F., & Zeisberg, M. (2006). Renal fibroblasts and myofibroblasts in chronic kidney disease. *The Journal of American Society of Nephrology*, Vol. 17, pp. 2992-2998, 1046-6673
- Takeji, M., Moriyama, T., Oseto, S., Kawada, N., Hori, M., Imai, E., & Miwa, T. (2006). Smooth muscle α -actin deficiency in myofibroblasts leads to enhanced renal tissue fibrosis. *Journal of Biological Chemistry*, Vol. 281, pp. 40193-40200, 1083-351X
- Tomasek, J.J., Gabbiani, G., Hinz, B., Chaponnier, C., & Brown, R.A. (2002). Myofibroblasts and mechano-regulation of connective tissue remodeling. *National Reviews Molecular Cell Biology*, Vol. 3, No. 5, pp. 349-363, 1471-0072
- Vartio, T., Laitinen, L., Narvanen, O., Cutolo, M., Thornell, L.E., Zardi, L., & Virtanen, I. (1987). Differential expression of the ED sequence-containing form of cellular fibronectin in embryonic and adult human tissues. *Journal of Cell Science*, Vol. 88, pp. 419-430, 1477-9137
- Wierzbicka-Patynowski, I., & Schwarzbauer, J.E. (2003). The ins and outs of fibronectin matrix assembly. *Journal of cell Science*, Vol. 116, pp. 3269-3276, 1477-9137

Zeisberg, E.M., Potenta, S.E., Sugimono, H., Zeisberg, M., & Kalluri, R. (2008). Fibroblasts in kidney fibrosis emerge via endothelial-to-mesenchymal transition. *The Journal of American Society of Nephrology*, Vol. 19, pp. 2282-2287, 1046-6673

IntechOpen

IntechOpen



Diabetic Nephropathy

Edited by Dr. John Chan

ISBN 978-953-51-0543-5

Hard cover, 166 pages

Publisher InTech

Published online 20, April, 2012

Published in print edition April, 2012

Internationally renowned experts have provided data on their own studies, and discuss the relative usefulness of their work in relation to diabetic nephropathy. The first section describes the novel role of intrarenal renin-angiotensin-aldosterone system (RAAS) and oxidative stress in the development of diabetic nephropathy and discusses the current and novel pharmacological interventions in the treatment of diabetic nephropathy. The second section discusses other important contributors outside of the RAAS in the pathogenesis of diabetic nephropathy including AGE/RAGE, epithelial-mesenchymal-transition (EMT) and immune cytokines. Features: Provides novel information on various pathophysiological determinants in the development of diabetic nephropathy Provides novel information on various pharmacological interventions of diabetic nephropathy

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Keisuke Ina, Hirokazu Kitamura, Shuji Tatsukawa and Yoshihisa Fujikura (2012). The Contribution of Fibronectin ED-A Expression to Myofibroblast Transdifferentiation in Diabetic Renal Fibrosis, Diabetic Nephropathy, Dr. John Chan (Ed.), ISBN: 978-953-51-0543-5, InTech, Available from: <http://www.intechopen.com/books/diabetic-nephropathy/contribution-of-fn-ed-a-expression-to-myofibroblast-transdifferentiation-in-diabetic-renal-fibrosis->

INTech
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](https://creativecommons.org/licenses/by/3.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen