

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

185,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Congenital Myasthenic Syndromes – Molecular Bases of Congenital Defects of Proteins at the Neuromuscular Junction

Kinji Ohno¹, Mikako Ito¹ and Andrew G. Engel²

¹*Division of Neurogenetics, Center for Neurological Diseases and Cancer,
Nagoya University Graduate School of Medicine, Nagoya,*

²*Department of Neurology, Mayo Clinic, Rochester, Minnesota,*

¹*Japan*

²*USA*

1. Introduction

Congenital myasthenic syndromes (CMS) are heterogeneous disorders caused by mutations in molecules expressed at the neuromuscular junction (NMJ) (Fig. 1). Each mutation affects the expression level or the functional properties or both of the mutant molecule. No fewer than 11 defective molecules at the NMJ have been identified to date. The mutant molecules include (i) acetylcholine receptor (AChR) subunits that forms nicotinic AChR and generate endplate potentials (Ohno *et al.*, 1995; Sine *et al.*, 1995), (ii) rapsyn that anchors and clusters AChRs at the endplate (Ohno *et al.*, 2002; Milone *et al.*, 2009), (iii) agrin that is released from nerve terminal and induces AChR clustering by stimulating the downstream LRP4/MuSK/Dok-7/rapsyn/AChR pathway (Huze *et al.*, 2009), (iv) muscle-specific receptor tyrosine kinase (MuSK) that transmits the AChR-clustering signal from agrin/LRP4 to Dok-7/rapsyn/AChR (Chevessier *et al.*, 2004; Chevessier *et al.*, 2008), (v) Dok-7 that interacts with MuSK and exerts the AChR-clustering activity (Beeson *et al.*, 2006; Hamuro *et al.*, 2008), (vi) plectin that is an intermediate filament-associate protein concentrated at sites of mechanical stress (Banwell *et al.*, 1999; Selcen *et al.*, 2011), (vii) glutamine-fructose-6-phosphate aminotransferase 1 encoded by *GFPT1*, the function of which at the NMJ has not been elucidated (Senderek *et al.*, 2011), (viii) skeletal muscle sodium channel type 1.4 (Na_v1.4) that spreads depolarization potential from endplate throughout muscle fibers (Tsujino *et al.*, 2003), (ix) collagen Q that anchors acetylcholinesterase (AChE) to the synaptic basal lamina (Ohno *et al.*, 1998; Ohno *et al.*, 1999; Kimbell *et al.*, 2004), (x) β2-laminin that forms a cruciform heterotrimeric lamins-221, -421, and -521 and links extracellular matrix molecules to the β-dystroglycan at the NMJ (Maselli *et al.*, 2009), (xi) choline acetyltransferase (ChAT) that resynthesizes acetylcholine from recycled choline at the nerve terminal (Ohno *et al.*, 2001). AChR (Lang & Vincent, 2009), MuSK (Hoch *et al.*, 2001; Cole *et al.*, 2008), and LRP4 (Higuchi *et al.*, 2011) are also targets of myasthenia gravis, in which autoantibody against each molecule impairs the neuromuscular transmission.

CMS are classified into three groups of postsynaptic, synaptic, and presynaptic depending on the localization of the defective molecules. Among the eleven molecules introduced

above, AChR, rapsyn, MuSK, Dok-7, plectin, and Nav1.4 are associated with the postsynaptic membrane. Agrin, ColQ, and β 2-laminin reside in the synaptic basal lamina. The only presynaptic disease protein identified to date is choline acetyltransferase (ChAT). A target molecule and its synaptic localization of glutamine-fructose-6-phosphate aminotransferase 1 (GFPT1) are still unresolved but the phenotypic consequence is the postsynaptic AChR deficiency. This chapter focuses on molecular bases of these three groups of CMS.




Fig. 1. Schematic of molecules expressed at the NMJ

2. Physiology of the NMJ

This section introduces molecular basis of development and maintenance of the NMJ, and physiological features of nicotinic muscle AChR.

2.1 NMJ synaptogenesis

At the NMJ, MuSK is an indirect receptor for agrin (Valenzuela *et al.*, 1995; Dechiara *et al.*, 1996). Agrin released from the nerve terminal binds to LRP4 on the postsynaptic membrane (Kim *et al.*, 2008; Zhang *et al.*, 2008). Binding of LRP4 to agrin phosphorylates MuSK. Phosphorylated MuSK recruits the noncatalytic adaptor protein Dok-7 (Okada *et al.*, 2006). Once recruited, Dok-7 further facilitates phosphorylation of MuSK, and induces clustering of rapsyn and AChR by phosphorylating the β subunit of AChR. Rapsyn self-associates and makes a homomeric cluster at the endplate, which serves as a scaffold for AChR. Rapsyn and AChR bind each other with a stoichiometry of 1:1. Rapsyn also binds to β -dystroglycan and links the rapsyn scaffold to the subsynaptic cytoskeleton (Froehner *et al.*, 1990; Cartaud *et al.*, 1998; Ramarao & Cohen, 1998; Ramarao *et al.*, 2001). Except for LRP4, each of the above molecules is a CMS target.

2.2 Physiology of the nicotinic muscle AChR

Nicotinic AChRs are pentameric ligand-gated ion channels. The family of pentameric ligand-gated ion channels includes cationic AChRs, cationic serotonergic receptors (5HT₃), anionic glycine receptors, and anionic GABA_A and GABA_C receptors (Keramidas *et al.*, 2004). Heteromeric neuronal nicotinic AChRs are comprised of various combinations of α ($\alpha 2$ - $\alpha 7$) and β subunits ($\beta 2$ - $\beta 4$), whereas homomeric AChRs are formed only by a single α subunit (e.g., $\alpha 7$ - $\alpha 9$) (Mihailescu & Drucker-Colin, 2000). On the other hand, nicotinic muscle AChRs have only two forms: fetal AChR that carries the α , β , δ , and γ subunits encoded by *CHRNA1*, *CHRNA1*, *CHRNA1*, *CHRNA1*, respectively, in the stoichiometry $\alpha_2\beta\delta\gamma$; and adult-type AChR that carries the ϵ subunit instead of the γ subunit in the stoichiometry $\alpha_2\beta\delta\epsilon$ (Mishina *et al.*, 1986). The ϵ subunit is encoded by *CHRNE*. Nicotinic muscle AChR harbors two binding sites for ACh at the interfaces between the α - δ and α - γ/α - ϵ subunits (Lee *et al.*, 2009; Mukhtasimova *et al.*, 2009). Binding of a single ACh molecule opens the channel pore but for a short time. Binding of two ACh molecules stabilizes the open state of AChR, and AChR stays open for a longer time. Only cations pass through the channel pore of nicotinic AChRs. Unlike sodium, potassium, or calcium channels, AChRs, in general, have no selectivity for cations, but $\alpha 7$ AChRs have 10-20 times higher permeability for Ca²⁺ than for Na⁺.

3. Postsynaptic CMS

Postsynaptic CMS is classified into four phenotypes: (i) endplate AChR deficiency due to defects in AChR, rapsyn, agrin, MuSK, Dok-7, plectin, glutamine-fructose-6-phosphate aminotransferase 1, (ii) slow-channel congenital myasthenic syndrome, (iii) fast-channel congenital myasthenic syndrome, and (iv) sodium channel myasthenia.

3.1 Endplate AChR deficiency

Endplate AChR deficiency is caused by defects in AChR, rapsyn, agrin, MuSK, Dok-7, plectin, and GFPT1.

3.1.1 Endplate AChR deficiency due to defects in AChR subunits

Endplate AChRs deficiency can arise from mutations in *CHRNA1*, *CHRNA1*, *CHRNA1*, and *CHRNE*, but not *CHRNA1*.

Two different groups of mutations of the AChR subunit genes cause endplate AChR deficiency. The first group includes null mutations in *CHRNE* encoding the ϵ subunit. The null mutations are caused by frameshifting DNA rearrangements, *de novo* creation of a stop codon, and frameshifting splice-site mutations, or mutations involving residues essential for subunit assembly. Large-scale in-frame DNA rearrangements also abolish expression of the AChR ϵ subunit (Abicht *et al.*, 2002). Mutations in the promoter region (Ohno *et al.*, 1999) and most missense mutations (Ohno *et al.*, 1997) do not completely abolish expression of the ϵ subunit but the molecular consequences are indistinguishable from those of null mutations. Lack of the ϵ subunit can be compensated for by the presence of the fetal γ subunit that is normally expressed in embryos (Engel *et al.*, 1996). The patients can survive with γ -AChR even in the absence of ϵ -AChR. If a null mutation resides in the other AChR

subunit genes, the affected individual will have no substituting subunit and cannot survive. Indeed, two homozygous missense low expressor or null mutations in *CHRNA1* and *CHRND* caused lethal fetal akinesia (Michalk *et al.*, 2008).

The second group of mutations affecting the AChR subunit genes includes missense mutations of *CHRNA1*, *CHRNA1*, and *CHRND*. These mutations compromise expression of the mutant subunit and/or the assembly of AChRs, but do not completely abolish AChRs expression. The main difference between mutations in *CHRNE* and those in *CHRNA1*, *CHRNA1*, and *CHRND* is tolerance to low or no expression of the ϵ subunit whereas similar mutations in other subunits generally have devastating consequences and cause high fatality. Some missense mutations in *CHRNA1*, *CHRNA1*, *CHRND*, and *CHRNE* also affect the AChR channel kinetics and vice versa. The kinetic effects will predominate if the second mutation is a low expressor, or if the kinetic mutation has slow-channel features with dominant gain-of function effects.

In endplate AChR deficiency, the postsynaptic membrane displays a reduced binding for peroxidase- or ^{125}I -labeled α -bungarotoxin and the synaptic response to ACh, reflected by the amplitude of the miniature endplate potential, endplate potential, and endplate current, is reduced. In some but not all cases the postsynaptic region is simplified. In most cases, the muscle fibers display an increased number of small synaptic contacts over an extended length of the muscle fiber. In some patients quantal release is higher than normal. In patients with null mutations in *CHRNE*, single channel recordings of AChRs at patient endplates reveal prolonged opening bursts that open to an amplitude of 60 pS, indicating expression of the fetal γ -AChR in contrast to the adult ϵ -AChR that has shorter opening bursts and opens to an amplitude of 80 pS. In contrast, in most patients with low-expressor mutations in the *CHRNA1*, *CHRNA1*, or *CHRND*, single channel recordings demonstrate no or minor kinetic abnormalities.

As in autoimmune myasthenia gravis, endplate AChR deficiency is generally well controlled by regular doses of anticholinesterases. Anticholinesterase medications inhibit the catalytic activity of AChE; this prolongs the dwell time of ACh in the synaptic space and allows each ACh molecule to bind repeatedly to AChR.

3.1.2 Endplate AChR deficiency due to defects in rapsyn

Congenital defects of rapsyn also cause endplate AChR deficiency. Rapsyn makes a homomeric cluster and binds to AChR as well as to β -dystroglycan, and forms AChR clusters at the endplate (Froehner *et al.*, 1990; Cartaud *et al.*, 1998; Ramarao & Cohen, 1998; Ramarao *et al.*, 2001). The structural domains of rapsyn include an N-terminal myristoylation signal required for membrane association (Ramarao & Cohen, 1998), seven tetratricopeptide repeats at codons 6 to 279 that subserve rapsyn self-association (Ramarao & Cohen, 1998; Ramarao *et al.*, 2001), a coiled-coil domain at codons 298 to 331 that binds to the long cytoplasmic loop of each AChR subunit (Bartoli *et al.*, 2001), a Cys-rich RING-H2 domain at codons 363-402 that binds to the cytoplasmic domain of β -dystroglycan (Bartoli *et al.*, 2001) and mediates the MuSK induced phosphorylation of AChR (Lee *et al.*, 2008), and a serine phosphorylation site at codon 406. Transcription of rapsyn in muscle is under the control of helix-loop-helix myogenic determination factors that bind to the *cis*-acting E-box sequence in the *RAPSN* promoter (Ohno *et al.*, 2003).

Loss-of-function mutations in *RAPSN* have been reported in the coding region (Ohno *et al.*, 2002; Burke *et al.*, 2003; Dunne & Maselli, 2003; Maselli *et al.*, 2003; Muller *et al.*, 2003; Banwell *et al.*, 2004; Yasaki *et al.*, 2004; Cossins *et al.*, 2006; Muller *et al.*, 2006) as well as in the promoter region (Ohno *et al.*, 2003). N88K in *RAPSN* is one of the most frequently observed mutations in CMS (Muller *et al.*, 2003; Richard *et al.*, 2003). We reported lack of a founder haplotype for N88K (Ohno & Engel, 2004), but analysis of markers closer to *RAPSN* later revealed possible presence of a shared haplotype (Muller *et al.*, 2004) suggesting that N88K is an ancient founder mutation but subsequent multiple recombination events and divergence of microsatellite markers have narrowed the shared haplotype region. Functional analysis L14P, N88K, and 553ins5 disclosed that these mutations have no effect on self-association of rapsyn but impair colocalization of rapsyn with AChR (Ohno *et al.*, 2002). Analysis of A25V, N88K, R91L, L361R, and K373del later revealed diverse molecular defects affecting colocalization of rapsyn with AChR, formation of agrin-induced AChR clusters, self-association of rapsyn, and expression of rapsyn (Cossins *et al.*, 2006). Although there are no genotype-phenotype correlations in mutations at the coding region, arthrogryposis at birth and other congenital malformations occurs in nearly a third of the patients. In addition, the -38A>G mutation affecting an E-box in the promoter region observed in Near-Eastern Jewish patients exhibits unique facial malformations associated with prognathism and malocclusion (Ohno *et al.*, 2003).

Most patients respond well to anticholinesterase medications. Some patients further improve with addition of 3,4-diaminopyridine, ephedrine, and albuterol (Banwell *et al.*, 2004). The drug 3,4-diaminopyridine blocks the presynaptic potassium channel, which slows the repolarization of the presynaptic membrane (Wirtz *et al.*, 2010) enhancing the influx of Ca^{2+} through the presynaptic voltage-gated P/Q-type and N-type channels. This, in turn, facilitates the exocytosis of synaptic vesicles and the quantal content of the endplate potential.

3.1.3 Endplate AChR deficiency due to a defect in agrin

Neural agrin released from the nerve terminal is a key mediator of synaptogenesis at the NMJ. A reported homozygous G1709R agrin mutation, however, did not cause AChR deficiency but mutations in agrin are potential causes of AChR deficiency by interfering with the activation of MuSK and by impeding synaptic maturation.

The patient harboring the G1709R mutation was a 42-year-old woman with right lid ptosis since birth, no oculoparesis, and mild weakness of facial, hip-girdle and anterior tibial muscles, and refractoriness to pyridostigmine or 3,4-diaminopyridine (Huze *et al.*, 2009). The mutation is in the laminin G-like 2 domain, upstream of the neuron-specific y and z exons that are required for MuSK activation and AChR clustering. AChR and agrin expression at the endplate were normal. Structural studies showed endplates with misshaped synaptic gutters partially filled by nerve endings and formation of new endplate regions. The postsynaptic regions were preserved. Expression studies in myotubes using a mini-agrin construct revealed the mutation did not affect MuSK activation or agrin binding to α -dystroglycan. Forced expression of the mutant mini-agrin gene in mouse soleus muscle induced changes similar to those at patient endplates. Thus, the observed mutation perturbs the maintenance of the endplate without altering the canonical function of agrin to induce development of the postsynaptic compartment.

3.1.4 Endplate AChR deficiency due to defects in MuSK

MuSK and LRP4 form a heteromeric receptor for agrin. Five *MUSK* mutations have been reported in three papers. The first report describes heteroallelic frameshift (220insC) and missense (V790M) mutations in a patient with respiratory distress in early life, mild ptosis, decreased upward gaze, and fatigable weakness of the cervical and proximal more than distal muscles. The symptoms were worsened by pregnancy. Treatment with pyridostigmine and 3,4-diaminopyridine was ineffective (Chevessier *et al.*, 2004). The frameshift mutation prevents MuSK expression and the missense mutation decreases MuSK expression and impairs its interaction with Dok-7. Forced expression of the mutant protein in mouse muscle decreased AChR expression at the endplate and caused aberrant axonal outgrowth (Chevessier *et al.*, 2004). Interestingly, mice homozygous for MuSK V789M (which corresponds to the human MuSK V790M) are normal but mice hemizygous for V789M are severely affected suggesting that MuSK V790M in humans is a haploinsufficient only when accompanied by a null mutation (Chevessier *et al.*, 2008).

A second report describes heteroallelic M605I and A727V mutations in MuSK in a patient with severe myasthenic symptoms since early life that improved after puberty but worsened after menstrual periods. The MEPP and MEPC amplitudes in anconeus muscle were reduced to about 30% of normal and the EPP quantal content was half-normal. Synaptic contacts were small and electron microscopy showed simplified postsynaptic regions with too few secondary synaptic clefts. The patient failed to respond to pyridostigmine, ephedrine or 3,4-diaminopyridine but responded partially to albuterol (Maselli *et al.*, 2010).

A third report describes a homozygous P31L mutation in the extracellular domain of MuSK in 5 patients in a consanguineous Sudanese kinship. The findings included ptosis from an early age, partial ophthalmoparesis, and weakness of torso and limb girdle muscles. Pyridostigmine therapy gave only slight benefit (Mihaylova *et al.*, 2009).

3.1.5 Endplate AChR deficiency due to defects in Dok-7

Phosphorylated MuSK recruits a noncatalytic adaptor protein, Dok-7. Recruited Dok-7 further facilitates phosphorylation of MuSK (Okada *et al.*, 2006). Dok-7 is highly expressed at the postsynaptic region of skeletal muscle and in heart. It harbors an N terminal pleckstrin homology domain (PH) important for membrane association, a phosphotyrosine-binding (PTB) domain, and C-terminal sites for phosphorylation. The PH and PTB domains are required for association with and phosphorylation of MuSK. Phosphorylation of two C terminal residues is a requisite for Dok-7 activation by Crk and Crk-L (Hallock *et al.*, 2010).

Numerous mutations have been identified in *DOK7* (Beeson *et al.*, 2006; Muller *et al.*, 2007; Anderson *et al.*, 2008; Selcen *et al.*, 2008; Vogt *et al.*, 2009; Ben Ammar *et al.*, 2010). Nearly all patients carry a common 1124_1127dupTGCC mutation in exon 7. This and other mutations upstream of the C-terminal phosphorylation sites abrogate the ability of Dok-7 to associate with Crk1/Crk1L and hence its activation (Hallock *et al.*, 2010; Wu *et al.*, 2010). Mutations disrupting or eliminating the PH and PTB domains of Dok-7 prevent dimerization and association of Dok-7 with MuSK (Bergamin *et al.*, 2010).

3.1.6 Endplate AChR deficiency due to defects in plectin

Plectin, encoded by *PLEC*, is a highly conserved and ubiquitously expressed intermediate filament-linking protein concentrated at sites of mechanical stress, such as the postsynaptic membrane of the endplate, the sarcolemma, Z-disks in skeletal muscle, hemidesmosomes in skin, and intercalated disks in cardiac muscle. Pathogenic mutations in *PLEC* result in epidermolysis bullosa simplex, a progressive myopathy (Smith *et al.*, 1996), and, in some patients, myasthenic syndrome (Banwell *et al.*, 1999; Selcen *et al.*, 2011). We reported two cases of CMS associated with plectin deficiency (Banwell *et al.*, 1999; Selcen *et al.*, 2011). The dystrophic changes in muscle are attributed to dislocation of the fiber organelles no longer anchored by the cytoskeletal intermediate filaments and to sarcolemmal defects allowing Ca^{2+} ingress into the muscle fibers. The myasthenic syndrome is attributed to destruction of the junctional folds lacking adequate cytoskeletal support.

3.1.7 Endplate AChR deficiency due to defects in glutamine-fructose-6-phosphate aminotransferase 1 (GFPT1)

Glutamine-fructose-6-phosphate transaminase 1, encoded by *GFPT1*, catalyzes transfer of an amino group from glutamine onto fructose-6-phosphate, yielding glucosamine-6-phosphate and glutamate. GFPT1 is a rate-limiting enzyme that controls the flux of glucose into the hexosamine biosynthesis pathway. GFPT1 thus initiates formation of UDP-N-acetylglucosamine (UDP-GlcNAc), which is a source of multiple glycosylation processes including addition of N-acetylglucosamine to serine or threonine residues (O-linked GlcNAc) (Wells *et al.*, 2001). The disease gene was discovered by linkage analysis and homozygosity mapping of 13 kinships with a limb-girdle CMS often associated with tubular aggregates in skeletal muscle (Senderek *et al.*, 2011). Immunoblots of muscle of affected patients revealed decreased expression of O-linked GlcNAc, but the responsible molecule(s) causing CMS remain elusive.

3.2 Slow-channel congenital myasthenic syndrome (SCCMS)

The second class of postsynaptic CMS due to mutations in the AChR subunit genes is SCCMS. SCCMS is an autosomal dominant disorder, in which a gain-of-function mutation on a single allele compromises the neuromuscular signal transduction (Ohno *et al.*, 1995). The mutation causes prolonged AChR channel openings and increases the synaptic response to ACh (Fig. 2). There is a single reported case of autosomal recessive SCCMS, in which an ϵL78P mutation minimally prolongs channel opening events but the mutant channel arising from a single allele is not sufficient to cause disease (Croxen *et al.*, 2002). In general, dominantly inherited disorders, including SCCMS, tend to present after adolescence and have a relatively mild course. Some patients with SCCMS, however, present early in life and become severely disabled even in the first decade.

In SCCMS, neuromuscular transmission is compromised by three distinct mechanisms. First, staircase summation of endplate potentials causes depolarization block of the postsynaptic membrane by rendering the voltage-gated skeletal muscle sodium channel go into an inactivated state and thereby inhibit action potential generation (Maselli & Soliven, 1991). Second, some mutant AChRs are prone to become desensitized (Milone *et al.*, 1997), which reduces the number of AChRs that respond to the released ACh quanta. Third,

prolonged opening of AChR causes excessive influx of extracellular calcium, which results in focal degeneration of the junctional folds as well as apoptosis of some of the junctional nuclei (Groshong *et al.*, 2007). In normal adult human ϵ -AChR, 7% of the synaptic current is carried by Ca^{2+} , which is higher than that carried by the human fetal γ -AChR or by muscle AChRs of other species (Fucile *et al.*, 2006). This predisposes endplate to Ca^{2+} overloading when the channel opening events are prolonged. In addition, at least two SCCMS mutations, ϵ T264P (Ohno *et al.*, 1995) and α V259F (Fidzianska *et al.*, 2005), increase the Ca^{2+} permeability 1.5- and 2-fold, respectively (Di Castro *et al.*, 2007).




Fig. 2. Slow channel CMS. (A) Schematic diagram of AChR subunits with SCCMS mutations. (B) Single channel currents from wild-type and slow channel (α V249F) AChRs expressed on HEK293 cells. (C) Miniature endplate current (MEPC) recorded from endplates of a control and a patient harboring α V249F. The patient's MEPC decays biexponentially (arrows) due to expression of both wild-type and mutant AChRs.

Slow channel mutations can be divided into two groups. The first group includes mutations at the extracellular domain like α G153S (Sine *et al.*, 1995), as well as at the N-terminal part of the first transmembrane domain like α N217K (Wang *et al.*, 1997) and ϵ L221F (Hatton *et al.*, 2003). These mutations increase the affinity for ACh binding, probably by retarding the dissociation of ACh from the binding site, which gives rise to repeated channel openings after a single event of ACh binding. The second group includes mutations at the second transmembrane domain (M2) that lines the ion channel pore. These mutations mostly introduce a bulky amino acid into the channel lining face, but ϵ T264P (Ohno *et al.*, 1995) introduces a kink into the channel pore, whereas β V266A (Shen *et al.*, 2003) and ϵ V265A (Ohno *et al.*, 1998) introduce a smaller amino acid into the pore. Mutations in M2 retard the channel closing rate α and variably enhance the channel opening rate β . Some mutations in M2 also increase affinity for ACh, which include α V249F (Milone *et al.*, 1997), ϵ L269F (Engel *et al.*, 1996), and ϵ T264P (Ohno *et al.*, 1995).

SCCMS can be treated with conventional doses of long-lived open channel blockers of AChR, such as the antiarrhythmic agent quinidine (Fukudome *et al.*, 1998; Harper & Engel, 1998) and the antidepressant fluoxetine (Harper *et al.*, 2003). Quinidine reduces the prolonged burst duration of SCCMS to the normal level at 5 μ M (Fukudome *et al.*, 1998). As the concentration of quinidine in the treatment of cardiac arrhythmia is 6-15 μ M, 5 μ M is readily attainable in clinical practice and indeed demonstrates significant effects (Harper & Engel, 1998). Similarly, fluoxetine reduces the prolonged burst duration to the normal level at 10 μ M, which is clinically attainable without adverse effects at 80 to 120 mg/day of fluoxetine (Harper *et al.*, 2003).

3.3 Fast-channel congenital myasthenic syndrome (FCCMS)

The third class of postsynaptic CMS due to mutations in AChR subunit genes is FCCMS. FCCMS is kinetically opposite to SCCMS (Fig. 3). In FCCMS, the closed state of AChR is stabilized compared to the open state which results in abnormally brief channel opening events which, in turn, reduces the amplitude of the endplate potential and impair the safety margin of neuromuscular transmission. The resulting pathophysiology is thus similar to endplate AChR deficiency, but abnormally small endplate potential is a qualitative instead of a quantitative defect in AChR.

FCCMS is an autosomal recessive disorder. One allele carries a missense mutation that confers a fast closure of AChRs, and the other allele usually harbors a low-expressor or null mutation, or the fast channel mutation occurs at homozygosity. As in heterozygous healthy parents of endplate AChR deficiency, we humans may completely lack 50% of each AChR subunit without any clinical symptoms. In FCCMS, a low-expressor or null mutation on one allele unmasks the deleterious effect of the fast-channel mutation on the second allele. Detailed kinetic analyses of FCCMS mutations have revealed special insights into the molecular architectures of the AChR subunits. Three such examples are presented here.

The ϵ 1254ins18 mutation causes a duplication of STRDQE codons at positions 413 to 418 close to the C-terminal end of the long cytoplasmic loop (LCP) linking the third (M3) and fourth (M4) transmembrane domains of the receptor. ϵ 1254ins18-AChR expressed on HEK293 cells opens in three different modes. The opening probabilities of normal AChRs are clustered into a single large peak, whereas the ϵ 1254ins18-AChR shows three different peaks (Milone *et al.*, 1998). In all the three modes, the AChR is activated slowly and inactivated rapidly, which gives rise to an inefficient synaptic response to ACh. Another FCCMS mutation, ϵ A411P in the LCP also destabilizes the channel opening kinetics. The channel opening probabilities of ϵ A411P-AChRs are widely distributed and do not form any discernible peaks (Wang *et al.*, 2000). Our analysis first disclosed that the function of LCP is to stabilize the open conformation of the AChR.

ϵ N436del is a deletion of Asn at the C-terminal end of the LCP. The deletion shortens the LCP and shifts a negatively charged Asp residue at codon 435 against M4. ϵ N436del-AChR decreases the duration of channel opening bursts 2.7-fold compared to the wild type due to a 2.3-fold decrease in gating efficiency and a 2.5-fold decrease in agonist affinity of the diliganded closed state. A series of artificial mutations established that the effects of ϵ N436del are not due to juxtaposition of a negative charge against M4 but to the shortening of the LCP. Deletion of the C-terminal residue of the LCP of the β and δ subunits also results in fast-

channel kinetics, but that in the α subunit dictates slow-channel kinetics. Thus, the LCPs of four AChR subunits contribute in an asymmetric manner to optimize the activation of AChRs through allosteric links to the channel and to the agonist binding sites (Shen *et al.*, 2005).

The mutation α V285I introduces a bulky amino acid into the M3 transmembrane domain and causes FCCMS (Fig. 3). Kinetic studies demonstrate that the mutation slows the channel opening rate β and speeds the channel closing rate α , resulting in a 15.1-fold reduction in the channel gating equilibrium constant θ ($= \beta/\alpha$). On the other hand, the mutation minimally affects affinity for ACh. The probability of channel openings decreased when we introduced Leu, a bulky amino acid, at position V285, but rather increased when we introduced smaller amino acids such as Thr and Ala. We observed similar effects when we introduced similar substitutions into the β , δ , and ϵ subunits. Thus, introduction of bulky amino acids narrows the channel pore, while introduction of smaller amino acids widens the channel pore. Our analysis thus revealed that the M3 domain backs up the channel-lining pore lined by the M2 transmembrane domains and has stereochemical effects on channel gating kinetics (Wang *et al.*, 1999).

FCCMS can be effectively treated with anticholinesterases and 3,4-diaminopyridine. The pharmacologic effects of these drugs were discussed in the section of endplate AChR deficiency (Section 3.1.2).




Fig. 3. Fast channel CMS. (A) Schematic diagram of AChR subunits with FCCMS mutations. (B) Single channel currents from wild-type and fast channel (α V285I) AChRs expressed on HEK293 cells. (C) Miniature endplate current (MEPC) recorded from endplates of a control and a patient harboring α V285I. The patient's MEPC decays faster than that of the normal control.

3.4 CMS due to defects in skeletal muscle sodium channel, Nav1.4

Another class of postsynaptic CMS is due to mutations in skeletal muscle sodium channel, Nav1.4, encoded by *SCN4A* (Tsujino *et al.*, 2003). Dominant gain-of-function mutations in this gene cause hyperkalemic periodic paralysis (Ptacek *et al.*, 1991), paramyotonia congenita (McClatchey *et al.*, 1992; Ptacek *et al.*, 1992), potassium-aggravated myotonia (Lerche *et al.*, 1993), and hypokalemic periodic paralysis type 2 (Bulman *et al.*, 1999). On the other hand, loss-of-function mutations cause a CMS.

Failure of normal-amplitude endplate potential depolarizing the resting potential to -40 mV in intercostal muscle of a CMS patient with episodes of apnea and myasthenic symptoms since birth prompted us to search for mutations in *SCN4A*. We identified two heteroallelic missense mutations, S246L and V1442E (Tsujino *et al.*, 2003). Activation kinetics of the mutant Nav1.4 was normal for both S246L and V1442E, but the fast inactivation curves were shifted to hyperpolarization by 7.3 mV for S246L and 33.2 mV for V1442E, indicating that both mutations enhance fast inactivation of the Nav1.4 immediately after it is activated. Moreover, a high proportion of the V1442 channel was in the inactivated state even at a normal resting membrane potential. Recovery from the fast-inactivated state was slowed for both mutations. This was in contrast to gain-of-function mutations in other diseases, which shift the fast inactivation curves to depolarization. Neither S246L nor V1442E affected slow inactivation. Analysis of use-dependent inactivation in HEK293 cells by stimulating at 50 Hz for 3 ms revealed that wild-type and S246L channels decreased the peak current only by 5% and V1442E channel decreased it by 30% during the first few pulses and suggested that the S246L mutation is relatively benign.

4. Synaptic CMS

Defects in three components of the synaptic basal lamina, AChE, $\beta 2$ laminin and neural agrin, are associated with CMS. The CMS caused by mutations in agrin was discussed above under the postsynaptic CMS (Section 3.1.3) because the site of action of agrin is the LRP4/MuSK complex at the endplate.

4.1 Endplate AChE deficiency due to defects in collagen Q

Three tetramers of catalytic AChE subunits are linked by a triple helical collagen Q (ColQ) to constitute an asymmetric ColQ-tailed AChE (Krejci *et al.*, 1997). ColQ carries three domains (i) an N-terminal proline-rich attachment domain (PRAD) that organizes the catalytic AChE subunits into a tetramer, (ii) a collagenic domain that forms a triple helix, and (iii) a C-terminal domain enriched in charged residues and cysteines. ColQ-tailed AChE is organized in the secretory pathway, excreted, and anchored into the synaptic basal lamina using two domains of ColQ (Fig. 4). First, the collagen domain harbors two heparan sulfate proteoglycan (HSPG) binding domains (Deprez *et al.*, 2003) that bind to HSPG, such as perlecan (Peng *et al.*, 1999). Second, the C-terminal domain binds to MuSK (Cartaud *et al.*, 2004).

Endplate AChE deficiency is caused by congenital defects of ColQ (Donger *et al.*, 1998; Ohno *et al.*, 1998; Ohno *et al.*, 2000). Congenital defects of ColQ cause endplate AChE deficiency. No mutations have been detected in a gene encoding the catalytic subunit of AChE in CMS

or in any other disease. There are three classes of ColQ mutations. First, mutations in the proline-rich attachment domain (PRAD) hinder binding of ColQ to AChE. Sedimentation analysis of AChE species of the patient muscle and transfected cells shows complete lack of ColQ-tailed AChE. Second, mutations in the collagen domain, most of which are truncation mutations, hinder formation of triple helix of ColQ. Sedimentation analysis of muscle and transfected cells demonstrate a truncated single-stranded ColQ associated with a homotetramer of AChE. Third, the mutations in the C-terminal domain have no deleterious effect on formation of the asymmetric ColQ-tailed AChE, but they compromise anchoring of ColQ-tailed AChE to the synaptic basal lamina as elegantly shown in vitro overlay binding of mutant and wild-type human recombinant ColQ-tailed AChE to the frog endplate (Kimbell *et al.*, 2004).




Fig. 4. ColQ anchors to the synaptic basal lamina by binding to perlecan and MuSK.

EMG studies show a decremental response as in other CMS. In addition, most patients have a repetitive CMAP response on a single nerve stimulus. The repetitive CMAP decrements faster than the primary CMAP. It can be overlooked unless a well rested muscle is tested by single nerve stimuli. The prolonged dwell time of unhydrolyzed ACh in the synaptic space prolongs the endplate potential; when this exceeds the absolute refractory period of the muscle fiber action potential, it elicits a repetitive CMAP. As mentioned above, a repetitive CMAP also occurs in slow channel syndrome.

Some aspects of the pathophysiology of endplate AChE deficiency resemble those of the SCCMS. As in the SCCMS, neuromuscular transmission is compromised by three distinct mechanisms. First, staircase summation of endplate potentials causes a depolarization block, which inactivates a proportion the voltage-gated skeletal sodium channel, $\text{Nav}1.4$. (Maselli & Soliven, 1991). Second, prolonged exposure of AChR to ACh during physiologic activity desensitizes a fraction of the available AChRs (Milone *et al.*, 1997). Third, repeated openings of AChR cause calcium overloading to the endplate, which culminates in an endplate myopathy (Groshong *et al.*, 2007). Unlike in the SCCMS, the nerve terminals are abnormally small and often encased by Schwann cells. This decreases the quantal content and hence the amplitude of the endplate potential (Engel *et al.*, 1977).

Anticholinesterase medications have no effect on neuromuscular transmission and can cause excessive muscarinic side effects. Quinidine (Fukudome *et al.*, 1997; Harper & Engel, 1997) and fluoxetine (Harper *et al.*, 2003), which shorten the open duration of the AChR channel and benefit the slow-channel syndrome, can increase muscle weakness. A respirator dependent infant with severe endplate AChE deficiency was improved by intermittent blockade of AChR by atracurium, an agent that protects AChR from overexposure to ACh (Brenningstall *et al.*, 1996). Ephedrine sulfate at a dose of 150 to 200 mg per day in adults is effective for myasthenic symptoms (Bestue-Cardiel *et al.*, 2005; Mihaylova *et al.*, 2008). Although high concentrations of ephedrine are able to block AChR openings (Milone & Engel, 1996), molecular bases of ephedrine effects in clinical practice remain elusive. As an alternative to ephedrine, albuterol sulfate 8 to 16 mg per day also shows benefit (Liewluck *et al.*, in press).

4.2 CMS due to a defect in $\beta 2$ laminin

Laminins are cruciform heterotrimeric glycoproteins composed of α , β , and γ chains and are assembled from products of five α , four β , and three γ genes. The laminin molecules are named according to their chain composition. For example, laminin-321 contains $\alpha 3$, $\beta 2$, and $\gamma 1$ chains (Aumailley *et al.*, 2005). Three laminins are present at the synaptic basal lamina, laminin-221, laminin-421, and laminin-521. Each contains the $\beta 2$ subunit. Laminin-421 is restricted to the primary synaptic cleft and promotes the precise alignment of pre- and postsynaptic specializations. Laminin-521 lines the primary and secondary clefts, promotes presynaptic differentiation, and prevents Schwann cells from entering the synaptic cleft. The synaptic laminins provide a stop signal for axons at developing endplates and organize presynaptic differentiation (Sanes, 1997). Mice deficient for *Lamb2* that encodes $\beta 2$ laminin show reduced terminal branching of presynaptic motor axons, with a decreased number of active zones, no clustering of the synaptic vesicles above the active zones, and extension of Schwann cell processes into the primary synaptic cleft, and decreased spontaneous and evoked quantal release (Noakes *et al.*, 1995; Patton *et al.*, 1998). In addition to its presence at the endplate, $\beta 2$ -laminin is also highly expressed in renal glomeruli and the eye. *LAMB2* mutations in humans cause Pierson syndrome characterized by ocular malformation including small non-reactive pupils, loss of accommodation, and abnormalities of the lens, cornea and retina and by fatal nephrotic syndrome that requires renal transplantation (Zenker *et al.*, 2004).

Maselli and coworkers reported a 20-year-old woman with Pierson syndrome caused by two heteroallelic frameshifting mutations (1478delG and 4804delC) in *LAMB2* who also had a severe CMS (Maselli *et al.*, 2009). The nephrotic syndrome was corrected by a renal transplant at age 15 months. The patient had respiratory distress in infancy, delayed motor milestones, a decremental EMG response, limited ocular ductions, bilateral ptosis, severe proximal limb weakness, scoliosis, and required assisted ventilation at night and sometimes during the day. AChE activity was spared at the NMJ. Electron microscopy of the NMJ showed small axon terminal size and encasement of nerve endings by the Schwann cell, widening of the primary synaptic clefts with invasion of the synaptic space by processes of Schwann cells, moderate simplification of postsynaptic membranes, and decreased number of synaptic vesicles. Both morphological and microelectrode studies were similar to those observed in *Lamb2*-mice (Noakes *et al.*, 1995). Notably, symptoms were worsened by pyridostigmine but were improved by ephedrine.

5. Presynaptic CMS

Choline acetyltransferase (ChAT) is the only presynaptic molecule that is known to be defective in CMS.

5.1 CMS with episodic apnea due to defects in choline acetyltransferase (ChAT)

ACh released from the nerve terminal is hydrolyzed into choline and acetate by AChE at the synaptic basal lamina. Choline is taken up by the nerve terminal by a high-affinity choline transporter on the presynaptic membrane (Apparsundaram *et al.*, 2000; Okuda *et al.*, 2000). ChAT resynthesizes ACh from choline and acetyl-CoA (Oda *et al.*, 1992). After the synaptic vesicles are acidified by the vesicular proton pump (Reimer *et al.*, 1998), the resynthesized cationic ACh is packed into a synaptic vesicle by the vesicular ACh transporter (vAChT) in exchange for protons (Erickson *et al.*, 1994).




Fig. 5. Choline acetyltransferase (ChAT). (A) Genomic structure of *CHAT* and identified mutations. A gene for vesicular acetylcholine transporter (vAChT) is in the first intron of *CHAT*. (B) Kinetics of wild-type and mutant ChAT enzymes. ChAT synthesizes acetylcholine using choline and acetyl-CoA. L210P abrogates an affinity of ChAT for acetyl-CoA (AcCoA), and R560H abolishes an affinity of ChAT for choline.

We determined the complete genomic structure of *CHAT* encoding ChAT, and identified ten mutations in five CMS patients with the characteristic clinical features of sudden episodes of apnea associated with variable myasthenic symptoms (Ohno *et al.*, 2001). Additional *CHAT* mutations were later reported by other groups (Maselli *et al.*, 2003; Schmidt *et al.*, 2003; Barisic *et al.*, 2005; Mallory *et al.*, 2009; Yeung *et al.*, 2009; Schara *et al.*, 2010). All of our patients showed a marked decrease of the endplate potential after subtetanic stimulation that recovered slowly over 5 to 10 min, which pointed to a defect in the resynthesis or

vesicular packaging of ACh at the nerve terminal. Kinetic studies of mutant ChAT enzymes disclosed variable decreases in affinity for choline and/or acetyl-CoA, as well as variable reduction the catalytic rate (Ohno *et al.*, 2001) (Fig. 5). Moreover, some recombinant mutants expressed at a reduced level in COS cells. Two patients carried a functionally null mutation on one allele, but ChAT encoded on the other allele was partially functional. Heterozygous parents that carried the null allele were asymptomatic indicating that humans can tolerate up to but not exceeding 50% reduction of presynaptic ChAT activity. None of our patients has autonomic symptoms or signs of central nervous system involvement other than that attributed to anoxic episodes. This suggests that the ChAT activity and/or substrate availability are rate limiting for ACh synthesis at the motor nerve but not at other cholinergic synapses. Indeed, stimulated quantal release at the endplate is higher than at other cholinergic synapses, which points to selective vulnerability of the NMJ to reduced ACh resynthesis. Crystal structure of ChAT resolved at 2.2 Å revealed that some of the reported *CHAT* mutations in CMS patients are not at the substrate-binding or the catalytic site of ChAT. Hence these mutation exert their effect by an allosteric mechanism or render the enzyme structurally unstable (Cai *et al.*, 2004).

In most patients, anticholinesterase medications are of benefit in ameliorating the myasthenic symptoms and preventing the apneic crises but few patients fail to respond to cholinergic therapy remaining permanently paralyzed and remain respirator dependent. Prophylactic anticholinesterase therapy is advocated even for patients asymptomatic between crises. Parents of affected children must be indoctrinated to anticipate sudden worsening of the weakness and possible apnea with febrile illnesses, excitement, or overexertion. Long-term nocturnal apnea monitoring is indicated in any patient in whom ChAT deficiency is proven or suspected (Byring *et al.*, 2002).

6. Conclusions

We reviewed the clinical and molecular consequences of defects in 11 genes associated with CMS. Molecular studies of CMS began with identification of a missense mutation in the AChR ϵ subunit in a SCCM patient (Ohno *et al.*, 1995). Since then, mutations in seven postsynaptic, three synaptic, and one presynaptic proteins have been discovered. In some CMS the disease gene has been elusive and await discovery. Resequencing analysis with the next generation sequencers may speed this effort.

7. Acknowledgments

Works in our laboratories were supported by Grants-in-Aid from the MEXT and the MHLW of Japan to K.O., and by NIH Grant NS6277 and a research Grant from Muscular Dystrophy Association to A.G.E.

8. References

- Abicht, A., Stucka, R., Schmidt, C., Briguet, A., Höpfner, S., Song, I.-H., Pongratz, D., Müller-Felber, W., Ruegg, M. A. & Lochmüller, H. (2002). A newly identified chromosomal microdeletion and an N-box mutation of the AChR gene cause a congenital myasthenic syndrome. *Brain*, Vol. 125, No., pp. 1005-1013, ISSN 0006-8950

- Anderson, J. A., Ng, J. J., Bowe, C., McDonald, C., Richman, D. P., Wollmann, R. L. & Maselli, R. A. (2008). Variable phenotypes associated with mutations in DOK7. *Muscle and Nerve*, Vol. 37, No. 4, pp. 448-456, ISSN 0148-639X
- Apparsundaram, S., Ferguson, S. M., George, A. L., Jr. & Blakely, R. D. (2000). Molecular cloning of a human, hemicholinium-3-sensitive choline transporter. *Biochemical and Biophysical Research Communications*, Vol. 276, No. 3, pp. 862-867, ISSN 0006-291X
- Aumailley, M., Bruckner-Tuderman, L., Carter, W. G., Deutzmann, R., Edgar, D., Ekblom, P., Engel, J., Engvall, E., Hohenester, E., Jones, J. C., Kleinman, H. K., Marinkovich, M. P., Martin, G. R., Mayer, U., Meneguzzi, G., Miner, J. H., Miyazaki, K., Patarroyo, M., Paulsson, M., Quaranta, V., Sanes, J. R., Sasaki, T., Sekiguchi, K., Sorokin, L. M., Talts, J. F., Tryggvason, K., Uitto, J., Virtanen, I., von der Mark, K., Wewer, U. M., Yamada, Y. & Yurchenco, P. D. (2005). A simplified laminin nomenclature. *Matrix Biology*, Vol. 24, No. 5, pp. 326-332, ISSN 0945-053X
- Banwell, B. L., Russel, J., Fukudome, T., Shen, X. M., Stilling, G. & Engel, A. G. (1999). Myopathy, myasthenic syndrome, and epidermolysis bullosa simplex due to plectin deficiency. *Journal of Neuropathology and Experimental Neurology*, Vol. 58, No. 8, pp. 832-846, ISSN 0022-3069
- Banwell, B. L., Ohno, K., Sieb, J. P. & Engel, A. G. (2004). Novel truncating RAPSN mutations causing congenital myasthenic syndrome responsive to 3,4-diaminopyridine. *Neuromuscular Disorders*, Vol. 14, No. 3, pp. 202-207, ISSN 0960-8966
- Barisic, N., Muller, J. S., Paucic-Kirincic, E., Gazdik, M., Lah-Tomulic, K., Pertl, A., Sertic, J., Zurak, N., Lochmuller, H. & Abicht, A. (2005). Clinical variability of CMS-EA (congenital myasthenic syndrome with episodic apnea) due to identical CHAT mutations in two infants. *Eur J Paediatr Neurol*, Vol. 9, No. 1, pp. 7-12, ISSN 1570-9639
- Bartoli, M., Ramarao, M. K. & Cohen, J. B. (2001). Interactions of the rapsyn RING-H2 domain with dystroglycan. *Journal of Biological Chemistry*, Vol. 276, No. 27, pp. 24911-24917, ISSN 0021-9258
- Beeson, D., Higuchi, O., Palace, J., Cossins, J., Spearman, H., Maxwell, S., Newsom-Davis, J., Burke, G., Fawcett, P., Motomura, M., Muller, J. S., Lochmuller, H., Slater, C., Vincent, A. & Yamanashi, Y. (2006). Dok-7 mutations underlie a neuromuscular junction synaptopathy. *Science*, Vol. 313, No. 5795, pp. 1975-1978, ISSN 0036-8075
- Ben Ammar, A., Petit, F., Alexandri, N., Gaudon, K., Bauche, S., Rouche, A., Gras, D., Fournier, E., Koenig, J., Stojkovic, T., Lacour, A., Petiot, P., Zagnoli, F., Viollet, L., Pellegrini, N., Orlikowski, D., Lazaro, L., Ferrer, X., Stoltenburg, G., Paturneau-Jouas, M., Hentati, F., Fardeau, M., Sternberg, D., Hantai, D., Richard, P. & Eymard, B. (2010). Phenotype genotype analysis in 15 patients presenting a congenital myasthenic syndrome due to mutations in DOK7. *J Neurol*, Vol. 257, No. 5, pp. 754-766, ISSN 0340-5354
- Bergamin, E., Hallock, P. T., Burden, S. J. & Hubbard, S. R. (2010). The cytoplasmic adaptor protein Dok7 activates the receptor tyrosine kinase MuSK via dimerization. *Molecular Cell*, Vol. 39, No. 1, pp. 100-109, ISSN 1097-2765
- Bestue-Cardiel, M., Saenz de Cabezón-Alvarez, A., Capablo-Liesa, J. L., Lopez-Pison, J., Pena-Segura, J. L., Martín-Martínez, J. & Engel, A. G. (2005). Congenital endplate

- acetylcholinesterase deficiency responsive to ephedrine. *Neurology*, Vol. 65, No. 1, pp. 144-146, ISSN 0028-3878
- Brenningstall, G. N., Kurachek, S. C., Fugate, J. H. & Engel, A. G. (1996). Treatment of congenital endplate acetylcholinesterase deficiency by neuromuscular blockade. *Journal of Child Neurology*, Vol. 11, No. 4, pp. 345-346, ISSN 0883-0738
- Bulman, D. E., Scoggan, K. A., van Oene, M. D., Nicolle, M. W., Hahn, A. F., Tollar, L. L. & Ebers, G. C. (1999). A novel sodium channel mutation in a family with hypokalemic periodic paralysis. *Neurology*, Vol. 53, No. 9, pp. 1932-1936, ISSN 0028-3878
- Burke, G., Cossins, J., Maxwell, S., Owens, G., Vincent, A., Robb, S., Nicolle, M., Hilton-Jones, D., Newsom-Davis, J., Palace, J. & Beeson, D. (2003). Rapsyn mutations in hereditary myasthenia: Distinct early- and late-onset phenotypes. *Neurology*, Vol. 61, No., pp. 826-828, ISSN 0028-3878
- Byring, R. F., Pihko, H., Tsujino, A., Shen, X.-M., Gustafsson, B., Hackman, P., Ohno, K., Engel, A. G. & Udd, B. (2002). Congenital myasthenic syndrome associated with episodic apnea and sudden infant death. *Neuromuscular Disorders*, Vol. 12, No. 6, pp. 548-553, ISSN 0960-8966
- Cai, Y., Cronin, C. N., Engel, A. G., Ohno, K., Hersh, L. B. & Rodgers, D. W. (2004). Choline acetyltransferase structure reveals distribution of mutations that cause motor disorders. *EMBO Journal*, Vol. 23, No. 10, pp. 2047-2058, ISSN 0261-4189
- Cartaud, A., Coutant, S., Petrucci, T. C. & Cartaud, J. (1998). Evidence for in situ and in vitro association between beta-dystroglycan and the subsynaptic 43k rapsyn protein - consequence for acetylcholine receptor clustering at the synapse. *Journal of Biological Chemistry*, Vol. 273, No. 18, pp. 11321-11326, ISSN 0021-9258
- Cartaud, A., Stochlic, L., Guerra, M., Blanchard, B., Lambergeon, M., Krejci, E., Cartaud, J. & Legay, C. (2004). MuSK is required for anchoring acetylcholinesterase at the neuromuscular junction. *Journal of Cell Biology*, Vol. 165, No. 4, pp. 505-515, ISSN 0021-9525
- Chevessier, F., Faraut, B., Ravel-Chapuis, A., Richard, P., Gaudon, K., Bauche, S., Prioleau, C., Herbst, R., Goillot, E., Ioos, C., Azulay, J. P., Attarian, S., Leroy, J. P., Fournier, E., Legay, C., Schaeffer, L., Koenig, J., Fardeau, M., Eymard, B., Pouget, J. & Hantai, D. (2004). MUSK, a new target for mutations causing congenital myasthenic syndrome. *Human Molecular Genetics*, Vol. 13, No. 24, pp. 3229-3240, ISSN 0964-6906
- Chevessier, F., Girard, E., Molgo, J., Bartling, S., Koenig, J., Hantai, D. & Witzemann, V. (2008). A mouse model for congenital myasthenic syndrome due to MuSK mutations reveals defects in structure and function of neuromuscular junctions. *Human Molecular Genetics*, Vol. 17, No. 22, pp. 3577-3595, ISSN 0964-6906
- Cole, R. N., Reddel, S. W., Gervasio, O. L. & Phillips, W. D. (2008). Anti-MuSK patient antibodies disrupt the mouse neuromuscular junction. *Annals of Neurology*, Vol. 63, No. 6, pp. 782-789, ISSN 0364-5134
- Cossins, J., Burke, G., Maxwell, S., Spearman, H., Man, S., Kuks, J., Vincent, A., Palace, J., Fuhrer, C. & Beeson, D. (2006). Diverse molecular mechanisms involved in AChR deficiency due to rapsyn mutations. *Brain*, Vol. 129, No. Pt 10, pp. 2773-2783, ISSN 0006-8950
- Croxen, R., Hatton, C., Shelley, C., Brydson, M., Chauplannaz, G., Oosterhuis, H., Vincent, A., Newsom-Davis, J., Colquhoun, D. & Beeson, D. (2002). Recessive inheritance

- and variable penetrance of slow-channel congenital myasthenic syndromes. *Neurology*, Vol. 59, No. 2, pp. 162-168, ISSN 0028-3878
- Dechiara, T. M., Bowen, D. C., Valenzuela, D. M., Simmons, M. V., Poueymirou, W. T., Thomas, S., Kinetz, E., Compton, D. L., Rojas, E., Park, J. S., Smith, C., Distefano, P. S., Glass, D. J., Burden, S. J. & Yancopoulos, G. D. (1996). The receptor tyrosine kinase MuSK is required for neuromuscular junction formation in vivo. *Cell*, Vol. 85, No. 4, pp. 501-512, ISSN 0092-8674
- Deprez, P., Inestrosa, N. C. & Krejci, E. (2003). Two different heparin-binding domains in the triple-helical domain of ColQ, the collagen tail subunit of synaptic acetylcholinesterase. *Journal of Biological Chemistry*, Vol. 278, No. 26, pp. 23233-23242, ISSN 0021-9258
- Di Castro, A., Martinello, K., Grassi, F., Eusebi, F. & Engel, A. G. (2007). Pathogenic point mutations in a transmembrane domain of the epsilon subunit increase the Ca²⁺ permeability of the human endplate ACh receptor. *J Physiol*, Vol. 579, No. Pt 3, pp. 671-677, ISSN 0022-3751
- Donger, C., Krejci, E., Pou Serradell, A., Eymard, B., Bon, S., Nicole, S., Chateau, D., Gary, F., Fardeau, M., J., M. & Guicheney, P. (1998). Mutation in the human acetylcholinesterase-associated collagen gene, *COLQ*, is responsible for congenital myasthenic syndrome with end-plate acetylcholinesterase deficiency (Type Ic). *Am J Hum Genet*, Vol. 63, No., pp. 967-975, ISSN 0002-9297
- Dunne, V. & Maselli, R. A. (2003). Identification of pathogenic mutations in the human rapsyn gene. *Journal of Human Genetics*, Vol. 48, No. 4, pp. 204-207, ISSN 1434-5161
- Engel, A. G., Lambert, E. H. & Gomez, M. R. (1977). A new myasthenic syndrome with end-plate acetylcholinesterase deficiency, small nerve terminals, and reduced acetylcholine release. *Annals of Neurology*, Vol. 1, No. 4, pp. 315-330, ISSN 0364-5134
- Engel, A. G., Ohno, K., Bouzat, C., Sine, S. M. & Griggs, R. C. (1996). End-plate acetylcholine receptor deficiency due to nonsense mutations in the epsilon subunit. *Annals of Neurology*, Vol. 40, No. 5, pp. 810-817, ISSN 0364-5134
- Engel, A. G., Ohno, K., Milone, M., Wang, H. L., Nakano, S., Bouzat, C., Pruitt, J. N., 2nd, Hutchinson, D. O., Brengman, J. M., Bren, N., Sieb, J. P. & Sine, S. M. (1996). New mutations in acetylcholine receptor subunit genes reveal heterogeneity in the slow-channel congenital myasthenic syndrome. *Human Molecular Genetics*, Vol. 5, No. 9, pp. 1217-1227, ISSN 0964-6906
- Erickson, J. D., Varoqui, H., Schafer, M. K., Modi, W., Diebler, M. F., Weihe, E., Rand, J., Eiden, L. E., Bonner, T. I. & Usdin, T. B. (1994). Functional identification of a vesicular acetylcholine transporter and its expression from a "cholinergic" gene locus. *Journal of Biological Chemistry*, Vol. 269, No. 35, pp. 21929-21932, ISSN 0021-9258
- Fidzianska, A., Ryniewicz, B., Shen, X. M. & Engel, A. G. (2005). IBM-type inclusions in a patient with slow-channel syndrome caused by a mutation in the AChR epsilon subunit. *Neuromuscular Disorders*, Vol. 15, No. 11, pp. 753-759, ISSN 0960-8966
- Froehner, S. C., Luetje, C. W., Scotland, P. B. & Patrick, J. (1990). The postsynaptic 43K protein clusters muscle nicotinic acetylcholine receptors in *Xenopus* oocytes. *Neuron*, Vol. 5, No. 4, pp. 403-410, ISSN 0896-6273

- Fucile, S., Sucapane, A., Grassi, F., Eusebi, F. & Engel, A. G. (2006). The human adult subtype ACh receptor channel has high Ca^{2+} permeability and predisposes to endplate Ca^{2+} overloading. *J Physiol*, Vol. 573, No. Pt 1, pp. 35-43, ISSN 0022-3751
- Fukudome, T., Ohno, K., Brengman, J. M. & Engel, A. G. (1997). Quinidine sulfate normalizes the open duration of slow channel congenital myasthenic syndrome acetylcholine receptor channels expressed in human embryonic kidney cells. *Neurology*, Vol. 48, No., pp. A72, ISSN 0028-3878
- Fukudome, T., Ohno, K., Brengman, J. M. & Engel, A. G. (1998). Quinidine normalizes the open duration of slow-channel mutants of the acetylcholine receptor. *Neuroreport*, Vol. 9, No. 8, pp. 1907-1911, ISSN 0959-4965
- Groshong, J. S., Spencer, M. J., Bhattacharyya, B. J., Kudryashova, E., Vohra, B. P., Zayas, R., Wollmann, R. L., Miller, R. J. & Gomez, C. M. (2007). Calpain activation impairs neuromuscular transmission in a mouse model of the slow-channel myasthenic syndrome. *Journal of Clinical Investigation*, Vol. 117, No. 10, pp. 2903-2912, ISSN 0021-9738
- Hallock, P. T., Xu, C. F., Park, T. J., Neubert, T. A., Curran, T. & Burden, S. J. (2010). Dok-7 regulates neuromuscular synapse formation by recruiting Crk and Crk-L. *Genes & Development*, Vol. 24, No. 21, pp. 2451-2461, ISSN 0890-9369
- Hamuro, J., Higuchi, O., Okada, K., Ueno, M., Iemura, S., Natsume, T., Spearman, H., Beeson, D. & Yamanashi, Y. (2008). Mutations causing DOK7 congenital myasthenia ablate functional motifs in Dok-7. *Journal of Biological Chemistry*, Vol. 283, No. 9, pp. 5518-5524, ISSN 0021-9258
- Harper, C. M. & Engel, A. G. (1997). Quinidine sulfate in the treatment of the slow channel congenital myasthenic syndrome. *Neurology*, Vol. 48, No., pp. A72, ISSN 0028-3878
- Harper, C. M. & Engel, A. G. (1998). Quinidine sulfate therapy for the slow-channel congenital myasthenic syndrome. *Annals of Neurology*, Vol. 43, No. 4, pp. 480-484, ISSN 0364-5134
- Harper, C. M., Fukudome, T. & Engel, A. G. (2003). Treatment of slow-channel congenital myasthenic syndrome with fluoxetine. *Neurology*, Vol. 60, No. 10, pp. 1710-1713, ISSN 0028-3878
- Hatton, C. J., Shelley, C., Brydson, M., Beeson, D. & Colquhoun, D. (2003). Properties of the human muscle nicotinic receptor, and of the slow-channel myasthenic syndrome mutant epsilonL221F, inferred from maximum likelihood fits. *J Physiol*, Vol. 547, No. Pt 3, pp. 729-760, ISSN 0022-3751
- Higuchi, O., Hamuro, J., Motomura, M. & Yamanashi, Y. (2011). Autoantibodies to low-density lipoprotein receptor-related protein 4 in myasthenia gravis. *Annals of Neurology*, Vol. 69, No. 2, pp. 418-422, ISSN 0364-5134
- Hoch, W., McConville, J., Helms, S., Newsom-Davis, J., Melms, A. & Vincent, A. (2001). Auto-antibodies to the receptor tyrosine kinase MuSK in patients with myasthenia gravis without acetylcholine receptor antibodies. *Nature Medicine*, Vol. 7, No. 3, pp. 365-368, ISSN 1078-8956
- Huze, C., Bauche, S., Richard, P., Chevessier, F., Goillot, E., Gaudon, K., Ben Ammar, A., Chaboud, A., Grosjean, I., Lecuyer, H. A., Bernard, V., Rouche, A., Alexandri, N., Kuntzer, T., Fardeau, M., Fournier, E., Brancaccio, A., Ruegg, M. A., Koenig, J., Eymard, B., Schaeffer, L. & Hantai, D. (2009). Identification of an agrin mutation

- that causes congenital myasthenia and affects synapse function. *American Journal of Human Genetics*, Vol. 85, No. 2, pp. 155-167, ISSN 0002-9297
- Keramidas, A., Moorhouse, A. J., Schofield, P. R. & Barry, P. H. (2004). Ligand-gated ion channels: mechanisms underlying ion selectivity. *Progress in Biophysics and Molecular Biology*, Vol. 86, No. 2, pp. 161-204, ISSN 0079-6107
- Kim, N., Stiegler, A. L., Cameron, T. O., Hallock, P. T., Gomez, A. M., Huang, J. H., Hubbard, S. R., Dustin, M. L. & Burden, S. J. (2008). Lrp4 Is a Receptor for Agrin and Forms a Complex with MuSK. *Cell*, Vol. 135, No. 2, pp. 334-342, ISSN 0092-8674
- Kimbell, L. M., Ohno, K., Engel, A. G. & Rotundo, R. L. (2004). C-terminal and heparin-binding domains of collagenic tail subunit are both essential for anchoring acetylcholinesterase at the synapse. *Journal of Biological Chemistry*, Vol. 279, No. 12, pp. 10997-11005, ISSN 0021-9258
- Krejci, E., Thomine, S., Boschetti, N., Legay, C., Sketelj, J. & Massoulié, J. (1997). The mammalian gene of acetylcholinesterase-associated collagen. *Journal of Biological Chemistry*, Vol. 272, No. 36, pp. 22840-22847, ISSN 0021-9258
- Lang, B. & Vincent, A. (2009). Autoimmune disorders of the neuromuscular junction. *Curr Opin Pharmacol*, Vol. 9, No. 3, pp. 336-340, ISSN 1471-4892
- Lee, W. Y., Free, C. R. & Sine, S. M. (2009). Binding to gating transduction in nicotinic receptors: Cys-loop energetically couples to pre-M1 and M2-M3 regions. *Journal of Neuroscience*, Vol. 29, No. 10, pp. 3189-3199, ISSN 0270-6474
- Lee, Y., Rudell, J., Yechikhov, S., Taylor, R., Swope, S. & Ferns, M. (2008). Rapsyn carboxyl terminal domains mediate muscle specific kinase-induced phosphorylation of the muscle acetylcholine receptor. *Neuroscience*, Vol. 153, No. 4, pp. 997-1007, ISSN 0306-4522
- Lerche, H., Heine, R., Pika, U., George, A. L., Jr., Mitrovic, N., Browatzki, M., Weiss, T., Rivet-Bastide, M., Franke, C., Lomonaco, M. & et al. (1993). Human sodium channel myotonia: slowed channel inactivation due to substitutions for a glycine within the III-IV linker. *J Physiol*, Vol. 470, No., pp. 13-22, ISSN 0022-3751
- Liewluck, T., Selcen, D. & Engel, A. G. (in press). Beneficial effects of albuterol in congenital endplate acetylcholinesterase deficiency and DOK-7 myasthenia. *Muscle and Nerve*, Vol., No., pp., ISSN 0148-639X
- Mallory, L. A., Shaw, J. G., Burgess, S. L., Estrella, E., Nurko, S., Burpee, T. M., Agus, M. S., Darras, B. T., Kunkel, L. M. & Kang, P. B. (2009). Congenital myasthenic syndrome with episodic apnea. *Pediatric Neurology*, Vol. 41, No. 1, pp. 42-45, ISSN 0887-8994
- Maselli, R. A. & Soliven, B. C. (1991). Analysis of the organophosphate-induced electromyographic response to repetitive nerve stimulation: paradoxical response to edrophonium and D-tubocurarine. *Muscle & Nerve*, Vol. 14, No. 12, pp. 1182-1188, ISSN 0148-639X
- Maselli, R. A., Chen, D., Mo, D., Bowe, C., Fenton, G. & Wollmann, R. L. (2003). Choline acetyltransferase mutations in myasthenic syndrome due to deficient acetylcholine resynthesis. *Muscle and Nerve*, Vol. 27, No. 2, pp. 180-187, ISSN 0148-639X
- Maselli, R. A., Dunne, V., Pascual-Pascual, S. I., Bowe, C., Agius, M., Frank, R. & Wollmann, R. L. (2003). Rapsyn mutations in myasthenic syndrome due to impaired receptor clustering. *Muscle and Nerve*, Vol. 28, No. 3, pp. 293-301, ISSN 0148-639X
- Maselli, R. A., Ng, J. J., Anderson, J. A., Cagney, O., Arredondo, J., Williams, C., Wessel, H. B., Abdel-Hamid, H. & Wollmann, R. L. (2009). Mutations in *LAMB2* causing a

- severe form of synaptic congenital myasthenic syndrome. *Journal of Medical Genetics*, Vol. 46, No. 3, pp. 203-208, ISSN 0022-2593
- Maselli, R. A., Arredondo, J., Cagney, O., Ng, J. J., Anderson, J. A., Williams, C., Gerke, B. J., Soliven, B. & Wollmann, R. L. (2010). Mutations in MUSK causing congenital myasthenic syndrome impair MuSK-Dok-7 interaction. *Human Molecular Genetics*, Vol., No., pp., ISSN 0964-6906
- McClatchey, A. I., McKenna-Yasek, D., Cros, D., Worthen, H. G., Kuncl, R. W., DeSilva, S. M., Cornblath, D. R., Gusella, J. F. & Brown, R. H., Jr. (1992). Novel mutations in families with unusual and variable disorders of the skeletal muscle sodium channel. *Nature Genetics*, Vol. 2, No. 2, pp. 148-152, ISSN 1061-4036
- Michalk, A., Stricker, S., Becker, J., Rupps, R., Pantzar, T., Miertus, J., Botta, G., Naretto, V. G., Janetzki, C., Yaqoob, N., Ott, C. E., Seelow, D., Wieczorek, D., Fiebig, B., Wirth, B., Hoopmann, M., Walther, M., Korber, F., Blankenburg, M., Mundlos, S., Heller, R. & Hoffmann, K. (2008). Acetylcholine receptor pathway mutations explain various fetal akinesia deformation sequence disorders. *American Journal of Human Genetics*, Vol. 82, No. 2, pp. 464-476, ISSN 0002-9297
- Mihailescu, S. & Drucker-Colin, R. (2000). Nicotine, brain nicotinic receptors, and neuropsychiatric disorders. *Archives of Medical Research*, Vol. 31, No. 2, pp. 131-144, ISSN 0188-4409
- Mihaylova, V., Muller, J. S., Vilchez, J. J., Salih, M. A., Kabiraj, M. M., D'Amico, A., Bertini, E., Wolfle, J., Schreiner, F., Kurlemann, G., Rasic, V. M., Siskova, D., Colomer, J., Herczegfalvi, A., Fabriciova, K., Weschke, B., Scola, R., Hoellen, F., Schara, U., Abicht, A. & Lochmuller, H. (2008). Clinical and molecular genetic findings in COLQ-mutant congenital myasthenic syndromes. *Brain*, Vol. 131, No. Pt 3, pp. 747-759, ISSN 0006-8950
- Mihaylova, V., Salih, M. A., Mukhtar, M. M., Abuzeid, H. A., El-Sadig, S. M., von der Hagen, M., Huebner, A., Nurnberg, G., Abicht, A., Muller, J. S., Lochmuller, H. & Guergueltcheva, V. (2009). Refinement of the clinical phenotype in musk-related congenital myasthenic syndromes. *Neurology*, Vol. 73, No. 22, pp. 1926-1928, ISSN 0028-3878
- Milone, M. & Engel, A. G. (1996). Block of the endplate acetylcholine receptor channel by the sympathomimetic agents ephedrine, pseudoephedrine, and albuterol. *Brain Research*, Vol. 740, No. 1-2, pp. 346-352, ISSN 0006-8993
- Milone, M., Wang, H. L., Ohno, K., Fukudome, T., Pruitt, J. N., Bren, N., Sine, S. M. & Engel, A. G. (1997). Slow-channel myasthenic syndrome caused by enhanced activation, desensitization, and agonist binding affinity attributable to mutation in the M2 domain of the acetylcholine receptor alpha subunit. *Journal of Neuroscience*, Vol. 17, No. 15, pp. 5651-5665, ISSN 0270-6474
- Milone, M., Wang, H.-L., Ohno, K., Prince, R., Fukudome, T., Shen, X.-M., Brengman, J. M., Griggs, R. C., Sine, S. M. & Engel, A. G. (1998). Mode switching kinetics produced by a naturally occurring mutation in the cytoplasmic loop of the human acetylcholine receptor epsilon subunit. *Neuron*, Vol. 20, No. 3, pp. 575-588, ISSN 0896-6273
- Milone, M., Shen, X. M., Selcen, D., Ohno, K., Brengman, J., Iannaccone, S. T., Harper, C. M. & Engel, A. G. (2009). Myasthenic syndrome due to defects in rapsyn: Clinical and

- molecular findings in 39 patients. *Neurology*, Vol. 73, No. 3, pp. 228-235, ISSN 0028-3878
- Mishina, M., Takai, T., Imoto, K., Noda, M., Takahashi, T., Numa, S., Methfessel, C. & Sakmann, B. (1986). Molecular distinction between fetal and adult forms of muscle acetylcholine receptor. *Nature*, Vol. 321, No. 6068, pp. 406-411, ISSN 0028-0836
- Mukhtasimova, N., Lee, W. Y., Wang, H. L. & Sine, S. M. (2009). Detection and trapping of intermediate states priming nicotinic receptor channel opening. *Nature*, Vol. 459, No. 7245, pp. 451-454, ISSN 0028-0836
- Muller, J. S., Mildner, G., Muller-Felber, W., Schara, U., Krampfl, K., Petersen, B., Petrova, S., Stucka, R., Mortier, W., Bufler, J., Kurlemann, G., Huebner, A., Merlini, L., Lochmuller, H. & Abicht, A. (2003). Rapsyn N88K is a frequent cause of congenital myasthenic syndromes in European patients. *Neurology*, Vol. 60, No. 11, pp. 1805-1810, ISSN 0028-3878
- Muller JS, Abicht A, Burke G, Cossins J, Richard P, Baumeister SK, et al. The congenital myasthenic syndrome mutation RAPSN N88K derives from an ancient Indo-European founder. *J Med Genet* 2004; 41: e104.
- Muller, J. S., Baumeister, S. K., Rasic, V. M., Krause, S., Todorovic, S., Kugler, K., Muller-Felber, W., Abicht, A. & Lochmuller, H. (2006). Impaired receptor clustering in congenital myasthenic syndrome with novel RAPSN mutations. *Neurology*, Vol. 67, No. 7, pp. 1159-1164, ISSN 0028-3878
- Muller, J. S., Herczegfalvi, A., Vilchez, J. J., Colomer, J., Bachinski, L. L., Mihaylova, V., Santos, M., Schara, U., Deschauer, M., Shevell, M., Poulin, C., Dias, A., Soudo, A., Hietala, M., Aarimaa, T., Krahe, R., Karcagi, V., Huebner, A., Beeson, D., Abicht, A. & Lochmuller, H. (2007). Phenotypical spectrum of DOK7 mutations in congenital myasthenic syndromes. *Brain*, Vol. 130, No. Pt 6, pp. 1497-1506, ISSN 0006-8950
- Noakes, P. G., Gautam, M., Mudd, J., Sanes, J. R. & Merlie, J. P. (1995). Aberrant differentiation of neuromuscular junctions in mice lacking s-laminin laminin beta 2. *Nature*, Vol. 374, No., pp. 258-262, ISSN 0028-0836
- Oda, Y., Nakanishi, I. & Deguchi, T. (1992). A complementary DNA for human choline acetyltransferase induces two forms of enzyme with different molecular weights in cultured cells. *Brain Research Molecular Brain Research*, Vol. 16, No. 3-4, pp. 287-294, ISSN 0006-8993
- Ohno, K., Hutchinson, D. O., Milone, M., Brengman, J. M., Bouzat, C., Sine, S. M. & Engel, A. G. (1995). Congenital myasthenic syndrome caused by prolonged acetylcholine receptor channel openings due to a mutation in the M2 domain of the epsilon subunit. *Proceedings of the National Academy of Sciences of the United States of America*, Vol. 92, No. 3, pp. 758-762, ISSN 0027-8424
- Ohno, K., Quiram, P. A., Milone, M., Wang, H.-L., Harper, M. C., Pruitt, J. N., 2nd, Brengman, J. M., Pao, L., Fischbeck, K. H., Crawford, T. O., Sine, S. M. & Engel, A. G. (1997). Congenital myasthenic syndromes due to heteroallelic nonsense/missense mutations in the acetylcholine receptor epsilon subunit gene: identification and functional characterization of six new mutations. *Human Molecular Genetics*, Vol. 6, No. 5, pp. 753-766, ISSN 0964-6906
- Ohno, K., Brengman, J., Tsujino, A. & Engel, A. G. (1998). Human endplate acetylcholinesterase deficiency caused by mutations in the collagen-like tail subunit

- (ColQ) of the asymmetric enzyme. *Proceedings of the National Academy of Sciences of the United States of America*, Vol. 95, No. 16, pp. 9654-9659, ISSN 0027-8424
- Ohno, K., Milone, M., Brengman, J. M., LoMonaco, M., Evoli, A., Tonali, P. A. & Engel, A. G. (1998). Slow-channel congenital myasthenic syndrome caused by a novel mutation in the acetylcholine receptor ϵ subunit. *Neurology*, Vol. 50 (Suppl. 4), No., pp. A432 (abstract), ISSN 0028-3878
- Ohno, K., Anlar, B. & Engel, A. G. (1999). Congenital myasthenic syndrome caused by a mutation in the Ets-binding site of the promoter region of the acetylcholine receptor epsilon subunit gene. *Neuromuscular Disorders*, Vol. 9, No. 3, pp. 131-135, ISSN 0960-8966
- Ohno, K., Brengman, J. M., Felice, K. J., Cornblath, D. R. & Engel, A. G. (1999). Congenital end-plate acetylcholinesterase deficiency caused by a nonsense mutation and an A->G splice-donor-site mutation at position +3 of the collagenlike-tail-subunit gene (COLQ): how does G at position +3 result in aberrant splicing? *American Journal of Human Genetics*, Vol. 65, No. 3, pp. 635-644, ISSN 0002-9297
- Ohno, K., Engel, A. G., Brengman, J. M., Shen, X.-M., Heidenrich, F. R., Vincent, A., Milone, M., Tan, E., Demirci, M., Walsh, P., Nakano, S. & Akiguchi, I. (2000). The spectrum of mutations causing endplate acetylcholinesterase deficiency. *Annals of Neurology*, Vol. 47, No., pp. 162-170, ISSN 0364-5134
- Ohno, K., Tsujino, A., Brengman, J. M., Harper, C. M., Bajzer, Z., Udd, B., Beyring, R., Robb, S., Kirkham, F. J. & Engel, A. G. (2001). Choline acetyltransferase mutations cause myasthenic syndrome associated with episodic apnea in humans. *Proceedings of the National Academy of Sciences of the United States of America*, Vol. 98, No. 4, pp. 2017-2022, ISSN 0027-8424
- Ohno, K., Engel, A. G., Shen, X.-M., Selcen, D., Brengman, J., Harper, C. M., Tsujino, A. & Milone, M. (2002). Rapsyn mutations in humans cause endplate acetylcholine-receptor deficiency and myasthenic syndrome. *American Journal of Human Genetics*, Vol. 70, No. 4, pp. 875-885, ISSN 0002-9297
- Ohno, K., Sadeh, M., Blatt, I., Brengman, J. M. & Engel, A. G. (2003). E-box mutations in the RAPSN promoter region in eight cases with congenital myasthenic syndrome. *Human Molecular Genetics*, Vol. 12, No. 7, pp. 739-748, ISSN 0964-6906
- Ohno, K. & Engel, A. G. (2004). Lack of founder haplotype for the rapsyn N88K mutation: N88K is an ancient founder mutation or arises from multiple founders. *Journal of Medical Genetics*, Vol. 41, No. 1, pp. e8, ISSN 0022-2593
- Okada, K., Inoue, A., Okada, M., Murata, Y., Kakuta, S., Jigami, T., Kubo, S., Shiraishi, H., Eguchi, K., Motomura, M., Akiyama, T., Iwakura, Y., Higuchi, O. & Yamanashi, Y. (2006). The muscle protein Dok-7 is essential for neuromuscular synaptogenesis. *Science*, Vol. 312, No. 5781, pp. 1802-1805, ISSN 0036-8075
- Okuda, T., Haga, T., Kanai, Y., Endou, H., Ishihara, T. & Katsura, I. (2000). Identification and characterization of the high-affinity choline transporter. *Nature Neuroscience*, Vol. 3, No. 2, pp. 120-125, ISSN 1097-6256
- Patton, B. L., Chiu, A. Y. & Sanes, J. R. (1998). Synaptic laminin prevents glial entry into the synaptic cleft. *Nature*, Vol. 393, No. 6686, pp. 698-701, ISSN 0028-0836
- Peng, H. B., Xie, H., Rossi, S. G. & Rotundo, R. L. (1999). Acetylcholinesterase clustering at the neuromuscular junction involves perlecan and dystroglycan. *Journal of Cell Biology*, Vol. 145, No., pp. 911-921, ISSN 0021-9525

- Ptacek, L. J., George, A. L., Jr., Griggs, R. C., Tawil, R., Kallen, R. G., Barchi, R. L., Robertson, M. & Leppert, M. F. (1991). Identification of a mutation in the gene causing hyperkalemic periodic paralysis. *Cell*, Vol. 67, No. 5, pp. 1021-1027, ISSN 0092-8674
- Ptacek, L. J., George, A. L., Jr., Barchi, R. L., Griggs, R. C., Riggs, J. E., Robertson, M. & Leppert, M. F. (1992). Mutations in an S4 segment of the adult skeletal muscle sodium channel cause paramyotonia congenita. *Neuron*, Vol. 8, No. 5, pp. 891-897, ISSN 0896-6273
- Ramarao, M. K., Bianchetta, M. J., Lanken, J. & Cohen, J. B. (2001). Role of rapsyn tetratricopeptide repeat and coiled-coil domains in self-association and nicotinic acetylcholine receptor clustering. *Journal of Biological Chemistry*, Vol. 276, No. 10, pp. 7475-7483, ISSN 0021-9258
- Ramarao, N. K. & Cohen, J. B. (1998). Mechanism of nicotinic acetylcholine receptor cluster formation by rapsyn. *Proceedings of the National Academy of Sciences of the United States of America*, Vol. 95, No. 7, pp. 4007-4012, ISSN 0027-8424
- Reimer, R. J., Fon, E. A. & Edwards, R. H. (1998). Vesicular neurotransmitter transport and the presynaptic regulation of quantal size. *Current Opinion in Neurobiology*, Vol. 8, No. 3, pp. 405-412, ISSN 0959-4388
- Richard, P., Gaudon, K., Andreux, F., Yasaki, E., Prioleau, C., S, B., Barois, A., Ioos, C., Mayer, M., Routon, M. C., Mokhtari, M., Leroy, J. P., Fournier, E., Hainque, B., Koenig, J., Fardeau, M., Eymard, B. & D, H. (2003). Possible founder effect of rapsyn N88K mutation and identification of novel rapsyn mutations in congenital myasthenic syndromes. *Journal of Medical Genetics*, Vol. 40, No. 6, pp. 81e, ISSN 0022-2593
- Sanes, J. R. (1997). Genetic analysis of postsynaptic differentiation at the vertebrate neuromuscular junction. *Current Opinion in Neurobiology*, Vol. 7, No. 1, pp. 93-100, ISSN 0959-4388
- Schara, U., Christen, H. J., Durmus, H., Hietala, M., Krabetz, K., Rodolico, C., Schreiber, G., Topaloglu, H., Talim, B., Voss, W., Pihko, H., Abicht, A., Muller, J. S. & Lochmuller, H. (2010). Long-term follow-up in patients with congenital myasthenic syndrome due to CHAT mutations. *Eur J Paediatr Neurol*, Vol. 14, No. 4, pp. 326-333, ISSN 1090-3798
- Schmidt, C., Abicht, A., Krampf, K., Voss, W., Stucka, R., Mildner, G., Petrova, S., Schara, U., Mortier, W., Bufler, J., Huebner, A. & Lochmuller, H. (2003). Congenital myasthenic syndrome due to a novel missense mutation in the gene encoding choline acetyltransferase. *Neuromuscular Disorders*, Vol. 13, No. 3, pp. 245-251, ISSN 0960-8966
- Selcen, D., Milone, M., Shen, X. M., Harper, C. M., Stans, A. A., Wieben, E. D. & Engel, A. G. (2008). Dok-7 myasthenia: phenotypic and molecular genetic studies in 16 patients. *Annals of Neurology*, Vol. 64, No. 1, pp. 71-87, ISSN 0364-5134
- Selcen, D., Juel, V. C., Hobson-Webb, L. D., Smith, E. C., Stickler, D. E., Bite, A. V., Ohno, K. & Engel, A. G. (2011). Myasthenic syndrome caused by plectinopathy. *Neurology*, Vol. 76, No. 4, pp. 327-336, ISSN 0028-3878
- Senderek, J., Muller, J. S., Dusl, M., Strom, T. M., Guergueltcheva, V., Diepolder, I., Laval, S. H., Maxwell, S., Cossins, J., Krause, S., Muelas, N., Vilchez, J. J., Colomer, J., Mallebrera, C. J., Nascimento, A., Nafissi, S., Kariminejad, A., Nilipour, Y., Bozorgmehr, B., Najmabadi, H., Rodolico, C., Sieb, J. P., Steinlein, O. K., Schlotter,

- B., Schoser, B., Kirschner, J., Herrmann, R., Voit, T., Oldfors, A., Lindbergh, C., Urtizberea, A., von der Hagen, M., Hubner, A., Palace, J., Bushby, K., Straub, V., Beeson, D., Abicht, A. & Lochmuller, H. (2011). Hexosamine biosynthetic pathway mutations cause neuromuscular transmission defect. *American Journal of Human Genetics*, Vol. 88, No. 2, pp. 162-172, ISSN 0002-9297
- Shen, X.-M., Ohno, K., Sine, S. M. & Engel, A. G. (2005). Subunit-specific contribution to agonist binding and channel gating revealed by inherited mutation in muscle acetylcholine receptor M3-M4 linker. *Brain*, Vol. 128, No., pp. 345-355, ISSN 0006-8950
- Shen, X. M., Ohno, K., Milone, M., Brengman, J. M., Tsujino, A. & Engel, A. G. (2003). Effect of residue side-chain mass on channel kinetics in second transmembrane domain of muscle AChR. *Molecular Biology of the Cell*, Vol. 14 (Suppl), No., pp. 223a (abstract), ISSN 1059-1524
- Sine, S. M., Ohno, K., Bouzat, C., Auerbach, A., Milone, M., Pruitt, J. N. & Engel, A. G. (1995). Mutation of the acetylcholine receptor alpha subunit causes a slow-channel myasthenic syndrome by enhancing agonist binding affinity. *Neuron*, Vol. 15, No. 1, pp. 229-239, ISSN 0896-6273
- Smith, F. J., Eady, R. A., Leigh, I. M., McMillan, J. R., Rugg, E. L., Kelsell, D. P., Bryant, S. P., Spurr, N. K., Geddes, J. F., Kirtschig, G., Milana, G., de Bono, A. G., Owaribe, K., Wiche, G., Pulkkinen, L., Uitto, J., McLean, W. H. & Lane, E. B. (1996). Plectin deficiency results in muscular dystrophy with epidermolysis bullosa. *Nature Genetics*, Vol. 13, No. 4, pp. 450-457, ISSN 1061-4036
- Tsujino, A., Maertens, C., Ohno, K., Shen, X.-M., Fukuda, T., Harper, C. M., Cannon, S. C. & Engel, A. G. (2003). Myasthenic syndrome caused by mutation of the SCN4A sodium channel. *Proceedings of the National Academy of Sciences of the United States of America*, Vol. 100, No., pp. 7377-7382, ISSN 0027-8424
- Valenzuela, D. M., Stitt, T. N., DiStefano, P. S., Rojas, E., Mattsson, K., Compton, D. L., Nunez, L., Park, J. S., Stark, J. L., Gies, D. R. & et al. (1995). Receptor tyrosine kinase specific for the skeletal muscle lineage: expression in embryonic muscle, at the neuromuscular junction, and after injury. *Neuron*, Vol. 15, No. 3, pp. 573-584, ISSN 0896-6273
- Vogt, J., Morgan, N. V., Marton, T., Maxwell, S., Harrison, B. J., Beeson, D. & Maher, E. R. (2009). Germline mutation in DOK7 associated with fetal akinesia deformation sequence. *Journal of Medical Genetics*, Vol. 46, No. 5, pp. 338-340, ISSN 0022-2593
- Wang, H.-L., Auerbach, A., Bren, N., Ohno, K., Engel, A. G. & Sine, S. M. (1997). Mutation in the M1 domain of the acetylcholine receptor alpha subunit decreases the rate of agonist dissociation. *Journal of General Physiology*, Vol. 109, No. 6, pp. 757-766, ISSN 0022-1295
- Wang, H.-L., Milone, M., Ohno, K., Shen, X.-M., Tsujino, A., Batocchi, A. P., Tonali, P., Brengman, J., Engel, A. G. & Sine, S. M. (1999). Acetylcholine receptor M3 domain: stereochemical and volume contributions to channel gating. *Nature Neuroscience*, Vol. 2, No. 3, pp. 226-233, ISSN 1097-6256
- Wang, H.-L., Ohno, K., Milone, M., Brengman, J. M., Evoli, A., Batocchi, A. P., Middleton, L. T., Christodoulou, K., Engel, A. G. & Sine, S. M. (2000). Fundamental gating mechanism of nicotinic receptor channel revealed by mutation causing a congenital

- myasthenic syndrome. *Journal of General Physiology*, Vol. 116, No. 3, pp. 449-462, ISSN 0022-1295
- Wells, L., Vosseller, K. & Hart, G. W. (2001). Glycosylation of nucleocytoplasmic proteins: signal transduction and O-GlcNAc. *Science*, Vol. 291, No. 5512, pp. 2376-2378, ISSN 0036-8075
- Wirtz, P. W., Titulaer, M. J., Gerven, J. M. & Verschuuren, J. J. (2010). 3,4-diaminopyridine for the treatment of Lambert-Eaton myasthenic syndrome. *Expert Rev Clin Immunol*, Vol. 6, No. 6, pp. 867-874, ISSN 1744-666X
- Wu, H., Xiong, W. C. & Mei, L. (2010). To build a synapse: signaling pathways in neuromuscular junction assembly. *Development*, Vol. 137, No. 7, pp. 1017-1033, ISSN 0950-1991
- Yasaki, E., Prioleau, C., Barbier, J., Richard, P., Andreux, F., Leroy, J. P., Darteville, P., Koenig, J., Molgo, J., Fardeau, M., Eymard, B. & Hantai, D. (2004). Electrophysiological and morphological characterization of a case of autosomal recessive congenital myasthenic syndrome with acetylcholine receptor deficiency due to a N88K rapsyn homozygous mutation. *Neuromuscular Disorders*, Vol. 14, No. 1, pp. 24-32, ISSN 0960-8966
- Yeung, W. L., Lam, C. W., Fung, L. W., Hon, K. L. & Ng, P. C. (2009). Severe congenital myasthenia gravis of the presynaptic type with choline acetyltransferase mutation in a Chinese infant with respiratory failure. *Neonatology*, Vol. 95, No. 2, pp. 183-186, ISSN 1661-7800
- Zenker, M., Aigner, T., Wendler, O., Tralau, T., Muntefering, H., Fenski, R., Pitz, S., Schumacher, V., Royer-Pokora, B., Wuhl, E., Cochat, P., Bouvier, R., Kraus, C., Mark, K., Madlon, H., Dotsch, J., Rascher, W., Maruniak-Chudek, I., Lennert, T., Neumann, L. M. & Reis, A. (2004). Human laminin beta2 deficiency causes congenital nephrosis with mesangial sclerosis and distinct eye abnormalities. *Human Molecular Genetics*, Vol. 13, No. 21, pp. 2625-2632, ISSN 0964-6906
- Zhang, B., Luo, S., Wang, Q., Suzuki, T., Xiong, W. C. & Mei, L. (2008). LRP4 serves as a coreceptor of agrin. *Neuron*, Vol. 60, No. 2, pp. 285-297, ISSN 0896-6273

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](https://creativecommons.org/licenses/by/3.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen