

# We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

185,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index  
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?  
Contact [book.department@intechopen.com](mailto:book.department@intechopen.com)

Numbers displayed above are based on latest data collected.  
For more information visit [www.intechopen.com](http://www.intechopen.com)



# Effect of Priming on Dormancy Breaking and Seedling Establishment of Caper (*Capparis spinosa* L.)

Saeed Khaninejad\*, Iman Hessam Arefi, and Mohammad Kafi

Department of Agronomy, Ferdowsi University of Mashhad, Iran

## Abstract

Caper (*Capparis spinosa*) has deep roots, drought tolerant species, and produces a satisfactory vegetative cover which protects soils from erosion; it can be highly useful for the prevention of land degradation. Domestication of capers as medicinal, vegetable or soil surface coverage plant is complicated by limited and variable seed germination under artificial conditions. In order to examine the role of different levels of  $\text{KNO}_3$  (0, 500, 1000, 2000, 4000 and 8000 ppm) and gibberellic acid ( $\text{GA}_3$ ) (0, 50, 100, 250, 500, 1000 and 2000 ppm) and durations (3, 12, 24 and 48 hr) on germination of Iranian caper seeds. In general 2000 mg/l gibberellic acid treatment resulted in more vigorous seed germination (42%) at any duration compared to any other concentration of the gibberellic acid. The highest seed germination of 26% was achieved when the seeds were treated 24 hour with 4000 ppm  $\text{KNO}_3$  solution but it was decreased in 8000 mg/l. The highest germination percentage (72%) was observed in seeds placed in filter papers wetted with in 250 ppm gibberellic acid after treatment with 8000 mg/l  $\text{KNO}_3$  for 24 hour (this duration was the best time span in two previous experiments). It seems that  $\text{GA}_3$  and  $\text{KNO}_3$  can replace partly to improve seed germination of caper. The highest seedling dry weight was achieved as seeds were treated in 100 ppm gibberellic acid plus 1000 ppm potassium nitrate. Therefore, it can be concluded that for best germination percentage of caper seeds, 250 ppm  $\text{GA}_3$  and 8000 ppm  $\text{KNO}_3$  and for the strongest seedling 100 ppm  $\text{GA}_3$  plus 1000 ppm  $\text{KNO}_3$  could be recommended.

**Keywords:** Caper, Dormancy,  $\text{GA}_3$ ,  $\text{KNO}_3$ , Medicinal Plant

## 1. Introduction

Caper (*Capparis spinosa* L.) plant is typical of the tropical Mediterranean areas as well as central Asia, Europe (Spain, Italy, Greece and Turkey) as well as North Africa and Middle East countries including Iran [9,11]. It is a perennial shrub, covering soil surface widely, and produces one of the deepest root systems [14]. The long roots and wide ecological amplitude allow it to withstand harsh environments. The species thus appears to be a suitable candidate for the protection of degraded areas in arid areas [14]. Apart from its roles in soil conservation, different parts of the caper plant is used in Iranian traditional medicine as a liver provoking agent, treatment of vessel clogs, anti-rheumatism and a diuretic agent [13]. Seed dormancy is an adaptive mechanism in many species particularly wild species to protect seedlings from freeze damage during the winter, or from drought stress in water shortage conditions [15]. Caper species is endangered in its natural habitats in Iran and many other countries due to climate change and overutilization.

Domestication of capers as medicinal, vegetable or soil surface coverage plant is complicated by limited and variable seed germination under artificial conditions. According to some researchers, there are germination obstacles in the caper seeds and; thus, there are propagation difficulties of caper seedlings [11]. To ensure high plantation and viability, a reliable and high germination percentage is required. Plant growth regulators such as gibberellic acid ( $GA_3$ ) [6] and chemicals such as  $KNO_3$  [5] have been recommended to break seed dormancy and enhance germination. Gibberellins promote growth by increasing extendibility of the cell wall followed by the hydrolysis of starch to sugars which reduces the potential in the cell, resulting in the entry of water into the cell causing elongation [2].  $KNO_3$  is the most widely used chemical for germination promoting [11]. Information on seed germination of capers is still limited. Therefore, it was thought that treatment of the seeds with plant growth regulators may influence rapid germination and root formation. The present study was conducted to examine the role of  $KNO_3$  and  $GA_3$  that might affect germination of Iranian capers seeds and study the possible advantages of in vitro germination over direct sowing of seeds in the soil.

### 1.1. Materials and Methods

In order to evaluate germination properties and effect of different treatments on dormancy breaking of *caparis spinosa*, these experiments were conducted at Physiological Research Laboratory, Ferdowsi University of Mashhad, Iran, in 2010. The seeds used in this study were collected from Shothern Khorasan province, Iran, where capers grows abundantly. The seeds were separated from the fruit material, rinsed in tap water, dried in shade and kept at room temperature in linen sacks until sowing.

The germination test conducted to find out germination properties of caper seeds in  $H_2O$  as control, concentrated gibberellic acid, potassium nitrate, gibberellic acid + potassium nitrate. A sample of 25 randomly selected seeds was used with four replications. Experiments were carried out in 90mm diameter Petri dishes using 90mm diameter Whatman No. 1 filter papers in the bottom and top of the petri dishes to cover the seeds. Experiment was continued for 21 days and every day the procedure of germination was monitored.

Similar to previous experiment, samples of twenty five randomly selected seeds with four replications were soaked in different doses (0, 50, 100, 250, 500, 1000 and 2000 ppm) and durations (3, 12, 24 and 48 hr) of gibberellic acid solution. No further water was added during the experiment period. For control treatment, Filter papers were moistened with distilled water.

Another 21 days experiment was conducted to find out potassium nitrate solution on germination. Seeds soaked in different doses (0, 500, 1000, 2000, 4000 and 8000 ppm) (Amri, 2010) and durations (3, 12, 24 and 48 hr) of potassium nitrate. Samples and germinated seeds were counted every day until the end of the experiment.

The seeds were placed in filter papers were soaked in 0, 50, 100, 250, 500, 1000 and 2000 ppm gibberellic acid solution after applying different doses (0, 500, 1000, 2000, 4000 and 8000 ppm) of potassium nitrate solution for 24 h. For control treatment, filter papers were moistened with distilled water. This experiment also continued for 21 days and seed germination was monitored every day.

The combined experiment of gibberellic acid and potassium nitrate was arranged in the polyethylene pots filled with growing medium composed of clay, sand and manure (1:1:1). Pots were kept in open air conditions after sowing. Seven different gibberellic acid treatment (0, 50, 100,

250, 500, 1000 and 2000 ppm) and six different potassium nitrate (0, 500, 1000, 2000, 4000 and 8000 ppm) treatment were applied on the seeds in the spring (March) of 2010. The experimental design was a factorial based on randomized complete block with three replications (35 pots for each block) for every treatment.

Analysis of variance performed using SAS 9.1 (SAS Institute, Cary, NC) and general linear models (PROC GLM) procedure. Multiple comparisons were conducted for significant effects with the least significant difference (LSD) test at  $\alpha = 0.05$  to determine the rate of seed germination for each patch and seed treatment.

## 2. Result and Discussions

There was no any symptom of germination of caper seeds in distilled water, Olmez *et al.*, (2004) reported only 3.67% germination in this plant using distilled water. The seed infection was started form the 4<sup>th</sup> day of the experiment and after 21 days, there were some damaged seeds in each petri dish.

### 2.1. Gibberellic acid treatment

The highest seed germination of 42% was achieved after 24 h of soaking in 2000 mg/l GA<sub>3</sub> (Table 1). However, 38% seed germination was achieved in the same duration at 1000 mg/l GA<sub>3</sub> (Table 1). In general 2000 mg/l GA<sub>3</sub> treatment resulted in more vigorous seed germination at any duration compared to any other concentration of the gibberellic acid. This indicates that the regulation of endogenous GA<sub>3</sub> levels after seed imbibitions is a crucial factor in determining seed germination. Duration of exposure of seeds to GA<sub>3</sub> is also important; for instance the germination percentage at 1000 mg/l of GA<sub>3</sub> for 6 hours was only 8% while at the same concentration of GA<sub>3</sub> it was 38% after 24 hours of GA<sub>3</sub> treatment. Germination percentage was reduced beyond 24 hours of soaking in gibberellic acid at all concentrations. Gibberellic acid is known to play an essential role in seed germination, leaf expansion, stem elongation, flowering and flower development [16]. Our results are in agreement with that of Negbi *et al.* [10] for *Hirschfeldia incana* and Orphanos [12] and Olmez *et al.*, [11] for *Capparis spinosa*, who found that the seed dormancy is mainly due to the seed coat that prevents germination and GA<sub>3</sub> has a positive effect on germination. Mayer and Shahin [8] observed that the gibberellins reduces oxygen requirement for germination. The caper germination percentage obtained in this experiment is higher than previously reported e.i. 27.4 % in Olmez *et al.*, [11] experiment.

Soaking time (hours)	GA <sub>3</sub> dose (mg/l)					
	50	100	250	500	1000	2000
6	6j*	6j	6j	10hi	8i	10hi
12	8i	8i	10hi	12gh	12gh	13fgh
24	24d	30c	30c	32c	38b	42a
48	14fg	16f	16f	24d	20e	22de

\*Values followed by different letters are significantly different at 0.05 level using LSD. There was no germination in distilled water (zero GA<sub>3</sub>)

**Table 1.** Effect of different duration and concentration of GA<sub>3</sub> on seed germination percentage of *C. spinosa*

## 2.2. Potassium nitrate treatment

Treatment with exogenous  $\text{KNO}_3$  stimulated of germination percentage of caper seeds. The highest seed germination of 26% was achieved when the seeds were treated 24 hour with 4000 mg/l  $\text{KNO}_3$  solution but it was decreased in 8000 mg/l. Moreover, germination percentage of caper seeds were dependence to concentrations up to 4000 mg/l and duration dependence up to 24 hours (Table 2). Several workers have reported that  $\text{KNO}_3$  improved the seed germination of many plants seed [3,11]. Potassium nitrate was found to be effective in breaking dormancy of many species [1]. Use of  $\text{KNO}_3$  has been an important seed treatment in seed-testing laboratories for many years without a good explanation for its action [5].

Soaking time	$\text{KNO}_3$ dose (mg/l)				
	500	1000	2000	4000	8000
6 hour	6c*	6c	8b	12c	10d
12 hour	12b	14fg	20cd	24ab	14fg
24 hour	16ef	18de	20cd	26a	22bc
48 hour	18de	16ef	18de	22bc	18de

\*Values followed by different letters are significantly different at 0.05 level using LSD. There was no germination in distilled water (zero  $\text{KNO}_3$ )

**Table 2.** Effect of different duration and concentration of  $\text{KNO}_3$  on seed germination percentage of *C. spinosa*

## 2.3. Potassium Nitrate + Gibberellic acid treatment:

The highest germination percentage (72%) was observed in seeds placed in filter papers wetted with in 250 ppm gibberellic acid after treatment with 8000 mg/l  $\text{KNO}_3$  for 24 hour (this duration was the best time span in two previous experiment), however there was no any significant difference between this germination percentage and that of 1000 ppm gibberellic acid after treatment with 4000 mg/lit  $\text{KNO}_3$  (Table 3). It seems that  $\text{GA}_3$  and  $\text{KNO}_3$  can replace partly to improve seed germination of caper. No germination in case of control was possibly due to the seed coat of the capers that forms mucilage on soaking in water [12]. The mucilage surrounding the seed is supposed to inhibit diffusion of oxygen to the embryos and prevent germination. Khan and Ungar [7] believed that vegetative hormones can break embryo dormancy and neutralize prevention role of Abscissas acid (ABA) directly or indirectly. But we could not find the synergistic effects of  $\text{GA}_3$  and  $\text{KNO}_3$  on germination of caper and other species in the literature.

$\text{GA}_3$ dose (mg/l)	$\text{KNO}_3$ dose (mg/l)					
	0	500	1000	2000	4000	8000
0	0	16u*	18tu	20st	22rs	22rs
50	24qr	22rs	26q	36lm	36lm	44hi
100	30op	30op	32no	42ij	42ij	64bc

GA <sub>3</sub> dose (mg/l)	KNO <sub>3</sub> dose (mg/l)					
	0	500	1000	2000	4000	8000
250	30op	40jk	34mn	48fg	52e	72a
500	32no	44hi	42ij	50ef	66b	46gh
1000	38kl	46gh	44hi	62c	70a	48fg
2000	42ij	46gh	56d	46gh	50ef	46gh

\*Values with in a column followed by different letters are significantly different at 0.05 level using LSD.

**Table 3.** Effect of different concentration of KNO<sub>3</sub> + GA<sub>3</sub> on seed germination percentage of *C. spinosa*

#### 2.4. Seedling establishment:

Significant effect of seed priming treatment was observed in the final seeding establishment. The highest seedling dry weight of 25 mg/plant was achieved when the seeds were treated with 100 ppm gibberellic acid treatment + 1000 ppm Potassium Nitrate (Table4). The germination phase of planted seeds is critical because it directly determines the density of a crop stand especially under arid conditions. Where dry soil may impair imbibitions of water and high temperature may affect seed viability and eventual density of a crop stand [4]. Hadas and Russo [4] further observed that a good stand can be ensured by a complete and fast germination and if germination seed is slow in taking up water, emergence is impaired and consequently the final stand is reduced.

GA <sub>3</sub> dose (mg/l)	KNO <sub>3</sub> dose (mg/l)					
	0	500	1000	2000	4000	8000
0		17ghi	18fgh	19efg	19efg	18fgh
50	18fgh*	19efg	19efg	20def	20def	19efg
100	20def	21cde	22bcd	23abc	23abc	22bcd
250	20def	23abc	25a	24ab	23abc	22bcd
500	19efg	20def	22bcd	21cde	20def	18fgh
1000	16hij	18fgh	18fgh	19efg	17ghi	15ijk
2000	12l	13kl	14jkl	14jkl	13kl	12l

\*Values followed by different letters are significantly different at 0.05 level using LSD.

**Table 4.** Effect of KNO<sub>3</sub> + GA<sub>3</sub> on seedling dry weight (mg) of *C. spinosa*

In general to improve caper seed germination, 2000 mg/l gibberellic acid treatment resulted in more vigorous seed germination (42%) at any duration compared to other concentrations. While, the highest seed germination of 26% was achieved when the seeds were treated 24 hour with 4000 ppm KNO<sub>3</sub> solution but it was decreased in 8000 mg/l. The highest germination percentage (72%) was observed in seeds placed in filter papers wetted with in 250 ppm gibberellic acid after treatment with 8000 mg/l KNO<sub>3</sub> for 24 hour (this duration was the best time span in two previ-

ous experiments). It seems that GA<sub>3</sub> and KNO<sub>3</sub> have a synergist effect and can replace partly to improve seed germination of caper. The highest seedling dry weight was achieved as seeds were treated in 100 ppm gibberellic acid + 1000 ppm potassium nitrate. Therefore, it can be concluded that for best germination percentage of caper seeds, 250 ppm GA<sub>3</sub> and 8000 ppm KNO<sub>3</sub> and for the strongest caper seedling 100 ppm GA<sub>3</sub> + 1000 ppm KNO<sub>3</sub> could be recommended for dormancy breaking of caper seeds.

### 3. Reference

- [1] Agrawal, P., K. and M. Dadlani. Techniques in seed science and technology. Second Edition. South Asian Publishers, New Delhi International Book Company Absecon Highlands: 1995. pp. 109–113.
- [2] Arteca, R., N. Plant growth substances principles and applications. Chapter 3: Chemistry, Biological Effects and Mechanism of Action, 1996. p: 66. Chapman & Hall 115 Fifty Avenue New York, NY 10003.
- [3] Cirak, C., A. K. Ayan and K. Kevseroglu. 2004. The effect of light and some presoaking treatments on germination rate of St. John's worth (*Hypericum perforatum* L.) seeds. Pakistan J. Biol. Sci., 7(2): 182-186.
- [4] Hadas, A. and D. Russo. 1974. Water uptake by seeds as affected by water stress, capillary conductivity and seed-soil water contact. Agron. J., 66: 643-647.
- [5] Hartmann H. T., D. E. Kester, F. Jr. Davies, and R. L. Geneve. 1997. Plant propagation principles and practices. Sixth Edition. New Jersey, Prentice Hall.
- [6] Iglesias, R. G., and M. J. Babiano. 1997. Endogenous abscisic acid during the germination of chickpea seed. Physiol. Planta., 100: 500–504.
- [7] Khan, M. A, and I. A. Ungar. 1985. The role of hormones in regulating the germination of polymorphic seeds and early seedling growth of *Atriplex triangularis* under saline conditions. Physiol. Planta., 63: 109–113.
- [8] Mayer, A. M. and Y. Shahin. 1974. Control of seed germination. Annual Rev. Plant Physiol., 25: 167–93.
- [9] Movafeghi A., Gh. Habibi and M. Aliasgarpoor. 2008. Plant regeneration of *Capparis spinosa* L. using hypocotyl explants. Iranian J. Biology, 21(2):289-297.
- [10] Negbi, M., E. Rushkin and D. Koller. 1966. Dynamic aspects of water relations in germination of *Hirschfeldia incana* seeds. Plant Cell Physiol., 7: 363–76.
- [11] Olmez, Z., Z. Yahyaoglu and A. Ö. Üçler. 2004. Effects of H<sub>2</sub>SO<sub>4</sub>, KNO<sub>3</sub> and GA<sub>3</sub> treatments on germination of caper (*Capparis ovata* Desf.) seeds. Pakistan Biol. Sci., 7 (6): 879-882.
- [12] Orphanos, P. I. 1983. Germination of caper (*Capparis spinosa* L.) seeds. J. Hort. Sci., 58: 267–70.
- [13] Ramazani M., M. Taghvaei, M. Masoudi, A. Riahi and N. Behbahani. 2009. The evaluation of drought and salinity effects on germination and seedling growth of Caper (*Capparis spinosa* L.). Iran. J. Rangeland, 2(4): 411-420.
- [14] Sakcali M. S., H. Bahadir and M. Ozturk. 2008. Eco-physiology of *Capparis spinosa* L. : a plant suitable for combating desertification Pak. J. Bot., 40(4): 1481-1486.
- [15] Shirazi, A. M. 2003. Standardizing methods for evaluating the chilling requirements to break dormancy in seeds and buds (including geophytes): introduction to the workshop. HortScience, 38: 334–335.
- [16] Yanmaguchi, S. and Y. Kamiya. 2002. Gibberellins and light stimulated seed germination. J. Plant Growth Regul., 20: 369–76.