

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

185,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Effect of Fungal Growth Inhibition from Pomegranate Flower and Peel Extracts

Mahsa Shafighi¹, Leila Amjad², Mahboubeh Madani³

1 Young Researchers Club, Falavarjan Branch, Islamic Azad University, Isfahan, Iran

2 Departments of Biology, Falavarjan Branch, Islamic Azad University, Isfahan, Iran

Departments of Microbiology, Falavarjan Branch, Islamic Azad University, Isfahan, Iran

Abstract

Candida species are now one of the most common organisms isolated from hospitalized patients. The range of antifungal agents available is limited, and some of the most effective agents are also toxic. In the other hand plants have been used for thousands of years to flavor and conserve food, to treat health disorders and to prevent diseases including epidemics. The aim of this study was to investigate the antifungal effect of petroleum ether, ethyl acetate and n-butanol fractions isolated from pomegranate pericarp and flower against *Candida albicans* (ATCC 3153). The maximum inhibition zone of *Candida albicans* was obtained by peel's n-butanol fraction, 35 mm. Petroleum ether fractions had no any antifungal activities.

Keywords: *Punica granatum* L., peel, flower, petroleum ether, ethyl acetate, n-butanol

1. Introduction

The *Punica granatum* L., is one of the oldest known edible fruits. It is native to Persia and from there it spreads into Asia, North Africa and Mediterranean Europe, including Turkey [1, 2]. According to Qur'an, Bible, Torah and Babylonian Talmud, pomegranate is a gift and heavenly fruit of God.

The extracts of traditional herbs have been shown to exert biological activity in vitro and in vivo. *Punica granatum* is employed in man medicine for the treatment of various diseases such as skin diseases, and wound healing, ulcers, fever, diarrhoea, and microbial infection. In the recently years, the *Punica granatum* has been the subject of much scientific research which have showed its antimicrobial, antioxidant and anti-cancer effects [3, 4].

The different types of phytochemical that have been showed identified from pomegranate pericarp (peel, rind) and pomegranate flower. Pomegranate pericarp's constituents are Luteolin, kaempferol, EA glycosides, EA, Pedunculagin, punicalin, phenolic punicalagins, Gallic acid and other fatty acids, catechin, EGCG, quercetin, rutin and other flavonols, flavones, flavonones, anthocyanidins. Pomegranate flower's constituents are Polyphenols, punicalagins punicalin, EA, Gallic acid, ursolic acid, triterpenoids, including maslinic and Asiatic acid [1, 5- 7].

Candida species are harmless saprophyte yeasts, a normal component of the human biota in the gastrointestinal tract and oral and vaginal mucosae. These yeasts can cause superficial infections

such as thrush and vaginitis; however, if the immune defences of the host become compromised, they can cause severe systemic infections. Risk factors for patients include infection by the human immunodeficiency virus (HIV), anticancer therapy, organ transplantation, abdominal surgery, catheters, diabetes and the use of broad-spectrum antibiotics [4, 11].

The aim of this study was to investigate the antifungal effect of some fractions isolated from pomegranate pericarp and flower against *Candida albicans*.

2. Materials and Methods

Samples of *P. Granatum* flowers and peels were collected and identification in June and September 2011 respectively from the Agricultural Research Centre in Isfahan, Iran.

The flowers and peels were air-dried in a low light at room temperature for 1 week. The material was thereafter ground in an electric grinder to produce a powder separately.

Peels and flowers powder extracted with 100% petroleum ether (F1) at room temperature for 24 hours respectively, the extracts were filtered. Then 20 gram of remains powders were extracted with 75% Ethanol by Soxhlet extraction for 8 hours.

The residues were dried over night and then evaporated by using a rotary evaporator. The dried extracts were suspended in distilled water (F2) and partition with ethyl acetate (F3) followed by n-butanol (F4). All fractions were frigid at -20°C until each experiment.

The yeast strain *C. albicans* (ATCC 3153) were used in this study. It obtained from Institute of Scientific and Industrial Researches, Tehran, Iran. At the first, it has been maintained at 4°C on Sabouraud dextrose agar (SDA) plates and sub cultured at 25°C in Sabouraud dextrose broth (SDB) before each experiment to ensure viability and purity.

Petri dishes contained 20 ml of SDA have been used for well-diffusion assay. Wells have been prepared in the SDA plates. In agar well diffusion, 10, 50, 100 and 200 µl of each fractions have been inoculated to each well separately. Then 100 ml of 10⁶ CFU/ml yeast suspension was spread uniformly onto the agar plate using cotton swabs. Diameters (in mm) of growth inhibition zones were measured after incubation at 25°C for 24- 48 hours.

3. Results and Discussion

Candida species are now the fourth most common organism recovered from the blood of hospitalized patients. Notwithstanding the increasing need for effective therapy, the range of antifungal agents available is limited, and some of the most effective agents are also toxic. In addition, although azoles have been used successfully for the treatment of *Candida* infections, numerous reports of treatment failures are now appearing in the literature [4]. In view of the lack of new classes of drugs or different molecular targets, drug combinations might be considered a viable strategy for therapy, considering the multiplicity of fungal targets against which current agents are effective.

The antimicrobial and antifungal effects of alcoholic extracts of different parts of pomegranate tree were previously studied [1, 8-11]. Nevertheless, few studies have showed the antifungal effect of different pomegranate fractions. This study showed the antifungal activity of 4 pomegran-

ate fractions. Antifungal effects of peel and flower fractions by well-diffusion assay method have been shown in Figure 1 and Figure 2 respectively.

The maximum inhibition zones of peel fractions against *C. albicans* ATCC 3153 were obtained in 200 μ l concentrations by n- butanol fraction, water fraction and ethyl acetate fraction respectively.

C. albicans ATCC 3153 only was sensitive to flower n- butanol fraction. In addition *C. Albicans* ATCC 3153 was resistant to peel and flower Petroleumether fractions.

The fractions had no any antifungal activity in 10 μ l, and there was direct relationship between concentration and inhibition zone.

Also, n- butanol, ethyl acetate, water and petroleumether had no any antifungal activities. In the other words, the extracted compounds of peel and flower were effective [4].




Figure 1. The inhibitory zons (mm) of peel’s fractions in 4 concentrations against *C. albicans* (ATCC 3153).
F1: Petroleumether Fraction, F2: Water Fraction, F3: Ethyl acetate Fraction, F4: n-butanol Frcation.




Figure 2. The inhibitory zones (mm) of flower’s fractions in 4 concentrations against *C. albicans* (ATCC 3153).
F1: Petroleumether Fraction, F2: Water Fraction, F3: Ethyl acetate Fraction, F4: n-butanol Frcation.

Acknowledgment

This work was supported by Young Researchers Club, Falavarjan Branch, Islamic Azad University. The authors also thank Dr. Ranjbar, Dr. Khatabakhsh and Mrs. Shahsar for their kindly aid.

4. References

- [1] Saad Sabbar Dahham, Mir Naiman Ali, Hajera Tabassum and Mazharuddin Khan, 2010. Studies on Antibacterial and Antifungal Activity of Pomegranate (*Punica granatum* L.). American-Eurasian J. Agric. & Environ. Sci., 9 (3): 273-281.
- [2] Ahmet D. Duman, Mustafa Ozgen, Kenan S. Dayisoğlu, Nurcan Erbil and Coskun Durgac, 2009. Antimicrobial Activity of Six Pomegranate (*Punica granatum* L.) Varieties and Their Relation to Some of Their Pomological and Phytonutrient characteristics. Molecules, 14: 1808-1817.
- [3] Kannan Ponnusamy, Chelladurai Petchiammal, Ramasamy Mohankumar, Waheeta Hopper, 2010. In vitro antifungal activity of indirubin isolated from a South Indian ethnomedicinal plant *Wrightia tinctoria* R. Br. Journal of Ethnopharmacology, 132: 349-354.
- [4] Eliana Harue Endo, Dio'genes Apar'cio Garcia Cortez, Ta'nia Ueda-Nakamura, Celso Vataru Nakamura, Benedito Prado Dias Filho, 2010. Potent antifungal activity of extracts and pure compound isolated from pomegranate peels and synergism with fluconazole against *Candida albicans*. Research in Microbiology, 161: 534-540.
- [5] M. Viuda-Martos, J. Fern'andez-L'opez, and J.A. P'erez-'Alvarez, 2010. Pomegranate and its Many Functional Components as Related to Human Health: A Review. Comprehensive Reviews in Food Science and Food Safety, 9: 635-654.
- [6] J. Pongpuntaruk, 2010. Chemical constituents from the stem of *Punica granatum* and the root of *mechelia alba*, 2010. Songkla University.
- [7] Ephraim P. Lansky, Robert A. Newmana, 2007. *Punica granatum* (pomegranate) and its potential for prevention and treatment of inflammation and cancer. Journal of Ethnopharmacology, 109: 177-206.
- [8] L. VASCONCELOS, F. C. SAMPAIO, M. C. SAMPAIO, M. S. Vieira PEREIRA, J. S. HIGINO, M. P. PEIXOTO, 2006. Minimum Inhibitory Concentration of Adherence of *Punica granatum* Linn (pomegranate) Gel against *S. mutans*, *S. mitis* and *C. albicans*. Braz Dent J, 17(3): 223-227.
- [9] Silva NCC, Fernandes Júnior, 2010. Biological properties of medicinal plants: a review of their antimicrobial activity. The Journal of Venomous Animals and Toxins including Tropical Diseases, 3(16): 402-413.
- [10] Abdollah Ghasemi Pirbalouti, Mahmood Bahmani, Majid Avijgan, 2009. Anti-*Candida* Activity of Some of the Iranian Medicinal Plants. Electronic Journal of Biology, 5(4): 85-88.
- [11] Maria G. Miguel, Maria A. Neves and Maria D. Antunes, 2010. Pomegranate (*Punica granatum* L.): A medicinal plant with myriad biological properties - A short review. Journal of Medicinal Plants Research, 4(25), 2836-2847.