We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Steroidogenic Enzyme 17,20-Lyase Activity in Cortisolsecreting and Non-Functioning Adrenocortical Adenomas

Hajime Ueshiba

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/53212>

1. Introduction

The frequency of adrenal incidentalomas has been reported to range between 0.7% and 4% in abdominal computed tomography [1-5], or from 1.4% to 8.7% in autopsy series [6-8]. The widespread application of high resolution imaging techniques, such as computed tomography, magnetic resonance imaging and ultrasonography, has led to an increasing frequency of discovering adrenal incidentalomas. The majority of these lesions are of cortical origin, include non-functioning adenomas, adenomas associated with pre-clinical Cushing's syndrome, pheochromocytomas, adrenocortical carcinomas, myelolipomas, ganglioneuromas, metastatic tumors and cysts [9-13]. Adrenal incidentalomas are usually asymptomatic, however many of these lesions may secrete weak precursor hormones or active hormones in insufficient amounts to cause clinically apparent disease. There are reports of several cases of adrenal incidentaloma who had no clinical evidence of Cushing's syndrome and normal basal steroid hormone secretion, but non-suppressible serum cortisol after dexamethasone administration. These cases were regarded as "pre-clinical Cushing's syndrome" [14-16] and many of them treated by adrenalectomy, which restored a normal cortisol suppression to dexamethasone.

To study the differences between non-functioning adenomas and those associated with pre-clinical Cushing's syndrome, we performed a comprehensive analysis of serum steroid hormone profiles in patients with such adrenocortical tumors using a combination method of high-performance liquid chromatography(HPLC) and radioimmunoassay(RIA) [17].

2. Materials and Methods

2.1. Subjects

We studied 22 patients with incidentally discovered adrenal masses. These lesions were seen on abdominal imaging and were diagnosed clinically and/or pathologically. From our adrenal incidentaloma cases, we selected for inclusion only patients with adrenocortical adenomas. Fifteen non-functioning adenoma cases (7 males and 8 females, mean age 59.3±13.7 yr) and seven pre-clinical Cushing's syndrome cases (2 males and 5 females, mean age 54.7±12.4 yr) were studied (Table 1). As the control group, sixteen healthy adults(8 males and 8 females, mean age 51.0±7.4 yr) were used.

Non-functioning	No	Age(yr)	Sex	Side	Size(mm)	BP(mmHg)	Operation
	1	71	M	right	25x15	150-80	(+)
	2	83	M	right	30x20	160-84	
	3	74	M	right	30x20	146-86	(+)
	4	61	M	left	8x8	130-70	
	5	49	M	right	22x13	134-78	
	6	42	M	left	18x12	138-80	
	7	53	M	left	19x17	142-82	
	8	87	F	right	32x15	100-80	
	9	57	F	left	12x12	140-80	
	10	53	F	right	22x18	104-60	(+)
	11	55	F	right	12x9	120-76	
	12	49	F	right	26x19	124-80	(+)
	13	42	F	left	21x18	144-84	
	14	56	F	left	32x21	128-72	(+)
	15	57	F	left	25x29	146-84	
Pre-clinical	1	41	M	left	15x10	146-88	
Cushing	2	54	M	left	16x11	148-84	
	3	61	F	right	18x18	130-84	(+)
	4	60	F	right	17x13	160-70	(+)
	5	59	F	right	17x15	130-82	
	6	72	F	right	18x12	156-86	(+)
	7	36	F	left	22x18	124-80	(+)

Table 1. Clinical features in cases of non-functioning adenoma and pre-clinical Cushing’s syndrome

2.2. Hormonal measurements

Blood samples were obtained from patients with adrenal masses and normal subjects between 8 and 9 a.m. after an overnight fast. Serum levels of 10 steroid hormones [pregnenolone (Preg), progesterone (P), deoxycorticosterone (DOC), corticosterone (B), 17-hydroxypregnenolone (17-OH-Preg), 17-hydroxyprogesterone (17-OHP), 11-deoxycortisol (S), cortisol (F), dehydroepiandrosterone (DHEA), Δ 4androstenedione (Δ 4A)] were determined by an HPLC/RIA method as previously described [17].

2.3. Statistical analyses

We analyzed product/precursor ratios as indices of the relative activities of adrenal steroidogenic enzymes. Data are shown as mean $\pm$ SD. Variables were compared by the Mann-Whitney U test with Bonferroni correction. P values less than 0.05 were considered statistically significant.

2.4. Results

Eleven cases of non-functioning adenomas (Group I) had normal levels in DHEA and Δ 4A compared to normal controls by simultaneous analysis (Fig.1). In 4 cases of non-functioning adenomas (Group II), however, serum levels of DHEA and Δ 4A were significantly decreased compared to normal controls by simultaneous analysis (Fig.2). Other steroid hormone levels were normal in all cases of non-functioning adenomas. Therefore, we divided these patients into two groups based on the relative activities of adrenal steroidogenic enzymes by calculating product/precursor ratios. In pre-clinical Cushing's syndrome, serum levels of DHEA and Δ 4A were significantly decreased compared to normal control values (Fig.3) and these results were similar to values in Group II patients.

17,20-Lyase activities assessed by DHEA/17-OH-Preg and Δ 4A/17-OHP ratios were significantly decreased in Group II of non-functioning adenoma and pre-clinical Cushing's syndrome compared to Group I patients and normal controls (Fig.4). 17-Hydroxylase activities assessed by 17-OH-Preg/Preg and 17-OHP/P ratios (Fig.5), 3 β -hydroxysteroid dehydrogenase activities assessed by P/Preg, 17-OHP/17-OH-Preg and Δ 4A/DHEA ratios (Fig.6), 21-hydroxylase activities assessed by DOC/P and S/17-OHP ratios (Fig.7) and 11 β -hydroxylase activities assessed by B/DOC and F/S ratios (Fig.8) had the same levels in all cases of non-functioning adenomas and pre-clinical Cushing's syndrome compared to normal control values.

Figure 1. Serum steroid hormone profiles in non-functioning adrenocortical adenoma Group I

Non-functioning adrenocortical adenoma (Group II)

Cholesterol

↓

Pregnenolone	→	17 α -hydroxypregnenolone	→	Dehydroepiandrosterone
0.83 $\pm$ 0.18 ng/ml		1.10 $\pm$ 0.31 ng/ml		1.5 $\pm$ 0.8 ng/ml*
(0.74 $\pm$ 0.14)		(0.99 $\pm$ 0.15)		(3.1 $\pm$ 0.5)

↓

↓

↓

Progesterone	→	17 α -hydroxyprogesterone	→	Δ 4androstenedione
0.22 $\pm$ 0.18 ng/ml		1.08 $\pm$ 0.19 ng/ml		0.32 $\pm$ 0.13 ng/ml*
(0.17 $\pm$ 0.03)		(0.94 $\pm$ 0.20)		(0.98 $\pm$ 0.29)

↓

↓

Deoxycorticosterone	11-deoxycortisol
0.072 $\pm$ 0.021 ng/ml	0.82 $\pm$ 0.18 ng/ml
(0.060 $\pm$ 0.013)	(0.76 $\pm$ 0.12)

↓

↓

Corticosterone	Cortisol
1.43 $\pm$ 0.29 ng/ml	9.8 $\pm$ 2.2 μ g/dl
(1.52 $\pm$ 0.37)	(9.7 $\pm$ 1.3)

↓

18-hydroxycorticosterone () : Normal control

↓

*p<0.05 vs Normal control

Aldosterone

Figure 2. Serum steroid hormone profiles in non-functioning adrenocortical adenoma Group II

Pre-clinical Cushing's syndrome

Cholesterol		
↓		
Pregnenolone	→ 17 α -hydroxypregnenolone	→ Dehydroepiandrosterone
0.79 $\pm$ 0.19 ng/ml	1.02 $\pm$ 0.27 ng/ml	1.7 $\pm$ 0.4 ng/ml*
(0.74 $\pm$ 0.14)	(0.99 $\pm$ 0.15)	(3.1 $\pm$ 0.5)
↓	↓	↓
Progesterone	→ 17 α -hydroxyprogesterone	→ Δ 4androstenedione
0.19 $\pm$ 0.08 ng/ml	1.12 $\pm$ 0.28 ng/ml	0.30 $\pm$ 0.14 ng/ml*
(0.17 $\pm$ 0.03)	(0.94 $\pm$ 0.20)	(0.98 $\pm$ 0.29)
↓	↓	
Deoxycorticosterone	11-deoxycortisol	
0.061 $\pm$ 0.013 ng/ml	0.90 $\pm$ 0.27 ng/ml	
(0.060 $\pm$ 0.013)	(0.76 $\pm$ 0.12)	
↓	↓	
Corticosterone	Cortisol	
1.49 $\pm$ 0.36 ng/ml	10.1 $\pm$ 1.9 μ g/dl	
(1.52 $\pm$ 0.37)	(9.7 $\pm$ 1.3)	
↓		
18-hydroxycorticosterone		() : Normal control
↓		*p<0.05 vs Normal control
Aldosterone		

Figure 3. Serum steroid hormone profiles in pre-clinical Cushing's syndrome

Figure 4. 17,20-Lyase activities

Figure 5. 17-Hydroxylase activities

Figure 6. 3β - Hydroxysteroid dehydrogenase activities

Figure 7. 21- Hydroxylase activities

Figure 8. 11 β -Hydroxylase activities

3. Discussion

We examined serum steroid hormone profiles of patients with non-functioning adrenocortical adenomas and pre-clinical Cushing's syndrome. We also assessed the steroidogenic enzymatic activities of these patients by analysis of the product/precursor ratios of C21 and C19 steroid hormones in the steroidogenic pathway. Patients with pre-clinical Cushing's syndrome and a subgroup of patients with non-functioning adrenocortical adenomas had distinctly decreased adrenal androgen secretion compared with a suppressed 17,20-lyase activity. The remaining patients with non-functioning adrenocortical adenomas had completely normal adrenal androgen values and 17,20-lyase activity.

Previous reports have shown an exaggerated response of 17-hydroxyprogesterone after ACTH stimulation in some patients with incidentally detected adrenal incidentalomas [18,19]. This response has been explained by existing 21-hydroxylase deficiency which may be a pathogenetic factor in the development of adrenal tumors. Macronodular adrenal hyperplasia is a frequent finding in patients with classic congenital adrenal hyperplasia, however, also about half of the heterozygous carriers in families with congenital adrenal hyperplasia were reported to have uni- or bilateral adrenal nodules on abdominal computed tomography in consecutive 4-mm scans [19]. These investigators concluded that mild 21-hydroxylase deficiency is associated with the formation of adrenal incidentaloma. Other investigators have reported that the ACTH-stimulated 17-hydroxyprogesterone levels were abnormally increased in more than 50% of patients with non-hyperfunctioning adrenal adenomas [20]. After unilateral adrenalectomy, this hormonal abnormality disappeared in most patients with adrenal tumors. The ACTH-stimulated 17-hydroxyprogesterone levels signifi-

cantly correlated with the size of the tumors. These results indicate that tumors themselves may be responsible for the increased ACTH-stimulated 17-hydroxyprogesterone levels in patients with non-hyperfunctioning adrenal adenomas.

In this study, we demonstrated that serum DHEA and $\Delta 4A$ levels were significantly decreased in some patients with non-functioning adrenocortical adenomas (Group II) and all those with pre-clinical Cushing's syndrome. 17,20-Lyase activities were significantly decreased in Group II of non-functioning adenoma and pre-clinical Cushing's syndrome compared to Group I patients and normal controls. We found no clinical or pathological differences between Group I patients and Group II patients who had normal adrenal androgen concentrations. We have no explanation as to why there was difference between these groups. Since serum steroid hormone profiles had a similar pattern between some non-functioning adrenocortical adenomas and adenomas of patients with pre-clinical Cushing's syndrome, we suggest that the former may in time progress into pre-clinical and clinical Cushing's syndrome. There are limitations in this study such as small sample size and no tissue investigation. In the future the activity of steroidogenesis enzyme should be measured in the specimens from adrenal tissues.

Author details

Hajime Ueshiba*

Address all correspondence to: ueshiba@med.toho-u.ac.jp

Department of Internal Medicine, Toho University School of Medicine, Japan

References

- [1] Belldgrun, A., Hussain, S., Seltzer, S. E., Loughlin, K. R., Gittes, R. F., & Richie, J. P. (1986). Incidentally discovered mass of the adrenal gland. *Surg Gynecol Obstet*, 163, 203-208.
- [2] Glazer, H. S., Weyman, P. J., Sagel, S. S., Levitt, R. G., & Mc Clennan, B. L. (1982). Nonfunctioning adrenal masses: incidental discovery on computed tomography. *Am J Roentgenol*, 139, 81-85.
- [3] Abecassis, M., Mc Loughlin, M. J., Langer, B., & Kudlow, J. E. (1985). Serendipitous adrenal masses: prevalence, significance, and management. *Am J Surg*, 149, 783-788.
- [4] Herrera, M. F., Grant, C. S., van Heerden, J. A., Sheedy, P. F., & Ilstrup, D. M. (1991). Incidentally discovered adrenal tumors: an institutional perspective. *Surgery*, 110, 1014-1021.

- [5] Peppercorn, P. D., Grossman, A. B., & Rezneck, R. H. (1988). Imaging of incidentally discovered adrenal masses. *Clin Endocrinol*, 48, 379-388.
- [6] Copeland, P. (1983). The incidentally discovered adrenal mass. *Ann Intern Med*, 98, 940-945.
- [7] Hedeland, H., Ostberg, G., & Hokfelt, B. (1968). On the prevalence of adrenocortical adenomas in an autopsy material in relation to hypertension and diabetes. *Acta Med Scand*, 184, 211-214.
- [8] Granger, P., & Genest, J. (1970). Autopsy study of adrenal glands in unselected normotensive and hypertensive patients. *Can Med Assoc J*, 103, 34-36.
- [9] Kloos, R. T., Gross, M. D., Francis, I. R., Korobkin, M., & Shapiro, B. (1995). Incidentally discovered adrenal masses. *Endo Rev*, 16, 460-484.
- [10] Bencsik, Z., Szabolcs, I., Kovacs, Z., Ferencz, A., Voros, A., Kaszas, I., Bor, K., Gonczi, J., Goth, M., Kovacs, L., Dohan, O., & Szilagyi, G. (1996). Low dehydroepiandrosterone sulfate (DHEA-S) level is not a good predictor of hormonal activity in nonselected patients with incidentally detected adrenal tumors. *J Clin Endocrinol Metab*, 81, 1726-1729.
- [11] Kasperlik-Zatuska, A., Rostonska, E., Stowinska-Srzednicka, J., Migdalska, B., Jeske, W., Makowska, A., & Snochowska, H. (1997). Incidentally discovered adrenal mass (incidentaloma): investigation and management of 208 patients. *Clin Endocrinol*, 46, 29-37.
- [12] Barzon, L., Scaroni, C., Sonino, N., Fallo, F., Paoletta, A., & Boscaro, M. (1999). Risk factors and long-term follow-up of adrenal incidentalomas. *J Clin Endocrinol Metab*, 84, 520-526.
- [13] Mantero, F., Terzolo, M., Arnaldi, G., Osella, G., Masini, A. M., Ali, A., Giovagnetti, M., Opocher, G., & Angeli, A. (2000). A survey on adrenal incidentaloma in Italy. *J Clin Endocrinol Metab*, 85, 637-644.
- [14] Bogner, U., Eggens, U., Hensen, J., & Oelkers, W. (1986). Incidentally discovered ACTH-dependent adrenal adenoma presenting as "pre-Cushing's syndrome". *Acta Endocrinol*, 111, 89-92.
- [15] Rosen, H. N., & Swartz, S. L. (1992). Subtle glucocorticoid excess in patients with adrenal incidentaloma. *Am J Med*, 92, 213-216.
- [16] Renicke, M., Nieke, J., Krestin, G. P., Saeger, W., Allolio, B., & Winkelmann, W. (1992). Preclinical Cushing's syndrome in adrenal "incidentaloma": comparison with adrenal Cushing's syndrome. *J Clin Endocrinol Metab*, 75, 826-832.
- [17] Ueshiba, H., Segawa, M., Hayashi, T., Miyachi, Y., & Irie, M. (1991). Serum profiles of steroid hormones in patients with Cushing's syndrome determined by a new HPLC/RIA method. *Clin Chem*, 37, 1329-1333.

- [18] Seppel, T., & Schlaghecke, R. (1994). Augmented 17-alpha-hydroxyprogesterone response to ACTH stimulation as evidence of decreased 21-hydroxylase activity in patients with incidentally discovered adrenal incidentalomas. *Clin Endo*, 41, 445-451.
- [19] Jaresch, S., Kornely, E., Kley, H. K., & Schlaghecke, R. (1992). Adrenal incidentaloma and patients with homozygous or heterozygous congenital adrenal hyperplasia. *J Clin Endocrinol Metab*, 74, 685-689.
- [20] Toth, M., Racz, K., Adleff, V., Futo, L., Jakab, C., Karlinger, K., Kiss, R., & Glaz, E. (2000). Comparative analysis of plasma 17-hydroxyprogesterone and cortisol responses to ACTH in patients with various adrenal tumors before and after unilateral adrenalectomy. *J Endocrinol Invest*, 23, 287-294.