

# We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

185,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index  
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?  
Contact [book.department@intechopen.com](mailto:book.department@intechopen.com)

Numbers displayed above are based on latest data collected.  
For more information visit [www.intechopen.com](http://www.intechopen.com)



---

# Mutational Analysis of Effectors Encoded by Monopartite Begomoviruses and Their Satellites

---

Muhammad Shafiq Shahid, Pradeep Sharma and Masato Ikegami

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/54518>

---

## 1. Introduction

The geminiviruses are plant-infecting viruses with genomes consisting of circular, single-stranded DNA (ssDNA) geminate particles [86]. Members of the family *Geminiviridae* have been grouped into four genera (*Begomovirus*, *Curtovirus*, *Mastrevirus* and *Topocuvirus*) based on genome organization, host range and insect vector [29, 87]. The majority of geminiviruses belong to the genus *Begomovirus*, are transmitted by whiteflies (*Bemisia tabaci*: Gennadius), and infect dicotyledonous plant species [85]. The monopartite begomovirus genome is ~2.8 kb nucleotides in length and encode genes both in complementary and virion sense from a non-coding intergenic region that contains promoter sequences and the origin (*ori*) of virion-strand DNA replication. The *ori* consists of a predicted hairpin structure that contains the absolutely conserved (for geminiviruses) nonanucleotide (TAATATTAC) loop sequence and repeated motifs upstream known as iterons.

## 2. Functions of effectors encoded by monopartite begomoviruses

### 2.1. Complementary-sense

The complementary-sense strand encodes the Rep proteins, also known as C1, AC1 and AL1, is a multifunctional protein and the only viral protein absolutely required for virus replication. Rep is encoded on the complementary sense strand (Fig. 1 DNA A). This protein is involved in several biological processes: initiation and termination of rolling circle replication (RCR) by nicking and religating the replication origin of viral DNA [45] and repression of its own gene transcription [19]. The Rep proteins of geminiviruses are closely related and show substantial sequence conservation. Four functional domains have been delineated for begomovirus Rep : the N-terminal domain (amino acids 1 to 120), which is involved in initiation by geminiviruses [63], AC1 protein initiates rolling circle replication

by a site-specific cleavage within the loop of the conserved nonamer sequence, TAATATTAC [30]. The AC1 protein binding site is located between the TATA box and the transcription start site for the *Rep* gene and acts as the origin recognition sequence and as a negatively regulatory element for *AC1* gene transcription [19], the oligomerization domain (121 to 180 aa), leading to interactions with itself and with host factors [28]. The AC1 protein alone can initiate RCR without requiring other accessory viral factors [34]. AC1 protein also has DNA helicase activity which depends upon the oligomeric state of the protein [14].

The transcriptional activator protein (TrAP); is also known as AC2, C2 and AL2. AC2 is a ~15-KD a transcriptional activator protein unique to begomoviruses because it is absent in mastreviruses and a related protein in curtoviruses, AC2 protein, seems to play a different role. In mastreviruses, AC1 protein provide the functions of AC2 [51]. TrAP is necessary for transactivation of late genes [90]. Recently, several researchers have shown that the AC2 gene of *Cabbage leaf curl virus* (CaLCuV) activates the CP promoter in mesophyll and acts to derepress the promoter in vascular tissue, similar to that observed for TGMV [44]. Further, since AC2 1-100 is as effective a suppressor as the full-length AC2 protein, activation and silencing suppression appear to be independent activities. For example Gopal et al. [26] showed that AC2 of *Bhindi yellow vein mosaic virus* (BYVMV) is involved in transactivation and only mildly in suppression of gene silencing of monopartite begomoviruses viruses and not in transmission.

The replication enhancer protein (REn); also named as AC3/AL3. AC3 is a ~16 KD a protein in curtoviruses and in begomoviruses. The AC3 protein greatly enhances viral DNA accumulation of curtoviruses and begomoviruses [22, 92] by interacting with Rep [81]. Experimental observations suggested that AC3 protein might increase the affinity of Rep for the origin. Complementation studies revealed that AC3 could act on heterologous viruses [93].

The C4 protein, for which the function remains unclear but for some viruses is a pathogenicity determinant and a suppressor of PTGS [73]. AC4 is highly variable among begomoviruses, which is expressed from an open reading frame (ORF) embedded in the Rep ORF.

## 2.2. Virion-sense

The virion-sense strand encodes the genes required for insect transmission and movement in plants, coat protein (CP) and V2 protein. Monopartite begomovirus capsids are composed of a single CP, encoded by the *V1* gene or (also known as AV1), depending on the geminivirus [107]. For monopartite geminiviruses, CP is essential for systemic spread through the plant [12]. Besides the encapsidation function, CP is also required for transmission of the virus between the plants. The CP of the monopartite begomoviruses facilitates the transfer of infecting viral DNA into the host cell nucleus and is essential for systemic virus movement [5, 46, 50, 104]. The CP also determines the vector specificity [10, 32, 33] and protects the viral ssDNA from degradation during transmission by the insect vector [3], or mechanical inoculation [24].

In contrast to New World (NW) begomoviruses, Old World (OW) begomoviruses have AV2/V2 and this is involved in the movement of monopartite viruses. A recent report shown that the V2 (a homolog of AV2) of a monopartite begomovirus is involved in overcoming host defenses mediated by post-transcriptional gene silencing as well as in movement [114, 115]. V2 targets a step in the RNA silencing pathway which is subsequent to the Dicer-mediated cleavage of dsRNA [109, 70].

### 3. Role of effectors encoded by satellites


#### 3.1. Betasatellite

Recently, the majority of the begomoviruses originating from the OW have been shown to be monopartite and to associate with a class of ssDNA satellites known as betasatellites (earlier known as DNA $\beta$ ) [9]. Betasatellites are approximately half the size of their helper begomoviruses (approx.1.4 kb) and are required by the helper virus to induce typical disease symptoms in their original hosts. The success of begomovirus-betasatellite disease complexes appears to be due to the promiscuous nature of betasatellites that allows them to be *trans*-replicated by several distinct begomoviruses [53, 59]. These begomovirus-betasatellite disease complexes are widespread throughout OW and outnumber bipartite begomoviruses whereas in the NW only bipartite begomoviruses are native. There have been recent reports which showed that betasatellite can complement the function of DNA B, suggesting that the betasatellite may provide movement functions to its helper begomovirus [74]. Betasatellite can be associated with distinct begomoviruses and it can interact and make new complex with diverse monopartite begomoviruses [110-112].

*Tomato leaf curl virus* (ToLCV), originating from Australia, was shown to be associated with a single-stranded DNA satellite molecule [18]. The ToLCV satellite (ToLCV-sat) is approximately 682 nt in length and sequence unrelated to ToLCV and it depends on ToLCV for replication and encapsidation. It has no discernable effects on viral replication or symptoms caused by ToLCV. Betasatellites have three structural features: a approx.115 bp highly conserved region,  $\beta C1$  gene and a region rich in adenine, [76, 108] (Fig. 1 betasatellite). This gene has the capacity to encode a 13-14-kDa protein comprising 118 amino acids, although some betasatellites have additional N-terminal amino acids [79, 108]. Recently it has been shown that betasatellite to be pathogenicity determinant and suppressor of RNA silencing [16, 66]. It also induced abnormal cell division in *N. benthamiana* [17]. Betasatellites do not contain the iterons of their helper begomoviruses, although betasatellite clearly must possess sequences that are recognized by the begomovirus-encoded Rep in order to allow transreplication of the betasatellite [76].

All the reported betasatellites [54] or defective betasatellites (half size of wild type betasatellite) [7] contain the A-rich region, the A-rich region may play biological role in betasatellites [95]. A-rich region is not required for *trans*-replication of betasatellite and not related with encapsidation also. However, the A-rich region deleted mutant caused milder symptom [95]. The begomovirus accumulates to normal levels in *Ageratum* in the presence

of betasatellite suggesting that the satellite functions either by facilitating the replication or movement of the begomovirus or by suppressing a host defense mechanism such as gene silencing. Recently it has been shown that a betasatellite can override the AC4 pathogenicity phenotype of TLCV and it can complement the function of DNA B [73]. Despite its importance to the disease phenotype, there is still no information available concerning even the most fundamental properties of the satellite.



**Figure 1.** Genome organization of monopartite begomoviruses-satellite complex. DNA-A (encoding replication-associated protein [Rep], coat protein [CP], replication enhancer protein [REn], transcriptional activator protein [TrAP] and proteins possibly involved in virus movement [AV2], pathogenicity determinant and a suppressor of RNA silencing [AC4], viral genome replication [AC5]) Alphasatellites are self-replicating molecules encoding their own Rep. Betasatellites are dependent on their helper viruses for their replication and encode a single protein,  $\beta$ C1, which upregulate replication of helper virus and suppress host defense. Both satellites have an A-rich region and in addition to this betasatellites have a region of sequence conserved between all examples known as the satellite conserved region (SCR).

In *Arabidopsis*, these pathways are affected by the DICER- like proteins (DCL1, DCL2, and DCL3) that are nuclear localized and are required for miRNA and siRNA biogenesis. Thus,  $\beta$ C1 protein may affect the activity of the DICER-like proteins in plants during nuclear activities that function in silencing suppression. The other possibilities are that  $\beta$ C1 protein could down-regulate transcription of a host protein that acts in the PTGS pathway in the cytoplasm or that  $\beta$ C1 protein could activate transcription of a host PTGS inhibitor [15].

#### 4. Alphasatellites

Many begomovirus betasatellite complexes are also associated with a third ssDNA component for which the collective term alphasatellite (earlier known as DNA 1; R.W. Briddon, manuscript in preparation). However, alphasatellites are dispensable for virus infection and appear to play no significant role in the etiology of the diseases with which they are associated [56]. Alphasatellite components are satellite-like, circular ssDNA molecules approx.1375 nucleotides in length (Fig.1 alphasatellite). They encode a single gene, a rolling circle replication initiator protein (Rep), and are capable of autonomous replication in plant cells. Closely related to the replication associated protein encoding components of nanoviruses (a second family of plant infecting ssDNA viruses), from which they are believed to have evolved, they require a helper begomovirus for movement within and between plants [56, 80].

Several alphasatellites are capable of replicating and systemically infecting their plant host in the presence of a helper begomovirus without a visible effect on symptom development or virulence [6, 40]. However AYWVSGA a different type of 'DNA-2'-class alphasatellite that ameliorates symptom severity in an infected host and also capable of reducing virulence and the relative accumulation of its associated Tomato leaf curl betasatellite (ToLCB) [1]. Alphasatellites have been acquired by helper begomoviruses to restrain virulence to achieve increased viral fitness [76, 105].

Recently, two 'DNA-1-type' alphasatellites *Gossypium mustelinum* symptomless alphasatellite (GMusSLA) and *Gossypium darwinii* symptomless alphasatellite (GDarSLA) phylogenetically divergent from the DNA-2-type alphasatellite have each been shown to attenuate symptoms caused by their helper begomovirus [60]. However [35] hypothesize that symptom attenuation and a relative reduction in betasatellite accumulation might result from DNA-2-mediated modulation of betasatellite activity. Possibly alphasatellite modulates begomovirus-betasatellite pathogenicity by interfering with  $\beta$ C1, a key virulence factor [8]. Also alphasatellite rep can interact with C4 of CLCuRaV that might be providing an additional possible mechanism for symptom amelioration by alphasatellites. Furthermore alpha-Rep down regulate betasatellite replication (In the field), and thus down-regulation of the manifestation of the pathogenicity determinant  $\beta$ C1 [60], moreover alpha-Rep proteins GMusSLA and GDarSLA can act as a strong suppression of posttranscriptional gene silencing (PTGS) [60].

#### 5. Post-transcriptional gene silencing (PTGS)

Post-transcriptional gene silencing (PTGS) which is initiated by double stranded RNA (dsRNA) is common in plant-virus interactions and is an evolutionarily conserved mechanism that protects host cells against invasive nucleic acids, such as viruses, transposons and transgenes [100]. As a counter to this host defense, most plant viruses encode proteins which act as suppressors of PTGS [71]. Viral suppressors of PTGS interfere with various steps of this pathway including initiation, maintenance or systemic silencing which are mainly downstream of dsRNA production [52, 57].



RNA silencing in plants operates as an antiviral defense response; to establish infection, viruses must suppress RNA silencing by the host [100]. Begomoviruses have been shown to induce PTGS in infected plants by producing virus specific siRNAs (21, 22 and 24 nt) [97]. To counteract this host defence, geminiviruses encode RNA silencing suppressors [4]. However, depending on each intrinsic virus and its interaction with plant host factors, the efficacy of virus-induced PTGS may vary [99]. At least three RNA-silencing suppressors have been reported in TYLCD-associated or related begomoviruses. Thus, the V2 protein of TYLCV functions as an RNA-silencing suppressor; it counteracts the innate immune response of the host plant by interacting with SISGS3, the tomato homolog of the Arabidopsis SGS3 protein involved in the RNA-silencing pathway. The TrAP protein of the related monopartite begomovirus, *Tomato yellow leaf curl China virus* (TYLCCNV) is also involved in suppression of RNA silencing [98], probably by activating transcription of host genes that control silencing [97]. The C4 protein of the monopartite begomoviruses ToLCV, *Ageratum yellow vein virus* (AYVV), and *Bhendi yellow vein mosaic virus* (BiYVMV) also have the ability to suppress RNA silencing [26, 84].

## 6. Mutagenesis of effectors encoded by monopartite begomoviruses

Little is known about gene function in monopartite begomoviruses. However, gene function has been studied extensively in other types of geminiviruses which share organization and nucleotide sequence similarities with TYLCV. Mutational analysis of few monopartite begomoviruses like TYLCV define similarities and differences between this single component geminivirus and bipartite geminiviruses in functions essential for systemic spread and infectivity [103]. The CP appeared to be required for systemic movement of TYLCV in *N. benthamiana* and tomato, consistent with those of mutation analyses obtained with other monopartite geminiviruses such as MSV, BCTV, and TLCV [5, 12, 69, 61] have shown that *Tomato yellow leaf curl virus*-Sardinia (TYLCV-Sar) C2 can form stable complexes with ssDNA (and less preferably with dsDNA) and that the binding is sequence nonspecific. AC2 of TYLCV have also been involved in the activation of other viral genes and was considered as a transcriptional activator [91]. However, transcription factors usually show high sequence specificity.

TYLCV ORF V1 truncated either 133 nt upstream or 19 nt downstream of the initiation codon of ORF V2 would altered the viral DNA forms, it suggested that the V1 protein may participate in the switch from dsDNA to ssDNA synthesis. Indeed, interaction between V1 and the CP has already been proposed, in view of the concerted evolution of these two protein sequences following a geo-graphical gradient of similarity [39], and the synergistic reduction in ssDNA levels of a TYLCV V1-V2 double mutant compared to single mutants [69]. Although TYLCV V1 mutants did not greatly overproduce dsDNA, the similarity of phenotype between BCTV V2 and TYLCV V1 mutants may indicated that the two corresponding gene products serve a related function. It has shown [69] (Table 1) has shown that disruption of the V1 gene in the monopartite Australian isolate of TLCV did not affect its ability to spread in tomato, although the infection was asymptomatic and the DNA levels reduced.

Virus	Accession number	Mutated ORF	Mutation Position	Type of mutation	References
TLCV-[AU]	AF084006	V1	N-terminal/ <i>Bam</i> HI	frameshift	Rigden et al., 1993
		V2	N-terminal/ <i>Bgl</i> II	frameshift	
		V/V2	N-terminal/ <i>Bgl</i> II/ <i>Bam</i> HI	frameshift	
		V/V2	C-terminal/ <i>Bgl</i> II/ <i>Bam</i> HI	deletion	
		V/V2	C-terminal/ <i>Bam</i> HI/ <i>Bgl</i> II	inversion	
TLCV-[AU]	AF084006	C4	N-terminal (at 2457&2463)	deletion	Rigden et al., 1993
		C4	N-terminal (at 2457&2463)	reversion	
TYLCV-Sar[ES:Psp95:93]	Z25751	C4	C>T at 2432	point	Jupin et al., 1994
TYLCV-Sar[ES:Psp95:93]	Z25751	C4	C>T at 2423		
		C2	$\Delta$ CC2>31	deletion	Noris et al., 1996
		C2	$\Delta$ NC2>33	deletion	
TYLCV-Sar[FR:98]	X61153	C2	$\Delta$ C2>33-104	deletion	Wartig et al.,1997
		C2	1523+GATC	frameshift	
		V1	156+GATC	frameshift	
		V1	324C>T	stop	
		V2	748>CTAG	stop	
TYLCV-Sic[IT:pSic36:95]	Z28390	CP	H134	substitution	Noris et al., 1998
TYLCV-Sar[ES:Psp95:93]	Z25751	RepC1 CP	L198 Q134H	substitution site-directed	Noris et al., 1998
TYLCV-DO[DO:99]	AF024715	CP	$\Delta$ NCP>180	deletion	Rojas et al., 2001
AYVV-[SG:pHN419:97]	X74516	CP	$\Delta$ CCP>150	deletion	
		V1	$\Delta$ NV1>63	deletion	
		V1	$\Delta$ CV1>84	deletion	
		C4	$\Delta$ NC4>54	deletion	
		C4	$\Delta$ CC4>60	deletion	
		C4	A>T at 2419 C4mut	substitution	
					Saunders et al., 2004



Virus	Accession number	Mutated ORF	Mutation Position	Type of mutation	References
PaLCuV-[PK:02]	AJ436992	V2	N-terminal (1-32)	deletion	Mubin et al., 2010
		V2	N-terminal (1-60)	deletion	
		V2	C-terminal (130-149)	deletion	
		V2	C-terminal (111-149)	deletion	
		V2	C-terminal (101-149)	deletion	
		V2	C-terminal (91-149)	deletion	
		V2	C-terminal (71-149)	deletion	
		V2	C-terminal (30-149)	deletion	
TYLCV-IL[IL:89]	X15656	CP	Lys-Thr CPmut3	substitution	Yaakov et al., 2011
		CP	Arg-Pro CPmut4	substitution	
TYLCV-Mld[ES:72:97]	AF071228	CP	Arg-Leu CPmut19	substitution	Tomas et al., 2011
		C4	C>G at 9 C4mut	substitution	
ToLCJV-A[ID]	AB100304	CP	CPΔ191-257	deletion	Sharma et al., 2009
		CP	CPΔ1-190	deletion	
		CP	CPΔ31-257	deletion	
		CP	CPΔ1-30/Δ50-257	deletion	
		CP	CPΔ1-30/Δ191-257	deletion	
		CP	CPΔ1-31	deletion	
		CP	CPΔ16-20	deletion	
		CP	CPΔ1-190/Δ245-250	deletion	
		CP	CPΔ1-30/Δ62-257	deletion	
ToLCJV-A[ID]	AB100304	V2	N-terminal (58aa)		Sharma et al., 2010
		V2	C-terminal (58aa)		
ToLCJV-A[ID]	AB100304	V2	pGEMV2ΔC		Sharma et al., 2011

**Table 1.** List of published studies reporting deletion mutants of monopartite begomoviruses

For example Noris et al. [62] suggested that the region of the CP between amino acids 129 and 134 is essential for both the correct assembly of virions and transmission by the insect vector. The genome of the SicRcv (infectious) had the same size as the original Sic DNA 9 (non-infectious) differed by only 2 nt. One change was at nt 205 (A instead of T in the plus strand), determining a CAC-to-CUC codon change in the RepC1 mRNA and an H198L

replacement in the RepC1 protein. The other mutation located at nt 708 (C instead of G), determining a CAG-to-CAC codon change in the CP mRNA and a Q134H replacement in the CP. This indicated that the Q134H mutation changed a viral DNA, only capable of replicating in single cells (Sic), into one that was systemically infectious, but not insect transmissible (SicRcv). Comparative analysis of Sic, SicRcv, and the hybrid genomes and showed that the mutation in the CP gene, not in the Rep gene, was responsible for restoring infectivity in SicRcv; however, it still did not result in a whitefly-transmissible TYLCV. In TYLCV-Sar, the two capsid protein alterations resulted in the same either non-infectious or non-transmissible phenotype. Mutants containing the combinations QQ, QH, and PH at positions 129 and 134 were infectious in plants, whereas those with PQ are not. The PQ mutants can replicate and accumulate CP and V2 protein in leaf discs, but appear unable to produce virus particles. Mutants having the PH combination at positions 129 and 134 infect plants and form apparently normal virions, but are not transmissible by whiteflies. Changing the amino acid at position 152 (D or E) does not influence the phenotype. Requirement of the CP for infection has been demonstrated previously [62] suggested that accurate particle assembly is also necessary. In fact, the PQ mutants, which are unable to assemble virions, accumulate CP in leaf discs, showing that its expression and stability were not altered. Another TYLCV protein, V2, for which a role in virus assembly has recently been, suggested [103].

For example Rojas et al. [70] has shown that C4, V1, and CP gene may function in TYLCV-DO movement. The CP localized to nuclei and nucleoli and was found to act as a nuclear shuttle, mediating the import and export of DNA [70]. It was consistent with results obtained for the TYLCV CP in heterologous experimental systems [43, 68]. Recently, Liu et al [49] also showed the same behavior for the CP of the monopartite mastrevirus, MSV. TYLCV CP was found to accumulate in the nucleolus and the absence of the N- and C-terminal CP mutants from the nucleolus implicates CP motifs in this localization. As the nucleolus is the site of rRNA synthesis and packaging of ribosomal proteins, it may also serve as the site of geminiviral replication/gene expression [70]. The TYLCV C4 targeted to the cell periphery and/or cell wall, consistent with a role in cell-to-cell movement of viral DNA [65, 75, 101].

Disruption of the AYVV C4 ORF (A>T at position 2419nt) alters the phenotype in agroinoculated *N. benthamiana* from upward leaf roll and vein swelling to downward leaf curl [79] (Table 2). Previously, [88] also found the identical functions of BCTV C4 ORF mutant in this host. The AYVV C4 proteins may perform partially redundant functions involving convergent pathways and the behavior of ACMV AC4 and TYLCCNV C4 is consistent with such a function [98].

For example Stanley and Latham [58] have shown that V2 protein of *Papaya leaf curl virus* (PaLCuV) is potentially involved in the elicitation of cell death response. The deletion mutants (having deletions of 32 and 60 amino acids, respectively, at the N-terminal end of V2) exhibited a systemic HR in *Nicotiana benthamiana* plants. While C-terminal end deletions of 60, 80 and 119 amino acids abolished the induction of HR, however 50 amino acids deletion

induced local necrosis, but not systemic. The mutants with 20 and 40 amino acids deletion produce HR both at the inoculated and in newly emerged leaves, although the systemic symptoms for the 40-amino-acid deletion mutant were delayed and were milder. The amino acid sequences between positions 92 and 101 are essential for the elicitation of HR, whereas those between 102 and 115 affect the timing and severity of the response. V2 of PaLCuV at amino acid positions 116 and 118 contains a conserved CxC. Mutations of this motif have been shown to abolish both the pathogenicity and suppressor of RNA silencing activities of the protein [64, 109]. Phosphorylation of MPs may also play a role in controlling the switch from viral replication to translation [36, 37]. Few earlier studies showed that for PaLCuV V2, deletion of sequences encompassing this motif abrogates the ability to induce HR [58].

The first 30 N-terminal amino acids of the TYLCV-IL CP are needed for nuclear import of the protein into the plant cell, suggesting the CP's involvement in nuclear shuttling of the virus genome [43]. This was confirmed by the finding of a strong interaction between the CP and the plant nuclear import receptor karyopherin  $\alpha 1$  (Kap  $\alpha 1$ ) [94]. The TYLCV CP has been found to interact with itself (CP-CP or homotypic interaction) which may be important for capsid assembly as it is made up solely of CP units serving as building blocks. Mutations in the TYLCV-IL V1 gene coding for the TYLCV-IL CP by replacing Lys with Thr, Arg with Pro, and Arg with Leu, according to the positions of amino acids mutated [31]. TYLCV CP mutated failed to interact with the w.t. CP, while the w.t. protein showed strong homotypic interaction. As the CP has been suggested to be a shuttle protein for the viral genome into the plant cell nucleus [43, 70], its interaction with the nuclear-transport mediator Kap  $\alpha 1$  is an important step and has been shown to occur at high affinity [94]. A mutation in the NLS domain, in particular at Arg19, disrupts the CP's interaction with proteins that are known to interact with the w.t. CP [106]. Earlier Sharma et al. [113-115] demonstrated by the constructed a series of single and double deletions into the coding sequence of *Tomato leaf curl Java virus* ToLCJAV-A[ID] CP and found that, amino acids (aa)16KVRRR20 in the N-terminal region of CP functioned as nuclear/nucleolar localization signals (NLSs). Further, the region from aa 52RKPR55 contained basic amino acid cluster was capable to redirect the CP to the nucleus. Deletion mutant analysis revealed that this property was attributed to a nuclear export signal (NES) sequence consisted of aa (245LKIRIY250) reside at C-terminal part of CP. Additionally ToLCJV V2 is a target of host defense responses. Deletion of 58 amino acids (aa) from the N-terminus did not affect the HR, suggesting that this region has no role in the HR, while deletion of 58 aa from the C-terminus of V2 abolished both the HR response and V2 silencing suppressor activity, suggesting that these sequences are required for the HR-like response and suppression of PTGS. He also demonstrated that ToLCJV V2 is a pathogenicity determinant that elicits an HR-like response. Further deletion analysis that fusion of Nterminal part of the V2, containing the nuclear export signals (NES), directed the accumulation of fluorescence towards the cell cytoplasm. Also V2 enhances the coat protein-mediated nuclear export of ToLCJV and is consistent with the model in which V2 mediates viral DNA export from the nucleus to the plasmodesmata.

Satellite	Accession number	region (nt.)	Type of mutation	References
AYVB-[SG:pBS-beta:99]	AJ252072	AT>TA at 547/ 548 $\beta$ C1mut1 G>T at 486 $\beta$ C1mut2		Saunders et al., 2004
TYLCCNB-[CN:Y10:01]	AJ421621	ATG>ATC (CIM-F) ACT>TGA (ACIM-S) ATG (2) >ATC (2) (CIM-B) GAA>TAG (CIM-T)	site-derected  site-derected site-derected  site-derected	Cui et al., 2004
TYLCCNB-[CN:Y10:01]	AJ421621	742-952	deletation	Xiaorong et al., 2004
TYLCCNB-[CN:Y10:01]	AJ421621	$\Delta$ C1 $\beta$	truncation	Qian and Zhou 2005
CLCuV $\beta$ -[PK:00]	AJ298903	195-484 504-596 586 ATG>ATA	site-derected stop frame-shift stop	Saeed et al., 2005
BYVMB-[IN:Muth:01]	AJ308425	51-140 $\Delta$ N $\beta$ C1 1-80 $\Delta$ C $\beta$ C1	deletation deletation	Kumar et al., 2006
TYLCV-satDNA-[AU:96]	U74627	$\Delta$ nt 35-146 (112nt)  $\Delta$ nt 146-296 (151nt) $\Delta$ nt 35-296 (262nt) $\Delta$ nt 296-420 (1251nt) $\Delta$ nt 296-492 (197nt) $\Delta$ nt 492-540 (49nt) $\Delta$ nt 540-641 (105nt)	deletation  deletation deletation deletation deletation deletation	Li et al., 2007
AYVJB-[ID:04]	AB162142	4>ATGtga	stop	Kont et al., 2007
AYVB-[SG:pBS-beta:99]	AJ252072	794-795 118-119 804-806 801-1047 1048-1051 1146-1147 1146-1150 1269-1271 1229-1234	deletation deletation deletation deletation deletation deletation deletation deletation	Saunders et al., 2008
TbCSVb-[CN:Y35:01]	AJ420318	DNA $\Delta$ C1 $\beta$		Qian et al., 2008

Satellite	Accession number	region (nt.)	Type of mutation	References
CLCuMA-[PK:2:99]	AJ132345	915-1117	deletion	Shahid et al., 2009
CLCuMuB-[PK:09]	FJ607041	Δ150-840	deletion	Nawaz-ul-Rehman et al., 2009
		Δ1130-116	deletion	
		Δ995-1095	deletion	
CLCuMB-[PK:00]	AJ298903	ΔC1β	deletion	Kharazmi et al., 2012
TYLCCNB-[CN:Y10:01]	AJ421621	N-terminal (NTG)	deletion	Cheng et al., 2011
		C-terminal (CTG)	deletion	
		βC1-ΔCTD	deletion	

**Table 2.** List of published studies reporting deletion mutants of DNA Satellites

## 7. Mutational analysis of effectors encoded by satellites

Betasatellite molecules have been associated with numerous monopartite begomoviruses in China, including *Tobacco curly shoot virus* (TbCSV) and TYLCCNV that infect tomato and tobacco field plants [108, 47]. TbCSB is not essential for infection but increases symptoms in some hosts [17, 47]. However in case of TYLCCNB which is essential for symptom induction, The βC1 gene of TYLCCNB is required for symptom induction but not for the replication of betasatellite. Also a mutated βC1 deleted is stably maintained in few hosts by TYLCCNV [17, 67]. TbCSB with the complete βC1 deleted (ΔβC1) returns to a size comparable to that of the intact betasatellite in few systemically infected *N. glutinosa*, *N. tabacum* Samsun and *P. hybrida* plants. The levels of accumulation of the size revertant betasatellite were similar to those of ΔβC1 in same hosts (*N. benthamiana* and *N. glutinosa*) plants showing size reversion of the betasatellite developed viral symptoms similar to those induced by TbCSV and DNAΔβC1 [67]. A βC1 gene frame-shift mutant of TYLCCNVB was unable to induce disease symptoms and consequently, did not play a role in silencing suppression [67]. The complete coding region of Y10βC1 (TYLCCNB), followed by N- and C-terminal deletion mutants showed multimerization mediated by amino acids between positions 60 and 100 [13]. Karyopherin-α, a transport receptor involved in nuclear import were reported to interact with the C-terminal sequences of BYVMB- βC1 [42]. A myristoylation-like motif (GMDVNE) positioned at the C-terminal of CLCuMB-βC1 (103 to 108aa) interacted with a ubiquitin-conjugating enzyme involved in targeting proteins for degradation by the 26S proteasome [20, 78] identified sequences on AYVB by deletion mutagenesis required for trans-replication by AYVV. βC1 of Cotton leaf curl Multan betasatellite (CLCuMuB) has been shown to have possible virus movement function [74]. Generally, sequences between the βC1 gene and the A-rich region are not essential for trans-replication by begomoviruses. Nevertheless, deletion of these sequences abolish the ability of the betasatellite to upregulate virus levels in plants and the symptoms expression [59]. For geminiviruses hairpin structure that contains the nonanucleotide sequence is an essential part of the virion sense origin of replication that is recognized and nicked (within



the nonanucleotide sequence) by Rep to initiate rolling-circle replication of the virion strand. Similarly, deletion of betasatellite sequences from 1130 to 116 that is conserved (between all betasatellites) stopped the betasatellite's ability to be trans-replicated and maintained by helper viruses both from OW (CLCuRaV) and New world and *Cabbage leaf curl virus* (CbLCuV). Trans-replication of CLCuMuB remained unaffected by deletion of the sequence between coordinates 995 and 1095 by CLCuRaV [59].

ToLCJAV alone can cause infection and displayed leaf curl symptoms. But, symptom expression of ToLCJAV in the presence of ToLCJAB is enhanced. In contrast, ToLCJAV and AYVB (mutated  $\beta$ C1) restored mild symptoms. It suggested that the  $\beta$ C1 protein was required for symptom induction and is a determinant of pathogenicity,  $\beta$ C1 protein expression in *N. benthamiana* plants and as a suppressor of PTGS [41].

For example Li et al. [47] have shown the deletion mutant of TYLCV sat-DNA (from 296-641nt) lacked the ability to replicate or replicated poorly by deleting of (region nt 35-296). Also sequence from nt 296-35 is to be essential for sat-DNA replication. The deletion of a 112 nt region downstream of the stem-loop from nt 35-146 and 151nt from 146-296 cannot effect on the replication of sat-DNA but reduced significantly. However, the deletion from nt 35-296 regions diminished sat-DNA replication these deletions loss of genomic sequences required for replication or due to changes in genome size. Heterologous non-viral DNA fragments can restore the wild-type 682 nt sat-DNA size and of replication when the replacement occurred in the region between nt 35 and 296. However, the sequence replacements in the region nt 35 to 296 of the sat-DNA improved the accumulation of sat-DNA considerably relative to the deleted constructs in this region. The sequence elements distributed within the entire sat-DNA molecule contribute to replication activity, but that sequence elements within the region from nt 35 to 296 are dispensable for replication.

For example Saeed et al. [72] used mutagenesis study of CLCuMB and tobacco was used as the host plant rather than cotton, the natural host of CLCuB. Few studies showed that it was symptomless when inoculated with *Cotton leaf curl Multan virus* (CLCuMV) alone but showed drastic symptoms when coinoculated with CLCuMB [9]. *Nicotiana benthamiana* showed a severe symptom on inoculated with CLCuMV with or without CLCuMB. Evidence for the involvement of the  $\beta$ C1 ORF in modulation of symptom expression also provided by [108] demonstrated few DNA- $\beta$  species associated with tomato and tobacco infecting begomoviruses and found that in-frame mutation of the  $\beta$ C1 initiation codon resulted in loss of symptom severity in *N. benthamiana*.

In recent studies Saunders et al. [79] have proved that disruption of the  $\beta$ C1 ORF prevented infection of the AYVB complex in *ageratum* and altered their phenotype in *N. benthamiana* to that produced by AYVV alone. For example Kumar et al. [42] tested the infectivity of two  $\beta$ C1 mutant constructs, first carrying a stop codon at amino acid position 41 and second with two stop codons at positions 9 and 41, and both resulted in loss of pathogenicity in tobacco plants on coinoculated with TLCV as helper virus. These mutation studies indicated that the  $\beta$ C1 ORF is involved in pathogenicity and that the expression of its N terminal 40 amino acids is not sufficient for its function.



Disruption of the  $\beta$ C1 ORF of AYVB by introducing an internal in-frame nonsense codon (G>T) did not prevent transreplication and systemic movement of the  $\beta$ C1 mutant by AYVV in lab host (*N.benthamiana*). The mutated  $\beta$ C1 removed the influence of the satellite on symptom development in this host and prevented symptomatic infection of ageratum. That suggested the  $\beta$ C1 protein is an important pathogenicity factor that plays an essential role in the proliferation of the AYVV-betasatellite complex in its real host. For example Saunders et al. [79] also shown that  $\beta$ C1 ORF initiation codon (AT) to a nonsense codon (TA) did not completely eliminate betasatellite activity. A similar mutation in the  $\beta$ C1 ORF of a satellite associated with TYLCCNB was shown previously [108]. The  $\beta$ C1 ORF encodes a pathogenicity determinant that suppressed a host defense mechanism [76].

For example [78] have demonstrated that the region of AYVB between the introduced nt 114 and 1047 sites is not required for betasatellite replication. This region includes the  $\beta$ C1 open reading frame (ORF), which encodes a gene essential for pathogenicity [79] and an A-rich region that may serve to maintain the size integrity of the satellite [76]. For example [78] found that the entire ORF is dispensable and is consistent with the findings of [67] for the betasatellite associated with TYLCCNV. In addition, removal of the A-rich region from TYLCCNB was tolerated, although the deletion mutant was associated with milder infection than those produced by the wild-type satellite [95]. In contrast, deletion of this region in AYVB did not affect the phenotype, at least in *N. benthamiana*. Maximum deletions within non coding regions of the begomovirus genome were not tolerated and the deletion mutants revert to wild type size by both intra-and intermolecular recombination during systemic movement [23, 25,]. For example Saunders et al. [78] also demonstrated removal of 361 nt of betasatellite representing 27% of the satellite and the region between nt 1047 and 1146 is important for betasatellite replication. It contained an inverted repeat flanking a sequence that is identical to the ToLCV iteron ToLCV sat-DNA [18]. Protein binding assays followed by mutagenesis have demonstrated that this motif in both ToLCV and sat-DNA represents a high affinity Rep binding site, although it is not required for replication of either the begomovirus or its satellite [48]. Saunders and associates [80] found that the region between nt 1146-1229 and sequences across the nt 1268 of AYVB are also required for replication.

SCR is highly conserved nature between distinct satellites [typically above 65% sequence identity with blocks of absolutely conserved sequence [7] strongly suggests that it also plays an important role in the virus replication cycle. In addition, the adjacent stem-loop and conserved nonanucleotide sequence would be expected to participate in replication. Approximately the 386 nt upstream of the stem-loop structure in ToLCV sat-DNA, as well as the stem-loop structure itself, are essential for replication [47].

$\beta$ C1 is a multi-functional protein encoded by betasatellites that are associated with the majority of monopartite begomoviruses [11]. For example Cheng et al. [13] proved by deletion mutants of Y10 $\beta$ C1 that multimerization was mediated by amino acids between positions 60 and 100. Previous studies say that the C-terminal sequences of BYVMB- $\beta$ C1 were interact with karyopherin  $\alpha$ , a transport receptor involved in nuclear import [42]. A myristoylation-like motif (GMDVNE) located at the C-terminal of CLCuMB- $\beta$ C1 (103-108aa)

interacted with a ubiquitin-conjugating enzyme involved in targeting proteins for degradation by the 26S proteasome [21]. It also seems to indicate interference with a functionality associated with the C terminus of Y10 $\beta$ C1.  $\beta$ C1 protein of AYVB, CLCuMB or BYVMB with GFP fused at the N-terminus also presented as granular spots in the cytoplasm and around the nucleus [42, 84].

TYLCCNB presumably has one or more *cis*-acting elements needed for replication and binds to TYLCCNV replication protein (Rep) for replication, and these elements are most probably located in the 115-nucleotide highly conserved region of betasatellite upstream of its stem-loop structure. Recently, Astroga [2] showed that a 5-bp core sequence (GGN1N2N3) is a typical constituent of Rep-binding iterons. Conserved GG motifs occur upstream of the 115-nucleotide highly conserved region of betasatellite. One or more of these GG motifs, combined with the 115 nucleotide highly conserved region, possibly responsible for Rep binding to betasatellite. However, the Rep binding activity of the TLCV-sat from Australia seems much less specific: TLCV-sat contains an A-rich region but lacks a  $\beta$ C1 gene and is believed to be a defective betasatellite molecule [48]. The effect of mutation of the conserved  $\beta$ C1 gene of TYLCCNVB indicated that the  $\beta$ C1 protein plays a key role in symptom induction.

The position and size of the  $\beta$ C1 gene of the betasatellite molecules are conserved in all betasatellite molecules, and the mutation of the start codon of C1 gene in TYLCCNB showed that it's a pathogenicity determinant [108, 6]. Few studies has been also shown that the  $\beta$ C1 protein of betasatellite associated with TYLCCNV or AYVV is an essential pathogenicity determinant [17, 79], it may act as suppressors of post-transcriptional gene silencing that interfering the host defense system, thus, the presence of C1 protein facilitates efficient infection of the virus in hosts [102]. For example Tau and Zhou [96] showed that  $\beta$ C1 gene was not required for the TYLCCNV and betasatellite replication and truncated betasatellite molecules with the deletion of the entire  $\beta$ C1 gene were stable in infected plants. Defective DNAs, betasatellite and alphasatellite associated with begomoviruses are maintained at approximately half the size of the genomic components [83, 89, 55, 80, 77]. Some proofs have been displayed that geminiviruses CP can encapsidate circular ssDNA molecules of about half or quarter the size of the genomic DNA [18, 51]. Immunocapture PCR indicated that the truncated TYLCCNB of about 1 kb in length may be encapsidated with TYLCCNV coat protein in vivo [67].

$\beta$ C1 of BYVMB have a nuclear export or peripheral localization function and  $\beta$ C1 interacts with itself, also with CP and the tomato protein karyopherin  $\alpha$ . Mutagenesis of  $\beta$ C1 protein showed that the domain of  $\beta$ C1 interacting with CP is at the N terminal half whereas the domain(s) of  $\beta$ C1 interacting with itself and karyopherin  $\alpha$  are at the C terminal half and the role of BYVMD  $\beta$ C1 as a suppressor of posttranscriptional gene silencing was explored [42]. Karyopherins are soluble transport receptors that interact with basic NLS sequences and help in nuclear import [27]. Full length betasatellite of CLCuMB can substitute for the movement function of the DNA B of a bipartite begomovirus *Tomato leaf curl New Dehli virus DNA-B* (ToLCNDV DNA-B). However, the betasatellite containing a disrupted  $\beta$ C1 ORF did not mobilize the DNA A for systemic infection, suggested that the  $\beta$ C1 protein was required for movement [74].

## 8. Potential of mutated satellites using a virus induced gene silencing vectors

### 8.1. Betasatellite

Betasatellites have about 200nt sequences (known as a-rich region) conserved among all that indicating may be these sequences have some biological roles in satellites. The role of A-Rich sequence may be to increase the required size of the molecule that is essential for encapsidation or systemic movement by the coat protein or movement protein encoded by begomovirus. TYLCCNB-Y10 could be infectious and mutant betasatellite (deleted a-rich region) could be encapsidated in the coat protein encoded by DNA-A that suggested may be A-Rich region is not required for trans-replication of TYLCCNB but only for size maintaining [95]. For example [20] reported that only a small region of the nucleotide sequence of CLCuMB upstream of the start codon of  $\beta C1$  (a 68-nt fragment), which contains a G-box, was important for  $\beta C1$  promoter activity. In addition to  $\beta C1$  ORF of CLCuMB delete a larger region (complete  $\beta C1$ ) to make it a gene delivery vector for plants. It can potentially tolerate the insertion of larger foreign sequences without affecting promoter activity [38]. Putative promoter and TATA box are located upstream of the  $\beta C1$  gene. Thus, the  $\beta C1$  gene of betasatellite could be replaced by a foreign gene and be modified to convert it into an expression vector [17]. The modified betasatellite might be an candidate gene silencing vector to study functional genomics in plants [54]. Also leaf curl symptoms in *Nicotiana* species can be brought by transgenic expression of the  $\beta C1$  gene of TYLCCNB that the severity of the symptoms parallels the level of  $\beta C1$  transcript in the transgenic plants and their ability to induce symptoms is abolished by mutation of the  $\beta C1$  gene. Possibly  $\beta C1$  gene of betasatellite may be replaced with a foreign gene and used as an expression vector for gene function analysis in plants [20].

Evidence has been shown that TYLCCNB modified by deletion of its  $\beta C1$  gene but retaining the  $\beta C1$  promoter and terminator, can be turned into a gene silencing vector. Also, insertion in the vector with fragments of endogenous plant genes or a transgene, in either the sense or antisense direction, can result in effective silencing of the cognate gene in plants [95]. The size of the mutated satellite DNA molecule significantly influences replication efficiency. TLCV sat-DNA can be used as a potential gene expression/silencing vector [47].

### 8.2. Alphasatellite

Also alphasatellite is a small molecule and easy to manipulate and have a wide host range and can apparently be maintained by a large number of distinct Begomovirus species. It has some sequences (A-rich approx.200 nt.), similar to betasatellite which can, potentially, be removed and still it can replicate autonomous. The A-rich deleted sequences of CLCuMA can not affect its ability to replicate autonomously and move, in trans, by a helper begomovirus that provide a space suitable for insertion of foreign sequences to increase its capacity to accept and maintain foreign gene sequences [82] (Table 2). This ability to amplify itself is useful for construction of VIGS vectors it will increase the copy number (and thus

also expression) of inserted sequences [82]. Rolling-circle replication initiator protein of GmusSA and GDarSLA act as a strong suppressor of PTGS.

## 9. Conclusions

The monopartite begomovirus associated with DNA-satellites (Betasatellite and Alphasatellite ) complex is in the norm throughout the Old World, particularly in South Asian countries. The epidemiology and evolution of this complex has been extensively analyzed since its first description. Monopartite begomovirus encoded all the genes needed to cause a successful infection. Many of these genes are coding for multifunctional proteins, adding another level of complexity in their interaction with host proteins, and their de novo creation. This shows the ability of begomoviruses and their associated satellites to rapidly evolve in response to selection pressures such as host plant resistance.

## Author details

Muhammad Shafiq Shahid and Masato Ikegami  
NODAI Research Institute, Tokyo University of Agriculture, Tokyo, Japan

Pradeep Sharma \*  
Division of Crop Improvement, Directorate of Wheat Research, Karnal, India

## 10. References

- [1] Amin, I., Patil, B., Briddon, R. W., Mansoor, S., and Fauquet, C. M. (2011). A common set of developmental miRNAs are upregulated in *Nicotiana benthamiana* by diverse begomoviruses. *Virology Journal* 8(1), 143.
- [2] Argüello-Astorga, G. R., and Ruiz, M. R. (2001). An iteron-related domain is associated to Motif 1 in the replication proteins of geminiviruses: identification of potential interacting amino acid-base pairs by a comparative approach. *Arch Virol* 146, 1465–1485.
- [3] Azzam, O., Frazer, J., de-la-Rosa, D., Beaver, J. S., Ahlquist, P., and Maxwell, D. P. (1994). Whitefly transmission and efficient ssDNA accumulation of *bean golden mosaic geminivirus* require functional coat protein. *Virology* 204(1), 289-96.
- [4] Bisaro, D. (2006). Silencing suppression by geminivirus proteins. *Virology* 344, 158 - 168.
- [5] Boulton, M. I., Steinkellner, H., Donson, J., Markham, P. G., King, D. I., and Davies, J. W. (1989). Mutational analysis of the virion-sense genes of *maize streak virus*. *J Gen Virol* 70(Pt 9), 2309-23.
- [6] Briddon, R. W. (2003). Cotton leaf curl disease, a multicomponent begomovirus complex. *Molecular Plant Pathol* 4, 427 - 434.
- [7] Briddon, R. W., Bull, S., Amin, I., Idris, A., Mansoor, S., Bedford, I., Dhawan, P., Rishi, N., Siwatch, S., and Abdel-Salam, A. (2003a). Diversity of DNA  $\beta$ : a satellite molecule associated with some monopartite begomoviruses. *Virology* 312, 106 - 121.

---

\* Corresponding Author

- [8] Briddon, R. W., Bull, S. E., Amin, I., Mansoor, S., Bedford, I. D., Dhawan, P., Rishi, N., Siwatch, S. S., Zafar, Y., Abdel-Salam, A. M., and Markham, P.G. (2004). Diversity of DNA 1: a satellite molecule associated with some monopartite begomoviruses-DNA  $\beta$  complex. *Virology* 324(2),462-474.
- [9] Briddon, R. W., Mansoor, S., Bedford, I. D., Pinner, M. S., Saunders, K., Stanley, J., Zafar, Y., Malik, K. A., and Markham, P. G. (2001). Identification of DNA components required for induction of cotton leaf curl disease. *Virology* 285, 234 - 243.
- [10] Briddon, R. W., Pinner, M., Stanley, J., and Markham, P. (1990). Geminivirus coat protein replacement alters insect specificity. *Virology* 177, 85 - 94.
- [11] Briddon, R. W., and Stanley, J. (2006). Sub-viral agents associated with plant-infecting single-stranded DNA viruses. *Virology* 344, 198 - 210.
- [12] Briddon, R. W., Watts, J., Markham, P. G., and Stanley, J. (1989). The coat protein of *beet curly top virus* is essential for infectivity. *Virology* 172(2), 628-633.
- [13] Cheng, X., Wang, X., Wu, J., Briddon, R. W., and Zhou, X. (2011).  $\beta$  C1 encoded by tomato yellow leaf curl China betasatellite forms multimeric complexes in vitro and in vivo. *Virology* 409(2), 156-162.
- [14] Clerot, D., and Bernardi, F. (2006). DNA helicase activity is associated with the replication initiator protein rep of *tomato yellow leaf curl geminivirus*. *J Virol* 80(22), 11322-30.
- [15] Cui, X., Li, G., Wang, D., Hu, D., and Zhou, X. (2005a). A begomovirus DNA betaencoded protein binds DNA, functions as a suppressor of RNA silencing, and targets the cell nucleus. *J Virol* 79, 10764 - 10775.
- [16] Cui, X., Li, G., Wang, D., Hu, D., and Zhou, X. (2005b). A begomovirus DNA  $\beta$ -encoded protein binds DNA, functions as a suppressor of RNA silencing, and targets the cell nucleus. *J Virol* 79, 10764 - 10775.
- [17] Cui, X., Tao, X., Xie, Y., Fauquet, C. M., and Zhou, X. (2004). A DNAbeta associated with *tomato yellow leaf curl China virus* is required for symptom induction. *J Virol* 78, 13966 - 13974.
- [18] Dry, I. B., Leslie, R., Krake, Justin, E., Rigden, and Rezaian, M. A. (1997). A novel subviral agent associated with a geminivirus: The first report of a DNA satellite. *Proceedings of the National Academy of Sciences* 94(13), 7088-7093.
- [19] Eagle, P. A., Orozco, B. M., and Hanley-Bowdoin, L. (1994). A DNA sequence required for geminivirus replication also mediates transcriptional regulation. *Plant Cell* 6(8), 1157-1170.
- [20] Eini, O., Akbar Behjatnia, S. A., Satish, D., Dry, I. B., Randles, J. W., and Rezaian, M. A. (2009a). Identification of sequence elements regulating promoter activity and replication of a monopartite begomovirus-associated DNA beta satellite. *J Gen Virol* 90, 253-260.
- [21] Eini, O., Dogra, S., Selth, L. A., Dry, I. B., Randles, J. W., and Rezaian, M. A. (2009b). Interaction with a host ubiquitin-conjugating enzyme is required for the pathogenicity of a geminiviral DNA  $\beta$  satellite. *Molecular Plant Microbe Interaction* 22(6), 737-46.
- [22] Elmer, J. S., Brand, L., Sunter, G., Gardiner, W. E., Bisaro, D. M., and Rogers, S. G. (1988). Genetic analysis of the *tomato golden mosaic virus*. II. The product of the AL1 coding sequence is required for replication. *Nucleic Acids Research* 16(14B), 7043-60.



- [23] Etesami, P., Watts, J., and Stanley, J. (1989). Size reversion of *African cassava mosaic virus* coat protein gene deletion mutants during infection of *Nicotiana benthamiana*. *J Gen Virol* 70(Pt 2), 277-89.
- [24] Frischmuth, T., and Stanley, J. (1998). Recombination between viral DNA and the transgenic coat protein gene of *African cassava mosaic geminivirus*. *J Gen Virol* 79(Pt 5), 1265-71.
- [25] Gilbertson, R. L., Sudarshana, M., Jiang, H., Rojas, M. R., and Lucas, W. J. (2003). Limitations on geminivirus genome size imposed by plasmodesmata and virusencoded movement protein: insights into DNA trafficking. *Plant Cell* 15(11), 2578-91.
- [26] Gopal, P., Kumar, P., Sinilal, B., Jose, J., Kasin Yadunandam, A., and Usha, R. (2007). Differential roles of C4 and  $\beta$  C1 in mediating suppression of post-transcriptional gene silencing: Evidence for transactivation by the C2 of *bhendi yellow vein mosaic virus*, a monopartite begomovirus. *Virus Res* 123, 9 - 18.
- [27] Gorlich, D., and Kutay, U. (1999). Transport between the cell nucleus and the cytoplasm. *Annual Review of Cell and Development Biology* 15, 607-60.
- [28] Hanley-Bowdoin, L., Settlage, S., and Robertson, D. (2004). Reprogramming plant gene expression: a prerequisite to geminivirus DNA replication. *Molecular Plant Pathol* 5, 149 - 156.
- [29] Hanley-Bowdoin, L., Settlage, S. B., Orozco, B. M., Nagar, S., and Robertson, D. (1999). Geminiviruses: Models for Plant DNA Replication, Transcription, and Cell Cycle Regulation. *Critical Reviews in Plant Sciences* 18(1), 71-106.
- [30] Heyraud-Nitschke, F., Schumacher, S., Laufs, J., Schaefer, S., Schell, J., and Gronenborn, B. (1995). Determination of the origin cleavage and joining domain of geminivirus Rep proteins. *Nucleic Acids Research* 23, 910-916.
- [31] Ho, S. N., Hunt, H. D., Horton, R. M., Pullen, J. K., and Pease, L. R. (1989). Sitedirected mutagenesis by overlap extension using the polymerase chain reaction. *Gene* 77(1), 51-9.
- [32] Hofer, P., Engel, M., Jeske, H., and Frischmuth, T. (1997). Nucleotide sequence of a new bipartite geminivirus isolated from the common weed *Sida rhombifolia* in Costa Rica. *J Gen Virol* 78, 1785 - 1790.
- [33] Höhnle, M., Höfer, P., Bedford, I. D., Briddon, R. W., Markham, P. G., and Frischmuth, T. (2001). Exchange of three amino acids in the coat protein results in efficient whitefly transmission of a non transmissible *abutilon mosaic virus* isolate. *Virology* 290, 164-171.
- [34] Hong, Y., Stanley, J., and van Wezel, R. (2003). Novel system for the simultaneous analysis of geminivirus DNA replication and plant interactions in *Nicotiana benthamiana*. *J Virology* 77(24), 13315-22.
- [35] Idris, M. A., Shahid, M. S., Briddon, R. W., Khan, A. J., Zhu, J. K., and Brown, J. K. (2011). An unusual alphasatellite associated with monopartite begomoviruses attenuates symptoms and reduces betasatellite accumulation. *J Gen Virol* 92, 706-717.
- [36] Karpova, O. V., Ivanov, K. I., Rodionova, N. P., Dorokhov Yu, L., and Atabekov, J. G. (1997). Nontranslatability and dissimilar behavior in plants and protoplasts of viral RNA and movement protein complexes formed in vitro. *Virology* 230(1), 11-21.
- [37] Karpova, O. V., Rodionova, N. P., Ivanov, K. I., Kozlovsky, S. V., Dorokhov, Y. L., and Atabekov, J. G. (1999). Phosphorylation of *Tobacco Mosaic Virus* Movement Protein Abolishes Its Translation Repressing Ability. *Virology* 261(1), 20-24.



- [38] Kharazmi, S., Behjatnia, S. A., Hamzehzarghani, H., and Niazi, A. (2012). Cotton leaf curl Multan betasatellite as a plant gene delivery vector trans-activated by taxonomically diverse geminiviruses. *Arch Virol* 5, 5.
- [39] Kheyr-Pour, A. B., M. Matzeit, V. Accotto, G. P. Crespi, S. Gronenborn, B. (1991). *tomato yellow leaf curl virus* from Sardinia is a whitefly-transmitted monopartite geminivirus. *Nucl Acids Res* 19(24), 6763–6769.
- [40] Kon, T., Rojas, M., Abdourhamane, I., and Gilbertson, R. (2009). Roles and interactions of begomoviruses and satellite DNAs associated with Okra leaf curl disease in Mali, West Africa. *J Gen Virol* 90, 1001 - 1013.
- [41] Kon, T., Sharma, P., and Ikegami, M. (2007). Suppressor of RNA silencing encoded by the monopartite *tomato leaf curl Java begomovirus*. *Arch Virol* 152(7), 1273-1282.
- [42] Kumar P, P., Usha, R., Zrachya, A., Levy, Y., Spanov, H., and Gafni, Y. (2006). Protein–protein interactions and nuclear trafficking of coat protein and  $\beta$ C1 protein associated with Bhendi yellow vein mosaic disease. *Virus Res* 122(1–2), 127-136.
- [43] Kunik, T., Palanichelvam, K., Czosnek, H., Citovsky, V., and Gafni, Y. (1998). Nuclear import of the capsid protein of *tomato yellow leaf curl virus* (TYLCV) in plant and insect cells. *Plant J* 13(3), 393-9.
- [44] Lacatus, G., and Sunter, G. (2008). Functional analysis of bipartite begomovirus coat protein promoter sequences. *Virology* 376(1), 79-89.
- [45] Laufs, J., Traut, W., Heyraud, F., Matzeit, V., Rogers, S. G., Schell, J., and Gronenborn, B. (1995). *In vitro* cleavage and ligation at the viral origin of replication by the replication protein of *tomato yellow leaf curl virus*. *PNAS* 92, 3879-3883.
- [46] Lazarowitz, S. G., Pinder, A. J., Damsteegt, V. D., and Rogers, S. G. (1989). *Maize streak virus* genes essential for systemic spread and symptom development. *EMBO J* 8(4), 1023-32.
- [47] Li, D., Akbar Behjatnia, S. A., Dry, I. B., Randles, J. W., Eini, O., and Rezaian, M. A. (2007). Genomic regions of *tomato leaf curl virus* DNA satellite required for replication and for satellite-mediated delivery of heterologous DNAs. *J Gen Virol* 88(7), 2073-2077.
- [48] Lin, B., Akbar Behjatnia, S. A., Dry, I. B., Randles, J. W., and Rezaian, M. A. (2003). High-Affinity Rep-Binding Is not Required for the Replication of a Geminivirus DNA and Its Satellite. *Virology* 305(2), 353-363.
- [49] Liu, H., Boulton, M. I., Oparka, K. J., and Davies, J. W. (2001). Interaction of the movement and coat proteins of *maize streak virus*: implications for the transport of viral DNA. *J Gen Virol* 82(1), 35-44.
- [50] Liu, H., Boulton, M. I., Thomas, C. L., Prior, D. A., Oparka, K. J., and Davies, J. W. (1999). *Maize streak virus* coat protein is karyophilic and facilitates nuclear transport of viral DNA. *Mol Plant Microbe Interact* 12(10), 894-900.
- [51] Liu, Y., Robinson, D. J., and Harrison, B. D. (1998). Defective forms of *cotton leaf curl virus* DNA-A that have different combinations of sequence deletion, duplication, inversion and rearrangement. *J Gen Virol* 79(P6), 1501-8.
- [52] Llave, C., Kasschau, K. D., Rector, M. A., and Carrington, J. C. (2002). Endogenous and Silencing-Associated Small RNAs in Plants. *The Plant Cell Online* 14(7), 1605-1619.
- [53] Mansoor, S., Briddon, R. W., Bull, S. E., Bedford, I. D., Bashir, A., Hussain, M., Saeed, M., Zafar, Y., Malik, K. A., Fauquet, C. M., and Markham, P. G. (2003a). Cotton leaf curl

- disease is associated with multiple monopartite begomoviruses supported by single DNA  $\beta$ . *Arch Virol* 148(10), 1969-1986.
- [54] Mansoor, S., Briddon, R. W., Zafar, Y., and Stanley, J. (2003b). Geminivirus disease complexes: an emerging threat. *Trends Plant Sci* 8, 128 - 134.
  - [55] Mansoor, S., Khan, S. H., Bashir, A., Saeed, M., Zafar, Y., Malik, K., Briddon, R. W., Stanley, J., and Markham, P. G. (1999a). Identification of a novel circular single-stranded DNA associated with cotton leaf curl disease in Pakistan. *Virology* 259, 190 -199.
  - [56] Mansoor, S., Khan, S. H., Bashir, A., Saeed, M., Zafar, Y., Malik, K. A., Briddon, R. W., Stanley, J., and Markham, P. G. (1999b). Identification of a novel circular singlestranded DNA associated with cotton leaf curl disease in Pakistan. *Virology* 259, 190 -199.
  - [57] Meister, G., and Tuschl, T. (2004). Mechanisms of gene silencing by double-stranded RNA. *Nature* 431(7006), 343-349.
  - [58] Mubin, M., Amin, I., Amrao, L., Briddon, R. W., and Mansoor, S. (2010). The hypersensitive response induced by the V2 protein of a monopartite begomovirus is countered by the C2 protein. *Mol Plant Pathol* 11(2), 245-254.
  - [59] Nawaz-ul-Rehman, M. S., Mansoor, S., Briddon, R. W., and Fauquet, C. M. (2009). Maintenance of an Old World betasatellite by a New World helper begomovirus and possible rapid adaptation of the betasatellite. *J Virol* 83, 9347 - 9355.
  - [60] Nawaz-ul-Rehman, M. S., Nahid, N., Mansoor, S., Briddon, R. W., and Fauquet, C. M. (2010). Post-transcriptional gene silencing suppressor activity of the alpha-Rep of non-pathogenic alphasatellites associated with begomoviruses. *Virology* 405, 300 -308.
  - [61] Noris, E., Jupin, I., Accotto, G. P., and Gronenborn, B. (1996). DNA-binding activity of the C2 protein of *tomato yellow leaf curl geminivirus*. *Virology* 217(2), 607-612.
  - [62] Noris, E. V., A. M. Caciagli, P. Masenga, V. Gronenborn, B. Accotto, G. P. (1998). Amino acids in the capsid protein of *tomato yellow leaf curl virus* that are crucial for systemic infection, particle formation, and insect transmission. *J Virol* 72 (12 ), 10050-10057.
  - [63] Orozco, B. M., Kong, L.-J., Batts, L. A., Elledge, S., and Hanley-Bowdoin, L. (2000). The multifunctional character of a geminivirus replication protein is reflected by its complex oligomerization properties. *J Biol Chem* 275, 6114-6122.
  - [64] Padidam, M., Beachy, R. N., and Fauquet, C. M. (1996). The role of AV2 ("precoat") and coat protein in viral replication and movement in *tomato leaf curl geminivirus*. *Virology* 224, 390 - 404.
  - [65] Pascal, E., Goodlove, P. E., Wu, L. C., and Lazarowitz, S. G. (1993). Transgenic tobacco plants expressing the geminivirus BL1 protein exhibit symptoms of viral disease. *Plant Cell* 5(7), 795-807.
  - [66] Qazi, J., Amin, I., Mansoor, S., Iqbal, M. J., and Briddon, R. W. (2007). Contribution of the satellite encoded gene  $\beta$  C1 to cotton leaf curl disease symptoms. *Virus Res* 128(1-2), 135-139.
  - [67] Qian, Y., and Zhou, X. (2005). Pathogenicity and stability of a truncated DNA  $\beta$  associated with *tomato yellow leaf curl China virus*. *Virus Res* 109(2), 159-163.
  - [68] Rhee, Y., Gurel, F., Gafni, Y., Dingwall, C., and Citovsky, V. (2000). A genetic system for detection of protein nuclear import and export. *Nature Biotechnology* 18(4), 433-437.
  - [69] Rigden, J. E., Dry, I. B., Mullineaux, P. M., and Rezaian, M. A. (1993). Mutagenesis of the virion-sense open reading frames of *tomato leaf curl geminivirus*. *Virology* 193(2), 1001-1005.

- [70] Rojas, M. R., Jiang, H., Salati, R., Xoconostle-Cazares, B., Sudarshana, M.R., Lucas, W.J., Gilbertson, R.L. (2001). Functional analysis of proteins involved in movement of the monopartite begomovirus, *tomato yellow leaf curl virus*. *Virology* 291, 110–125.
- [71] Ruiz, M. T., Voinnet, O., and Baulcombe, D. C. (1998). Initiation and Maintenance of Virus-Induced Gene Silencing. *Plant Cell* 10(6), 937-946.
- [72] Saeed, M., Behjatnia, S., Mansoor, S., Zafar, Y., Hasnain, S., and Rezaian, M. (2005). A single complementary-sense transcript of a geminiviral DNA  $\beta$  satellite is determinant of pathogenicity. *Mol Plant-Microbe Interact* 18, 7 - 14.
- [73] Saeed, M., Mansoor, S., Rezaian, M. A., Briddon, R. W., and Randles, J. W. (2008). Satellite DNA beta overrides the pathogenicity phenotype of the C4 gene of *tomato leaf curl virus*, but does not compensate for loss of function of the coat protein and V2 genes. *Arch Virol* 153, 1367 - 1372.
- [74] Saeed, S., Zafar, Y., Randles, J. W., and Rezaian, M. A. (2007). A monopartite begomovirus-associated DNA beta satellite substitutes for the DNA B of a bipartite begomovirus to permit systemic infection. *J Gen Virol* 88, 2881 - 2889.
- [75] Sanderfoot, A. A., and Lazarowitz, S. G. (1995). Cooperation in Viral Movement: The Geminivirus BL1 Movement Protein Interacts with BR1 and Redirects It from the Nucleus to the Cell Periphery. *Plant Cell* 7(8), 1185–1194.
- [76] Saunders, K., Bedford, I., Briddon, R., Markham, P., Wong, S., and Stanley, J. (2000). A unique virus complex causes Ageratum yellow vein disease. *PNAS* 97, 6890 - 6895.
- [77] Saunders, K., Bedford, I. D., and Stanley, J. (2001). Pathogenicity of a natural recombinant associated with ageratum yellow vein disease: implications for geminivirus evolution and disease aetiology. *Virology* 282(1), 38-47.
- [78] Saunders, K., Briddon, R. W., and Stanley, J. (2008). Replication promiscuity of DNAbetasatellites associated with monopartite begomoviruses; deletion mutagenesis of the *ageratum yellow vein virus* DNA  $\beta$  satellite localises sequences involved in replication. *J Gen Virol* 89, 3165 - 3172.
- [79] Saunders, K., Norman, A., Gucciardo, S., and Stanley, J. (2004). The DNA beta satellite component associated with ageratum yellow vein disease encodes an essential pathogenicity protein ( $\beta$  C1). *Virology* 324, 37 - 47.
- [80] Saunders, K., and Stanley, J. (1999). A nanovirus-like DNA component associated with yellow vein disease of Ageratum conyzoides: evidence for interfamilial recombination between plant DNA viruses. *Virology* 264, 142 - 152.
- [81] Settlege, S. B., See, R. G., and Hanley-Bowdoin, L. (2005). Geminivirus C3 protein: replication enhancement and protein interactions. *J Virol* 79, 9885 - 9895.
- [82] Shahid, M. S., Ali, L., Andleeb, S. (2009). The function of the a-rich region of the alphasatellite associated with the cotton leaf curl disease in Pakistan. *EurAsia J BioSci* 3, 152-156.
- [83] Sharma, A. M., A. Osaki, T. Ikegami, M. (1998). Characterization of virus-specific DNA forms from tomato tissues infected by *tobacco leaf curl virus*: evidence for a single genomic component producing defective DNA molecules. *Plant Pathol* 47, 787-793.
- [84] Sharma, P., Ikegami, M., and Kon, T. (2010). Identification of the virulence factors and suppressors of posttranscriptional gene silencing encoded by *ageratum yellow vein virus*, a monopartite begomovirus. *Virus Res* 149(1), 19-27.

- [85] Stanley, J., Bisaro, D. M., Briddon, R. W., Brown, J. K., Fauquet, C. M., Harrison, B. D., Rybicki, E. P., and Stenger, D. C. (2005a). Geminiviridae. *Virus Taxonomy, VIIIth Report of the ICTV*, 301 - 326.
- [86] Stanley, J., Bisaro, D. M., Briddon, R. W., Brown, J. K., Fauquet, C. M., Harrison, B. D., Rybicki, E. P., and Stenger, D. C. (2005b). Geminiviridae. *Virus Taxonomy, VIIIth Report of the ICTV*, 301 - 326.
- [87] Stanley, J., Bisaro, D. M., Briddon, R. W., Brown, J. K., Fauquet, C. M., Harrison, B. D., Rybicki, E. P., Stenger, D. C., by, E., Fauquet, C. M., Mayo, M. A., Maniloff, J., Desselberger, J., and Ball, L. A. (2005c). Geminiviridae. In *Virus Taxonomy, 5th report of the ICTV*. London: Elsevier/Academic Press, pp 301–326.
- [88] Stanley, J., and Latham, J. R. (1992). A symptom variant of *beet curly top geminivirus* produced by mutation of open reading frame C4. *Virology* 190(1), 506-509.
- [89] Stanley, J., Saunders, K., Pinner, M. S., and Wong, S. M. (1997). Novel defective interfering dnas associated with *ageratum yellow vein geminivirus* infection of *ageratum conyzoides*. *Virology* 239(1), 87-96.
- [90] Sunter, G., and Bisaro, D. (1997). Regulation of a geminivirus coat protein promoter by AL2 protein (TrAP): evidence for activation and derepression mechanisms. *Virology* 232, 269 - 280.
- [91] Sunter, G., and Bisaro, D. M. (1991). Transactivation in a geminivirus: AL2 gene product is needed for coat protein expression. *Virology* 180(1), 416-9.
- [92] Sunter, G., Hartitz, M. D., Hormuzdi, S. G., Brough, C. L., and Bisaro, D. M. (1990). Genetic analysis of *tomato golden mosaic virus*: ORF AL2 is required for coat protein accumulation while ORF AL3 is necessary for efficient DNA replication. *Virology* 179(1), 69-77.
- [93] Sunter, G., Stenger, D. C., and Bisaro, D. M. (1994). Heterologous complementation by geminivirus AL2 and AL3 genes. *Virology* 203(2), 203-10.
- [94] Talya Kunik, L. M., Vitaly Citovsky and Yedidya Gafni (1999). Characterization of a tomato karyopherin  $\alpha$  that interacts with the *tomato yellow leaf curl virus* (TYLCV) capsid protein. *Journal of Experimental Botany* 50(334), 731-732.
- [95] Tao, X., Qing, L., and Zhou, X. (2004). Function of A-Rich region in DNA  $\beta$  associated with *tomato yellow leaf curl China virus*. *Chinese Sci Bull* 49(14), 1490-1493.
- [96] Tao, X., and Zhou, X. (2004). A modified viral satellite DNA that suppresses gene expression in plants. *Plant J* 38(5), 850-60.
- [97] Trinks, D. R., R. Shivaprasad, P. V. Akbergenov, R. Edward J. Oakeley, K. Veluthambi, Thomas Hohn, Pooggin, M. M. (2005). Suppression of RNA silencing by a *geminivirus* nuclear protein, ac2, correlates with transactivation of host genes. *J Virol* 79(4), 2517-2527.
- [98] Van Wezel, R., Dong, X., Liu, H., Tien, P., Stanley, J., and Hong, Y. (2002). Mutation of three cysteine residues in *tomato yellow leaf curl China virus* C2 protein causes dysfunction in pathogenesis and posttranscriptional gene silencing suppression. *Mol Plant Microbe Interaction* 15, 203 - 208.
- [99] Vanitharani, R., Chellappan, P., Pita, J. S., and Fauquet, C. M. (2004). Differential roles of AC2 and AC4 of cassava geminiviruses in mediating synergism and suppression of posttranscriptional gene silencing. *J Virol* 78, 9487 - 9498.
- [100] Voinnet, O. (2005). Induction and suppression of RNA silencing: insights from viral infections. *Nature Rev Genet* 6(3), 206-220.



- [101] Von Arnim, A., Frischmuth, T., and Stanley, J. (1993). Detection and possible functions of *African cassava mosaic virus* DNA B gene products. *Virology* 192(1), 264-72.
- [102] Wang, M. B., X. Wu, L. Liu, L. Smith, N. A. Isenegger, D. Wu, R. Masuta, C. Vance, V. B. Watson, J. Rezaian M.A. (2004). On the role of RNA silencing in the pathogenicity and evolution of viroids and viral satellites. *PNAS* 101(9), 3275-3280.
- [103] Wartig, L., Kheyr-Pour, A., Noris, E., De Kouchkovsky, F., Jouanneau, F., Gronenborn, B., and Jupin, I. (1997). Genetic analysis of the monopartite *tomato yellow leaf curl geminivirus*: roles of V1, V2, and C2 orfs in viral pathogenesis. *Virology* 228(2), 132-140.
- [104] Woolston, C. J., Reynolds, H. V., Stacey, N. J., and Mullineaux, P. M. (1989). Replication of wheat dwarf virus DNA in protoplasts and analysis of coat protein mutants in protoplasts and plants. *Nucl Acids Res* 17(15), 6029-6041.
- [105] Wu, P., and Zhou, X. (2005). Interaction between a nanovirus-like component and the *tobacco curly shoot virus*/satellite complex. *Acta Biochim Biophys Sin* 37, 25 - 31.
- [106] Yaakov, N., Levy, Y., Belausov, E., Gaba, V., Lapidot, M., and Gafni, Y. (2011). Effect of a single amino acid substitution in the NLS domain of *tomato yellow leaf curl virus*-Israel (TYLCV-IL) capsid protein (CP) on its activity and on the virus life cycle. *Virus Res* 158(1-2), 8-11.
- [107] Zhang, S. C., Wege, C., and Jeske, H. (2001). Movement proteins (BC1 and BV1) of *abutilon mosaic geminivirus* are cotransported in and between cells of sink but not of source leaves as detected by green fluorescent protein tagging. *Virology* 290(2), 249-60.
- [108] Zhou, X., Xie, Y., Tao, X., Zhang, Z., Li, Z., and Fauquet, C. M. (2003). Characterization of DNA $\beta$  associated with begomoviruses in China and evidence for co-evolution with their cognate viral DNA-A. *J Gen Virol* 84, 237 - 247.
- [109] Zrachya, A., Glick, E., Levy, Y., Arazi, T., Citovsky, V., and Gafni, Y. (2006). Suppressor of RNA silencing encoded by *tomato yellow leaf curl virus*-Israel. *Virology* 358, 159 - 165.
- [110] Kon, T., Hidayat, S.H., Hase, S., Takahashi, H., and Ikegami, M. (2006). The natural occurrence of two distinct begomoviruses associated with DNA  $\beta$  and a recombinant DNA in a tomato plant from Indonesia. *Phytopathology* 96(5), 517-525.
- [111] Ito, T., Kimbara, J., Sharma, P., and Ikegami, M. (2009). Interaction of tomato yellow leaf curl virus with diverse betasatellites enhances symptom severity. *Arch Virol* 154, 1233-1239.
- [112] Ogawa, T., Sharma, P. and Ikegami, M. (2008). The begomoviruses Honeysuckle yellow vein mosaic virus and Tobacco leaf curl Japan virus with DNA  $\beta$  satellites cause yellow dwarf disease of tomato. *Virus Res* 137(2), 235-244.
- [113] Sharma, P., and Ikegami, M. (2009). Characterization of signals that dictate nuclear/nucleolar and cytoplasmic shuttling of the capsid protein of Tomato leaf curl Java virus associated with DNA  $\beta$  satellite. *Virus Res* 144, 145-153.
- [114] Sharma, P., and Ikegami, M. (2010). Tomato leaf curl virus V2 protein is a determinant of virulence, hypersensitive response and suppression of posttranscriptional gene silencing. *Virology* 396, 85-93.
- [115] Sharma, P., Gaur, R.K. and Ikegami, M. (2011). Subcellular localization of V2 protein of Tomato leaf curl Java virus by using green fluorescent protein and yeast hybrid system. *Protoplasma* 248, 281-288.