We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists



186,000

200M



Our authors are among the

TOP 1% most cited scientists





WEB OF SCIENCE

Selection of our books indexed in the Book Citation Index in Web of Science™ Core Collection (BKCI)

Interested in publishing with us? Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected. For more information visit www.intechopen.com



The Role of the Coagulation System in Preterm Parturition

Vered Klaitman, Ruth Beer-Wiesel, Tal Rafaeli, Moshe Mazor and Offer Erez

Additional information is available at the end of the chapter

http://dx.doi.org/10.5772/54843

1. Introduction

Term and preterm parturition have a common pathway that includes irregular uterine contractions, cervical effacement and dilatation, along with decidual activation and rupturing of the chorioamniotic membrane. This pathway is observed in the physiologic labor at term as well as in the pathological processes leading to premature delivery. Indeed, the clinical presentation of preterm parturition involves all components of this common pathway: 1. Preterm contractions in women with spontaneous preterm labor with intact membranes; 2.cervical effacement and dilatation in women with cervical insufficiency; and/or 3.decidual activation and rupture of membranes in those with preterm PROM.

The syndrome of preterm parturition is the clinical presentation of several underlying mechanisms, not all of them being fully understood. Among the well establishes etiologies for preterm parturition are: intra amniotic infection/inflammation, cervical insufficiency, increased thrombin generation and vascular pathology of the placenta, as well as multiple gestations. The study of the maternal fetal interface and the placenta contributes to the deciphering of the mechanism leading to preterm birth. Placental and decidual vascular lesions have been reported in about 20-30% of patients who deliver preterm. There are accumulating evidence that preterm parturition is associated with an increased activation of maternal hemostatic system which also interacts with the acute inflammatory processes observed in this syndrome. Moreover, higher rates of fetal growth restriction and placental vascular lesions were observed among women with preterm labor who delivered at term suggesting that some vascular insults may not be severe enough to cause preterm birth but still inflict some effect on fetal growth.



© 2013 Klaitman et al.; licensee InTech. This is an open access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/3.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

The following chapter will summarize the changes in maternal hemostatic system during normal pregnancy and those associated with preterm labor and preterm PROM. In addition, we will review the vascular changes associated with preterm parturition. The last section of this chapter will address the role of the hemostatic and angiogenic markers for the prediction of spontaneous preterm birth.

2. The physiology of hemostasis

Hemostasis is crucial for the maintenance of the vascular tree integrity.

The major components of the hemostatic system, which function in concert, are the following: 1) the vessel wall and endothelium; 2) platelets and other formed elements of blood, such as monocytes and red cells; and 3) plasma proteins (the coagulation and fibrinolytic factors and inhibitors). These components act altogether in a synchronize fashion to generate the hemostatic plug, preserve the integrity of the vascular tree in the body, and avoid uncontrolled clot formation and thrombosis (See figure 1).



Figure 1. Initiation of coagulation by endothelial injury and exposure of tissue factor to the extrinsic pathway. Then, the platelets aggregate by activation of intrinsic pathway and platelet activation. And finally propagation and clot stabilization by thrombin – fibrin balance formation (from reference #12 with permission).

The vessel wall: Endothelial damage or activation is a crucial event that launches the cascade of reactions leading to thrombin generation and fibrin clot formation. The vascular endothelium is the focal point for both initiation and inhibition of the coagulation process. During endothelial damage the sub endothelial tissue factor molecules are exposed and initiate coagulation by activation of the extrinsic pathway. This is relevant also for the maternal fetal

interface since the villous and exteravillos trophoblast of the human placenta adopt the properties of endothelium and vessel wall in order to allow laminar flow of maternal blood through the placental bed without unnecessary activation of the coagulation cascade; and any damage in their integrity will activate the coagulation cascade.

Platelets activation and plaque formation: Following vascular injury platelets initially adhere to sub-endothelial collagen via von Willebrand factor (vWF) (figure 2). These vWF "bridges" are anchored at one end to sub-endothelial type IV collagen molecules and at the other end to the platelet GPIb/IX/V receptor [1,2]. The adherent platelets can also attach to other sub-endothelial extracellular matrix proteins (e.g. laminin, fibronectin, and vitronectin) via cell-membrane bound integrins [3]. The binding of these receptors activates the platelets through calcium-dependent cytoskeletal changes.



Figure 2. Mediators of platlet activation and agregation (from reference #5 with permission).

Activated platelets form pseudopodia that further enhance vWF coupling to the sub endothelium. Moreover, ADP induces a conformational change in the GPIIb/IIIa receptor on the platelet membrane causing platelet aggregation via the formation of a high affinity fibrinogen bridges anchored at either end by GPIIb/IIIa receptors on 2 different platelets [4]. Thus, Platelet activation converts the normally inactive Gp IIb/IIIa receptor into an active one, enabling binding to fibrinogen and VWF. Because the surface of each platelet has about 50,000 Gp IIb/ IIIa-binding sites, numerous activated platelets recruited to the site of vascular injury can rapidly form an occlusive aggregate by means of a dense network of intercellular fibrinogen bridges. Since this receptor is the key mediator of platelet aggregation, it has become an effective target for antiplatelet therapy [5].

2.1. Plasma proteins

Coagulation factors: Plasma coagulation proteins (*clotting factors*) normally circulate in plasma in their inactive forms. The sequence of coagulation protein reactions that culminate in the formation of fibrin was originally described as a *waterfall* or a *cascade*. Two pathways of blood coagulation have been described in the past: the extrinsic, or tissue factor, pathway and the intrinsic or contact activation, pathway (figure 3). However, the current approach is a more unify view in which the coagulation cascade is normally initiated through tissue factor exposure and activation of the *extrinsic pathway* that generates thrombin and activates the elements of the classic *intrinsic pathway*. These reactions take place on phospholipid surfaces, usually on activated platelets.

The initial phase of coagulation is the exposure of tissue factor to coagulation factors, caused either by endothelial damage or activation [6]. Tissue factor (TF) is a 47kDa cell bound trans-membrane glycoprotein and member of class 2 cytokine superfamily [7], that functions as: 1) a receptor, with signal transduction resulting in the induction of genes involved in inflammation, apoptosis, embryonic development and cell migration[8]; and 2])as an activator and cofactor for factors VII/VIIa in the coagulation cascade. It is constitutively expressed by many extravascular tissues, especially perivascular ones, and it is highly expressed in the brain, heart, lungs, kidneys, testis and placenta [9-11], reflecting the importance of these tissues to the organism[12]. TF expression can be induced in monocytes and platelets, and has been detected on circulating microparticles (MP) derived from these and other cell types [7,13]. The expression of this coagulation factor can also be induced on endothelium in response to inflammatory stimuli including exposure to: 1) bacterial lipopolysaccharide (LPS) in sepsis; 2) adhesion molecules (P-selectin expressed on platelets, CD40 ligand expressed on white blood cells); and 3) inflammatory cytokines (interleukin-6, tumor necrosis factor), and oxidized low-density lipoprotein(LDL) [14].

Tissue factor activates the coagulation cascade by binding to the serine protease factor VIIa; the complex of TF+FVIIa activates factor X to factor Xa and initiating the converting factor IX to factor IXa in the intrinsic system leading to further formation of factor Xa. Thus, this factor is formed through the actions of either the tissue factor/factor VIIa complex or factor IXa (with factor VIIIa as a cofactor), and converts pro-thrombin to thrombin, the pivotal protease of the coagulation system. Thrombin is a multifunctional enzyme that converts soluble plasma fibrinogen to an insoluble fibrin matrix. Thrombin also activates factor XIII (fibrin-stabilizing factor) to factor XIIIa, which covalently cross-links and thereby stabilizes the fibrin clot.



Figure 3. The extrinsic and intrinsic pathways of coagulation are a model for simplifying the coagulation system. This model is reflected in the laboratory PT and PTT measurements (from reference #12 with permission).

An additional mechanism in which the activation of the coagulation cascade can take place is through microparticles. Circulating microparticles are an area of intense research, as increased levels of them were shown to be pro-coagulant, even in the absence of TF expression [14-17]. These microparticles are tiny (<1 mm) membrane-bound vesicles and express membrane antigens that reflect their cellular origin[18]. The concentration of circulating microparticles is increased during platelet activation, inflammation, or apoptosis. Elevated concentrations of microparticles are encountered in diseases with vascular involvement and hypercoagulability such as disseminated intravascular coagulation, diabetes, and immune-mediated thrombosis [12]. Anti-coagulation protein:

Thrombin also plays a crucial role in activating the inhibition of the coagulation cascade.

Following its activation thrombin binds to thombomodulin causing a conformational change that activates the endothelial receptor of protein c, which in turns activates protein C. The latter is bound to its cofactor protein s together this complex inactivates factor Va and factor VIIIa of the intrinsic pathway, reducing substantially thrombin generation13. In addition to protein c and protein s there is the tissue factor pathway inhibitor (TFPI) which is the main inhibitor of the extrinsic pathway of coagulation, this protein inhibits the activity of factor VIIa and factor Xa reducing by this thrombin generation by the extrinsic pathway. the complex of protein z and Protein Z-dependent protease inhibitor (ZPI) inactivates factor Xa, and can also directly inhibit factor XIa. However, by far the most active inhibitor of both factor Xa and thrombin is antithrombin (AT) (previously known as antithrombin III). The AT molecule binds to either thrombin or factor Xa and then complexes with vitronectin which causes a conformational change that facilitates heparin binding. The resultant quaternary structure augments thrombin inactivation 1000-fold. The function of these anticoagulation protein is essential for maintaining the homeostasis between coagulation and adequate blood flow in the vascular tree, and deficiency in these protein is associated with increase risk for thromboembolic diseases and other complications.

Fibrinolysis factors and inhibitors: The process of fibrinolysis (i.e., clot lysis) is also crucial to the prevention of thrombosis (Figure 4). Fibrin is degraded to its degradation products (FDPs) by plasmin [19], that is generated from plasminogen by the action of tissue-type plasminogen activator (tPA) embedded in fibrin. This process is accelerated when plasminogen itself is bound to fibrin. Endothelial cells produce a second plasminogen activator, urokinase-type plasminogen activator (uPA). The latter's activation requires high molecular weight kininogen, kallikrein, and plasmin. This helps explain why deficiency of the former two "intrinsic pathway" clotting factors (ie, activators of factor XI) paradoxically lead to thrombosis and not bleeding.

There are series of inhibitors of premature fibrinolysis and, thus, hemorrhage. Plasmin is directly inhibited by a2-plasmin inhibitor. This inhibitor is also bound to the fibrin clot where it is positioned to prevent premature fibrinolysis. Type-1 plasminogen activator inhibitor (PAI-1) is synthesized by endothelial cells and platelets in response to thrombin binding to (protease activated receptor) PARs. In pregnancy, the decidua is a rich source of PAI-1, while the placenta produces PAI-1 and PAI -2, and serve as the primary source for the latter [20,21]. In the initial stages of platelet plug and fibrin clot formation, endothelial cells release PAI-1 but after a delay, endothelial cells release tPA and uPA to promote fibrinolysis.

Thrombin-activatable fibrinolysis inhibitor (TAFI) is another antifibrinolytic factor which acts by cleaving the C-terminal lysine in fibrin, to render it resistant to cleavage by plasmin [22,23]. Levels of TAFI are increased in the third trimester [23]. Interestingly, TAFI is also activated by the thrombin-thrombomodulin complex, once again implicating thrombin as the ultimate arbiter of hemostasis. The fibrinolytic system can influence coagulation in several ways. For example, FDPs inhibit thrombin action, a major source of hemorrhage in disseminated intravascular coagulation. In addition, PAI-1 bounded to vitronectin and also to heparin can directly inhibit thrombin and factor Xa activity [5,24].



Figure 4. Fibrinolysis factors and inhibitors. The following scheme shows the process of fibrin degradation and plasmin generation (from reference #5 with permission).

3. Pregnancy associated changes in the hemostatic system

Pregnancy is a challenging time period for the hemostatic system. The demands from this system changes in different sites and are somewhat contradictory. The formation of the placental bed in which the maternal blood is running outside the maternal vessels necessitate the mother to address two challenges, the first one is to protect herself from a life threatening bleeding, and the second is to enable a continues blood flow through the placental bed outside the maternal blood vessels without activating the coagulation cascade. These challenges have been addressed by the formation of three compartments: 1) The maternal compartment which becomes adaptive and pro-coagulant to prevent severe bleeding during delivery; 2) The fetus that develops his coagulation system during gestation while floating in the pro-coagulant amniotic fluid; and 3) The maternal fetal interface of which the intervillous space is hypocoa-

gulated in order to ensure the extravascular laminar flow of maternal blood and the maternal decidua is rich with tissue factor to prevent bleeding.

During gestation, changes in the coagulation system are considered to be adaptive to prevent hemorrhage at the time of delivery [25-29]. Indeed, normal pregnancy has been associated with excessive maternal thrombin generation [28,30] and a tendency for platelets to aggregate in response to agonists [31,32] (TABLE 1). Pregnancy is accompanied by 2 to 3-fold increase in fibrinogen concentrations and 20% to 1000% increase in factors VII, VIII, IX, X, and XII, all of which peak at term [33]. The concentrations of vWF increase up to 400% by term [33]. By contrast, those of pro-thrombin and factor V remain unchanged while the concentrations of factors XIII and XI decline modestly [34]. Indeed there is evidence of chronic low-level thrombin and fibrin generation throughout normal pregnancy as indicated by enhanced concentrations of pro-thrombin fragment 1.2, thrombin-antithrombin (TAT III) complexes, and soluble fibrin polymers[23]. Free protein S concentration declines significantly (up to 55%) during pregnancy due to increased circulating complement 4B-binding protein its molecular carrier. Protein S nadir at delivery and this reduction is exacerbated by cesarean delivery and infection [33,35]. As a consequence, pregnancy is associated with an increase in resistance to activated protein C [23,33]. The concentrations of PAI-1 increase by 3 to 4-folds during pregnancy while plasma PAI-2 values, which are negligible before pregnancy reach concentrations of 160 mg/L at delivery [33]. Thus, pregnancy is associated with increased clotting potential, as well as decreased anticoagulant properties, and fibrinolysis [5]. Therefore, it can be defined as a prothrombotic state.

		Increased	Decreased	No change
Systemic changes	Procoagulant factors	I,V,VII,VIII,IX,X	XI	
	Anticoagulant factors	Soluble TM	PS	
	Adhesive proteins	vWF		6
	Fibrinolytic proteins	PAI-1,PAI-2	t-PA	TAFI
	Microparticles and APLA	MP		APLA
Local placental changes		TF	TFPI	

Table 1. Hemostatic changes in pregnancy [42].

In contrast to the maternal circulation and the decidua, the establishment of the uteroplacental circulation challenges the hemostatic system. Indeed, fibrin deposition sites were identified in

decidual veins at sites of trophoblast invasion, where villi are implanted into veins [36]. Compared with endothelial vasculature, the trophoblasts lining decidual spiral arteries have a reduced capacity to lyse fibrin, and recent studies have shown that this is caused by high concentrations of plasminogen activator inhibitors [37] that affects it fybrinolytic capabilities. In addition, perivascular decidualized human endometrial stromal cells are ideally positioned to prevent postimplantation haemorrhage during endovascular trophoblast invasion by expressing TF, which is a primary cellular mediator of hemostasis [38,39]. In vivo and in vitro studies have demonstrated that estradiol (E2) enhances TF expression during progesterone induced decidualization. It was demonstrated that paracrine factors, such as endothelial growth factor (EGF) are involved with steroid-enhancing TF expression in decidualized human endometrial stromal cells through the EGF receptor [40]. The trophoblast and the placenta have distinct anticoagulation properties that aimed on one hand to prevent bleeding and on the other hand to allow laminar flow of maternal blood through the intervillous space [41,42]. Accumulating evidence suggest that the trophoblast acquires properties of vascular epithelium and expresses coagulation inhibitors such as tissue factor pathway inhibitor 2[43-45] (also known as placental protein 5 [46,47]), heparin co-factor II, dermatan sulfate [48, 49], and thrombomodulin [50-53] as well as pro-coagulant proteins such as tissue factor [38,54]. Moreover, a knockout mouse model for the endothelial receptor of protein C was lethal inutero and the embryos died on the 10.5 day of gestation, and fibrin deposition was found around their giant trophoblast cells of these embryos [55]. Moreover, a recent report demonstrated that the placenta is an extra-hepatic source of the anti-coagulant proteins including protein C, protein S, as well as protein Z, and their expression is constitutive irrespective of obstetrical conditions [85].

4. The hemostatic system and preterm parturition

The involvement of the hemostatic system in the pathophysiology of preterm parturition is becoming more and more apparent. Indeed, changes in maternal and fetal gens, abnormal placental vascular finding and increased thrombin generation in the maternal circulation were all reported in association with preterm parturition.

Genetic studies: During parturition there is remodeling of the extra cellular matrix of the uterine cervix. Recently increased expression of the tissue type plasminogen activator gene was reported during labor at term [56]. This finding was supported by the role of plasminogen activation in the remodeling of the extracellular matrix in human amnion, chorio-decidua, and placenta during and after labor [57]. Moreover, in a genetic association study that tested maternal and fetal genes in women with preterm labor reported this gene to be highly express in fetuses of Hispanic patients who delivered preterm.

Single nucleotide polymorphisms of the coagulation genes are associated with increased risk for preterm birth. Indeed, in a study that aimed to identify the impact of genetic polymorphisms with pro-thrombotic and anti-thrombotic effects on the occurrence of preterm birth in a large cohort of very-low-birth-weight (VLBW)-infants and their mothers, and term-borninfants and their mothers, maternal factor VII-121del/ins and the infant's factor VII-121del/ins polymorphisms were more frequent in the group of singleton VLBW and their mothers. Furthermore, the frequency of the factor XIII-Val34Leu polymorphism was significantly lower in singleton VLBW than in term infant controls; and in a multivariate regression analysis, previous preterm delivery, the maternal carrier status of the factor-VII-121del/ins polymorphism (OR=1.7, 95% CI: 1.12-2.5, p=0.007) and the lower frequency of infant's factor-XIII-Val34Leu polymorphism (OR=0.53; 95% CI: 0.29-0.96; p=0.038) were found to be independently associated with preterm delivery [58]. The association between Polymorphisms in factor VII and preterm birth was also reported among Caucasian in the USA. This study included maternal and fetal DNA from 370 patients. For maternal data the strongest associations were found in genes in the complement-coagulation pathway related to decidual hemorrhage in preterm birth. In this pathway 3 of 6 genes examined had SNPs significantly associated with preterm birth, including factor V, factor VII, and tissue plasminogen activator. The single strongest effect was observed in tPA marker rs879293 with a significant allelic and genotypic association with preterm delivery (OR- 2.80, CI 1.77–4.44, for a recessive model). Finally, exploratory multi-locus analyses in the complement and coagulation pathway were performed and revealed a potentially significant interaction between a marker in Factor V (rs2187952) and Factor VII (rs3211719) (p<0.001); the authors concluded that "These results support a role for genes in both the coagulation and inflammation pathways, and potentially different maternal and fetal genetic risks for preterm birth"[59].

Collectively the evidence brought here suggest that genetic polymorphism of the coagulation genes may predispose a subset of women to an increased risk for preterm birth. What is the role of gene environmental interaction and in what mechanisms these polymorphisms of the coagulation genes affect the risk for preterm parturition are still unknown and are an area of future research.

Changes in maternal circulation: Increased thrombin generation in the maternal circulation, above that reported during normal pregnancy, has been reported in all the great obstetrical syndromes including preeclampsia [60-66], fetal growth restriction [61,62,67,68], fetal death [69], preterm labor (PTL) [30,70], and preterm PROM [30,69,71]. There are several possible explanations for the increased thrombin generation reported in women with preterm parturition: 1) increased activation of coagulation cascade in the maternal circulation due to pathological processes including bleeding or inflammation; and 2) depletion of anticoagulation proteins that subsequently leads to increased thrombin generation.

Increased activation of the coagulation cascade among women with preterm parturition is well supported by the current literature. Indeed, women with preterm PROM and preterm labor have a higher median maternal plasma concentration of thrombin-anti-thrombin (TAT) III complexes [30,70]. In addition, in women with preterm labor elevated maternal plasma TAT III concentration was associated with a higher chance to deliver within <7 days from admission [69] (figure 5, and figure 6). Median maternal plasma Tissue factor, concentration is higher in women with preterm PROM, but not in those with PTL, than in those with normal pregnancies [72]. Nevertheless, women with preterm labor as well as those with preterm PROM had both increased tissue factor activity in comparison to normal pregnant women.



Figure 5. Thrombin–antithrombin III (*TAT*) levels in control patients, patients with preterm labor who delivered within 3 weeks, and patients with preterm labor who delivered after 3 weeks. *Open diamonds,* Mean levels; *black error bars,* SD. **P* <.05, Student-Newman-Keuls method (from reference #70 with permission).



Figure 6. Maternal plasma TAT III concentration in women with preterm labor (PTL) and those with a Normal pregnancy (from reference #30 with permission).

The activation of the coagulation system in the placental and maternal compartment of patients with preterm parturition can result from the following underlying mechanisms: 1(decidual hemorrhage that leads to a retro-placental clot formation [73]; 2) intra-amniotic infection which can induce decidual bleeding and sub-clinical abruption [74], as well as increased intraamniotic TAT complexes [69]; and 3) an increased maternal systemic inflammatory response [75] that may activate the extrinsic pathway of coagulation due to the expression and release of tissue factor (TF) by activated monocytes [76]. These mechanisms result in an increased thrombin generation, which has been associated with the following: 1) stimulation of decidual cell secretion of matrix metalloproteinase (MMP) (i.e. MMP-1 and MMP-3) that can degrade the extracellular matrix of the chorioamniotic membranes [77,78] (as is in preterm PROM); and 2) myometrial activation and uterine contractions generation that may lead to preterm labor with or without rupture of membranes and a subsequent preterm delivery [70,79,80]. While thrombin is generated as a consequence of activation of the coagulation cascade, TF, the most powerful natural pro-coagulant, is abundant in the uterine decidua in the normal state [81,82], as part of an efficient hemostatic mechanism in the uterine wall, which is activated in the course of normal pregnancy during implantation[83] and after delivery[84]. However, this hemostatic mechanism may also be activated due to pathological decidual bleeding in pregnancies complicated by placental abruption [73,85] and intra-amniotic infection [74].

A novel mechanism that may lead to an increased thrombin generation in women with preterm parturition is depleted or insufficient anticoagulant proteins concentration. Indeed, women with preterm labor without intra-amniotic infection or inflammation and those with vaginal bleeding who delivered preterm had a lower median maternal plasma protein *Z*, a co-factor that participate in the inhibition of factor Xa, concentration than women with normal pregnancy and those with vaginal bleeding who delivered at term [86]. Moreover, both patients with preterm labor and preterm PROM had a lower median maternal plasma concentration of total tissue factor pathway inhibitor (TFPI), the main physiological inhibitor of the TF pathway, regardless of the presence of infection or gestational age at delivery. These observations suggest that the increased thrombin generation observed among these patients may derive not only from an increased activation of the hemostatic system, but also from insufficient anti-coagulation. The latter can be due to either low concentrations of the anticoagulant proteins, or as a result of an abnormal balance between coagulation factors and their inhibitors.

The overall balance between the concentration and activity of the coagulation factors and the anti-coagulation proteins is one of the determining factors of thrombin generation. In the normal state, the immunoreactive concentrations of TFPI in the plasma are 500 to 1000 times higher than that of TF [87], suggesting that an excess of anti-coagulant proteins closely controls the coagulation cascade activity [87]. Although preterm labor was not associated with a significant change in the median maternal plasma tissue factor concentration, the TFPI/TF ratio was lower than that of normal pregnant women, mainly due to decreased TFPI concentrations. Along with the reports that patients with preterm PROM [72], as well as those with preeclampsia [88], have a lower median maternal plasma TFPI/TF ratio than that of normal pregnant women. The lower TFPI/TF ratio in patients with preeclampsia occurs despite the increase in the median maternal plasma TFPI concentration observed in these patients. This

suggests that the balance between TF and its natural inhibitor may better reflect the overall activity of the TF pathway of coagulation, than the individual concentrations of TF or TFPI. Collectively, these observations suggest that our attention should be focused not only on the coagulation protein but also on their inhibitors since an imbalance between them may contribute to increased thrombin generation leading to the activation of preterm parturition.

Inflammation is a major process in term and pre term parturition. In recent years it has become apparent that tight and reciprocal interactions exist between coagulation and inflammation [90]. Originally, much attention was given to mechanisms by which inflammatory mediators, most notably cytokines, can activate coagulation. More recent investigations have revealed that, in turn, mediators involved in the regulation of coagulation and anticoagulation have major effects on the inflammatory processes.

Inflammation elicits coagulation primarily by activating the tissue factor pathway [89] and the generation of thrombin and fibrin. The pivotal role of tissue factor in activation of coagulation during a systemic inflammatory response syndrome, such as produced by endotoxemia or severe sepsis, is well established, and attenuation of the activation of the tissue factor/factor VIIa pathway in endotoxemic humans and chimpanzees and in bacteremic baboons abrogated the activation of the common pathway of coagulation [90-92] and decreased the morbidity associated with these systemic inflammatory conditions.

During preterm PROM and preterm labor, there is a moderate maternal systemic inflammation that results in monocyte and granulocyte activation [75]. Activated monocytes express TF on their membrane [93-97] and shed micro-particles containing TF into the plasma [93]. In addition, the lack of association between intra-amniotic infection/inflammation, as well as the placental histologic findings and median maternal plasma concentrations of TF and TFPI, suggest that the pro-coagulant changes observed in patients with preterm PROM may be due to a systemic rather than a local (i.e. placental, intrauterine) inflammatory process.

Moreover, preterm PROM is associated with an increased activation of the decidual component of the common pathway of parturition [98]. Thus, in pregnancies complicated by abnormal placentation or intrauterine infection, decidual bleeding may lead to a higher expression of TF and activation of the coagulation cascade, resulting in increased thrombin generation. The latter has uterotonic properties that may generate uterine contractions that could initiate labor [70,79,80]. Moreover, thrombin can mediate the activation of MMP-1 [78], MMP-3 [77], and MMP-9[99] that can digest components of the extracellular matrix, weaken the chorioamniotic membranes and predispose to preterm PROM.

The mechanisms described above are localized to the maternal-fetal interface. The lack of association between median maternal plasma TF concentrations and the presence of intraamniotic infection/inflammation or vaginal bleeding in patients with preterm PROM suggest that the systemic maternal inflammatory response during preterm PROM[75] may contribute the increase median maternal plasma TF concentration in these patients regardless to the presence of infection or inflammation in the amniotic cavity or the occurrence of vaginal bleeding [100]. In addition to the maternal circulation intra-amniotic infection and/or inflammation is associated with an increased amniotic fluid TAT III complexes (figure 7). This is important since it represent an increased thrombin generation in the amniotic cavity during infection and or inflammation that may contribute to uterine contractility and the development of preterm birth [99]. Of interest, elevated intra-amniotic TAT III concentrations were associated with a shorter amniocentesis to delivery interval and an earlier gestational age at delivery only inpatients with preterm labor without intra-amniotic infection or inflammation [99]. This observation suggest that in a subset of patients with preterm labor activation of the coagulation system can generate preterm parturition and delivery; while in those with intra-amniotic infection and or inflammation the activation of the coagulation and thrombin generation is a byproduct of the inflammatory process leading to preterm birth.



Figure 7. The effect of amniotic fluid thrombin-antithrombin (TAT) III concentrations on gestational age at delivery (from reference #69 with permission)

5. Placental vascular changes in women with preterm parturition

Accumulating evidence from studies of the placenta [101,102], uterine artery Doppler scans [103], and animal experiments [104], suggest a role for uteroplacental ischemia in preterm birth. Indeed, Arias et al reported that about 20% of the placentas of patients who delivered preterm following preterm labor or preterm PROM had vascular lesions [105].

The invasion of trophoblast cells into the decidual and myometrial segments of the spiral arteries is a key point of normal placentation. This process results in reversible changes of the normal spiral arteries wall architecture [106]. The "disappearance of the normal muscular and elastic structures of the arteries and their replacement by fibrinoid material in which trophoblast cells are embedded" was originally termed physiologic transformation by Brosens et al in 1967. This process progressively remodeled, starting from the end of the first trimester onward, the uterine spiral arteries to form dilated conduits lacking maternal vasomotor control, ensuring the delivery of a constant supply of blood to the maternal-fetal interface at an optimal velocity for nutrient exchange [107]. Notably, several coagulation components, such as TF and thrombomodulin, are involved not only in hemostasis but also with placental blood vessel differentiation [38,42], affecting thereby the generation of different pathological condition affecting pregnancy and parturition; A higher rate of failure of transformation of the spiral arteries was reported in placentas of patients with preterm labor and preterm PROM than in those of normal pregnant women. This lesion has been implicated in the increased vascular resistance in the placental bed and the reduction of blood flow to the intervillous space [108,109], this is considered as a marker for defective placentation. Failure of transformation of the spiral artery was first reported in women with preeclampsia [110]. Indeed, the extent of this lesion in placenta of women with preeclampsia is more extensive than what is detected in women with preterm labor or preterm PROM [111-113], suggesting when extensive failure of physiologic transformation is present, narrowed uteroplacental arteries predispose to reduced perfusion of the intervillous space, ischemia, and compensatory maternal hypertension. If these lesions are less extensive, the degree of ischemia may be insufficient to induce maternal hypertension, but may predispose to preterm labor/preterm delivery by itself or in association with other pathologic processes, such as intrauterine infection. A report that support the concept that clinical presentation is somewhat reflecting the extent of the disease is the study by Espinoza et al who found that women with an episode of preterm labor who delivered at term had a higher rate of SGA neonates and increased frequency of placental vascular lesions in comparison to those with preterm labor who delivered preterm [114]. This finding suggests that in some cases the vascular lesions that lead to development of preterm labor is not severe enough to cause preterm birth, however the "price" for the continuum of the pregnancy to term is decrease fetal growth.

6. The future — Hemostatic markers for preterm parturition?

In light of the association between maternal plasma TAT III concentrations in women with preterm labor and preterm birth within a week; and the association between amniotic fluid TAT III concentrations, the interval from amniocentesis to delivery, and gestational age at delivery. The question of the role of hemostatic markers as predictors for preterm birth is relevant. A preliminary report by Vidaeff et al found that increased concentrations of amniotic fluid TAT III concentration during mid-trimester amniocentesis of asymptomatic patients is associated with subsequent preterm delivery [115]. Aside the amniotic fluid new assays that

study the thrombin generation potential in maternal blood may offer similar answer in less invasive methods [116].

7. Conclusions

The understanding of the homeostasis system and coagulation process is crucial for understanding the physiological and pathological parturition. The significant impact of placentation abnormalities caused by those same changes in the haemostatic system, on maternal and fetal wellbeing is yet to be studied.

The aim of this chapter was to provide a window to the complexity of the normal homeostasis and pregnancy and a view of the different pathological conditions that may emerge during parturition. The inflammation as well as the coagulation and placental implantation are all part of the total picture of parturition.

Author details

Vered Klaitman, Ruth Beer-Wiesel, Tal Rafaeli, Moshe Mazor and Offer Erez

*Address all correspondence to: erezof@bgu.ac.il

Department of Obstetrics and Gynecology "B", Soroka University Medical Center, School of Medicine, Faculty of Health Sciences, Ben Gurion University of the Negev, Beer Sheva, Israel

References

- Rand JH, Wu XX, Potter BJ, Uson RR, Gordon RE. Co-localization of von Willebrand factor and type VI collagen in human vascular subendothelium. Am J Pathol 1993 Mar; 142(3):843-850.
- [2] Ruggeri ZM, Dent JA, Saldívar E. Contribution of Distinct Adhesive Interactions to Platelet Aggregation in Flowing Blood. Blood 1999 July 01;94(1):172-178.
- [3] Nurden AT, Nurden P. A review of the role of platelet membrane glycoproteins in the platelet-vessel wall interaction. Baillieres Clin Haematol 1993 Sep;6(3):653-690.
- [4] Pytela R, Pierschbacher MD, Ginsberg MH, Plow EF, Ruoslahti E. Platelet membrane glycoprotein IIb/IIIa: member of a family of Arg-Gly-Asp--specific adhesion receptors. Science 1986 Mar 28;231(4745):1559-1562.
- [5] Lockwood CJ. Pregnancy-associated changes in the hemostatic system. Clin Obstet Gynecol 2006 Dec;49(4):836-843.

- [6] Greer P., Forster J., Lukens JN., Rodgers GM., Paraskevas F., Glader B. Wintrobs clinical hematology. : Lippincott, Williams and Wilkins; 2004.
- [7] Key NS, Geng JG, Bach RR. Tissue factor; from Morawitz to microparticles. Trans Am Clin Climatol Assoc 2007;118:165-173.
- [8] Rao LVM, Pendurthi UR. Tissue Factor–Factor VIIa Signaling. Arteriosclerosis, Thrombosis, and Vascular Biology ;25(1):47-56.
- [9] Semeraro N, Colucci M. Tissue factor in health and disease. Thromb Haemost 1997 Jul; 78(1):759-764.
- [10] Mackman N, Sawdey MS, Keeton MR, Loskutoff DJ. Murine tissue factor gene expression in vivo. Tissue and cell specificity and regulation by lipopolysaccharide. Am J Pathol 1993 Jul;143(1):76-84.
- [11] Drake TA, Morrissey JH, Edgington TS. Selective cellular expression of tissue factor in human tissues. Implications for disorders of hemostasis and thrombosis. Am J Pathol 1989 May;134(5):1087-1097.
- [12] ADAMS RLC, BIRD RJ. Review article: Coagulation cascade and therapeutics update: Relevance to nephrology. Part 1: Overview of coagulation, thrombophilias and history of anticoagulants. Nephrology 2009;14(5):462-470.
- [13] Guller S, Tang Z, Ma YY, Di Santo S, Sager R, Schneider H. Protein composition of microparticles shed from human placenta during placental perfusion: Potential role in angiogenesis and fibrinolysis in preeclampsia. Placenta 2011 Jan;32(1):63-69.
- [14] Piccin A, Murphy WG, Smith OP. Circulating microparticles: pathophysiology and clinical implications. Blood Rev 2007 May;21(3):157-171.
- [15] Sturk-Maquelin KN, Nieuwland R, Romijn FP, Eijsman L, Hack CE, Sturk A. Pro- and non-coagulant forms of non-cell-bound tissue factor in vivo. J Thromb Haemost 2003 Sep;1(9):1920-1926.
- [16] Marlar R, Kleiss A, Griffin J. An alternative extrinsic pathway of human blood coagulation. Blood 1982 December 01;60(6):1353-1358.
- [17] Mackman N, Tilley RE, Key NS. Role of the Extrinsic Pathway of Blood Coagulation in Hemostasis and Thrombosis. Arteriosclerosis, Thrombosis, and Vascular Biology 2007 August 01;27(8):1687-1693.
- [18] Greenwalt TJ. The how and why of exocytic vesicles. Transfusion 2006;46(1):143-152.
- [19] Ranby M, Brandstrom A. Biological control of tissue plasminogen activator-mediated fibrinolysis. Enzyme 1988;40(2-3):130-143.
- [20] Schatz F, Lockwood CJ. Progestin regulation of plasminogen activator inhibitor type 1 in primary cultures of endometrial stromal and decidual cells. Journal of Clinical Endocrinology & Metabolism 1993 September 01;77(3):621-625.

- [21] Lanir N, Aharon A, Brenner B. Procoagulant and anticoagulant mechanisms in human placenta. Semin Thromb Hemost 2003 Apr;29(2):175-184.
- [22] Bouma BN, Meijers JC. New insights into factors affecting clot stability: A role for thrombin activatable fibrinolysis inhibitor (TAFI; plasma procarboxypeptidase B, plasma procarboxypeptidase U, procarboxypeptidase R). Semin Hematol 2004 Jan;41(1 Suppl 1):13-19.
- [23] Ku DH, Arkel YS, Paidas MP, Lockwood CJ. Circulating levels of inflammatory cytokines (IL-1 beta and TNF-alpha), resistance to activated protein C, thrombin and fibrin generation in uncomplicated pregnancies. Thromb Haemost 2003 Dec;90(6): 1074-1079.
- [24] Urano T, Ihara H, Takada Y, Nagai N, Takada A. The inhibition of human factor Xa by plasminogen activator inhibitor type 1 in the presence of calcium ion, and its enhancement by heparin and vitronectin. Biochim Biophys Acta 1996 Dec 5;1298(2):199-208.
- [25] Yuen PM, Yin JA, Lao TT. Fibrinopeptide A levels in maternal and newborn plasma. Eur J Obstet Gynecol Reprod Biol 1989 Mar;30(3):239-244.
- [26] Sorensen JD, Secher NJ, Jespersen J. Perturbed (procoagulant) endothelium and deviations within the fibrinolytic system during the third trimester of normal pregnancy. A possible link to placental function. Acta Obstet Gynecol Scand 1995 Apr;74(4): 257-261.
- [27] Walker MC, Garner PR, Keely EJ, Rock GA, Reis MD. Changes in activated protein C resistance during normal pregnancy. Am J Obstet Gynecol 1997 Jul;177(1):162-169.
- [28] Bellart J, Gilabert R, Miralles RM, Monasterio J, Cabero L. Endothelial cell markers and fibrinopeptide A to D-dimer ratio as a measure of coagulation and fibrinolysis balance in normal pregnancy. Gynecol Obstet Invest 1998;46(1):17-21.
- [29] de Boer K, ten Cate JW, Sturk A, Borm JJ, Treffers PE. Enhanced thrombin generation in normal and hypertensive pregnancy. Am J Obstet Gynecol 1989 Jan;160(1):95-100.
- [30] Chaiworapongsa T, Espinoza J, Yoshimatsu J, Kim YM, Bujold E, Edwin S, et al. Activation of coagulation system in preterm labor and preterm premature rupture of membranes. J Matern Fetal Neonatal Med 2002 11(6):368-373.
- [31] Yoneyama Y, Suzuki S, Sawa R, Otsubo Y, Power GG, Araki T. Plasma adenosine levels increase in women with normal pregnancies. Am J Obstet Gynecol 2000 May;182(5): 1200-1203.
- [32] Sheu JR, Hsiao G, Luk HN, Chen YW, Chen TL, Lee LW, et al. Mechanisms involved in the antiplatelet activity of midazolam in human platelets. Anesthesiology 2002 Mar; 96(3):651-658.
- [33] Bremme KA. Haemostatic changes in pregnancy. Best Pract Res Clin Haematol 2003 Jun;16(2):153-168.

- [34] Eichinger S, Weltermann A, Philipp K, Hafner E, Kaider A, Kittl EM, et al. Prospective evaluation of hemostatic system activation and thrombin potential in healthy pregnant women with and without factor V Leiden. Thromb Haemost 1999 Oct;82(4):1232-1236.
- [35] Faught W, Garner P, Jones G, Ivey B. Changes in protein C and protein S levels in normal pregnancy. Am J Obstet Gynecol 1995 Jan;172(1 Pt 1):147-150.
- [36] Craven CM, Chedwick LR, Ward K. Placental basal plate formation is associated with fibrin deposition in decidual veins at sites of trophoblast cell invasion. Am J Obstet Gynecol 2002 Feb;186(2):291-296.
- [37] Sheppard BL, Bonnar J. Uteroplacental hemostasis in intrauterine fetal growth retardation. Semin Thromb Hemost 1999;25(5):443-446.
- [38] Lockwood CJ, Krikun G, Runic R, Schwartz LB, Mesia AF, Schatz F. Progestinepidermal growth factor regulation of tissue factor expression during decidualization of human endometrial stromal cells. J Clin Endocrinol Metab 2000 Jan;85(1):297-301.
- [39] Beller FK, Ebert C. The coagulation and fibrinolytic enzyme system in pregnancy and in the puerperium. Eur J Obstet Gynecol Reprod Biol 1982 May;13(3):177-197.
- [40] Schatz F, Krikun G, Caze R, Rahman M, Lockwood CJ. Progestin-regulated expression of tissue factor in decidual cells: implications in endometrial hemostasis, menstruation and angiogenesis. Steroids 2003 Nov;68(10-13):849-860.
- [41] Clark P, Brennand J, Conkie JA, McCall F, Greer IA, Walker ID. Activated protein C sensitivity, protein C, protein S and coagulation in normal pregnancy. Thromb Haemost 1998 Jun;79(6):1166-1170.
- [42] Brenner B. Haemostatic changes in pregnancy. Thromb Res 2004;114(5–6):409-414.
- [43] Udagawa K, Miyagi Y, Hirahara F, Miyagi E, Nagashima Y, Minaguchi H, et al. Specific expression of PP5/TFPI2 mRNA by syncytiotrophoblasts in human placenta as revealed by in situ hybridization. Placenta 1998 Mar-Apr;19(2-3):217-223.
- [44] Udagawa K, Yasumitsu H, Esaki M, Sawada H, Nagashima Y, Aoki I, et al. Subcellular localization of PP5/TFPI-2 in human placenta: a possible role of PP5/TFPI-2 as an anticoagulant on the surface of syncytiotrophoblasts. Placenta 2002 Feb-Mar;23(2-3): 145-153.
- [45] Salem HT, Menabawey M, Seppala M, Shaaban MM, Chard T. Human seminal plasma contains a wide range of trophoblast-'specific' proteins. Placenta 1984 Sep-Oct;5(5): 413-417.
- [46] Salem HT, Chard T. Placental protein 5 (PP5): biological and clinical studies. Placenta Suppl 1982;4:103-114.
- [47] Lee JN, Salem HT, Chard T, Huang SC, Ouyang PC. Circulating placental proteins (hCG, SP1 and PP5) in trophoblastic disease. Br J Obstet Gynaecol 1982 Jan;89(1):69-72.

- [48] Rohde LH, Carson DD. Heparin-like glycosaminoglycans participate in binding of a human trophoblastic cell line (JAR) to a human uterine epithelial cell line (RL95). J Cell Physiol 1993 Apr;155(1):185-196.
- [49] Gonzalez M, Neufeld J, Reimann K, Wittmann S, Samalecos A, Wolf A, et al. Expansion of human trophoblastic spheroids is promoted by decidualized endometrial stromal cells and enhanced by heparin-binding epidermal growth factor-like growth factor and interleukin-1 beta. Mol Hum Reprod 2011 Jul;17(7):421-433.
- [50] Maruyama I, Bell CE, Majerus PW. Thrombomodulin is found on endothelium of arteries, veins, capillaries, and lymphatics, and on syncytiotrophoblast of human placenta. J Cell Biol 1985 Aug;101(2):363-371.
- [51] Leach L, Bhasin Y, Clark P, Firth JA. Isolation of endothelial cells from human term placental villi using immunomagnetic beads. Placenta 1994 Jun;15(4):355-364.
- [52] Fazel A, Vincenot A, Malassine A, Soncin F, Gaussem P, Alsat E, et al. Increase in expression and activity of thrombomodulin in term human syncytiotrophoblast microvilli. Placenta 1998 May;19(4):261-268.
- [53] Sood R, Weiler H. Embryogenesis and gene targeting of coagulation factors in mice. Best Pract Res Clin Haematol 2003 Jun;16(2):169-181.
- [54] Reverdiau P, Jarousseau AC, Thibault G, Khalfoun B, Watier H, Lebranchu Y, et al. Tissue factor activity of syncytiotrophoblast plasma membranes and tumoral trophoblast cells in culture. Thromb Haemost 1995 Jan;73(1):49-54.
- [55] Crawley JT, Gu JM, Ferrell G, Esmon CT. Distribution of endothelial cell protein C/ activated protein C receptor (EPCR) during mouse embryo development. Thromb Haemost 2002 Aug;88(2):259-266.
- [56] Hassan SS, Romero R, Tarca AL, Draghici S, Pineles B, Bugrim A, et al. Signature pathways identified from gene expression profiles in the human uterine cervix before and after spontaneous term parturition. Am J Obstet Gynecol 2007 Sep;197(3): 250.e1-250.e7.
- [57] Tsatas D, Baker MS, Rice GE. Differential expression of proteases in human gestational tissues before, during and after spontaneous-onset labour at term. J Reprod Fertil 1999 May;116(1):43-49.
- [58] Yu Y, Tsai HJ, Liu X, Mestan K, Zhang S, Pearson C, et al. The joint association between F5 gene polymorphisms and maternal smoking during pregnancy on preterm delivery. Hum Genet 2009 Jan;124(6):659-668.
- [59] Velez DR, Fortunato SJ, Thorsen P, Lombardi SJ, Williams SM, Menon R. Preterm birth in Caucasians is associated with coagulation and inflammation pathway gene variants. PLoS One 2008 Sep 26;3(9):e3283.

- [60] Schjetlein R, Abdelnoor M, Haugen G, Husby H, Sandset PM, Wisloff F. Hemostatic variables as independent predictors for fetal growth retardation in preeclampsia. Acta Obstet Gynecol Scand 1999 Mar;78(3):191-197.
- [61] Chaiworapongsa T, Yoshimatsu J, Espinoza J, Kim YM, Berman S, Edwin S, et al. Evidence of in vivo generation of thrombin in patients with small-for-gestational-age fetuses and pre-eclampsia. J Matern Fetal Neonatal Med 2002 Jun;11(6):362-367.
- [62] Hayashi M, Numaguchi M, Ohkubo N, Yaoi Y. Blood macrophage colony-stimulating factor and thrombin-antithrombin III complex concentrations in pregnancy and preeclampsia. Am J Med Sci 1998 Apr;315(4):251-257.
- [63] Kobayashi T, Terao T. Preeclampsia as chronic disseminated intravascular coagulation. Study of two parameters: thrombin-antithrombin III complex and D-dimers. Gynecol Obstet Invest 1987;24(3):170-178.
- [64] Kobayashi T, Tokunaga N, Sugimura M, Suzuki K, Kanayama N, Nishiguchi T, et al. Coagulation/fibrinolysis disorder in patients with severe preeclampsia. Semin Thromb Hemost 1999;25(5):451-454.
- [65] Kobayashi T, Sumimoto K, Tokunaga N, Sugimura M, Nishiguchi T, Kanayama N, et al. Coagulation index to distinguish severe preeclampsia from normal pregnancy. Semin Thromb Hemost 2002 Dec;28(6):495-500.
- [66] Hayashi M, Inoue T, Hoshimoto K, Negishi H, Ohkura T, Inaba N. Characterization of five marker levels of the hemostatic system and endothelial status in normotensive pregnancy and pre-eclampsia. Eur J Haematol 2002 Nov-Dec;69(5-6):297-302.
- [67] Hayashi M, Ohkura T. Elevated levels of serum macrophage colony-stimulating factor in normotensive pregnancies complicated by intrauterine fetal growth restriction. Exp Hematol 2002 May;30(5):388-393.
- [68] Ballard HS, Marcus AJ. Primary and secondary platelet aggregation in uraemia. Scand J Haematol 1972;9(3):198-203.
- [69] Erez O, Romer R, Vaisbuch E, Chaiworapongsa T, Kusanovic JP, Mazaki-Tovi S, et al. Changes in amniotic fluid concentration of thrombin–antithrombin III complexes in patients with preterm labor: Evidence of an increased thrombin generation. J Matern Fetal Neonatal Med 2009 22(11):971-982.
- [70] Elovitz MA, Baron J, Phillippe M. The role of thrombin in preterm parturition. Am J Obstet Gynecol 2001 Nov;185(5):1059-1063.
- [71] Rosen T, Kuczynski E, O'Neill LM, Funai EF, Lockwood CJ. Plasma levels of thrombinantithrombin complexes predict preterm premature rupture of the fetal membranes. J Matern Fetal Med 2001 Oct;10(5):297-300.
- [72] Erez O, Espinoza J, Chaiworapongsa T, Gotsch F, Kusanovic JP, Than NG, et al. A link between a hemostatic disorder and preterm PROM: a role for tissue factor and tissue factor pathway inhibitor. J Matern Fetal Neonatal Med 2008 21(10):732-744.

- [73] Lockwood CJ, Toti P, Arcuri F, Paidas M, Buchwalder L, Krikun G, et al. Mechanisms of abruption-induced premature rupture of the fetal membranes: thrombin-enhanced interleukin-8 expression in term decidua. Am J Pathol 2005 Nov;167(5):1443-1449.
- [74] Gomez R, Romero R, Nien JK, Medina L, Carstens M, Kim YM, et al. Idiopathic vaginal bleeding during pregnancy as the only clinical manifestation of intrauterine infection.J Matern Fetal Neonatal Med 2005 Jul;18(1):31-37.
- [75] Gervasi MT, Chaiworapongsa T, Naccasha N, Pacora P, Berman S, Maymon E, et al. Maternal intravascular inflammation in preterm premature rupture of membranes. J Matern Fetal Neonatal Med 2002 Mar;11(3):171-175.
- [76] Osterud B, Bjorklid E. Sources of tissue factor. Semin Thromb Hemost 2006 Feb;32(1): 11-23.
- [77] Mackenzie AP, Schatz F, Krikun G, Funai EF, Kadner S, Lockwood CJ. Mechanisms of abruption-induced premature rupture of the fetal membranes: Thrombin enhanced decidual matrix metalloproteinase-3 (stromelysin-1) expression. Am J Obstet Gynecol 2004 Dec;191(6):1996-2001.
- [78] Rosen T, Schatz F, Kuczynski E, Lam H, Koo AB, Lockwood CJ. Thrombin-enhanced matrix metalloproteinase-1 expression: a mechanism linking placental abruption with premature rupture of the membranes. J Matern Fetal Neonatal Med 2002 Jan;11(1): 11-17.
- [79] Elovitz MA, Ascher-Landsberg J, Saunders T, Phillippe M. The mechanisms underlying the stimulatory effects of thrombin on myometrial smooth muscle. Am J Obstet Gynecol 2000 Sep;183(3):674-681.
- [80] Elovitz MA, Saunders T, Ascher-Landsberg J, Phillippe M. Effects of thrombin on myometrial contractions in vitro and in vivo. Am J Obstet Gynecol 2000 Oct;183(4): 799-804.
- [81] Lockwood CJ, Krikun G, Schatz F. Decidual cell-expressed tissue factor maintains hemostasis in human endometrium. Ann N Y Acad Sci 2001 Sep;943:77-88.
- [82] Lockwood CJ, Krikun G, Schatz F. The decidua regulates hemostasis in human endometrium. Semin Reprod Endocrinol 1999;17(1):45-51.
- [83] Lockwood CJ, Schatz F. A biological model for the regulation of peri-implantational hemostasis and menstruation. J Soc Gynecol Investig 1996 Jul-Aug;3(4):159-165.
- [84] Hahn L. On fibrinolysis and coagulation during parturition and menstruation. Acta Obstet Gynecol Scand Suppl 1974;28:7-40.
- [85] Major CA, de Veciana M, Lewis DF, Morgan MA. Preterm premature rupture of membranes and abruptio placentae: is there an association between these pregnancy complications? Am J Obstet Gynecol 1995 Feb;172(2 Pt 1):672-676.
- [86] Kusanovic JP, Espinoza J, Romero R, Hoppensteadt D, Nien JK, Kim CJ, et al. Plasma protein Z concentrations in pregnant women with idiopathic intrauterine bleeding and

in women with spontaneous preterm labor. J Matern Fetal Neonatal Med 2007 Jun;20(6): 453-463.

- [87] Shimura M, Wada H, Wakita Y, Nakase T, Hiyoyama K, Nagaya S, et al. Plasma tissue factor and tissue factor pathway inhibitor levels in patients with disseminated intravascular coagulation. Am J Hematol 1997 Aug;55(4):169-174.
- [88] Erez O, Romero R, Hoppensteadt D, Than NG, Fareed J, Mazaki-Tovi S, et al. Tissue factor and its natural inhibitor in pre-eclampsia and SGA. J Matern Fetal Neonatal Med 2008 Dec;21(12):855-869.
- [89] Levi M, van der Poll T. Inflammation and coagulation. Crit Care Med 2010 Feb;38(2 Suppl):S26-34.
- [90] van der Poll T, Levi M, Hack CE, ten Cate H, van Deventer SJ, Eerenberg AJ, et al. Elimination of interleukin 6 attenuates coagulation activation in experimental endotoxemia in chimpanzees. J Exp Med 1994 Apr 1;179(4):1253-1259.
- [91] van der Poll T, de Boer JD, Levi M. The effect of inflammation on coagulation and vice versa. Curr Opin Infect Dis 2011 Jun;24(3):273-278.
- [92] Schouten M, Wiersinga WJ, Levi M, van der Poll T. Inflammation, endothelium, and coagulation in sepsis. J Leukoc Biol 2008 Mar;83(3):536-545.
- [93] Butenas S, Bouchard BA, Brummel-Ziedins KE, Parhami-Seren B, Mann KG. Tissue factor activity in whole blood. Blood 2005 Apr 1;105(7):2764-2770.
- [94] Osterud B. Cellular interactions in tissue factor expression by blood monocytes. Blood Coagul Fibrinolysis 1995 Jun;6 Suppl 1:S20-5.
- [95] Rivers RP, Hathaway WE, Weston WL. The endotoxin-induced coagulant activity of human monocytes. Br J Haematol 1975 Jul;30(3):311-316.
- [96] Bach RR, Moldow CF. Mechanism of tissue factor activation on HL-60 cells. Blood 1997 May 1;89(9):3270-3276.
- [97] Egorina EM, Sovershaev MA, Bjorkoy G, Gruber FX, Olsen JO, Parhami-Seren B, et al. Intracellular and surface distribution of monocyte tissue factor: application to intersubject variability. Arterioscler Thromb Vasc Biol 2005 Jul;25(7):1493-1498.
- [98] Romero R, Espinoza J, Kusanovic JP, Gotsch F, Hassan S, Erez O, et al. The preterm parturition syndrome. BJOG 2006 Dec;113 Suppl 3:17-42.
- [99] Stephenson CD, Lockwood CJ, Ma Y, Guller S. Thrombin-dependent regulation of matrix metalloproteinase (MMP)-9 levels in human fetal membranes. J Matern Fetal Neonatal Med 2005 Jul;18(1):17-22.
- [100] Erez O, Romero R, Vaisbuch E, Kusanovic JP, Mazaki-Tovi S, Chaiworapongsa T, et al. High tissue factor activity and low tissue factor pathway inhibitor concentrations in patients with preterm labor. J Matern Fetal Neonatal Med 2010 23(1):23-33.

- [101] Germain AM, Carvajal J, Sanchez M, Valenzuela GJ, Tsunekawa H, Chuaqui B. Preterm labor: placental pathology and clinical correlation. Obstet Gynecol 1999 Aug;94(2): 284-289.
- [102] Arias F, Rodriquez L, Rayne SC, Kraus FT. Maternal placental vasculopathy and infection: two distinct subgroups among patients with preterm labor and preterm ruptured membranes. Am J Obstet Gynecol 1993 Feb;168(2):585-591.
- [103] Strigini FA, Lencioni G, De Luca G, Lombardo M, Bianchi F, Genazzani AR. Uterine artery velocimetry and spontaneous preterm delivery. Obstet Gynecol 1995 Mar;85(3): 374-377.
- [104] Combs CA, Katz MA, Kitzmiller JL, Brescia RJ. Experimental preeclampsia produced by chronic constriction of the lower aorta: validation with longitudinal blood pressure measurements in conscious rhesus monkeys. Am J Obstet Gynecol 1993 Jul;169(1): 215-223.
- [105] Arias F, Victoria A, Cho K, Kraus F. Placental histology and clinical characteristics of patients with preterm premature rupture of membranes. Obstet Gynecol 1997 Feb; 89(2):265-271.
- [106] Pijnenborg R, Bland JM, Robertson WB, Brosens I. Uteroplacental arterial changes related to interstitial trophoblast migration in early human pregnancy. Placenta 1983 Oct-Dec;4(4):397-413.
- [107] Brosens I, Robertson WB, Dixon HG. The physiological response of the vessels of the placental bed to normal pregnancy. J Pathol Bacteriol 1967 Apr;93(2):569-579.
- [108] Kim YM, Bujold E, Chaiworapongsa T, Gomez R, Yoon BH, Thaler HT, et al. Failure of physiologic transformation of the spiral arteries in patients with preterm labor and intact membranes. Am J Obstet Gynecol 2003 Oct;189(4):1063-1069.
- [109] Kim YM, Chaiworapongsa T, Gomez R, Bujold E, Yoon BH, Rotmensch S, et al. Failure of physiologic transformation of the spiral arteries in the placental bed in preterm premature rupture of membranes. Am J Obstet Gynecol 2002 Nov;187(5):1137-1142.
- [110] Brosens IA, Robertson WB, Dixon HG. The role of the spiral arteries in the pathogenesis of preeclampsia. Obstet Gynecol Annu 1972;1:177-191.
- [111] Khong Y, Brosens I. Defective deep placentation. Best Pract Res Clin Obstet Gynaecol 2011 Jun;25(3):301-311.
- [112] Espinoza J, Romero R, Mee Kim Y, Kusanovic JP, Hassan S, Erez O, et al. Normal and abnormal transformation of the spiral arteries during pregnancy. J Perinat Med 2006;34(6):447-458.
- [113] Brosens I, Pijnenborg R, Vercruysse L, Romero R. The "Great Obstetrical Syndromes" are associated with disorders of deep placentation. Am J Obstet Gynecol 2011 Mar; 204(3):193-201.

- [114] Espinoza J, Kusanovic JP, Kim CJ, Kim YM, Kim JS, Hassan SS, et al. An episode of preterm labor is a risk factor for the birth of a small-for-gestational-age neonate. Am J Obstet Gynecol 2007 Jun;196(6):574.e1-5; discussion 574.e5-6.
- [115] Vidaeff AC, Monga M, Saade G, Bishop K, Ramin SM. Prospective investigation of second-trimester thrombin activation and preterm birth. Am J Obstet Gynecol 2012
 Apr;206(4):333.e1-333.e6.
- [116] Hackney DN, Catov JM, Simhan HN. Low concentrations of thrombin-inhibitor complexes and the risk of preterm delivery. Am J Obstet Gynecol 2010 Aug;203(2): 184.e1-184.e6.





IntechOpen