

# We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

185,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index  
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?  
Contact [book.department@intechopen.com](mailto:book.department@intechopen.com)

Numbers displayed above are based on latest data collected.  
For more information visit [www.intechopen.com](http://www.intechopen.com)



---

## Non-Thyroidal Illness: Physiopathology and Clinical Implications

---

Antonio Mancini, Sebastiano Raimondo,  
Chantal Di Segni, Mariasara Persano and  
Alfredo Pontecorvi

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/55644>

---

### 1. Introduction

In critical illness, several abnormalities in thyroid hormone (TH) secretion, metabolism and action have been described in patients without previous diagnosis of intrinsic thyroid disease and are collectively called “Non thyroidal syndrome” (NTIS) [1]; this term is now largely employed, in the place of “euthyroid sick syndrome” [2-4] or “low- $T_3$  syndrome”, due to the most common abnormality, a decreased level of serum total triiodothyronine ( $T_3$ ), which can be detected very early, within 2 hours after the onset of severe physical stress [5-7]. However,  $T_3$  lowering is only one of the endocrine picture described in such a situation; therefore the term NTIS seems to be more appropriate, also strengthening its extrathyroidal source.

NTIS has been depicted in about 70% of hospitalized patients for different diseases [8-10]. Moreover, the severity of morbidity and outcome in patients studied in intensive care unit (ICU) has been correlated with the alteration in thyroid function [11,12]. The hormonal response exhibits different pattern in acute and chronic phase, since in the first phase the alteration predominates in peripheral metabolism of TH, while in the latter central mechanisms controlling thyroid secretion progressively arise [13,14].

Since there is no clear evidence of tissue hypothyroidism, such a condition seems to be an adaptive response, and thyroid replacement therapy is not usually required, but this topic is still debated, since indirect signs of true hypothyroidism at tissue levels have been showed [15]. The question is open and different reviews have been published on this topic [1, 16-20]; but, very recently different molecular mechanisms have been shown to gain insight into the complex situation of NTIS. The role of intracellular oxidative stress (OS) has

been underlined. Therefore we present a review of these recent results and some personal data in patients affected by chronic obstructive pulmonary disease and patients studied after major cardiovascular surgery.

## 2. Clinical observations

A low  $T_3$  state has been described in a variety of clinical situations, such as starvation [21], sepsis [22], surgery [23], trauma [24], myocardial infarction and heart failure [25,26], cardiopulmonary bypass [27], respiratory failure [28], bone marrow transplantation [29], other severe illness [30]. In a very recent paper in unselected ICU patients, free  $T_3$  ( $fT_3$ ) was the most powerful and the only independent predictor of ICU mortality, with a prognostic improving value when added to APACHE II score [31]. A retrospective study in a large group of patients treated with mechanical ventilation (MV) confirmed that NTIS represents a risk factor for prolonged MV [32].

Due to the importance of TH in **cardiac function**, it is not surprising that cardiac patients have been extensively studied under this profile. TH influence cardiac function with different mechanisms: inotropic and chronotropic positive effect via nuclear and non-nuclear pathways in cardiomyocytes, increase in cardiac contractility through augmented tissue oxygen delivery and consumption; decrease in systemic vascular resistance, through direct TH action on vascular smooth muscle cells; other endocrine effects are exerted on renin-angiotensin-aldosterone axis and on erythropoietin secretion [19, 33].

One of the early studies was performed in patients serially followed after acute myocardial infarction; a sustained and prolonged decrease of total  $T_3$  ( $TT_3$ ) and  $fT_3$  was described, while  $TT_4$  but not  $fT_4$  showed a transient decrease; thyroxine binding globulin (TBG) levels remained unchanged, while thyroxine binding prealbumin (TBPA) and albumin exhibited a prolonged fall. TSH, despite low  $T_3$ , did not increase, remaining inappropriately low [34]. In this sense, the increase of TSH was shown to be correlated with a good prognosis [35].

It has been reported that patients with heart failure have low  $T_3$  serum concentrations, which correlate with cardiac function [36]. In advanced heart failure, a low  $fT_3$  index/reverse  $T_3$  ratio was associated with higher right atrial pulmonary artery and pulmonary capillary wedge and lower ejection fraction [26].

Low  $T_3$  syndrome has been considered a strong predictor of death and directly implicated in poor prognosis of cardiac patients in a large group of patients admitted in a cardiology department [37].

TH are implicated in metabolic function of myocardial cells; they have been shown to inversely correlate with Coenzyme  $Q_{10}$  ( $CoQ_{10}$ ), a component of mitochondrial respiratory chain, also endowed with powerful antioxidant properties [38]. Preliminary data of our group in patients studied after major heart surgery showed low  $T_3$  levels concurrently with signs of tissue hypothyroidism (elevated  $CoQ_{10}$  levels) [39]. In fact we found  $CoQ_{10}$  levels, evaluated by high

performance liquid chromatography (HPLC), in the hypothyroid range, despite the fact cardiac diseases are well known to be associated with low CoQ<sub>10</sub>.

The studies in **pulmonary disorders** have not been so extensively investigated [40-42]. In the just cited paper, low T<sub>3</sub> state was again considered a predictor of outcome in respiratory failure [28]. Among chronic conditions, no conclusive data are reported on chronic obstructive pulmonary disease (COPD), as reported in a recent review [43]. No clear evidence of thyroid function alteration has been reported in such a condition [44], although in patients with severe hypoxemia a strong positive correlation between total T<sub>3</sub>/total T<sub>4</sub> ratio (TT<sub>3</sub>/TT<sub>4</sub>) and PaO<sub>2</sub> has been described [45]. Increased fT<sub>3</sub> concentrations have been reported in stable COPD, with a positive association to PaCO<sub>2</sub> [46], while others reported lower total T<sub>3</sub>, fT<sub>3</sub> and TT<sub>3</sub>/TT<sub>4</sub> ratios in patients with severe hypoxemia [47]. Low Forced Expiratory Volume at 1<sup>st</sup> second (FEV<sub>1</sub>) is associated with low basal and stimulated levels of thyroid stimulating hormone (TSH) [48]; however the impact of hypoxemia on TSH response to exogenous thyrotropin releasing hormone (TRH) is controversial [45,46].

We have recently studied patients with COPD, evaluating lung parameters and antioxidant parameters, due to a possible involvement of OS in NTIS (see below). COPD is a complex condition, which cannot be considered a lung-related disorder, but rather a systemic disease also associated to increased oxidative stress. We evaluated thyroid hormones and antioxidant systems, the lipophilic CoQ<sub>10</sub> and total antioxidant capacity (TAC) in COPD patients to reveal the presence of a low-T<sub>3</sub> syndrome in COPD and investigate the correlation between thyroid hormones, lung function parameters and antioxidants. The evaluation of CoQ<sub>10</sub> was particularly interesting, also for the energetic role of this molecule, which is a component of the mitochondrial respiratory chain, as above stated; its concentrations were also corrected for cholesterol, due to its lipophilic nature. We studied 32 COPD patients and 45 controls; CoQ<sub>10</sub> was assayed by HPLC; TAC by the metmyoglobin-ABTS method and expressed as latency time (LAG) in radical species appearance. We found significantly lower LAG values, fT<sub>3</sub> and fT<sub>4</sub> levels and significantly higher TSH in COPD patients vs controls. LAG values significantly correlated with fT<sub>3</sub> concentration. Twelve out of 32 patients exhibited fT<sub>3</sub> levels lower than normal range. When dividing COPD patients in two groups on the basis of the fT<sub>3</sub> concentration (normal fT<sub>3</sub> COPD and low fT<sub>3</sub> COPD), we observed lower LAG values in normal fT<sub>3</sub>-COPD, compared to healthy subjects, with a further significant reduction in low fT<sub>3</sub>-COPD patients. Moreover higher TSH concentrations were present in normal fT<sub>3</sub>-COPD, compared to healthy subjects, with a further significant increase in low fT<sub>3</sub>-COPD patients. CoQ<sub>10</sub>/cholesterol ratio was higher in low fT<sub>3</sub>-COPD vs normal fT<sub>3</sub>-COPD, with a nearly significant difference. These data seem to indicate an increased oxidative stress in low fT<sub>3</sub>-COPD and a role of fT<sub>3</sub> in modulating antioxidant systems. However low fT<sub>3</sub> levels are joined to metabolic indexes of true hypothyroidism, suggesting that elevated CoQ<sub>10</sub> expresses a reduced tissue utilization. Interestingly, there was no significant difference in lung parameters when comparing normal- or low-fT<sub>3</sub> COPD patients, according to the definition of COPD as a systemic disease, with respiratory parameters unable to define the severity of disease. In fact metabolic dysfunctions (i.e. osteoporosis, vascular and cardiac involvement, muscle impairment) play a role in the natural history of disease but were found poorly related to respiratory impairment,

underlying the need of indexes related to a real tissue condition; the pattern of  $fT_3$  could indicate such a situation, as reinforced by the pattern of  $CoQ_{10}$  levels; decreased plasma antioxidant capacity and increased  $CoQ_{10}$  levels in low  $fT_3$ -COPD again suggested a possible condition of hypothyroidism at tissue levels [49].

The thyroid function has been investigated in patients with acute **kidney injury**. TSH levels inversely correlated with urea concentrations. 82.9% of patients exhibited alteration in thyroid function, especially low- $T_3$ . This picture was ameliorated by improvement of renal function. No prognostic role was attributed to this dysfunction [50].


Primary hypothyroidism (non-autoimmune) is often observed in patients with chronic kidney disease (CKD); in particular the prevalence of subclinical hypothyroidism is related to GRF decline [51]. The earliest and the most common thyroid function abnormality in CKD patients is a low  $T_3$  level (especially  $TT_3$  than  $fT_3$ ) [52]. The mechanisms for  $T_3$  decrease in this condition are: fasting, chronic metabolic acidosis and chronic protein malnutrition, influencing  $T_4$  deiodination, as well as protein binding of  $T_3$ . Moreover, inflammatory cytokines such as tumor necrosis factor (TNF)- $\alpha$  and interleukin (IL)-1 inhibit the expression of type 1 5'-deiodinase (see below), which is responsible for peripheral conversion of  $T_4$  to  $T_3$  [53]. Alteration of renal handling of iodine can increase serum iodine levels, causing a prolonged Wolff – Chaikoff effect [54]. A prognostic role has been attributed to the hormonal marker: the low  $fT_3$  levels in CKD patients have been correlated with higher levels of markers of inflammation [highly sensitive C-reactive protein (hsCRP), IL-6, etc.], malnutrition (lower prealbumin, IGF-1), increased endothelial dysfunction, poorer cardiac function, poor survival, and higher all-cause as well as cardiovascular mortality in some studies [53, 55].

Little is known in TH alterations in acute **liver failure** (ALF) [56]. An animal model was investigated (pigs subjected to surgical liver devascularisation). In this case serum  $T_4$  and  $T_3$  levels were markedly decreased, but  $fT_3$  and TSH did not change. The downregulation of  $T_4$  and  $T_3$  levels during ALF seems to correlate well with the severity of disease and was also related to alteration in parameters of inflammation, oxidative stress and myocardial thyroid receptors; thus the mechanisms in this case seem to be very complex. In humans acute liver failure (ALF) is accompanied by hormonal implications, as has been recently shown for the hepatoadrenal syndrome [57], since an unexpected incidence of adrenal failure was discovered in ALF and post-transplantation patients; a glucocorticoid treatment can influence outcome. Thyroid function alterations have been described during chronic liver failure [58-60]; a low  $T_4$ -variant of NTIS has been described in a subgroup of patients with cirrhosis at risk for decreased survival [58]; serum levels of  $fT_3$  and  $TT_4$  (but not  $TT_3$  and  $fT_4$ ) were significantly lower in patients with hepatic encephalopathy compared to decompensated cirrhotic patients without encephalopathy [60]. Much less clear data are available for ALF [61, 62]. In cirrhotic and also in acutely ill patients from various etiologies, derangements of thyroid hormones are common (up to 79% in the latter group, as reported from autoptic observations) [63].

Finally, during **starvation** (especially carbohydrate deprivation) deiodination of  $T_4$  to  $T_3$  is rapidly inhibited, causing the low- $T_3$  syndrome [1, 21, 64]. Interestingly, caloric deprivation can be also a major factor influencing TH in severe illness, as demonstrated in bacterial sepsis [65].

A particular model is that of eating disorders, especially anorexia nervosa, in which low- $T_3$  is accompanied by a constellation of hormone alteration, index of hypothalamic derangement [66]. Other psychiatric models should be considered with caution when evaluating thyroid alterations, due to other interfering factors, such as the underlying psychiatric disorder, substance abuse or other medications [67].

On the basis of the reported studies and other reviews [68, 69] we can summarize the main variations in the pituitary-thyroid axis as reported in Fig. 1, according to the severity of NTIS.



**Figure 1.**

### 3. Physiopathological mechanisms

Various mechanisms are responsible for the TH pattern observed in different situations, keeping in mind the difference between “acute” and “chronic” phases and possible differences related to the underlying diseases. They can summarize in four categories: central TSH regulation, TH blood transportation, peripheral metabolism by deiodinases, actions at receptorial and post-receptorial levels.

#### a. Central regulation of TSH

Basal TSH levels are usually normal or low, but not extremely inhibited [1, 70, 71]; in most cases they are inadequate in respect to thyroid hormone levels. The response to TRH is variable, ranging from blunted to normal response [72, 73]; the response to TRH, even in presence of

low basal level, can be interpreted as a sign of hypothalamic dysfunction, according to data concerning other hormones (gonadotropins, ACTH-cortisol axis) [13, 74]. Absence of circadian rhythm has been reported [75]. The variation of glycosilation is responsible for reduced bioactivity [76]. The finding that TH alterations are partially reversed by the combined infusion of TRH and GH secretagogues [77] reinforces the role of central component of NTIS.

#### **b. Transportation**

Also transportation of thyroid hormones is altered; Thyroxine binding globulin (TBG) has been shown to be reduced, probably for increased cleavage by proteases. The binding to transport protein is also negatively influenced by inhibitors (not only in serum, but also in tissues), therefore influencing the metabolism of TH [1]. Recently, decrease of TBG, determined by RIA or radioimmunoassay, albumin and transthyretin (TTR) have been described in septic patients [78]; therefore the total binding power of serum is low, in the view of authors, without the need to postulate the effects of additional factors, such as binding inhibitors or modification of binding affinity.

#### **c. Deiodinase**

A lot of studies concern the activity of deiodinases, the main group of enzymes, which by removal of iodine, catalyze activation or inactivation of TH. They are selenoproteins, members of the thioredoxin family, and require a thiol cofactor for their activity [20,79]. The activation of prohormone  $T_4$  into the biologically active hormone  $T_3$  is catalyzed by type 1 (D1, encoded by DIO1) and type 2 (D2, encoded by DIO2) via deiodination of the outer ring; on the contrary, the removal of inner ring iodine is catalyzed by type 3 (D3, encoded by DIO3), causing inactivation of both  $T_4$  and  $T_3$  [80]. In humans, 80% of circulating  $T_3$  comes from deiodination by D1 and D2, while the other 20% comes directly from thyroid secretion. The most common alteration in NTIS patients is a decrease in  $T_3$ , caused by reduced conversion of  $T_4$  to  $T_3$  [81]. The Deiodinase 1 is down regulated, as demonstrated in liver, causing reduced  $T_3$  generation [1]. Deiodinase 3 is instead increased, as observed in liver and muscle, especially in the case of low tissue perfusion, and the conversion of  $T_3$  to reverse- $T_3$  ( $rT_3$ ) is a mechanism reinforcing the low  $T_3$  levels [80]. However, central and peripheral deiodinases are differently regulated;  $T_3$  in the pituitary are normal since local deiodination is enhanced, thus the pituitary is actually euthyroid and therefore TSH circulating levels inappropriate to other tissue  $fT_3$  levels [1]. The role of D3 has recently been reviewed [82]. Moreover more recent studies focused on modulation of deiodinases activity, rather than their levels (see below).

#### **d. Thyroid receptors and Postreceptorial mechanisms**

It has been shown that thyroid hormone receptors (TR) are down-regulated in skeletal muscle of patients with non-septic shock; in particular they showed lower expression of TR- $\beta$ , TR- $\alpha_1$  and their nuclear partner retinoid X receptor  $\gamma$  (RXRG) [83]. The RXRA gene expression was higher, even if its protein was lower, suggesting the existence of post-transcriptional mechanisms that down-regulate protein levels. Nuclear factor of kappa light chain enhancer of activated B cells (NkFB), a transcriptional factor involved in immune and inflammatory response, attenuates the induction of DIO1 by  $T_3$  [84]; however NkFB1 activation was not

different in comparison to control subjects. However the results are not unequivocal, since there results were not reproducible in cultures of human smooth muscle cells (HSkMC) incubated with the patients' serum [83].

Molecular mechanisms of thyroid action in NTIS have been recently investigated in other models, studying, other than TR, also the transporters, which allow TH to be transported across the plasma membrane in order to be metabolized and interact with their receptors. Monocarboxylate transporter 8 (MCT8) has been shown to be a very active and specific transporter [85]. Moreover, other proteins modulate the transcription function of TR, acting as coactivators or corepressors; among the latter the silencing mediator of retinoid and TR (SMRT) via histone deacetylation [86]. In patients with septic shock, skeletal muscle expression of TR- $\beta$ 1, RXRG and D2 was lower than in control group and RXRA was higher. In subcutaneous adipose tissue, the authors found lower MCT8, TRHB1, THRA1, RXRG and SMRT and higher UCP3 expression, suggesting decreased thyroid hormone action [87].

Interestingly, the reduced expression of TH transporters has been considered a compensatory mechanism (rather than a cause of low-  $T_3$ ), strongly suggesting a real hypothyroidism at tissue levels in such a condition [88].

#### 4. The role of cytokines

The role of cytokines, as key molecules involved in coordinating the hormone, immune and inflammatory response to a variety of stressful stimuli, has been largely investigated [1].

In a series of septic patients studied shortly after admission to an ICU,  $TT_4$ ,  $fT_4$ ,  $TT_3$  and TSH were depressed, and IL-1B, sIL-2R and  $TNF\alpha$  were elevated [89] suggesting central suppression of TSH, even if the relationship with cytokines was not so clear. The hypothalamic-pituitary-adrenal axis was activated as expected. It has been shown that continuous infusion of IL-1 in rats cause suppression of TSH,  $T_3$  and  $fT_4$ ; higher doses of IL-1 were accompanied by a febrile reaction and suppression of food intake, with a cascade of events altering thyroid hormone economy [90], but IL-1 did not reproduce the decrease in hepatic 5'-deiodinase activity believed to be characteristic of NTIS.

$TNF$  is another proinflammatory cytokine that is thought to be involved in many of the illnesses associated with NTI. Infusion of r $TNF$  in man produced a decrease in serum  $T_3$  and TSH and increase in r  $T_3$  [91]. These studies suggest that  $TNF$  could be involved in the IL-6-mediated activation of hypothalamic-pituitary axis. Also in this case other data did not confirm the role of  $TNF$ , since the effects of endotoxin of TH in humans were not counteracted by the  $TNF\alpha$  blockade by specific IgG fusion proteins [92].  $TNF\alpha$  was found during in vitro studies to activate  $NkFB$  [93], which in turn inhibits the  $T_3$ -induced expression of D1 as above reported.

On the contrary, an important role has been attributed to IL-6, which is often elevated in serum of NTIS patients [94] and its level is inversely related to  $T_3$  levels [95]. Short term infusion of rIL-6 to human volunteers [96] caused a suppression of TSH, but daily injections over 42 days cause only a modest decrease in  $T_3$  and a transient increase in r  $T_3$  and in  $fT_4$  concentrations.

More recent evidences on the role of IL-6 have been reported by studies in human cell lines: the effects of IL-6 on both endogenous cofactor-mediated and dithiotreitol-stimulated cell sonicate deiodinase activity have been studied [80]. In this model T<sub>3</sub> generation by D1 and D2 was suppressed by IL-6, despite an increase in sonicate deiodinases (and mRNAs): this inhibitory action was prevented by addition of N-acetyl-cysteine (NAC), an antioxidant that restores intracellular glutathione (GSH) concentrations. The interest of the paper is also the link of deiodinase activity and OS (see below).

Finally, the potential interaction between the complex network of cytokines and the hypothalamic pituitary thyroid axis, even if is not possible to build a simplistic model, probably plays a pathogenetic role in NTIS [1]. The role of cytokines in eating disorders and related TH alterations has also been reviewed [97].

## 5. Oxidative stress in NTIS

Previous studies have shown that both hyperthyroidism and hypothyroidism are associated with enhanced oxidative stress involving enzymatic and non-enzymatic antioxidants [98]. Besides, some complications of hyperthyroidism are due just to the oxidative stress in target tissues [99]. Thyroid hormones *per se* can act as oxidants and produce DNA-damage (contrasted by catalase), probably through the phenolic group, similar to that of steroidal estrogens [100]. Many other mechanisms, reviewed by Venditti & Di Meo [101], can be involved, with a specificity in tissue response. We recently reviewed the relationships between thyroid hormone, OS and reproduction [102].


At a systemic level, also in humans, hyperthyroidism has been associated with reduced circulating levels of alpha-tocopherol [103, 104] and Coenzyme Q<sub>10</sub> [38, 104]. Coenzyme Q<sub>10</sub> showed a trend to increase in hypothyroidism [38]; it appeared to be a sensitive index of tissue effect of thyroid hormones, in situations in which drug interference, such as amiodarone [105] or systemic illness inducing a low-T<sub>3</sub> conditions [106] complicate the interpretation of thyroid hormone levels. However, data on hypothyroidism in humans are conflicting [102]. Baskol et al showed in a group of 33 patients with primary hypothyroidism elevated malondialdehyde (MDA) and nitric oxide (NO) levels and low paraoxonase (PON1) activity, while superoxide dismutase (SOD) was not different from controls. Interestingly, thyroid treatment decreased MDA and increased PON1, without reaching levels observed in controls [107]. They concluded that a prooxidant environment in hypothyroidism could play a role in the pathogenesis of atherosclerosis in such patients. Elevated MDA levels were also shown in subclinical hypothyroidism [108]; the increased in OX was attributed to lack of antioxidants but also to altered lipid metabolism, since MDA showed a correlation with LDL-cholesterol, total cholesterol and triglycerides. Total antioxidant status (TAS) was similar in overt hypothyroidism, subclinical hypothyroidism and controls.

Another study [109] showed increased levels of thiobarbituric acid reactive substances (TBARS), but also of antioxidants, such as SOD, catalase (CAT) and Vitamin E. All these parameters correlated with T<sub>3</sub>; moreover the correlation between T<sub>3</sub> and CAT remained

significant also when corrected with total cholesterol. This datum was not confirmed by other authors [110, 111]. We showed low Total Antioxidant Capacity (TAC) levels in hypothyroid patients and increased CoQ<sub>10</sub> levels also in secondary hypothyroidism (mainly due to its metabolic role in mitochondrial respiratory chain and therefore underutilized in hypothyroid tissue). In the last case, hypothyroidism has a predominant effect on the possible decreasing effect of OS [112].

Different conditions with NTIS are associated to OS, due to augmented production of radical oxygen species (ROS) or nitrogen species [113]; since thyroid hormones, as above stated, can increase ROS generation, OS could be viewed as a compensatory mechanism since, decreasing metabolic rate, could protect against further radical generation. A reducing environment is maintained in the cytosol by intracellular thiols, especially GSH and Thioredoxin (TRX), which, as we have seen, are cofactors for deiodinases. Therefore their depletion, due to buffering effect against radical propagation, could interfere with the conversion of T<sub>4</sub> to T<sub>3</sub> [79]. Moreover, another reported mechanism is the nuclear sequestration of the SECIS binding protein 2 (SPB2), which reduces incorporation of selenocysteine residues in the selenoproteins [114]. IL-6 is known to induce OS, therefore an unifying hypothesis is cytokine-induced OS and a secondary alteration of expression and activity of deiodinases [79]. However further studies can clarify these complex interaction and especially the potential role of antioxidant in protecting against OS in NTIS.

On the basis of the physiopathological studies above reported, we can conclude that the alterations of pituitary-thyroid axes do not only depend from the severity of the disease, but also from the nutritional status of the patients and their inflammatory response, also related to oxidative stress (see fig. 2).



**Figure 2.** Interaction of factors influencing pituitary-thyroid axis

## 6. Treatment

Controversial results have been reported on the topic of replacement therapy. The replacement therapy with 1.5 µg/Kg BW L-thyroxine iv was able to restore normal  $T_4$  levels, but not  $T_3$  levels, without effect on mortality, which remained at 80% both in treated patients and control groups [115]. Similarly, another study in burns, using 200 µg  $T_3$ /daily, did not show significant benefits [116].

Despite studies in animals were in favour of a positive effect in experimental renal failure [1], in humans an increased mortality was showed in a group of acute renal failure treated with L-thyroxine and no beneficial effect of  $T_3$  was observed in transplanted patients [117,118].

Studies in humans showed a slight cardiovascular benefit in patients with shock, respiratory disease, coronary artery bypass graft, premature infants [1, 119].

The discrepancies in the reported study, however, can be attributed to different severity of low- $T_3$ , different schedule of treatment, clinical situations with very different physiopathology, so that it is difficult to obtain a definitive conclusion.

Other interventional studies are reviewed by Bello et al. [19], showing in their complex a beneficial effects on cardiovascular parameters, but not unequivocal benefit of patients' outcome. In fact, in patients with dilated cardiomyopathy, the administration of TH significantly increased left ventricular end-diastolic volume and stroke volume while decreased heart rate [120]. In patients studied after coronary artery bypass surgery, the administration of intravenous  $T_3$  or placebo produced an increase in cardiac output and lowered systemic vascular resistance, without influencing the patients' outcome and therapeutic schedules [121]. In contrast, another study [122] performed after elective coronary artery bypass grafting showed a beneficial effect of intravenous  $T_3$  administration on incidence of postoperative myocardial ischemia and on need for pacemakers or mechanical cardiac support devices. It must be reminded that the administration of TH can directly influence myocardial oxygen supply and demand, causing myocardial ischemic events, even in the absence of coronary artery stenosis or spasms, as reported in some cases [123].

Similar conclusions, biochemical rather than clinical advantage, were drawn in a group of patients after acute burn injuries [124].

## 7. Conclusion

In conclusion, we cannot answer the dilemma, just posed by eminent authors [125,126], about the treatment of low  $T_3$  in NTIS. Some data argue in favour of a real hypothyroidism at tissue level in NTIS; therefore this condition cannot be simply considered an adaptive response. Probably, a full understanding of molecular mechanisms, which cause or are a consequence of low  $T_3$  levels, will allow choosing patients who can really have a benefit from replacement therapy and the appropriate schedule of treatment.

## Author details

Antonio Mancini, Sebastiano Raimondo, Chantal Di Segni, Mariasara Persano and  
Alfredo Pontecorvi

Dept of Internal Medicine, Division of Endocrinology, Catholic University School of Medicine,  
Rome, Italy

## References

- [1] De Groot LJ. Non-thyroidal illness syndrome is a manifestation of hypothalamic-pituitary dysfunction, and in view of current evidence, should be treated with appropriate replacement therapies. *Crit Care Clin.* 2006;22(1) 57-86, vi.
- [2] Rubenfeld S. Euthyroid sick syndrome. *N Engl J Med* 1978;299 1414.
- [3] Wartofsky L, Burman KD. Alterations in thyroid function in patients with systemic illness: the "Euthyroid sick syndrome". *Endocr Rev* 1982;3 164-217.
- [4] Chopra IJ. Euthyroid sick syndrome: abnormalities in circulating thyroid hormones and thyroid hormone physiology in illness (NTI). *Med Grand Rounds* 1982;1 201-212.
- [5] Chopra IJ, Chopra U, Smith SR, Reza M, Solomon DH. Reciprocal changes in serum concentrations of 3,3',5-triiodothyronine (T3) in systemic illnesses. *J Clin Endocrinol Metab* 1975;41 1043-1049.
- [6] Umpierrez GE. Euthyroid Sick Syndrome, *South Med J* 2002;95(5) 506-513.
- [7] Michalaki M, Vagenakis AG, Makri M, Kalfarentzos F, Kyriazopoulou V. Dissociation of the early decline in serum T(3) concentration and serum IL-6 rise and TNF-alpha in illness syndrome induced by abdominal surgery. *J Clin Endocrinol Metab* 2001;86 4198-4205.
- [8] Chopra IJ. Clinical review 86: Euthyroid sick syndrome: is it a misnomer? *J Clin Endocrinol Metab* 1997;82 329-334.
- [9] Bermudez F, Surks MI, Oppenheimer JH. High incidence of decreased serum triiodothyronine concentration in patients with non thyroidal disease. *J Clin Endocrinol Metab* 1975; 41 27-40.
- [10] Kaplan MM, Larsen PR, Crantz FR, Dzau VJ, Rossing TH, Haddow JE. Prevalence of abnormal thyroid function test results in patients with acute medical illnesses. *Am J Med* 1982;72 9-16.
- [11] Marx C, Petros S, Bornstein SR, Weise M, Wendt M, Menschikowski M, Engelmann L, Hoffken G. Adrenocortical hormones in survivors and nonsurvivors of severe sepsis: diverse time course of dehydroepiandrosterone, dehydroepiandrosterone-sulfate, and cortisol. *Crit Care Med* 2003;31 1382-1388.

- [12] Schuetz P, Muller B, Nusbaumer C, Wieland M, Christ-Crain M. Circulating levels of GH predict mortality and complement prognostic scores in critically ill medical patients. *Eur J Endocrinol* 2009;160 157-163.
- [13] Van den Berghe G, De Zegher F, Bouillon R. Acute and prolonged critical illness as different neuroendocrine paradigms. *J Clin Endocrinol Metab* 1998;83 1827-1834.
- [14] Van den Berghe G. Dynamic neuroendocrine responses to critical illness. *Frontiers Neuroendocrinol* 2002;23 370-391.
- [15] Mancini A, Corbo GM, Gaballo A, Valente S, Gigliotti P, Cimino V, De Marinis L, Principi F, Littarru GP. Relationships between plasma CoQ10 levels and thyroid hormones in chronic obstructive pulmonary disease. *Biofactors* 2005;25 (1-4) 201-204.
- [16] Peeters RP, Wouters PJ, Kaptein E, van Toor H, Visser TJ, Van den Berghe G. Reduced activation and increased inactivation of thyroid hormone in tissues of critically ill patients. *J Clin Endocrinol Metab.* 2003; 88 (7) 3202-3211.
- [17] Luca F, Goichot B, Brue T. Non thyroidal illnesses (NTIS). *Ann Endocrinol (Paris)* 2010;71 (Suppl 1) S 13- 24.
- [18] Economidou F, Douka E, Tzanela M, Nanas S, Kotanidou A. Thyroid function during critical illness. *Hormones* 2011;10 (2) 117-124.
- [19] Bello G, Ceaichisciuc I, Silva S, Antonelli M. The role of thyroid dysfunction in the critically ill: a review of literature. *Minerva Anesthesiol* 2010;76 (11) 919-928.
- [20] Magagnin Wajner S, Maia AL. New insights toward the acute non thyroidal illness syndrome. *Front Endocrinol* 2012;3 (8) 1-7 Epub 26 Jan 2012
- [21] Henneman G, Docter R, Krenning EP. Causes and effects of the low T3 syndrome during caloric deprivation and non-thyroidal illness: an overview. *Acta Med Austriaca* 1988;15 42-45.
- [22] Monig H, Arendt T, Meyer M, Kloehn S, Bewig B. Activation of the hypothalamo-pituitary-adrenal axis in response to septic or non-septic diseases- implications for the euthyroid sick syndrome. *Intensive Care Med* 1999;25 1402-1406.
- [23] Cherem HJ, Nellen HH, Barabejski FG, Chong MBA, Lifshits GA. Thyroid function and abdominal surgery. A longitudinal study. *Arch Med Res* 1992;23 143-147.
- [24] Ilias I, Stamoulis K, Armaganidis A, Lyberopoulos P, Tzanela M, Orfanos S, Theodorakopoulou M, Tsagarakis S, Dimopoulou I. Contribution of endocrine parameters in predicting outcome of multiple trauma patients in an intensive care unit. *Hormones* 2007;6 218-226.
- [25] Vardarli I, Schmidt R, Wdowinski JM, Teuber J, Schwedes U, Usadel KH. The hypothalamo-hypophyseal thyroid axis, plasma protein concentration and the hypophyseogonadal axis in low T3 syndrome following acute myocardial infarct. *Klinische Wochenschrift* 1987;65 129-133.

- [26] Hamilton MA, Stevenson LW, Lun M, Walden JA. Altered thyroid hormone metabolism in advanced heart failure. *J Am Coll Cardiol.* 1990;16 (1) 91-95.
- [27] Holland FW, Brown PS, Weintraub BD, Clark RE. Cardiopulmonary bypass and thyroid function: an "euthyroid sick syndrome". *Ann Thorac Surg* 1991;52 46-50.
- [28] Scoscia E, Baglioni S, Eslami A, Iervasi G, Monti S, Todisco T. Clinical study "Low triiodothyronine (T3) state: a predictor of outcome in respiratory failure? Results of a clinical pilot study. *Eur J Endocrinol* 2004;151 557-560.
- [29] Vexiau P, Perez- Castiglioni P, Socie G, Devergie A, Toubert ME, Aractingi S, Gluckmann E. The " Euthyroid sick syndrome": incidence, risk factors and prognostic value soon after allogenic bone marrow transplantation. *Br J Hematol* 1993;85 778-782.
- [30] Kaptein EM. Clinical relevance of thyroid hormone alterations in non-thyroidal illness. *Thyroid International* 1997;4 22-25.
- [31] Wang F, Pan W, Wang H, Wang S, Pan S, Ge J. Relationship between thyroid function and ICU mortality: a prospective observation study. *Crit Care* 2012;16 (1) R11.
- [32] Bello G, Pennisi MA, Montini L, Silva S, Maviglia R, Cavallaro F, Bianchi A, De Marinis L, Antonelli M. Nonthyroidal illness syndrome and prolonged mechanical ventilation in patients admitted to the ICU. *Chest* 2009;135 (6) 1448-1454.
- [33] Klein I, Ojamaa K. Thyroid hormone and the cardiovascular system. *N Engl J Med* 2001;344 501-509.
- [34] Franklyn JA, Gammage MD, Ramsden DB, Sheppard MC. Thyroid status in patients after acute myocardial infarction. *Clin Sci (Lond)* 1984;67 585-590.
- [35] De Marinis L, Mancini A, Masala R, Torlontano M, Sandric S, Barbarino A. Evaluation of pituitary-thyroid axis response to acute myocardial infarction. *J Endocrinol Invest* 1985;8 507-511.
- [36] Opasich C, Pacini F, Ambrosino N. Sick euthyroid syndrome in patients with moderate to severe chronic heart failure. *Eur Heart J* 1996;17 1860-1866.
- [37] Iervasi G, Pingitore A, Landi P, Raciti M, Ripoli A, Scarlattini M, L'Abbate A, Donato L. Low- T3 syndrome: a strong prognostic predictor of death in patients with heart disease. *Circulation* 2003;107 (5) 708-713.
- [38] Mancini A, Festa R, Di Donna V, Leone E, Littarru GP, Silvestrini A, Meucci E, Pontecorvi A. Hormones and antioxidant systems: role of pituitary and pituitary-dependent axes. *J Endocrinol Invest* 2010;33 (6) 422-433.
- [39] Mancini A, Corbo GM, Scapigliati A, Leone E, Conti M, Littarru GP, Meucci E, De Marinis L, Pontecorvi A. Low-T3 syndrome in chronic obstructive pulmonary disease and heart surgery patients: evaluation of plasma antioxidant systems. *Endocrine Abstracts* 2008;16 752.

- [40] Chow CC, Mak TW, Chan CH & Cockram CS. Euthyroid sick syndrome in pulmonary tuberculosis before and after treatment. *Ann Clin Biochem* 1995;32 385-391.
- [41] Kawakami M, Usami I, Kuroki H & Goto M. Thyroid hormones in patients with clinical stable pneumoconiosis. *Nihon Kyobu Shikkan Gakkai Zasshi* 1993;31 1215-1219.
- [42] Wawrzynska L, Sakowicz A & Filipecki S. Euthyroid sick syndrome in patients with respiratory failure. *Pneumol Alergol Pol* 1996;64 (Suppl 2) 193-199.
- [43] Laghi F, Adiguzel N, Tobin MJ. Endocrinological derangements in COPD. *Eur Respir J* 2009;34 975-996.
- [44] Creutzberg EC, Casaburi R. Endocrinological disturbances in chronic obstructive pulmonary disease. *Eur Respir J* 2003;22 (Suppl 46) 76s-80s.
- [45] Dimopolou I, Ilias I, Mastorakos G, Mantzos E, Roussos C, Koutras DA. Effects of severity of chronic obstructive pulmonary disease on thyroid function. *Metabolism Clin Exper* 2001;50 1397-1401.
- [46] Okutan O, Kartaloglu Z, Onde ME, Bozkanat E, Kunter E. Pulmonary function tests and thyroid hormone concentrations in patients with chronic obstructive pulmonary disease. *Med Princ Pract* 2004;13 126-128.
- [47] Karadag F, Ozcan H, Karul AB, Yilmaz M, Cildag O. Correlates of non-thyroidal illness syndrome in chronic obstructive pulmonary disease. *Respir Med* 2007;101 1439-1446.
- [48] Bratel T, Wennlund A, Carlstrom K. Impact of hypoxaemia on neuroendocrine function and catecholamine secretion in chronic obstructive pulmonary disease (COPD): Effects of long-term oxygen treatment. *Respir Med* 2000;94 1221-1228.
- [49] Mancini A, Corbo GM, Gaballo A, Raimondo S, Di Segni C, Gigliotti P, Silvestrini A, Valente S, Littarru GP, Pontecorvi A, Meucci E. Relationship between plasma antioxidants and thyroid hormones in chronic obstructive pulmonary disease. *Exp Clin Endocrinol Diab* 2012;120 623-628.
- [50] Iglesias P, Olea T, Vega-Cabrera C, Heras M, Bajo MA, Del Peso G, Arias MJ, Selgas R, Díez JJ. Thyroid function tests in acute kidney injury. *J Nephrol*. 2012;0. doi: 10.5301/jn.5000106.
- [51] Lo JC, Chertow GM, Go AS, Hsu CY. Increased prevalence of subclinical and clinical hypothyroidism in persons with chronic kidney disease. *Kidney Int* 2005;67 1047-1052.
- [52] Wiederkehr MR, Kalogiros J, Krapf R. Correction of metabolic acidosis improves thyroid and growth hormone axes in haemodialysis patients. *Nephrol Dial Transplant* 2004;19 1190-1197.
- [53] Zoccali C, Tripepi G, Cutrupi S, Pizzini P, Mallamaci F. Low triiodothyronine: A new facet of inflammation in end-stage renal disease. *J Am Soc Nephrol* 2005;16 2789-2795.

- [54] Bando Y, Ushiogi Y, Okafuji K, Toya D, Tanaka N, Miura S. Non-autoimmune primary hypothyroidism in diabetic and non-diabetic chronic renal dysfunction. *Exp Clin Endocrinol Diabetes* 2002;110 408-415.
- [55] Carrero JJ, Qureshi AR, Axelsson J, Yilmaz MI, Rehnmark S, Witt MR, Barany P, Heimbürger O, Suliman ME, Alvestrand A, Lindholm B, Stenvinkel P. Clinical and biochemical implications of low thyroid hormone levels (total and free forms) in euthyroid patients with chronic kidney disease. *J Intern Med* 2007;262 690-701
- [56] Kostopanagiotou G, Kalimeris K, Mourouzis I, Costopanagiotou C, Arkadopoulos N, Panagopoulos D, Papoutsidakis N, Chranioti A, Pafiti A, Spanou D, Smyrniotis V, Pantos C. Thyroid hormones alterations during acute liver failure: possible underlying mechanisms and consequences. *Endocrine* 2009;36 (2) 198-204.
- [57] Marik PE, Gayowski T, Starzl TE. The hepatoadrenal syndrome: a common yet unrecognized clinical condition. *Crit Care Med* 2005;33 1254-1259
- [58] Caregaro L, Alberino F, Amodio P, Merkel C, Angeli P, Plebani M, Gatta A. Nutritional and prognostic significance of serum hypothyroxinemia in hospitalized patients with liver cirrhosis. *J Hepatol* 1998;28 115-121
- [59] Zietz B, Lock G, Plach B, Drobnik W, Grossmann J, Scholmerich J, Straub RH. Dysfunction of the hypothalamic-pituitary-glandular axes and relation to Child-Pugh classification in male patients with alcoholic and virus-related cirrhosis. *Eur J Gastroenterol Hepatol* 2003;15 495-501.
- [60] Kayacetin E, Kisakol G, Kaya A. Low serum total thyroxine and free triiodothyronine in patients with hepatic encephalopathy due to non-alcoholic cirrhosis. *Swiss Med Wkly* 2003;133 210-213.
- [61] Bratusch-Marrain P, Vierhapper H, Grubeck-Loebenstien B, Waldhausl W, Kleinberger G. Pituitary-thyroid dysfunction in severe non-thyroidal disease: "low-T4 syndrome". *Endokrinologie* 1982;80 207-212.
- [62] Kaptein EM, Robinson WJ, Grieb DA. Peripheral serum thyroxine, triiodothyronine and reverse triiodothyronine kinetics in the low state of acute nonthyroidal illnesses. *J Clin Invest* 1982;69 526-535.
- [63] Arem R, Wiener GJ, Kaplan SG, Kim HS, Reichlin S, Kaplan MM. Reduced tissue thyroid hormone levels in fatal illness. *Metabolism* 1993;42 1102-1108.
- [64] Hennemann G, Docter R, Krenning EP. Causes and effects of the low-T3 syndrome during caloric deprivation and non-thyroidal illness: an overview. *Acta Med Austriaca* 1988;15 (suppl 1) 42-45.
- [65] Richmand DA, Molitch ME, O'Donnell TF. Altered thyroid hormone levels in bacterial sepsis: the role of nutritional adequacy. *Metabolism* 1980;29 936-942.

- [66] Mancini A, Di Donna V, Leone E, Giacchi E. Endocrine alterations in anorexia nervosa. In: Mancini A, Daini S, Caruana L (eds) *Anorexia nervosa: a multidisciplinary approach*. New York: Nova Science Pub Inc; 2010. p3-30.
- [67] Dickerman AL, Barnhill JW. Abormal thyroid function tests in psychiatric patients: a red herring? *Am J Psychiatry* 2012;169 (2) 127-133.
- [68] Salvatore D, Davies TF, Schumberger M, Hay ID, Larsen PR. Thyroid physiology and diagnostic evaluation of patients with thyroid disorders. In: Melmed S, Polonsky KS, Larsen PR, Kronenberg HM (Eds) *Williams Textbook of Endocrinology* 12<sup>th</sup> Ed. Philadelphia: Elsevier Saunders; 2011. p 327-361.
- [69] Warner MH, Beckett GJ. Mechanisms behind the non-thyroidal illness syndrome: an update. *J Endocrinol* 2010;205 1-13
- [70] Melmed S, Geola FL, Reed AW, Pekary AE, Park J, Hershman JM. A comparison of methods for assessing thyroid function in illness. *J Clin Endocrinol Metab* 1982;54 300-306.
- [71] Docter R, Krenning EP, de Jong M, Hennemann G. The sick euthyroid syndrome: changes in thyroid hormone serum parameters and hormone metabolism. *Clin Endocrinol* 1993;39 499-518.
- [72] Vierhapper H, Laggner A, Waldhausl W, Grubeck-Loebenstein B, Kleinberger G. Impaired secretion of TSH in critically ill patients with 'low T4-syndrome'. *Acta Endocrinol* 1982;101 542-549.
- [73] Faber J, Kirkegaard C, Rasmussen B, Westh H, Busch-Sorensen M, Jensen IW. Pituitary-thyroid axis in critical illness. *J Clin Endocrinol Metab* 1987;65 315-320.
- [74] Van den Berghe G, Weekers F, Baxter RC, Wouters P, Iranmanesh A, Bouillon R, Veldhuis JD. Five-day pulsatile gonadotropin-releasing hormone administration unveils combined hypothalamic-pituitary-gonadal defects underlying profound hypoandrogenism in man with prolonged critical illness. *J Clin Endocrinol Metab* 2001;86 3217-3226.
- [75] Arem R, Deppe S. Fatal nonthyroidal illness may impair nocturnal thyrotropin levels. *Am J Med* 1990;88 258-262.
- [76] Lee H- Y, Suhl J, Pekary AE, Hershman JM. Secretion of thyrotropin with reduced concanavalin-A-binding activity in patients with severe nonthyroid illness. *J Clin Endocrinol Metab* 1987;65 942.
- [77] Van den Berghe G, De Zegher F, Baxter RC, Veldhuis JD, Wouters P, Schetz M, Verwaest C, Van der Vorst E, Lauwers P, Bouillon R, Bowers CY. Neuroendocrinology of prolonged critical illness: effects of exogenous thyrotropin-releasing hormone and its combination with growth hormone secretagogues. *J Clin Endocrinol Metab* 1998;83 309-319.

- [78] Afandi B, Vera R, Schussler GC, Yap MG. Concordant decreases of thyroxine and thyroxine binding protein concentrations during sepsis. *Metabolism* 2000;49 (6) 753-754.
- [79] Maia AL, Kim BW, Huang SA, Harney JW, Larsen PR. Type 2 iodothyronine deiodinase is the major source of plasma T3 in euthyroid humans. *J Clin Invest* 2005;115 (9) 2524-2533.
- [80] Wajner SM, Goemann IM, Bueno AL, Larsen PR, Maia AL. IL-6 promotes nonthyroidal illness syndrome by blocking thyroxine activation while promoting thyroid hormone inactivation in human cells. *J Clin Invest*. 2011;121 (5) 1834-1845.
- [81] Chopra IJ. Clinical review 86: Euthyroid sick syndrome: is it a misnomer? *J Clin Endocrinol Metab* 1997;82 (2) 329-334.
- [82] Dentice M, Domenico S. Deiodinases: the balance of thyroidal hormone. Local impact of thyroid hormone inactivation. *J Endocrinol* 2011;209 273-282.
- [83] Lado-Abeal J, Romero A, Castro-Piedras I, Rodriguez-Perez A, Alvarez-Escudero J. Thyroid hormone receptors are down-regulated in skeletal muscle of patients with non-thyroidal illness syndrome secondary to non-septic shock. *Eur J Endocrinol* 2010;163 (5) 765-73.
- [84] Nagaya T, Fujieda M, Otsuka G, Yang JP, Okamoto T & Seo H. A potential role of activated NF- $\kappa$ B in the pathogenesis of euthyroid sick syndrome. *J Clin Invest* 2000;106 393-402.
- [85] Friesema EC, Ganguly S, Abdalla A, Manning Fox JE, Halestrap AP & Visser TJ. Identification of monocarboxylate transporter 8 as a specific thyroid hormone transporter. *J Biol Chem* 2003;278 40128-40135.
- [86] Goodson M, Jonas BA, Privalsky MA. Corepressors: custom tailoring and alterations while you wait. *Nucl Recept Signal* 2005;3 e003.
- [87] Rodriguez-Perez A, Palos-Paz F, Kaptein E, Visser TJ, Dominguez-Gerpe L, Alvarez-Escudero J, Lado-Abeal J. Identification of molecular mechanisms related to nonthyroidal illness syndrome in skeletal muscle and adipose tissue from patients with septic shock. *Clin Endocrinol (Oxf)* 2008;68 (5) 821-827.
- [88] Mebis L, Paletta D, Debaveye Y, Ellger B, Langouche L, D'Hoore A, Darras VM, Visser TJ, Van den Berghe G. Expression of thyroid hormone transporters during critical illness. *Eur J Endocrinol* 2009;161 (2) 243-250.
- [89] Monig H, Arendt T, Meyer M, Kloehn S, Bewig B. Activation of the hypothalamic-pituitary-adrenal axis in response to septic or non-septic disease-implications for the euthyroid sick syndrome. *Intensive Care Med* 1999;25 1402-1406.
- [90] Hermus RM, Sweep CGJ, Van Der Meer MJM, Ross HA, Smals AGH, Benraad TJ, Kloppenborg PWC. Continuous infusion of interleukin-1 induces a nonthyroidal illness syndrome in the rat. *Endocrinology* 1992;131 2139-2146.

- [91] Van der Poll T, Romijn JA, Wiersinga WM, Saurwein HP. Tumor necrosis factor: a putative mediator of the sick euthyroid syndrome in man. *J Clin Endocrinol Metab* 1990;71 1567-1572.
- [92] Van der Poll T, Endert E, Coyle SM, Agosti JM, Lowry SF. Neutralization of TNF does not influence endotoxin induced changes in thyroid hormone metabolism in humans. *Am J Physiol* 1999;276 R357-362.
- [93] Nagaya T, Fujieda M, Otsuka G, Yang JP, Okamoto T, Seo H. A potential role of activated NF-kappa B in the pathogenesis of euthyroid sick syndrome. *J Clin Invest* 2000;106 393-402.
- [94] Bartalena L, Brogioni S, Grasso L, Velluzzi F, Martino E. Relationship of the increased serum interleukin-6 concentration to changes of thyroid function in nonthyroidal illness. *J Endocrinol Invest* 1994;17 269-274.
- [95] Boelen A, Platvoet-ter Schiphorst MC, Wiersinga WM. Association between serum interleukin-6 and serum 3,5,3'-triiodothyronine in nonthyroidal illness. *J Clin Endocrinol Metab* 1993;77 1695-1699.
- [96] Stouthard JML, Van Der Poll T, Endert E, Bakker PJM, Veenhof CHN, Sauerwein HP, Romijn JA. Effects of acute and chronic interleukin-6 administration on thyroid hormone metabolism in humans. *J Clin Endocrinol Metab* 1994;79 1342-1346.
- [97] Mancini A, Leone E, Di Donna V, Festa R. Anorexia nervosa and cytokines. In: Mancini A, Daini S, Caruana L (eds) *Anorexia nervosa: a multidisciplinary approach*. New York: Nova Science Pub Inc; 2010. p 31-49.
- [98] Resch U, Helsel G, Tatzber F & Sinzinger H. Antioxidant status in thyroid dysfunction. *Clin Chem Lab Med* 2002;40 1132-1134.
- [99] Asayama K, Kato K. Oxidative muscular injury and its relevance to hyperthyroidism. *Free Radic Biol Med* 1990;8 293-303.
- [100] Dobrzynska MM, Baumgartner A & Andersin D. Antioxidants modulate thyroid hormone- and noradrenaline-induced DNA damage in human sperm. *Mutagenesis* 2004;19 (49) 325-330.
- [101] Venditti P, Di Meo S. Thyroid hormone-induced oxidative stress. *Cell Mol Life Sci* 2006;63 (4) 414-434.
- [102] Mancini A, Giacchi E, Raimondo S, Di Segni C, Silvestrini A, Meucci E. Hypothyroidism, oxidative stress and reproduction. In: Springer D (ed) *Hypothyroidism-Influences and treatments*. Rijeka: InTech; 2012. p 117-134.
- [103] Ademoglou E, Gokkusu C, Yarman S & Azizlerli H. The effect of Methimazol on oxidant and antioxidant system in patients with hyperthyroidism. *Pharmacol Res* 1998;3 93-96.

- [104] Bianchi G, Solaroli E, Zaccheroni V, Grossi G, Bargossi AM & Melchionda N. Oxidative stress and anti-oxidant metabolites in patients with hyperthyroidism: effect of treatment. *Horm metab res* 1990;31 620-624.
- [105] Mancini A, De Marinis L, Calabrò F, Sciuto R, Oradei A, Lippa S, Sandric S, Littarru GP, Barbarino A. Evaluation of metabolic status in amiodarone-induced thyroid disorders: plasma Coenzyme Q10 determination. *J Endocrinol Invest* 1989;12 511-516.
- [106] Mancini A, Corbo GM, Gaballo A, Valente S, Gigliotti P, Cimino V, De Marinis L, Principi F, Littarru GP. Relationships between plasma Coenzyme Q10 levels and thyroid hormones in chronic obstructive pulmonary disease. *Biofactors* 2005;25 (1-4) 201-204.
- [107] Baskol G, Atmaca H, Tanriverdi F, Baskol M, Kocer D & Bayram F. Oxidative stress and enzymatic antioxidant status in patients with hypothyroidism before and after treatment. *Exp Clin Endocrinol Diab* 2007;115 (8) 522-526.
- [108] Torun AN, Kulaksizoglu S, Kulaksizoglu M, Pamuk BO, Isbilen E & Tutuncu NB. Serum total antioxidant status and lipid peroxidation marker malondialdehyde levels in overt and subclinical hypothyroidism. *Clin Endocrinol (Oxf)* 2009;70 (3) 469-474.
- [109] Santi A, Duarte MM, Moresco RN, Menezes C, Bagatini MD, Schetinger MR & Loro VL. Association between thyroid hormones, lipids and oxidative stress biomarkers in overt hypothyroidism. *Clin Chem Lab Med* 2010;48 (11) 1635- 1639.
- [110] Coria MJ, Pastràn AI, Gimenez MS. Serum oxidative stress parameters of women with hypothyroidism. *Acta Biomed* 2009;80 135-139.
- [111] Kebapcilar L, Akinci B, Bayraktar F, Comlekci A, Solak A, Demir T, Yener S, Küme T, Yesil S. Plasma thiobarbituric acid-reactive substance levels in subclinical hypothyroidism. *Med princ pract* 2007;16 432-436.
- [112] Mancini A, Leone E, Silvestrini A, Festa R, Di Donna V, De Marinis L, Pontecorvi A, Littarru GP, Meucci E. Evaluation of antioxidant systems in pituitary-adrenal axis diseases. *Pituitary* 2010;13 (2) 138-145.
- [113] Abilés J, de la Cruz AP, Castaño J, Rodríguez-Elvira M, Aguayo E, Moreno-Torres R, Llopis J, Aranda P, Argüelles S, Ayala A, de la Quintana AM, Planells EM. Oxidative stress is increased in critically ill patients according to antioxidant vitamins intake, independent of severity: a cohort study. *Crit Care* 2006;10 (5) R146.
- [114] Papp LV, Lu J, Striebel F, Kennedy D, Holmgren A, Khanna KK. The redox state of SECIS binding protein 2 controls its localization and selenocysteine incorporation function. *Mol Cell Biol* 2006;26 (13) 4895-910.
- [115] Brent GA, Hershman JM. Thyroxine therapy in patients with severe illnesses and lower serum thyroxine concentration. *J Clin Endocrinol Metab* 1986;63 1-8.

- [116] Becker RA, Vaughan GM, Ziegler MG, Seraile LG, Goldfarb W, Mansour EH, McManus WF, Pruitt BA, Mason AD. Hypermetabolic low triiodothyronine syndrome of burn injury. *Crit Care Med* 1982;10 870-875.
- [117] Acker CG, Singh AR, Flick RP, Bernardini J, Greenberg A, Johnson JP. A trial of thyroxine in acute renal failure. *Kidney Int* 2000;57 293-298.
- [118] Acker CG, Flick R, Shapiro R, Scantlebury VP, Jordan ML, Vivas C, Greenberg A, Johnson JP. Thyroid hormone in the treatment of post-transplant acute tubular necrosis (ATN). *Am J Transplant* 2002;2 57-61.
- [119] Schoenberger W, Grimm W, Emmrich P, Gempp W. Thyroid administration lowers mortality in premature infants. *Lancet* 1979;2 1181.
- [120] Pingitore A, Galli E, Barison A, Iervasi A, Scarlattini M, Nucci D, L'Abbate A, Mariotti R, Iervasi G. Acute effects of triiodothyronine (T3) replacement therapy in patients with chronic heart failure and low-T3 syndrome: a randomized, placebo-controlled study. *J Clin Endocrinol Metab* 2008;93 1351-1358.
- [121] Klemperer JD, Klein I, Gomez M, Helm RE, Ojamaa K, Thomas SJ, Isom OW, Krieger K. Thyroid hormone treatment after coronary-artery bypass surgery. *N Engl J Med* 1995;333 1522-1527.
- [122] Mullis-Jansson SL, Argenziano M, Corwin S, Homma S, Weinberg AD, Williams M, Rose EA, Smith CR. A randomized double-blind study of the effect of triiodothyronine on cardiac function and morbidity after coronary bypass surgery. *J Thorac Cardiovasc Surg* 1999;117 1128-1134.
- [123] Bergeron GA, Goldsmith R, Schiller NB. Myocardial infarction, severe reversible ischemia, and shock following excess thyroid administration in a woman with normal coronary arteries. *Arch Intern Med* 1988;148 1450-1453.
- [124] Becker RA, Vaughan GM, Zeigler MG, et al. Hypermetabolic low triiodothyronine syndrome of burn injury. *Crit Care Med* 1982;10 870-875.
- [125] De Groot LJ. Dangerous dogmas in medicine: the nonthyroidal illness syndrome. *J Clin Endocrinol Metab* 1999;84 151-164.
- [126] Chopra IJ. Nonthyroidal illness syndrome or euthyroid sick syndrome? *Endocr Pract* 1996;2 45-52.