

# We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

186,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index  
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?  
Contact [book.department@intechopen.com](mailto:book.department@intechopen.com)

Numbers displayed above are based on latest data collected.  
For more information visit [www.intechopen.com](http://www.intechopen.com)



---

# Heterogeneity and Plasticity of Multiple Myeloma

---

Hana Šváchová, Sabina Sevcikova and Roman Hájek


Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/55365>

---

## 1. Introduction

Modern molecular and cytogenetic approaches have furthered progress in our understanding of MM biology and have led to the development of targeted therapy that has improved management of this incurable disease. Novel agents such as bortezomib, lenalidomide or thalidomide, have increased median survival rates and improved prospects for MM patients resistant to conventional therapy [1, 2]. Despite these therapeutic advances, MM remains a very difficult disease to treat still accompanied by the threat of repeated relapses with a fatal ending. These observations indicate that at least some of the MM cells are not targeted efficiently by current drug therapies. The existence of such persistent populations, called myeloma stem cells (MSC) or myeloma-initiating cells (MIC) has been suspected for more than two decades. However, the cells of origin remain elusive [3-9]. Timeline of growing knowledge about putative MSC is displayed in **Figure 1**. Discrepancies among myeloma stem cell concepts have arisen in parallel with the high phenotypic heterogeneity of clonal PCs that might be another factor contributing to the failure of therapies and identification of the population responsible for relapse. Myeloma PCs strongly depends on the supportive role of the bone marrow (BM) microenvironment (MEV) – it is a source of essential growth factors, supports survival and dissemination of pathological PCs [10-14]. Furthermore, hypoxic conditions of tumor microenvironments support tumor progression by inducing angiogenesis, maintaining the malignant phenotype and stimulating osteoclastogenesis [15-18]. There is growing evidence that signals from pathological microenvironments can (reversibly) alter the phenotype of PCs. Such plasticity of PCs might result in obvious heterogeneity of MM and generate inconsistencies among myeloma stem cell concepts.



**Figure 1.** Timeline of myeloma stem cell concepts

**2. Myeloma stem cell concepts**

A number of laboratories have tried to identify a biologically distinct population of so called myeloma precursors or myeloma stem cells (MSC) which are responsible for the incurability of MM (see **Table 1**). However, none of these concepts have been unambiguously proven until now. With regards to the fact that abnormal PCs of MM show features of advanced differentiation and mature morphology, the population responsible for the origin and sustainability of tumor mass has been suspected in the minor population of clonotypic or clonogenic CD138<sup>+</sup> cells retaining key stem cell properties, tumor-initiating potential, self-renewal and resistance to chemotherapy [6, 19-25]. However, it was demonstrated that even the dominant population of human CD138<sup>+</sup> PCs contained clonogenic cells; these cells show plasticity potential that might be responsible for dedifferentiation and acquiring of stem cell properties [8, 26-30].

	Characteristics of the cells	Citation
Clonotypic B cells	PB CD19, CD38, CD10, CD11b, CD34 (HPCA-1), variable CD20, PCA-1, CD45RO, variable CD45, and CD56	[19-23, 31-34]
	PB CD34 <sup>+</sup> , CD38 <sup>+</sup> , CD184 <sup>+</sup> , CD31 <sup>+</sup> , CD50 <sup>+</sup> , CD138 <sup>-</sup> , CD19 <sup>-</sup> , CD20 <sup>-</sup>	[35]
	PB CD19 <sup>+</sup> CD27 <sup>+</sup> CD38 <sup>-</sup> memory B-like cells	[24]
Clonotypic pre-switch B cells	pre-switch somatically hypermutated clonotypic cells (VDJ sequence still joined to C $\mu$ gene)	[36-38]
	PB and BM CD19, HLA class II or surface IgM	
Clonogenic CD138 <sup>-</sup> cells	BM CD20 <sup>+</sup> CD27 <sup>+</sup> CD34 <sup>-</sup> CD138 <sup>-</sup> clonogenic myeloma stem cells	[6, 39-41]
	capacity to exclude dyes sideways from the diagonal in FACS analysis plots CD138 <sup>-</sup> and CD138 <sup>+</sup>	[42, 43]
Side population	BM CD38 <sup>+</sup> CD45 <sup>-</sup> PC (SCID-hu model)	[7, 26, 28]
	BM clonogenic CD38 <sup>+</sup> CD138 <sup>-</sup> CD45 <sup>-</sup> PC (SCID-rab model)	
Clonogenic plasma cells	BM CD38 <sup>+</sup> CD138 <sup>-</sup> CD45 <sup>-</sup> CD19 <sup>-</sup> CD34 <sup>-</sup> PC (osteoclast coculture)	[27]
	CD19 <sup>-</sup> CD45 <sup>low/-</sup> CD38 <sup>high</sup> /CD138 <sup>+</sup> PCs	[30]
Intermediate PC precursors	CD138 <sup>+</sup> CD44 <sup>+</sup> give rise to short-lived PC	[44-46]
	CD138 <sup>+</sup> CD44 <sup>-</sup> /CD138 <sup>-</sup> CD44 <sup>+</sup> give rise to long-lived PC in murine model candidates for the normal counterpart of transformed MM cells	

**Table 1.** Candidates for putative neoplastic PC precursors/MS.

## 2.1. Clonotypic B cells

Myeloma precursors that share identical variable diversity joining (VDJ) regions, rearrangements of immunoglobulin heavy chain gene (*IgH*) with patient's tumor PCs and show a pre-plasma cell phenotype are referred to as clonotypic B cells (CBL). They have been identified in peripheral blood, lymph nodes and bone marrow [32, 36, 37, 47-49]. An extensive accumulation of somatic mutations in *IgH* gene and an absence of intraclonal variation suggest that CBLs originate from post-germinal center B cells [37, 50]. Although the concept of clonotypic B lymphocytes as neoplastic PC precursors is widely accepted, other studies suggest the existence of MM precursors in other compartments (see Table 1). Additionally, phenotypic profile and amount of CBLs vary among studies – e.g., it is not clear if these cells resemble CD19<sup>+</sup>CD27<sup>+</sup>CD38<sup>+</sup> memory B-cells or carry a marker of hematopoietic stem cells, CD34 with or without expression of the CD19 surface marker [21, 23, 35]. Furthermore, McSweeney *et al.* (1996) did not find bone marrow CD19<sup>+</sup> cells to be clonally restricted to kappa/lambda or revealed any deviations of expression patterns of B-cell maturation markers from normal B-cell components [51]. His results do not support a hypothesis of disturbed B-cell maturation and development of the disease from early stages of B-cell ontogeny. Moreover, DNA analysis proved that CD38<sup>++</sup>CD19<sup>+</sup> cells were aneuploid in most cases with a typical cell cycle profile indicating the presence of a proliferating population, while cells expressing CD19 were diploid. This suggests the existence of self-replicating plasma cell compartments that have a capacity to replenish the tumor without the involvement of early B lymphocyte progenitors.

A number of studies that confirmed the presence of clonotypic B cells have been based mainly on the detection of the same *IgH* rearrangements by PCR-based methods. The percentage of MM patients with *IgH*-positive clonotypic B-cells ranged from 40% and 87% [36, 48, 52]. Limited dilution PCR assays detected the abundance of CBLs in 0.24% - 25% of peripheral blood mononuclear cells (PBMC) and 66% of all peripheral B cells [32, 36, 52, 53]. This wide range of occurrence may result from methodological errors of the PCR technique or expression of atypical, non-clinical *Ig* transcripts [53]. The major disadvantage of the PCR method is that this approach does not allow the morphological identification of cells of interest or sort cells for subsequent functional analyses. In addition, it is important to take into account that the presence of the same *IgH* rearrangements is reliable marker of clonality; it does not confirm the malignancy of cells *per se*. Therefore, the detection of *IgH* rearrangements in CD19<sup>+</sup> cells could also indicate that a small clone of premalignant or premyelomatous B cells persists despite the transformation of other cells into malignant clones but this population does not actively contribute to myeloma clones [51]. Rasmussen *et al.* (2010) suggests that peripheral blood memory B cells represent pre-malignant, partially transformed remnants that could have some proliferative advantage over normal memory B cells. They questioned the involvement of CBLs in MM maintenance due to the fact these cells expressed specific 'early' oncogenes (*FGFR3*, *MMSET*, *CCND1*) that were deregulated by an *IgH* translocation but lacked 'late' oncogene (*KRAS*) mutations [54].

A large interpatient and interstudy variability led Trepel *et al.*, (2012) to establish patient-individual ligands mimicking the epitope recognized by the myeloma immunoglobulin to specifically target clonotypic surface *IgH*-positive B cells of MM patients [53]. In a cohort of 15

MM patients, semi-nested PCR using HCDR3-specific patient-individual primers detected CBLs in 50% of peripheral blood and/or bone marrow samples. This frequency is line with published data. However, a new flow cytometric protocol detected clonotypic B cells only in one patient with a sensitivity of  $10^{-3}$ , ie. less than one clonotypic B cell per 1000 PBMCs. These cells accounted for about 0.15% of PBMCs and 5% of B cells in this patient. Surprising discrepancies between these two approaches could indicate nonspecific annealing of the CDR3-primer leading to false positive PCR results. Conclusions of this study suggest that the abundance of CBLs is exceedingly low and has been enormously overestimated in previous studies. Similarly to findings of Rasmussen *et al.* (2010), authors consider that unlike those CBLs they are true “feeder” cells for malignant PC compartments and an essential prerequisite of myeloma maintenance and progression [54]. In regard to the rare occurrence of CBLs in MM, it is more likely that these cells are non-malignant remnants, which may be a part of the malignant plasma cell clone but do not participate in tumor maintenance [53].

## 2.2. Clonotypic pre-switch B cells

An identical rearrangement of the IgH VDJ region with a consistent pattern of hypermutations within the clone is a signature of CBLs. MM plasma cell clones generally express monoclonal Igs of these isotypes: IgG, IgA, IgD, IgM, in rare cases kappa or lambda light chains and non-secretory myeloma is detected [55]. Clonotypic isotypes with identical Vh gene sequences and patterns of hypermutations linked to different classes of Ig heavy-chain constant region genes (including C $\mu$  gene) compared to clinical isotypes have been identified in some MM patients [36-38, 56]. B cells with “non-clinical” isotype have been called pre-switch (IgM<sup>+</sup>) clonotypic B cells. Particular classes of Ig are generated by mechanism of class switch recombination (CSR) resulting in the formation of a hybrid switch region composed of switch region S $\mu$  and a respective isotype switch region (C $\delta$ , C $\gamma$ 3, C $\gamma$ 1, C $\alpha$ 1, C $\gamma$ 2, C $\gamma$ 4, C $\epsilon$  and C $\alpha$ 2). Consequently, immunoglobulin production is ‘switched’ from IgM to IgG and IgA (occasionally to IgD or IgE). Sequences between the switch regions are cut out as a deletion loop after double-stranded breaks without changing the VDJ sequence [57-59]. The presence of CSR is a hallmark of “post-switch” B cells. However, pre-switch B cells do not seem to undergo CSR because IgH transcript is composed of the clonotypic VDJ sequence still joined to C $\mu$  gene [56].

Preswitch (IgM<sup>+</sup>) CBLs have been identified in BM and PB of most MM patients; however their frequency is very low. Palumbo *et al.* (1992) have already suggested that MM may originate from pre-switch cells [60]. Billadeau *et al.* (1993), using the allele-specific oligonucleotide PCR, demonstrated the existence of the pre-switch isotype species that were clonally related to the myeloma tumor. Another study also proved the evidence of pre-switch cells with somatically hypermutated clonotypic VDJ region [37]. These cells exhibited expression of CD19 and HLA class II or surface bound IgM. Reiman *et al.* (2001) described the presence of clinical and nonclinical clonotypic isotypes in PB, BM and G-CSF – mobilized blood autografts of MM patients. Expression of preswitch clonotypic transcripts persisted in the blood despite high-dose chemotherapy with stem cell support suggesting drug resistance of this pre-switch population. The persistence of preswitch clonotypic isotypes was associated with reduced survival and with a more advanced disease at the time of diagnosis. Moreover, both pre-



switched and postswitched cells were able to engraft in the NOD/SCID mice indicating their potential clinical relevance. Authors considered the fact that MM is the disease of post-switch cells, IgH isotype switching in MM may accompany worsening disease [38]. Despite these facts, Taylor *et al.* (2008) questioned the role of pre-switch (IgM<sup>+</sup>) clonotypic cells as a progenitor pool for postswitch MM-PCs. They hypothesized that if they are progenitors for MM-PCs, multiple clonotypic switch junctions, or changes in the switch junction, are expected in the postswitch progeny. However, results of specific clonotypic-switch PCR determined the presence of a single, unchanged clonotypic switch junction. Thus, postswitch MM-PCs most likely originate from a single CSR event and pre-switch IgM<sup>+</sup> cells do not represent MM-PC progenitors [52].

Interestingly, pre-switch (non-clonotypic) IgM<sup>+</sup> CD27<sup>+</sup> cells with mutated IgH V region have been detected in various immunodeficiencies and autoimmune diseases but also in healthy donors [61, 62]. Expression of surface marker CD27 suggests that these cells maybe IgM<sup>+</sup> CD27<sup>+</sup> memory B cells. On the other hand, considering the occurrence of IgM<sup>+</sup> CD27<sup>+</sup> cells in humans who cannot form germinal centers (GC), it was presumed that these cells: 1) are generated independently of germinal centers and therefore cannot represent a subset of memory B cells; 2) undergo SHM during generation of the preimmune repertoire; and 3) they mediate responses to T cell-independent antigen. The evidence for these presumptions came from the studies of patients with hyper-IgM syndrome, X-linked lymphoproliferative disease or common variable immunodeficiency that are characterized by some type of defect in GC formation. However, quantification of IgM<sup>+</sup> CD27<sup>+</sup> cells in these diseases showed that their numbers reached approximately 20–40% of the number observed in normal individuals. These results indicate that germinal centers may play a role in the development of IgM<sup>+</sup> CD27<sup>+</sup> cells. IgM<sup>+</sup> memory B cells may be generated at the centroblast stage and undergo SHM but leave GC before the onset of isotype switching. Whether IgM<sup>+</sup> CD27<sup>+</sup> cells with mutated IgH V region might be a source of pre-switch (IgM<sup>+</sup>) CBLs, it remains a matter of further debate and research.

### 2.3. Clonogenic CD138<sup>+</sup> cells

Matsui *et al.*, (2004) suggested that the source of MSC responsible for the initiation and maintenance of MM might be due to a minor population of less differentiated cells reminiscent of memory B-lymphocytes with surface markers CD20<sup>+</sup>CD27<sup>+</sup>CD34<sup>+</sup>CD138<sup>+</sup>. They showed that CD138<sup>+</sup>/CD34<sup>+</sup> cells derived from MM cell lines RPMI 8226, NCI-H929 and primary clinical samples were clonogenic *in vitro*. Cloning efficiency correlated with the disease stage [6]. The depletion of either CD20<sup>+</sup> or CD27<sup>+</sup> cells from the CD138<sup>+</sup>/CD34<sup>+</sup> population significantly limited clonogenic growth of MM, therefore the phenotype of MM cells with *in vitro* clonogenic potential were suggested to be characterized by a pattern of surface markers CD20<sup>+</sup>CD27<sup>+</sup>CD138<sup>+</sup> [39]. Clonogenic potential of CD138<sup>+</sup>/CD34<sup>+</sup> cells was evaluated by successful engraftment of non-obese diabetic/severe combined immunodeficiency (NOD/SCID) mice during both primary and secondary transplantation. CD138<sup>+</sup>/CD34<sup>+</sup> were unable to form colonies *in vitro* and human engraftment was not detected in any of the mice injected with CD138<sup>+</sup> cells. A chimeric anti-CD20 monoclonal antibody, rituximab, was showed to

inhibit clonogenic growth of CD138<sup>+</sup> cells *in vitro* [6,39]. However, clinical trials failed to confirm an effect of rituximab as a useful maintenance therapy for MM [63, 64].

Paino *et al.*, (2011) reevaluated the presence and function of CD20<sup>+</sup> putative MSC in a panel of myeloma cell lines [65]. Although, Matsui *et al.* (2004) described a small population (2-5%) of CD138<sup>+</sup>CD20<sup>+</sup> cells in NCI-H929 and RPMI-8226 cell lines, Paino *et al.* (2011) were not able to detect CD20 by flow cytometry in the majority of tested MM cell lines [6, 65]. Only RPMI-8226 cell lines contained a small population of CD20<sup>dim+</sup> cells (0.3%). These data are consistent with the report of Rossie *et al.* (2010) that showed that U266, NCI-H929 and RPMI-8226 MM cell lines are CD20<sup>+</sup> [66]. Despite previous results, memory B-cell phenotype of putative CD20<sup>+</sup> MSC was also not confirmed. On the contrary, CD20<sup>dim+</sup> cells displayed a myelomatous plasma cell phenotype: CD38<sup>+</sup>CD138<sup>+</sup>CD19<sup>+</sup>CD27<sup>+</sup>CD45<sup>+</sup>. Additionally, CD20<sup>dim+</sup> cells did not exhibit stem cell properties and compared to the CD20<sup>+</sup> population they showed a lower level of self-renewal potential. Both populations developed plasmacytomas when they were injected into CD17-SCID mice suggesting that CD20<sup>dim+</sup> cells are not essential for tumor formation. Furthermore, sorted and plated CD20<sup>dim+</sup> cells did not differentiate into CD20<sup>+</sup> cells. However, CD20<sup>+</sup> cells give rise to CD20<sup>dim+</sup> cells indicating a hierarchical order of differentiation from CD20<sup>+</sup> to CD20<sup>dim+</sup> cells. Overall, these results do not support CD20 as a marker associated with MSC phenotype [65].

Other drugs, such as dexamethasone, lenalidomide, bortezomib, or 4-hydroxycyclophosphamide did not significantly affect CD138<sup>+</sup> cells indicating resistance to some conventional or novel therapy, a characteristic feature of CSC [39]. All four agents significantly inhibited the clonogenic growth of CD138<sup>+</sup> cells isolated from MM cells. Nevertheless, it could not be evaluated in CD138<sup>+</sup> PCs from MM patients because they lack *in vitro* clonogenic activity. Detection of a small CD138<sup>+</sup> population of MM cell lines (< 2%) that displayed stem cell properties mediating drug resistance, such as the capacity to efflux the DNA binding dye Hoechst 33342 and higher relative levels of aldehyde dehydrogenase (ALDH) activity further supports the existence of a resistant MSC compartment in MM. Moreover, CD138<sup>+</sup> cells of MM cell lines also exhibited cellular quiescence similar to adult stem cells – almost all CD138<sup>+</sup> cells were shown to remain in G<sub>0</sub> – G<sub>1</sub> phase and less than 1,5 % in S phase, compared to two thirds of CD138<sup>+</sup> cells in G<sub>0</sub> – G<sub>1</sub> and about 20% in S phase [39]. On the other hand, results of the first study proved that CD138<sup>+</sup> cells isolated from the same MM cell lines expressed higher levels of the proliferation marker Ki67 than CD138<sup>+</sup> cells [6]. These discrepancies might come from different passages of cell lines or culture conditions but cannot conclusively prove a quiescent state of CD138<sup>+</sup> cells.

Besides the clonogenic population of CD138<sup>+</sup> cells, Matsui *et al.* (2004) also isolated circulating clonotypic CD19<sup>+</sup>CD27<sup>+</sup> B cells in peripheral blood of MM patients which were successfully engrafted into NOD/SCID mice. However, it remains unclear whether CD138<sup>+</sup> clonogenic cells are identical to clonotypic CD19<sup>+</sup> B cells. Hosen *et al.* (2012) examined 16 MM samples and found that only CD138<sup>+</sup>CD19<sup>+</sup>CD38<sup>++</sup> cells formed colonies *in vitro* whereas CD19<sup>+</sup> B cells did not [67]. Moreover, CD138<sup>+</sup>CD19<sup>+</sup>CD38<sup>++</sup> cells engrafted into SCID-rab mouse models in 3 cases out of 9, whilst there was no detection of CD19<sup>+</sup> B cell engraftment. Neither CD19<sup>+</sup> B cells transplanted into NOD/SCID IL2R $\gamma$ c(-/-) mice propagated into MM. Surprisingly, CD138<sup>+</sup> PCs



also give rise of MM, but more slowly than CD138<sup>+</sup> cells. Thus, CD138-negative clonogenic cells might represent a population which has the potential to give rise to MM but does not overlap with the population of CD19<sup>+</sup> B cells. These results indicate that CD138-negative clonogenic cells are more PCs than B cells but this does not necessarily mean that Matsui's myeloma stem cell concept is wrong. Although, MSCs phenotypically resemble memory B cells, they may be modified PCs and CD138-positive PCs might represent some "transit" population that subsequently lost its mature phenotype (this will be discussed later). However, Christensen *et al.* (2012) reexamined CD138<sup>+</sup> population of so called MSC and obtained surprising results which were strongly controversial to both previous mentioned reports [68]. An analysis of primary CD138<sup>+</sup> PCs of MM patients showed that the number of CD138<sup>+</sup> cells increased in parallel with increasing time from sampling to analysis. In regards to the fact that myeloma PCs loses expression of the surface antigen CD138 in apoptosis, Annexin V was included in all analyses to monitor apoptotic cells [69]. Expectedly, if a CD138<sup>+</sup> population was detected, these cells were positive for Annexin V. Similar results were also obtained by Chiron *et al.* (2011). Furthermore, qPCR techniques confirmed similarly high levels of CD138 mRNA in both CD138<sup>+</sup> and CD138<sup>+</sup> MM cells. CD138<sup>+</sup> and CD138<sup>+</sup> subpopulations varied neither in expression of CD19 nor in expression of CD20. These contradictory results would imply that the CD138-negative population may represent only cells undergoing apoptosis as a consequence of previous sample handling [68]. Nevertheless, these new findings cannot completely deny results of previous studies that showed the potential of CD138<sup>+</sup> cells to form colonies *in vitro* and engraft mouse models.

## 2.4. Side population

Side population (SP) cells were defined based on their capacity to exclude dyes such as Hoechst 33342 [70]. Hoechst 33342 dye binds to the AT-rich regions found in the minor groove of DNA. Upon UV excitation, cells with this efflux capacity can be identified as the minor population of the positively stained cells sideways from the diagonal in FACS analysis plots. SP cells have been detected in various cancer cell lines as well as primary tumors (rev. in 71). Side population possesses CSC characteristics such as the capacity for regrowth of the tumor, expression of stem cell-like genes and resistance to chemotherapy. This is why SP cells are believed to be the true population responsible for tumor maintenance.

The presence of SP cells were investigated in four MM cell lines, RPMI 8226, U266, OPM2 and KMS-11, and primary MM samples [42]. SP was defined using control cells stained with both Hoechst and verapamil, L-type calcium channel blocker, to establish the SP gate[72]. Reduction of SP cells was demonstrated in all tested cell lines. In 18 of 21 bone marrow samples from MM patients the percentage of SP cells ranged from 0 to 4.9% compared to 0.05% of SP in normal bone marrow. There was neither a significant difference between treated and non-treated MM patients based on the percentage of SP nor a correlation between the percentage of SP and the paraprotein concentration or disease stage. 0.18–0.83% of SP cells express the clonal surface immunoglobulin light chain restriction that matched 89–97% of each patient's PCs. SP cells were also analyzed for the expression of CD138. A mean of 96.1% of SP cells were found to be CD138<sup>+</sup>. However the CD138<sup>+</sup> fraction also contained SP indicating that they are present in

both CD138<sup>-</sup> and CD138<sup>+</sup> compartments. Jakubikova *et al.* (2011) also demonstrated that the SP fraction of MM cell lines expressed CD138 antigen [43]. Moreover, these results did not prove a correlation between expression of CD19, CD20, or CD27 and the proportion of SP cells. Conversely, SP cells showed more clonogenic potential and proliferation index than the main population. Furthermore, adherence to stromal cells increased percentage, viability and proliferation potential of SP cells. Supportive role of BM microenvironment was attenuated by lenalidomide and thalidomide. Lenalidomide itself directly decreased the percentage and clonogenicity of SP cells. This study demonstrated innovative and promising strategies for targeting putative myeloma-initiating cells and prevention of relapse.

Although SPs exhibit stem cell properties and might represent a “feeder” population responsible for the relapse of the disease, others question the method for detection of SP. Hoechst staining binds to DNA resulting in toxicity to live cells, that is why SP cells might represent only a population that survived the lethal effect of Hoechst. SP phenotype might also be affected by staining time, dye concentration or cellular concentration. The problem also lies in cytometric approaches that showed inconsistencies among gating strategies and might lead to contamination of the SP fraction by non-SP cells. Furthermore, usage of verapamil as an inhibitor of efflux was also criticized because verapamil-sensitive cells were detected in the negative or SP gate [73]. These problematic findings require more stringent gating strategies to clear doubts about SP.

## 2.5. Clonogenic plasma cells

The clonogenic potential of primary MM cells was first demonstrated by Hamburger & Salmon (1977). They showed that freshly explanted human myeloma cells are able to form colonies of monoclonal PCs. Colonies consisted of immature plasmablasts and mature PCs. Drewinko *et al.*, (1981) investigated the growth fraction of MM cells. In untreated and nonresponsive patients the growth fraction represented 4% of MM cells. Patients in relapse had the growth fraction ranging from 14% to 83%. Nonproliferating fraction contained true quiescent cells, some proliferating cells with very long intermitotic times, and some proliferating cell that have entered the maturation phase. Although these two reports demonstrated clonogenic and proliferative capacity of primary MM tumors, these cells were not phenotypically defined as whole mononuclear fraction of BM was used for analyses. Therefore, results of these studies cannot answer what kind of cells represents true growth fraction.

First experiments demonstrating the clonogenic growth of phenotypically defined PCs were carried out by Yaccoby and Epstein (1999). They proved that CD38<sup>++</sup>CD45<sup>-</sup> PCs derived from PB and BM are able to engraft severe combined immunodeficiency (SCID)-hu host system with implanted human bone. Circulating clonal PCs grew more rapidly in SCID-hu hosts than those in the BM suggesting that this may represent a subpopulation with a higher growth potential. In contrast to previous reports, PC-depleted blood cells did not give rise to MM in SCID-hu hosts [7]. Yata & Yaccoby, 2004 presented an alternative model for study of myeloma-initiating cells (MIC) that uses rabbit bones implanted subcutaneously in unconditioned SCID mice. The SCID-rab model was also successfully engrafted with CD138<sup>+</sup> PCs of MM patients. Although, these two models did not show a serial engraftment of the disease, they strongly indicate a

critical role of specific bone marrow microenvironments for plasma cell survival and proliferation [28].

A dependence of myeloma PCs on human BM microenvironment was clearly demonstrated in an experiment with NOD/SCID/common cytokine receptor  $\gamma$  chain-deficient (NSG) and recombina-activating gene 2/common cytokine receptor  $\gamma$  chain-deficient (RAG2-/ $\gamma$  c-) mice [30]. Results proved that CD138<sup>+</sup>/CD38<sup>high</sup> cells from MM patients led to a repopulation of CD19<sup>+</sup>CD38<sup>low</sup> or CD138<sup>+</sup>CD38<sup>+</sup> B-lineage cells in human bone-bearing mice but no engraftments were detected in human bone-free mice even after orthotropic intrafemoral injection. Moreover, serially xenotransplanted CD19<sup>+</sup>CD138<sup>+</sup> cells preferably engrafted from a human bone graft but were not detected in any mouse hematopoietic tissues. All grafts derived from CD138<sup>+</sup>/CD38<sup>high</sup> cells were clonally related to myeloma PCs, whereas engraftments of CD19<sup>+</sup>CD38<sup>low</sup> B cells were polyclonal CD19<sup>+</sup>CD38<sup>low</sup> cells. Further fractionation of CD138<sup>+</sup>/CD38<sup>high</sup> cells and their subsequent xenotransplantation showed that CD45<sup>low</sup> or CD19<sup>+</sup>CD38<sup>high</sup>/CD138<sup>+</sup> cells had a higher engraftment ability than CD45<sup>high</sup> or CD19<sup>+</sup> plasmablasts. In line with the above mentioned results, it was concluded that CD19<sup>+</sup>CD45<sup>low</sup>CD38<sup>high</sup>/CD138<sup>+</sup> PCs rather than CD19<sup>+</sup> B cells or plasmablasts enrich for MICs. In addition, analysis of clonogenic potentials of CD138<sup>+</sup> and CD138<sup>-</sup> populations derived from plasma cell leukemia patients strongly favored these findings [29]. This study demonstrated that, though, CD138<sup>+</sup> PCs derived from plasma cell leukemia patients formed colony forming units (CFU) in a very low frequency, no CFU were found when CD138<sup>-</sup> cells were seeded. Harvested cells of CFUs derived from CD138<sup>+</sup> PCs were strongly positive for CD138 but negative for CD20. There was no evidence that they can be CD138<sup>-</sup>CD19<sup>+</sup>CD20<sup>-</sup> plasmablasts or CD19<sup>+</sup>CD20<sup>+</sup>CD138<sup>-</sup> B-cells. These findings further underline an important role of CD20<sup>-</sup>CD138<sup>+</sup> population as a conceivable reservoir of clonal PCs at least in plasma cell leukemia.

Against the statement that CD138-positive PCs represent a “feeder” population responsible for incurability of MM is the fact that abnormal PCs show a low proliferation potential with a plasma cell labeling index (PCLI) ranging from 0.5% in MGUS to 1% in early MM, PCLI. Therefore, the growth fraction of progenitor cells giving rise to MM has been expected in less differentiated stages of B cell development. Despite low proliferative activity of malignant PCs, PCLI is one of the most important prognostic factor with a strong impact on overall survival [98, 99]. Additionally, limited proliferation potential of PCs could be explained by error reduction during DNA synthesis that was proposed for normal adult stem cells that are highly clonogenic but proliferatively silenced [100].

## 2.6. Intermediate plasma cell precursors

O'Connor *et al.* (2002) identified post-GC-precursors in a mouse model that might contribute to long-lived humoral immunity. These cells were distinct from splenic B cells, mature or memory B cells as well as mature PCs. Mediate levels of CD138 indicated that PC precursors might represent a transition state of PC development. PC precursors were demonstrated to migrate to the BM where they proliferate and persist for a long period of time and consequently differentiate to mature PCs without antigen stimulation. CD138<sup>+</sup>CD44<sup>+</sup> give rise to short-lived PCs, CD138<sup>+</sup>CD44<sup>-</sup> or CD138<sup>-</sup>CD44<sup>+</sup> differentiated to long-lived PCs. Two roles of PC precu-

sors were suggested; (i) they might either serve as a reservoir of PCs upon PC attrition or (ii) contribute to post-GC affinity maturation of the humoral immune response. Due to similarities of phenotype, proliferative potential, and differential capacity of the PC precursors to putative MM progenitors, authors suggested that these cells are candidates for the normal counterpart of transformed MM progenitors [74].

The true origin of MM remains a matter for further debate. An unanswered question persists whether clonotypic CD138<sup>+</sup> cells or side populations represents myeloma precursors or phenotypic variants of the MM tumor cells that have lost some plasma cell markers and gained some B-cell markers as Yaccoby (2005) suggested [27, 54]. He presented a new model for myelomagenesis; that mature CD45<sup>low/intermediate</sup> CD38<sup>high</sup> CD138<sup>high</sup>, CD19<sup>+</sup>CD34<sup>+</sup> PC situated closely with osteoclasts in the lytic bone lesions reverse senescence, acquire a stem cell phenotype, (CD45<sup>intermediate/high</sup> CD19<sup>low</sup> and CD34<sup>low</sup>), and become quiescent and apoptosis-resistant. This indicates that myeloma PCs have plasticity that allows them to reprogram, dedifferentiate and acquire autonomous survival properties which are responsible for drug-resistance and relapse of MM patients. Similar results were also obtained by Kukreja *et al.* (2006) when cultured U266 myeloma cell line and primary myeloma PCs with dendritic cells (DC). The coculture of U266 cells and DCs led to an increase in the proportion of cells lacking CD138; a marker of terminally differentiated PCs, and induction of B cell lymphoma 6 (BCL6) expression. It was suggested that BCL6 plays an important role in survival and self-renewal of germinal center B cells and that suppression of BCL6 is a critical feature of normal PC differentiation. Furthermore, a presence of DCs in the culture enhanced clonogenic growth of the myeloma cell line and as well as primary myeloma PCs [75]. Just as in the previous report, this data suggests that the differentiation state of myeloma cells is plastic and can be modified in the presence of DCs. Both reports emphasize an importance of tumor microenvironments for myeloma plasticity. Whether myeloma PCs exhibit plasticity that might be responsible for phenotypic heterogeneity of MMs, and also form inconsistencies among myeloma stem cell concepts, it will be discussed further.

### 3. Phenotypic heterogeneity of multiple myeloma

The high phenotypic heterogeneity of clonal PCs is a hallmark of MM; abnormal PCs frequently express a wide spectrum of multi-lineage antigens such as myeloid, T-cell and natural killer-associated antigens. Compared to normal polyclonal PCs which show high CD38 and CD138 expression along with the B-lymphocyte marker CD19 abnormal PCs generally lack CD19 expression, show variable expression of CD45, dim expression of CD38 and heterogeneous signals of CD138. Further, they have weaker expression of CD27, increased expression of CD28, CD33 and CD56 and variable expression of CD20 and CD117 (rev. in [76, 77]). Causes of phenotypic heterogeneity remain unclear.

PCs belong to the B-cell lineage that expresses paired box protein 5 (PAX5) and its target, CD19. PAX5 activation and inactivation is essential for early B cell commitment as well as for maintenance of the functional identity of B cells throughout B cell development. PAX5



represses B lineage 'inappropriate' genes and simultaneously activates B lineage-specific genes [78, 79]. Conditional PAX5 reduction in late B lymphocytes promotes development to the mature B cell stage. PAX5 and CD19 expression are considered to be downregulated or lost but it has been reported that expression of PAX5 and CD19 is restored in normal PCs of pleural effusion, ascitic fluid and BM aspirate. In contrast to normal PCs, clonal PCs of MM do not express PAX5 or CD19 (> 95% of cases) [80, 81].

Phenotypic heterogeneity of abnormal PCs might reflect a certain degree of dedifferentiation of these cells. It has been previously shown that elimination of PAX5 shifts B cells to multipotency [82]. Therefore expression of multi lineage markers on clonal PCs might be due to a lack of PAX5 gene expression but not only as a consequence of malignant transformation. Cell heterogeneity is the phenomenon that is commonly seen in many cancer types. The cause of this variability might be due to the loss of the lineage master gene which can lead to dedifferentiation or transdifferentiation into cells of other lineages [83].

The ability to differentiate into multiple lineages is a characteristic feature of pluripotent embryonic stem cells. However, pluripotency does not necessarily have to be limited to a population of undifferentiated stem cells of the early embryo and lost irreversibly upon terminal differentiation [84]. Fully differentiated somatic cells can be reprogrammed into inducible pluripotent stem cells (iPSC) by 'forced' expression of pluripotency/reprogramming factors: OCT4, SOX2, c-Myc and KLF4 or NANOG, LIN28, c-Myc and KLF4 [85, 86]. Human somatic cells from all three germ layers, including human differentiated mature B lymphocytes, have been successfully reprogrammed [87-93]. Reprogramming of mature B cells required additional 'sensitization' by the myeloid transcription factor CCAAT/enhancer-binding protein- $\alpha$  (C/EBP $\alpha$ ) that causes a disruption of PAX5 functions, or a specific knock-down of the B cell transcription factor PAX5. This indicates that the loss of PAX5 expression might be associated with gain of stem cell features for mature PCs and represent one of the key events in the pathogenesis of MM.

#### 4. Plasticity of multiple myeloma

There is growing evidence that MM encompasses a certain degree of plasticity that might be responsible for expression of a wide spectrum of multi lineage markers and discrepancies among myeloma stem cell theories. Two facts support this hypothesis:

- i. There is a strong link between induction of pluripotency and tumor progression; several human cancers acquire stem cell-like plasticity upon (re)expression of reprogramming factors. It raises the possibility that dedifferentiation is a key mechanism for the generation of tumor-initiating cells in human cancer [84]. The process of dedifferentiation is suggested to be under control of tumor microenvironment. Interaction of tumor cells with their microenvironment might induce altered differentiation; epithelial-mesenchymal transition (EMT) which has been observed in some solid tumors [94, 95]. Similarly to MM, phenotypic plasticity of myeloma PCs were observed in long-term co-culture with osteoclasts; myeloma PCs lost their




mature phenotype and dedifferentiated to an immature, resilient, apoptosis-resistant phenotype. In addition, CD138<sup>+</sup> PCs cocultured with DCs lost expression of CD138 and increased the clonogenic potential [75]. It raises the possibility that signals from the microenvironment can (reversibly) alter the phenotype of PCs and cause obvious heterogeneity of MM. This regulation might keep dynamic equilibrium between CSC and non-CSC compartments [96].

Dynamic state between CSC and non-CSC like compartments were also observed in a subpopulation of differentiated basal-like human mammary epithelial cells that spontaneously converted to stem-like cells *in vitro* and *in vivo*. Moreover, oncogenic transformation enhances this spontaneous conversion. These findings indicate that normal and CSC-like cells can arise *de novo* and indicate the importance of the differentiation state of cells-of-origin as a critical factor determining the phenotype of their transformed derivatives [97]. Whether mature PCs exhibit a plasticity potential responsible for dedifferentiation into cells with stem cell like properties is questionable but highly attractive. Confirmation that mature PCs have the capacity to convert into CSCs could resolve current inconsistencies among myeloma stem cell theories and help to target the population responsible for relapse of the disease.

- ii. Despite of mature phenotype, myeloma PCs have been reported to express pluripotency factors (SOX2, c-Myc or KLF4) and stem/progenitor markers, such as germline stem cell markers of the MAGE family, the hematopoietic progenitor marker CD117 or neural stem cell marker nestin [83, 101-105]. Some of these factors provide valuable prognostic information. Currently, the role of c-myc is growing as the key transforming factor in the progression of asymptomatic MGUS to a symptomatic disease [101]. SOX2 and MAGE have been demonstrated to be relevant targets for immunotherapy due to their immunogenicity. Interestingly, MGUS patients frequently mount a humoral and cellular immune response against SOX2 but MM patients lack anti-SOX2 immunity [104, 106]. CD117 (c-kit) is an essential hematopoietic growth factor receptor with tyrosine-kinase activity. Aberrant expression of CD117 is detected on a subset of MGUS and MM and is associated with a favorable outcome for MM patients [107, 108]. KLF4 was described in regulation of apoptosis, proliferation and differentiation of B cells and B-cell malignancies. Strong up-regulation of KLF4 was detected in MM cell lines in the process of apoptosis suggesting a role of KLF4 in MM progression [105]. Nestin is a remarkable protein that is found in rapidly dividing cells of developing and regenerating tissues [109]. Therefore, it is surprising that this gene and protein expression of the stem cell marker nestin has been detected in terminally differentiated PCs of MM patients [83, 110]. Moreover, our recent work proved that nestin protein is a tumor specific marker for CD138<sup>+</sup>38<sup>+</sup>PC of MM [110]. Expression of nestin, a marker of stem/progenitor cells in malignant PCs that are considered to be terminally differentiated, is highly controversial and indicates that nestin might play an exceptional role in the pathology of MM. However, biological or clinical implications of nestin have not been determined in monoclonal gammopathies or other hematological malignancies so far.

Summarizing these findings supports the hypothesis that stem cell-like features are not rare in monoclonal gammopathies and might indicate the existence of inducible stem cell properties in more differentiated cells than was initially thought [27, 111] (see Fig. 2).




**Figure 2.** Model of inducible stem cell properties in myeloma

## 5. Putative role of nestin in myeloma plasticity

Neural stem cell marker nestin was first identified in neuroepithelial stem/progenitor cells of the rat central nervous system (CNS) by Hockfield and McKay in 1985 [112]. Nestin is detected in a wide range of undifferentiated tissues under normal and pathological conditions [113-116]. Expression of nestin is a common feature of multipotent proliferative progenitor cells with self-renewal and regeneration potentials. During terminal differentiation, nestin expression is silenced but can be reactivated upon injury or other pathological conditions such as neoplastic transformation. The human nestin gene is located at 1q23 locus and is composed of four exons and three introns. Nestin expression is driven by a minimal promoter that is activated by transcriptional factor Sp1 [117]. Moreover, epigenetic regulation was also demonstrated. Results indicated that histone acetylation might be sufficient to mediate activation of nestin transcription [118].

Nestin protein belongs to a large family of intermediate filament (IF) proteins that are encoded by more than 70 genes expressed in a time and site-specific manner in metazoan cells. Members of the IF family are divided into six classes of proteins according to their structure, properties and localization [119]. Nestin, a class VI protein, is characterized by a  $\alpha$ -helical central 'rod' domain which is typical for all IF, that contains repeated hydrophobic heptad motifs, a short N-terminus (head) and a very long C-terminus (tail) (Fig. 3). C-terminus is suggested to function as a linker or cross-bridge between intermediate filaments, microfilaments and

microtubules [120]. The molecular weight of human nestin protein is ~ 200 – 220 kDa; more often it is found in its glycosylated form with molecular weight of 240kDa [121, 122]. Nestin does not fold by itself most likely because of its very short N-terminus. Therefore, nestin requires the presence of other IF proteins such as type III vimentin, desmin or type IV  $\alpha$ -internexin [109, 123, 124].



**Figure 3.** Structure of intermediate filaments (adopted and modified from Michalzyk & Ziman, [2005])

### 5.1. Cellular roles of nestin

An important regulator of nestin organization and dynamics during mitosis is cdc2-mediated phosphorylation. Phosphorylation/dephosphorylation of nestin may modulate disassembly and assembly of intermediate filaments [125]. These processes might play a role during increased cytoplasmic trafficking in progenitor cells undergoing division or in migrating interphase cells [126-128]. Nestin was shown to participate in asymmetric redistribution of cytoskeletal proteins and other factors to daughter neuroepithelial cells [129]. Nestin structure serves also as a scaffold for cdk5/p35 activity resulting in a cytoprotective effect against stress induced cell death in neural progenitor cells [122, 130]. Moreover, nestin is supposed to be a major determinant in suppression of anti-proliferative activity of glucocorticoid receptors (GR) in undifferentiated tissues by anchoring GRs in cytoplasm but cells lacking nestin accumulate GRs in the nucleus [131]. This mechanism might be responsible for glucocorticoid resistance seen in many cancers.


### 5.2. Nestin in multiple myeloma

Biological or clinical implications of nestin have not been systematically studied in monoclonal gammopathies or other hematological malignancies so far. The first mention of nestin expression in MM was described in malignant PCs of 5 MM patients and two myeloma cell lines (NOP2 and Liu01) by Liu et al. (2007). Authors referred to the existence of CD56<sup>+</sup> primary MM cells expressing neuronal markers, such as nestin, neuron-specific enolase and  $\beta$ -tubulin III [83]. Expression of nestin was reported to be stimulated by the Notch signaling pathway in human gliomas [132]. The Notch pathway plays also an important role in survival and proliferation of malignant PCs and negatively affects bone disease in MM [133]. Our recent

work proved that nestin protein is a tumor-specific marker for CD138<sup>+</sup>38<sup>+</sup>PC of MM patients [110]. Regarding the fact that myeloma PCs appear to be mature, terminally differentiated cells with a low proliferative potential, the presence of nestin is very surprising. Cells expressing nestin may represent a transient population undergoing dedifferentiation into CSC-like phenotype. Providing that PCs possess a truly dedifferentiation potential, the direction of dedifferentiate needs to be elucidated and whether or not this process is reversible.

Unexpectedly, our results showed that nestin was present in one-third of cases in 50% to almost 100% of PCs. Also expression of MAGE genes was found in majority of PCs [102, 134]. Thus, if nestin is expressed in most cells, it means that majority of PCs might gain stem cell features and could repopulate the tumor after therapy. These findings take into question the hierarchical model of myelomagenesis based on the presumption that MM originates from a minor population of MSC. Conversely, stochastic clonal evolution model better describes mechanisms responsible for recurrence of MM. This model suggests that majority of tumor cells have a character of stem cells and may repopulate tumor cells after treatment [135]. Plasticity of PCs better fits the clonal evolution model because weakly proliferating PCs would not be able to quickly repopulate the tumor, therefore we hypothesized that they need to convert into more rapidly dividing cells.

Association of nestin with plasticity of terminally differentiated cell types were demonstrated in the study of metaplastic conversion of mature pancreatic epithelial cells [136] Metaplastic conversion is defined as the replacement of one differentiated cell type by another mature cell type and is frequently associated with an increased risk of subsequent neoplasia. Causes of these changes remains unclear but one possible mechanism may be transdifferentiation of one mature cell type to another one either directly or via an undifferentiated transient cells (rev. in [137]). Means *et al.* (2005) showed that mature acinar cells can convert into ductal epithelia under EGFR signaling. Metaplastic changes were accompanied by occurrence of nestin-positive intermediates similar to nestin-positive precursors observed during early pancreatic development. Results of this study proved a real trans-differentiation potential of mature mammalian cells and indicated that plasticity of mature cell types may play a role in the generation of neoplastic precursors. Whether nestin-positive myeloma PCs might represent particular intermediates undergoing changes leading to the occurrence of cells with a changed (transdifferentiation) or less differentiated phenotype (dedifferentiation), it remains a matter for future research. On the other hand, nestin expression might represent only a byproduct of tumor transformation without any association with myeloma heterogeneity. However, the dynamic character of nestin network plays an important role in the key cell processes such as proliferation, migration and cell survival. Nestin polymerization/ depolymerization influences intracellular signaling, it is likely responsible for rapid redistribution of intracellular proteins, cytoskeletal remodeling and/or might function as a scaffold for protein interactions [122, 125, 129, 130] (**Fig. 4**). All these properties might represent an important prerequisite for myeloma plasticity.



**Figure 4.** Nestin remodeling and cellular functions

**6. Cancer stem cell phenotype and tumor microenvironment**

Recent knowledge supports the hypothesis that altered BM microenvironments participate in both mechanisms leading to tumor progression; induction of stem cell features and stimulation of angiogenesis. Cancer stem cell phenotypes may be a plastic state induced in cancer cells depending upon microenvironmental signals, such as hypoxia. Hypoxia has a great impact on the production of angiogenic factors but it is also a crucial regulator of the stem cell phenotype. Several reports have shown that hypoxia and HIFs are involved in maintaining a stem-like state in normal tissues [138]. One example can represent hematopoietic stem cells that reside in regions regulated by oxygen tension. It is hypothesized that undifferentiated phenotypes of these cells relies on HIF activity in hypoxic areas. Hypoxic areas in tumors might be an analog to stem cell niches in normal tissues. Furthermore, Yoshida *et al.* (2009) have shown that hypoxic conditions significantly improve generation of iPSC [139]. Growing knowledge of cancer stem cell biology suggests that hypoxia may act as a critical regulator of the cancer stem cell phenotype.



Series of experiments have demonstrated that hypoxia is responsible for altering the cellular phenotype by causing an increase in proliferation, self-renewal and upregulation of stem cell genes in both CSC and non-CSC. Several groups have shown that hypoxia can regulate histone methylation and thus alter the epigenetic status of cancer cells [140-142]. Tumor hypoxia also correlates with poor outcome of patients. HIFs were shown to induce the embryonal stem cell-like transcriptional program, including *OCT4*, *NANOG*, *SOX2*, *KLF4*, *MYC*, and microRNA-302 in cancer cell lines of prostate, brain, kidney, cervix, lung, colon, liver and breast tumors [143]. Hypoxic microenvironment potentiates biological effect of Notch signaling in adenocarcinoma of the lung or alters gene expression of neuroblastoma cells to induce more immature phenotype [144, 145]. CD133, a cancer stem cell marker, has been reported by several groups to be upregulated under hypoxic conditions [146, 147]. McCord et al. [2009] showed that hypoxia not only increased the sub-population of glioblastoma cells positive for CD133, but also enhanced expression of other stem cell markers, such as *SOX2*, *OCT4* and *nestin* [148]. Low oxygen levels induced also HIF-2 $\alpha$  expression that can increase the expression of stem cell-associated genes and confer tumorigenic potential to non-CSC [140].

The BM microenvironment of MM is also hypoxic, and myeloma PCs are long term exposed to low oxygen levels. Tumor adaptation to hypoxia is mediated by the production of HIF-1 [149]. Both HIF-1 $\alpha$  and HIF-2 $\alpha$  have been reported to be activated in MM patients resulting in stimulation of angiogenesis [150]. Although a role of BM microenvironment is generally recognized as a crucial factor affecting myeloma development and support progression, it is surprising that an importance of hypoxic microenvironment for modulation of plasma cell phenotype have never been studied in MM.

## 7. Conclusion

Despite the achievements of currently used therapy, MM remains difficult to treat. Novel agents such as inhibitors of proteasome or immunomodulatory drugs have prolonged survival of patients with MM and even some patients persist in long term remission. However, there is a little known about mechanisms of myeloma development or the population responsible for the origin and relapse of the disease. Many scientists have tried to explain causes of the relapse but none of their theories have been conclusively confirmed so far. In this review, we explain inconsistencies among particular concepts and the inability to detect cells of origin by the plasticity potential of myeloma PCs. Plasticity of myeloma PCs might be a cause of vast phenotypic heterogeneity of MM and different characteristics of putative myeloma precursors. Under specific signals from aberrant microenvironments, PCs might undergo dedifferentiation/transdifferentiation changing their phenotype profile, and acquire stem cell-like properties to ensure survival. Therefore, an effort to target the specific cell type based only on surface markers is not sufficient. Instead, it is necessary to concentrate on pathologic mechanisms responsible for the transition from non-CSC to CSC like cells. Additional focus on adjacent microenvironments and specific prevention of stem cell-like conversion might increase success of future therapy.

## Acknowledgements

This work was supported by grants from The Ministry of Education, Youth and Sports: MSM0021622434 and Czech Science Foundation GAP304/10/1395. The authors would like to thank Andrea Knight for proof reading the manuscript.

## Author details

Hana Šváchová<sup>1</sup>, Sabina Sevcikova<sup>1</sup> and Roman Hájek<sup>1,2</sup>

\*Address all correspondence to: [roman.hajek@fno.cz](mailto:roman.hajek@fno.cz)

1 Babak Myeloma Group, Department of Pathological Physiology, Faculty of Medicine, Masaryk University, Czech Republic

2 Faculty of Medicine, University of Ostrava and University Hospital of Ostrava, Czech Republic,, Czech Republic

## References

- [1] Rajkumar SV, Kyle RA. Multiple myeloma: diagnosis and treatment. *Mayo Clin Proc.* 2005 Oct;80(10):1371-82.
- [2] Richardson P, Anderson K. Thalidomide and dexamethasone: a new standard of care for initial therapy in multiple myeloma. *J Clin Oncol.* 2006 Jan;24(3):334-6.
- [3] Hamburger A, Salmon SE. Primary bioassay of human myeloma stem cells. *J Clin Invest.* 1977 Oct;60(4):846-54.
- [4] Bergsagel DE, Valeriote FA. Growth characteristics of a mouse plasma cell tumor. *Cancer Res.* 1968 Nov;28(11):2187-96.
- [5] Park CH, Bergsagel DE, McCulloch EA. Mouse myeloma tumor stem cells: a primary cell culture assay. *J Natl Cancer Inst.* 1971 Feb;46(2):411-22.
- [6] Matsui W, Huff CA, Wang Q, Malehorn MT, Barber J, Tanhehco Y, et al. Characterization of clonogenic multiple myeloma cells. *Blood.* 2004 Mar;103(6):2332-6.
- [7] Yaccoby S, Epstein J. The proliferative potential of myeloma plasma cells manifest in the SCID-hu host. *Blood.* 1999 Nov;94(10):3576-82.
- [8] Pfeifer S, Perez-Andres M, Ludwig H, Sahota SS, Zojer N. Evaluating the clonal hierarchy in light-chain multiple myeloma: implications against the myeloma stem cell hypothesis. *Leukemia.* 2011 Jul;25(7):1213-6.

- [9] Mellstedt H, Hammarström S, Holm G. Monoclonal lymphocyte population in human plasma cell myeloma. *Clin Exp Immunol*. 1974 Jul;17(3):371-84.
- [10] Vacca A, Ribatti D. Bone marrow angiogenesis in multiple myeloma. *Leukemia*. 2006 Feb;20(2):193-9.
- [11] Laroche M, Brousset P, Ludot I, Mazières B, Thiechart M, Attal M. Increased vascularization in myeloma. *Eur J Haematol*. 2001 Feb;66(2):89-93.
- [12] Sezer O, Niemöller K, Eucker J, Jakob C, Kaufmann O, Zavrski I, et al. Bone marrow microvessel density is a prognostic factor for survival in patients with multiple myeloma. *Ann Hematol*. 2000 Oct;79(10):574-7.
- [13] Rajkumar SV, Leong T, Roche PC, Fonseca R, Dispenzieri A, Lacy MQ, et al. Prognostic value of bone marrow angiogenesis in multiple myeloma. *Clin Cancer Res*. 2000 Aug;6(8):3111-6.
- [14] Rajkumar SV, Mesa RA, Fonseca R, Schroeder G, Plevak MF, Dispenzieri A, et al. Bone marrow angiogenesis in 400 patients with monoclonal gammopathy of undetermined significance, multiple myeloma, and primary amyloidosis. *Clin Cancer Res*. 2002 Jul;8(7):2210-6.
- [15] Liao D, Johnson RS. Hypoxia: a key regulator of angiogenesis in cancer. *Cancer Metastasis Rev*. 2007 Jun;26(2):281-90.
- [16] Brahimi-Horn MC, Chiche J, Pouyssegur J. Hypoxia and cancer. *J Mol Med (Berl)*. 2007 Dec;85(12):1301-7.
- [17] Colla S, Storti P, Donofrio G, Todoerti K, Bolzoni M, Lazzaretti M, et al. Low bone marrow oxygen tension and hypoxia-inducible factor-1 $\alpha$  overexpression characterize patients with multiple myeloma: role on the transcriptional and proangiogenic profiles of CD138(+) cells. *Leukemia*. 2010 Nov;24(11):1967-70.
- [18] Knowles HJ, Athanasou NA. Hypoxia-inducible factor is expressed in giant cell tumour of bone and mediates paracrine effects of hypoxia on monocyte-osteoclast differentiation via induction of VEGF. *J Pathol*. 2008 May;215(1):56-66.
- [19] Bergsagel PL, Masellis Smith A, Belch AR, Pilarski LM. The blood B-cells and bone marrow plasma cells in patients with multiple myeloma share identical IgH rearrangements. *Curr Top Microbiol Immunol*. 1995;194:17-24.
- [20] Bergsagel PL, Smith AM, Szczeppek A, Mant MJ, Belch AR, Pilarski LM. In multiple myeloma, clonotypic B lymphocytes are detectable among CD19+ peripheral blood cells expressing CD38, CD56, and monotypic Ig light chain. *Blood*. 1995 Jan;85(2):436-47.
- [21] Szczeppek AJ, Bergsagel PL, Axelsson L, Brown CB, Belch AR, Pilarski LM. CD34+ cells in the blood of patients with multiple myeloma express CD19 and IgH mRNA

and have patient-specific IgH VDJ gene rearrangements. *Blood*. 1997 Mar;89(5):1824-33.

- [22] Jensen GS, Belch AR, Mant MJ, Ruether BA, Yacyszyn BR, Pilarski LM. Expression of multiple beta 1 integrins on circulating monoclonal B cells in patients with multiple myeloma. *Am J Hematol*. 1993 May;43(1):29-36.
- [23] Pilarski LM, Hipperson G, Seeberger K, Pruski E, Coupland RW, Belch AR. Myeloma progenitors in the blood of patients with aggressive or minimal disease: engraftment and self-renewal of primary human myeloma in the bone marrow of NOD SCID mice. *Blood*. 2000 Feb;95(3):1056-65.
- [24] Rasmussen T, Jensen L, Johnsen HE. Levels of circulating CD19+ cells in patients with multiple myeloma. *Blood*. 2000 Jun;95(12):4020-1.
- [25] Kubagawa H, Vogler LB, Capra JD, Conrad ME, Lawton AR, Cooper MD. Studies on the clonal origin of multiple myeloma. Use of individually specific (idiotype) antibodies to trace the oncogenic event to its earliest point of expression in B-cell differentiation. *J Exp Med*. 1979 Oct;150(4):792-807.
- [26] Yaccoby S, Pennisi A, Li X, Dillon SR, Zhan F, Barlogie B, et al. Atacicept (TACI-Ig) inhibits growth of TACI(high) primary myeloma cells in SCID-hu mice and in coculture with osteoclasts. *Leukemia*. 2008 Feb;22(2):406-13.
- [27] Yaccoby S. The phenotypic plasticity of myeloma plasma cells as expressed by dedifferentiation into an immature, resilient, and apoptosis-resistant phenotype. *Clin Cancer Res*. 2005 Nov;11(21):7599-606.
- [28] Yata K, Yaccoby S. The SCID-rab model: a novel in vivo system for primary human myeloma demonstrating growth of CD138-expressing malignant cells. *Leukemia*. 2004 Nov;18(11):1891-7.
- [29] Chiron D, Surget S, Maïga S, Bataille R, Moreau P, Le Gouill S, et al. The peripheral CD138+ population but not the CD138- population contains myeloma clonogenic cells in plasma cell leukaemia patients. *Br J Haematol*. 2012 Mar;156(5):679-83.
- [30] Kim D, Park CY, Medeiros BC, Weissman IL. CD19(-)CD45(low/-)CD38(high)/CD138(+) plasma cells enrich for human tumorigenic myeloma cells. *Leukemia*. 2012 May.
- [31] Pilarski LM, Masellis-Smith A, Szczepiek A, Mant MJ, Belch AR. Circulating clonotypic B cells in the biology of multiple myeloma: speculations on the origin of myeloma. *Leuk Lymphoma*. 1996 Aug;22(5-6):375-83..
- [32] Szczepiek AJ, Seeberger K, Wizniak J, Mant MJ, Belch AR, Pilarski LM. A high frequency of circulating B cells share clonotypic Ig heavy-chain VDJ rearrangements with autologous bone marrow plasma cells in multiple myeloma, as measured by single-cell and in situ reverse transcriptase-polymerase chain reaction. *Blood*. 1998 Oct;92(8):2844-55.

- [33] Pilarski LM, Jensen GS. Monoclonal circulating B cells in multiple myeloma. A continuously differentiating, possibly invasive, population as defined by expression of CD45 isoforms and adhesion molecules. *Hematol Oncol Clin North Am.* 1992 Apr; 6(2):297-322.
- [34] Jensen GS, Mant MJ, Pilarski LM. Sequential maturation stages of monoclonal B lineage cells from blood, spleen, lymph node, and bone marrow from a terminal myeloma patient. *Am J Hematol.* 1992 Nov;41(3):199-208.
- [35] Conway EJ, Wen J, Feng Y, Mo A, Huang WT, Keever-Taylor CA, et al. Phenotyping studies of clonotypic B lymphocytes from patients with multiple myeloma by flow cytometry. *Arch Pathol Lab Med.* 2009 Oct;133(10):1594-9.
- [36] Billadeau D, Ahmann G, Greipp P, Van Ness B. The bone marrow of multiple myeloma patients contains B cell populations at different stages of differentiation that are clonally related to the malignant plasma cell. *J Exp Med.* 1993 Sep;178(3):1023-31.
- [37] Bakkus MH, Van Riet I, Van Camp B, Thielemans K. Evidence that the clonogenic cell in multiple myeloma originates from a pre-switched but somatically mutated B cell. *Br J Haematol.* 1994 May;87(1):68-74.
- [38] Reiman T, Seeberger K, Taylor BJ, Szczepek AJ, Hanson J, Mant MJ, et al. Persistent pre-switch clonotypic myeloma cells correlate with decreased survival: evidence for isotype switching within the myeloma clone. *Blood.* 2001 Nov;98(9):2791-9.
- [39] Matsui W, Wang Q, Barber JP, Brennan S, Smith BD, Borrello I, et al. Clonogenic multiple myeloma progenitors, stem cell properties, and drug resistance. *Cancer Res.* 2008 Jan;68(1):190-7.
- [40] Peacock CD, Wang Q, Gesell GS, Corcoran-Schwartz IM, Jones E, Kim J, et al. Hedgehog signaling maintains a tumor stem cell compartment in multiple myeloma. *Proc Natl Acad Sci U S A.* 2007 Mar;104(10):4048-53.
- [41] Huff CA, Matsui W. Multiple myeloma cancer stem cells. *J Clin Oncol.* 2008 Jun; 26(17):2895-900.
- [42] Loh YS, Mo S, Brown RD, Yamagishi T, Yang S, Joshua DE, et al. Presence of Hoechst low side populations in multiple myeloma. *Leuk Lymphoma.* 2008 Sep;49(9):1813-6.
- [43] Jakubikova J, Adamia S, Kost-Alimova M, Klippel S, Cervi D, Daley JF, et al. Lenalidomide targets clonogenic side population in multiple myeloma: pathophysiologic and clinical implications. *Blood.* 2011 Apr;117(17):4409-19.
- [44] Munshi NC, Anderson KC, Bergsagel PL, Shaughnessy J, Palumbo A, Durie B, et al. Consensus recommendations for risk stratification in multiple myeloma: report of the International Myeloma Workshop Consensus Panel 2. *Blood.* 2011 May;117(18): 4696-700.



- [45] O'Connor BP, Gleeson MW, Noelle RJ, Erickson LD. The rise and fall of long-lived humoral immunity: terminal differentiation of plasma cells in health and disease. *Immunol Rev.* 2003 Aug;194:61-76.
- [46] O'Connor BP, Cascalho M, Noelle RJ. Short-lived and long-lived bone marrow plasma cells are derived from a novel precursor population. *J Exp Med.* 2002 Mar;195(6):737-45.
- [47] Caligaris-Cappio F, Bergui L, Gregoret MG, Gaidano G, Gaboli M, Schena M, et al. Role of bone marrow stromal cells in the growth of human multiple myeloma. *Blood.* 1991 Jun;77(12):2688-93.
- [48] Chen BJ, Epstein J. Circulating clonal lymphocytes in myeloma constitute a minor subpopulation of B cells. *Blood.* 1996 Mar;87(5):1972-6.
- [49] Rasmussen T, Lodahl M, Hancke S, Johnsen HE. In multiple myeloma clonotypic CD38- /CD19+ / CD27+ memory B cells recirculate through bone marrow, peripheral blood and lymph nodes. *Leuk Lymphoma.* 2004 Jul;45(7):1413-7.
- [50] Vescio RA, Cao J, Hong CH, Lee JC, Wu CH, Der Danielian M, et al. Myeloma Ig heavy chain V region sequences reveal prior antigenic selection and marked somatic mutation but no intraclonal diversity. *J Immunol.* 1995 Sep;155(5):2487-97.
- [51] McSweeney PA, Wells DA, Shults KE, Nash RA, Bensinger WI, Buckner CD, et al. Tumor-specific aneuploidy not detected in CD19+ B-lymphoid cells from myeloma patients in a multidimensional flow cytometric analysis. *Blood.* 1996 Jul;88(2):622-32.
- [52] Taylor BJ, Kriangkum J, Pittman JA, Mant MJ, Reiman T, Belch AR, et al. Analysis of clonotypic switch junctions reveals multiple myeloma originates from a single class switch event with ongoing mutation in the isotype-switched progeny. *Blood.* 2008 Sep;112(5):1894-903.
- [53] Trepel M, Martens V, Doll C, Rahlff J, Gösch B, Loges S, et al. Phenotypic detection of clonotypic B cells in multiple myeloma by specific immunoglobulin ligands reveals their rarity in multiple myeloma. *PLoS One.* 2012;7(2):e31998.
- [54] Rasmussen T, Haaber J, Dahl IM, Knudsen LM, Kerndrup GB, Lodahl M, et al. Identification of translocation products but not K-RAS mutations in memory B cells from patients with multiple myeloma. *Haematologica.* 2010 Oct;95(10):1730-7.
- [55] Basak GW, Carrier E. The search for multiple myeloma stem cells: the long and winding road. *Biol Blood Marrow Transplant.* 2010 May;16(5):587-94.
- [56] Corradini P, Voena C, Omedé P, Astolfi M, Boccadoro M, Dalla-Favera R, et al. Detection of circulating tumor cells in multiple myeloma by a PCR-based method. *Leukemia.* 1993 Nov;7(11):1879-82.

- [57] Bergsagel PL, Chesi M, Nardini E, Brents LA, Kirby SL, Kuehl WM. Promiscuous translocations into immunoglobulin heavy chain switch regions in multiple myeloma. *Proc Natl Acad Sci U S A*. 1996 Nov;93(24):13931-6.
- [58] Fenton JA, Pratt G, Rawstron AC, Morgan GJ. Isotype class switching and the pathogenesis of multiple myeloma. *Hematol Oncol*. 2002 Jun;20(2):75-85.
- [59] Nemec P, Kuglík P, Hájek R. (The origin and formation of chromosomal translocations in multiple myeloma). *Klin Onkol*. 2008;21(2):53-8.
- [60] Palumbo A, Battaglio S, Astolfi M, Frieri R, Boccadoro M, Pileri A. Multiple independent immunoglobulin class-switch recombinations occurring within the same clone in myeloma. *Br J Haematol*. 1992 Dec;82(4):676-80.
- [61] Souto-Carneiro MM, Mahadevan V, Takada K, Fritsch-Stork R, Nanki T, Brown M, et al. Alterations in peripheral blood memory B cells in patients with active rheumatoid arthritis are dependent on the action of tumour necrosis factor. *Arthritis Res Ther*. 2009;11(3):R84.
- [62] Tangye SG, Good KL. Human IgM+CD27+ B cells: memory B cells or "memory" B cells? *J Immunol*. 2007 Jul;179(1):13-9.
- [63] Lim SH, Zhang Y, Wang Z, Varadarajan R, Periman P, Esler WV. Rituximab administration following autologous stem cell transplantation for multiple myeloma is associated with severe IgM deficiency. *Blood*. 2004 Mar;103(5):1971-2.
- [64] Musto P, Carella AM, Greco MM, Falcone A, Sanpaolo G, Bodenizza C, et al. Short progression-free survival in myeloma patients receiving rituximab as maintenance therapy after autologous transplantation. *Br J Haematol*. 2003 Nov;123(4):746-7.
- [65] Paíno T, Ocio EM, Paiva B, San-Segundo L, Garayoa M, Gutiérrez NC, et al. CD20 positive cells are undetectable in the majority of multiple myeloma cell lines and are not associated with a cancer stem cell phenotype. *Haematologica*. 2012 Jul;97(7):1110-4.
- [66] Rossi EA, Rossi DL, Stein R, Goldenberg DM, Chang CH. A bispecific antibody-IFN- $\alpha$ 2b immunocytokine targeting CD20 and HLA-DR is highly toxic to human lymphoma and multiple myeloma cells. *Cancer Res*. 2010 Oct;70(19):7600-9.
- [67] Hosen N, Matsuoka Y, Kishida S, Nakata J, Mizutani Y, Hasegawa K, et al. CD138-negative clonogenic cells are plasma cells but not B cells in some multiple myeloma patients. *Leukemia*. 2012 Sep;26(9):2135-41.
- [68] Christensen JH, Jensen PV, Kristensen IB, Abildgaard N, Lodahl M, Rasmussen T. Characterization of potential CD138 negative myeloma "stem cells". *Haematologica*. 2012 Jun;97(6):e18-20.

- [69] Jourdan M, Ferlin M, Legouffe E, Horvathova M, Liautard J, Rossi JF, et al. The myeloma cell antigen syndecan-1 is lost by apoptotic myeloma cells. *Br J Haematol*. 1998 Mar;100(4):637-46.
- [70] Goodell MA, Brose K, Paradis G, Conner AS, Mulligan RC. Isolation and functional properties of murine hematopoietic stem cells that are replicating in vivo. *J Exp Med*. 1996 Apr;183(4):1797-806.
- [71] Wu C, Alman BA. Side population cells in human cancers. *Cancer Lett*. 2008 Sep;268(1):1-9.
- [72] Srinivasan V, Sivaramakrishnan H, Karthikeyan B. Detection, isolation and characterization of principal synthetic route indicative impurities in verapamil hydrochloride. *Sci Pharm*. 2011 Sep;79(3):555-68.
- [73] Montanaro F, Liadaki K, Schiend J, Flint A, Gussoni E, Kunkel LM. Demystifying SP cell purification: viability, yield, and phenotype are defined by isolation parameters. *Exp Cell Res*. 2004 Aug;298(1):144-54.
- [74] O'Connor BP, Cascalho M, Noelle RJ. Short-lived and long-lived bone marrow plasma cells are derived from a novel precursor population. *J Exp Med*. 2002 Mar;195(6):737-45.
- [75] Kukreja A, Hutchinson A, Dhodapkar K, Mazumder A, Vesole D, Angitapalli R, et al. Enhancement of clonogenicity of human multiple myeloma by dendritic cells. *J Exp Med*. 2006 Aug;203(8):1859-65.
- [76] Kumar S, Kimlinger T, Morice W. Immunophenotyping in multiple myeloma and related plasma cell disorders. *Best Pract Res Clin Haematol*. 2010 Sep;23(3):433-51.
- [77] Paiva B, Almeida J, Pérez-Andrés M, Mateo G, López A, Rasillo A, et al. Utility of flow cytometry immunophenotyping in multiple myeloma and other clonal plasma cell-related disorders. *Cytometry B Clin Cytom*. 2010 Jul;78(4):239-52.
- [78] Busslinger M. Transcriptional control of early B cell development. *Annu Rev Immunol*. 2004;22:55-79.
- [79] Calame K. MicroRNA-155 function in B Cells. *Immunity*. 2007 Dec;27(6):825-7.
- [80] Harada H, Kawano MM, Huang N, Harada Y, Iwato K, Tanabe O, et al. Phenotypic difference of normal plasma cells from mature myeloma cells. *Blood*. 1993 May;81(10):2658-63.
- [81] Mahmoud MS, Huang N, Nobuyoshi M, Lisukov IA, Tanaka H, Kawano MM. Altered expression of Pax-5 gene in human myeloma cells. *Blood*. 1996 May;87(10):4311-5.
- [82] Nutt SL, Heavey B, Rolink AG, Busslinger M. Commitment to the B-lymphoid lineage depends on the transcription factor Pax5. *Nature*. 1999 Oct;401(6753):556-62.

- [83] Liu S, Otsuyama K, Ma Z, Abroun S, Shamsasenjan K, Amin J, et al. Induction of multilineage markers in human myeloma cells and their down-regulation by interleukin 6. *Int J Hematol*. 2007 Jan;85(1):49-58.
- [84] Allan AL. *Cancer Stem Cells in Solid Tumors*. Stem Cell Biology and Regenerative Medicine: Springer, 2011; 2011. 475 p.
- [85] Takahashi K, Yamanaka S. Induction of pluripotent stem cells from mouse embryonic and adult fibroblast cultures by defined factors. *Cell*. 2006 Aug;126(4):663-76.
- [86] Yu J, Vodyanik MA, Smuga-Otto K, Antosiewicz-Bourget J, Frane JL, Tian S, et al. Induced pluripotent stem cell lines derived from human somatic cells. *Science*. 2007 Dec;318(5858):1917-20.
- [87] Lowry WE, Richter L, Yachechko R, Pyle AD, Tchieu J, Sridharan R, et al. Generation of human induced pluripotent stem cells from dermal fibroblasts. *Proc Natl Acad Sci U S A*. 2008 Feb;105(8):2883-8.
- [88] Park IH, Lerou PH, Zhao R, Huo H, Daley GQ. Generation of human-induced pluripotent stem cells. *Nat Protoc*. 2008;3(7):1180-6.
- [89] Takahashi K, Tanabe K, Ohnuki M, Narita M, Ichisaka T, Tomoda K, et al. Induction of pluripotent stem cells from adult human fibroblasts by defined factors. *Cell*. 2007 Nov;131(5):861-72.
- [90] Hanna J, Markoulaki S, Schorderet P, Carey BW, Beard C, Wernig M, et al. Direct reprogramming of terminally differentiated mature B lymphocytes to pluripotency. *Cell*. 2008 Apr;133(2):250-64.
- [91] Aasen T, Raya A, Barrero MJ, Garreta E, Consiglio A, Gonzalez F, et al. Efficient and rapid generation of induced pluripotent stem cells from human keratinocytes. *Nat Biotechnol*. 2008 Nov;26(11):1276-84.
- [92] Maherali N, Hochedlinger K. Guidelines and techniques for the generation of induced pluripotent stem cells. *Cell Stem Cell*. 2008 Dec;3(6):595-605.
- [93] Aoi T, Yae K, Nakagawa M, Ichisaka T, Okita K, Takahashi K, et al. Generation of pluripotent stem cells from adult mouse liver and stomach cells. *Science*. 2008 Aug;321(5889):699-702.
- [94] Mani SA, Guo W, Liao MJ, Eaton EN, Ayyanan A, Zhou AY, et al. The epithelial-mesenchymal transition generates cells with properties of stem cells. *Cell*. 2008 May;133(4):704-15.
- [95] Prindull G, Zipori D. Environmental guidance of normal and tumor cell plasticity: epithelial mesenchymal transitions as a paradigm. *Blood*. 2004 Apr;103(8):2892-9.
- [96] Dhodapkar MV. Immunity to stemness genes in human cancer. *Curr Opin Immunol*. 2010 Apr;22(2):245-50.

- [97] Chaffer CL, Brueckmann I, Scheel C, Kaestli AJ, Wiggins PA, Rodrigues LO, et al. Normal and neoplastic nonstem cells can spontaneously convert to a stem-like state. *Proc Natl Acad Sci U S A*. 2011 May;108(19):7950-5.
- [98] Bergsagel PL, Kuehl WM. Chromosome translocations in multiple myeloma. *Oncogene*. 2001 Sep;20(40):5611-22.
- [99] Greipp PR, San Miguel J, Durie BG, Crowley JJ, Barlogie B, Bladé J, et al. International staging system for multiple myeloma. *J Clin Oncol*. 2005 May;23(15):3412-20.
- [100] Clevers H. Stem cells, asymmetric division and cancer. *Nat Genet*. 2005 Oct;37(10):1027-8.
- [101] Chesi M, Robbiani DF, Sebag M, Chng WJ, Affer M, Tiedemann R, et al. AID-dependent activation of a MYC transgene induces multiple myeloma in a conditional mouse model of post-germinal center malignancies. *Cancer Cell*. 2008 Feb;13(2):167-80.
- [102] Jungbluth AA, Ely S, DiLiberto M, Niesvizky R, Williamson B, Frosina D, et al. The cancer-testis antigens CT7 (MAGE-C1) and MAGE-A3/6 are commonly expressed in multiple myeloma and correlate with plasma-cell proliferation. *Blood*. 2005 Jul;106(1):167-74.
- [103] Schoenhals M, Kassambara A, De Vos J, Hose D, Moreaux J, Klein B. Embryonic stem cell markers expression in cancers. *Biochem Biophys Res Commun*. 2009 May;383(2):157-62.
- [104] Spisek R, Kukreja A, Chen LC, Matthews P, Mazumder A, Vesole D, et al. Frequent and specific immunity to the embryonal stem cell-associated antigen SOX2 in patients with monoclonal gammopathy. *J Exp Med*. 2007 Apr;204(4):831-40.
- [105] Zhu L, Somlo G, Zhou B, Shao J, Bedell V, Slovak ML, et al. Fibroblast growth factor receptor 3 inhibition by short hairpin RNAs leads to apoptosis in multiple myeloma. *Mol Cancer Ther*. 2005 May;4(5):787-98.
- [106] Dhodapkar MV, Dhodapkar KM. Spontaneous and therapy-induced immunity to pluripotency genes in humans: clinical implications, opportunities and challenges. *Cancer Immunol Immunother*. 2011 Mar;60(3):413-8.
- [107] Bataille R, Pellat-Deceunynck C, Robillard N, Avet-Loiseau H, Harousseau JL, Moreau P. CD117 (c-kit) is aberrantly expressed in a subset of MGUS and multiple myeloma with unexpectedly good prognosis. *Leuk Res*. 2008 Mar;32(3):379-82..
- [108] Schmidt-Hieber M, Pérez-Andrés M, Paiva B, Flores-Montero J, Perez JJ, Gutierrez NC, et al. CD117 expression in gammopathies is associated with an altered maturation of the myeloid and lymphoid hematopoietic cell compartments and favorable disease features. *Haematologica*. 2011 Feb;96(2):328-32.



- [109] Michalczyk K, Ziman M. Nestin structure and predicted function in cellular cytoskeletal organisation. *Histol Histopathol.* 2005 Apr;20(2):665-71.
- [110] Svachova H, Pour L, Sana J, Kovarova L, Raja KR, Hajek R. Stem cell marker nestin is expressed in plasma cells of multiple myeloma patients. *Leuk Res.* 2011 Aug;35(8):1008-13.
- [111] Blau HM, Brazelton TR, Weimann JM. The evolving concept of a stem cell: entity or function? *Cell.* 2001 Jun;105(7):829-41.
- [112] Hockfield S, McKay RD. Identification of major cell classes in the developing mammalian nervous system. *J Neurosci.* 1985 Dec;5(12):3310-28.
- [113] About I, Laurent-Maquin D, Lendahl U, Mitsiadis TA. Nestin expression in embryonic and adult human teeth under normal and pathological conditions. *Am J Pathol.* 2000 Jul;157(1):287-95.
- [114] Ha Y, Kim TS, Yoon DH, Cho YE, Huh SG, Lee KC. Reinduced expression of developmental proteins (nestin, small heat shock protein) in and around cerebral arteriovenous malformations. *Clin Neuropathol.* 2003 Sep-Oct;22(5):252-61.
- [115] Yamada H, Takano T, Ito Y, Matsuzuka F, Miya A, Kobayashi K, et al. Expression of nestin mRNA is a differentiation marker in thyroid tumors. *Cancer Lett.* 2009 Jul;280(1):61-4.
- [116] Wiese C, Rolletschek A, Kania G, Blyszczuk P, Tarasov KV, Tarasova Y, et al. Nestin expression--a property of multi-lineage progenitor cells? *Cell Mol Life Sci.* 2004 Oct;61(19-20):2510-22.
- [117] Cheng L, Jin Z, Liu L, Yan Y, Li T, Zhu X, et al. Characterization and promoter analysis of the mouse nestin gene. *FEBS Lett.* 2004 May;565(1-3):195-202.
- [118] Han DW, Do JT, Araúzo-Bravo MJ, Lee SH, Meissner A, Lee HT, et al. Epigenetic hierarchy governing Nestin expression. *Stem Cells.* 2009 May;27(5):1088-97.
- [119] Guérette D, Khan PA, Savard PE, Vincent M. Molecular evolution of type VI intermediate filament proteins. *BMC Evol Biol.* 2007;7:164.
- [120] Marvin MJ, Dahlstrand J, Lendahl U, McKay RD. A rod end deletion in the intermediate filament protein nestin alters its subcellular localization in neuroepithelial cells of transgenic mice. *J Cell Sci.* 1998 Jul;111 ( Pt 14):1951-61.
- [121] Grigelionienė G, Blennow M, Török C, Fried G, Dahlin I, Lendahl U, et al. Cerebrospinal fluid of newborn infants contains a deglycosylated form of the intermediate filament nestin. *Pediatr Res.* 1996 Dec;40(6):809-14.
- [122] Sahlgren CM, Mikhailov A, Hellman J, Chou YH, Lendahl U, Goldman RD, et al. Mitotic reorganization of the intermediate filament protein nestin involves phosphorylation by cdc2 kinase. *J Biol Chem.* 2001 May;276(19):16456-63.

- [123] Sjöberg G, Jiang WQ, Ringertz NR, Lendahl U, Sejersen T. Colocalization of nestin and vimentin/desmin in skeletal muscle cells demonstrated by three-dimensional fluorescence digital imaging microscopy. *Exp Cell Res*. 1994 Oct;214(2):447-58.
- [124] Coulombe PA, Wong P. Cytoplasmic intermediate filaments revealed as dynamic and multipurpose scaffolds. *Nat Cell Biol*. 2004 Aug;6(8):699-706.
- [125] Steinert PM, Chou YH, Prahlad V, Parry DA, Marekov LN, Wu KC, et al. A high molecular weight intermediate filament-associated protein in BHK-21 cells is nestin, a type VI intermediate filament protein. Limited co-assembly in vitro to form heteropolymers with type III vimentin and type IV alpha-internexin. *J Biol Chem*. 1999 Apr;274(14):9881-90.
- [126] Lendahl U, Zimmerman LB, McKay RD. CNS stem cells express a new class of intermediate filament protein. *Cell*. 1990 Feb;60(4):585-95.
- [127] Kachinsky AM, Dominov JA, Miller JB. Intermediate filaments in cardiac myogenesis: nestin in the developing mouse heart. *J Histochem Cytochem*. 1995 Aug;43(8):843-7.
- [128] Vaittinen S, Lukka R, Sahlgren C, Hurme T, Rantanen J, Lendahl U, et al. The expression of intermediate filament protein nestin as related to vimentin and desmin in regenerating skeletal muscle. *J Neuropathol Exp Neurol*. 2001 Jun;60(6):588-97.
- [129] Chou YH, Khuon S, Herrmann H, Goldman RD. Nestin promotes the phosphorylation-dependent disassembly of vimentin intermediate filaments during mitosis. *Mol Biol Cell*. 2003 Apr;14(4):1468-78.
- [130] Sahlgren CM, Pallari HM, He T, Chou YH, Goldman RD, Eriksson JE. A nestin scaffold links Cdk5/p35 signaling to oxidant-induced cell death. *EMBO J*. 2006 Oct;25(20):4808-19.
- [131] Reimer R, Helmbold H, Szalay B, Hagel C, Hohenberg H, Deppert W, et al. Nestin modulates glucocorticoid receptor function by cytoplasmic anchoring. *PLoS One*. 2009;4(6):e6084.
- [132] Shih AH, Holland EC. Notch signaling enhances nestin expression in gliomas. *Neoplasia*. 2006 Dec;8(12):1072-82.
- [133] Schwarzer R, Kaiser M, Acikgoez O, Heider U, Mathas S, Preissner R, et al. Notch inhibition blocks multiple myeloma cell-induced osteoclast activation. *Leukemia*. 2008 Dec;22(12):2273-7.
- [134] Atanackovic D, Hildebrandt Y, Jadczyk A, Cao Y, Luetkens T, Meyer S, et al. Cancer-testis antigens MAGE-C1/CT7 and MAGE-A3 promote the survival of multiple myeloma cells. *Haematologica*. 2010 May;95(5):785-93.
- [135] Adams JM, Strasser A. Is tumor growth sustained by rare cancer stem cells or dominant clones? *Cancer Res*. 2008 Jun;68(11):4018-21.

- [136] Means AL, Meszoely IM, Suzuki K, Miyamoto Y, Rustgi AK, Coffey RJ, et al. Pancreatic epithelial plasticity mediated by acinar cell transdifferentiation and generation of nestin-positive intermediates. *Development*. 2005 Aug;132(16):3767-76.
- [137] Tosh D, Slack JM. How cells change their phenotype. *Nat Rev Mol Cell Biol*. 2002 Mar;3(3):187-94.
- [138] Heddlestone JM, Li Z, Lathia JD, Bao S, Hjelmeland AB, Rich JN. Hypoxia inducible factors in cancer stem cells. *Br J Cancer*. 2010 Mar;102(5):789-95.
- [139] Yoshida Y, Takahashi K, Okita K, Ichisaka T, Yamanaka S. Hypoxia enhances the generation of induced pluripotent stem cells. *Cell Stem Cell*. 2009 Sep;5(3):237-41.
- [140] Heddlestone JM, Li Z, McLendon RE, Hjelmeland AB, Rich JN. The hypoxic microenvironment maintains glioblastoma stem cells and promotes reprogramming towards a cancer stem cell phenotype. *Cell Cycle*. 2009 Oct;8(20):3274-84.
- [141] Pan ZH, Chen ZD. (Hypoxia-inducible factor 1 and prostate cancer). *Zhonghua Nan Ke Xue*. 2007 Apr;13(4):356-9.
- [142] Xia X, Lemieux ME, Li W, Carroll JS, Brown M, Liu XS, et al. Integrative analysis of HIF binding and transactivation reveals its role in maintaining histone methylation homeostasis. *Proc Natl Acad Sci U S A*. 2009 Mar;106(11):4260-5.
- [143] Mathieu J, Zhang Z, Zhou W, Wang AJ, Heddlestone JM, Pinna CM, et al. HIF induces human embryonic stem cell markers in cancer cells. *Cancer Res*. 2011 Jul;71(13):4640-52.
- [144] Chen YQ, Zhao CL, Li W. Effect of hypoxia-inducible factor-1alpha on transcription of survivin in non-small cell lung cancer. *J Exp Clin Cancer Res*. 2009;28:29.
- [145] Jögi A, Øra I, Nilsson H, Lindeheim A, Makino Y, Poellinger L, et al. Hypoxia alters gene expression in human neuroblastoma cells toward an immature and neural crest-like phenotype. *Proc Natl Acad Sci U S A*. 2002 May;99(10):7021-6.
- [146] Seidel S, Garvalov BK, Wirta V, von Stechow L, Schänzer A, Meletis K, et al. A hypoxic niche regulates glioblastoma stem cells through hypoxia inducible factor 2 alpha. *Brain*. 2010 Apr;133(Pt 4):983-95.
- [147] Soeda A, Park M, Lee D, Mintz A, Androutsellis-Theotokis A, McKay RD, et al. Hypoxia promotes expansion of the CD133-positive glioma stem cells through activation of HIF-1alpha. *Oncogene*. 2009 Nov;28(45):3949-59.
- [148] McCord AM, Jamal M, Shankavaram UT, Shankavarum UT, Lang FF, Camphausen K, et al. Physiologic oxygen concentration enhances the stem-like properties of CD133+ human glioblastoma cells in vitro. *Mol Cancer Res*. 2009 Apr;7(4):489-97.
- [149] Hickey MM, Simon MC. Regulation of angiogenesis by hypoxia and hypoxia-inducible factors. *Curr Top Dev Biol*. 2006;76:217-57.

- [150] Martin SK, Diamond P, Williams SA, To LB, Peet DJ, Fujii N, et al. Hypoxia-inducible factor-2 is a novel regulator of aberrant CXCL12 expression in multiple myeloma plasma cells. *Haematologica*. 2010 May;95(5):776-84.

IntechOpen

IntechOpen

