

# We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

186,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index  
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?  
Contact [book.department@intechopen.com](mailto:book.department@intechopen.com)

Numbers displayed above are based on latest data collected.  
For more information visit [www.intechopen.com](http://www.intechopen.com)



---

# SUMO Tasks in Chromatin Remodeling

---

Garcia-Dominguez Mario

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/55395>

---

## 1. Introduction

Post-translational modifications implicate attachment of diverse molecules to proteins after translation. These modifications are essential for many biological processes as they are involved in their regulation. From relatively simple molecules to small polypeptides are common covalent modifiers of proteins. Sumoylation consists in the post-translational modification of proteins by attachment of the small polypeptide SUMO (small ubiquitin-like modifier). This post-translational modification was identified two decades ago and has been very actively investigated to date. Sumoylation has consequences on protein structure and regulation. This modification controls many processes in the eukaryotic cell and is essential for viability of all the organisms studied so far.

From the discovery of ubiquitin in 1975, a number of ubiquitin-like proteins (UBLs) have been identified in eukaryotes and it has been shown that many of them are able to covalently attach to other proteins (reviewed in [1]). Several aspects are common to most UBLs: they are small polypeptides (less than 200 amino acids) capable of attaching to other macromolecules in a covalent way, present common structural features and use similar modification pathways. These characteristics strongly support duplication and diversification during evolution as the origin of the different pathways. Ubiquitin and UBLs are characterized by the presence of the  $\beta$ -grasp fold, which also appears in ubiquitin-like domains of several other proteins of the ubiquitin system and in numerous non-related proteins (reviewed in [2]). The  $\beta$ -grasp fold seems to have emerged in prokaryotes as a translation-related RNA-binding module, which diversified structurally and biochemically before to dramatically expand in eukaryotes [2]. Besides ubiquitin and SUMO, examples of UBLs are NEDD8, FUBI, FAT10, ISG15, UFM1, Atg8, Atg12 and Urm1 (reviewed in [1]).


The first report of a protein being modified by SUMO occurred in the nineties and concerned the mammalian nuclear pore-associated GTPase activating protein RanGAP1 [3, 4]. Subse-

quently, more than a hundred proteins have been identified as SUMO substrates. Although similarities with ubiquitin are notable [5], SUMO plays many regulatory functions in the cell that significantly differ from the major role displayed by ubiquitin: labeling proteins to target them for proteasomal degradation [6]. A variety of consequences derived from protein sumoylation (new interaction surfaces, modulation of protein affinity and binding capacities to other molecules, modulation of protein activity, blocking of protein domains, steric hindrance, crosstalk or interference with other post-translational modifications) account for the many roles attributed to SUMO (reviewed in [7]). A major role of SUMO is associated with RanGAP1 and thereby with the nuclear pore complex. Thus, involvement of SUMO in nucleocytoplasmic transport of proteins has been well established [8]. SUMO has been also implicated in chromosome dynamics in mitosis and meiosis (condensation, cohesion, separation) and genome integrity, as many proteins involved in DNA replication, repair and recombination are modulated by SUMO modification (reviewed in [7]). Other roles attributed to SUMO are related to enzyme regulation, protein stability and cellular structure (reviewed in [9, 10]). However, the most prominent function of SUMO concerns transcriptional regulation, and specially transcription repression (reviewed in [11, 12]). The role of SUMO in transcription, in the context of chromatin structure and dynamics, is analyzed in this chapter.

## 2. The modification pathway

### 2.1. Enzymes involved

Modification by SUMO involves the ATP-dependent activation of mature SUMO (C terminus of SUMO needs to be excised by proteolysis) by the E1 enzyme, transfer to the E2 enzyme UBC9 and conjugation to the target protein, often mediated by a SUMO ligase or E3 (Figure 1 and Table 1) (reviewed in [9]). Maturation of the SUMO precursor, as well as removal of SUMO from targets is displayed by SUMO specific proteases.



**Figure 1.** The sumoylation pathway. Cleavage of the SUMO C terminus enables ATP-mediated activation and binding to the E1 enzyme to be transferred to the E2 conjugating enzyme UBC9, which mediates target modification with the concurrence of an E3 SUMO ligase. Recycling of SUMO is performed by the same proteases involved in maturation.

SUMO E1 activity is performed by the SAE1/UBA2 heterodimer in human, in contrast to the ubiquitination pathway where the E1 activity is displayed by a monomeric enzyme. However, the SAE1 subunit is homologous to the N-terminal part of ubiquitin E1, while the UBA2 subunit

is homologous to its C terminus [13]. Both monomers work together and are not found separately [14]. E1 activation of mature SUMO involves ATP hydrolysis and formation of a thiolester bond between E1 and the C terminus of SUMO before being transferred to the E2. While several E2 have been described for ubiquitination, UBC9 is the only E2 known for sumoylation [15, 16]. Thus, UBC9 is the conjugating enzyme directly involved in attachment of SUMO to the different substrates. This second step of the sumoylation reaction involves the formation of a thiolester bond between SUMO and UBC9 upon transfer from the E1. The region surrounding the active site cysteine (C93 in mammals) in UBC9 is able to directly interact with sumoylation consensus sequence (see below) in target proteins [17-19].

Enzyme	Protein	Activity	References
E1 (activating)	AOS1/UBA2	ATP-mediated activation of SUMO	[13]
E2 (conjugating)	UBC9	SUMO conjugation to target	[15, 16]
E3 (ligase)	PIAS1-4	Facilitates transfer to target	[20, 21]
	RanBP2		[22]
	Polycomb-2 (Pc2)		[23]
	TOPORS		[24]
	Class IIa HDACs		[25]
	KAP-1		[26]
	RHES		[27]
	Krox20		[28]
protease	SEN1-3, 5-7	Maturation/recycling	[29-31]
	DeSI-1		[32]

**Table 1.** Enzymes involved in SUMO conjugation.

SUMO ligases are involved in facilitating the SUMO attachment to substrates (reviewed in [33]). To date, few ligases have been described for sumoylation, in contrast to ubiquitination, where lots of them are known to play an essential role and mediate substrate specificity. In fact, SUMO ligases were undervalued at the beginning, since certain substrates are sumoylated in vitro, provided that E1 and E2 are present at the adequate concentrations. Since UBC9 is able to directly interact with sumoylation consensus sequence in substrates, it is able to render sumoylation in the absence of a ligase. However, a number of proteins, which augmented the efficiency of SUMO conjugation, were identified. The list of SUMO ligases progressively increases and essential roles for these have been described in vivo (see [34]). Although mechanisms of action of SUMO ligases have not been completely elucidated, it is obvious that many ligases facilitate transfer of SUMO by bringing together SUMO-loaded UBC9 and the target protein. Thus, similar to the RING domain-containing E3 ligases involved in ubiquitination, SUMO ligases do not establish a covalent bond with SUMO. In this context, a SUMO ligase should normally i) interact with the substrate, ii) interact with UBC9, iii) facilitate SUMO

transfer to the substrate. Ligases of the PIAS family (PIAS1 to 4) have been extensively studied [35]. They present a type of RING finger domain, the SP-RING (Siz/PIAS RING), for UBC9 interaction, although Ubc9 binding to a PHD domain in plant PIAS proteins has been also described [36]. Other SUMO ligases described so far are RanBP2 [22], the Polycomb-2 (Pc2) protein [23], class IIa histone deacetylases (HDACs) [25], topoisomerase I-binding RING finger protein (TOPORS) [24], the PHD containing protein KAP-1 [26], Ras homologue enriched in striatum (RHES) [27] and the transcription factor Krox20 [28]. In contrast to most ubiquitin ligases, SUMO ligases may display significant promiscuity, as many of them enhance sumoylation of a variety of substrates.

SUMO proteases are involved in maturation of the SUMO precursor by exposing two glycine residues at the C terminus for binding to E1 [37]. In addition, they are also involved in SUMO recycling by excising the SUMO moiety from substrates. Yeast Ulp1p was the first SUMO protease identified [29]. Sequence analysis revealed that it corresponded to a protease of the C48 cysteine group, not related to deubiquitylating enzymes but similar to adenovirus proteases. Mammalian SUMO proteases are represented by the SENP (sentrin-specific protease) family. It comprises six members, SENP1 to 3, and SENP5 to 7 [38]. A seventh member, initially identified as SENP4, resulted to actually correspond to SENP3. Besides SENP1 to 7, an additional family member has been reported, SENP8. However, this protease does not act on SUMO, but on another UBL, NEDD8 [39, 40]. Very recently, a new type of SUMO protease has been described, the desumoylating isopeptidase 1 (DeSI-1) [32]. The different SUMO proteases show diverse cellular localization and different specificities for the various SUMO molecules and substrates (reviewed in [38]).

## 2.2. SUMO molecules

Four different SUMO molecules have been described in mammals: SUMO1 to 4. SUMO1 has been implicated in regulation of many processes, while SUMO2 and SUMO3 are highly related with the response to stress. Consequently, a significant pool of free SUMO2 and SUMO3 is detected in the cell, which is rapidly mobilized after exposure to a variety of stress conditions. In contrast, most of SUMO1 appears conjugated to proteins [41]. SUMO2 and SUMO3 are usually referred as SUMO2/3, as they share 97% identity and antibodies hardly differentiate the two forms. By contrast, SUMO1 only shares about 50% identity with SUMO2/3. Despite the low similarity showed between ubiquitin and SUMO (about 18% identity with SUMO1), structurally they are quite similar, excepting the N-terminal region of SUMO not present in ubiquitin [42]. A remarkable difference between SUMO1 and SUMO2/3 is the ability of this last to form poly-SUMO chains *in vitro* as well as *in vivo*, due to the presence of a sumoylation consensus sequence in the molecule [43]. SUMO4 is the last SUMO molecule identified. It shows a restricted expression pattern [44] and several data bring into question its capacity to be conjugated to proteins [45]. However, a polymorphism found in human SUMO4 correlates with type 1 diabetes [46]. The different SUMO molecules share a common modification pathway and the existence of functional redundancy has been suggested. However, specific modification by the different SUMO paralogs has been implicated in the regulation of a variety



of processes. Thus, the modification pathway is able to differentially conjugate the various SUMO molecules depending on the substrate or the regulatory process [47].

### 2.3. Sumoylation consensus motifs and SUMO interacting motifs

Covalent attachment of SUMO occurs through the  $\epsilon$ -amino group of a lysine residue in target proteins. In many cases the Lys (K) residue is the core of the consensus sequence  $\Psi KxE$ , being  $\Psi$  a large hydrophobic residue and  $x$  any amino acid. Extended consensus (phosphorylation-dependent SUMO motif (PDSM) and negatively charged residues-dependent SUMO motif (NDSM)) and variations have been described as well (Table 2) (reviewed in [34]). However, sumoylation also occurs at non-consensus sequences. As mentioned above, the consensus sequence is directly contacted by UBC9. Thus, it is possible that when sumoylation occurs at non-consensus sequences, certain amino acid residues, otherwise dispersed in the primary structure of the target protein, bring together in the three-dimensional structure to mimic a consensus-like environment. It is worth noting that conversely, sumoylation consensus sequences in a protein are not always substrate for SUMO attachment, indicating that additional structural features regulate and enable modification by SUMO. Besides covalent attachment of SUMO, many proteins can associate with SUMO in a different way involving a non-covalent interaction (reviewed in [48]). This occurs through SUMO interacting motifs (SIMs) in proteins. SIMs are usually characterized by the presence of a short hydrophobic region surrounded by negatively charged residues (Table 2) [49]. The non-covalent interaction of proteins with SUMO has been revealed essential in the regulation of several processes. In a variety of cases function of the system relies in the combinatorial occurrence of sumoylation sites and SIMs in a given protein or in different subunits of a complex, which determines its macromolecular architecture (Figure 2 and see below). This situation is exemplified by the promyelocytic leukaemia protein (PML), in which combination of sumoylation sites and SIMs dictates the formation of PML nuclear bodies and the recruitment of additional proteins [48].

### 2.4. Regulation of sumoylation

Despite that certain SUMO targets appear constitutively sumoylated, it is obvious that sumoylation, as a signaling pathway needs to be regulated. A striking feature of SUMO modification consists in the so-called “SUMO enigma” [9]. It has been observed that many SUMO targets are difficult to detect at the sumoylated state, but mutation of the acceptor lysine has severe consequences in the process involved. In other words, at the steady state, only a low proportion of the whole pool of a given target appears sumoylated, although sumoylation results essential for function of the target. Thus, sumoylation has been suggested to be a highly dynamic and transient modification that permanently marks targets for specific fates even though the SUMO moiety has been removed [9]. This can be explained by viewing sumoylation as a temporal facilitator for the establishment of protein interactions, other protein modifications, or sub-cellular localization (Figure 2).

Sumoylation can be regulated at different levels (reviewed in [34, 50]). First level of regulation in SUMO modification relies in the nature of target proteins, as target sequence, structural features, and other protein modifications affect attachment of SUMO. The other way to

SUMO binding	Type		Sequence
Sumoylation site	Consensus		ΨKxE
	Extended consensus	PDSM	ΨKxE <sub>x</sub> SP
		NDSM	ΨKxE <sub>x</sub> [D/E] <sub>n</sub>
	Inverted consensus		ExKΨ
	Phosphorylated Ser		ΨKxS
SIM	ZNF198		DDDDDDDD VVFI
	PIAS1		VEVI DLTl DSSSDDEEEE
	SP100		IIVI SSEDSEGSTDVD
	PML		EE R VVVI SSEDSD
	RanBP2		SDSPSDDD VLIV
	CoREST1		EESEDELEE ANGNNP IDIEV


**Table 2.** Sumoylation sites and SUMO interacting motifs (SIMs). Ψ represents a large hydrophobic residue. Sumoylated Lys (K) is frequently close to a hydrophobic residue and to negatively charged environment, either acidic residues Asp/Glu (D/E) or phosphorylation sites (SP). SIMs usually consist in a stretch of 4 amino acids, containing at least 3 hydrophobic residues, close to an acidic region (Asp/Glu) (D/E) or putative phosphorylation sites (Ser/Thr) (S/T). Examples of SIMs with acidic/phosphorylation region N-terminal to the hydrophobic core (ZNF198), C-terminal (with spacer (PIAS1), without spacer (SP100)), at both sides (PML), SUMO1 specific (RanBP2) and SUMO2/3 specific (CoREST1), are shown.

regulate sumoylation depends on the modification pathway. Availability of the different SUMO paralogs when sumoylation is required, or acting on the E1 and E2 enzymes, represents a global way to regulate sumoylation. For instance, stress conditions normally leads to SUMO2/3 conjugation, as SUMO2/3 is freely available in the cell [41]. On the other hand, it has been shown that expression of the Gam1 protein by the CELO adenovirus leads to E1 and E2 degradation and thereby to inhibition of sumoylation [51]. Finally, a more selective way of regulating sumoylation is given through the activity of SUMO ligases and proteases. Thus, localization or spatiotemporal regulation of the expression of these proteins has consequences in target sumoylation.

### 3. SUMO in transcription

#### 3.1. Transcription repression

It is of significance that the sumoylation consensus sequence, before being established to be the site for SUMO attachment, was initially identified as a negative regulatory sequence in several transcription factors [52]. This scenario is exemplified by the transcription factor Elk-1 (Ets (E twenty-six)-like kinase 1), where a repressive domain, the R motif, was identified as an acceptor region for SUMO attachment [53]. Targeting SUMO or the SUMO conjugation enzyme UBC9 to promoters through a Gal4-based system efficiently represses transcription [54-56],



**Figure 2.** Sumoylation and SIMs are involved in complex architecture and function. Schematic representation of some examples for SUMO-SIM interactions involved in recruiting proteins to a particular subcellular localization, in the architecture of PML aggregates and association to Daxx, and in recruitment of different repressor complexes to the chromatin through sumoylated Sp3 for transcription repression.

indicating that sumoylation mainly associates with transcription repression. Examples from different organisms have argued in favor of such a role. A characteristic of silenced genes is that they correlate with low levels of histone acetylation, while active genes usually display high histone acetylation. It has been described in yeast that temperature-sensitive mutation in Ubc9 leads to an increase in global histone acetylation [57]. In addition, in fission yeast, it has been shown that SUMO is required for the maintenance of heterochromatin stability [58]. Early evidence of the involvement of the SUMO pathway in maintenance of the heterochromatin came from *Drosophila*, as a PIAS mutant was identified as a suppressor of position effect variegation, that is, as a mutant releasing heterochromatin-induced gene silencing [59]. A mechanism that clearly account for the repressive role of SUMO is explained by the ability of SUMO to recruit histone deacetylases [56]. For many transcription factors sumoylation has been linked to transcription repression. Additional examples to Elk-1 are NAB proteins [28], c-Jun [60], p53 [61], I $\kappa$ B $\alpha$  [62], C/EBP [63], Sp3 [64] and MEF2 [65]. It is worth noting that in many cases sumoylation turns activators into repressors, as it is the case of p300 and CREB binding protein (CBP) [66, 67]. However, beyond SUMO modification of transcription factors, SUMO association with architecture and function of chromatin-associated repressor complexes is recently getting increased importance. This has been reviewed in [68, 69] and is described below.

### 3.2. Transcription activation

Despite the clear association of SUMO with gene repression, several reports illustrate the involvement of sumoylation in transcription activation. Examples of transcription factors whose activity is stimulated by SUMO are TCF4 [70], GATA4 [71], Pax6 [72], p45 [73], Smad4



[74], Oct4 [75], p53 [76], myocardin [77], PEA3 [78], NFAT1 [79] and HSF1 and 2 [80, 81]. Intriguingly, p53 has been reported both to be activated and repressed by SUMO [76, 82]. Since sumoylation may compete other post-translational modifications, a mechanism proposed for SUMO-mediated activation of transcription consists in avoiding degradation, and thereby in stabilization, of the transcription factor, as it has been proposed for Oct4 [83]. Otherwise, SUMO modification may interfere with association of repressors with the transcription factor, as occurs for Ikaros, whose sumoylation avoid interaction with histone deacetylase complexes [84]. Recently, two publications have brought into consideration the general assumption that SUMO globally associates with transcription repression. It has been reported in yeast that SUMO is detected at all the constitutively transcribed genes tested and in inducible genes upon activation [85]. However, Ubc9 inactivation results in increased transcription of inducible genes, although sumoylation at promoters is reduced, suggesting a role for SUMO in the silencing of inducible genes. In sum, authors conclude that while SUMO associates with repression in some contexts, other properties of SUMO come into play under normal constitutive transcription [85]. More recently, a study performed in HeLa cells has revealed that from G1 to S phase of the cell cycle SUMO1 marks chromatin at the proximal promoter region on many of the most active housekeeping genes [86]. SUMO1 depletion results in reduced expression of these genes. However, this occurs for half of the active genes and the nature of the sumoylated proteins at the promoters remains unknown [86]. Taken together, all these data indicate that although SUMO may intrinsically associate with transcription repression, many other general processes, including constitutive transcription, may also depend on sumoylation, structurally or as a signaling pathway.

## 4. Histone modification and chromatin remodeling

### 4.1. Histone sumoylation

Regarding histone modification, sumoylation has been implicated in both, direct modification of histones and deposition/recognition of other histone marks, such as acetylation and methylation. Histone sumoylation has been demonstrated in both yeast and mammal cells [55, 57]. All core histones and the H2A.Z variant have been shown to be sumoylated in yeast [57, 87], while work on mammal cells has been centered on histone H4 [55]. The N-terminal tail of canonical histones is the target for sumoylation, indicating that sumoylation may interplay with other histone modifications at this region, like acetylation, methylation and phosphorylation. Interfering with the sumoylation pathway significantly reduces the level of histone sumoylation in yeast [57]. Histone sumoylation has been associated with transcription repression. Indeed, mutation of sumoylation sites in histone H2B in yeast leads to increased basal expression of several non-induced genes [57]. A more specific role in Rad51-labeling of persistent DNA double strand breaks has been attributed to sumoylation of the histone variant H2A.Z in yeast [87]. However, which is the real impact of histone sumoylation in transcription in vivo and whether it is a common feature all along the genome need to be clarified.

#### 4.2. Involvement of SUMO in recognition of histone modifications

As explained before, sumoylation of histone tails may affect the way in which different proteins recognize other histone modifications. Conversely, sumoylation of a chromatin-associated factor may modulate its capacity to recognize a specific histone modification. For instance, it has been reported that sumoylation of the bromodomain GTE3 protein, a BET (bromodomain and extra terminal domain) family member, interferes with the capacity of this protein to associate with acetylated histone tails [36]. A surprising link between the sumoylation pathway and recognition of histone modifications is illustrated by a recent and intriguing report describing the capacity of the PHD domain of plant PIAS proteins to directly recognize histone modifications such as methylated Lys4 and Arg2 on histone H3 (methyl-H3K4 and methyl-H3R2) [88].

Polycomb group (PcG) proteins are involved in regulation of gene transcription and chromatin structure especially during development. These transcriptional repressors regulate lineage choice during development and differentiation by establishing long-term heritable gene silencing of relevant genes, for instance *Hox* genes. Thus, they are tightly linked to stem cell biology and cancer [89]. Two main complexes assembling PcG proteins have been described [90]. The polycomb repressive complex 2 (PRC2) contains the histone methyl transferase Enhancer of Zeste (EZH2) and is involved in methylation of H3K27. The PRC1 complex contains the Polycomb protein, which is involved in recognition of the repressive mark trimethyl-H3K27 through a chromodomain. Recruitment of PRC1 to the chromatin results in ubiquitination of histone H2A. Hence, coordinated action of both complexes is involved in the establishment of a compact chromatin structure, which results in gene silencing. One of the mammalian orthologs of *Drosophila* Polycomb is Polycomb-2 (Pc2), which has been shown to display SUMO ligase activity, as previously mentioned [23]. Interestingly, two SIMs have been described in Pc2, one of them has been shown to be relevant for the several functions attributed to Pc2 [91]. Among the SUMO substrates identified for Pc2 are the kinase HIPK2 and the corepressor CtBP1 (see also below), sumoylation of which results in enhanced transcription repression [92-94]. CtBP has been shown to colocalize with Pc2 in nuclear foci called PcG bodies, which contain several PcG proteins [95]. Other Pc2 substrates for sumoylation are ZEB2, DNMT3A and centrin-2 [96, 97]. It has been recently reported that Pc2 mediates sumoylation and recruitment of BMI1 at sites of DNA lesions, linking Pc2 ligase activity with the DNA damage response [98]. Several polycomb subunits have been shown to be sumoylated, for instance SUZ12, EZH2 and YY1, although Pc2 has not been involved in the process [99, 100]. A clear role of sumoylation in PcG proteins-mediated repression came from studies in *C. elegans*. The SOP-2 protein is related to *Drosophila* and vertebrate PRC1-associated PcG proteins Polyhomeotic and Sex combs on midleg (Scm). It has been shown that sumoylation of SOP-2 is required for repression of *Hox* genes in *C. elegans* [101]. Indeed, impaired sumoylation leads to ectopic *Hox* gene expression and homeotic transformations, resulting in a phenotype similar to that provoked by *sop-2* mutations. Additional evidence of SUMO involvement in PcG-mediated repression in vertebrates has been more recently reported. It was previously shown that Pc2 is a target of SUMO [23]. Later, Kang et al demonstrated that sumoylated Pc2 is a target for the SUMO protease SENP2 [102]. In *Senp2* knockout mice,

sumoylated Pc2 accumulates, resulting in increased occupancy at promoters of PcG target genes, such as *Gata4* and *Gata6*. As a result, expression of these genes is reduced during development, which leads to embryonic heart defects among other disorders [102]. Chromatin occupancy by PRC2 subunits and levels of trimethyl-H3K27 seem not to be affected, suggesting that Pc2 sumoylation has a role in recognition of H3K27 methylation, which is released by SENP2.

### 4.3. SUMO-mediated regulation of histone modifications

As previously mentioned, the major impact of sumoylation on histone modification is linked to the role of SUMO in the architecture and function of several chromatin-associated complexes involved in histone modification. Sumoylation by itself may condition the way other histone marks are deposited. However, it has been unambiguously demonstrated that sumoylation is essential for function of a variety of complexes implicated in histone modification, which mostly associate with transcription repression [68, 69]. It has been previously indicated that SUMO is required for the maintenance of constitutive heterochromatin in fission yeast [58]. However, increased evidence of SUMO involvement in the establishment of heterochromatin-like structures in euchromatin loci (facultative heterochromatin) has emerged during the last years. Facultative heterochromatin, besides displaying significant DNA methylation, is characterized by low levels of histone acetylation and histone H3 methylated at Lys4 (H3K4), and high levels of histone H3 methylated at Lys27 (H3K27) and Lys9 (H3K9, di- or tri-methylated), and histone H4 methylated at Lys20 (H4K20, mono-, di- or tri-methylated) [103]. Some of the complexes involved in the establishment of these marks are compiled in Table 3 and described below.

#### 4.3.1. Histone methylation

The histone methyltransferase SETDB1 is involved in tri-methylation of H3K9, a repressive histone mark. The methyl CpG binding protein MBD1 and MCAF1 associate to SETDB1 in a complex, linking DNA methylation to histone methylation. This complex is recruited to the KAP-1 (KRAB associated protein-1) corepressor in a SUMO-dependent manner [26]. In its turn, sumoylated KAP-1 recruits the SETDB1 complex to the chromatin through the zinc finger protein KRAB. This is mediated by a SIM in SETDB1 [26]. In addition, another SIM has been reported in MCAF1, and both MCAF1 and MBD1 are sumoylated [104, 105]. Interestingly, a PHD domain in KAP-1 displays an E3 ligase activity, which promotes intramolecular sumoylation of the adjacent bromodomain [26]. The SETDB1 complex, as explained below, is also recruited to the transcription factor Sp3 in a SUMO dependent manner for transcription repression [106].

Recently, the SUMO ligase PIAS1 has been involved in maintaining an epigenetic repressive state, as studied at the *Foxp3* locus, that restricts differentiation of natural occurring thymus-derived regulatory T cells [107]. Knocking down of *PIAS1* leads to reduced DNA methylation and loss of the repressive mark methyl-H3K9 on the *Foxp3* promoter. A prominent role of PIAS1 in recruitment and association to the DNA methyltransferases DNMT3A and DNMT3B

is also reported. In correlation with loss of H3K9 methylation, HP1 $\gamma$  disappears from the *Foxp3* promoter in the absence of PIAS1 [107].

Complex (subunits)	Activity	Recruiting factor
LSD1/CoREST (LSD1, CoREST, BHC80, HDAC1/2, BRAF35, ZEB1, ZNF217/198)	H3K4 demethylation (LSD1) Histone deacetylation (HDAC1/2)	CtBP1
NurD (CHD3/4, HDAC1/2, RbAp46/48, MTA1/2, MBD3/2)	Nucleosome remodeling (CHD3) Histone deacetylation (HDAC1)	KAP-1
SETDB1 (SETDB1, MBD1, MCAF1)	H3K9 tri-methylation (SETDB1)	KAP-1 Sp3
L3MBTL1 (L3MBTL1, HP1)	Methyl-histone recognition (L3MBTL1)	Sp3
dMEC (dMi-2, dMEP-1)	Nucleosome remodeling (dMi-2)	Sp3
PCR2 (EZH2, EED, SUZ12, RbAp46/48)	H3K27 methylation (EZH2)	various
PCR1 (Pc2, PHC, RNF1/2, SCM1)	trimethyl-H3K27 recognition (Pc2)	–

**Table 3.** SUMO associated repressor complexes. Table summarizes some repressor complexes whose function depends on sumoylation. Examples of different transcription factors involved in recruitment of these complexes are also shown.

#### 4.3.2. Histone demethylation

The histone demethylase LSD1 mediates gene repression by removing methyl groups from mono- or di-methyl-H3K4, which are marks of active transcription [108]. LSD1 works in a corepressor complex together with HDACs and CoREST1 [109, 110]. It has been shown that the LSD1/CoREST complex mediates SUMO-dependent repression of neuronal-specific genes, such as *SCN1A* and *SCN3A*, in non-neuronal cells [111]. Recruitment to the chromatin and repression depends on SUMO2/3 and is mediated by a specific SIM in CoREST. SUMO deconjugation by the SUMO protease SENP3 provokes increased levels of di-methyl-H3K4 and acetyl-H3, which leads to gene activation. Different subunits of the LSD1/CoREST complex have been shown to be sumoylated and/or to contain SIMs (reviewed in [68]). It has been recently shown that sumoylation of the LSD1/CoREST complex subunit BRAF35 controls neuronal differentiation [112]. Overexpression of BRAF35, but not of a sumoylation mutant, strongly impairs neuronal differentiation promoted by neurogenic factors in the vertebrate neural tube. Interestingly, iBRAF, a paralogue of BRAF35 occasionally associated to the LSD1/CoREST complex, is not sumoylated but is able to dimerize with BRAF35, inhibiting BRAF35 sumoylation and binding to the LSD1/CoREST complex. The LSD1/CoREST complex usually



associates with the corepressor CtBP (C-terminal binding protein), which in turn is recruited to the chromatin by a variety of transcription factors [113]. Two CtBPs have been reported in vertebrates, CtBP1 and CtBP2. CtBP1 mediates repression by recruiting a number of repression factors that in addition to LSD1 and HDACs, includes the H3K9 histone methyl-transferase G9a. CtBP1-mediated repression depends on sumoylation [92]. Besides direct interaction of CtBP1 with UBC9, CtBP1 sumoylation is also determined by the SUMO ligases PIAS1, PIAS2 and Pc2 [23, 92, 114]. One of the transcription factors recruiting CtBP1 to the chromatin is the Krüpel-like zinc finger DNA-binding repressor ZEB1, which is also a target for sumoylation [115]. Attachment of SUMO to ZEB1 is required for this factor to display full repression activity [94]. Another zinc finger protein that has been associated with the LSD1/CoREST complex is ZNF198. This factor is both able to be sumoylated and to non-covalently interact with SUMO through a SIM [116-118]. Altogether, these data indicate that SUMO is involved on several functional aspects of the LSD1/CoREST complex: it mediates recruitment of the complex to the chromatin, but also is involved in the architecture of the complex, as different subunits associate to the complex in a sumoylation/SIM-dependent manner.

#### 4.3.3. Histone deacetylation

It has been indicated that repression activity of the LSD1/CoREST complex is in part displayed through HDACs. Indeed, HDAC1 and HDAC2 are components of the LSD1/CoREST complex [110]. Another complex involved in HDAC recruitment to the chromatin is the NuRD (nucleosome remodeling and deacetylation) complex [119]. The core component of the mammalian NuRD complex is the ATP-dependent nucleosome remodeling enzyme CHD3. In addition, this complex includes one or two type I HDACs, histone binding proteins RbAp46 and RbAp48, a methylated DNA-binding protein (MBD2 or MBD3), and members of the MTA and p66 families of proteins [119]. A screening in *Drosophila* cell cultures identified the CHD3 homologue dMi-2, as a factor required for SUMO dependent repression by Sp3 [64]. dMi-2/CHD3 both sumoylates and is able to interact with SUMO-modified transcription factors through a SIM [26, 64]. Thus, CHD3 also interacts with sumoylated KAP-1 [26]. However, it has been demonstrated that phosphorylation of Ser824 in the C terminus of KAP-1, directly impairs interaction between the CHD3 SIM and the SUMO molecule attached to KAP-1 [120]. Therefore, KAP-1 sumoylation is not affected, but recognition of SUMO by the SIM in CHD3. KAP-1 phosphorylation has a role in double-strand break repair, as displaces the chromatin barrier imposed by CHD3-dependent nucleosome-remodeling activity. Additional components of the NuRD complex have been shown to be sumoylated and/or to contain SIMs: MTA1/2, HDAC1, RbAp48 and p66 [111, 121-123]. Interestingly, phenotype of certain vulval mutants in *C. elegans*, which associate with genes coding for NuRD components [124], is quite similar to that of *SUMO* and *UBC9* mutants [125], indicating that function of the NuRD complex is linked to sumoylation.

SUMO directly associates with HDACs in a variety of ways. As previously indicated a well-established link between SUMO and HDACs is illustrated by the SIM-mediated recruitment of HDACs to sumoylated proteins [56, 121]. It was first demonstrated for HDAC2 recruitment to sumoylated Elk-1 [56], and subsequently for HDACs 1, 3, 4, 5 and 6, and



class III SIRT1 deacetylases to several factors. Sumoylation of the coactivator p300 mediates recruitment of class II HDAC6 and class III SIRT1 deacetylases [66, 126]. HDAC1 recruitment to sumoylated Groucho, p68 and reptin has also been described [121, 127, 128]. Moreover, a SUMO-histone H4 fusion has been shown to precipitate HDAC1 [55]. Despite these data, it is not clear at present whether SUMO-dependent recruitment of HDACs involves direct binding of HDAC to SUMO or whether HDACs associate through cofactors recruited in a SUMO-dependent manner, as indicated for the LSD1/CoREST and NuRD complexes. Another example of SUMO-dependent recruitment of HDAC is depicted by the Daxx-mediated recruitment of HDAC2 to sumoylated CBP [67]. In this context, it is worth noting that in a variety of cases, HDAC recruitment does not account for full repression activity mediated by SUMO, as inhibition of HDACs does not relieve SUMO-dependent repression as expected. For instance, it has been shown in a reporter system that repression mediated by a Gal4-SUMO fusion is not sensitive to HDAC inhibition [56], as also occurs for SUMO-dependent Sp3-mediated repression [64, 129]. Despite HDAC2 recruitment by sumoylated Elk-1, HDAC2 knockdown only partially alleviates SUMO-dependent Elk-1-mediated repression [56]. Therefore, histone deacetylases are recruited in a SUMO-dependent manner through repressor complexes, together with additional repressor components, to account for full repression activity of the complex. Conversely, HDAC displacement by target sumoylation has been less reported, but examples have been described. Thus, sumoylation of the Prospero-related homeobox 1 (Prox1) and the de novo DNA methyltransferase DNMT3A disrupts association to HDAC3 and HDAC1/2, respectively [130, 131]. On the other hand, HDACs have also been shown to be substrates for SUMO, which regulates HDAC activity. Then, mutation of the sumoylation sites in HDAC1 has been shown to dramatically reduce its repression activity in a reporter assay [122]. It has been reported that the protease SENP1 is able to remove SUMO from sumoylated HDAC1, which leads to enhanced transcription activity by the androgen receptor [132]. Interestingly, the viral protein Gam1 interferes with HDAC1 sumoylation [133]. The RanBP2 ligase has been demonstrated to promote sumoylation of HDAC4 [134], and a relevant role for SUMO chain formation on HDAC4 has been attributed to the non-covalent interaction between SUMO and UBC9 [135]. Paradoxically, while HDAC1 sumoylation seems to be essential for its repression activity [122], SUMO attachment to HDAC1 impairs association to the CoREST repressor [116]. As previously mentioned, class IIa HDACs have been reported as SUMO ligases. HDAC4 and other class IIa HDACs promote SUMO2/3 attachment to the myocyte enhancer factor 2 family members MEF2D and MEF2C, which leads to repression of target genes [25]. Conversely, ligase activity is inhibited by HDAC4 sumoylation. HDAC4 ligase activity has been also demonstrated on liver X receptors sumoylation by SUMO2/3 [136] and on HIC1 sumoylation by SUMO1 [137], while enhanced sumoylation of PML protein has been attributed to HDAC7 [138].

#### **4.4. Multiple complexes contribute to SUMO-dependent Sp3-mediated repression**

Sp3 belongs to the specificity protein (Sp) family of transcription factors, which regulate multiple genes involved in housekeeping, development and cell cycle. Sp3 is expressed ubiquitously and can act either as an activator or a repressor depending on the promoter

context [106, 139]. Sp3-mediated repression depends on Sp3 sumoylation, and as previously indicated, this repression activity is not affected by HDAC inhibitors [129, 140].

A genome-wide RNAi screen in *Drosophila* cell cultures revealed that multiple complexes were involved in SUMO-dependent repression by Sp3 [64]. Among the genes identified whose knockdown impaired SUMO-dependent transcription repression were genes encoding the ATP-dependent chromatin remodeler dMi-2, the *Drosophila* ortholog of the nematode protein MEP-1 and the polycomb protein Sfmbl. Biochemical analyses indicated that dMi-2, MEP-1 and Sfmbl interacted with each other, bound to SUMO and were recruited to the chromatin in a SUMO-dependent manner. In addition, chromatin immunoprecipitation experiments showed that sumoylated Sp3 recruits a number of heterochromatin associated proteins, including dMi-2, the H3K9 histone methyl transferase (HMT) SETDB1, the H4K20 histone methyl transferase SUV4-20H, heterochromatin protein 1 (HP1)  $\alpha$ ,  $\beta$  and  $\gamma$ , and MBT-domain proteins [141].

It has been previously indicated that dMi-2 is the core component of NuRD complex, a complex with associated HDACs. However, Sp3-SUMO-mediated repression is not sensitive to HDACs inhibitors, indicating either that dMi-2 mediates repression outside the NuRD complex or that there is a redundancy in the mechanisms driving Sp3-SUMO-mediated repression. In fact, several data indicate that dMi-2 is also part of another complex lacking HDAC activity. This complex, dMec, is composed by dMi-2 and the *Drosophila* homolog of the *C. elegans* protein MEP-1 (dMEP-1), and works as a corepressor of proneural genes [142]. Knockdown of dMEP-1 leads to derepression of Sp3 target genes, which is in contrast to functional redundancy among the different repression mechanisms recruited to Sp3 [64]. It is worth noting that MEP-1 was previously shown to contribute to SUMO-dependent repression in *C. elegans* [143]. Thus, sumoylated LIN-1 recruits MEP-1 for repression and inhibition of vulval cell fate. As LIN-1 is homologous to the human Elk-1, it is tempting to speculate that a similar mechanism may account for the SUMO-mediated HDAC-independent repression by Elk-1, despite the absence of a clear MEP-1 homolog in vertebrates.

As formerly mentioned, two HMTs were also recruited to SUMO-Sp3: SETDB1 and SUV4-20H, while the HMT SUV39H1 was not associated [141]. These HMTs were shown to be recruited to the *Dhfr* promoter in a sumoylatable Sp3-dependent manner. Knocking down of *SETDB1* and *SUV4-20H* resulted in reduced trimethylation of H3K9 and H4K20 at the *Dhfr* promoter.

Finally, polycomb protein Sfmbl and the corresponding mammalian orthologs L3MBTL1 and L3MBTL2 also associate to sumoylated Sp3 [64, 141]. These proteins contain repeats of the malignant brain tumor (MBT) domain, which is structurally related to the chromodomain and the Tudor domain, and like these, is able to recognize methylated histones. However, MBTs associate with higher affinity to mono- and di- than to trimethylated histones [144]. It has been shown that L3MBTL1 binds HP1 $\gamma$  and compacts chromatin in a mono- and dimethylated H4K20 and H1bK26-dependent manner [145]. Therefore, this association provides a way to explain L3MBTL1-mediated repression. Binding of HP1 $\alpha$ ,  $\beta$  and  $\gamma$  to Sp3 depends on sumoylation [141]. Sumoylated histone H4 recruits HP1 $\gamma$  [55], and HP1 $\alpha$  has also been shown to preferentially bind sumoylated SP100 [146], suggesting that, as occurs for HDACs, SUMO mediates HP1 recruitment. As Sfmbl, the PRC2-associated PcG protein Scm also contains MBT

repeats. In contrast to Scm, Sfmbt together with Pleiohomeotic, integrates in the polycomb complex PhoRC. Thus, different polycomb complexes include MBT-containing subunits, which might be involved in recognition of mono- and dimethylated histones to facilitate trimethylation by recruiting other subunits with histone methyltransferase activity.

In sum, Sp3 constitutes a paradigm of SUMO-dependent transcription repression through a variety of factors and chromatin-associated complexes. Clear evidence of SUMO involvement in Sp3-mediated repression came from the generation of knock-in mice with a non-sumoylatable version of Sp3 [147]. As Lys residues are targets for other modifications different of sumoylation, for instance acetylation, authors, instead of mutating core Lys551 to Arg changed the acidic residue at the sumoylation site. Interestingly, they substituted Glu by Asp, which abrogated sumoylation, despite for many authors it is assumed the consensus  $\Psi KxE/D$ . Mutation did not affect Sp3 protein levels. However, spermatocyte-specific genes *Dmc1* and *Dnahc8*, and neuronal genes *Paqr6*, *Rims3* and *Robo3* appeared derepressed in non-testicular and extra-neural tissues and in mouse embryonic fibroblasts [147]. This correlated with loss of the repressive heterochromatin marks trimethyl-H3K9 and trimethyl-H4K20 and affected recruitment of repressor proteins, such as HP1, SETDB1, CHD3, and L3MBTL1/2, to the corresponding promoters. Surprisingly, homozygous knock-in mice born at expected mendelian frequency, were fertile and exhibited no obvious phenotype, in contrast to mice lacking Sp3 [148], suggesting that additional mechanisms may control protein expression from the aberrantly induced transcripts.

## 5. Conclusions

Sumoylation results essential for development and growth of all the investigated eukaryotes. In mice, embryos lacking the SUMO conjugating enzyme Ubc9 die at the early postimplantation stage, highlighting the relevance of SUMO conjugation during development [149]. The SUMO pathway is conserved from yeast to human and, together with ubiquitination, appears to be the most utilized pathway in post-translational modifications by UBLs. Despite similar structural features and a common evolutionary origin of SUMO and ubiquitin, they have significantly diverged from a functional point of view. In fact, a complete machinery has evolved around SUMO for specific conjugation/deconjugation of this molecule. Compared with ubiquitin, about 20 N-terminal extra amino acids are present in SUMO, which should account for the different and specific SUMO roles. From the many examples of protein modification by SUMO, structural, regulatory, signaling, and scaffold roles are inferred for this molecule. All these aspects convene to reveal SUMO modification as an important post-translational modification involved in transcription repression. Therefore, SUMO prefigures as an adaptor molecule essential for correct assembly and function of a variety of chromatin-associated repressor complexes. This does not exclude that involvement of SUMO in various systems results in transcriptional activation. A number of SUMO-dependent histone modifications and chromatin remodeling activities have been summarized in this chapter (Table 3). They include, HDACs, HMTs and histone demethylase activities, associated to the NuRD, LSD1/CoREST, SETDB1, dMec, L3MBTL1 and Polycomb complexes, which result in chromatin

compaction and gene silencing. However, many questions remain open. For instance, whether proteins with intrinsic repression activity like HDACs are directly recruited by SUMO or instead, relevant repression activity in vivo results from association of HDACs to repressor complexes recruited in a SUMO-dependent manner, needs to be clarified. In addition, although HDACs have intrinsic repression activity, it has been shown that sumoylation of HDAC1 accounts for its full repression activity [122], raising the question whether SUMO modulates its activity or is recruiting additional repressors. Another intriguing aspect concerns functional redundancy among the different repressors recruited to a locus via SUMO. A number of repressors are recruited to the chromatin through a Gal4-SUMO2 fusion [123], but it has been shown that individually knocking down of these factors has little consequences in SUMO2 displayed repression, which may be explained by functional redundancy of the multiple repressors associated. In a similar way, downregulation of CHD3 (mammalian dMi-2) or L3MBTL1/2 does not impair Sp3-SUMO-mediated repression in vertebrate cells [64, 141]. However, mutation of *dMi-2* or *Sfmbt* in *Drosophila* has a significant impact in Sp3-SUMO-dependent repression [64], suggesting that promoter context and local features account for the level of functional redundancy of SUMO-associated repressors. In addition, an important aspect of the SUMO modification concerns the fleeting nature of the modification in many cases, which means that SUMO-SIM interactions may have permanent consequences despite they are not further detected, a notion that implies a kind of memory and that thereby links SUMO to epigenetics. Interestingly, mutation of the SUMO2 SIM in CoREST is sufficient to abrogate repression of some neuronal specific genes in non-neuronal cells [111], highlighting the relevance of the non-covalent interaction of proteins with SUMO in regulating SUMO-dependent repression. In this context, SIMs and sumoylation sites have been described in many subunits within a repressor complex (reviewed in [68]), which rises the question about how the appropriate connections are established.

## Abbreviations

CBP, CREB binding protein

DeSI-1, desumoylating isopeptidase-1

EZH2, Enhancer of Zeste

HDAC, histone deacetylase

HMT, histone methyl transferase

KAP-1, KRAB associated protein-1

NDSM, negatively charged residues-dependent SUMO motif

Pc2, polycomb-2

PcG, polycomb group

PCR1/2, polycomb repressive complex 1/2

PDSM, phosphorylation-dependent SUMO motif

PIAS, protein inhibitor of activated STAT

PML, promyelocytic leukaemia protein

SEN1, sentrin-specific protease

SIM, SUMO interacting motif

Sp3, specificity protein 3

SP-RING, Siz/PIAS-RING

SUMO, small ubiquitin-like modifier

UBLs, ubiquitin-like proteins

## Acknowledgements

Work in the Garcia-Dominguez laboratory is supported by the Spanish National Ministry of Economy and Competitiveness grant BFU2012-37304/BFI. I thank JC Reyes, P Garcia-Gutierrez and F Juarez-Vicente for critical reading of this chapter.

## Author details

Garcia-Dominguez Mario

Address all correspondence to: [mario.garcia@cabimer.es](mailto:mario.garcia@cabimer.es)

Stem Cells Department. Andalusian Center for Molecular Biology and Regenerative Medicine (CABIMER) & High Council for Scientific Research (CSIC), Seville, Spain

## References

- [1] Hochstrasser M (2009) Origin and function of ubiquitin-like proteins. *Nature* 458: 422-429.
- [2] Burroughs AM, Iyer LM, Aravind L (2012) The natural history of ubiquitin and ubiquitin-related domains. *Frontiers in Bioscience* 17: 1433-1460.
- [3] Mahajan R, Delphin C, Guan T, Gerace L, Melchior F (1997) A small ubiquitin-related polypeptide involved in targeting RanGAP1 to nuclear pore complex protein RanBP2. *Cell* 88: 97-107.



- [4] Matunis MJ, Coutavas E, Blobel G (1996) A novel ubiquitin-like modification modulates the partitioning of the Ran-GTPase-activating protein RanGAP1 between the cytosol and the nuclear pore complex. *The Journal of Cell Biology* 135: 1457-1470.
- [5] Muller S, Hoege C, Pyrowolakis G, Jentsch S (2001) SUMO, ubiquitin's mysterious cousin. *Nature Reviews* 2: 202-210.
- [6] Weissman AM (2001) Themes and variations on ubiquitylation. *Nature Reviews* 2: 169-178.
- [7] Seeler JS, Dejean A (2003) Nuclear and unclear functions of SUMO. *Nature Reviews* 4: 690-699.
- [8] Palancade B, Doye V (2008) Sumoylating and desumoylating enzymes at nuclear pores: underpinning their unexpected duties? *Trends in Cell Biology* 18: 174-183.
- [9] Hay RT (2005) SUMO: a history of modification. *Molecular Cell* 18: 1-12.
- [10] Johnson ES (2004) Protein modification by SUMO. *Annual Review of Biochemistry* 73: 355-382.
- [11] Gill G (2005) Something about SUMO inhibits transcription. *Current Opinion in Genetics & Development* 15: 536-541.
- [12] Lyst MJ, Stancheva I (2007) A role for SUMO modification in transcriptional repression and activation. *Biochemical Society Transactions* 35: 1389-1392.
- [13] Johnson ES, Schwiienhorst I, Dohmen RJ, Blobel G (1997) The ubiquitin-like protein Smt3p is activated for conjugation to other proteins by an Aos1p/Uba2p heterodimer. *The EMBO Journal* 16: 5509-5519.
- [14] Azuma Y, Tan SH, Cavenagh MM, Ainsztein AM, Saitoh H, Dasso M (2001) Expression and regulation of the mammalian SUMO-1 E1 enzyme. *Faseb Journal* 15: 1825-1827.
- [15] Desterro JM, Thomson J, Hay RT (1997) Ubch9 conjugates SUMO but not ubiquitin. *FEBS Letters* 417: 297-300.
- [16] Johnson ES, Blobel G (1997) Ubc9p is the conjugating enzyme for the ubiquitin-like protein Smt3p. *The Journal of Biological Chemistry* 272: 26799-26802.
- [17] Sampson DA, Wang M, Matunis MJ (2001) The small ubiquitin-like modifier-1 (SUMO-1) consensus sequence mediates Ubc9 binding and is essential for SUMO-1 modification. *The Journal of Biological Chemistry* 276: 21664-21669.
- [18] Tatham MH, Chen Y, Hay RT (2003) Role of two residues proximal to the active site of Ubc9 in substrate recognition by the Ubc9.SUMO-1 thiolester complex. *Biochemistry* 42: 3168-3179.

- [19] Tatham MH, Kim S, Yu B, Jaffray E, Song J, Zheng J, Rodriguez MS, Hay RT, Chen Y (2003) Role of an N-terminal site of Ubc9 in SUMO-1, -2, and -3 binding and conjugation. *Biochemistry* 42: 9959-9969.
- [20] Kotaja N, Karvonen U, Janne OA, Palvimo JJ (2002) PIAS proteins modulate transcription factors by functioning as SUMO-1 ligases. *Molecular and Cellular Biology* 22: 5222-5234.
- [21] Sachdev S, Bruhn L, Sieber H, Pichler A, Melchior F, Grosschedl R (2001) PIASy, a nuclear matrix-associated SUMO E3 ligase, represses LEF1 activity by sequestration into nuclear bodies. *Genes & Development* 15: 3088-3103.
- [22] Pichler A, Gast A, Seeler JS, Dejean A, Melchior F (2002) The nucleoporin RanBP2 has SUMO1 E3 ligase activity. *Cell* 108: 109-120.
- [23] Kagey MH, Melhuish TA, Wotton D (2003) The polycomb protein Pc2 is a SUMO E3. *Cell* 113: 127-137.
- [24] Weger S, Hammer E, Heilbronn R (2005) Topors acts as a SUMO-1 E3 ligase for p53 in vitro and in vivo. *FEBS Letters* 579: 5007-5012.
- [25] Gregoire S, Yang XJ (2005) Association with class IIa histone deacetylases upregulates the sumoylation of MEF2 transcription factors. *Molecular and Cellular Biology* 25: 2273-2287.
- [26] Ivanov AV, Peng H, Yurchenko V, Yap KL, Negorev DG, Schultz DC, Psulkowski E, Fredericks WJ, White DE, Maul GG, Sadofsky MJ, Zhou MM, Rauscher FJ, 3rd (2007) PHD domain-mediated E3 ligase activity directs intramolecular sumoylation of an adjacent bromodomain required for gene silencing. *Molecular Cell* 28: 823-837.
- [27] Subramaniam S, Sixt KM, Barrow R, Snyder SH (2009) Rhes, a striatal specific protein, mediates mutant-huntingtin cytotoxicity. *Science* 324: 1327-1330.
- [28] Garcia-Gutierrez P, Juarez-Vicente F, Gallardo-Chamizo F, Charnay P, Garcia-Dominguez M (2011) The transcription factor Krox20 is an E3 ligase that sumoylates its Nab coregulators. *EMBO Reports* 12: 1018-1023.
- [29] Li SJ, Hochstrasser M (1999) A new protease required for cell-cycle progression in yeast. *Nature* 398: 246-251.
- [30] Li SJ, Hochstrasser M (2000) The yeast ULP2 (SMT4) gene encodes a novel protease specific for the ubiquitin-like Smt3 protein. *Molecular and Cellular Biology* 20: 2367-2377.
- [31] Nishida T, Tanaka H, Yasuda H (2000) A novel mammalian Smt3-specific isopeptidase 1 (SMT3IP1) localized in the nucleolus at interphase. *European Journal of Biochemistry / FEBS* 267: 6423-6427.
- [32] Shin EJ, Shin HM, Nam E, Kim WS, Kim JH, Oh BH, Yun Y (2012) DeSUMOylating isopeptidase: a second class of SUMO protease. *EMBO Reports* 13: 339-346.

- [33] Melchior F, Schergaut M, Pichler A (2003) SUMO: ligases, isopeptidases and nuclear pores. *Trends in Biochemical Sciences* 28: 612-618.
- [34] Gareau JR, Lima CD (2010) The SUMO pathway: emerging mechanisms that shape specificity, conjugation and recognition. *Nature Reviews* 11: 861-871.
- [35] Rytinki MM, Kaikkonen S, Pehkonen P, Jaaskelainen T, Palvimo JJ (2009) PIAS proteins: pleiotropic interactors associated with SUMO. *Cellular and Molecular Life Sciences* 66: 3029-3041.
- [36] Garcia-Dominguez M, March-Diaz R, Reyes JC (2008) The PHD domain of plant PIAS proteins mediates sumoylation of bromodomain GTE proteins. *The Journal of Biological Chemistry* 283: 21469-21477.
- [37] Yeh ET (2009) SUMOylation and De-SUMOylation: wrestling with life's processes. *The Journal of Biological Chemistry* 284: 8223-8227.
- [38] Mukhopadhyay D, Dasso M (2007) Modification in reverse: the SUMO proteases. *Trends in Biochemical Sciences* 32: 286-295.
- [39] Gan-Erdene T, Nagamalleswari K, Yin L, Wu K, Pan ZQ, Wilkinson KD (2003) Identification and characterization of DEN1, a deneddylase of the ULP family. *The Journal of Biological Chemistry* 278: 28892-28900.
- [40] Wu K, Yamoah K, Dolios G, Gan-Erdene T, Tan P, Chen A, Lee CG, Wei N, Wilkinson KD, Wang R, Pan ZQ (2003) DEN1 is a dual function protease capable of processing the C terminus of Nedd8 and deconjugating hyper-neddylated CUL1. *The Journal of Biological Chemistry* 278: 28882-28891.
- [41] Saitoh H, Hinchey J (2000) Functional heterogeneity of small ubiquitin-related protein modifiers SUMO-1 versus SUMO-2/3. *The Journal of Biological Chemistry* 275: 6252-6258.
- [42] Bayer P, Arndt A, Metzger S, Mahajan R, Melchior F, Jaenicke R, Becker J (1998) Structure determination of the small ubiquitin-related modifier SUMO-1. *Journal of Molecular Biology* 280: 275-286.
- [43] Tatham MH, Jaffray E, Vaughan OA, Desterro JM, Botting CH, Naismith JH, Hay RT (2001) Polymeric chains of SUMO-2 and SUMO-3 are conjugated to protein substrates by SAE1/SAE2 and Ubc9. *The Journal of Biological Chemistry* 276: 35368-35374.
- [44] Bohren KM, Nadkarni V, Song JH, Gabbay KH, Owerbach D (2004) A M55V polymorphism in a novel SUMO gene (SUMO-4) differentially activates heat shock transcription factors and is associated with susceptibility to type I diabetes mellitus. *The Journal of Biological Chemistry* 279: 27233-27238.
- [45] Owerbach D, McKay EM, Yeh ET, Gabbay KH, Bohren KM (2005) A proline-90 residue unique to SUMO-4 prevents maturation and sumoylation. *Biochemical and Biophysical Research Communications* 337: 517-520.

- [46] Wang CY, She JX (2008) SUMO4 and its role in type 1 diabetes pathogenesis. *Diabetes/Metabolism Research and Reviews* 24: 93-102.
- [47] Zhu J, Zhu S, Guzzo CM, Ellis NA, Sung KS, Choi CY, Matunis MJ (2008) Small ubiquitin-related modifier (SUMO) binding determines substrate recognition and paralogue-selective SUMO modification. *The Journal of Biological Chemistry* 283: 29405-29415.
- [48] Kerscher O (2007) SUMO junction-what's your function? New insights through SUMO-interacting motifs. *EMBO Reports* 8: 550-555.
- [49] Minty A, Dumont X, Kaghad M, Caput D (2000) Covalent modification of p73alpha by SUMO-1. Two-hybrid screening with p73 identifies novel SUMO-1-interacting proteins and a SUMO-1 interaction motif. *The Journal of Biological Chemistry* 275: 36316-36323.
- [50] Bossis G, Melchior F (2006) SUMO: regulating the regulator. *Cell Div* 1: 13.
- [51] Boggio R, Colombo R, Hay RT, Draetta GF, Chiocca S (2004) A mechanism for inhibiting the SUMO pathway. *Molecular Cell* 16: 549-561.
- [52] Iniguez-Lluhi JA, Pearce D (2000) A common motif within the negative regulatory regions of multiple factors inhibits their transcriptional synergy. *Molecular and Cellular Biology* 20: 6040-6050.
- [53] Yang SH, Jaffray E, Hay RT, Sharrocks AD (2003) Dynamic interplay of the SUMO and ERK pathways in regulating Elk-1 transcriptional activity. *Molecular Cell* 12: 63-74.
- [54] Holmstrom S, Van Antwerp ME, Iniguez-Lluhi JA (2003) Direct and distinguishable inhibitory roles for SUMO isoforms in the control of transcriptional synergy. *Proceedings of the National Academy of Sciences USA* 100: 15758-15763.
- [55] Shiio Y, Eisenman RN (2003) Histone sumoylation is associated with transcriptional repression. *Proceedings of the National Academy of Sciences USA* 100: 13225-13230.
- [56] Yang SH, Sharrocks AD (2004) SUMO promotes HDAC-mediated transcriptional repression. *Molecular Cell* 13: 611-617.
- [57] Nathan D, Ingvarsdottir K, Sterner DE, Bylebyl GR, Dokmanovic M, Dorsey JA, Whelan KA, Krsmanovic M, Lane WS, Meluh PB, Johnson ES, Berger SL (2006) Histone sumoylation is a negative regulator in *Saccharomyces cerevisiae* and shows dynamic interplay with positive-acting histone modifications. *Genes & Development* 20: 966-976.
- [58] Shin JA, Choi ES, Kim HS, Ho JC, Watts FZ, Park SD, Jang YK (2005) SUMO modification is involved in the maintenance of heterochromatin stability in fission yeast. *Molecular Cell* 19: 817-828.

- [59] Hari KL, Cook KR, Karpen GH (2001) The *Drosophila* Su(var)2-10 locus regulates chromosome structure and function and encodes a member of the PIAS protein family. *Genes & Development* 15: 1334-1348.
- [60] Muller S, Berger M, Lehembre F, Seeler JS, Haupt Y, Dejean A (2000) c-Jun and p53 activity is modulated by SUMO-1 modification. *The Journal of Biological Chemistry* 275: 13321-13329.
- [61] Wu SY, Chiang CM (2009) Crosstalk between sumoylation and acetylation regulates p53-dependent chromatin transcription and DNA binding. *The EMBO journal*.
- [62] Desterro JM, Rodriguez MS, Hay RT (1998) SUMO-1 modification of IkappaBalpha inhibits NF-kappaB activation. *Molecular Cell* 2: 233-239.
- [63] Kim J, Cantwell CA, Johnson PF, Pfarr CM, Williams SC (2002) Transcriptional activity of CCAAT/enhancer-binding proteins is controlled by a conserved inhibitory domain that is a target for sumoylation. *The Journal of Biological Chemistry* 277: 38037-38044.
- [64] Stielow B, Sapetschnig A, Kruger I, Kunert N, Brehm A, Boutros M, Suske G (2008) Identification of SUMO-dependent chromatin-associated transcriptional repression components by a genome-wide RNAi screen. *Molecular Cell* 29: 742-754.
- [65] Shalizi A, Gaudilliere B, Yuan Z, Stegmuller J, Shirogane T, Ge Q, Tan Y, Schulman B, Harper JW, Bonni A (2006) A calcium-regulated MEF2 sumoylation switch controls postsynaptic differentiation. *Science* 311: 1012-1017.
- [66] Girdwood D, Bumpass D, Vaughan OA, Thain A, Anderson LA, Snowden AW, Garcia-Wilson E, Perkins ND, Hay RT (2003) P300 transcriptional repression is mediated by SUMO modification. *Molecular Cell* 11: 1043-1054.
- [67] Kuo HY, Chang CC, Jeng JC, Hu HM, Lin DY, Maul GG, Kwok RP, Shih HM (2005) SUMO modification negatively modulates the transcriptional activity of CREB-binding protein via the recruitment of Daxx. *Proceedings of the National Academy of Sciences USA* 102: 16973-16978.
- [68] Garcia-Dominguez M, Reyes JC (2009) SUMO association with repressor complexes, emerging routes for transcriptional control. *Biochimica et Biophysica Acta* 1789: 451-459.
- [69] Ouyang J, Gill G (2009) SUMO engages multiple corepressors to regulate chromatin structure and transcription. *Epigenetics* 4: 440-444.
- [70] Yamamoto H, Ihara M, Matsuura Y, Kikuchi A (2003) Sumoylation is involved in beta-catenin-dependent activation of Tcf-4. *The EMBO Journal* 22: 2047-2059.
- [71] Wang J, Feng XH, Schwartz RJ (2004) SUMO-1 modification activated GATA4-dependent cardiogenic gene activity. *The Journal of Biological Chemistry* 279: 49091-49098.
- [72] Yan Q, Gong L, Deng M, Zhang L, Sun S, Liu J, Ma H, Yuan D, Chen PC, Hu X, Liu J, Qin J, Xiao L, Huang XQ, Zhang J, Li DW (2010) Sumoylation activates the transcrip-



tional activity of Pax-6, an important transcription factor for eye and brain development. *Proceedings of the National Academy of Sciences USA* 107: 21034-21039.

- [73] Shyu YC, Lee TL, Ting CY, Wen SC, Hsieh LJ, Li YC, Hwang JL, Lin CC, Shen CK (2005) Sumoylation of p45/NF-E2: nuclear positioning and transcriptional activation of the mammalian beta-like globin gene locus. *Molecular and Cellular Biology* 25: 10365-10378.
- [74] Lin X, Liang M, Liang YY, Brunicardi FC, Feng XH (2003) SUMO-1/Ubc9 promotes nuclear accumulation and metabolic stability of tumor suppressor Smad4. *The Journal of Biological Chemistry* 278: 31043-31048.
- [75] Wei F, Scholer HR, Atchison ML (2007) Sumoylation of Oct4 enhances its stability, DNA binding, and transactivation. *The Journal of Biological Chemistry* 282: 21551-21560.
- [76] Gostissa M, Hengstermann A, Fogal V, Sandy P, Schwarz SE, Scheffner M, Del Sal G (1999) Activation of p53 by conjugation to the ubiquitin-like protein SUMO-1. *The EMBO Journal* 18: 6462-6471.
- [77] Wang J, Li A, Wang Z, Feng X, Olson EN, Schwartz RJ (2007) Myocardin sumoylation transactivates cardiogenic genes in pluripotent 10T1/2 fibroblasts. *Molecular and Cellular Biology* 27: 622-632.
- [78] Guo B, Sharrocks AD (2009) Extracellular signal-regulated kinase mitogen-activated protein kinase signaling initiates a dynamic interplay between sumoylation and ubiquitination to regulate the activity of the transcriptional activator PEA3. *Molecular and Cellular Biology* 29: 3204-3218.
- [79] Terui Y, Saad N, Jia S, McKeon F, Yuan J (2004) Dual role of sumoylation in the nuclear localization and transcriptional activation of NFAT1. *The Journal of Biological Chemistry* 279: 28257-28265.
- [80] Goodson ML, Hong Y, Rogers R, Matunis MJ, Park-Sarge OK, Sarge KD (2001) Sumo-1 modification regulates the DNA binding activity of heat shock transcription factor 2, a promyelocytic leukemia nuclear body associated transcription factor. *The Journal of Biological Chemistry* 276: 18513-18518.
- [81] Hong Y, Rogers R, Matunis MJ, Mayhew CN, Goodson ML, Park-Sarge OK, Sarge KD (2001) Regulation of heat shock transcription factor 1 by stress-induced SUMO-1 modification. *The Journal of Biological Chemistry* 276: 40263-40267.
- [82] Wu SY, Chiang CM (2009) Crosstalk between sumoylation and acetylation regulates p53-dependent chromatin transcription and DNA binding. *The EMBO Journal* 28: 1246-1259.
- [83] Zhang Z, Liao B, Xu M, Jin Y (2007) Post-translational modification of POU domain transcription factor Oct-4 by SUMO-1. *Faseb Journal* 21: 3042-3051.

- [84] Gomez-del Arco P, Koipally J, Georgopoulos K (2005) Ikaros SUMOylation: switching out of repression. *Molecular and Cellular Biology* 25: 2688-2697.
- [85] Rosonina E, Duncan SM, Manley JL (2010) SUMO functions in constitutive transcription and during activation of inducible genes in yeast. *Genes & Development* 24: 1242-1252.
- [86] Liu HW, Zhang J, Heine GF, Arora M, Gulcin Ozer H, Onti-Srinivasan R, Huang K, Parvin JD (2012) Chromatin modification by SUMO-1 stimulates the promoters of translation machinery genes. *Nucleic Acids Research*.
- [87] Kalocsay M, Hiller NJ, Jentsch S (2009) Chromosome-wide Rad51 spreading and SUMO-H2A.Z-dependent chromosome fixation in response to a persistent DNA double-strand break. *Molecular Cell* 33: 335-343.
- [88] Shindo H, Suzuki R, Tsuchiya W, Taichi M, Nishiuchi Y, Yamazaki T PHD finger of the SUMO ligase Siz/PIAS family in rice reveals specific binding for methylated histone H3 at lysine 4 and arginine 2. *FEBS Letters* 586: 1783-1789.
- [89] Bracken AP, Helin K (2009) Polycomb group proteins: navigators of lineage pathways led astray in cancer. *Nature Reviews Cancer* 9: 773-784.
- [90] Lanzuolo C, Orlando V (2012) Memories from the Polycomb Group Proteins. *Annual Review of Genetics* 46: 559-587.
- [91] Yang SH, Sharrocks AD The SUMO E3 ligase activity of Pc2 is coordinated through a SUMO interaction motif. *Molecular and Cellular Biology* 30: 2193-2205.
- [92] Lin X, Sun B, Liang M, Liang YY, Gast A, Hildebrand J, Brunicardi FC, Melchior F, Feng XH (2003) Opposed regulation of corepressor CtBP by SUMOylation and PDZ binding. *Molecular Cell* 11: 1389-1396.
- [93] Roscic A, Moller A, Calzado MA, Renner F, Wimmer VC, Gresko E, Ludi KS, Schmitz ML (2006) Phosphorylation-dependent control of Pc2 SUMO E3 ligase activity by its substrate protein HIPK2. *Molecular Cell* 24: 77-89.
- [94] Wang J, Scully K, Zhu X, Cai L, Zhang J, Prefontaine GG, Krones A, Ohgi KA, Zhu P, Garcia-Bassets I, Liu F, Taylor H, Lozach J, Jayes FL, Korach KS, Glass CK, Fu XD, Rosenfeld MG (2007) Opposing LSD1 complexes function in developmental gene activation and repression programmes. *Nature* 446: 882-887.
- [95] van der Vlag J, Otte AP (1999) Transcriptional repression mediated by the human polycomb-group protein EED involves histone deacetylation. *Nat Genet* 23: 474-478.
- [96] Klein UR, Nigg EA (2009) SUMO-dependent regulation of centrin-2. *Journal of Cell Science* 122: 3312-3321.
- [97] Wotton D, Merrill JC (2007) Pc2 and SUMOylation. *Biochemical Society Transactions* 35: 1401-1404.

- [98] Ismail IH, Gagne JP, Caron MC, McDonald D, Xu Z, Masson JY, Poirier GG, Hendzel MJ CBX4-mediated SUMO modification regulates BMI1 recruitment at sites of DNA damage. *Nucleic Acids Research* 40: 5497-5510.
- [99] Deng Z, Wan M, Sui G (2007) PIASy-mediated sumoylation of Yin Yang 1 depends on their interaction but not the RING finger. *Molecular and Cellular Biology* 27: 3780-3792.
- [100] Riising EM, Boggio R, Chiocca S, Helin K, Pasini D (2008) The polycomb repressive complex 2 is a potential target of SUMO modifications. *PLoS ONE* 3: e2704.
- [101] Zhang H, Smolen GA, Palmer R, Christoforou A, van den Heuvel S, Haber DA (2004) SUMO modification is required for in vivo Hox gene regulation by the *Caenorhabditis elegans* Polycomb group protein SOP-2. *Nature Genetics* 36: 507-511.
- [102] Kang X, Qi Y, Zuo Y, Wang Q, Zou Y, Schwartz RJ, Cheng J, Yeh ET (2010) SUMO-specific protease 2 is essential for suppression of polycomb group protein-mediated gene silencing during embryonic development. *Molecular Cell* 38: 191-201.
- [103] Trojer P, Reinberg D (2007) Facultative heterochromatin: is there a distinctive molecular signature? *Molecular Cell* 28: 1-13.
- [104] Lyst MJ, Nan X, Stancheva I (2006) Regulation of MBD1-mediated transcriptional repression by SUMO and PIAS proteins. *The EMBO Journal* 25: 5317-5328.
- [105] Uchimura Y, Ichimura T, Uwada J, Tachibana T, Sugahara S, Nakao M, Saitoh H (2006) Involvement of SUMO modification in MBD1- and MCAF1-mediated heterochromatin formation. *The Journal of Biological Chemistry* 281: 23180-23190.
- [106] Valin A, Gill G (2007) Regulation of the dual-function transcription factor Sp3 by SUMO. *Biochemical Society Transactions* 35: 1393-1396.
- [107] Liu B, Tahk S, Yee KM, Fan G, Shuai K (2010) The ligase PIAS1 restricts natural regulatory T cell differentiation by epigenetic repression. *Science* 330: 521-525.
- [108] Shi Y, Lan F, Matson C, Mulligan P, Whetstine JR, Cole PA, Casero RA (2004) Histone demethylation mediated by the nuclear amine oxidase homolog LSD1. *Cell* 119: 941-953.
- [109] Lee MG, Wynder C, Cooch N, Shiekhhattar R (2005) An essential role for CoREST in nucleosomal histone 3 lysine 4 demethylation. *Nature* 437: 432-435.
- [110] Shi YJ, Matson C, Lan F, Iwase S, Baba T, Shi Y (2005) Regulation of LSD1 histone demethylase activity by its associated factors. *Molecular Cell* 19: 857-864.
- [111] Ouyang J, Shi Y, Valin A, Xuan Y, Gill G (2009) Direct binding of CoREST1 to SUMO-2/3 contributes to gene-specific repression by the LSD1/CoREST1/HDAC complex. *Molecular Cell* 34: 145-154.
- [112] Ceballos-Chavez M, Rivero S, Garcia-Gutierrez P, Rodriguez-Paredes M, Garcia-Dominguez M, Bhattacharya S, Reyes JC (2012) Control of neuronal differentiation

- by sumoylation of BRAF35, a subunit of the LSD1-CoREST histone demethylase complex. *Proceedings of the National Academy of Sciences USA* 109: 8085-8090.
- [113] Chinnadurai G (2007) Transcriptional regulation by C-terminal binding proteins. *The International Journal of Biochemistry and Cell Biology* 39: 1593-1607.
  - [114] Kagey MH, Melhuish TA, Powers SE, Wotton D (2005) Multiple activities contribute to Pc2 E3 function. *The EMBO Journal* 24: 108-119.
  - [115] Kuppuswamy M, Vijayalingam S, Zhao LJ, Zhou Y, Subramanian T, Ryerse J, Chinnadurai G (2008) Role of the PLDLS-binding cleft region of CtBP1 in recruitment of core and auxiliary components of the corepressor complex. *Molecular and Cellular Biology* 28: 269-281.
  - [116] Gocke CB, Yu H (2008) ZNF198 stabilizes the LSD1-CoREST-HDAC1 complex on chromatin through its MYM-type zinc fingers. *PLoS ONE* 3: e3255.
  - [117] Gocke CB, Yu H, Kang J (2005) Systematic identification and analysis of mammalian small ubiquitin-like modifier substrates. *The Journal of Biological Chemistry* 280: 5004-5012.
  - [118] Kunapuli P, Kasyapa CS, Chin SF, Caldas C, Cowell JK (2006) ZNF198, a zinc finger protein rearranged in myeloproliferative disease, localizes to the PML nuclear bodies and interacts with SUMO-1 and PML. *Experimental Cell Research* 312: 3739-3751.
  - [119] Denslow SA, Wade PA (2007) The human Mi-2/NuRD complex and gene regulation. *Oncogene* 26: 5433-5438.
  - [120] Goodarzi AA, Kurka T, Jeggo PA (2011) KAP-1 phosphorylation regulates CHD3 nucleosome remodeling during the DNA double-strand break response. *Nature Structural & Molecular Biology* 18: 831-839.
  - [121] Ahn JW, Lee YA, Ahn JH, Choi CY (2009) Covalent conjugation of Groucho with SUMO-1 modulates its corepressor activity. *Biochemical and Biophysical Research Communications* 379: 160-165.
  - [122] David G, Neptune MA, DePinho RA (2002) SUMO-1 modification of histone deacetylase 1 (HDAC1) modulates its biological activities. *The Journal of Biological Chemistry* 277: 23658-23663.
  - [123] Rosendorff A, Sakakibara S, Lu S, Kieff E, Xuan Y, DiBacco A, Shi Y, Shi Y, Gill G (2006) NXP-2 association with SUMO-2 depends on lysines required for transcriptional repression. *Proceedings of the National Academy of Sciences USA* 103: 5308-5313.
  - [124] Solari F, Ahringer J (2000) NURD-complex genes antagonise Ras-induced vulval development in *Caenorhabditis elegans*. *Current Biology* 10: 223-226.
  - [125] Poulin G, Dong Y, Fraser AG, Hopper NA, Ahringer J (2005) Chromatin regulation and sumoylation in the inhibition of Ras-induced vulval development in *Caenorhabditis elegans*. *The EMBO Journal* 24: 2613-2623.



- [126] Bouras T, Fu M, Sauve AA, Wang F, Quong AA, Perkins ND, Hay RT, Gu W, Pestell RG (2005) SIRT1 deacetylation and repression of p300 involves lysine residues 1020/1024 within the cell cycle regulatory domain 1. *The Journal of Biological Chemistry* 280: 10264-10276.
- [127] Jacobs AM, Nicol SM, Hislop RG, Jaffray EG, Hay RT, Fuller-Pace FV (2007) SUMO modification of the DEAD box protein p68 modulates its transcriptional activity and promotes its interaction with HDAC1. *Oncogene* 26: 5866-5876.
- [128] Kim JH, Choi HJ, Kim B, Kim MH, Lee JM, Kim IS, Lee MH, Choi SJ, Kim KI, Kim SI, Chung CH, Baek SH (2006) Roles of sumoylation of a reptin chromatin-remodelling complex in cancer metastasis. *Nature Cell Biology* 8: 631-639.
- [129] Ross S, Best JL, Zon LI, Gill G (2002) SUMO-1 modification represses Sp3 transcriptional activation and modulates its subnuclear localization. *Molecular Cell* 10: 831-842.
- [130] Ling Y, Sankpal UT, Robertson AK, McNally JG, Karpova T, Robertson KD (2004) Modification of de novo DNA methyltransferase 3a (Dnmt3a) by SUMO-1 modulates its interaction with histone deacetylases (HDACs) and its capacity to repress transcription. *Nucleic Acids Research* 32: 598-610.
- [131] Shan SF, Wang LF, Zhai JW, Qin Y, Ouyang HF, Kong YY, Liu J, Wang Y, Xie YH (2008) Modulation of transcriptional corepressor activity of prospero-related homeobox protein (Prox1) by SUMO modification. *FEBS Letters* 582: 3723-3728.
- [132] Cheng J, Wang D, Wang Z, Yeh ET (2004) SENP1 enhances androgen receptor-dependent transcription through desumoylation of histone deacetylase 1. *Molecular and Cellular Biology* 24: 6021-6028.
- [133] Colombo R, Boggio R, Seiser C, Draetta GF, Chiocca S (2002) The adenovirus protein Gam1 interferes with sumoylation of histone deacetylase 1. *EMBO Reports* 3: 1062-1068.
- [134] Kirsh O, Seeler JS, Pichler A, Gast A, Muller S, Miska E, Mathieu M, Harel-Bellan A, Kouzarides T, Melchior F, Dejean A (2002) The SUMO E3 ligase RanBP2 promotes modification of the HDAC4 deacetylase. *The EMBO Journal* 21: 2682-2691.
- [135] Knipscheer P, van Dijk WJ, Olsen JV, Mann M, Sixma TK (2007) Noncovalent interaction between Ubc9 and SUMO promotes SUMO chain formation. *The EMBO Journal* 26: 2797-2807.
- [136] Ghisletti S, Huang W, Ogawa S, Pascual G, Lin ME, Willson TM, Rosenfeld MG, Glass CK (2007) Parallel SUMOylation-dependent pathways mediate gene- and signal-specific transrepression by LXRs and PPARgamma. *Molecular Cell* 25: 57-70.
- [137] Stankovic-Valentin N, Deltour S, Seeler J, Pinte S, Vergoten G, Guerardel C, Dejean A, Leprince D (2007) An acetylation/deacetylation-SUMOylation switch through a phylogenetically conserved psiKXEP motif in the tumor suppressor HIC1 regulates transcriptional repression activity. *Molecular and Cellular Biology* 27: 2661-2675.



- [138] Gao C, Ho CC, Reineke E, Lam M, Cheng X, Stanya KJ, Liu Y, Chakraborty S, Shih HM, Kao HY (2008) Histone deacetylase 7 promotes PML sumoylation and is essential for PML nuclear body formation. *Molecular and Cellular Biology* 28: 5658-5667.
- [139] Li L, He S, Sun JM, Davie JR (2004) Gene regulation by Sp1 and Sp3. *Biochemistry and Cell Biology* 82: 460-471.
- [140] Sapetschnig A, Rischitor G, Braun H, Doll A, Schergaut M, Melchior F, Suske G (2002) Transcription factor Sp3 is silenced through SUMO modification by PIAS1. *The EMBO Journal* 21: 5206-5215.
- [141] Stielow B, Sapetschnig A, Wink C, Kruger I, Suske G (2008) SUMO-modified Sp3 represses transcription by provoking local heterochromatic gene silencing. *EMBO Reports* 9: 899-906.
- [142] Kunert N, Wagner E, Murawska M, Klinker H, Kremmer E, Brehm A (2009) dMec: a novel Mi-2 chromatin remodelling complex involved in transcriptional repression. *The EMBO Journal* 28: 533-544.
- [143] Leight ER, Glossip D, Kornfeld K (2005) Sumoylation of LIN-1 promotes transcriptional repression and inhibition of vulval cell fates. *Development* 132: 1047-1056.
- [144] Kim J, Daniel J, Espejo A, Lake A, Krishna M, Xia L, Zhang Y, Bedford MT (2006) Tudor, MBT and chromo domains gauge the degree of lysine methylation. *EMBO Reports* 7: 397-403.
- [145] Trojer P, Li G, Sims RJ, 3rd, Vaquero A, Kalakonda N, Boccuni P, Lee D, Erdjument-Bromage H, Tempst P, Nimer SD, Wang YH, Reinberg D (2007) L3MBTL1, a histone-methylation-dependent chromatin lock. *Cell* 129: 915-928.
- [146] Seeler JS, Marchio A, Losson R, Desterro JM, Hay RT, Chambon P, Dejean A (2001) Common properties of nuclear body protein SP100 and TIF1alpha chromatin factor: role of SUMO modification. *Molecular and Cellular Biology* 21: 3314-3324.
- [147] Stielow B, Kruger I, Diezko R, Finkernagel F, Gillemans N, Kong-a-San J, Philipsen S, Suske G (2010) Epigenetic silencing of spermatocyte-specific and neuronal genes by SUMO modification of the transcription factor Sp3. *PLoS Genetics* 6: e1001203.
- [148] Bouwman P, Gollner H, Elsasser HP, Eckhoff G, Karis A, Grosveld F, Philipsen S, Suske G (2000) Transcription factor Sp3 is essential for post-natal survival and late tooth development. *The EMBO Journal* 19: 655-661.
- [149] Nacerddine K, Lehembre F, Bhaumik M, Artus J, Cohen-Tannoudji M, Babinet C, Pandolfi PP, Dejean A (2005) The SUMO pathway is essential for nuclear integrity and chromosome segregation in mice. *Developmental Cell* 9: 769-779.