

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

186,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Sleep and Cognition in Developmental Age

Marco Carotenuto and Maria Esposito

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/58967>

1. Introduction

Cognitive development is the construction of thought processes, including remembering, problem solving, and decision-making, from childhood to adulthood. Moreover, it refers to how a person perceives, thinks, and gains understanding of his or her world through the interaction of genetic and learned factors. Among the areas of cognitive development are information processing, intelligence, reasoning, language development, and memory.

Historically, the cognitive development of children has been studied in a variety of ways. The oldest is through the Intelligence Quotient (IQ) based on the concept of "mental age" according to which the scores of a child of average intelligence match his or her age, while a gifted child's performance is comparable to that of an older child, and a slow learner's scores are similar to those of a younger child. IQ tests are used worldwide, but they have come under increasing criticism for defining intelligence too narrowly and for being biased with regard to race and gender.

Therefore, the study and knowledge of the various exogenous/environmental factors that could influence the cognitive development could be considered mandatory for the comprehension of the childhood general developing.

2. Sleep and cognitive processes in children

In healthy children, disturbed sleep has been associated with behavioural impairments (e.g., hyperactivity, aggression, anxiety, etc.) [1-7] and reduced neurocognitive performance (e.g., lower IQ, impaired memory, reduced academic performance, reduced attentive ability, etc.) [1,2,6,8].

Among pediatric age, growing evidence suggests the role of sleep habits as disturbing factor for influencing the cognitive development. The sleep disturbance can impact cognition as shown by the negative effects of sleep breathing disorders both in adults [9] and children [10] and as pinpointed by studies on the interaction between specific sleep stages in declarative memory functioning [11] and learning disabilities [12].

Since 1980, [13] more studies have suggested that sleep is associated with IQ levels in children, although the underlying mechanism remains still unknown. Studies involving children and adults have identified a significant relationship between poor or insufficient sleep and decreased cognitive capacity [14-26]. Furthermore, longer habitual sleep duration in healthy school-aged participants has been associated with better performance on measures of perceptual reasoning and overall IQ [23]. These findings reveal an association between sleep duration and performance on IQ tests. Actually, the DSM-5 [27] highlighted the importance of general mental abilities and of the adaptive functioning, beyond the IQ scores for the assessment of individual cognition. In this light, the social and practical domains appear to be more relevant, although the role of sleep cannot be excluded by these aspects, as showed by reports among subjects affected by intellectual disabilities (28-32).

Several Authors have proposed that sleep spindles may physiologically underpin intelligence or high-level general mental ability [33-35]. Sleep spindles are a feature of (predominantly) stage 2 Non-Rapid Eye Movement (NREM) sleep, and are characterized by recurrent and brief bursts of spindle-like EEG activity.

Moreover, spindles may be classified as either slow (10–13 Hz) or fast (13–16 Hz), with different EEG scalp topographies [36], and are both co-active also with hemodynamic responses in different cortical regions [37] and playing a strong role in the reprocessing of previously encoded information [38]. Moreover, Mednick et al in 2013 have been showed as the pharmacological increase induction in spindle occurrence tend to improve memory, providing even stronger support for a mechanistic role of spindles in memory consolidation [39]. Specifically, retention of verbal informations are related to spindles recorded over frontal brain regions [40], while parietal spindles correlate with spatial memory [41].

In general, in humans the maturation of cognitive skills seems to be linked to a subsequent period of slow-wave activity (SWA), that undergoes maturation in parallel with cortical morphology [42,43] and sleep spindles (10–14 Hz) directly involved in synaptic remodeling, leading to alterations in synaptic strength and synchronized neuronal firing [44-47]

On the other hand, sleep spindle frequency in healthy school-age children seems to be negatively associated with performance on the working memory and perceptual reasoning modules of the Wechsler Intelligence Scale for Children-IV (WISC-IV) [48]. Moreover, lower sleep spindle frequency may be associated with better performance on the Intelligence perceptual reasoning and working memory WISC-IV scales, although sleep spindle amplitude, duration and density could be not directly associated with performance on the IQ test. [48]

In general in paediatric age children, sleep disturbances have been often considered as the epiphenomena of an underlying maturational disorder leading to cognitive impairments. However, cortical maturation and sleep-dependent mechanisms of brain plasticity seem to

follow similar developmental trajectories, suggesting closer interactions between these two dynamic processes [42,43,49,50]. In this light, abnormal sleep activity in children might be a causal, or at least a contributing factor in cognitive and learning impairments [51,52].

Specifically, converging evidence demonstrated that sleep plays a critical role in the 'evolution' of memories [53]. In fact, once encoded, sleep-dependent memory processing can not only stabilize memories – a process classically referred to as memory consolidation – but can also enhance them and integrate them into existing memory networks, extracting key elements for retention, abstracting the gist from multi-item memories, discovering the rules governing such collections of item memories, and even modifying them in ways that facilitate the subsequent discovery of creative insights [53].

Conversely, in the declarative memory domain (i.e. the memory for facts and events, usually verbalizable and explicit), beneficial effects of post-learning sleep on performance have been highlighted using mostly verbal (word pairs) and visuospatial (e.g. memory for pictures or objects' location, virtual navigation) hippocampus-dependent learning tasks. For instance, cued recall of learned pairs of words was consistently shown better after post-learning sleep than after a similar period of time spent awake [54-61]. Additionally, sleep may help protecting recently learned memories against retroactive interference due to the acquisition of a novel and related verbal material [62,63]. Therefore, sleep would participate in memory consolidation processes more than by merely protecting novel memories from ongoing, non-specific daytime interferences and memory decay as previously advocated [64]. Likewise in the non-declarative or procedural memory domain, beneficial effects of post-learning sleep have been evidenced for the consolidation of novel skills and habits, although results are more controversial. Sleep-dependent improvements in visual discrimination skills have been consistently demonstrated using the texture discrimination task (TDT) [65-70]. Furthermore in this task, performance deteriorates over repeated practice sessions within a day reflecting the saturation of the underlying neural circuits, unless sleep is allowed between sessions [71].

In this light, performance stabilizes or even increases depending on the duration of the sleep episode and the availability of slow wave sleep (SWS) and rapid eye movement (REM) sleep [72].

A most characteristic electrophysiological feature of non-rapid eye movement (NREM) sleep is the slow oscillation, visible on scalp electroencephalography (EEG) as a biphasic wave of high amplitude and a fundamental frequency of around 1 Hz [73]. This slow oscillation is the result of the alternation of periods of extended synchronization and desynchronization of the membrane potentials of numerous cerebral cortical neurons [74].

During the hyperpolarized phase, often called "down state", neurons remain silent for up to a few hundred milliseconds. During the depolarized phase, also called "up state", neuronal spike activity takes place, often including burst firing [74]. The "up states" seem to be associated with complex and widespread neuronal network activity throughout the brain [75], including high-frequency oscillations. Especially these oscillations, and their coalescence with

slow oscillations, have been implicated in network communication and systems consolidation of memory traces [76-79].

During the up states of slow oscillations, newly encoded memory representations are thought to be reactivated and redistributed, enabling a shift from temporary storage to long-term storage. Crucial for the dynamic formation of neuronal ensembles and altering of the synaptic connections during the up state is the co-occurring thalamo-cortical and cortico-cortical neuronal activity in higher frequency bands, notably the 10–15 Hz sleep spindles [80] and the >30 Hz gamma oscillations [81-83].

Over past decades, it has been evidenced that sleep can contribute to the consolidation of declarative memories in children. How and whether sleep helps in consolidating verbal and non-verbal procedural skills in this population remains a matter of debate and deserves further investigations. Dedicated studies combining comprehensive behavioural measures, neurophysiological and/or neuroimaging recordings in healthy and pathological populations are crucially needed to unravel the mechanisms underlying the evolution of sleep-dependent memory consolidation processes during childhood. Moreover, we could speculate that neurophysiological and neuroimaging investigations may contribute to enlighten the pathophysiological associations linking abnormal sleep patterns, cognitive disturbances and impaired sleep-dependent plasticity processes throughout the developmental phase. These investigations should be conducted in parallel with the study of pathological conditions in which children present abnormal sleep patterns and cognitive deficits, such as, for a few instances, ADHD, specific language impairments and epileptic syndromes. In this framework, comparing the development of sleep-dependent plasticity markers [84] in children with or without cognitive disorders, and how this evolution interacts with cognitive functioning and/or cortical maturation, constitutes a promising field of research to understand the pathophysiological conditions subtending the long-term disruption of cerebral plasticity processes involved in memory consolidation during sleep [84].

Moreover, the well known relationship between sleep and cognition in all ages of life suggests a key role of sleep in cognitive impairment conditions such as mental retardation [85-89], borderline intellectual functioning [32], learning [12,90], memory [91,92] and executive functions disabilities [93-95]. The approaching to the intellectual disabilities could be difficult, particularly in developmental age. In this framework sleep neurophysiology may help the knowledge and comprehension for the functional interrelationships between the cerebral areas.

In general, the decreasing of sleep efficiency and decreased REM ratio were reported as characteristic neurophysiological signs in several developmental disabilities like Down syndrome [96,97], autism [96], Angelman syndrome [98] and in ADHD [99].

Moreover, lower sleep efficiency, higher WASO, increases in NREM sleep EEG (relative) delta and region-dependent decreases in sigma/high frequency activities were reported in subjects with Asperger syndrome [100].

Finally, reduced total sleep time, decreased sleep efficiency percentage, higher WASO, increases in frontally measured NREM sleep EEG delta power and SWS time, as well as region-

dependent decreases in sigma power and reduced REM sleep percent were reported in Williams syndrome [101]. Thus, several papers are reporting similar sleep-EEG alterations in different conditions affecting intellectual functioning.

3. From childhood to adulthood: Differences and similarities in the developmental course

The differences between children and adults are legion, and how they approach and learn from new situations is clearly one of them. Purely psychological studies, ranging from the work of Piaget in the 1950s and 1960s to the ongoing work of Spelke and Carey [102], have focused on the developmental trajectory of learning capacities and the dependence of each incremental improvement on the ones preceding it. Other studies focused on the continuing development of the cerebral cortex as key to changes in learning style and intellectual development [103]. In their recent study, Wilhelm et al. suggest that at least some of the differences in how adults and children process newly acquired information result from age-dependent differences in the forms of sleep-dependent processing applied to such memories [104]. Specifically, their findings suggest that children, 8–11 years of age, show greater sleep-dependent extraction of explicit, or declarative, knowledge of the rules that govern an implicit procedural task than do adults, 18–35 years old [104].

In general, not every memory undergoes all of these forms of sleep-dependent processing, and the mechanisms that determine which ones are employed for a given memory remain poorly understood. [104].

A possible explanation of this age difference in declarative knowledge is found in the structure of children' sleep. Children not only obtained significantly more sleep than the adults (9.8 vs 6.5 hr), but spent more than twice as much of that time in deep, slow wave sleep (SWS; 39% vs 17%; 217 vs 64 min)). [105].

The suggestion that increased SWS in children might lead to better extraction or maintenance of declarative as opposed to non-declarative (e.g., procedural) knowledge has its counterpart in the suggestion found in a recent report [106] that further decreases in SWS with aging might underlie the difficulty to retain new declarative memories experienced by the elderly.

Even childhood naps may be part of this story. Among 15-month-old infants, only those who napped after a learning task retained knowledge of it the next morning [107]. Moreover, they suggested that the developmental changes in sleep architecture, with more naps, SWS, and REM sleep in children than adults, reflects parallel changes in how sleep guides the evolution of memories across the life cycle, in part enhancing explicit fact memory in children, but more abstract knowledge in adults. Perhaps sleep makes children smarter, but adults wiser [105].

The expression of slow waves undergoes remarkable changes during development, both with respect to their topographical distribution [43, 108-110], as well as with respect to their amplitude [111-113]. The amplitude of slow oscillations increases during childhood to peak shortly before puberty [112]. Conversely, a steep drop occurs during adolescence, decelerating

at the age of about 17 years, after which the amplitude declines only slowly [111]. The amplitude of slow oscillations reflects the degree of synchronization by which cortical neurons switch between up and down states [88]. Although receiving much less attention, the capacity of a densely connected neuronal network to synchronize its activity may not only be reflected in the amplitude of slow oscillations, but might as well lead to more pronounced oscillations in frequency bands other than the 0.5–4 Hz range. Indeed, power in the theta (4–8 Hz) range declines across puberty and early adolescence [113]. Gaudreau et al. [114] investigated NREM sleep EEG power in a wider range of frequency bands across the age range of 6 to 60 years. They report a much higher absolute power of theta (4.0–7.75 Hz), alpha (8.0–12.0 Hz) and beta (15.25–31.0 Hz) in the group of children in the range of 6 to 10 years, as compared to the groups of adolescents (range 14 to 16 years), young adults (range 19 to 29 years) and middle aged adults (range 36 to 60 years). The largest values for spindle-range power (12.25–15.0 Hz) were found in the adolescent group, suggestive of an inverted-U shape peaking somewhere between the age of about 10 years and late adolescence. Jenni and Carskadon [115] investigated developmental changes across the 0.6 to 25 Hz NREM-sleep power spectrum and found that children aged 9.6–12.9 years, as compared to children aged 11.8–15.9 years, had significantly higher absolute power not only in the low frequencies up to about 7 Hz, but also in the 12–13 Hz sigma range and 16–17 Hz low beta range. Recently, both Tarokh et al. [109, 110, 116] and Baker et al. [117] applied within-subject follow-up design rather than the above-mentioned cross-sectional approaches, to confirm that changes in the sleep EEG across adolescence were not restricted to the lower frequency bands, neither to NREM sleep only. Across adolescence, the sleep EEG power decreases over a wide range of frequencies, up to the beta range for at least some derivations. In summary, the above mentioned developmental studies suggest that a wide range of cortical oscillations measured in the scalp EEG show their maximal signal-to-noise ratio in late childhood, around the age of 11, where the signal of interest is the amplitude of the oscillations and the noise reflects the noise floor of scalp EEG assessment [118]. In contrast the gamma power increased on the rising slope and positive peak of the slow wave, with strongly suppression of both gamma and spindle activities during the negative peak, independently by external stimulation (i.e. acoustic) [118].

Recently, the topographic distribution of slow wave activity (SWA; EEG power between 0.75 and 4.5 Hz) during non-rapid eye movement (NREM) sleep was proposed to parallel cortical maturation from childhood through adolescence [43]. High density sleep EEG recordings in children and adolescents between 2 and 20 years of age showed that SWA exhibits a regional, age specific predominance with a developmental shift from occipital to frontal regions reaching frontal derivations only during adolescence. Strikingly, the local SWA maxima paralleled the time course of cortical gray matter [119, 120] and behavioural maturation [121] indicating that SWA may be a marker of brain maturation. This interpretation seems to be in line with the increasing number of reports showing a direct relationship between sleep slow waves and plastic cortical processes [122, 123]. More specifically, it has been hypothesized that wakefulness is associated with a significant increasing in synaptic strength, which is homeostatically rebalanced during sleep. This hypothesis was confirmed in various species examining markers of synaptic strength. A close relationship between SWA and cortical synapses has been proposed early on [124]. Although direct evidence is lacking, recent findings from animal

studies or humans using current in vivo measures for cortical structure and activity support the suggestion that synaptic strength is reflected in deep sleep slow waves [125]. Several studies have shown that slow wave characteristics (SWA, topography, slope, amplitude) are closely related to maturational alterations in the cortex [42,43,49,50]. Moreover, slow waves represent synchronized activity among cortical neurons, as shown by multiunit recordings in the rat [123]. Thus, neurons show synchronized activity, the larger is the amplitude of slow waves displayed by this network. Increased synchronization is achieved by stronger synaptic connections and/or a denser network (i.e. more connections) [125].

In general it has been stated that the SWA is related to cortical plasticity [123, 126-130] (e.g., a change in strength and/or number of synapses) not only occurs because of learning processes, but also in the course of brain maturation. In fact, it was shown that SWA is not equally distributed across the scalp in children and adolescents, but exhibits local age-specific maxima [43]. Furthermore, the location of maximal SWA seems to parallel the time course of cortical maturation along the posteroanterior axis [131]. Thus, the topography of SWA may reflect cortical plasticity during development [125].

Regarding the effect of sleep deprivation, children demonstrate difficult behaviors when sleep deprived that can be stressful and impact quality of life for the entire family. Connecting sleep problems with daytime behavioral challenges may not be intuitive to parents. Adults manifest different symptoms when sleep deprived such as daytime sleepiness, psychomotor slowing and impairments in cognitive processing and memory [132]. In comparison, sleep deprivation in children is more likely to be associated with a range of emotional/behavioral disturbances, including problematic behaviors [133], attention problems [134-136], anxiety/depression [137], and hyperactivity [138, 139]. Brain maturation is a complex process [140] that begins prenatally with neural proliferation and migration and synapse formation continuing till two years of age. Myelination is an important process that begins prenatally as well but continues into adolescence with different systems myelinating at different times. The determinants of neurodevelopment and behavior rely on complex neural circuits that connect neural substrates to serve a specific function. The development of these neural circuits is still a mystery and influenced by genetic, sociocultural, medical and environmental factors [141]. The neuroanatomic substrates involved in neurobehavioral functioning span cortical, subcortical and brainstem regions and formulate complex networks which include the prefrontal cortex, amygdala and striatum. Executive functioning is highly localized to the prefrontal cortex. The amygdala is of great importance to emotional reactivity and affect and striatum to reward seeking behavior. Neuroimaging techniques reveal complex patterns of neuroanatomical functioning during specific sleep stages.

During NREM slow wave sleep, the brainstem, thalamus, basal ganglia, and prefrontal and temporal lobe regions all appear to undergo reduced activity [142]. In REM sleep, significant levels of activity are reported in the pontine tegmentum, thalamic nuclei, occipital cortex, mediobasal prefrontal lobes together with affect related regions including the amygdala, hippocampus, and anterior cingulate cortex [142]. The prefrontal cortex is relatively inactive all through sleep in contrast to its high activity during waking states [143]. This inactivity is reflected by the high voltage and slow brain wave oscillations in NREM sleep in the frontal

lobes, relative to other cortical regions, suggesting that the thalamocortical input is disabled and a lower level of metabolism in the frontal lobes during NREM sleep stages is present [144]. Several investigators have suggested that sleep is particularly important for restoring prefrontal cortical activity [145-148] however, this restorative process remains poorly understood. Neuroimaging studies showed profound effects of one night's total sleep deprivation on the blood flow to prefrontal areas which correspond to the deteriorations in daytime prefrontal task performance [149,150].

On the other hand, sleep deprivation also impacts neural circuitry underlying regulation of emotions, impulsivity and reward seeking behavior. Sleep deprived adult volunteers viewing emotional images have increased activation of the amygdala on functional neuroimaging yet weaker connection between the prefrontal cortex and the amygdala [151]. This scenario allows for uncontrolled, increased emotional response. Likewise, neurocognitive functions that involve the striatum and basal ganglia such as risk avoidance and responsiveness to rewards are also impacted by sleep deprivation. For instance, sleep deprived adults take greater risks and are less concerned about consequences of their behavior [152]. Such findings have also been noted in adolescents aged 11–13 years using functional magnetic resonance imaging (fMRI) and a guessing task with monetary rewards [153]. During reward anticipation, less activation in the caudate nucleus (part of the ventral striatum) was associated with reduced sleep time, later sleep onset time, and lower self-reported sleep quality. During reward outcome, less caudate activation was seen with later sleep onset time, earlier sleep offset time, and lower sleep quality. This findings suggested that sleep deprivation could contribute to low reactivity in reward-related brain areas in adolescents and may lead to compensatory increases in reward-driven behavior. Such findings have significant public health implications when one considers that reward seeking behaviors are associated with depressive symptoms, sensation seeking, and substance abuse in adolescents [137, 154].

4. Conclusions

The relationship between sleep and cognition is intriguing and not yet well understood.

Investigation into sleep habits in the young and the neurophysiological study of sleep (e.g. sleep macrostructure, microstructure, power spectra and CAP) may be considered as mandatory in the future for a better knowledge and comprehension of cognition development.

Author details

Marco Carotenuto* and Maria Esposito

*Address all correspondence to: marco.carotenuto@unina2.it

Sleep Clinic for Developmental Age, Department of Mental Health, Physical and Preventive Medicine, Second University of Naples, Italy

References

- [1] Buckhalt, J.A., El-Sheikh, M., Keller, P.S., Kelly, R.J., 2009. Concurrent and longitudinal relations between children's sleep and cognitive functioning: the moderating role of parent education. *Child Development* 80, 875–892
- [2] Hiscock, H., Canterford, L., Ukomunne, O.C., Wake, M., 2007. Adverse associations of sleep problems in Australian preschoolers: national population study. *Pediatrics* 119, 86–93.
- [3] Huang, Y.S., Chen, N.H., Li, H.Y., Wu, Y.Y., Chao, C.C., Guilleminault, C., 2004. Sleep disorders in Taiwanese children with attention deficit/hyperactivity disorder. *Journal of Sleep Research* 13, 269–277.
- [4] O'Brien, L.M., Ivanenko, A., Crabtree, V.M., Holbrook, C.R., Bruner, J.L., Klaus, C.J., Gozal, D., 2003. Sleep disturbances in children with attention deficit hyperactivity disorder. *Pediatric Research* 54, 237–243.
- [5] Owens, J.A., Maxim, R., Nobile, C., McGuinn, M., Msall, M., 2000. Parental and self-report of sleep in children with attention-deficit/hyperactivity disorder. *Archives of Pediatrics & Adolescent Medicine* 154, 549–555
- [6] Sadeh, A., Gruber, R., Raviv, A., 2002. Sleep, neurobehavioural functioning, and behaviour problems in school-age children. *Child Development* 73, 403–417
- [7] Willoughby, M.T., Angold, A., Egger, H.L., 2008. Parent-reported attention-deficit/hyperactivity disorder symptomatology and sleep problems in a preschool-age pediatric clinic sample. *Journal of the American Academy of Child and Adolescent Psychiatry* 47, 1086–1094
- [8] Randazzo, A.C., Muehlbach, M.J., Schweitzer, P.K., Walsh, J.K., 1998. Cognitive function following acute sleep restriction in children ages 10–14. *Sleep* 21, 861–868.
- [9] Verstraeten E, Neurocognitive effects of obstructive sleep apnea syndrome, *Curr. Neurol. Neurosci. Rep.* 7 (2007) 161–166
- [10] Kheirandish L, Gozal D, Neurocognitive dysfunction in children with sleep disorders, *Dev. Sci.* 9 (4) (2006) 388–399
- [11] Backhaus J, Junghanns K, Born J, Hohaus K, Faasch F, Hohagen F, Impaired declarative memory consolidation during sleep in patients with primary insomnia: influence of sleep architecture and nocturnal cortisol release, *Biol. Psychiatry* 60 (2006) 1324–1330
- [12] Bruni O, Ferri R, Novelli L, Finotti E, Terribili M, Troianiello M, Valente D, Sabatello U, Curatolo P. Slow EEG amplitude oscillations during NREM sleep and reading disabilities in children with dyslexia. *Dev Neuropsychol.* 2009;34(5):539-51. doi: 10.1080/87565640903133418

- [13] Borrow SJ, Adam K, Chapman K, Oswald I, Hudson L, Idzikowski CJ. REM sleep and normal intelligence. *Biol Psychiatry*. 1980 Feb;15(1):165-9
- [14] Beebe, D.W., 2011. Cognitive, behavioral, and functional consequences of inadequate sleep in children and adolescents. *Pediatric Clinics of North America* 58, 649–665
- [15] Bell-McGinty, S., Habeck, C., Hilton, H., Rakitin, B., Scarmeas, N., Zarahn, E., et al., 2004. Identification and differential vulnerability of a neural network in sleep deprivation. *Cerebral Cortex* 14, 496–502
- [16] Busby, K., Pivik, R.T., 1983. Sleep patterns in children of superior intelligence. *Journal of Child Psychology and Psychiatry* 24, 587–600
- [17] Caldwell, J.A., Mu, Q., Smith, J.K., Mishory, A., Caldwell, J.L., Peters, G., et al., 2005. Are individual differences in fatigue vulnerability related to baseline differences in cortical activation? *Behavioral Neuroscience* 119, 694–707
- [18] Chee, M.W., Choo, W.C., 2004. Functional imaging of working memory after 24 h of total sleep deprivation. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience* 24, 4560–4567
- [19] Chee, M.W., Chuah, L.Y., Venkatraman, V., Chan, W.Y., Philip, P., Dinges, D.F., 2006. Functional imaging of working memory following normal sleep and after 24 and 35 h of sleep deprivation: correlations of fronto-parietal activation with performance. *NeuroImage* 31, 419–428
- [20] Choo, W.C., Lee, W.W., Venkatraman, V., Sheu, F.S., Chee, M.W., 2005. Dissociation of cortical regions modulated by both working memory load and sleep deprivation and by sleep deprivation alone. *NeuroImage* 25, 579–587
- [21] Dewald, J.F., Meijer, A.M., Oort, F.J., Kerkhof, G.A., Bogels, S.M., 2010. The influence of sleep quality, sleep duration and sleepiness on school performance in children and adolescents: a meta-analytic review. *Sleep Medicine Reviews* 14, 179–189
- [22] Gozal, D., Kheirandish-Gozal, L., 2007. Neurocognitive and behavioral morbidity in children with sleep disorders. *Current Opinion in Pulmonary Medicine* 13, 505–509
- [23] Gruber, R., Laviolette, R., Deluca, P., Monson, E., Cornish, K., Carrier, J., 2010. Short sleep duration is associated with poor performance on IQ measures in healthy school age children. *Sleep Medicine* 11, 289–294
- [24] Habeck, C., Rakitin, B.C., Moeller, J., Scarmeas, N., Zarahn, E., Brown, T., et al., 2004. An event-related fMRI study of the neurobehavioral impact of sleep deprivation on performance of a delayed-match-to-sample task. *Brain Research. Cognitive Brain Research* 18, 306–321
- [25] Voderholzer, U., Piosczyk, H., Holz, J., Landmann, N., Feige, B., Loessl, B., 2011. Sleep restriction over several days does not affect long-term recall of declarative and procedural memories in adolescents. *Sleep Medicine* 12, 170–178

- [26] Wilhelm, I., Prehn-Kristensen, A., Born, J., 2012. Sleep-dependent memory consolidation—what can be learnt from children? *Neuroscience and Biobehavioral Reviews* 36, 1718–1728
- [27] American Psychiatric Association: Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition. Arlington, VA, American Psychiatric Association, 2013
- [28] Mason TB, Arens R, Sharman J, Bintliff-Janisak B, Schultz B, Walters AS, Cater JR, Kaplan P, Pack AI. Sleep in children with Williams Syndrome. *Sleep Med.* 2011 Oct; 12(9):892-7. doi: 10.1016/j.sleep.2011.05.003
- [29] Annaz D, Hill CM, Ashworth A, Holley S, Karmiloff-Smith A. Characterisation of sleep problems in children with Williams syndrome. *Res Dev Disabil.* 2011 Jan-Feb; 32(1):164-9. doi: 10.1016/j.ridd.2010.09.008
- [30] Miano S, Bruni O, Aricò D, Elia M, Ferri R. Polysomnographic assessment of sleep disturbances in children with developmental disabilities and seizures. *Neurol Sci.* 2010 Oct;31(5):575-83. doi: 10.1007/s10072-010-0291-8
- [31] Kronk R, Bishop EE, Raspa M, Bickel JO, Mandel DA, Bailey DB Jr. Prevalence, nature, and correlates of sleep problems among children with fragile X syndrome based on a large scale parent survey. *Sleep.* 2010 May;33(5):679-87
- [32] Esposito M, Carotenuto M. Borderline intellectual functioning and sleep: the role of cyclic alternating pattern. *Neurosci Lett.* 2010 Nov 19;485(2):89-93. doi:10.1016/j.neulet.2010.08.062.
- [33] Bodizs, R., Kis, T. Lazar, A.S., Havran, L., Rigo, P., Clemens, Z., et al., 2005. Prediction of general mental ability based on neural oscillation measures of sleep. *Journal of Sleep Research* 14, 285–292
- [34] Brière, M., Forest, G., Lussier, I., Godbout, G., 2000. Implicit verbal recall correlates positively with EEG sleep spindle activity. *Sleep* 23, A219
- [35] Fogel, S.M., Nader, R., Cote, K.A., Smith, C.T., 2007. Sleep spindles and learning potential. *Behavioral and Neurosciences* 121, 1–10
- [36] Zeitlhofer, J., Gruber, G., Anderer, P., Asenbaum, S., Schimicek, P., Saletu, B., 1997. Topographic distribution of sleep spindles in young healthy subjects. *J. Sleep Res.* 6 (3), 149–155
- [37] Schabus, M., Dang-Vu, T.T., Albouy, G., Balteau, E., Boly, M., Carrier, J., Darsaud, A., Degueldre, C., Desseilles, M., Gais, S., et al., 2007. Hemodynamic cerebral correlates of sleep spindles during human non-rapid eye movement sleep. *Proc. Natl. Acad. Sci. U. S. A.* 104 (32), 13164–13169
- [38] Cox R, Hofman WF, de Boer M, Talamini LM. Local sleep spindle modulations in relation to specific memory cues. *NeuroImage* 99 (2014) 103–110. <http://dx.doi.org/10.1016/j.neuroimage.2014.05.028>

- [39] Mednick, S.C., McDevitt, E.A., Walsh, J.K., Wamsley, E., Paulus, M., Kanady, J.C., Drummond, S.P., 2013. The critical role of sleep spindles in hippocampal-dependent memory: a pharmacology study. *J. Neurosci.* 33 (10), 4494–4504
- [40] Clemens, Z., Fabó, D., Halász, P., 2005. Overnight verbal memory retention correlates with the number of sleep spindles. *Neuroscience* 132 (2), 529–535
- [41] Clemens, Z., Fabó, D., Halász, P., 2006. Twenty-four hours retention of visuospatial memory correlates with the number of parietal sleep spindles. *Neurosci. Lett.* 403 (1), 52–56
- [42] Buchmann, A.; Ringli, M.; Kurth, S.; Schaerer, M.; Geiger, A.; Jenni, O.G.; Huber, R. EEG sleep slow-wave activity as a mirror of cortical maturation. *Cereb. Cortex* 2011, 21, 607–615
- [43] Kurth, S.; Ringli, M.; Geiger, A.; LeBourgeois, M.; Jenni, O.G.; Huber, R. Mapping of cortical activity in the first two decades of life: A high-density sleep electroencephalogram study. *J. Neurosci.* 2010, 30, 13211–13219
- [44] Lante, F.; Toledo-Salas, J.C.; Ondrejcek, T.; Rowan, M.J.; Ulrich, D. Removal of synaptic Ca^{2+} -permeable AMPA receptors during sleep. *J. Neurosci.* 2011, 31, 3953–3961
- [45] Birtoli, B.; Ulrich, D. Firing mode-dependent synaptic plasticity in rat neocortical pyramidal neurons. *J. Neurosci.* 2004, 24, 4935–4940
- [46] Vyazovskiy, V.V.; Harris, K.D. Sleep and the single neuron: The role of global slow oscillations in individual cell rest. *Nat. Rev. Neurosci.* 2013, 14, 443–451
- [47] Kurth S, Achermann P, Rusterholz T, LeBourgeois MK. Development of Brain EEG Connectivity across Early Childhood: Does Sleep Play a Role? *Brain Sci.* 2013, 3, 1445–1460; doi:10.3390/brainsci3041445
- [48] Gruber, R., et al., The association between sleep spindles and IQ in healthy school-age children, *International Journal of Psychophysiology* (2013), <http://dx.doi.org/10.1016/j.ijpsycho.2013.03.018>
- [49] Kurth, S., Jenni, O.G., Riedner, B.A., Tononi, G., Carskadon, M.A., Huber, R., 2010. Characteristics of sleep slow-waves in children and adolescents. *Sleep* 33, 475–480
- [50] Kurth, S., Ringli, M., Lebourgeois, M.K., Geiger, A., Buchmann, A., Jenni, O.G., Huber, R., 2012. Mapping the electrophysiological marker of sleep depth reveals skill maturation in children and adolescents. *NeuroImage* 63, 959–965
- [51] Chan, S., Baldeweg, T., Cross, J.H., 2011. A role for sleep disruption in cognitive impairment in children with epilepsy. *Epilepsy & Behavior* 20, 435–440
- [52] Van Bogaert, P., Urbain, C., Galer, S., Ligot, N., Peigneux, P., De Tiège, X., 2011. Impact of focal interictal epileptiform discharges on behaviour and cognition in children. *Neurophysiologie Clinique/Clinical Neurophysiology* 42, 53–58

- [53] Stickgold, R. and Walker, M.P. (2013) Sleep-dependent memory triage: evolving generalization through selective processing. *Nat. Neurosci.* 16, 139–145
- [54] Gais, S., Born, J., 2004. Declarative memory consolidation: mechanisms acting during human sleep. *Learning & Memory* (Cold Spring Harbor, N.Y.) 11, 679–685
- [55] Gais, S., Lucas, B., Born, J., 2006. Sleep after learning aids memory recall. *Learning & Memory* 13, 259–262
- [56] Wilhelm, I., Diekelmann, S., Born, J., 2008. Sleep in children improves memory performance on declarative but not procedural tasks. *Learning & Memory* 15, 373–377
- [57] Diekelmann, S., Wilhelm, I., Born, J., 2009. The whats and whens of sleep-dependent memory consolidation. *Sleep Medicine Reviews* 13, 309–321
- [58] Diekelmann, S., Buchel, C., Born, J., Rasch, B., 2011. Labile or stable: opposing consequences for memory when reactivated during waking and sleep. *Nature Neuroscience* 14, 381–386
- [59] Peigneux, P., Laureys, S., Fuchs, S., Destrebecqz, A., Collette, F., Delbeuck, X., Phillips, C., Aerts, J., Del Fiore, G., Degueldre, C., Luxen, A., Cleeremans, A., Maquet, P., 2003. Learned material content and acquisition level modulate cerebral reactivation during post training rapid-eye-movements sleep. *NeuroImage* 20, 125–134
- [60] Peigneux, P., Laureys, S., Fuchs, S., Collette, F., Perrin, F., Reggers, J., Phillips, C., Degueldre, C., Del Fiore, G., Aerts, J., Luxen, A., Maquet, P., 2004 Are spatial memories strengthened in the human hippocampus during slow wave sleep? *Neuron* 44, 535–545
- [61] Peigneux, P., Schmitz, R., Urbain, C., 2010. Sleep and forgetting. In: Della Sala, S. (Ed.), *Forgetting*. Psychology Press, Hove, UK
- [62] Ellenbogen, J.M., Hulbert, J.C., Stickgold, R., Dinges, D.F., Thompson-Schill, S.L., 2006. Interfering with theories of sleep and memory: sleep, declarative memory, and associative interference. *Current Biology* 16, 1290–1294
- [63] Ellenbogen, J.M., Hulbert, J.C., Jiang, Y., Stickgold, R., 2009. The sleeping brain's influence on verbal memory: boosting resistance to interference. *PLoS One* 4, e4117
- [64] Wixted, J.T., 2004. The psychology and neuroscience of forgetting. *Annual Review of Psychology* 55, 235–269
- [65] Gais, S., Plihal, W., Wagner, U., Born, J., 2000. Early sleep triggers memory for early visual discrimination skills. *Nature Neuroscience* 3, 1335–1139
- [66] Gais, S., Rasch, B., Wagner, U., Born, J., 2008. Visual-procedural memory consolidation during sleep blocked by glutamatergic receptor antagonists. *Journal of Neuroscience* 28, 5513–5518

- [67] Karni, A., Tanne, D., Rubenstein, S., Askenasy, J.J.M., Sagi, D., 1994. Dependence on REM sleep of overnight improvement of a perceptual skill. *Science* 265, 679–682
- [68] Karni, A., Meyer, G., Rey-Hipolito, C., Jezzard, P., Adams, M.M., Turner, R., Ungerleider, L.G., 1998. The acquisition of skilled motor performance: fast and slow experience-driven changes in primary motor cortex. *Proceedings of the National Academy of Sciences of the United States of America* 95, 861–868
- [69] Stickgold, R., 2005. Sleep-dependent memory consolidation. *Nature* 437, 1272–1278
- [70] Stickgold, R., Whidbee, D., Schirmer, B., Patel, V., Hobson, J.A., 2000. Visual discrimination task improvement: a multi-step process occurring during sleep. *Journal of Cognitive Neuroscience* 12, 246–254
- [71] Mednick, S.C., Nakayama, K., Cantero, J.L., Atienza, M., Levin, A.A., Pathak, N., Stickgold, R., 2002. The restorative effect of naps on perceptual deterioration. *Nature Neuroscience* 5, 677–681
- [72] Mednick, S.C., Nakayama, K., Stickgold, R., 2003. Sleep-dependent learning: a nap is as good as a night. *Nature Neuroscience* 6, 697–698
- [73] Achermann, P., Borbély, A.A., 1997. Low-frequency (b1 Hz) oscillations in the human sleep electroencephalogram. *Neuroscience* 81, 213–222
- [74] Steriade, M., Nuñez, A., Amzica, F., 1993. A novel slow (b1 Hz) oscillation of neocortical neurons in vivo: depolarizing and hyperpolarizing components. *Journal of Neuroscience* 13, 3252–3265
- [75] Volgushev, M., Chauvette, S., Timofeev, I., 2011. Long-range correlation of the membrane potential in neocortical neurons during slow oscillation. *Progress in Brain Research* 193, 181–199
- [76] Diekelmann, S., Born, J., 2010. The memory function of sleep. *Nature Reviews. Neuroscience* 11, 114–126
- [77] Mölle, M., Born, J., 2011. Slow oscillations orchestrating fast oscillations and memory consolidation. *Progress in Brain Research* 193, 93–110
- [78] Schwindel, C.D., McNoughton, B.L., 2011. Hippocampal–cortical interactions and the dynamics of memory trace reactivation. *Progress in Brain Research* 193, 163–177
- [79] Van Someren, E.J.W., Van Der Werf, Y.D., Roelfsema, P.R., Mansvelder, H.D., Lopes da Silva, F.H., Da Silva, F.H.L., 2011. Slow brain oscillations of sleep, resting state, and vigilance. *Progress in Brain Research* 193, 3–15
- [80] Rosanova, M., Ulrich, D., 2005. Pattern-specific associative long-term potentiation induced by a sleep spindle-related spike train. *Journal of Neuroscience* 25, 9398–9405

- [81] Steriade, M., Amzica, F., Contreras, D., 1996. Synchronization of fast (30–40 Hz) spontaneous cortical rhythms during brain activation. *Journal of Neuroscience* 16, 392–417
- [82] Mena-Segovia, J., Bolam, J.P., 2011. Phasic modulation of cortical high-frequency oscillations by pedunculopontine neurons. *Progress in Brain Research* 193, 85–92
- [83] Mena-Segovia, J., Sims, H.M., Magill, P.J., Bolam, J.P., 2008. Cholinergic brainstem neurons modulate cortical gamma activity during slow oscillations. *Journal of Physiology (London)* 586, 2947–2960
- [84] Urbain, C., et al., Pathophysiology of sleep-dependent memory consolidation processes in children, *International Journal of Psychophysiology* (2013), <http://dx.doi.org/10.1016/j.ijpsycho.2013.06.022>
- [85] Stores G, Stores R. Sleep disorders and their clinical significance in children with Down syndrome. *Dev Med Child Neurol.* 2013 Feb;55(2):126-30. doi:10.1111/j.1469-8749.2012.04422.x.
- [86] Anders T, Iosif AM, Schwichtenberg AJ, Tang K, Goodlin-Jones B. Sleep and daytime functioning: a short-term longitudinal study of three preschool-age comparison groups. *Am J Intellect Dev Disabil.* 2012 Jul;117(4):275-90. doi: 10.1352/1944-7558-117.4.275.
- [87] Miamoto CB, Pereira LJ, Ramos-Jorge ML, Marques LS. Prevalence and predictive factors of sleep bruxism in children with and without cognitive impairment. *Braz Oral Res.* 2011 Sep-Oct;25(5):439-45
- [88] Ringli M, Huber R. Developmental aspects of sleep slow waves: linking sleep, brain maturation and behavior. *Prog Brain Res.* 2011;193:63-82. doi:10.1016/B978-0-444-53839-0.00005-3
- [89] Buckley AW, Rodriguez AJ, Jennison K, Buckley J, Thurm A, Sato S, Swedo S. Rapid eye movement sleep percentage in children with autism compared with children with developmental delay and typical development. *Arch Pediatr Adolesc Med.* 2010 Nov;164(11):1032-7. doi: 10.1001/archpediatrics.2010.202
- [90] Mercier L, Pivik RT, Busby K. Sleep patterns in reading disabled children. *Sleep.* 1993 Apr;16(3):207-15.
- [91] Dijk DJ. Sleep in children, sleep spindles, and the metrics of memory. *J Sleep Res.* 2013 Apr;22(2):119-20. doi: 10.1111/jsr.12052
- [92] Kopasz M, Loessl B, Hornyak M, Riemann D, Nissen C, Piosczyk H, Voderholzer U. Sleep and memory in healthy children and adolescents-a critical review. *Sleep Med Rev.* 2010 Jun;14(3):167-77. doi: 10.1016/j.smrv.2009.10.006

- [93] Bernier A, Beauchamp MH, Bouvette-Turcot AA, Carlson SM, Carrier J. Sleep and Cognition in Preschool Years: Specific Links to Executive Functioning. *Child Dev.* 2013 Feb 24. doi: 10.1111/cdev.12063
- [94] Hahn C, Cowell JM, Wiprzycka UJ, Goldstein D, Ralph M, Hasher L, Zelazo PD. Circadian rhythms in executive function during the transition to adolescence: the effect of synchrony between chronotype and time of day. *Dev Sci.* 2012 May;15(3):408-16. doi: 10.1111/j.1467-7687.2012.01137.x
- [95] Barnes ME, Gozal D, Molfese DL. Attention in children with obstructive sleep apnoea: an event-related potentials study. *Sleep Med.* 2012 Apr;13(4):368-77. doi: 10.1016/j.sleep.2011.06.007
- [96] Diomedi M, Curatolo P, Scalise A, Placidi F, Caretto F, Gigli GL. (1999) Sleep abnormalities in mentally retarded autistic subjects: Down's syndrome with mental retardation and normal subjects. *Brain Dev.* 21(8):548-53
- [97] Miano S, Bruni O, Elia M, Scifo L, Smerieri A, Trovato A, et al. (2008) Sleep phenotypes of intellectual disability: a polysomnographic evaluation in subjects with Down syndrome and Fragile-X syndrome. *Clin Neurophysiol.*;119(6):1242-7
- [98] Miano S, Bruni O, Leuzzi V, Elia M, Verrillo E, Ferri R. (2004) Sleep polygraphy in Angelman syndrome. *Clin Neurophysiol.* 115(4):938-45
- [99] Sobanski E, Schredl M, Kettler N, Alm B. (2008) Sleep in adults with attention deficit hyperactivity disorder (ADHD) before and during treatment with methylphenidate: a controlled polysomnographic study. *Sleep.* 31(3):375-81
- [100] Lazar AS, Lazar ZI, Biro A, Gyori M, Tarnok Z, Prekop C, Keszei A, Stefanik K, Gardos J, Halasz P, Bodizs R. (2010) Reduced fronto-cortical brain connectivity during NREM sleep in Asperger syndrome: an EEG spectral and phase coherence study. *Clin Neurophysiol.* 121(11):1844-54
- [101] Gombos F, Bodizs R, Kovacs I. (2011) Atypical sleep architecture and altered EEG spectra in Williams syndrome. *J Intellect Disabil Res*;55(3):255-62
- [102] Spaepen, E. et al. (2011) Number without a language model. *Proc. Natl. Acad. Sci. U.S.A.* 108, 3163–3168
- [103] Shaw, P. et al. (2006) Intellectual ability and cortical development in children and adolescents. *Nature* 440, 676–679
- [104] Wilhelm, I. et al. (2013) The sleeping child outplays the adult's capacity to convert implicit into explicit knowledge. *Nat. Neurosci.* 16, 391–393
- [105] Stickgold R. Early to bed: how sleep benefits children's memory. *Trends Cogn Sci.* 2013 Jun;17(6):261-2. doi: 10.1016/j.tics.2013.04.006
- [106] Mander, B.A. et al. (2013) Prefrontal atrophy, disrupted NREM slow waves and impaired hippocampal-dependent memory in aging. *Nat. Neurosci.* 16, 357–364

- [107] Hupbach, A. et al. (2009) Nap-dependent learning in infants. *Dev. Sci.* 12, 1007–1012
- [108] Jenni, O.G., Van Reen, E., Carskadon, M.A., 2005. Regional differences of the sleep electroencephalogram in adolescents. *Journal of Sleep Research* 14, 141–147
- [109] Tarokh, L., Carskadon, M.A., 2010. Developmental changes in the human sleep EEG during early adolescence. *Sleep* 33, 801–809
- [110] Tarokh, L., Carskadon, M.A., Achermann, P., 2010. Developmental changes in brain connectivity assessed using the sleep EEG. *Neuroscience* 171, 622–634
- [111] Feinberg, I., Campbell, I.G., 2010. Sleep EEG changes during adolescence: an index of a fundamental brain reorganization. *Brain and Cognition* 72, 56–65
- [112] Feinberg, I., Thode, H.C., Chugani, H.T., March, J.D., 1990. Gamma distribution model describes maturational curves for delta wave amplitude, cortical metabolic rate and synaptic density. *Journal of Theoretical Biology* 142, 149–161
- [113] Feinberg, I., De Bie, E., Davis, N.M., Campbell, I.G., 2011. Topographic differences in the adolescent maturation of the slow wave EEG during NREM sleep. *Sleep* 34, 325–333
- [114] Gaudreau, H., Carrier, J., Montplaisir, J., 2001. Age-related modifications of NREM sleep EEG: from childhood to middle age. *Journal of Sleep Research* 10, 165–172
- [115] Jenni, O.G., Carskadon, M.A., 2004. Spectral analysis of the sleep electroencephalogram during adolescence. *Sleep* 27, 774–783
- [116] Tarokh, L., Reen, E. Van, Lebourgeois, M., Seifer, R., Carskadon, M.A., Van Reen, E., 2011. Sleep EEG provides evidence that cortical changes persist into late adolescence. *Sleep* 34, 1385–139
- [117] Baker, F.C., Turlington, S.R., Colrain, I., 2012. Developmental changes in the sleep electroencephalogram of adolescent boys and girls. *Journal of Sleep Research* 21, 59–67
- [118] Piantoni, G., et al., Modulation of gamma and spindle-range power by slow oscillations in scalp sleep EEG of children, *International Journal of Psychophysiology* (2013), <http://dx.doi.org/10.1016/j.ijpsycho.2013.01.017>
- [119] Gogtay, N., Giedd, J.N., Lusk, L., Hayashi, K.M., Greenstein, D., Vaituzis, A.C., Nugent, T.F., Herman, D.H., Clasen, L.S., Toga, A.W., Rapoport, J.L., Thompson, P.M., 2004. Dynamic mapping of human cortical development during childhood through early adulthood. *Proceedings of the National Academy of Sciences of the United States of America* 101, 8174–8179
- [120] Sowell, E.R., Thompson, P.M., Leonard, C.M., Welcome, S.E., Kan, E., Toga, A.W., 2004. Longitudinal mapping of cortical thickness and brain growth in normal children. *Journal of Neuroscience* 24, 8223–8231

- [121] Luna, B., Sweeney, J.A., 2004. The emergence of collaborative brain function: FMRI studies of the development of response inhibition. *Annals of the New York Academy of Sciences* 1021, 296–309
- [122] Tononi, G., Cirelli, C., 2006. Sleep function and synaptic homeostasis. *Sleep Medicine Reviews* 10, 49–62
- [123] Vyazovskiy, V.V., Olcese, U., Lazimy, Y.M., Faraguna, U., Esser, S.K., Williams, J.C., Cirelli, C., Tononi, G., 2009. Cortical firing and sleep homeostasis. *Neuron* 63, 865–878
- [124] Feinberg, I., 1982. Schizophrenia: caused by a fault in programmed synaptic elimination during adolescence? *Journal of Psychiatric Research* 17, 319–334
- [125] Ringli, M., et al., The sleep EEG topography in children and adolescents shows sex differences in language areas, *International Journal of Psychophysiology* (2013), <http://dx.doi.org/10.1016/j.ijpsycho.2013.04.008>
- [126] Huber, R., Graf, T., Cote, K.A., Wittmann, L., Gallmann, E., Matter, D., Schuderer, J., Kuster, N., Borbely, A.A., Achermann, P., 2000. Exposure to pulsed high-frequency electromagnetic field during waking affects human sleep EEG. *NeuroReport* 11, 3321–3325
- [127] Huber, R., Ghilardi, M.F., Massimini, M., Tononi, G., 2004. Local sleep and learning. *Nature* 430, 78–81
- [128] Huber, R., Ghilardi, M.F., Massimini, M., Ferrarelli, F., Riedner, B.A., Peterson, M.J., Tononi, G., 2006. Arm immobilization causes cortical plastic changes and locally decreases sleep slow wave activity. *Nature Neuroscience* 9, 1169–1176
- [129] Huber, R., Mäki, H., Rosanova, M., Casarotto, S., Canali, P., Tononi, G., Massimini, M., 2013. Human cortical excitability increases with time awake. *Cerebral Cortex* 23, 332–338
- [130] Vyazovskiy, V.V., Cirelli, C., Pfister-Genskow, M., Faraguna, U., Tononi, G., 2008. Molecular and electrophysiological evidence for net synaptic potentiation in wake and depression in sleep. *Nature Neuroscience* 11, 200–208
- [131] Shaw, P., Kabani, N.J., Lerch, J.P., Eckstrand, K., Lenroot, R., Gogtay, N., Greenstein, D., Clasen, L., Evans, A., Rapoport, J.L., Giedd, J.N., Wise, S.P., 2008. Neurodevelopmental trajectories of the human cerebral cortex. *Journal of Neuroscience* 28, 3586–3594
- [132] O'Brien, L.M., Holbrook, C.R., Mervis, C.B., Klaus, C.J., Bruner, J.L., Raffield, T.J., Rutherford, J., Mehl, R.C., Wang, M., Tuell, A., Hume, B.C., Gozal, D., 2003. Sleep and neurobehavioral characteristics of 5-to 7-year-old children with parentally reported symptoms of attention-deficit/hyperactivity disorder. *Pediatrics* 111, 554–563

- [133] Aronen, E.T., Paavonen, E.J., Fjallberg, M., Soininen, M., Torronen, J., 2000. Sleep and psychiatric symptoms in school-age children. *Journal of the American Academy of Child and Adolescent Psychiatry* 39, 502–508
- [134] Fallone, G., Acebo, C., Seifer, R., Carskadon, M.A., 2005. Experimental restriction of sleep opportunity in children: effects on teacher ratings. *Sleep* 28, 1561–1567
- [135] Sadeh, A., Gruber, R., Raviv, A., 2003. The effects of sleep restriction and extension on schoolage children: what a difference an hour makes. *Child Development* 74, 444–455
- [136] Sadeh, A., Pergamin, L., Bar-Haim, Y., 2006. Sleep in children with attention-deficit hyperactivity disorder: ameta-analysis of polysomnographic studies. *Sleep Medicine Reviews* 10, 381–398
- [137] Forbes, E.E., Bertocci, M.A., Gregory, A.M., Ryan, N.D., Axelson, D.A., Birmaher, B., Dahl, R.E., 2008. Objective sleep in pediatric anxiety disorders and major depressive disorder. 47, 148–155
- [138] Touchette, E., Petit, D., Seguin, J.R., Boivin, M., Tremblay, R.E., Montplaisir, J.Y., 2007. Associations between sleep duration patterns and behavioral/cognitive functioning at school entry. *Sleep* 30, 1213–1219
- [139] Touchette, E., Cote, S.M., Petit, D., Liu, X., Boivin, M., Falissard, B., Tremblay, R.E., Montplaisir, J.Y., 2009. Short nighttime sleep-duration and hyperactivity trajectories in early childhood. *Pediatrics* 124, e985–993
- [140] Giedd, J., 1999. Brain development, IX: human brain growth. *The American Journal of Psychiatry* 156, 4
- [141] Rice, D., Barone Jr., S., 2000. Critical periods of vulnerability for the developing nervous system: evidence from humans and animal models. *Environmental Health Perspectives* 108 (Suppl. 3), 511–533
- [142] Nofzinger, E.A., 2005. Functional neuroimaging of sleep. *Seminars in Neurology* 25, 9–18
- [143] Binder, J.R., Frost, J.A., Hammeke, T.A., Bellgowan, P.S., Rao, S.M., Cox, R.W., 1999. Conceptual processing during the conscious resting state. A functional MRI study. *Journal of Cognitive Neuroscience* 11, 80–95
- [144] Maquet, P., Degueldre, C., Delfiore, G., Aerts, J., Peters, J.M., Luxen, A., Franck, G., 1997. Functional neuroanatomy of human slow wave sleep. *Journal of Neuroscience* 17, 2807–2812
- [145] Dahl, R.E., 1996. The impact of inadequate sleep on children's daytime cognitive function. *Seminars in Pediatric Neurology* 3, 44–50
- [146] Dahl, R.E., 1996. The regulation of sleep and arousal: development and psychopathology. *Development and Psychopathology* 8, 3–27

- [147] Horne, J.A., 1993. Human sleep, sleep loss and behaviour. Implications for the prefrontal cortex and psychiatric disorder. *The British Journal of Psychiatry* 162, 413–419
- [148] Muzur, A., Pace-Schott, E.F., Hobson, J.A., 2002. The prefrontal cortex in sleep. *Trends in Cognitive Science* 6, 475–481
- [149] Drummond, S.P., Brown, G.G., Stricker, J.L., Buxton, R.B., Wong, E.C., Gillin, J.C., 1999. Sleep deprivation-induced reduction in cortical functional response to serial subtraction. *Neuroreport* 10, 3745–3748
- [150] Thomas, M., Sing, H., Belenky, G., Holcomb, H., Mayberg, H., Dannals, R., Wagner, H., Thorne, D., Popp, K., Rowland, L., Welsh, A., Balwinski, S., Redmond, D., 2000. Neural basis of alertness and cognitive performance impairments during sleepiness. I. Effects of 24 h of sleep deprivation on waking human regional brain activity. *Journal of Sleep Research* 9, 335–352
- [151] van der Helm, E., Gujar, N., Walker, M.P., 2010. Sleep deprivation impairs the accurate recognition of human emotions. *Sleep* 33, 335–342
- [152] Dijk, D.J., 2011. Risk-taking and other effects of sleep loss on brain function and behaviour. *Journal of Sleep Research* 20, 375–376
- [153] Holm, S.M., Forbes, E.E., Ryan, N.D., Phillips, M.L., Tarr, J.A., Dahl, R.E., 2009. Reward related brain function and sleep in pre/early pubertal and mid/late pubertal adolescents. *The Journal of Adolescent Health* 45, 326–334
- [154] Nelson, E.E., Leibenluft, E., McClure, E.B., Pine, D.S., 2005. The social re-orientation of adolescence: a neuroscience perspective on the process and its relation to psychopathology. *Psychological Medicine* 35, 163–174