

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

186,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Developing Tree Biomass Models for Eight Major Tree Species in China

WeiSheng Zeng

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/65664>

Abstract

In the context of climate change, estimating forest biomass for large regions is key to national carbon stocks, but few models have been developed at regional level. Based on mensuration data from large samples (4818 and 1626 trees for above- and belowground biomass, respectively) of eight major tree species in China, the author developed one- and two-variable compatible integrated model systems for aboveground and belowground biomass, biomass conversion factor (*BCF*) and root-to-shoot ratio (*RSR*), using the error-in-variable simultaneous equations. Furthermore, the differences of aboveground and belowground biomass among various species were analyzed using the dummy variable approach. The results indicated that (1) two-variable models were almost better than one-variable models for aboveground biomass estimation, while the two model systems were not significantly different for belowground biomass estimation; (2) the eight species can be ranked in terms of aboveground biomass from *Quercus* (largest), *Betula*, *Populus*, *Pinus massoniana*, *Picea*, *Larix*, *Abies* to *Cunninghamia lanceolata* and in terms of belowground biomass from *Quercus* (largest), *Betula*, *Larix*, *Picea*, *Populus*, *P. massoniana*, *C. lanceolata* to *Abies*; (3) mean prediction errors (*MPEs*) of aboveground biomass models for the species were less than 5%, whereas *MPEs* of belowground biomass equations were less than 10%, except for *Abies*.

Keywords: aboveground biomass, belowground biomass, biomass conversion factor, root-to-shoot ratio, error-in-variable simultaneous equations

1. Introduction

Increasingly, governments worldwide attach considerable importance to estimating biomass and carbon storage of forest ecosystems in the context of global climate change. To help countries conduct national greenhouse gas inventories, forest biomass estimation and carbon stock assessment, the Intergovernmental Panel on Climate Change (IPCC) provided such carbon-accounting parameters as biomass expansion factors (*BEF*) and root-to-shoot ratios (*RSR*) for estimating different geographic zones in 2003 [1]. However, it probably has great

uncertainty to apply these parameters for biomass estimation. Developing individual tree biomass models and parameters for national monitoring and assessment of biomass and carbon storage of forest ecosystems has become fundamentally important.

The earliest research on forest biomass abroad can be traced to the 1870s [2]. In recent years, biomass models for major tree species in America, Canada and some European countries have been developed or improved [3–11]. Their purpose was to assess and monitor forest biomass and carbon storage and to provide a basis for evaluating the contribution of forest ecosystems to the global carbon cycle. Studies on forest biomass in China have only been implemented since the late 1970s when some related articles were published [12, 13], i.e., a century after the earliest study abroad. Due to special historical reasons, China did not participate in the International Biological Program (IBP), initiated by the International Union of Forest Research Organizations (IUFRO), during the period of 1964–1974 and thus missed the golden development stage of forest biomass research [14].

Reviewing the development of forest biomass modeling near 40 years in China, three stages could be classified: the first is estimating biomass and productivity of major forest types toward the end of the twentieth century [13, 15–30]; the second is assessing carbon storage in Chinese forest ecosystems since the beginning of the current century [31–37]; and the third is the new development stage for monitoring and assessing forest biomass and carbon storage at provincial and national levels [14, 38]. To monitor forest biomass and carbon storage in the National Forest Inventory (NFI) system, the National Forest Biomass Modeling Program has been implemented since early 2009. Up to now, many papers on modeling individual tree biomass have been published [39–51], which classified 70 modeling populations for developing individual tree biomass models, determined the sample structure of each population and studied the modeling methods including nonlinear error-in-variable simultaneous equations, mixed-effects modeling, dummy variable modeling and segmented modeling approaches. Also, logarithmic regression and weighted regression were analyzed [52] and goodness evaluation and precision analysis of biomass models were studied [53]. Based on the studying achievements, two ministerial standards on technical regulations and five ministerial standards on biomass models have been approved for application [54–60]. In the near future, more ministerial standards on biomass models for other tree species would be published.

From the published papers and ministerial standards, we could find that the aboveground and belowground biomass models were developed separately owing to the unequal sample sizes and most of the studies were only based on sample trees of one tree species. In this chapter, the author will use the mensuration data of aboveground and belowground biomass from 4818 to 1626 destructive sample trees of eight major tree species, respectively. The main purpose was to develop an integrated individual tree model system for aboveground and belowground biomass, biomass conversion factor (*BCF*) and root-to-shoot ratio (*RSR*), using the approach of nonlinear error-in-variable simultaneous equations with dummy variable. The system could assure aboveground biomass models compatible with stem volume models and *BCF* models and belowground biomass models compatible with aboveground biomass models and *RSR* models. Secondly, the generalized dummy-variable models of aboveground and belowground biomass for eight major tree species were established and compared and the ranks of eight

species for aboveground and belowground biomass estimation were provided respectively from the species-specific parameter estimates.

2. Materials and methods

2.1. Data

During the 5 years between 2009 and 2013, a total amount of 4818 sample trees for 31 modeling populations of eight major tree species or species groups, namely, *Picea* spp., *Abies* spp., *Betula* spp., *Quercus* spp., *Populus* spp., *Larix* spp., *Cunninghamia lanceolata* and *Pinus massoniana*, which occupied more than 60% of forest volume in China [39], were felled for aboveground biomass mensuration. The sample trees were evenly distributed in ten diameter classes of 2, 4, 6, 8, 12, 16, 20, 26, 32 and more than 38 cm for each modeling population, and about 15 sample trees in each diameter class were selected by height class as evenly as possible. For example, if three height classes were defined, i.e., low, intermediate and high, then five sample trees should be selected in each height class. For each sample tree, the diameter at breast height of stem was measured in the field. After the tree was felled, total trunk length (tree height, from ground level to the top) and live crown length were also measured. The trunk was divided into 11 sections at points corresponding to 0, 0.05, 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8 and 0.9 of tree height. Base diameters of all sections were measured and the tree volume was computed using Smalian's formula [61], which referred to total volume over bark. Specifically, the formula was written as $V = (A_1 + A_2)/2 \times L$ with V as the volume of a section of tree trunk, A_1 and A_2 as two areas of the small and large ends of the section and L as the section length. The fresh weights of stem, branch and foliage were also measured; subsamples were selected and weighed in the field [54]. Among all sample trees, about one third (1626 trees) were selected for measuring both aboveground and belowground biomass. The whole roots were excavated out, fresh weights of stump, coarse roots (more than 10 mm) and small roots (2–10 mm, not including fine roots less than 2 mm) were measured, respectively and subsamples were selected. After being taken into the laboratory, all subsamples were oven-dried at 85°C until a constant weight was reached. According to the ratio of dry weight to fresh weight, each component biomass was computed and the aboveground biomass of the tree was obtained by summation [54]. **Table 1** shows the general situation for biomass samples of eight major tree species or groups.

2.2 Model construction

The general form of individual tree biomass and stem volume models is as follows [45, 62]:

$$y = \beta_0 x_1^{\beta_1} x_2^{\beta_2} \cdots x_j^{\beta_j} + \varepsilon \quad (1)$$

where y is biomass (kg), x_j are predictive biometric variables, which reflect the dimensions of a tree, such as diameter at breast height D (cm) and tree height H (m), β_j are parameters and ε is the error term. Because the biomass data are significantly heteroscedastic, some measures should be taken to eliminate heteroscedasticity prior to parameter estimation. In this paper, weighted regression was applied and the specific weight functions were derived from the

Species	Samples	Variables	Mean	Min	Max	S.D.	CV (%)
<i>Picea</i> spp.	900/295	Diameter D (cm)	17.0	1.0	65.5	12.8	75.6
		Height H (m)	12.3	1.4	46.9	8.1	66.4
		Stem volume V (dm ³)	343.0	0.6	6770.7	609.9	177.8
		Aboveground biomass M_a (kg)	174.5	0.4	1668.9	251.3	143.9
		Belowground biomass M_b (kg)	41.2	0.1	289.1	61.3	148.8
<i>Abies</i> spp.	751/249	Diameter D (cm)	17.1	1.1	68.0	13.0	76.6
		Height H (m)	11.9	1.5	39.0	7.4	62.7
		Stem volume V (dm ³)	352.4	0.5	4525.0	589.5	167.3
		Aboveground biomass M_a (kg)	168.9	0.3	1817.0	262.7	155.6
		Belowground biomass M_b (kg)	29.0	0.1	393.4	52.4	180.7
<i>Betula</i> spp.	690/236	Diameter D (cm)	15.9	1.0	60.8	11.8	73.7
		Height H (m)	11.3	1.9	33.0	6.2	55.1
		Stem volume V (dm ³)	235.0	0.3	2782.7	345.9	147.2
		Aboveground biomass M_a (kg)	167.4	0.2	1671.0	240.6	143.7
		Belowground biomass M_b (kg)	45.0	0.1	343.6	67.0	148.8
<i>Quercus</i> spp.	670/228	Diameter D (cm)	16.1	1.5	54.0	11.6	72.1
		Height H (m)	10.9	1.4	28.6	6.3	57.6
		Stem volume V (dm ³)	253.2	0.2	2487.1	370.9	146.5
		Aboveground biomass M_a (kg)	208.2	0.3	1664.1	295.2	141.8
		Belowground biomass M_b (kg)	51.4	0.1	385.9	71.6	139.4
<i>Populus</i> spp.	602/207	Diameter D (cm)	16.4	1.2	48.9	11.9	72.3
		Height H (m)	12.9	2.4	31.1	6.9	53.6
		Stem volume V (dm ³)	281.4	0.3	2228.4	385.3	136.9
		Aboveground biomass M_a (kg)	174.1	0.2	1065.1	241.3	138.6
		Belowground biomass M_b (kg)	35.6	0.1	384.5	54.3	152.7
<i>Larix</i> spp.	602/199	Diameter D (cm)	16.7	1.5	54.2	12.3	73.7
		Height H (m)	12.6	1.4	37.5	7.6	60.0
		Stem volume V (dm ³)	316.6	0.6	3016.6	471.7	149.0
		Aboveground biomass M_a (kg)	160.4	0.2	1301.9	231.1	144.1
		Belowground biomass M_b (kg)	41.0	0.1	300.0	61.8	150.9
<i>Cunninghamia lanceolata</i>	302/108	Diameter D (cm)	16.4	1.8	42.0	11.8	71.8
		Height H (m)	11.5	1.9	33.0	7.1	61.7
		Stem volume V (dm ³)	293.7	0.6	1815.2	409.7	139.5
		Aboveground biomass M_a (kg)	75.6	0.3	644.9	105.5	139.5
		Belowground biomass M_b (kg)	25.9	0.1	174.9	37.7	145.8

Species	Samples	Variables	Mean	Min	Max	S.D.	CV (%)
<i>Pinus massoniana</i>	301/104	Diameter D (cm)	16.5	1.2	47.2	11.9	72.4
		Height H (m)	12.1	1.6	30.3	7.2	59.4
		Stem volume V (dm ³)	300.8	0.3	1825.4	405.7	134.9
		Aboveground biomass M_a (kg)	125.1	0.1	1079.3	171.6	137.2
		Belowground biomass M_b (kg)	35.5	0.1	285.0	53.7	151.6

Min—minimum, Max—maximum, S.D.—standard deviation, and CV—coefficient of variation. The sample sizes are for aboveground and belowground biomass mensuration, respectively.

Table 1. General situation of biomass samples for eight major tree species.

residuals of independently fitted models by ordinary least squares regression [62, 63]. Since models based on one (D) or two variables (D and H) have been commonly used, this paper develops both one- and two-variable models. The aboveground biomass, belowground biomass and stem volume models based on two variables can be expressed respectively as:

$$M_a = a_0 D^{a_1} H^{a_2} + \varepsilon \quad (2)$$

$$M_b = b_0 D^{b_1} H^{b_2} + \varepsilon \quad (3)$$

$$V = c_0 D^{c_1} H^{c_2} + \varepsilon \quad (4)$$

where M_a and M_b are aboveground and belowground biomass (kg), respectively; V is stem volume (dm³); a_i , b_i and c_i are parameters; and other symbols are the same as above.

2.2.1 Integrated compatible model systems

The aboveground biomass is correlated to stem volume through biomass conversion factor (BCF), which is equal to biomass expansion factor (BEF) multiplied by basic wood density following the IPCC's approach [64]. Because the BCF is an important parameter for forest biomass estimation [65], it is very common to develop both an aboveground biomass model and a BCF model that are compatible with stem volume model [45, 51]. Similarly, belowground biomass is connected with aboveground biomass model through root-to-shoot ratio (RSR) [66, 67]. Because the RSR model is also an important parameter for forest biomass estimation, generally both belowground biomass model and RSR model compatible with aboveground biomass model are developed simultaneously [44]. Therefore, we can develop an integrated aboveground and belowground biomass model system through using the nonlinear error-in-variable simultaneous equation approach [51, 68]. Because the belowground biomass observations were only 1/3 of the aboveground biomass observations, a dummy variable (x) was required for those trees for which no belowground biomass observation was available, i.e., 1 for the trees with belowground biomass observation and 0 for the trees with no belowground biomass observation [69]. The system can ensure the compatibility between

aboveground biomass, belowground biomass, stem volume, *BCF* and *RSR*. The one- and two-variable integrated systems are as follows, respectively:

$$\begin{cases} M_a = a_0 D^{a_1} + \varepsilon \\ M_b = b_0 D^{b_1} x + \varepsilon \\ V = c_0 D^{c_1} + \varepsilon \\ BCF = a_0 D^{a_1} / c_0 D^{c_1} + \varepsilon \\ RSR = b_0 D^{b_1} x / a_0 D^{a_1} + \varepsilon \end{cases} \quad (5)$$

$$\begin{cases} M_a = a_0 D^{a_1} H^{a_2} + \varepsilon \\ M_b = b_0 D^{b_1} H^{b_2} x + \varepsilon \\ V = c_0 D^{c_1} H^{c_2} + \varepsilon \\ BCF = a_0 D^{a_1} H^{a_2} / c_0 D^{c_1} H^{c_2} + \varepsilon \\ RSR = b_0 D^{b_1} H^{b_2} x / a_0 D^{a_1} H^{a_2} + \varepsilon \end{cases} \quad (6)$$

where, M_a , M_b , V , *BCF* and *RSR* are aboveground biomass, belowground biomass, stem volume, biomass conversion factor and root-to-shoot ratio, respectively, which are regarded as error-in-variables; D and H are diameter at breast height and tree height, which are regarded as error-free variables; x is a dummy variable to distinguish if belowground biomass is available; and a_i , b_i and c_i are parameters.

Various methods have been attempted to estimate the parameters of the simultaneous equations. Parresol [63] used the seemingly unrelated regression (SUR) for solving the additivity of simultaneous biomass equations. Tang et al. [70] further developed an error-in-variable modeling approach to estimate the parameters of simultaneous equations, which has been widely used in recent years [40, 45, 49, 51]. In this study, the error-in-variable simultaneous equation approach was used to estimate the parameters of the integrated systems based on maximum likelihood estimation through ForStat software (statistical software with analytical tools for forestry as well as general statistical procedures, developed in the Chinese Academy of Forestry, Beijing, China) [68].

In addition, the weighted regression method was used to eliminate the heteroscedasticity commonly exhibited in biomass and volume data by using specific weight functions, which were derived from the residuals of biomass or volume equations fitted through the ordinary least square (OLS) technique [52, 62]. For biomass conversion factor and root-to-shoot ratio modeling, the OLS regression technique was directly used to estimate the parameters because the *BCF* and *RSR* data mostly exhibited homoscedasticity.

2.2.2 Generalized dummy variable models

The one-variable biomass equation was the most widely used model in estimating individual tree biomass [3, 7]. The power function of one-variable aboveground biomass equation was based on the WBE theory for the origin of allometric scaling laws [71, 72]. According to the results from Zeng and Tang [73], the generalized one-variable aboveground biomass model can be expressed as:

$$M_a = aD^{7/3} + \varepsilon \quad (7)$$

That is, the power parameter of the allometric model is constantly equal to $7/3$ (≈ 2.33), only the parameter a depends on tree species. If a variable vector z was defined as dummy variable to indicate tree species, then the generalized model (7) could be expressed as:

$$M_a = (a + v_a z) D^{7/3} + \varepsilon \quad (8)$$

where a is the global parameter and v_a is tree species-specific parameter vector. The dummy variable vector z includes seven elements, indicating the eight tree species by the following combinations:

$z_1 = 1, z_2 = 0, z_3 = 0, z_4 = 0, z_5 = 0, z_6 = 0$ and $z_7 = 0$ for *Picea* spp.

$z_1 = 0, z_2 = 1, z_3 = 0, z_4 = 0, z_5 = 0, z_6 = 0$ and $z_7 = 0$ for *Abies* spp.

$z_1 = 0, z_2 = 0, z_3 = 1, z_4 = 0, z_5 = 0, z_6 = 0$ and $z_7 = 0$ for *Betula* spp.

$z_1 = 0, z_2 = 0, z_3 = 0, z_4 = 1, z_5 = 0, z_6 = 0$ and $z_7 = 0$ for *Quercus* spp.

$z_1 = 0, z_2 = 0, z_3 = 0, z_4 = 0, z_5 = 1, z_6 = 0$ and $z_7 = 0$ for *Populus* spp.

$z_1 = 0, z_2 = 0, z_3 = 0, z_4 = 0, z_5 = 0, z_6 = 1$ and $z_7 = 0$ for *Larix* spp.

$z_1 = 0, z_2 = 0, z_3 = 0, z_4 = 0, z_5 = 0, z_6 = 0$ and $z_7 = 1$ for *C. lanceolata*

$z_1 = 0, z_2 = 0, z_3 = 0, z_4 = 0, z_5 = 0, z_6 = 0$ and $z_7 = 0$ for *P. massoniana*

Consequently, from comparing the estimated values of species-specific parameter vector v_a , the differences among various tree species could be analyzed.

2.3 Model evaluation

Many statistical indices could be used to evaluate individual tree biomass models [63]. According to the study results from Zeng and Tang [53], the following six statistical indices, namely, the coefficient of determination (R^2), standard error of estimate (SEE), mean prediction error (MPE), total relative error (TRE), average systematic error (ASE) and mean percent standard error ($MPSE$), were very important for assessing biomass models. In this study, the same six statistical indices were used for model evaluation [50, 51]:

$$R^2 = 1 - \frac{\sum (y_i - \hat{y}_i)^2}{\sum (y_i - \bar{y})^2} \quad (9)$$

$$SEE = \sqrt{\frac{\sum (y_i - \hat{y}_i)^2}{n-p}} \quad (10)$$

$$TRE = \frac{\sum (y_i - \hat{y}_i)}{\sum \hat{y}_i} \times 100 \quad (11)$$

$$ASE = \frac{\sum (y_i - \hat{y}_i)}{\hat{y}_i} / n \times 100 \quad (12)$$

$$MPE = t_\alpha \cdot (SEE / \bar{y}) / \sqrt{n} \times 100 \quad (13)$$

$$MPSE = \frac{\sum |(y_i - \hat{y}_i) / \hat{y}_i|}{n} \times 100 \quad (14)$$

where y_i are observed values, \hat{y}_i are estimated values, \bar{y} is mean value of samples, n is the number of samples, p is the number of parameters and t_α is the t -value at confidence level α with $n-p$ degrees of freedom.

3. Results and analysis

The one- and two-variable integrated systems (Eqs. (5) and (6)) for eight tree species or groups were estimated using the error-in-variable simultaneous equation approach through ForStat (Tables 2 and 3). The six fitting statistics, R^2 , SEE , TRE , ASE , MPE and $MPSE$, were calculated and could be used for evaluating the goodness-of-fit of the three models (Table 4). From the

Species	Aboveground biomass models		Belowground biomass models		Stem volume models	
	a_0	a_1	b_0	b_1	c_0	c_1
Pi	0.17417	2.2270	0.04853	2.1954	0.1528	2.4548
Ab	0.10195	2.3676	0.02873	2.2452	0.1297	2.5106
Be	0.13392	2.3401	0.05767	2.2039	0.1712	2.3653
Qu	0.16592	2.3409	0.10619	2.0373	0.1448	2.4351
Po	0.09198	2.4490	0.02958	2.3200	0.1410	2.4702
La	0.12473	2.3190	0.03154	2.3355	0.1464	2.4737
Cl	0.09782	2.3099	0.02853	2.2500	0.1144	2.5421
Pm	0.13771	2.3243	0.01959	2.4400	0.1514	2.4655

Pi—*Picea* spp., Ab—*Abies* spp., Be—*Betula* spp., Qu—*Quercus* spp., Po—*Populus* spp., La—*Larix* spp., Cl—*Cunninghamia lanceolata*, and Pm—*Pinus massoniana*. Same in Tables 3–7.

Table 2. The parameter estimates of the one-variable integrated system (Eq. (5)).

Species	Aboveground biomass models			Belowground biomass models			Stem volume models		
	a_0	a_1	a_2	b_0	b_1	b_2	c_0	c_1	c_2
Pi	0.11007	2.1369	0.2615	0.03284	2.3516	-0.0527	0.07763	1.7758	-1.0122
Ab	0.06720	2.0221	0.5442	0.02412	2.5974	-0.3600	0.07429	1.8135	0.9975
Be	0.08322	2.0749	0.4844	0.04531	2.1630	0.1401	0.08383	1.8246	0.8965
Qu	0.10520	1.9808	0.5939	0.09338	2.1694	-0.1091	0.07796	1.8607	0.9115
Po	0.06304	2.2460	0.3588	0.03216	2.5313	-0.2697	0.07611	1.9503	0.7927
La	0.07437	2.0003	0.5438	0.02195	2.2354	0.2369	0.07610	1.8067	0.9827
Cl	0.06740	1.9253	0.5765	0.02252	2.5080	-0.2072	0.07417	1.7949	1.0121
Pm	0.10462	2.1591	0.2857	0.01744	2.5697	-0.1028	0.09393	1.8696	0.8451

Table 3. The parameter estimates of the two-variable integrated system (Eq. (6)).

Species	Systems	Items	R^2	SEE	MPE (%)	TRE (%)	ASE (%)	MPSE (%)
Pi	(5)	AB	0.9109	75.05	2.81	1.31	-2.44	24.21
		BB	0.7842	28.55	5.51	-0.25	-3.75	40.29
		SV	0.8380	245.64	4.68	2.63	0.19	27.65
	(6)	AB	0.9061	77.10	2.89	0.37	7.68	25.26
		BB	0.7751	29.19	5.63	0.65	9.28	43.70
		SV	0.9744	97.70	1.86	0.44	6.01	15.87
Ab	(5)	AB	0.9223	73.29	3.10	0.28	-1.24	22.19
		BB	0.5474	35.29	11.97	-1.15	-2.74	49.06
		SV	0.9222	164.52	3.34	-0.60	-0.29	20.79
	(6)	AB	0.9434	62.61	2.65	0.24	4.97	22.50
		BB	0.5547	35.08	11.89	0.37	7.64	52.51
		SV	0.9800	83.41	1.69	0.25	4.34	13.05
Be	(5)	AB	0.9139	70.63	3.15	2.02	0.11	22.52
		BB	0.7734	31.95	6.16	2.47	-1.15	37.64
		SV	0.9118	102.74	3.26	2.87	7.19	26.85
	(6)	AB	0.9332	62.27	2.78	1.05	2.47	21.30
		BB	0.7741	31.97	6.17	0.18	1.16	38.38
		SV	0.9566	72.13	2.29	0.49	6.53	19.60
Qu	(5)	AB	0.9030	91.99	3.35	2.67	0.93	27.82
		BB	0.8168	30.73	5.23	2.21	-1.83	40.37
		SV	0.9262	100.81	3.01	4.11	4.36	27.33
	(6)	AB	0.9285	79.04	2.87	1.67	2.81	24.66
		BB	0.8133	31.09	5.29	1.11	2.35	41.08
		SV	0.9790	53.88	1.61	1.07	3.71	16.72
Po	(5)	AB	0.9379	60.17	2.76	1.51	-0.86	17.55
		BB	0.8440	21.51	5.47	2.21	-2.71	32.15
		SV	0.9539	82.81	2.35	-0.41	1.51	15.87
	(6)	AB	0.9506	53.72	2.47	1.17	1.25	17.11
		BB	0.8618	20.29	5.15	1.77	1.15	32.11
		SV	0.9842	48.49	1.38	0.83	1.04	9.48
La	(5)	AB	0.9123	68.47	3.41	1.26	1.15	23.18
		BB	0.6773	35.22	8.65	-0.52	0.14	37.37
		SV	0.9016	148.10	3.74	0.10	4.26	25.04
	(6)	AB	0.9432	55.14	2.75	0.22	6.91	24.68
		BB	0.6675	35.83	8.81	-0.83	6.78	40.08
		SV	0.9770	71.63	1.81	0.21	5.58	15.05

Species	Systems	Items	R^2	SEE	MPE (%)	TRE (%)	ASE (%)	MPSE (%)
Cl	(5)	AB	0.9614	30.62	2.98	3.24	1.03	22.99
		BB	0.8414	15.08	7.43	1.66	1.75	38.39
		SV	0.9474	94.14	3.62	1.99	2.28	17.25
	(6)	AB	0.9774	23.47	2.28	1.47	6.10	23.66
		BB	0.8342	15.50	7.64	-0.88	9.87	41.43
		SV	0.9931	34.10	1.31	0.50	5.21	11.30
Pm	(5)	AB	0.9542	48.45	3.21	0.91	0.25	17.29
		BB	0.8509	20.85	7.64	1.55	-2.12	39.21
		SV	0.9503	90.62	3.40	0.17	0.34	20.87
	(6)	AB	0.9572	46.89	3.10	0.41	3.76	16.30
		BB	0.8546	20.69	7.59	-0.63	1.44	39.73
		SV	0.9846	50.59	1.90	0.64	1.13	12.51

AB—aboveground biomass, BB—belowground biomass, SV—stem volume, R^2 —coefficient of determination, SEE—standard error of estimate, MPE—mean prediction error, TRE—total relative error, ASE—average systematic error, and MPSE—mean percent standard error.

Units of SEE: dm^3 for volume and kg for biomass.

Table 4. The fitting statistics of two integrated systems (Eqs. (5) and (6)).

fitting results of integrated systems (Eqs. (5) and (6)), the parameter estimates of the BCF and RSR models could be obtained (Table 5).

From comparison of the fitting statistics of two integrated systems (Eqs. (5) and (6)) in Table 4, we can find that for aboveground biomass estimation, two-variable models were better than one-variable models except *Picea*. For belowground biomass estimation, one- and two-variable models were not significantly different, even some of one-variable models were slightly better than two-variable models, such as *Picea*, *Quercus*, *Larix* and *C. lanceolata*. Considering that tree height measurement is time consuming and two-variable biomass models are not significantly different from one-variable models, especially for belowground biomass estimation, it was commended to apply one-variable models in forestry practice such as National Forest Inventory.

From Table 2, it was found that the estimates of parameter a_1 were approximately equal to 7/3, confirming the results of an earlier study [73]. To analyze the difference among various tree species, the dummy model (8) was fitted using the aboveground biomass data of all eight species (Table 6).

According to the parameter estimates in Table 6, we could rank the eight tree species by aboveground biomass estimates in descending order as *Quercus*, *Betula*, *Populus*, *P. massoniana*, *Picea*, *Larix*, *Abies* and *C. lanceolata*. That is, *Quercus* had the largest aboveground biomass, whereas *C. lanceolata* had the smallest one for the same diameter trees. The aboveground biomass estimates of the dummy model (Eq. (8)) for *Quercus*, *Betula*, *Populus*, *P. massoniana*,

Species	Systems	BCF models	RSR models
Pi	(5)	$BCF = 1.1401 D^{-0.2278}$	$RSR = 0.2786 D^{-0.0316}$
	(6)	$BCF = 1.4183 D^{0.3611} H^{-0.7507}$	$RSR = 0.2983 D^{0.2147} H^{-0.3142}$
Ab	(5)	$BCF = 0.7858 D^{-0.1430}$	$RSR = 0.2818 D^{-0.1224}$
	(6)	$BCF = 0.9046 D^{0.2086} H^{-0.4533}$	$RSR = 0.3589 D^{0.5753} H^{-0.9041}$
Be	(5)	$BCF = 0.7821 D^{-0.0252}$	$RSR = 0.4307 D^{-0.1362}$
	(6)	$BCF = 0.9928 D^{0.2504} H^{-0.4121}$	$RSR = 0.5445 D^{0.0880} H^{-0.3443}$
Qu	(5)	$BCF = 1.1456 D^{-0.0942}$	$RSR = 0.6400 D^{-0.3036}$
	(6)	$BCF = 1.3494 D^{0.1201} H^{-0.3177}$	$RSR = 0.8877 D^{0.1887} H^{-0.7030}$
Po	(5)	$BCF = 0.6522 D^{-0.0212}$	$RSR = 0.3216 D^{-0.1290}$
	(6)	$BCF = 0.8283 D^{0.2958} H^{-0.4339}$	$RSR = 0.5102 D^{0.2853} H^{-0.6285}$
La	(5)	$BCF = 0.8522 D^{-0.1547}$	$RSR = 0.2528 D^{0.0165}$
	(6)	$BCF = 0.9773 D^{0.1936} H^{-0.4389}$	$RSR = 0.2951 D^{0.2351} H^{-0.3068}$
Cl	(5)	$BCF = 0.8554 D^{-0.2321}$	$RSR = 0.2917 D^{-0.0599}$
	(6)	$BCF = 0.9087 D^{0.1303} H^{-0.4356}$	$RSR = 0.3341 D^{0.5828} H^{-0.7836}$
Pm	(5)	$BCF = 0.9096 D^{-0.1412}$	$RSR = 0.1422 D^{0.1157}$
	(6)	$BCF = 1.1138 D^{0.2894} H^{-0.5593}$	$RSR = 0.1667 D^{0.4106} H^{-0.3885}$

Table 5. The simultaneously estimated BCF and RSR models.

Species	Global parameter (<i>a</i>)	Species-specific parameters (<i>v_a</i>)
Pi	0.13485	-0.01084
Ab		-0.01959
Be		0.00443
Qu		0.03908
Po		0.00106
La		-0.01441
Cl		-0.04254
Pm		0.00000

Table 6. The parameter estimates of dummy aboveground biomass model (Eq. (8)).

Picea, *Larix* and *Abies* were 88%, 51%, 47%, 46%, 34%, 30% and 25% larger, respectively, than that for *C. lanceolata* (see **Figure 1**).

Similarly, for one-variable belowground biomass models, it was found that the estimates of parameter b_1 for eight species were not significantly different. To analyze the difference of belowground biomass estimation among various tree species, we fitted the following dummy model:




Figure 1. Comparison of aboveground biomass models for eight tree species.

Global parameters			
Species	b_0	b_1	Species-specific parameters (v_b)
Pi	0.03551	2.2544	0.00424
Ab			-0.00792
Be			0.01437
Qu			0.01835
Po			0.00091
La			0.00583
Cl			-0.00761
Pm			0.00000

Table 7. The parameter estimates of dummy belowground biomass model (Eq. (15)).

$$M_b = (b_0 + v_b z) D^{b_1} + \varepsilon \tag{15}$$

where b_0 and b_1 are global parameters and v_b is species-specific parameter vector. The parameter estimates of dummy model (Eq. (15)) are listed in Table 7.

According to the parameter estimates in Table 7, we could rank the eight tree species by below-ground biomass estimates in descending order as *Quercus*, *Betula*, *Larix*, *Picea*, *Populus*, *P.*




Figure 2. Comparison of belowground biomass models for eight tree species.

massoniana, *C. lanceolata* and *Abies*. That is, *Quercus* had the largest belowground biomass, while *Abies* had the smallest one for the same diameter trees. The belowground biomass estimates of the dummy model (Eq. (15)) for *Quercus*, *Betula*, *Larix*, *Picea*, *Populus*, *P. massoniana* and *C. lanceolata* were 95%, 81%, 50%, 44%, 32%, 29% and 1% larger, respectively, than that for *Abies* (see **Figure 2**).

4. Discussion and conclusion

In this study, data on above- and belowground biomass from 4818 to 1626 sample trees, respectively, for eight major tree species in China were used to develop compatible individual tree biomass models. The models included aboveground biomass equations and *BCF* equations compatible with stem volume equations and belowground biomass equations and *RSR* models compatible with aboveground biomass equations. To solve compatibility of the biomass models, the nonlinear error-in-variable simultaneous equations were applied and to solve the issue of unequal sample sizes for above- and belowground biomass, the dummy-variable model approach was used. In the technical regulation on methodology for tree biomass modeling [55], the segmented modeling approach was recommended when the biomass estimate of small trees was obviously biased [43, 46]. Furthermore, for the tree species distributed in various regions, it was generally needed to develop biomass models for different regions. For example, according to the population classification on modeling of single-tree biomass equations [39], it was necessary to establish five sets of biomass models for both *Abies* and *Picea*. But in this study, the segmented modeling approach was not used to develop biomass models for large and small trees, respectively and the differences among various

regions were not taken into account, only one set of biomass models, including one- and two-variable models, was developed for each tree species.

The data of three tree species, i.e., *C. lanceolata*, *P. massoniana* and *Larix* spp., were used or partly used to develop biomass models, which were published as original papers [40–51] or ministerial standards [56, 57]. Comparing with the study results by Zeng et al. [47], the parameter estimates and fitness indices of aboveground biomass and volume models are very close to those for *C. lanceolata* in this study. From the achievements by Zeng and Tang [45], we can find that the parameter estimates of aboveground biomass and volume models are not significantly different from those for *P. massoniana* in this chapter, but this study provided better models considering the statistical indices of goodness-of-fit. Comparing with the biomass models published as ministerial standards [56, 57], the developed models in this study are more generalized and simpler for application in national and regional biomass estimation. There are four sets of biomass models in total for trees (dbh \geq 5 cm) and saplings (dbh < 5 cm) for two modelling populations of each tree species in the ministerial standards [56, 57] and here we have only one set of biomass models which are suitable for both trees and saplings and for the whole country.

The results indicated that two-variable models were almost better than one-variable models for aboveground biomass estimation, while the two model systems were not significantly different for belowground biomass estimation. The mean prediction errors (*MPEs*) of aboveground biomass models for the eight species were less than 5%, whereas *MPEs* of belowground biomass equations were less than 10%, except for *Abies*. The models developed in this study can provide a basis for estimating biomass for the eight major tree species in China and will fill in the lack for China on the web platform GlobAllomeTree [74]. Also, they will have the potential to support the implementation of policies and mechanisms designed to mitigate climate change (e.g., CDM and REDD+) and to calculate costs and benefits associated with forest carbon projects. In addition, the overall modeling methodology presented in this study can be taken into consideration in any case that involves individual tree biomass modeling.

Acknowledgements

The author acknowledges the National Biomass Modeling Program in Continuous Forest Inventory (NBMP-CFI), which was funded by the State Forestry Administration of China, for providing the mensuration biomass data of eight tree species. This study was financially supported by the National Natural Science Foundation of China (Grant No. 31370634).

Author details

WeiSheng Zeng

Address all correspondence to: zengweisheng@forestry.gov.cn

Academy of Forest Inventory and Planning, State Forestry Administration, Beijing, China

References

- [1] IPCC. Good practice guidance for land use, land use change and forestry. The Institute for Global Environmental Strategies for the IPCC, Japan; 2003
- [2] Ebermeyr E. A comprehensive study of forest litter with regard to silvicultural effects on soil chemistry. Springer, Berlin; 1876
- [3] Jenkins JC, Chojnacky DC, Heath LS, Birdsey RA. National-scale biomass estimators for United States tree species. *For Sci.* 2003; 49(1):12–35
- [4] Bi H, Turner J, Lambert MJ. Additive biomass equations for native eucalypt forest trees of temperate Australia. *Trees.* 2004; 18:467–479
- [5] Lambert MC, Ung CH, Raulier F. Canadian national tree aboveground biomass equations. *Can J For Res.* 2005; 35:1996–2018
- [6] Snorrason A, Einarsson SF. Single-tree biomass and stem volume functions for eleven tree species used in Icelandic forestry. *Icelandic Agric Sci.* 2006; 19:15–24
- [7] Muukkonen P. Generalized allometric volume and biomass equations for some tree species in Europe. *Eur J For Res.* 2007; 126:157–166
- [8] Návar J. Allometric equations for tree species and carbon stocks for forests of northwestern Mexico. *For Ecol Manage.* 2009; 257: 427–434
- [9] Blujdea VNB, Pilli R, Dutca I, Ciuvat L, Abrudan IV. Allometric biomass equations for young broadleaved trees in plantations in Romania. *For Ecol Manage.* 2012; 264:172–184
- [10] Fayolle A, Doucet JL, Gillet JF, Bourland N, Lejeune P. Tree allometry in central Africa: testing the validity of pantropical multi-species allometric equations for estimating biomass and carbon stocks. *For Ecol Manage.* 2013; 305:29–37
- [11] Sileshi GW. A critical review of forest biomass estimation models, common mistakes and corrective measures. *For Ecol Manage.* 2014; 3: 237–254
- [12] Li WH. Concept of forest biomass productivity and its basic studying approach. *Nat Res.* 1978; 1:71–92
- [13] Pan WC, Li LC, Gao ZH. Biomass and nutrient elements distribution of two different forest types of Chinese fir. *Hunan For Sci Technol.* 1980; 4:1–14
- [14] Zeng WS. Development of monitoring and assessment of forest biomass and carbon storage in China. *For Ecosyst.* 2014; 1:20. DOI: 10.1186/s40663-014-0020-5
- [15] Li WH, Deng KM, Li F. Research on biomass productivity of major ecosystems in the Changbai Mountain. *For Ecosyst Res.* 1981; (Test issue):34–50
- [16] Feng ZW, Chen CY, Zhang JW, Wang KP, Zhao JL, Gao H. Determination of biomass of *Pinus massoniana* stand in Huitong county, Hunan province. *Sci Silv Sin.* 1982; 18(2):127–134

- [17] Ye JZ. Annual dynamic of the biomass of Chinese fir forests on Yangkou forestry farm, J Nanjing For Univ. 1984; 4:1–9
- [18] Chen LZ, Chen QL, Bao XC, Ren JK, Miu YG, Hu YH. Studies on Chinese arborvitae (*Platycladus orientalis*) forest and its biomass in Beijing. J Plant Ecol. 1986; 10(1):17–25
- [19] Xu ZB. Biomass productivity of major forest types in Daxinganling. Chin J Ecol. 1988; 7 (Sp):49–54
- [20] Ma QY. A study on the biomass of Chinese pine forests. J Beijing For Univ. 1989; 11(4):1–10
- [21] Chen CG, Zhu JF. Manual of tree biomass for main species in northern China. Chinese Forestry Press, Beijing; 1989.
- [22] Liu SR, Chai YX, Cai TJ, Peng CH. Research on biomass and net primary productivity of *Larix gmelini* plantations. J Northeast For Univ. 1990; 18(2): 40–46
- [23] Liu ZG. Research on biomass and productivity of *Larix principis-rupprechtii* plantations. J Beijing For Univ. 1992; 14(Sp.1):114–123
- [24] Liu XZ. Study on biomass of Masson pine forests at different ages. For Res Manage. 1993; 2: 77–80
- [25] Fang JY, Liu GH, Xu SL. Biomass and net production of forest vegetation in China. Acta Ecol Sin. 1996; 16(4): 497–508
- [26] Luo TX. Patterns of net primary productivity for Chinese major forest types and their mathematical models. Dissertation, Commission for Integrated Survey of Natural Resources, the Chinese Academy of Sciences and State Planning Commission, Beijing; 1996.
- [27] Tian DL, Pan HH, Kang WX, Fang HB Study on biomass of second generation of Chinese fir plantations. J Cent S Univ For Technol. 1998; 18(3):14–19
- [28] Feng ZW, Wang XK, Wu G. The biomass and productivity of forest ecosystem in China. Science Press, Beijing; 1999
- [29] Zeng WS, Luo QB, He DB. Study on compatible nonlinear tree biomass models. Chin J Ecol. 1999; 18(4):19–24
- [30] Tang SZ, Zhang HR, Xu H. Study on establishing and estimating method of compatible biomass model. Sci Silv Sin. 2000; 36(Sp.1):19–27
- [31] Liu GH, Fu BJ, Fang JY. Carbon dynamics of Chinese forests and its contribution to global carbon balance. Acta Ecol Sin. 2000; 20(5):733–740
- [32] Zhou YR, Yu ZL, Zhao SD. Carbon storage and budget of major Chinese forest types. Acta Phytoecologica Sinica. 2000; 24(5):518–522
- [33] Fang JY, Chen AP, Peng CH, Zhao SQ, Ci LJ. Changes in forest biomass carbon storage in China between 1949 and 1998. Science. 2001; 292:2320–2322

- [34] Xu B, Guo ZD, Piao SL, Fang JY. Biomass carbon stocks in China's forests between 2000 and 2050: a prediction based on forest biomass–age relationships. *Sci China Life Sci.* 2010; 53(7):776–783. DOI: 10.1007/s11427-010-4030-4
- [35] Li HK, Lei YC. Estimation and evaluation of forest biomass carbon storage in China. Chinese Forestry Press, Beijing; 2010, 60 p.
- [36] Li HK, Lei YC, Zeng WS. Forest carbon storage in China estimated using forest inventory data. *Sci Silv Sin.* 2011; 47(7):7–12
- [37] Li HK, Zhao PX, Lei YC, Zeng WS. Comparison on estimation of wood biomass using forest inventory data. *Sci Silv Sin.* 2012; 48(5):44–52
- [38] Zheng DX, Liao XL, Li CW, Ye QL, Chen PL. Estimation and dynamic change analysis of forest carbon storage in Fujian province. *Acta Agric Univ Jiangxiensis.* 2013; 35(1):112–116
- [39] Zeng WS, Tang SZ, Huang GS, Zhang M. Population classification and sample structure on modeling of single-tree biomass equations for national biomass estimation in China. *For Res Manage.* 2010; 3:16–23
- [40] Zeng WS, Tang SZ. Using measurement error modeling method to establish compatible single-tree biomass equations system. *For Res.* 2010; 23(6):797–803
- [41] Zeng WS, Zhang HR, Tang SZ. Using the dummy variable model approach to construct compatible single-tree biomass equations at different scales—a case study for Masson pine (*Pinus massoniana*) in southern China. *Can J For Res.* 2011; 41(7): 1547–1554. DOI: 10.1139/X11-068
- [42] Zeng WS. Methodology on modeling of single-tree biomass equations for national biomass estimation in China [thesis]. Chinese Academy of Forestry, Beijing; 2011.
- [43] Zhang LJ, Zeng WS, Tang SZ. Comparison of nonlinear regression equation with intercept and segmented modeling approach for estimation of single tree biomass. *For Res.* 2011; 24(4): 453–457
- [44] Zeng WS, Tang SZ. Establishment of belowground biomass equations for larch in north-eastern and Masson pine in southern China. *J Beijing For Univ.* 2011; 33(2):1–6
- [45] Zeng WS, Tang SZ. Modeling compatible single-tree biomass equations of Masson pine (*Pinus massoniana*) in southern China. *J For Res.* 2012; 23(4): 593–598. DOI: 10.1007/s11676-012-0299-4
- [46] Dang YF, Wang XJ, Zeng WS. Using segmented modeling approach to construct tree volume and biomass equations for larch in northeastern China. *For Res.* 2012; 25(5): 558–563
- [47] Zeng M, Nie XY, Zeng WS. Compatible tree volume and aboveground biomass equations of Chinese fir in China. *Sci Silv Sin.* 2013; 49(10):74–79. DOI: 10.11707/j.1001-7488.20131012

- [48] Fu LY, Zeng WS, Zhang HR, Wang GX, Lei YC, Tang SZ. Generic linear mixed-effects individual-tree biomass models for *Pinus massoniana* Lamb. in southern China. *South For.* 2014; 76(1):47–56. DOI: 10.2989/20702620.2013.870389
- [49] Zou WT, Zeng WS, Zhang LJ, Zeng M. Modeling crown biomass for four pine species in China. *Forests.* 2015; 6(2):433–449. DOI: 10.3390/f6020433
- [50] Zeng WS. Using nonlinear mixed model and dummy variable model approaches to construct origin-based single tree biomass equations. *Trees Struct Funct.* 2015; 29(1): 275–283. DOI: 10.1007/s00468-014-1112-0
- [51] Zeng WS. Integrated individual tree biomass simultaneous equations for two larch species in northeastern and northern China. *Scand J For Res.* 2015; 30(7): 594–604. DOI: 10.1080/02827581.2015.1046481
- [52] Zeng WS, Tang SZ. Bias correction in logarithmic regression and comparison with weighted regression for non-linear models. *Nat Prec.* 2011; Available from <http://dx.doi.org/10.1038/npre.2011.6708.1>
- [53] Zeng WS, Tang SZ. Goodness evaluation and precision analysis of tree biomass equations. *Sci Silvae Sin.* 2011; 47(11): 106–113
- [54] State Forestry Administration of China. Technical regulation on sample collections for tree biomass modeling. China Standard Press, Beijing; 2015; 11 p.
- [55] State Forestry Administration of China. Technical regulation on methodology for tree biomass modeling. China Standard Press, Beijing; 2015; 7 p.
- [56] State Forestry Administration of China. Tree biomass models and related parameters to carbon accounting for *Cunninghamia lanceolata*. China Standard Press, Beijing; 2015; 14 p.
- [57] State Forestry Administration of China. Tree biomass models and related parameters to carbon accounting for *Pinus massoniana*. China Standard Press, Beijing; 2015; 14 p.
- [58] State Forestry Administration of China. Tree biomass models and related parameters to carbon accounting for *Pinus yunnanensis*. China Standard Press, Beijing; 2015; 9 p.
- [59] State Forestry Administration of China. Tree biomass models and related parameters to carbon accounting for *Pinus tabulaeformis*. China Standard Press, Beijing; 2015; 9 p.
- [60] State Forestry Administration of China. Tree biomass models and related parameters to carbon accounting for *Pinus elliottii*. China Standard Press, Beijing; 2015; 9 p.
- [61] Jayaraman K. A statistical manual for forestry research. FAO Regional Office for Asia and the Pacific, Bangkok, 1999; 231 p.
- [62] Parresol BR. Assessing tree and stand biomass: a review with examples and critical comparisons. *For Sci.* 1999; 45: 573–593
- [63] Parresol BR. Additivity of nonlinear biomass equations. *Can J For Res.* 2001; 31: 865–878

- [64] IPCC. IPCC guidelines for national greenhouse gas inventories—agriculture, forestry and other land use (Volume 4). Published: IGES, Japan, 2006.
- [65] Pajtík J, Konôpka B, Lukac M. Biomass functions and expansion factors in young Norway spruce (*Picea abies* L. Karst) trees. For Ecol Manage. 2008; 256: 1096–1103
- [66] Wang XP, Fang JY, Zhu BA. Forest biomass and root–shoot allocation in northeast China. For Ecol Manage. 2008; 255: 4007–4020
- [67] Mugasha WA, Eid T, Bollandas OM, Malimbwi RE, Chamshama SAO, Zahabu E, Katani JZ. Allometric models for prediction of above- and belowground biomass of trees in the miombo woodlands of Tanzania. For Ecol Manage. 2013; 310: 87–101
- [68] Tang SZ, Lang KJ, Li HK. Statistics and computation of biomathematical models. Science Press, Beijing; 2008. 584 p.
- [69] Crecente-Campo F, Soares P, Tomé M, Diéguez-Aranda U. Modelling annual individual-tree growth and mortality of Scots pine with data obtained at irregular measurement intervals and containing missing observations. For Ecol Manage. 2010; 260: 1965–1974
- [70] Tang SZ, Li Y, Wang YH. Simultaneous equations, error-in-variable models and model integration in systems ecology. Ecol Model. 2001; 142: 285–294
- [71] West GB, Brown JH, Enquist BJ. A general model for the origin of allometric scaling laws in biology. Science. 1997; 276(5309): 122–126
- [72] West GB, Brown JH, Enquist BJ. A general model for the structure and allometry of plant vascular systems. Nature. 1999; 400: 664–667
- [73] Zeng WS, Tang SZ. A new general allometric biomass model. Nat Prec. 2011; Available from <http://dx.doi.org/10.1038/npre.2011.6704.2>
- [74] Henry M, Bombelli A, Trotta C, et al. GlobAllomeTree: international platform for tree allometric equations to support volume, biomass and carbon assessment. iForest (early view): e1–e5 [Internet]. 2013. Available from: <http://www.sisef.it/iforest/contents/?id=ifor0901-006> [Assessed 2016-09-04]

